Tgfbr1 controls developmental plasticity between the hindlimb and external genitalia by remodeling their regulatory landscape

Anastasiia Lozovska[1], Artemis G. Korovesi[1], André Dias[1#], Alexandre Lopes[1], Donald A. Fowler[1], Gabriel G. Martins[1], Ana Nóvoa[1], and Moisés Mallo[1*]

1 Instituto Gulbenkian de Ciência, Rua da Quinta Grande 6, 2780-156 Oeiras, Portugal

\# Current address: Department of Experimental and Health Sciences, Universitat Pompeu Fabra, Barcelona, Spain.

* Lead author: mallo@igc.gulbenkian.pt

Key words: Tgfbr1, hindlimb, external genitalia, chromatin conformation, mouse

The hindlimb and external genitalia of present-day tetrapods are thought
to derive from an ancestral common primordium that evolved to generate a
wide diversity of structures adapted for efficient locomotion and mating in
the ecological niche conquered by the species. We show that despite long
evolutionary distance from the ancestral condition, the early primordium of
the mouse external genitalia preserved the capacity to take hindlimb fates. In
the absence of *Tgfbr1*, the pericloacal mesoderm generates an extra pair of
hindlimbs at the expense of the external genitalia. It has been shown that the
hindlimb and the genital primordia share many of their key regulatory factors.
***Tgfbr1* controls the response to those factors by modulating the accessibility**
status of regulatory elements that control the gene regulatory networks
leading to the formation of genital or hindlimb structures. Our work uncovers
a remarkable tissue plasticity with potential implications in the evolution of
the hindlimb/genital area of tetrapods, and identifies a novel mechanism for
***Tgfbr1* activity that might also contribute to the control of other physiological**
or pathological processes.
36

Introduction

The vertebrate body is built sequentially in a head to tail progression. While this is a continuous process, it involves two transitions entailing major changes in gene regulatory mechanisms. The first transition results in the switch from a head to a trunk developmental program [1], marking the start of axial extension and the layout of the primordia for most of the organ systems involved in vital and reproductive functions. The second transition organizes the end of trunk structures and activates the tail developmental program [2]. The transition from trunk to tail entails significant reorganization of embryonic structures involving derivatives from all germ layers. The neuromesodermal competent (NMC) progenitors [3,4] relocate from the epiblast to the tail bud, from where they keep extending the body axis [5,6]. The progenitors for the lateral plate mesoderm, involved in the formation and vascularization of the trunk-resident organs [7], undergo terminal differentiation leading to the formation of the hindlimbs and of the ventral lateral mesoderm that will then become the pericloacal mesoderm [2]. Reciprocal interactions between the pericloacal mesoderm and the endodermal cloaca will then generate the genital tubercle (GT), the precursor of the external genitalia, while also organizing the exit channels of the digestive and excretory systems with the formation of the rectus/anus, the bladder and the urethra [8–12]. Although the hindlimb shares many of the patterning and morphogenetic processes with the forelimb [13], genetic analyses indicate major regulatory differences controlling the earliest stages of development of the two appendages [14–16]. Interestingly, molecular studies revealed that the hindlimb and the GT share many of their regulatory processes [9,17–19], indicating that they might be closely connected developmentally. Indeed, it has been reported that in some tetrapod species hindlimbs and genitals share a common primordium, which might represent the ancestral condition [20]. In mammals, it has been suggested that the posterior displacement of the cloaca relative to the hindlimb bud disconnected the limb primordium from the influence of the endodermal signals promoting genital fates, thus allowing formation of both legs and external genitalia [20].

Genetic studies, mostly in mouse embryos, revealed that the control of the early stages of hindlimb and cloacal/pericloacal development requires functional input from several regulatory factors. These include signaling pathways, like those of the transforming growth factor β/bone morphogenetic protein (Tgfβ/BMP)

superfamily [21–25], fibroblast growth factors (FGFs) [26–28], sonic hedgehog (Shh) [29–31]

and both canonical and non-canonical WNTs [10,11,32], as well as several transcription

factors, including Hoxa13, Hoxd13 [33,34] and Isl1[35]. Importantly, while the same

regulatory factors play essential roles in both structures [9,18], their functional

outcome differ substantially in the hindlimb and the GT, indicating the existence of

mechanisms canalizing distinct patterns of response in the hindlimb and the

pericloacal mesoderm when exposed to a particular regulatory factor. Shh provides

paradigmatic example of such functional duality. It determines anterior/posterior

limb bud polarity from the posterior limb bud mesenchyme regulating the identity of

the limb skeletal elements [36]. Conversely, from its expression domain in the

endodermal cloaca it controls the morphogenesis of the GT, as well as the

organization of the different outlets for the urogenital and intestinal systems [29,30].

The mechanisms regulating this distinct tissue response to common regulatory

factors are largely unknown.

Genetic analyses have shown that the Tgfβ receptor 1 (*Tgfbr1*) (also known

as *Alk5*) plays a key role in the activation of the trunk to tail transition, also initiating

the regulatory sequence controlling formation of the hindlimb and external genitalia

[2,37,38]. Indeed, *Tgfbr1* null mutant embryos fail to induce the primordia of the

hindlimb or cloacal/pericloacal structures, as shown by the absence of early

markers or morphological landmarks for these tissues [37]. Conversely, premature

activation of Tgfbr1 signaling in the axial progenitors resulted in earlier induction of

these structures, as estimated by gene marker expression and the substantial

anterior displacement of the hindlimb buds [2].

It has been shown that the Growth and Differentiation factor 11 (Gdf11) is a

physiological ligand for Tgfbr1 in its role at the trunk to tail transition [38]. Indeed,

Gdf11 mouse mutant embryos show a significant delay in the transition to tail

development [2,39]. Expression and functional analyses in several other vertebrates

indicate that this role for Gdf11 is not restricted to the mouse but might be shared

by most other tetrapods [40,41]. However, contrary to *Tgfbr1* mutant embryos, *Gdf11*

mutants still activate the transition to tail development, although with a significant

delay, indicating that other ligands in the Tgfβ/BMP family might cooperate with

Gdf11 in triggering Tgfbr1 activity in axial tissues. One such factor is Gdf8 as

revealed by the enhanced trunk size observed in *Gdf11*/*Gdf8* compound mutants

[42]. Bmp4 and Bmp7 also stand out as potential candidates to mediate some of the

Tgfbr1 activities in the caudal body. Both *Bmp7/Tsg* mutants and compound *Bmp7*

homozygous/*Bmp4* heterozygous conditional inactivation in the region of the trunk

to tail transition resulted in the fusion of the hindlimbs (sirenomelia) due to the

absence of ventral lateral mesoderm [22,23] and, thus, of the pericloacal mesoderm

that builds the GT [20], which is also absent from *Tgfbr1* null mutant embryos [21].

Although BMPs are not considered as canonical Tgfbr1 ligands [43], given the high

complexity of interactions between different receptors of the Tgfβ/BMP superfamily

[44–47], functional interactions between Tgfbr1 and Bmp4/Bmp7 cannot be ruled out,

most particularly because the bona fide receptors for those BMPs in the

pericloacal/hindlimb area await identification.

The essential role of Tgfbr1 signaling in the induction of both hindlimb and GT

primordia despite generating morphologically and functionally distinct structures,

prompted us to evaluate whether it also played a role at later stages of hindlimb

and/or GT development. For this we inactivated *Tgfbr1* in the caudal embryonic

region after the primordia for the hindlimb and GT were already laid down. We found

that in the absence of *Tgfbr1* the pericloacal mesoderm is converted into a second

set of hindlimbs. By comparing the chromatin accessibility profiles of wild type and

mutant tissues, we could observe that the mutant pericloacal mesoderm acquires a

significant number of limb-type signatures, while losing many of those specifically

associated with the GT. Our results indicate that Tgfbr1 modulates the type of

response of the pericloacal mesoderm to patterning signals from the endoderm by

remodeling the regulatory landscape of the chromatin, thus defining a limb or a GT-

type of gene activation.

Results

Generation of Tgfbr1-cKO developmental model.

For the conditional *Tgfbr1* inactivation approach we used three *Tgfbr1* alleles,

namely the *Tgfbr1* null allele [37] and either *Tgfbr1^flox^*, carrying LoxP sites flanking

Exon 3 [48], or *Tgfbr1^3ex3-flox^*, containing a triplicated Exon 3 each surrounded by LoxP

sites (supplementary Fig. 1). Characterization of mice homozygous for the

*Tgfbr1^3ex3-flox^* allele revealed that this allele generates a transcript containing a

tandem of three Exon 3 sequences in frame with the flanking Exons 2 and 4

(supplementary Fig. 1C). Expression in HEK293T cells of the $Tgfbr1^{3ex3}$ mRNA

isolated from the mouse embryos confirmed that it generates a protein product with

the expected size of Tgfbr1 containing triplicated Exon 3 sequences (supplementary

Fig. 1D). Exon 3 encodes the transmembrane domain and major part of the GS

domain of the protein, including the serin-rich region phosphorylated upon ligand

binding and implicated in signal transduction [43,49]. Homozygous $Tgfbr1^{3ex3-flox}$

animals were not viable, and $Tgfbr1^{3ex3-flox/-}$ fetuses phenocopied the knock out of

the Tgfbr1 ligand $Gdf11$ [2,39] (supplementary Fig. 1B, B'). Together, the above results

strongly suggest that $Tgfbr1^{3ex3-flox}$ produces a hypomorphic receptor.

$Tgfbr1$ inactivation was promoted using the $Cdx2Cre^{ERT}$ transgenic driver [2],

triggering Cre activity in caudal embryonic tissues by tamoxifen administration

(supplementary Fig. 1E). Experiments using either $Tgfbr1$ floxed allele gave similar

phenotypes, but while $Tgfbr1^{flox/-};Cdx2Cre^{ERT+/0}$ embryos required two consecutive

tamoxifen doses at embryonic stage (E) 6.75 and E7.25, a single tamoxifen injection

at E7.25 was sufficient to obtain the same mutant phenotype from for $Tgfbr1^{3ex3-flox/-}$

$;Cdx2Cre^{ERT+/0}$ embryos, thus reducing the frequency of tamoxifen-derived late fetal

miscarriage. Our tamoxifen treatment scheme inactivated $Tgfbr1$ in the tissues

caudal from the forelimb bud, as confirmed by RT-qPCR analyses on RNA

extracted from the posterior part of E10.5 embryos, including the hindlimbs

(Supplementary Fig. 1F, G). Tamoxifen-treated embryos and fetuses with either the

$Tgfbr1^{flox/-};Cdx2Cre^{ERT+/0}$ or the $Tgfbr1^{3ex3-flox/-};Cdx2Cre^{ERT+/0}$ genotype will be

hereafter referred to as $Tgfbr1$-cKO.

Tgfbr1-cKO exhibit multiple developmental malformations

The most prominent feature of $Tgfbr1$-cKO mutants was a duplication of their

hindlimbs [4/5 at E16.5] (Fig. 1A, B). The fifth fetus analyzed at this stage lacked

overt hindlimb duplication but still contained small protrusions posterior to the

hindlimbs, a phenotype likely derived from less efficient recombination. The limb

identity of the duplicated structures was confirmed by the presence of skeletal

structures that, although variable in morphology, were clearly identified as

belonging to limbs (Fig. 1C; supplementary Fig. 2; Supplementary movie 1). For

instance, a close look at one such duplicated hindlimb showed the presence

stylopod, zeugopod and autopod elements. When these last two sections were

analyzed in more detail using light sheet microscopy, in addition to the presence of

two skeletal elements in the zeugopod, we could identify individual phalanxes in
several of the fingers, although with somewhat abnormal patterns, as well as
polydactyly (supplementary Fig. 2).

Another prominent phenotype of *Tgfbr1-cKO* fetuses was the presence of an
omphalocele [5/5 at E16.5] (Fig. 1B). This phenotype, but not the duplicated
hindlimbs, was also observed in *Tgfb2/Tgfb3* double mutants [50]. This indicates both
that Tgfbr1 might mediate Tgfb2 and Tgfb3 activity in the body wall and that different
ligands should trigger Tgfbr1 signaling in the body wall and the pericloacal/hindlimb
mesoderm. The three *Tgfbr1-cKO* mutants analyzed either by optical projection
tomography (OPT) or by histological sections lacked kidneys (Fig. 1D-E';
Supplementary movies 2 and 3), a trait also observed in mutant fetuses for *Gdf11*
[2,39]. The cloacal-derived tissues were also strongly compromised in these mutant
embryos (Fig. 1D-E'; supplementary Fig. 3C,C'; Supplementary movies 2 and 3).
The gut failed to generate normal rectal-anal structures. Instead, the caudal end of
the intestinal tube merged with a structure that could be identified as a small bladder
connected to a hypomorphic urethra, thus resembling a persistent cloaca. The
external genitalia were also virtually absent from *Tgfbr1-cKO* fetuses. These
phenotypes are likely to result from interference with the functional interactions
between the cloacal endoderm and the pericloacal mesoderm driving normal
morphogenesis of the cloacal region [9,11,18,19,31]. Indeed, endodermal malformations
in this area were already evident in *Tgfbr1-cKO* mutants at mid gestation. At E10.5
and E11.5, the mutant cloaca was larger than in control embryos, lacked features
associated with normal septation, and often contained protrusions entering the
intra-cloacal lumen (2/4 embryos, supplementary Fig. 3a-B'). Remains of these
protrusions persisted later in development at E16.5 in the malformed bladder
(supplementary Fig. 3C, C').

The extra set of hindlimbs in Tgfbr1-cKO mutants derives from the pericloacal
mesoderm

Morphological alterations in the hindlimb/pericloacal area of *Tgfbr1-cKO*
embryos were already visible at mid-gestation (Fig. 2). *Fgf8* expression in E10.5
embryos revealed a posterior and medial expansion of the apical ectodermal ridge
(AER) of the hindlimb bud almost reaching the ventral midline of the embryo (Fig.
2A, A'), suggesting an extension of the hindlimb field into the pericloacal mesoderm

of *Tgfbr1-cKO* embryos. This was associated with several expression changes in the pericloacal region. For instance, hindlimb specific markers, like *Pitx1* and *Lin28a* were at E10.5 ectopically activated in the pericloacal region (Fig. 2B-C'). Conversely, *Tbx5* expression, normally observed in the pericloacal mesoderm, and later in the GT, but not in the hindlimb, could not be detected in this region of E11.5 *Tgfbr1-cKO* embryos (Fig 2G, G'), and *Fgf8* expression was absent from the cloacal endoderm of the mutant embryos (Fig 2A, A'). Together these observations suggest a change of identity in the pericloacal mesoderm, becoming incorporated into the hindlimb field instead of entering its normal GT developmental fate.

Expression of genes active in both the hindlimb and the pericloacal region/GT, was also consistent with the pericloacal tissues taking a hindlimb fate. For instance, *Wnt5a*, which in wild type embryos is expressed in the limb buds beneath the AER and in the pericloacal region and GT, was not detected next to the cloacal endoderm of E10.5 *Tgfbr1-cKO* embryos, being instead observed beneath the AER along the extended hindlimb domain into the pericloacal region (Fig. 2E, E'). This pattern also revealed that the hindlimb domain of *Tgfbr1-cKO* embryos become split in two prominences, the first indication of the generation of two individual hindlimb structures from extended hindlimb bud of the mutant embryos. *Fgf8* expression in this region of *Tgfbr1-cKO* mutants at E11.5 was also consistent with the split of the hindlimb bud in two fields, as it was observed through the apical border of the hindlimb region but demarcating two apparent prominences separated by a valley (Fig. 3A, A'). Additional support for the conversion of pericloacal mesoderm into hindlimb identity was provided by genes showing dynamic expression patterns in these structures. In wild type embryos, *Isl1* is initially activated in the hindlimb and GT primordia, becoming later downregulated in the hindlimb while keeping strong expression in the GT [51] (Fig 2H). At E11.5 *Tgbfr1-cKO* mutants showed residual *Isl1* expression mainly in the expanded endodermal cloaca (note that *Isl1* normal expression profile includes the cloacal endoderm [35]) but became downregulated in the adjacent mesoderm in the area likely to generate the extra hindlimb bud (Fig 2H'). Conversely, *Fgf10*, an essential gene for limb induction [52], whose expression in the GT is observed only at later stages [26], was strongly activated already at E10.5 following limb-like patterns, reaching mesodermal tissue adjacent to the endoderm (Fig. 2D, D').

Although clearly not as well defined as in wild type embryos, a degree of

anterior-posterior patterning could still be observed in the hindlimb region of *Tgfbr1-*

cKO embryos, consistent with changes in digit identity observed in the autopod

skeleton of the E16.5 fetuses (supplementary Fig. 2). For instance, at E10.5, *Hand2*

was expressed in the mutants as a continuous domain that run medially into

pericloacal tissue, showing a slightly stronger expression in the posterior area of

the two emerging prominences of the hindlimb bud, although not as clearly

restricted to the posterior limb bud mesenchyme observed in wild type embryos

(Fig. 2F, F'). A similar trend was observed at E11.5, when *Hand2* expression

followed a slight anterior-posterior distribution, being excluded from the anterior-

most region of the anterior limb bud prominence (Fig. 3B, B') and showed a slight

stronger expression in the posterior part of the caudal prominence of the mutant

limb bud (Fig. 3B, B'). Expression of the anterior marker *Pax9* was observed in the

anterior border of the anterior prominence of the limb bud in late E11.5 *Tgfbr1-cKO*

embryos (Fig. 3C, C'). However, we could not detect *Pax9* signal associated with

the posterior prominence. Whether this indicates real absence of *Pax9* expression

in this region or just delayed onset of expression remains to be determined.

Interestingly, the posterior prominence of the mutant hindlimb bud retained a

considerable degree of dorso-ventral patterning, as revealed by *En1* and *Lmx1b*

expression at E10.5 that followed typical limb patterns [53,54], although they were not

as well defined as in control hindlimb buds (Fig. 3D-e').

It has been shown that regulation of GT development relies largely on signals

from the cloacal endoderm, with Shh playing a central role in this process [30,31]. The

observation that *Wnt5a*, a known Shh target in the GT [31], was not activated in the

pericloacal mesoderm of *Tgfbr1-cKO* despite strong endodermal *Shh* expression

(Fig. 4A, A'; supplementary Fig. 3a-B'), suggested that either this mesenchyme

became refractory to Shh activity or that it changed its response profile. The

expression of the Shh target *Gli1* was consistent with the second possibility, as it

showed equivalent patterns in the pericloacal mesenchyme in both wild type and

Tgfbr1-cKO embryos (Fig. 4B, B'), thus indicating that this tissue still responds to

the Shh signal from the endoderm.

Reorganization of the chromatin regulatory landscape in the pericloacal mesoderm

of Tgfbr1-cKO embryos

We explored the mechanisms modulating the pericloacal mesoderm response
to Shh (and most likely also to other regulatory factors) by comparing the genomic
accessibility profiles in the posterior hindlimb prominences (extra hindlimbs) and
region adjacent to the cloaca (we will refer to it as mutant GT) of *Tgfbr1-cKO* with
those of hindlimb buds and GT of control embryos by ATAC-seq [55] (Fig. 4C).
Principal component (PC) analysis of those profiles revealed two main features.
PC1 grouped the tissues from *Tgfbr1-cKO* embryos together with the wild type GT
and clearly apart from the control hindlimb, consistent with the pericloacal origin of
the extra hindlimb of the mutant embryos. PC2, on the contrary, grouped the extra
hindlimb of the mutant embryos together the wild type hindlimb, separated from the
control GT (Fig. 4D).

We further explored the PC1 and PC2 by annotating 500 top peaks
contributing for each of the PCs (top loadings). For both PC1 and PC2, most of the
top loadings were in distal intergenic regions (46,2% and 48,6%, respectively), while
only small proportion of the peaks (6,8% for PC1 and 8,6% for PC2) were attributed
to potential promoter regions (<= 5 kb upstream of TSS) (Fig. 4E). This suggested
that most of the genomic regions whose accessibility patterns are influenced by the
Tgfbr1 might represent distal regulatory elements.

Hierarchical clustering of the top loadings contributing to the PC1 reveled the
existence of two distinct clusters: cluster 1 represent peaks less accessible in the
control limb than in other conditions, while cluster 2 contains the peaks more
accessible in the control limb than in other conditions (Fig 4F). These two clusters
represent limb specific regions regulated by Tgfbr1 signaling. Top loadings of PC2
formed 4 clusters based of their accessibility patterns (Fig. 4G). Clusters 1 and 3
are particularly interesting, because they represent the peaks with accessibility
patterns common between control and mutant limbs, and different from wild type
and mutant GT. These clusters, therefore, are likely to contain the regulatory
elements specifying limb fates and being responsible for canalizing the mutant
pericloacal mesenchyme towards the limb fate.

To further elaborate on the potential significance of conserved patterns within
top PC loadings clusters, we performed a series of complementary differential
analyses on our samples. We identified two accessibility patterns suggestive of their
involvement in the differential control of cell fate decisions of the pericloacal
mesoderm. First, we identified a set of peaks more accessible in the GT than in the

hindlimb samples from wild type embryos, and that showed significantly reduced

accessibility in the tissues from *Tgfbr1-cKO* embryos (1650 regions with

$Log_2FC>1.5$; FDR< $1e^{-3}$) (Fig. 4H, H'). These elements might represent enhancers

involved in the regulation of genes differentially expressed in the pericloacal region

of wild type embryos. The loss of accessibility to these elements in the *Tgfbr1-cKO*

mutants could thus contribute significantly to the inability of the pericloacal

mesoderm of the mutant embryos to enter their normal GT fate.

Another group of chromatin regions were significantly more accessible in

hindlimb-generating tissues of wild type and *Tgfbr1-cKO* embryos than in GT

tissues (526 peaks with $Log_2FC>1.5$; FDR<$1e^{-3}$ are shown in Fig. 5A, A'). These

regions represent a pattern analog to that observed in clusters 1 and 3 of PC2

whose accessibility patterns did not diverge between mutant and control hindlimb

($Log_2FC<0.5$; FDR>0.1), indicating that they might represent the most relevant

elements driving the mutant pericloacal mesenchyme towards limb identity. For

further description we will refer to the GT-specific elements as pattern 1 elements

and to those associated with hindlimb tissues pattern 2 elements.

We identified pattern 1 elements within the *Tbx5, Isl1* and *Wnt5a* genomic

regions, three of the pericloacal mesoderm markers down-regulated in the

pericloacal region of *Tgfbr1-cKO* mutants (Fig. 4I, J, K), which will serve as a proof

of principle for the involvement of pattern 1 elements in GT expression. These

elements mapped to highly conserved genomic areas, suggesting their possible

involvement in regulatory processes. When tested using a transgenic reporter

assay, one of the elements associated with *Tbx5* showed activity in the pericloacal

region [n=6/14] (Fig 4I'). Interestingly, it did not activate overt forelimb expression

(supplementary Fig. 4A), the major expression domain of this gene [14], further

indicating specificity for the pericloacal region. Similarly, the element identified

downstream of *Isl1* also largely reproduced *Isl1* expression in the GT when tested

in transgenic reporter assays [n=5/12] (Fig. 4J', Fig. 6B, and supplementary Fig.

4C). An element associated with *Wnt5a* was also able to activate expression in the

GT, although much less frequently than the other elements [n=2/18] (Fig 4K'). This

element was also active in the tail, but it did not promote expression in the hindlimb.

Together, these observations are consistent with these elements indeed

representing enhancers potentially involved in the expression of the relevant genes

in the pericloacal area.

We found pattern 2 elements in regions associated with genes playing
essential roles during the earliest stages of limb development, which could thus play
a relevant role in promoting limb fates from the pericloacal mesoderm (Fig. 5C, D).
As for pattern 1 elements, these regions were also highly conserved among
vertebrates. We found pattern 2 elements in the *Fgf10* genomic region (Fig. 5C), a
gene activated in the mesenchyme adjacent to the cloaca entering hindlimb fate in
the mutant embryo (Fig. 2D, D'). Interestingly, published ChIP-seq data from
forelimb buds [56,57] showed binding of Gli3 and Hoxa13 to one of these elements,
located within an intronic region of this gene (Fig. 5C), indicating that it might
respond to Shh and/or Hoxa13 activities in the developing limb buds. The lack of
accessibility of this element in the GT of wild type embryos might thus suggest that
Tgfbr1 signaling renders this enhancer blind to endodermal Shh and/or the strong
pericloacal Hoxa13 (and maybe also Hoxd13) expression [34], consistent with absent
Fgf10 at early stages in this tissue (Fig. 2D). Rather surprisingly, this element failed
to generate reporter activity in transgenic embryos [n=10]. We still do not
understand whether this reflects real lack of activity or the absence of proper
genomic context for their activity.

Another potentially relevant pattern 2 element was located within the *Fmn1*
locus, matching the position of one of the enhancers (element 7 in [58]) controlling
Grem1 in the limb buds (Fig 5D) that had been shown to respond to Shh activity
from the zone of polarizing activity (ZPA) [58,59]. In wild type embryos *Grem1*
expression is not detected in the pericloacal mesoderm or in the GT (Fig. 5E),
indicating that in these tissues *Tgfbr1* might render the *Grem1* regulatory elements
blind to endodermal Shh. The inaccessible state of this *Grem1* enhancer in the GT
(Fig. 5D) is consistent with this hypothesis. Conversely, the gain of accessibility of
the *Grem1* enhancer in the extra hindlimb suggests that in the absence of *Tgfbr1*
the pericloacal mesoderm could become responsive to the endodermal Shh.
Considering the capacity of *Grem1* to expand the expression of AER markers [60–62],
this gene is a candidate to contribute to the observed AER extension in the hindlimb
primordium of *Tgfbr1-cKO* embryos. *Grem1* expression analyses in *Tgfbr1-cKO*
embryos revealed unexpected complexities. Some features of the expression
pattern were consistent with *Grem1* becoming responsive to endodermal Shh, as
expression was detected in the caudal-most end of the expanded hindlimb next to
the endoderm (Fig. 5E'). Rather surprisingly, however, *Grem1* expression was

strongly reduced or absent throughout the mesoderm of the extended hindlimb bud

of *Tgfbr1-cKO* embryos (Fig. 5E'). *Grem1* limb expression depends on Shh activity

from the ZPA [58,63,64], suggesting that the absent *Grem1* expression resulted from

the abnormal *Shh* expression observed in the hindlimb buds of *Tgfbr1-cKO*

mutants, which followed spotty patterns beneath the AER instead of forming a well-

defined ZPA in the posterior limb bud mesenchyme (Fig. 4A', supplementary Fig.

3A').

We explored whether this abnormal *Shh* expression resulted from a functional

impact of the absence of Tgfbr1 activity on the ZRS, the regulatory element

controlling *Shh* expression in the ZPA [65]. As expected, this region was accessible

in wild type hindlimbs (Fig. 5F). Conversely, it was inaccessible in the wild type GT,

consistent with the absence of the *Shh* expression in the pericloacal mesoderm or

the GT mesoderm, despite the presence of relevant activators of this enhancer, like

Hand2, Hoxa13 and Hoxd13 [17,34,66,67] [the endodermal *Shh* expression is under the

control of a different enhancer [68], which was accessible in both GT samples

(supplementary Fig. 3D), consistent with conserved endodermal *Shh* expression in

the *Tgfbr1-cKO* embryos]. Remarkably, the ZRS was also non-accessible in the

hindlimb region of *Tgfbr1-cKO* embryos, consistent with the observed abnormal

Shh limb expression (Fig. 5F). Therefore, Tgfbr1 seems to also control ZRS

accessibility to their regulatory transcription factors. Interestingly, this enhancer is

part of an additional pattern (pattern 3) consisting of elements requiring Tgfbr1

activity to become accessible in the limb bud. Analysis of the ATAC-seq data

revealed that 5764 ($Log_2FC>1.5$; $FDR<1e^{-3}$) elements fit the criteria of this pattern

(Fig. 5B, B'). The complementary tissue specificities of patterns 1 and 3 indicates

context dependency for Tgfbr1 activity in the modulation of genomic configuration.

Footprinting analysis on elements differentially accessible in the GT.

The canonical Tgfbr1 signaling pathway involves activation of Smad2 or

Smad3 that recruit Smad4 and enter the nucleus where they interact with Smad-

responsive elements to regulate gene expression [69]. It has been shown that Smads

can interact with chromatin remodelers, like Foxh1 or components of the SWI/SNF

complex to promote accessibility of enhancers that are later regulated by Smad

transcriptional activity [70–72]. To understand whether a similar mechanism might be

involved in the reorganization of the regulatory landscape in the pericloacal

mesoderm we performed a transcription factor footprint analysis and subsequently

motif enrichment on the ATAC-seq profiles using HINT-ATAC [73]. We did not find

enrichment of Smad footprints in the accessible regions of the pericloacal

mesoderm of wild type embryos (supplementary Table 1). Similarly, the

transcription factor Foxh1, previously shown to interact with Smad proteins was also

not found enriched in these elements. These results indicate that the mechanism

by which Tgfbr1 signaling modulates accessibility of regulatory regions is likely to

be different from that described for the regulation by Tgfβ or Nodal in cell lines [70–

72]. The lack of enrichment for Smad footprints in the elements specifically accessible

in the GT of wild type embryos indicate that Smad-dependent signaling is unlikely

to be a major component participating in the promotion of genital fates in the

pericloacal mesoderm.

Additional analysis of these footprints revealed enrichment in signals

associated with factors known to play significant roles in the development of the

external genitalia, including Hoxa13, Hoxd13, Isl1, Six2 or effectors of the canonical

Wnt pathway like Lef1, Tcf7 or Tcf7l. As a proof of principle to understand the

relevance of these footprints in the regulation of pattern 1 elements, we focused on

a Lef1 binding motif identified in the TF footprint analysis within the *CR-Isl1*

chromatin region analyzed above (green rectangle in Fig. 6A). We tested the

functional relevance of this potential Lef1 binding site, by generating a mutant

version of this enhancer lacking this site (*CR-Isl1$^{\Delta Lef1}$*) and testing its activity in the

β-galactosidase transgenic reporter assay (Fig. 6C and supplementary Fig. 4D).

When compared to the wild type enhancer (Fig. 6B, Fig. 4J' and supplementary Fig.

4C) *CR-Isl1$^{\Delta Lef1}$* transgenic embryos showed reduced activity in the GT

mesenchyme, and no activity in the posterior part of the hindlimb typically observed

in the control embryos [6/15] (Fig. 6C and supplementary Fig. 4D). In addition, *CR-*

Isl1$^{\Delta Lef1}$ activated β-galactosidase expression in the otic vesicle (Fig. 6C and

supplementary Fig. 4D), likely derived from a conserved Gata3 motif within the TF

footprint (red rectangle in Fig. 6A) at level comparable to that observed in *CR-Isl1*

transgenics (Fig. 6B, supplementary Fig. 4C), which served as an internal control

to evaluate reporter expression. This is consistent with potential relevance of this

Lef1 binding site for the activity of this *Isl1* associated regulatory element, and thus,

to respond to canonical Wnt signaling in the GT. It is possible that the residual

activity of *CR-Isl1^{ΔLef1}* in the GT derived from the presence of an additional Lef1

binding motif slightly upstream on the TF footprint (blue rectangle in Fig. 6A). We

therefore tested the activity of a deletion mutant of this enhancer including both Lef1

binding motifs and the region between them (*CR-Isl1^{ΔΔLef1}*). Transgenic embryos for

this construct had no sign of activity in the GT, limbs, or the otic vesicle [2/7],

promoting instead extensive reporter activation throughout the embryo (Fig. 6D,

supplementary Fig. 4E). This pattern contrasts with those obtained with *CR-Isl1* or

CR-Isl1^{ΔLef1}, which were consistently almost restricted to the GT and otic vesicle.

Together, these results are compatible with the interpretation that Tgfbr1 signaling

renders the *CR-Isl1* enhancer accessible to its regulation by canonical Wnt

signaling in the pericloacal region.

Discussion

The capacity of the pericloacal mesoderm of mouse embryos to generate both

external genitalia and hindlimb structures reveals a remarkable degree of

developmental plasticity in this tissue. This finding provides a new dimension to the

developmental connection between the hindlimb and the GT, previously suggested

by the considerable overlap in their regulatory networks [9,17–19]. It should be noted

that *Tgfbr1-cKO* embryos develop two independent hindlimb structures despite

being first induced as a single bud extending into the pericloacal mesoderm. This

suggests differential segregation of the tissues originating from the pericloacal

mesoderm and the somatopleure-derived limb field, despite both entering limb

differentiation paths, which might indicate that limb and pericloacal mesodermal

tissues contain distinct cell affinity properties that keep each primordium as discrete

developmental units independently of the differentiation path they enter to generate

mature structures. Consistent with this interpretation the PC analysis of ATAC-seq

profiles showed that the tissue eventually generating the extra hindlimb keeps a

high degree of similarity with the GT of wild type embryos (PC1 in our analysis),

despite the simultaneous loss of some GT signatures and the acquisition of several

limb-like characteristics (PC2 in our analysis). Further experiments will be required

to explore the validity of this hypothesis.

The Tgfbr1-dependent differentiation plasticity of the pericloacal mesoderm,

together with its apparent inability to mix with the primordial limb mesoderm has

important implications for both the development and evolution of the limb and

genital area of tetrapods. It has been suggested that in mice the posterior

displacement of the developing cloaca relative to the hindlimb was key to

disconnect the hindlimb field from the influence of the cloacal signals that induce

the GT, thus allowing the somatic lateral plate mesoderm and the ventral lateral

mesoderm entering hindlimb and genital fates, respectively [20]. This contrasts with

the position of those structures in squamates, in which the cloaca is formed just

adjacent to the hindlimb field, a configuration that has been suggested to promote

development of paired external genitalia (hemipenises) instead of hindlimbs from

these primordia in snakes [20]. However, similar proximity was also observed in

lizards that develop both paired external genitalia and hindlimbs [20]. In addition, and

most importantly, the phenotype of the *Tgfbr1-cKO* embryos shows that the

mesoderm adjacent to the cloaca can form limb structures, indicating that the

intrinsic properties of the mesoderm are key determinants of the differentiation route

it will enter. It will be therefore interesting to determine whether a mechanism related

to the developmental plasticity uncovered by our work, including changes in cell

affinity properties of prospective hindlimb and genital primordia, could help

explaining the absence of hindlimbs in snakes but their presence in most lizards.

We showed that Tgfbr1 modifies the profile of chromatin accessibility in the

pericloacal mesoderm. The involvement of Tgfβ/BMP signaling in chromatin

remodeling has been reported before for Tgfβ and Nodal signals [70–72]. In those

cases, this resulted from the interaction of Smads with chromatin remodelers like

Foxh1 or components of the SWI/SNF complex, which would turn Smad-responsive

enhancers accessible and enable the Smad transcriptional activity [70–72]. Although a

similar mechanism could contribute to Tgfbr1 activity in the pericloacal and limb

mesoderm, it is unlikely to represent a major component of this activity. First, many

chromatin regions gain accessibility in the absence of *Tgfbr1*, indicating that Tgfbr1

activity can also render chromatin regions non-accessible. In addition, we did not

detect enrichment of Smad binding to elements rendered accessible by *Tgfbr1*

using HINT-ATAC [73] on the ATAC-seq profiles. Instead, this analysis showed some

enrichment in footprints for factors known to play important roles in GT

development. Even considering the limitations of the bioinformatic approach, these

observations indicate that Tgfbr1 modulates the catalog of regions accessible to

regulatory factors, not restricted to the downstream effectors of the Tgfbr1 pathway.
Our work thus suggests a model according to which *Tgfbr1* controls the differential
morphogenesis of the limb and GT primordia by determining the set of enhancers
accessible or inaccessible to mediate the activity of patterning inputs. This
regulatory mode gains relevance when considering that many of these inputs are
common to both structures, despite generating mature structures as anatomically
different as the hindlimbs and external genital structures [9,17–19]. This idea can be
illustrated by two examples. One of them concerns the *Grem1* enhancer, involved
in *Grem1* activation in the limb bud in response to Shh from the ZPA [58]. This
enhancer is non accessible in the pericloacal mesoderm of wild type embryos, fitting
with the lack of Grem1 expression in the pericloacal mesoderm of wild type embryos
despite the strong Shh expression in the endodermal cloaca. It however became
accessible in the Tgbfr1-negative mesoderm generating extra hindlimb, and *Grem1*
expression was detected adjacent to the Shh source in the endoderm. The second
example refers to the *Isl1* GT enhancer identified in this work. While we found no
evidence of direct involvement of effectors of the Tgfbr1 pathway in its activation,
our data indicate that Tgfbr1 renders this enhancer accessible to regulation by the
canonical Wnt signaling, a pathway that has been shown to participate in GT
development [11].

The identification of chromatin elements uniquely associated with hindlimb
and GT fates might guide the search for the molecular signature that specifies
hindlimb or GT development. The large number of elements in each group suggest
that these signatures might be quite complex, likely involving the combined activity
of several factors. Identification of those factors will be challenging, considering that
in mammals enhancers are typically located far from the genes they regulate, which
are often not even the closest transcription unit [74].

An intriguing observation from our experiments was that *Tgfbr1-cKO* embryos
were still able to generate mature limb structures that, although malformed, still
contained most limb elements despite *Shh* being downregulated in the hindlimb
bud. This phenotype contrasts with the absence of most distal elements in *Shh*
mutant embryos [75]. It has been shown that a brief pulse of Shh activity during the
initial bud stage is enough to promote normal limb development [59]. As the ZRS
seems to be silent in *Tgfbr1-cKO* mutants, it is possible that the initial Shh input
required to set limb development in motion is provided by the endoderm, facilitated

by the extension of the hindlimb field into the pericloacal mesoderm. This would

argue in favor of limb polarizing activity intrinsic to cloacal endoderm [31], and against

exclusively pro-genital regulatory potential of cloaca [20].

A challenging question left open by our work is deciphering the mechanism(s)

by which Tgfbr1 activity controls such large-scale remodeling of the regulatory

landscape in the target tissues. The lack of enrichment in footprints for Tgfbr1

effectors in the target regions makes a direct activity of the canonical Tgfbr1

pathway in this process unlikely. Instead, it is more probable that this regulation is

indirect. Whatever the mechanism or mechanisms, it requires a high degree of

coordination, which would still be more challenging if several mechanisms are

involved in the generation of the different patterns (e.g. if the control of accessibility

and inaccessibility relies on independent mechanisms). Identification of those

mechanisms and determining whether they also operate in other physiological and

pathological processes under the control of members of the Tgfβ/BMP signaling

family might have far reaching implications for our understanding of morphogenetic

processes and disease.

Materials and methods

Ethics statement

All animal procedures were performed in accordance with Portuguese

(Portaria 1005/92) and European (directive 2010/63/EU) legislations and guidance

on animal use in bioscience research. The project was reviewed and approved by

the Ethics Committee of "Instituto Gulbenkian de Ciência" and by the Portuguese

National Entity "Direcção Geral de Alimentação Veterinária" (license reference:

014308).

Mice

Generation of the *Tgfbr1^Flox^* mouse lines was previously reported [48]. *Tgfbr1^3ex3-*

flox resulted from the same experiments in which *Tgfbr1^flox^* was generated, identified

upon sequencing through the *Tgfbr1* locus, which revealed that the exon 3 had been

triplicated and the presence of LoxP sites in the introns between the triplicated

exons and in those connecting them with exon 2 and exon 4. The *Cdx2Cre^ERT^* mice

were previously reported [2]. Mice were genotyped from ear or digit biopsies.

Samples were incubated in 50 μL of the PBND buffer (50 mM KCl, 10 mM Tris-HCl,

pH 8.3, 2.5 mM MgCl$_2$, 0.1 mg/mL gelatin, 0.45% NP40, 0.45% Tween 20)

supplemented with 100 μg/mL of proteinase K at 55°C overnight. Proteinase K was

inactivated at 95°C for 15 minutes and 1 μL of genomic DNA was used for

genotyping by PCR. *Cdx2Cre^{ERT};Tgfbr1$^{+/-}$* mice were genotyped for the *Cre* allele

and *Tgfbr1* null allele; *Tgfbr1$^{flox/+}$* and *Tgfbr1$^{3ex3-flox/+}$* mice were genotyped with

primers surrounding 3'-LoxP (primers listed in supplementary table 2).

Tgfbr1-cKO (*Cdx2Cre^{ERT};Tgfbr1$^{flox/-}$* or *Cdx2Cre^{ERT};Tgfbr1$^{3ex3-flox/-}$*) embryos

were obtained by crossing *Tgfbr1$^{flox/flox}$* or *Tgfbr1$^{3ex3-flox/+}$* females with *Tgfbr1$^{+/-}$*

;Cdx2Cre^{ERT} males. Noon of the day of the plug was considered E0.5. To induce

recombination pregnant females were treated with tamoxifen (Sigma, T5648)

dissolved in corn oil. Tamoxifen was administered at 0.1 mg per gram of body

weight by oral gavage either once at E6.75 when *Tgfbr1$^{3ex3-flox}$* females were

involved, or twice, at E6.75 and E7.25 when *Tgfbr1^{flox}* females were involved.

Embryos were obtained by cesarean section and processed for further

analyses (see below). They were genotyped from their yolk sacs for Cre, *Tgfbr1$^-$*

and *Tgfbr1^{flox}* when analyzing *Tgfbr1-cKO* embryos or for β-galactosidase when

analyzing the reporter transgenic embryos (primers listed in supplementary table

2). Yolk sacs were collected in 50 μL of yolk sac lysis buffer (50 mM KCl, 10 mM

Tris-HCl, pH8.3, 2 mM MgCl$_2$, 0.45% Tween-20, 0.45% NP40) supplemented with

100 μg/mL of proteinase K and incubated at 55°C overnight. Samples were heat-

deactivated as previously described and used for PCR.

Transgenic embryos

To generate transgenic constructs, the relevant enhancers (genomic

coordinates shown in supplementary table 3) were amplified by PCR (primers in

supplementary table 2) from genomic DNA and cloned into a vector containing the

adenovirus major late promoter, the coding region of the β-galactosidase gene and

the SV40 polyadenylation signal [2]. The enhancers were confirmed by direct

sequencing. The constructs were isolated from the plasmid backbone, gel-purified

using the NZYGelpure (NZYTech #MB01102) and eluted from the columns with 50

μL of the kit's elution buffer. The purified constructs were diluted in microinjection

buffer (10 mM Tris.HCl, 0.25 mM EDTA, pH 7.5) at 2 ng/μL and microinjected into

the pronucleus of fertilized FVB/N oocytes according to standard protocols [76].

Microinjected oocytes were transferred into the uteri of pseudopregnant NMRI

females and embryos were recovered at E10.5, E11.5 or E12.0 and stained for β-

galactosidase activity.

RT-qPCR

To assess recombination efficiency, the posterior part of the E10.5 *Tgfbr1-*

cKO embryos, including their hindlimbs, were dissected in PBS (137 mM NaCl, 2.7

mM KCl, 10 mM Na_2HPO_4, and 1.8 mM KH_2PO_4) on ice. Whole E10.5 *Tgfbr1^{-/-}*

embryos were used as controls. RNA was extracted from fresh tissue using TRI

reagent (Sigma, #T9424) according to manufacturer's protocol. 200 ng of RNA from

each sample was used to prepare complementary DNA using a random hexamer

mix (NZYTech #MB12901) and following the protocol of the NZY Reverse

Transcriptase enzyme (NZYTech #MB12401). All cDNA samples were diluted 1:5

and 1 μL was used in quantitative PCR (qPCR) reactions with iTaq Universal SYBR

Green Supermix (Bio-Rad, #1725124), according to manufacturer's instruction and

using the Applied Biosystems QuantStudio 7 flex qPCR System. Primers were

localized in exon 3 and exon 4 of *Tgfbr1* mRNA (listed in supplementary table 2),

so that in case of successful recombination the product is not amplified.

PCR product concentration in each sample was calculated using the standard

curve with following formula *Quantity = 10^(mean Ct-b/m)* (where "m" is slope and

"b" is an intercept of the standard curve), and *Tgfbr1* expression level was

normalized to *β-Actin* expression. Linear model was fitted (Tgfbr1 normalized

expression ~ Genotype) with R built-in function (lm). Analysis of variance (ANOVA)

was performed on the linear model with the built-in function in R (anova).

Significance levels between groups of samples were assessed by Tukey method

using the "glht" function from "multicomp" package in R. Differences were

considered significant at $*p<0.05$, $**p<0.01$ and $***p<0.001$. Boxplots were

generated with "ggplot2" and "ggdignif" packages in R.

Cell culture and transfection

HEK 293T cells (ATCC #CRL-3216) were cultured in Dulbecco's modified

Eagle's medium (DMEM, Life Technologies) supplemented with 10% fetal bovine

serum (FBS, Life Technologies) and 1% penicillin/streptomycin at 37 °C in a 5% CO_2 atmosphere. Cells were seeded in 35 mm plates and when they reached 70% confluency, the medium was changed to transfection medium (DMEM supplemented with 10% FBS). Expression vectors were constructed by amplifying the coding region of the *Tgfbr1* transcript isolated from wild type or *Tgfbr1*^{3ex3-flox} homozygous embryos by RT-PCR (primers in supplementary table 2) and cloned into the pRK5 expression vector. Constructs were verified by sequencing. 3 μg of the plasmids was transfected into the 393T cells using Lipofectamine 2000 (Thermo Fisher Scientific #11668027) according to the manufacturer's instructions. After 24 hours incubation, cells were processed for Western blot analysis.

Protein extraction and western blot

The transfected cells were washed twice with ice-cold PBS and then 150 μL of lysis buffer (50 mM Tris-HCl, 150 mM NaCl and 1% Triton X-100, pH8.0) was added to each plate and incubated on ice for 10 minutes. Lysates were scraped into microcentrifuge tubes on ice and centrifuged at 20000 rcf for 10 min at 4˚C. The supernatant was collected, frozen on dry ice and stored at -80˚C. Protein lysates were mixed (2:1) with 3X loading buffer (549 mM Tris HCl, 563 mM Tris Base, 2.05 mM EDTA, 8 % LDS (Lithium dodecyl sulfate), 10 % Glycerol, 0.75 % SERVA Blue G250, 0.25% Phenol red, 160 mM DTT). Samples were incubated for 15 minutes at 65˚C and resolved by SDS-PAGE in a 10% polyacrylamide gel. Proteins were transferred to PVDF membranes in 20% methanol, 25 mM Tris, 200 mM glycine at 200 mA for 1 hour. The membranes were blocked in blocking solution [5% dry milk dissolved in PBS containing 0.1% Tween-20 (PBT)] for 1 hour at room temperature and then incubated with primary antibodies overnight at 4˚C. Primary antibodies were anti-Tgfbr1 (Sigma-Aldrich #HPA056473, 1:1000 in blocking solution) and anti-actin (Abcam #ab179467, 1:1000 in blocking solution). Membranes were then washed in PBT and incubated with HRP conjugated-anti-rabbit IgG (GE Healthcare # NA9340, 1:5000 in blocking solution) at room temperature for 1 hour and washed twice in PBT. Signals were developed by chemiluminescence using Amersham ECL Prime Western Blotting Detection Reagent (GE Healthcare #RPN2232) and images were captured using the GE Amersham Imager 600.

Skeletal preparation

E17.5 fetuses were recovered from pregnant females by cesarean section and dissected from the extraembryonic membranes, eviscerated, the skin removed and then fixed in 100% ethanol for 2 days. Cartilages are then stained by incubation with 450 mg/L of alcian blue (Sigma, #A5268) in 80% ethanol/20% acetic acid for one day. Fetuses were then postfixed in 100% ethanol overnight. Tissue was cleared by incubation in 2% KOH for 6 hours, followed by staining of ossified bones with a 50 mg/l of alizarin red S (Sigma, #130-22-3) solution in 2% KOH for 3 hours. Specimens were then further incubated in 2% KOH until tissues were fully cleared. Skeletons were stored in 25% glycerol in water. All incubations were performed at room temperature with rolling. Genotyping was performed on gut tissue by incubating in Laird's buffer (100 mM Tris-HCl pH 8.5, 5 mM EDTA, 0.2% SDS, 200 mM NaCl) containing 10 μg/mL proteinase K at 55ºC overnight. Genomic DNA was then recovered by precipitation with isopropanol (1:1, vol:vol) and transferred to TE pH 8.0. Genotyping was then performed by PCR using oligos specified in supplementary table 2.

OPT and SPIM imaging

Optical projection tomography (OPT) and SPIM (Selective Illumination Plane Microscopy) were used to image E16.5 fetuses as previously described [77] with minor modifications. Briefly, fetuses were recovered by cesarean section in ice cold PBS, washed several times in PBS and fixed in 4% paraformaldehyde (PFA) in PBS at 4ºC for several days. After several washes with demineralized water fetuses were dehydrated by sequential incubation in demineralized water with increasing concentrations of methanol (10% increases) and then twice in 100% methanol. Fetuses were then bleached in a three-day process with a sequence of 2.5%, 5% and 10% hydrogen peroxide in methanol at room temperature. Fetuses were then rehydrated through a reverse methanol/demineralized water series and embedded in a 0,7% low-melting agarose. Clearing was done with a 1:2 solution of Benzoic Alcohol:Benzyl Benzoate (BABB) using a BABB/methanol series (25% BABB increases). 100% BABB was introduced on day 3 and replaced every day for the next 4 days, until the fetuses became completely transparent. Anatomical datasets were then obtained using a custom built OPT/SPIM scanner and procedures;

briefly, for OPT green auto-fluorescence was acquired on 1600 sequential angles

for a full revolution, the raw dataset was pre-processed with a custom-built ImageJ

macro, and then back-projection reconstructed using SkyScan's "nrecon" as in [78],

and then post-processed to reduce noise and enhance contrast in ImageJ. 3D

reconstruction and visualization of internal organs was performed using the Amira

software (Thermo Fisher Scientific). The autopods were also imaged with SPIM at

higher magnification to be able to reconstruct the digits and bones, also using

Amira. To facilitate the segmentation of the bones we used the Biomedisa online

tool for machine-learning-assisted interpolation after manual segmentation of a few

sparse sections [79]. The correlated OPT and SPIM datasets, and their segmented

volumes, were then assembled into a correlative dataset in Amira, to produce the

final 3D reconstructions.

Histological analysis

E16.5 fetuses were fixed in Bouin's fixative at room temperature for 2 days,

dehydrated by repeat washes in 100% ethanol for 5 days and embedded in paraffin.

Paraffin blocks were then sectioned at 10 µm and stained with Masson's trichrome

according to standard methods.

Whole mount in situ hybridization and sectioning

Embryos were fixed in 4% PFA overnight, then dehydrated through 25%, 50%

and 75% series of methanol in PBT (PBS, 0,1% Tween 20), then incubated in 100%

methanol. Embryos were then rehydrated through a reverse methanol/PBT series

and incubated three times in PBT for at least 5 min. each at room temperature.

Embryos were then bleached for 1 hour in 6% hydrogen peroxide in PBT and

permeabilized in 10 µg/mL of proteinase K in PBT for a period that depended on

the embryo size. The reaction was then quenched with a 2 mg/mL solution of

glycine in PBT, washed twice in PBT and postfixed in a 4% PFA and 0,2%

glutaraldehyde mix for 20 minutes, followed by two washes in PBT. Hybridization

was performed at 65°C overnight in hybridization solution (50% formamide, 1.3 x

SSC pH 5.5 [20 x SSC is 3M NaCl, 300 mM sodium citrate], 5 mM EDTA, 0.2 %

Tween 20, 50 µg/mL yeast tRNA, 100 µg/mL heparin) containing the relevant

digoxigenin-labelled antisense RNA probes. The probes used in this work are listed

in supplementary table 2. RNA probes were transcribed in vitro from the vector for

1.5 hours at 37˚C with relevant RNA polymerase and the DIG RNA Labeling Mix

(Roche #11277073910). The reaction product was verified in 0,8% agarose gel,

precipitated with ethanol in the presence of 0.3M sodium acetate, pH 5.3 and

resuspended in 100 μL of TE pH 8.0. Probes were diluted in hybridization solution

(6 μL in 1 ml of solution). After hybridization, embryos were washed twice with

hybridization solution without tRNA and heparin, at 65˚C for 30 min each, then in a

1:1 mix of hybridization solution and TBST (25 mM Tris.HCl, pH 8.0, 140 mM NaCl,

2.7 mM KCl, 0.1% Tween 20) for 30 minutes at 65ºC, and finally washed three times

with TBST at room temperature. Embryos were then equilibrated in MABT (100 mM

Maleic acid, 150 mM NaCl, 0.1 % Tween-20, pH 7.5) and blocked in MABT blocking

buffer [MABT containing 1% blocking reagent (Roche #11096176001)] with 10%

sheep serum for 2,5 hours at room temperature. Embryos were then incubated

overnight at 4ºC with a 1:2000 dilution of alkaline phosphatase-conjugated anti-

digoxigenin antibody (Roche #11093274910) in MABT blocking buffer with 1%

sheep serum. After extensive washes with MABT at room temperature, embryos

were equilibrated in NTMT buffer (100 mM Tris HCl, pH 9.5, 50 mM $MgCl_2$, 100 mM

NaCl, 0.1% Tween-20) and developed with a 1:50 dilution of NBT/BCIP solution

(Roche #11681451001) in NTMT at room temperature in the dark. The reaction was

stopped with PBT, and embryos postfixed with 4% PFA overnight at room

temperature and stored in PBT. Pictures were taken with a SWIFTCAM SC1803

connected to a stereoscope. Stained embryos were embedded in a mix containing

0,45% gelatin, 27% bovine serum albumin (BSA), 18% sucrose, jellified with 1,75%

glutaraldehyde and sectioned at 35 μm on a Leica Vibratome VT 1000 S.

β-galactosidase staining

Embryos were dissected out in ice cold PBS and fixed in 4% PFA at 4ºC for

30 minutes. They were then washed three times in β-gal wash buffer (PBS plus

0.02% Tween 20) for 10 minutes at room temperature and β-galactosidase activity

developed in PBS containing 5 mM $K_3Fe(CN)_6$, 5 mM $K_4Fe(CN)_6$, 2 mM $MgCl_2$,

0.02% Tween 20, 0.4 mg/ml X-gal (Promega #V3941) overnight in the dark at 37ºC.

The reaction was stopped with β-gal wash buffer, embryos postfixed in 4% PFA

overnight at room temperature and stored in PBS.

ATAC-Seq

ATAC-seq was performed as in [55] with minor modifications. E11.25 mouse embryos were dissected on ice in DMEM High Glucose medium (Biowest #L0102-500) containing 10% FBS (this will be referred to as medium in the rest of the protocol). Hindlimbs and genital tubercles were dissected out and collected on ice-cold media. Single cell suspension was prepared by treating the tissue with 500 μL of Accutase (Sigma #A6964-500ML) for 30 minutes at 37˚C with shaking at 600 rpm. Single cells were pelleted at 6000 rcf for 5 minutes at 4˚C, resuspended in 200 μL of media and counted. 50000 viable cells from each sample were used for nuclei extraction. Cells were incubated with 50 μL of ATAC resuspension buffer (10 mM Tris HCl pH 7.4, 10 mM NaCl, 3 mM $MgCl_2$, 0.1% NP-40, 0.1% Tween-20 (Sigma-Aldrich, P7949), 0.01% Digitonin (Target Mol #282T2721-1ml/10mM in DMSO) for 3 minutes on ice. Lysis was quenched by adding 1 ml of ATAC resuspension buffer without NP-40 and Digitonin. Nuclei were pelleted by centrifugation at 500 rcf for 10 minutes at 4˚C. The pellet was then resuspended in 50 μL of transposition buffer [for 50 μL: 25 μL 2x TD buffer, 2.5 μL Tn5 transposition enzyme (100 nM final) (Illumina #15028212), 16.5 μL PBS, 0.5 μL 1% digitonin, 0.5 μL 10% Tween-20, 5 μL H_2O] and incubated at 37˚C for 30 minutes with shaking at 1000 rpm. The DNA was purified with Qiagen MinElute PCR purification kit (Qiagen #50928004) and eluted in 20 μL of the kit's elution buffer. Libraries were amplified by PCR with NEBNext High-Fidelity 2X PCR Master Mix (New England Biolabs #174M0541S) for 9 cycles in 96-well Thermal cycler (Biorad) and purified with Qiagen MinElute PCR purification kit. Tagmentation efficiency was assessed on TapeStation 4200 (Agilent). Double size selection to remove primer dimers and fragments exceeding 1 kb was performed using SPRIselect beads (Beckman Coulter). Another quality control was performed with High Sensitivity DNA assay using the Fragment Analyzer (Agilent). The 4nM libraries pool was sequenced with Illumina NextSeq 2000 (100 cycles, Pair-end 50 bp). Two biological replicates were performed per tissue.

The data obtained in this work has been deposited in the GEO repository, (accession number GSE231592).

ATAC-Seq bioinformatic and statistical analysis

Data bioinformatic analysis was performed on the Galaxy server [80]. Raw sequencing fastq files for each library were assessed for quality, adapter content and duplication rates with FastQC (v0.11.9) (http://www.bioinformatics.babraham.ac.uk/projects/fastqc). Adapters were trimmed and reads with length < 20 bp were removed using cutadapt (v1.16.5)[81] (3´ adapter sequence: CTGTCTCTTATACACATCT). The trimmed reads were aligned to the mouse reference genome (GRCm38/mm10 Dec. 2011) using Bowtie2 (v2.4.2)[82] [parameters paired-end options (-X/–maxins 1000, --fr, --dovetail), --very-sensitive]. The aligned reads were filtered using BamTools Filter (v2.4.1) [83] (parameters --isProperPair true, --mapQuality ≥ 30, and --reference!=chrM) and the duplicate reads were removed using Picard MarkDuplicates (v2.18.2.2) (http://broadinstitute.github.io/picard/). The resulting BAM files were converted to BED and the reads that overlap to the blacklisted regions listed in https://www.encodeproject.org/files/ENCFF547MET/ were removed using Bedtools (v2.30.0) [84]. The filtered BED files were used as inputs for peak calling. Peaks were called from individual replicates using MACS2 callpeak (v2.1.1) [85] (parameters --format single-end BED, --nomodel, --extsize 200, --shift -100, --qvalue 0.05).

BAM files were then converted to BigWig with "deeptools" [86] normalizing to 1x effective genome size (2652783500 for GRCm38/mm10). BigWig files were used for data visualization in IGV with the addition of the conservation scoring by phyloP (phylogenetic p-values)[87] for 60 vertebrate genomes from the UCSC genome browser (mm10.60way.phyloP60way.bw file downloaded from http://hgdownload.cse.ucsc.edu/goldenpath/mm10/phyloP60way/) and ChIP-seq data from Hoxa13 (GSE81356) [56] and Gli3 (GSE133710) [57].

The raw count matrix was created by merging peaks across samples as in [88,89]. DESeq2 object was built from the raw count matrix and blind dispersion estimated with the "vst" function from the "DESeq2" package in R [90]. Top 2500 most variable peaks estimated with the "rowVars" function in R were selected for the principal component analysis (PCA). PCA was performed using the built-in "prcomp" function in R. 500 top loadings contributing to PC1 and PC2 were annotated using ChIPseeker package in R [91]. Hierarchical clustering was performed on the Z-score normalized reads. Correlation distances between the peaks were

computed with "as.dist" function and clustering was performed using complete

linkage method with "hclust" function. Clusters were defined using a threshold h =

0.6. For statistical analysis raw reads were analyzed using EdgeR [92]. ANOVA-like

test was performed on all samples and pairwise comparisons between the samples

were made by analyzing individual contrasts. Venn diagrams were created from the

lists of selected differentially accessible peaks with "eulerr" package in R. Heat

Maps were made from data frames containing log2 normalized counts across

samples with the "pheatmap" package (https://CRAN.R-

project.org/package=pheatmap).

Footprinting analysis of ATAC-Seq data

We further used the program RGT HINT-ATAC [73] and motif matching and

enrichment to identify, plot and compare transcription factors (TF) footprints in the

different samples. In detail, the BAM files containing the filtered aligned reads for

each biological replicate were merged and used as a matrix for the footprinting

analysis for the regions corresponding to the peaks called by MACS2 (BED file)

(described above). The BAM and BED files for the peaks called in the control GT

and the subset of the BAM and BED files corresponding to the regions that were

found accessible only in GT (pattern 1 peaks) were used for the footprint analysis

and then tested for motif matching to the Hocomoco database[93] for finding the motif-

predicted binding sites (mpbs) independently. Motif enrichment analysis was

performed on the footprinting results for the peaks of pattern 1, using as a

background the footprinting results for all the peaks called in control GT. Position

frequency matrices for Lef1 (MA0768.2) and Gata3 (MA0037.4) binding motifs in

Mus Musculus were obtained from Jaspar database.

Acknowledgements

We would like to thank the members of the Mallo lab, especially Ana Casaca

for continuous support at different stages of this project, the IGC mouse facility for

their help with animal housing, histopathology facility for their help with histology

procedures, and genomics facility for helping with the ATAC-seq procedures. This

project was funded by Fundação para a Ciência e a Tecnologia (FCT) grants

2022.01629.PTDC to MM, PTDC/BII-BTI/32375/2017 to GGM, and PhD fellowships

PD/BD/128437/2017 to AL and PD/BD/128426/2017 to AD, by the PhD fellowship

Author contribution

Conceptualization: A.L. and M.M.; Formal Analysis: A.L. and A.K.**;** Funding
Acquisition: M.M.; Investigation: A.L., A.K. and M.M.; Methodology: A.L., A.D.
G.G.M. A.N., A.Lo. and D.F.; Writing – Original Draft Preparation: A.L. and M.M.;
Writing – Review & Editing: A.L., A.K. and M.M.

Competing interest declaration

The authors declare no competing interests.

References

1.   Duarte, P., Brattig Correia, R., Nóvoa, A. & Mallo, M. Regulatory changes associated
with the head to trunk developmental transition. *BMC Biol* **21**, 170 (2023).
2.   Jurberg, A. D., Aires, R., Varela-Lasheras, I., Nóvoa, A. & Mallo, M. Switching axial
progenitors from producing trunk to tail tissues in vertebrate embryos. *Dev Cell* **25**,
451–462 (2013).
3.   Binagui-Casas, A., Dias, A., Guillot, C., Metzis, V. & Saunders, D. Building consensus
in neuromesodermal research: Current advances and future biomedical
perspectives. *Curr Opin Cell Biol* **73**, 133–140 (2021).
4.   Tzouanacou, E., Wegener, A., Wymeersch, F. J., Wilson, V. & Nicolas, J. F. Redefining
the Progression of Lineage Segregations during Mammalian Embryogenesis by
Clonal Analysis. *Dev Cell* **17**, 365–376 (2009).
5.   Wymeersch, F. J., Wilson, V. & Tsakiridis, A. Understanding axial progenitor biology
in vivo and in vitro. *Development* **148**, dev180612 (2021).
6.   Wilson, V., Olivera-Martinez, I. & Storey, K. G. Stem cells, signals and vertebrate
body axis extension. *Development* **136**, 2133–2133 (2009).
7.   Prummel, K. D., Nieuwenhuize, S. & Mosimann, C. The lateral plate mesoderm.
Development (Cambridge) **147**, dev175059 (2020).
8.   Seifert, A. W., Harfe, B. D. & Cohn, M. J. Cell lineage analysis demonstrates an
endodermal origin of the distal urethra and perineum. *Dev Biol* **318**, 143–152
(2008).
9.   Yamada, G. *et al.* Molecular genetic cascades for external genitalia formation: An
emerging organogenesis program. *Developmental Dynamics* **235**, 1738–1752
(2006).
10. Miyagawa, S. *et al.* Disruption of the temporally regulated cloaca endodermal β-
catenin signaling causes anorectal malformations. *Cell Death Differ* **21**, 990–997
(2014).

11. Miyagawa, S. *et al.* Dosage-dependent hedgehog signals integrated with Wnt/β-
catenin signaling regulate external genitalia formation as an appendicular program.
Development **136**, 3969–3978 (2009).

12. Huang, Y. C., Chen, F. & Li, X. Clarification of mammalian cloacal morphogenesis
using high-resolution episcopic microscopy. *Dev Biol* **409**, 106–113 (2016).

13. Tickle, C. How the embryo makes a limb: Determination, polarity and identity. *J
Anat* **227**, 418–430 (2015).

14. Rallis, C. *et al.* Tbx5 is required for forelimb bud formation and continued
outgrowth. *Development* **130**, 2741–2751 (2003).

15. Naiche, L. a & Papaioannou, V. E. Tbx4 is not required for hindlimb identity or post-
bud hindlimb outgrowth. *Development* **134**, 93–103 (2007).

16. Naiche, L. A. & Papaioannou, V. E. Loss of Tbx4 blocks hindlimb development and
affects vascularization and fusion of the allantois. *Development* **130**, 2681–2693
(2003).

17. Kondo, T., Zakany, J., Innis, J. W. & Duboule, D. Of fingers, toes and penises [1].
Nature **390**, 29 (1997).

18. Cohn, M. J. Development of the external genitalia: Conserved and divergent
mechanisms of appendage patterning. *Developmental Dynamics* **240**, 1108–1115
(2011).

19. Lin, C. *et al.* Delineating a Conserved Genetic Cassette Promoting Outgrowth of
Body Appendages. *PLoS Genet* **9**, e1003231 (2013).

20. Tschopp, P. *et al.* A relative shift in cloacal location repositions external genitalia in
amniote evolution. *Nature* **516**, 391–394 (2014).

21. Lozovska, A. *et al.* Tgfbr1 regulates lateral plate mesoderm and endoderm
reorganization during the trunk to tail transition. *BioRxiv* (2023)
doi:10.1101/2023.08.22.554351.

22. Zakin, L., Reversade, B., Kuroda, H., Lyons, K. & De Robertis, E. Sirenomelia in Bmp7
and Tsg compound mutant mice: requirement for Bmp signaling in the
development of ventral posterior mesoderm. *Development* **132**, 2489–2499
(2005).

23. Suzuki, K. *et al.* Reduced BMP Signaling Results in Hindlimb Fusion with Lethal
Pelvic/Urogenital Organ Aplasia: A New Mouse Model of Sirenomelia. *PLoS One* **7**,
e43453 (2012).

24. Suzuki, K. *et al.* Regulation of outgrowth and apoptosis for the terminal appendage:
External genitalia: Development by concerted actions of BMP signaling.
Development **130**, 6209–6220 (2003).

25. Kajioka, D. *et al.* Bmp4 is an essential growth factor for the initiation of genital
tubercle (GT) outgrowth. *Congenit Anom (Kyoto)* **60**, 15–21 (2020).

26. Satoh, Y. *et al.* Regulation of external genitalia development by concerted actions
of FGF ligands and FGF receptors. *Anat Embryol (Berl)* **208**, 479–486 (2004).

27. Haraguchi, R. *et al.* Molecular analysis of external genitalia formation: The role of
fibroblast growth factor (fgf) genes during genital tubercle formation. *Development*
127, 2471–2479 (2000).

28. Seifert, A. W., Yamaguchi, T. & Cohn, M. J. Functional and phylogenetic analysis
shows that Fgf8 is a marker of genital induction in mammals but is not required for
external genital development. *Development* **136**, 2643–2651 (2009).

29. Seifert, A. W., Bouldin, C. M., Choi, K. S., Harfe, B. D. & Cohn, M. J. Multiphasic and
tissue-specific roles of sonic hedgehog in cloacal septation and external genitalia
development. *Development* **136**, 3949–3957 (2009).

30. Haraguchi, R. *et al.* Unique functions of sonic hedgehog signaling during external
genitalia development. *Development* **128**, 4241–4250 (2001).

31. Perriton, C. L., Powles, N., Chiang, C., Maconochie, M. K. & Cohn, M. J. Sonic
hedgehog signaling from the urethral epithelium controls external genital
development. *Dev Biol* **247**, 26–46 (2002).

32. Yamaguchi, T. P., Bradley, A., McMahon, A. P. & Jones, S. A Wnt5a pathway
underlies outgrowth of multiple structures in the vertebrate embryo. *Development*
126, 1211–1223 (1999).

33. Warot, X., Fromental-Ramain, C., Fraulob, V., Chambon, P. & Dollé, P. Gene dosage-
dependent effects of the Hoxa-13 and Hoxd-13 mutations on morphogenesis of the
terminal parts of the digestive and urogenital tracts. *Development* **124**, 4781–4791
(1997).

34. Scott, V., Morgan, E. A. & Stadler, H. S. Genitourinary functions of Hoxa13 and
Hoxd13. *Journal of Biochemistry* vol. 137 671–676 Preprint at
https://doi.org/10.1093/jb/mvi086 (2005).

35. Su, T. *et al.* LIM homeodomain transcription factor Isl1 affects urethral epithelium
differentiation and apoptosis via Shh. *Cell Death Dis* **10**, 713 (2019).

36. Tickle, C. & Towers, M. Sonic hedgehog signaling in limb development. *Front Cell
Dev Biol* **5**, 14 (2017).

37. Dias, A. *et al.* A Tgfbr1/Snai1-dependent developmental module at the core of
vertebrate axial elongation. *Elife* **9**, e56615 (2020).

38. Andersson, O., Reissmann, E. & Ibáñez, C. F. Growth differentiation factor 11 signals
through the transforming growth factor-beta receptor ALK5 to regionalize the
anterior-posterior axis. *EMBO Rep* **7**, 831–837 (2006).

39. McPherron, A. C., Lawle, A. M. & Lee, S.-J. Regulation of anterior/posterior
patterning of the axial skeleton by growth/differentiation factor 11. *Nat Genet* **22**,
260–264 (1999).

40. Matsubara, Y. *et al.* Anatomical integration of the sacral-hindlimb unit coordinated
by GDF11 underlies variation in hindlimb positioning in tetrapods. *Nat Ecol Evol* **1**,
1392–1399 (2017).

41. Aires, R. *et al.* Oct4 Is a Key Regulator of Vertebrate Trunk Length Diversity. *Dev Cell*
38, 262–274 (2016).

42. McPherron, A. C., Huynh, T. V & Lee, S.-J. Redundancy of myostatin and
growth/differentiation factor 11 function. *BMC Dev Biol* **9**, 24 (2009).

43. Heldin, C.-H. & Moustakas, A. Signaling receptors for TGF-β family members. *Cold
Spring Harb Perspect Biol* **8**, 1–33 (2016).

44. Goumans, M.-J. *et al.* Activin Receptor-like Kinase (ALK)1 Is an Antagonistic
Mediator of Lateral TGF/ALK5 Signaling. *Mol Cell* **12**, 817–828 (2003).

45. Carvalho, R. L. C. *et al.* Compensatory signalling induced in the yolk sac vasculature
by deletion of TGFβ receptors in mice. *J Cell Sci* **120**, 4269–4277 (2007).

46. Ramachandran, A. *et al.* TGF-β uses a novel mode of receptor activation to
phosphorylate SMAD1/5 and induce epithelial-to-mesenchymal transition. *Elife* **7**,
e31756 (2018).

47. Wang, W. *et al.* The TGF β type I receptor TGF β RI functions as an inhibitor of BMP
signaling in cartilage. *Proceedings of the National Academy of Sciences USA* **116**,
15570–15579 (2019).
48. Pineault, K. M., Novoa, A., Lozovska, A., Wellik, D. M. & Mallo, M. Two
CRISPR/Cas9-mediated methods for targeting complex insertions, deletions, or
replacements in mouse. *MethodsX* **6**, 2088–2100 (2019).
49. Wieser, R., Wrana, J. L. & Massague, J. GS domain mutations that constitutively
activate T beta R-I, the downstream signaling component in the TGF-beta receptor
complex. *Embo J* **14**, 2199–2208 (1995).
50. Dünker, N. & Krieglstein, K. Tgfβ2-/-Tgfβ3-/- double knockout mice display severe
midline fusion defects and early embryonic lethality. *Anat Embryol (Berl)* **206**, 73–
83 (2002).
51. Yang, L. *et al.* Isl1 Cre reveals a common Bmp pathway in heart and limb
development. *Development* **133**, 1575–1585 (2006).
52. Min, H. *et al.* Fgf-10 is required for both limb and lung development and exhibits
striking functional similarity to Drosophila branchless. *Genes Dev* **12**, 3156–3161
(1998).
53. Davis, C. A., Holmyard, D. P., Millen, K. J. & Joyner, A. L. Examining pattern
formation in mouse, chicken and frog embryos with an En-specific antiserum.
Development **111**, 287–298 (1991).
54. Riddle, R. D. *et al.* Induction of the LIM Homeobox Gene Lmx7 by WNT7a
Establishes Dorsoventral Pattern in the Vertebrate Limb. *Cell* **83**, 631–640 (1995).
55. Corces, M. R. *et al.* An improved ATAC-seq protocol reduces background and
enables interrogation of frozen tissues. *Nat Methods* **14**, 959–962 (2017).
56. Sheth, R. *et al.* Distal Limb Patterning Requires Modulation of cis-Regulatory
Activities by HOX13. *Cell Rep* **17**, 2913–2926 (2016).
57. Lex, R. K. *et al.* GLI transcriptional repression regulates tissue-specific enhancer
activity in response to hedgehog signaling. *Elife* **9**, e50670 (2020).
58. Malkmus, J. *et al.* Spatial regulation by multiple Gremlin1 enhancers provides digit
development with cis-regulatory robustness and evolutionary plasticity. *Nat
Commun* **12**, 5557 (2021).
59. Zhu, J., Patel, R., Trofka, A., Harfe, B. D. & Mackem, S. Sonic hedgehog is not a limb
morphogen but acts as a trigger to specify all digits in mice. *Dev Cell* **57**, 2048–2062
(2022).
60. Michos, O. *et al.* Gremlin-mediated BMP antagonism induces the epithelial-
mesenchymal feedback signaling controlling metanephric kidney and limb
organogenesis. *Development* **131**, 3401–3410 (2004).
61. Nissim, S., Hasso, S. M., Fallon, J. F. & Tabin, C. J. Regulation of Gremlin expression
in the posterior limb bud. *Dev Biol* **299**, 12–21 (2006).
62. Farin, H. F. *et al.* Tbx2 Terminates Shh/Fgf Signaling in the Developing Mouse Limb
Bud by Direct Repression of Gremlin1. *PLoS Genet* **9**, e1003467 (2013).
63. Zuniga, A. *et al.* Mouse limb deformity mutations disrupt a global control region
within the large regulatory landscape required for Gremlin expression. *Genes Dev*
18, 1553–1564 (2004).
64. Li, Q. *et al.* A Gli silencer is required for robust repression of gremlin in the
vertebrate limb bud. *Development (Cambridge)* **141**, 1906–1914 (2014).

65. Lettice, L. A. *et al.* A long-range Shh enhancer regulates expression in the
developing limb and fin and is associated with preaxial polydactyly. *Hum Mol Genet*
12, 1725–1735 (2003).

66. Galli, A. *et al.* Distinct roles of Hand2 in initiating polarity and posterior Shh
expression during the onset of mouse limb bud development. *PLoS Genet* **6**,
e1000901 (2010).

67. Leal, F. & Cohn, M. J. Loss and Re-emergence of Legs in Snakes by Modular
Evolution of Sonic hedgehog and HOXD Report Loss and Re-emergence of Legs in
Snakes by Modular Evolution of Sonic hedgehog and HOXD Enhancers. *Current*
Biology **26**, 1–8 (2016).

68. Sagai, T. *et al.* A cluster of three long-range enhancers directs regional Shh
expression in the epithelial linings. *Development* **136**, 1665–1674 (2009).

69. Jia, S. & Meng, A. TGF β family signaling and development. *Development* **148**,
dev188490 (2021).

70. Coda, D. M. *et al.* Distinct modes of SMAD2 chromatin binding and remodeling
shape the transcriptional response to NODAL/Activin signaling. *Elife* **6**, 1–31 (2017).

71. Coda, D. M. *et al.* A network of transcription factors governs the dynamics of
NODAL/Activin transcriptional responses. *J Cell Sci* **135**, jjcs259972 (2022).

72. Xi, Q., He, W., Zhang, X. H. F., Le, H. Van & Massague, J. Genome-wide impact of
the BRG1 SWI/SNF chromatin remodeler on the transforming growth factor β
transcriptional program. *Journal of Biological Chemistry* **283**, 1146–1155 (2008).

73. Li, Z. *et al.* Identification of transcription factor binding sites using ATAC-seq.
Genome Biol **20**, 45 (2019).

74. Liang, L. *et al.* Complementary Alu sequences mediate enhancer–promoter
selectivity. *Nature* **619**, 868–875 (2023).

75. Chiang, C. *et al.* Manifestation of the limb prepattern: Limb development in the
absence of sonic hedgehog function. *Dev Biol* **236**, 421–435 (2001).

76. Hogan, B., Beddington, R., Constantini, F. & Lacy, E. *Manipulating the mouse*
embryo: a laboratory manual. (Cold Spring Harbour Laboratory Press, 1994).

77. Dias, A., Martins, G. G., Lopes, A. & Mallo, M. Three and Four-Dimensional
Visualization and Analysis Approaches to Study Vertebrate Axial Elongation and
Segmentation. *Journal of Visualized Experiments* **168**, e62086 (2021).

78. Martins, G. G. *et al.* Optical projection tomography. in *Imaging Modalities for*
Biological and Preclinical Research: A Compendium, Volume 1: Part I: Ex vivo
biological imaging (eds. Walter, A., Mannheim, J. G. & Caruana, C. J.) i.2.f1-i.2.f11
(IOP Publishing, 2021). doi:10.1088/978-0-7503-3059-6ch12.

79. Lösel, P. D. *et al.* Introducing Biomedisa as an open-source online platform for
biomedical image segmentation. *Nat Commun* **11**, 5577 (2020).

80. Afgan, E. *et al.* The Galaxy platform for accessible, reproducible and collaborative
biomedical analyses: 2016 update. *Nucleic Acids Res* **44**, W3–W10 (2016).

81. Martin, M. Cutadapt removes adapter sequences from high-throughput
sequencing reads. *EMBnet J* **17**, 10–12 (2011).

82. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat*
Methods **9**, 357–359 (2012).

83. Barnett, D. W., Garrison, E. K., Quinlan, A. R., Strömberg, M. P. & Marth, G. T.
Bamtools: A C++ API and toolkit for analyzing and managing BAM files.
Bioinformatics **27**, 1691–1692 (2011).

84.   Quinlan, A. R. & Hall, I. M. BEDTools: A flexible suite of utilities for comparing
1100   genomic features. *Bioinformatics* **26**, 841–842 (2010).

1101   85.   Feng, J., Liu, T., Qin, B., Zhang, Y. & Liu, X. S. Identifying ChIP-seq enrichment using
1102   MACS. *Nat Protoc* **7**, 1728–1740 (2012).

1103   86.   Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing
1104   data analysis. *Nucleic Acids Res* **44**, W160–W165 (2016).

1105   87.   Pollard, K. S., Hubisz, M. J., Rosenbloom, K. R. & Siepel, A. Detection of nonneutral
1106   substitution rates on mammalian phylogenies. *Genome Res* **20**, 110–121 (2010).

1107   88.   Rausch, T., Hsi-Yang Fritz, M., Korbel, J. O. & Benes, V. Alfred: Interactive multi-
1108   sample BAM alignment statistics, feature counting and feature annotation for long-
1109   and short-read sequencing. *Bioinformatics* **35**, 2489–2491 (2019).

1110   89.   Erarslan-Uysal, B. *et al.* Chromatin accessibility landscape of pediatric T-
1111   lymphoblastic leukemia and human T-cell precursors. *EMBO Mol Med* **12**, e12104
1112   (2020).

1113   90.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
1114   dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

1115   91.   Wang, Q. *et al.* Exploring Epigenomic Datasets by ChIPseeker. *Curr Protoc* **2**, e585
1116   (2022).

1117   92.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor package for
1118   differential expression analysis of digital gene expression data. *Bioinformatics* **26**,
1119   139–140 (2009).

1120   93.   Kulakovskiy, I. V. *et al.* HOCOMOCO: Towards a complete collection of transcription
1121   factor binding models for human and mouse via large-scale ChIP-Seq analysis.
1122   *Nucleic Acids Res* **46**, D252–D259 (2018).

1123   94.   Davis, C. A. & Joyner, A. L. Expression patterns of the homeo box-containing genes
1124   and the proto-oncogene int-1 diverge during mouse development. *Genes Dev* **2**,
1125   1736–1744 (1988).

1126   95.   Crossley, P. H. & Martin, G. R. The mouse Fgf8 gene encodes a family of
1127   polypeptides and is expressed in regions that direct outgrowth and patterning in
1128   the developing embryo. *Development* **121**, 439–451 (1995).

1129   96.   Srivastava, D. *et al.* Regulation of cardiac mesodermal and neural crest
1130   development by the bHLH transcription factor, dHAND. *Nat Genet* **16**, 154–160
1131   (1997).

1132   97.   Aires, R. *et al.* Tail Bud Progenitor Activity Relies on a Network Comprising Gdf11,
1133   Lin28, and Hox13 Genes. *Dev Cell* **48**, 383–395 (2019).

1134   98.   Szeto, D. P. *et al.* Role of the Bicoid-related homeodomain factor Pitx1 in specifying
1135   hindlimb morphogenesis and pituitary development. *Genes Dev* **13**, 484–494
1136   (1999).

1137   99.   Echelard, Y. *et al.* Sonic hedgehog, a member of a family of putative signaling
1138   molecules, is implicated in the regulation of CNS polarity. *Cell* **75**, 1417–1430
1139   (1993).

1140   100.   Chapman, D. L. *et al.* Expression of the T-box family genes, Tbx1-Tbx5, during early
1141   mouse development. *Developmental Dynamics* **206**, 379–390 (1996).

1142

1143
1144

Figures

| Control | *Tgfbr1-cKO* | *Tgfbr1-cKO* | Control | *Tgfbr1-cKO* |

Figure 1. Malformations in the E16.5 *Tgfbr1-cKO* fetuses. (A, B) Images of fixed control (A) and *Tgfbr1-cKO* (B) fetuses. The mutant fetus shows the presence of an omphalocele (asterisk) and hindlimb duplication. (C) 3D reconstruction of the limb skeleton of a *Tgfbr1-cKO* fetus obtained by OPT and after segmentation of the limb skeleton. Extra hindlimbs are in magenta. Ossification shown in yellow. (D–E') 3D reconstruction of organs and the excretory outlets of the control (D, D') and *Tgfbr1-cKO* (E, E') fetuses. Images were obtained by OPT, followed by segmentation of the relevant structures. D' E' show virtual sections of the segmented 3D specimen. Asterisk in E' shows the gut-bladder connection in the mutant. K: kidney; G: gut; Bl: bladder; Ov: ovary Ut: uterus; V: vagina; Ur: urethra.

1159
Figure 2. The pericloacal mesenchyme of the *Tgfbr1-cKO* embryos adopts hindlimb fate.
Ventral views of the hindlimb/cloacal region of wild type (A-H) or *Tgfbr1-cKO* embryos (A'-
H') stained for *Fgf8* (E10.5)(A, A'), *Lin28a* (E10.5) (B, B'), *Pitx1* (E10.5) (n=2; C, C'), *Fgf10*
(E10.5) (D, D'), *Wnt5a* (E11.0) (E, E'), *Hand2* (E11.0) (F, F'), *Tbx5* (E11.5) (G, G') or *Isl1*
(E11.5) (H, H'). *Fgf8* expression shows posterior and medial extension of the AER (black
arrow) and the absence of expression in the cloacal endoderm (black arrowhead). Hindlimb
markers *Lin28a*, *Pitx1* and *Fgf10* extend into the pericloacal region *Tgfbr1-cKO* embryos
(white arrows). *Wnt5a* and *Hand2* are expressed in the posterior prominence of the *Tgfbr1-*
cKO hindlimbs, while forming clear separation of the hindlimb and GT domains in the
control embryos (yellow arrowheads). Pericloacal expression of *Wnt5a*, *Tbx5* and *Isl1* is
lost in the mutant embryos (yellow arrows). *Isl1* expression in cloacal endoderm is marked
by white arrowhead. At least n=3 embryos were analyzed per probe, unless stated
otherwise.
1173

Figure 3. Limb patterning in the *Tgfbr1-cKO* embryos. (A, A') Expression of *Fgf8* in E11.5 control (A) and *Tgfbr1-cKO* (A') embryos. The black arrow shows separation between the two hindlimb prominences. (B, B') Expression of *Hand2* in E11.5 control (B) and *Tgfbr1-cKO* (B') embryos. White arrow in (B') shows a posterior expression domain of the prospective mutant extra hindlimb. (C, C') Anterior *Pax9* expression in E11.5 control (C) and *Tgfbr1-cKO* (C') hindlimbs. A single expression signal is observed in the hindlimb region of *Tgfbr1-cKO* embryo corresponding to the most anterior part of the structure (white arrowhead). (D-e') Dorsal-ventral polarity in the *Tgfbr1-cKO* mutant's hindlimb. Expression of the ventral limb marker *En1* in E10.5 control (D, d) and *Tgfbr1-cKO* (D', d') embryos. (D, D') show lateral views of the hindlimb region in whole mount embryos; (d, d') show transversal sections through the region indicated in D, D'. Black arrowheads show *En1* expression in the ventral ectoderm of the limb bud. Expression of the dorsal mesodermal limb marker *Lmx1b* in E10.5 control (E, e) and *Tgfbr1-cKO* (E', e') embryos. (E, E') show lateral views of the hindlimb region in whole mounted embryos; (e, e') show transversal sections through the region indicated in E, E'. Yellow arrowheads show *Lmx1b* expression in the dorsal mesenchyme of the limb bud. n=3 embryos were analyzed per probe.

Figure 4. Tgfbr1 regulates the chromatin landscape in the pericloacal region. (A–B') The pericloacal mesenchyme of *Tgfbr1-cKO* embryos responds to Shh signaling from the cloacal endoderm. *Shh* expression in E11.5 control (A) and *Tgfbr1-cKO* (A') embryos. *Gli1* expression in E10.5 control (B) and *Tgfbr1-cKO* (B') embryos (n=3 embryos were analyzed per probe). (C) Schematic representation of the tissue dissected for the ATAC-seq experiments in control (top) and mutant (bottom) embryos. (D) PCA plot of the ATAC-seq samples. (E) Frequencies at the annotated genomic regions of 500 top loadings from the PC1 (grey) and PC2 (yellow). (F, G) Hierarchical clustering (HC) of PC1 (F) and PC2 (G) 500 top loadings. Heatmaps show scaled normalized counts. (H, H') Chromatin regions following pattern 1. (H) Venn diagram showing interception of the regions tested more accessible in control GT than in other tissues by pairwise comparisons [FDR<0.001, logFC>1.5]. (H') Heatmap showing log2 normalized counts across samples of the regions following pattern 1. (I, J, K) ATAC-seq tracks showing accessible chromatin in control GT, but not in the other samples in a region 93.7 kb upstream of the *Tbx5* TSS (I), 24.0 kb downstream of the *Isl1* TSS (J), and 91.7 kb upstream of the *Wnt5a* TSS (K). The regions highlighted with the blue shade are highly conserved among placental animals (bottom track). Two independent biological replicates were analyzed per tissue in the ATAC-seq experiments. (I', J', K') The highlighted elements drive β-galactosidase expression in the GT region (white arrows). Hoxa13 and Gli3 ChiP-Seq tracks from forelimb buds (GSE81356 and GSE133710, respectively) are shown in yellow and magenta, respectively. None of the analyzed enhancers showed enrichment in Hoxa13 or Gli3 binding activity in the forelimbs, consistent with the GT specificity of those elements. cHL: control hindlimb; mExHL: mutant extra hindlimb; cGT: control GT; mGT: mutant GT; CR: conserved region.

1219
Figure 5. (A, A') Chromatin regions following pattern 2 and pattern 3. (A) Venn diagram
showing interception between the regions more accessible in mutant extra limb than in
mutant GT [FDR<0.001, logFC>1.5] and more accessible in control hindlimb than in control
GT [FDR<0.001, logFC>1.5], but at the same time not differentially accessible between
mutant extra hindlimb and control hindlimb [FDR>0.01, logFC<0.5]. (A') Heatmap showing
log2 normalized counts from genomic regions following pattern 2 across the samples. (B,
B') Chromatin regions following pattern 3. (B) Venn diagram showing interception between
regions tested more accessible in control hindlimb than in other tissues by pairwise
comparisons [FDR<0.001, logFC>1.5]. (B') Heatmap showing log2 normalized counts for
the chromatin regions following pattern 3 across samples. (C, D) ATAC-Seq profiles
showing *Fgf10* putative regulatory element (C) and *Grem1* enhancer (D) that gains
accessibility in the mutant extra hindlimb (highlighted with the blue shadow). Also shown
are ChiP-Seq tracks of ChIP seq data from wild type forelimbs for Gli3 (yellow) and Hoxa13
(magenta) (obtained from GSE81356 and GSE133710, respectively), indicating their
binding to those elements in the forelimb bud. The lower track shows conservation in
placental mammals. (E, E') *Grem1* expression in E10.5 wild type (E) or *Tgfbr1-cKO* (E')
embryos showing the absence of pericloacal expression in wild type embryos and ectopic
activation in the caudal-most region of the pericloacal mesoderm of the mutant embryo

(white arrow). *Grem1* is expressed in the developing limb bud of the control embryo, but it
is almost not detectable in the developing limb buds of the mutant embryo (white
arrowheads) (n=3 embryos are analyzed per probe). (F) ATAC-seq data through the ZRS
(highlighted with the blue shadow), showing that it is accessible in the control limb but not
in any of the other samples. Two independent biological replicates were analyzed for all
ATAC-seq experiments. CR: conserved region; cHL: control hindlimb; mExHL: mutant extra
hindlimb; cGT: control GT; mGT: mutant GT.
1245

1246
Figure 6. *CR-Isl1* is a potential Wnt-responsive element. (A) Schematic representation of
the *CR-Isl1* region of Fig. 4J. The top panel shows the ATAC-seq signal in control GT and
the placental conservation score. The footprint corresponded to a ~65 bp region containing
a Gata3 binding motif (red), and Lef1 binding motif (green). An additional Lef1 binding motif
(blue) was identified 21 bp upstream of the Gata3 motif, although it was not included within
the footprint. The genomic sequences of these motifs are shown along with position
frequency matrices (PFMs). (B-D) Activation of the β-galactosidase reporter by the *CR-Isl1*
(B), *CR-Isl1$^{\Delta Lef1}$* (C), and *CR-Isl1$^{\Delta\Delta Lef1}$* (D) elements in E11.5 embryos. Black arrowheads
show expression in otic vesicle; white arrow indicates expression in the posterior limb bud;
black arrow shows expression in the GT.
1257

A

$Tgfbr1^{flox}$

$Tgfbr1^{3ex3-flox}$

$Tgfbr1^{3ex3}$ mRNA

Control $Tgfbr1^{3ex3/-}$

B   **B'**

T13

L6

T16

L9

C $Tgfbr1^{3ex3}$ Wild type

2000
1500

D $Tgfbr1^{3ex3}$ $Tgfbr1^{wt}$ Untr.

anti-Tgfbr1

anti-Actin

E $Cdx2Cre^{ERT};Rosa26R$

F

Tgfbr1 normalized expression

***   **   NS   NS

0.008
0.006
0.004
0.002

$Tgfbr1^{-/-}$
$Cdx2Cre;Tgfbr1^{fl/-}$
$Cdx2Cre;Tgfbr1^{fl/+}$
$Tgfbr1^{fl/-}$
$Tgfbr1^{fl/+}$

G

Tgfbr1 normalized expression

NS   ***   *

4e-04
3e-04
2e-04
1e-04

$Tgfbr1^{-/-}$
$Cdx2Cre;Tgfbr1^{3ex3fl/-}$
$Tgfbr1^{3ex3fl/+}$
Wild type

1258
1259
Supplementary figure 1. Generation on the $Tgfbr1$-cKO. (A) Scheme showing $Tgfbr1^{flox}$
and $Tgfbr1^{3ex3-flox}$ alleles. Highlighted is the region containing exons 2, 3 and 4 (colored
boxes) and the LoxP sites (black triangles). Also represented is the generation of a mRNA

with a triplicated exon 3 after splicing of the transcript from the $Tgfbr1^{3ex3\text{-}flox}$ allele. (B, B')

Skeletal staining of a E17.5 wild type (B) and $Tgfbr1^{3ex3\text{-}flox/\text{-}}$ fetus (B'), showing the presence

of additional thoracic (T) and lumbar (L) vertebrae, resembling a $Gdf11$ mutant phenotype.

(C) RT-PCR fragment spanning the whole $Tgfbr1$ coding region using RNA isolated from

wild type (1,5 kb) and $Tgfbr1^{3ex3\text{-}flox/3ex3\text{-}flox}$ (2 kb) embryos. (D). Western blot analysis of

protein extracts from HEK293T cells transfected with mammalian expression vectors

carrying either the $Tgfbr1^{3ex3}$ or the $Tgfbr1$ coding regions, showing that the $Tgfbr1^{3ex3}$

transcript produces a larger protein than its wild type counterpart. (E) Activation of β-

galactosidase expression in a tamoxifen-treated E10.5 $Cdx2Cre^{ERT};Rosa26R$ embryo

following the scheme used in this study. (F) RT-qPCR showing recombination efficiency in

$Tgfbr1^{flox/\text{-}};Cdx2Cre^{ERT+/0}$ embryos. Normalized $Tgfbr1$ expression level in $Tgfbr1^{flox/\text{-}}$

$::Cdx2Cre^{ERT+/0}$ is not different from $Tgfbr1^{\text{-}/\text{-}}$. Level of normalized expression in

$Tgfbr1^{+/flox};Cdx2Cre^{ERT+/0}$ equals that of $Tgfbr1^{flox/\text{-}}$ and half of $Tgfbr1^{flox/+}$ control, consistent

with it carrying only one functional allele. (G) RT-qPCR showing recombination efficiency

in $Tgfbr1^{3ex3\text{-}flox/\text{-}};Cdx2Cre^{ERT+/0}$ embryos. Normalized $Tgfbr1$ expression level in $Tgfbr1^{3ex3\text{-}}$

$^{flox/\text{-}};Cdx2Cre^{ERT+/0}$ embryos is significantly lower than in control embryos, although not

reaching $Tgfbr1^{\text{-}/\text{-}}$ level, consistent with lower amount of tamoxifen given compared to the

$Tgfbr1^{flox}$ line. NS: nonsignificant, *p=0.018, **p<0.01, ***p<0.001.

1281

1282

1283

Supplementary Figure 2. Skeletal phenotype of the *Tgfbr1-cKO* extra hindlimb. (A) Whole
mount image of the OPT acquired E16.5 fetus. (B, B') Whole mount (B) and 3D
reconstruction of the skeleton (B') of the right extra hindlimb. The autopod contained 5
digits, most of them with two phalanges, except for digit V, which only generated one
phalanx and digit II, which contained two adjacent distal phalanges (white arrow in B and
B'). (C, C') Whole mount (C) and 3D reconstruction of the skeleton (C') of the left extra
hindlimb. The autopod contained 6 metatarsal bones. Anterior digits developed two
phalanges, except for digit II, which generated two adjacent distal phalanges (white arrow
in C and C'); the posterior digits developed one phalanx, except for one digit that did not
develop any distal phalanges (red arrowhead in C').

1294

1295
Supplementary figure 3. Urogenital malformations in the *Tgfbr1-cKO.* (A-a') *Shh*
expression in E11.5 control (A, a) and *Tgfbr1-cKO* (A', a') embryos. (a, a') show transversal
sections through the cloacal region at the level indicated by the dashed line in A and A'. (B,
B') Sagittal sections through E10.5 control (B) and *Tgfbr1-cKO* (B') embryos stained with a
Shh probe. (C, C') Sagittal histological sections through bladder region of E16.5 control (C)
and *Tgfbr1-cKO* (C') fetuses. Black arrows in a, a', B, B' indicate the cloaca; arrowheads in
a', B', C' indicate intra-cloacal protrusions; bl: bladder; GT: genital tubercle; k: kidney; u:
ureter; ur: urethra. (D) ATAC-Seq profiles through the region containing the enhancer
regulating endodermal *Shh* expression (blue shadow). Bottom track shows conservation in
placental mammals. cHL: control hindlimb; mExHL: mutant extra hindlimb; cGT: control GT;
mGT: mutant GT.
1307

1308
Supplementary Figure 4. (A) The putative *Tbx5* GT enhancer shown in Fig. 4I' does not
drive β-galactosidase expression in the forelimb (black arrowhead, forelimb outlined with
the dashed line). (B) β-galactosidase expression driven by the *Grem1* enhancer in Fig. 5D
using a transgenic reporter assay [n=2/8] (limbs outlined with the dashed line). (C-E)
Activation of the β-galactosidase reporter by the *CR-Isl1* (C), *CR-Isl1$^{\Delta Lef1}$* (D), and *CR-*
Isl1$^{\Delta\Delta Lef1}$ (E) region in E12.0 embryos. Of note, the embryo shown in D is the one with the
strongest reporter expression in the GT region of all *Isl1$^{\Delta Lef1}$* transgenic embryos analyzed.

Top panels show schematic representation of DNA region: blue rectangle: 5' Lef1 binding motif; red rectangle: Gata3 binding motif; green rectangle: 3' Lef1 binding motif within the TF footprint observed in the *Isl1*-related Pattern 1 element. Black arrowheads show expression in otic vesicle; white arrow shows expression in posterior limb bud; black arrow indicates expression in GT. CR: conserved region.

Supplementary movie 1. OPT image from a E16.5 *Tgfbr1-cKO* fetus highlighting the limbs. The extra hindlimbs are shown in magenta.

Supplementary movie 2. OPT image from a E16.5 wild type fetus highlighting the caudal end of the intestine (red), the kidneys (cold green), the ovaries and uterus (bright green), the vagina (yellow), the urinary bladder (dark blue) and the urethra (light blue).

Supplementary movie 3. OPT image from a E16.5 *Tgfbr1-cKO* fetus highlighting the caudal end of the intestine (red), the urinary bladder (dark blue) and the urethra (light blue).

Supplementary Table 1. Transcription factor footprint analysis of pattern 1 elements. Column A: number of pattern 1 regions with at least 1 occurrence of the specified transcription factor; column B: number of pattern1 regions with no occurrence of the specified transcription factor; C: number of background regions with at least 1 occurrence of the specified transcription factor; D: number of background regions with no occurrence of the specified transcription factor.

Supplementary table 2. Oligonucleotides used in this work.

Genotyping primers		
Tgfbr1-flox	Forward	TTGAGCTTGCTGTCTGACTGGATAG
	Reverse	CTGTGGTTGGCAGGCATGTG
Cre	Forward	CGAGTGATGAGGTTCGCAAG
	Reverse	CCTGATCCTGGCAATTTCGGCT
Tgfbr1-null	Forward	CTACTGTGTTTCAAATGGGAGGGC
	Reverse	GGCCTGTCGGATCCTATCATC
β-galactosidase	Forward	AGCAGTTTTTCCAGTTCCGTTTATC
	Reverse	AGCGGCGTCAGCAGTTGTTTTTTAT
Cloning of regulatory elements		
CR-Isl1	Forward	GACTCGAGTCTGTGATACAAAACAATATATC

		Reverse	GAGGATCCAATCTTTCAAAGACATGGAGGG
CR-Isl1 - Left1 sites deletions		Primer1	GCGAATTCGGCTGAATGGAGGTCATTGGA
		Primer2	GCGAATTCTTCTGGCTTCAGTTCCTGAGAC
		Primer3	GCGAATTCCAGGTATGAAAAGCTGTGC
CR-Tbx5		Forward	GACTCGAGCTTTATGTATCTGAGCACACTG
		Reverse	GAAGATCTTCCTTCAACAAACCAT CCAC
CR-Wnt5a		Forward	CTGCTCGAGACTAGTACCTGCTGCTTATCAACCTAGTG
		Reverse	CAGCCATGGCTCCTACCTCACTGACTATCCAG
CR-Fgf10		Forward	CAGCTCGAGACTAGTAATCTGAGGTGGTTGCTCACTC
		Reverse	CATCCATGGCTATGGAAGGCTTATGTATCAC
Grem1-Enh		Forward	GACGTCGACACTAGTGTCATGTGTTCCTGTGATTCTG
		Reverse	GAGCCTGTATAAGAAGTTCAGGGCCA TGGCAG
Other primers			
qPCR-Tgfbr1		Forward	ACCGTGTGCCAAATGAAGAGG
		Reverse	CATCTAGATCTTGTAACACAATGGTCCTGGC
qPCR-Actin		Forward	ATGAAGATCCTGACCGAGCG
		Reverse	TACTTGCGCTCAGGAGGAGC
Tgfbr1-CDS		Forward	GAGAATTCCGGGCCACAAACAGTGGC
		Reverse	GAGTCGACCATTTTGATGCCTTCCTGT TGGC
Oligonucleotides for generating in situ probes			
Lmx1b		Forward	CTGCTGTGCAAGGGTGACTATGAG
		Reverse	GAGGCAAAGTAGGAGCTCTGCATG
Fgf10		Forward	TTCCTAGAAGTTATGGATGTTG
		Reverse	GTACTGCATCCACCAACAGTG
Grem1		Forward	AATGAATCGCACCGCATACAC
		Reverse	GACTAATACGACTCACTATAGGGAAGCAACTGCTGGTTCTTCTG
In situ probes			
En1	Davis and Joyner, 1988, ref [94]		
Fgf10	Whole coding region cloned into TOPO. Oligonucleotides listed above		
Fgf8	Crossley and Martin, 1995, ref [95]		
Grem1	PCR fragment with the whole coding region and T7 promoter at the 3' end. Oligonucleotides listed above		
Hand2	Srivastava et al, 1997, ref [96]		
Isl1	Jurberg et al 2013, ref [2]		
Lin28a	Aires et al., 2019, ref [97]		
Lmx1b	PCR fragment encompassing nucleotides 504 to 1202 of the mRNA cloned into pKS bluescript. Oligonucleotides listed above.		
Pitx1	Szeto et al, 1999, ref [98]		
Shh	Echelard et al, 1993, ref [99]		
Tbx5	Chapman et al, 1996, ref [100]		
Wnt5a	Yamaguchi et al, 1999, ref [32]		

1344

1345   **Supplementary table 3.** Coordinates of the genomic elements used in the

1346   reporter assays (from the GRCm38/mm10 reference genome).

1347

Element	Coordinates
CR-Isl1	chr13:116,284,356-116,285,685
CR-Tbx5	chr5:119,701,323-119,702,815
CR-Wnt5a	chr14:28,412,522-28,413,803
CR-Fgf10	chr13:118,745,999-118,747,518
Grem1 enhancer	chr2: 113,580,982-113,582,008

1348