1

Cell-penetrating peptides stimulate protein transport on the Twin-arginine translocation pathway

7
8
9

Robert McNeilage, Iniyan Ganesan, Johnathan Keilman, and Steven M. Theg*

Department of Plant Biology, University of California – Davis, Davis, California 95616 USA

14
15
16
17

*Corresponding author, smtheg@ucdavis.edu

19
20

22
23

Keywords: Protein transport, Twin arginine translocation pathway, Thylakoids, Inverted membrane vesicles, Toroidal pores,

Abstract

The Tat pathway is essential for photosynthetic protein transport across plant thylakoid membranes and is also ubiquitous throughout prokaryotes and archaea. The Tat pathway is unique amongst protein translocation pathways as it specializes in transporting folded proteins driven by a proton motive force. Mechanistic details of the actual translocation step (s) of the pathway remain elusive. Here, we show that membrane thinning stimulates Tat transport and, conversely, membrane strengthening abolishes Tat transport. We draw parallels from the Tat transport mechanism to that of cell penetrating peptides and propose that the Tat pore could be toroidal in shape and lined by lipids, as in those formed by cell penetrating peptides.

Significance Statement

Protein translocation across membranes is a significant cellular activity in both prokaryotes and eukaryotes. The Tat pathway for protein translocation operates in bacteria, archaea, chloroplasts, and plant mitochondria. Its mechanism of action has been difficult to decipher, but recent evidence suggests it does not use a conical proteinaceous transport channel. Instead, it has been suggested to translocate proteins through lipid-lined toroidal pores set up by membrane thinning. This work supports that hypothesis by showing that membrane-thinning cell-penetrating peptides stimulate the Tat pathway in both chloroplasts and bacterial plasma membranes, and that membrane stabilization blocks the pathway. We believe this is the most direct evidence to date of the toroidal pore mechanism operating in the Tat pathway.

Introduction

The Twin-Arginine Translocation (Tat) pathway transports folded proteins across the thylakoid membrane of chloroplasts, the inner membrane in plant mitochondria, and the periplasmic membranes of bacteria and archaea utilizing a proton motive force (pmf) as its sole energy source (Hamsanathan & Musser, 2018, New et al., 2018, Palmer & Stansfeld, 2020). Because both photosynthetic proteins and bacterial virulence factors(De Buck et al., 2008) pass through the Tat pathway, it is an important process to understand from multiple human health perspectives, both in nutrition and disease. In most instances, Tat transport is mediated by three proteins: Tha4, Hcf106, and cpTatC in thylakoids; TatA, TatB, and TatC in bacteria. Tha4 and Hcf106 are structurally similar, being composed of an N-terminal transmembrane helix (TMH), followed by a hinge region, an amphipathic helix (APH) that lies along the surface of the membrane, and an unstructured C-terminal domain. The Hcf106 APH and C-terminal domains are longer than their Tha4 counterparts(Porcelli et al., 2002). Interestingly, the NMR structures of TatA and TatB show that the TMHs are shorter than the hydrophobic portion of the lipid bilayer(Rodriguez et al., 2013a, Zhang et al., 2014), which creates a hydrophobic mismatch. TatC has six transmembrane domains with a topology where both termini face the stroma(Gouffi et al., 2002, Rollauer et al., 2012). Two of the TatC TMHs are also short, again leading to hydrophobic mismatch(Ramasamy et al., 2013, Rollauer et al., 2012). This hydrophobic mismatch has recently been shown to be important for the efficient operation of the Tat pathway(Hao et al., 2022a, Mehner-Breitfeld et al., 2022, Stockwald et al., 2022). Translocation of proteins along this pathway requires the precursor protein to possess an N-terminal signal peptide that enables the precursor to be recognized and bound by the receptor complex. The signal peptide contains the essential and eponymous RR (twin arginine) motif(Stanley et al.,

2000). Oligomers of Hcf106 and TatC in a 1:1 ratio compose the receptor complex for binding

of the precursor signal sequence(Bolhuis et al., 2001) in an energy independent manner(Mori &

Cline, 2002). After precursor binding and in the presence of a pmf, the receptor complex recruits

multimers of Tha4 and translocation occurs(Mori & Cline, 2002). After the translocation event,

the active translocon dissociates into the Hcf106-cpTatC receptor complex and oligomers of

Tha4(Mori & Cline, 2002). Sometime during or after the translocation event, the signal peptide

is cleaved. The detailed mechanism of the actual translocation event following Tha4 recruitment

is not yet understood.

Currently, there are three competing models proposed to accomplish protein transport on

the Tat pathway: the protein iris model in which Tha4 joins the translocon in varying

stoichiometry to form a tightly sealing pore around the incoming protein, the co-enzyme model,

in which TatA is thought to act as a co-enzyme that accumulates at the translocation site to

activate a translocon built from a complex consisting of TatB and TatC (Hauer et al., 2017,

Hauer et al., 2013), and the membrane defect model, in which the membrane is proposed to

form toroidal pores in response to bilayer destabilizing forces and through which proteins

traverse the membranes. A recent variation of the latter model posits that the transmembrane

helix (TMH) of TatA flips to lie along the trans-side of the membrane surface, contributing to the

bilayer instability (Stockwald et al., 2022).

The protein iris model faces several structural challenges: the pore would be either lined

with the hydrophobic TMHs, or the APHs of Tha4 would have to fold back in a hairpin onto the

TMDs. The hairpin fold is not favored by experimental evidence(Alcock et al., 2017, Aldridge et

al., 2012, Koch et al., 2012). Despite the Tat pathway translocating precursors of different

shapes and sizes, the membranes of the thylakoids remain tightly sealed during translocation and

do not allow for leakage of ions, which seems unlikely with the proteinaceous pores just

described(Asher & Theg, 2021, Teter & Theg, 1998). These findings, coupled with the transient

assembly of the active translocon, have led to the suggestion that the Tat pathway does not

utilize a proteinaceous pore to achieve translocation(Asher & Theg, 2021, Bruser & Sanders,

2003, Hao et al., 2022a). It has been proposed that instead, under the influence of the Tat

machinery subunits, the transported substrate, and membrane energization the membrane is

locally thinned to an extent that defects in the bilayer develop that allow a precursor to pass

through the membrane itself via lipid lined toroidal pores (reviewed in(Patel et al., 2014)). A

key component in this model is the thylakoid membrane thinning that was observed almost 50

years ago to occur in the light (Murakami & Packer, 1970a), and which has recently been

repeated in two different laboratories (Johnson et al., 2011a, Johnson et al., 2011b, Kirchhoff et

al., 2011, Li et al., 2020). We believe this provides a compelling mechanism for coupling

between the pmf and cpTat transport as the former causes membrane thinning and the latter

depends on it.

Cell Penetrating Peptides (CPPs) are short amphipathic peptides that interact with and

can induce toroidal pore formation in membranes (Huang & Li, 2023, Leveritt Iii et al., 2015).

Many CPPs adopt an α-helical configuration when interacting with membranes. The α-helices

are usually amphipathic, with positively charged residues, often arginine, on one side of the helix

and hydrophobic residues on the other side. The hydrophobic residues interface with the fatty

acid side chains, while the positively charged residues interact with the polar or negatively

charged lipid head groups of membranes. CPPs have generated considerable interest for their

ability to spontaneously cross membranes by causing membrane thinning and the formation of

toroidal pores(Herce & Garcia, 2007, Huang & Li, 2023, Madani et al., 2011). The ability to

thin membranes to the point of pore formation by CPPs suggests a possible mechanism for Tat

protein translocation, where the potential membrane thinning effect of the pmf combined with

destabilizing effects of the Tat component subunits, i.e., hydrophobic mismatch, might lead to

local and transient toroidal pore formation.

In this study, we used CPPs to initiate the thinning of thylakoid and *E. coli*

membranes(Chen et al., 2003, Mecke et al., 2005). We explored the idea that the CPPs can

provide some of the energy normally generated by the pmf to the Tat system by pre-thinning the

membrane. We demonstrate that CPPs specifically stimulate Tat transport and that the opposite

effect of strengthening the membrane abolishes transport. We believe that these experiments

establish the central role of membrane thinning in the Tat mechanism.

Results

Cell Penetrating Peptides stimulate Tat protein translocation

The mechanism through which CPPs cross membranes is proposed to involve actively

thinning the membrane to the point that the bilayer breaks down with the formation of toroidal

pores(Chen et al., 2003, Herce & Garcia, 2007). If the Tat pathway operates under similar

principles, that is, through membrane thinning, then we reasoned that CPPs should stimulate the

transport of proteins on the Tat pathway by contributing to the membrane thinning process. To

test this hypothesis, different CPPs were added to thylakoid Tat pathway transport reactions. We

chose four peptides from unrelated sources with widely differing sequences and related only in

their ability to spontaneously penetrate membranes. Fig. 1 shows that each peptide tested

stimulated the transport of iOE17, a model Tat substrate in thylakoids(Alder & Theg, 2003).

Because CPPs are known to form pores in membranes(Herce & Garcia, 2007), we tested

the possibility of passive diffusion of precursors into the thylakoids through a CPP induced pore.

Additionally, CPPs are the subject of investigation for delivering drugs and proteins across

membranes(Ramsey & Flynn, 2015), therefore, we address the possibility of precursors

hitchhiking on CPPs simultaneously with the previous concern. To these ends, we asked

whether CPPs could facilitate transport either using a transport-incompetent precursor or in the

absence of an energy source. If CPPs promote spontaneous diffusion of precursors across

thylakoids membranes, then they should cause translocation in the absence of a pmf.

Thylakoids, CPPs, and iOE17 were incubated together in the dark at ambient temperature for

30min. CPPs did not allow for translocation without the light-generated pmf (Fig. 1B). In the

presence of a light source, translocation occurred and was stimulated by CPPs. We then asked if

the pmf and CPPs would be able to cause the translocation of a transport-incompetent Tat

substrate. We used a mutated iOE17 with RR to KK substitutions in the signal sequence

(iOE17KK), which renders the precursor transport-incompetent(Ize et al., 2002). iOE17KK was

not transported with or without CPPs present (Fig 1C). These results preclude the possibility that

either precursors or the thylakoid processing protease can either diffuse through pores created by

CPPs or be carried by them across the thylakoid membrane in a Tat-independent or energy-

independent manner.

In these two controls, the precursor was found with reisolated thylakoids. This indicates

that being present at the membrane is not sufficient for CPPs to facilitate passive transport of

precursors. These results also suggest that if precursors travel through any pore in the

membrane, it was induced and controlled by the Tat machinery, which would have been inactive

in these controls.

To determine the specificity of CPP stimulation of the Tat pathway, we monitored the

effect of CPPs on protein translocation across thylakoids via the Sec pathway, which occurs

through a proteinaceous pore (Li et al., 2016). Transport of the model thylakoid Sec substrate,

iOE33, was not significantly altered by HIVTat (Fig. 1D). This clearly establishes that CPPs do

not universally stimulate protein translocation.

Figure 1a shows that stimulation of the Tat pathway drops at higher concentrations of

melittin and penetratin. We reasoned that these CPPs caused enough ion leakage to overwhelm

the ability of the thylakoids to maintain the pmf, leading to the decline in transport stimulation.

To investigate this possibility, we monitored the effect of higher concentrations of melittin on

both the electrical and chemical components of the pmf. Melittin at 10 μM caused a faster

dissipation of the light-induced $\Delta\psi$ across the thylakoid membrane compared with controls (Fig.

1E). Similarly, melittin at 10 μM resulted in a reduced steady-state trans-membrane ΔpH (Fig.

1F). Both of these observations are consistent with CPPs mediating proton and other ion leakage

through pores induced at these higher concentrations. Taken together, these results indicate that

high concentrations of CPPs compromise the integrity of the thylakoid membranes, likely by

inducing transient pores(Herce & Garcia, 2007). This is not surprising because inducing pores in

and causing the rupturing of membranes is a well-documented effect of some CPPs(Herce &

Garcia, 2007, Yang et al.), leading them also to be described as antimicrobial peptides (Huang &

Li, 2023).

Effects of CPPs on the energetics of Tat protein transport

Next, we further evaluated the mechanism of CPP-mediated Tat transport stimulation.

We reasoned that the light- and pH-induced membrane thinning of thylakoids observed in the

literature(Kirchhoff et al., 2011, Li et al., 2020, Murakami & Packer, 1970b) could provide a

mechanism through which the pmf could be coupled to protein translocation on the Tat pathway

if the latter depended on membrane thinning. This would also provide an explanation of the

stimulatory effect of CPPs, since they are proposed to act via membrane thinning(Herce &

Garcia, 2007, Madani et al., 2011). Specifically, if both CPPs and the pmf contribute to Tat

protein transport through their effects on membrane thickness, then we would expect that CPPs

would lower the threshold energy needed to promote protein transport by providing some

thinning before application of the ΔpH. To test this idea, we measured translocation as a

function of ΔpH to elucidate the threshold ΔpH at which translocation begins(Rollauer et al.,

2012).

In chemiosmotic systems, plotting the work performed by the pmf as a function of the

driving force results in a biphasic plot. Initially, there is not enough energy in the gradient to

perform the work queried. At some point a threshold is reached, above which the output rises

linearly with the driving force. This thermodynamic threshold represents the minimum energy

required for the work queried, in this case protein transport (Alder & Theg, 2003). Figure 2

shows that the stimulatory effect of both melittin and penetratin is caused by their lowering of

the ΔpH threshold for Tat protein transport. This is consistent with the idea that the CPPs

predispose the membranes for protein transport by mediating part of the essential thinning that is

then finished by the pmf.

CPPs interact with the thylakoid membrane, causing a change in density

The ΔpH and Δψ measurements and the ΔpH threshold experiments suggest that CPPs

change the structure of the thylakoids. We suspected that the thylakoids were reorganized in

response to interaction with the CPPs. Previously it has been observed that the vesicle inducing

protein in plastid1 (VIPP1) stimulates Tat protein translocation and also causes a change in

thylakoid density (Lo & Theg, 2012a). We asked whether CPPs could lead to a similar change

in the density of the thylakoids. To this end, we performed thylakoid pelleting assays(Lo &

Theg, 2012b), observing how much chlorophyll remains in the supernatant after low-speed

centrifugation. As seen in Fig. 3, thylakoids generally pellet more readily upon incubation with

CPPs, and the response is dose dependent. This suggests that CPPs interact directly with the

thylakoid membrane, causing a reorganization of the thylakoid super-structure, which leads to

their increased density.

CPPs stimulate Tat transport in bacterial inverted membrane vesicles

It is widely held that the mechanism of protein transport on the Tat pathway is the same

in all Tat systems found in the different domains of life (Celedon & Cline, 2012a, Palmer &

Berks, 2012), and this view has gained further support from recent work in our lab (Zhou et al.,

2023). Accordingly, we would predict that the stimulation of the Tat pathway by CPPs

demonstrated above in chloroplasts would apply to the bacterial Tat pathway as well. To test

this, we examined the effect of CPPs on the transport of the model substrate pre-SufI-IAC into

inverted membrane vesicles (IMVs) prepared from JM109 cells overexpressing

TatABC(Bageshwar & Musser, 2007, Bageshwar et al., 2009). As in thylakoids, we again found

that CPPs stimulated Tat transport in the bacterial system (Fig. 4). This result suggests that CPPs

can act universally in stimulating Tat transport, implying a shared mechanism across systems.

The CPP stimulation effect is lost at higher concentrations of CPPs, likely due to pmf uncoupling

as in thylakoids (Fig. 1E,F).

The amphipathic domains of Hcf106 stimulates Tat protein translocation

One of the biophysical characteristics linking different CPPs is their abilities to form

amphipathic helices. All TatA- and TatB-family member proteins, including Tha4 and Hcf106,

also have amphipathic helices, essential for their function, and the APH of Hcf106 is particularly

enriched in arginines, as are many CPPs. Given the similarities of these APHs to CPPs we tested

their ability to stimulate translocation. While the Tha4 APH peptide did not increase transport of

iOE17 (Fig. 5A), the Hcf106 APH peptide did stimulate translocation at a level comparable to

CPPs (Fig. 5E). As with the CCPs, we noted a decline in transport stimulation at 20 µM Hcf106

APH peptide. However, unlike the situation with CCPs, we observed that the Hcf106 APH

peptide neither compromised membrane integrity nor increased the thylakoid membrane density

at this concentration (Fig. 5B-D). The reason for this decrease in iOE17 transport at 20 µM

Hcf106 APH peptide remains unknown; we did not probe this phenomenon further.

242   Interestingly, the mature substrate protein formed in the presence of the Hcf106 APH

peptide at 10 and 20 µM was not fully protected from digestion with thermolysin, suggesting that

some fraction had not completely crossed the thylakoid membrane (Fig. 5F). The surprising

appearance of a matured but protease-sensitive Tat substrate protein is not new in the literature

but is very unusual(Di Cola & Robinson, 2005, Frobel et al., 2012, Leheny et al., 1998). Fröbel

et al.(Frobel et al., 2012) proposed that in the bacterial system a precursor could be bound by

TatC in such a manner that allows the signal peptide to cross the membrane without full

translocation of the passenger protein. Whether we are observing the same phenomenon here

and what role the Hcf106 APH peptide plays in it, remain to be elucidated.

251   **Membrane thickening abolishes Tat transport**

252   Having established that CPPs and the Hcf106 APH peptide stimulate Tat protein

transport, presumably through membrane thinning and destabilization, we next determined the

effect of membrane thickening. To thicken, or in other words strengthen, the thylakoid

membrane, we employed trifluoroethanol (TFE). TFE inserts into the membrane at the

hydrophobic/hydrophilic interface and increases the lateral membrane pressure, which in effect

stabilizes the lipid bilayer(van den Brink-van der Laan et al., 2004, Yuan et al., 2017). We found

that Tat transport is very sensitive to TFE, with a precipitous drop in transport at 1% (v/v) TFE

(Fig. 6). The ΔpH was not greatly affected by TFE, ruling out the possibility of an indirect effect

of TFE on transport through a loss of the pmf. As before, to compare an effect on Tat transport

to that with a proteinaceous pore translocon, we examined the effect of TFE on Sec transport.

TFE did reduce Sec transport, but still supported Sec transport at concentrations 2.7-fold higher

than that required to abolish Tat transport (Fig. 6). Considering the complementary effects of

TFE and CPPs, along with reports on lengthening the TatA and TatB TMHs (Hao et al., 2022a,

Mehner-Breitfeld et al., 2022, Stockwald et al., 2022), it is apparent that membrane structure, in

particular, membrane thinning, is critical for the Tat translocation mechanism.

268 **Discussion**

In this work, we demonstrated that CPPs stimulate Tat transport (Fig. 1) by interacting

with and changing the thylakoid membrane (Fig. 3) and thereby lowering the ΔpH threshold for

translocation (Fig. 2).

Since CPPs are known to cause thinning in membranes(Chen et al., 2003, Herce &

Garcia, 2007, Madani et al., 2011, Mecke et al., 2005), it is likely that they also act to stimulate

the Tat pathway by thinning the thylakoid membrane. This mechanism also successfully predicts

the decrease in the threshold ΔpH for protein transport mediated by melittin and penetratin.

Given that the bacterial Tat system is also stimulated by CPPs, we can extrapolate that the

membrane thinning mechanism of action is conserved across systems. This concurs with recent

observations that *E. coli* TatA causes membrane defects in coordination with the substrate

protein(Hou et al., 2018).

Membrane thinning provides an interesting mechanism for translocation in the context of

coupling of the pmf to protein transport. Assuming a random distribution of CPPs in the

thylakoid membrane, the global thinning caused by the pmf(Kirchhoff et al., 2011) is likely

mimicked by the action of the CPPs. This idea is further supported by CPP stimulation of the

bacterial pathway because CPP interaction with model bacterial membranes is more widely

studied(Lee et al., 2005). We would expect that the thinning could reach an extreme locally as

the translocon subunits each contribute to narrowing the membrane through hydrophobic

mismatch in their TMD regions (Hao et al., 2022a, Mehner-Breitfeld et al., 2022, Stockwald et

al., 2022) and/or potential contributions from their APH regions. Should this local thinning draw

further Tha4 subunits into the vicinity, then the membrane destabilization calculated by

Rodrigiez et al.(Rodriguez et al., 2013b) that occurred when the TatA mulitmer exceeds nine

subunits would provide the conditions for the formation of a toroidal pore. However, one could

expect the Tat machinery to induce controlled pores that would not result in massive ion

leakage(Asher & Theg, 2021, Teter & Theg, 1998) by forming a sphincter, having a very short

life, or some combination of these two.

In the toroidal pore model, the head groups of lipids would be expected to interact with

the substrate. While lipid-substrate interactions have been demonstrated(Bageshwar et al., 2009,

Rodriguez et al., 2013b), cross links between the two have not been investigated and may be a

worthy subject of study to determine which lipids comprise the toroidal pore during transport,

since some lipids are essential to Tat transport (Rathmann et al., 2017). One would expect to

find the pore consisting of lipids that introduce curvature in the membrane. Interestingly,

branched and phospholipids of the thylakoid membrane tend to interact with photosynthetic

membrane proteins (Mizusawa & Wada, 2012). If this data can be extrapolated to more

membrane proteins, then TatA recruitment may serve as a means of collecting the lipids that

induce the curvature necessary for toroidal pore formation.

A further driver for membrane thinning could be provided by attraction of the charged

arginine residues in the Tat substrates' signal peptide to the trans-side of the membrane, as is

hypothesized for the mechanism of membrane crossing for arginine-rich CPPs (Herce & Garcia,

2007). However, this view is tempered by the recognition that the signal sequence arginines are

likely buried in the receptor complex and interacting with TatC (Rollauer et al., 2012, Zoufaly et

al., 2012). Thus, if they play a role, it may be upon release of the precursor from the receptor

complex. Substrate proteins poised at the pore site could then traverse the membrane while the

pore is open with the signal acting as an anchor. Such pores would be expected to close quickly

once the substrate is cleared as the membrane-thinning pmf would be locally decreased due to

the concurrent movement of protons through the pore. This would be a natural consequence of

the large number of protons observed to be released from the proton gradient per protein

translocated(Alder & Theg, 2003).

A toroidal pore itself does not provide directionality. Since transport occurs *in vitro*

without soluble factors, directionality would be provided by the translocon. Alternatively,

substrate exit from the translocon to the trans side of the membrane could be random. In the case

of a failed translocation attempt, rebinding of the substrate and reassembly of the active

translocon may be necessary. We recognize that a mechanism requiring multiple attempts might

explain, in part, the extraordinarily high energetic cost(Alder & Theg, 2003) and time(Celedon &

Cline, 2012b, Musser & Theg, 2000, Teter & Theg, 1998) of the translocation reaction, as only

successful events would be scored as protein transport..

Here, we have not directly addressed whether the Tat pore is proteinaceous or toroidal.

However, membrane thinning is fundamental to toroidal pore formation, which is generally not

the case for proteinaceous pores. An example of an exception would be proteinaceous pores

formed by mechanosensitive ion channels(Louhivuori et al., 2010). Therefore, it remains

possible that membrane thinning causes a conformational switch in TatA that induces assembly

of a proteinaceous pore. Again, we disfavor this model due to the problem of the substrate

contacting the hydrophobic TMHs of the Tat machinery. A solution to the substrate interaction

with TMHs has been proposed in the charged zipper model wherein the C-terminal domain of

the TatA family proteins are proposed to line a TatA pore(Walther et al., 2013). This model is

not supported by the evidence {Alcock, 2017 #26358;Rodriguez, 2013 #26297;Berks, 2015

#26291}.

It is interesting that the Hcf106 APH peptide caused Tat transport stimulation, while the

Tha4 APH peptide did not. This may be because Tha4 APH may require a higher concentration

to act on the membrane. Indeed, an actively transporting Tat translocase contains a higher

stoichiometry of Tha4 than Hcf106(Celedon & Cline, 2012b) and only one Hcf106 subunit

within the complex may be involved per substrate transported. Additionally, the full length Tha4

APH may be more active than the shorter peptide we used, despite having a reduced hydrophobic

moment. We also note that the APH of That4 has one arginine, whereas the APH of Hcf106 has

five. This residue is found in abundance in CPPs, which have in fact been referred to as

arginine-rich peptides (Futaki et al., 2013, Hao et al., 2022b); a titration showed that the optimal

number for maximum CPP activity is eight arginines (Futaki et al., 2001).

In summary, we have shown here that protein translocation on the Tat pathway is

stimulated by CPPs and CPP-like peptides (the Hcf106 APH) in both thylakoids and bacteria.

We interpret this as additional evidence that Tat protein transport involves membrane thinning

and ultimately lipid bilayer breakdown to form toroidal pores which allow protein substrates to

traverse the biological membranes. The transient and local nature of such pores has rendered

them undetectable to date, and this will be the challenge for our future studies of this enigmatic

pathway.

Materials and Methods

Reagents

Cell Penetrating Peptides were purchased from Anaspec. Tha4 APH

(KKLPEVGRSIGQTVKSFQQAAK) and Hcf106 APH

(KGLAEVARNLGKTLREFQPTIREIQDVSREFKSTLER) were purchased from Lifetien

at >90% purity. Peas were purchased form Harris Seed.

Buffers. Grinding buffer (GB): .05M (K)HEPES, .33M Sorbitol, 1mM $MgCl_2$, $MnCl_2$,

2mM Na_2EDTA, pH to 7.3 then .1%BSA. Import Buffer (IB): .05M K-Tricine, .33M Sorbitol,

3mM $MgCl_2$ pH 8. 2x Sample Buffer (2xSB): .125M Tris (pH 6.8), 4% SDS, 20%

glycerol, .005% Bromophenol Blue, 10% BME.

Methyl viologen (MV), valinomycin, N,N Naphthylethylene diamine (NED),

Phosphatidyl glycerol 18:1-06:0 2- (4-nitro-2,1,3-benzoxadiazol-7-yl) (PG-NBD), and nigericin

were stored in 100% EtOH.

Plant growth and thylakoid isolation.

Peas were grown on soil as described previously (Lo & Theg, 2011). Pea leaves and

shoots from 10 to 14-day old plants were blended in short pulses in GB. Blendate was

centrifuged for 5 min at 4250 rpm. The pellet was resuspended in GB and passed through a

Percoll gradient (50:50 percoll:2xGB). The lower band of intact chloroplast was then washed

twice with IB. Thylakoids were obtained by lysing chloroplasts in water supplemented with 5

mM MgCl$_2$. Lysis was stopped by adding an equal volume of 2xIB. Isolations were quantified

by chlorophyll content as described(Lo & Theg, 2011).

Thylakoid Tat Translocation assays.

Translation. In vitro translations were performed with wheat germ extract as directed by

the manufacturer (Promega or tRNA Probes) with ^{3}H leucine or ^{35}S methionine. Plasmid

encoding iOE17 was described previously(Alder & Theg, 2003).

Translocation. Transport reactions (60 µl) contained 2 µl translation product, CPPs at

indicated concentrations, and 0.33 mg/mL chlorophyll-equivalent thylakoids in 1xIB. Reactions

were initiated by addition of thylakoids, incubated 6 min at room temperature under illumination,

and stopped with 10 volumes of cold IB and transferred to the dark on ice. Thylakoids were

reisolated by pelleting at 3,000 rpm for 5 min. The pellet was resuspended in SB, boiled for 10

min, and analyzed by SDS-PAGE and fluorography.

Thermolysin digestion. After stopping a transport reaction, 200 µg/mL thermolysin and 5

mM CaCl$_2$ were added where noted. After a 30-minute incubation on ice, EDTA was added to a

final concentration of 25 mM. Reisolation of thylakoids and proceeding followed as above.

Thylakoid Sec Translocation assays.

A plasmid containing the thylakoid Sec substrate iOE33 was a gift from Ken Cline; the

protein was in vitro translated as described above. SecA was purified from plasmid pET28a-

cpSecA1 as described(Endow et al., 2015).

Thylakoids were pretreated with SecA for 15 min at room temperature. Transport

reactions (60 µl) containing 4 µl radiolabeled substrate, 5 mM ATP, 25 µg/mL SecA, and 0.33

mg/mL thylakoids were illuminated at room temperature for 30 min. Samples were subsequently

treated and analyzed as described for Tat transport.

Measurement of Δψ. Measurements of the Δψ-reporting carotenoid electrochromic shift

(ECS) were performed on a JTS-10 spectrophotometer. MV (20 µM), thylakoids (20 µg Chl), the

given concentration of melittin were brought to a total volume of 1mL in IB. The rate of decay

of the ECS signal was monitored after delivering two 9 ms flashes 100 ms apart.

Measurement of the thermodynamic threshold for Tat protein transport. ΔpH

measurements were made on a Horiba Flouromax fluorometer. 20 mM MV, 4 mM NED, 10 µL

of radiolabeled precursor, 120 µg thylakoids, and the given concentration of CPPs were made up

to a 2 mL total volume with IB in a 2 mL microcentrifuge tube. After 10 s of dark the excitation

beam was applied. After 1 minute from experiment start, an actinic light of varying intensity,

depending on desired ΔpH, was applied for 6 minutes. The transport reaction was stopped with

1 mM final concentrations of nigericin and valinomycin. Thylakoids were immediately

reisolated, resuspended in SB and analyzed by SDS-PAGE and fluorography.

Measurement of relative thylakoid density. As in Lo and Theg(Lo & Theg, 2012b), 60

µL of 0.33 mg/mL chlorophyll-equivalent thylakoids were incubated with CPPs in the dark at

ambient temperatures for 10 min and then centrifuged at 100 rpm for 1 min. 20 µL of

supernatant was removed and diluted into 180 µL 89% acetone and the concentration of

chlorophyll was calculated from absorbance as previously described(Lo & Theg, 2012b).

E. coli inverted membrane vesicle (IMV) transport

The pre-SufI-IAC protein was overexpressed in *E. coli* from plasmid pSufI-

IAC(Bageshwar et al., 2009), which encodes SufI precursor with substituted native cysteines and

a single C-terminal cysteine. Protein was purified under native conditions as previously

described for pre-SufI(Bageshwar & Musser, 2007). Eluted protein (6.5 µM) was biotinylated by

addition of 0.7 mM TCEP and 500 µM biotin maleimide and incubation on ice for 30 min. The

labeling reaction was quenched with 2 mM DTT.

IMV isolation from JM109 overexpressing pTatABC and transport assays were

essentially as previously described(Bageshwar & Musser, 2007). Briefly, 110 nM biotinylated

pre-SufI-IAC was mixed with IMVs (A_{280}=5), 4 mM NADH, and indicated concentrations of

CPPs. Reactions were incubated at 37˚C for 8 min and stopped on ice. Samples were

subsequently treated with 0.3 mg/mL Proteinase K for 40 min at RT, quenched with 3 mM

PMSF on ice, and diluted with an equal volume of 2xSB. Samples were analyzed by SDS-PAGE

and α-biotin Western blotting as described previously(Ganesan et al., 2018).

Acknowledgements

429   We thank Siegfried Musser and Shruthi Hamsanathan for pSufI-IAC, Tracy Palmer and Jon

430   Cherry for JM109 expressing pTatABC, and Ken Cline for the iOE33 clone. We gratefully

431   acknowledge support from the Division of Chemical Sciences, Geosciences, and Biosciences,

432   Office of Basic Energy Sciences of the U.S. Department of Energy through Grants DE-FG02-

433   03ER15405 and DE-SC0020304 to SMT.

Author Contributions

Robert McNeilage: Conceptualization; performed experiments in Figures 1, 2, 3 and 5; writing

– original draft, review and editing. **Iniyan Ganesan:** Conceptualization; performed

experiments in Figure 3 and 6; writing – original draft, review and editing. Johnathan Keilman:

Performed experiment in Figure 6; writing – review and editing. Steven M. Theg:

Conceptualization; supervision; funding acquisition; project administration; writing – original

draft, review and editing.

Disclosure and competing interests statement

444   The authors declare that they have no conflict of interest.

Data Availability

447   This study includes no data deposited in external repositories.

References

Alcock F, Damen MPM, Levring J, Berks BC (2017) In vivo experiments do not support the charge zipper model for Tat translocase assembly. eLife 6: e30127

Alder NN, Theg SM (2003) Energetics of protein transport across biological membranes. a study of the thylakoid DeltapH-dependent/cpTat pathway. Cell 112: 231-42

Aldridge C, Storm A, Cline K, Dabney-Smith C (2012) The chloroplast twin arginine transport (Tat) component, Tha4, undergoes conformational changes leading to Tat protein transport. J Biol Chem 287: 34752-63

Asher AH, Theg SM (2021) Electrochromic shift supports the membrane destabilization model of Tat-mediated transport and shows ion leakage during Sec transport. Proc Natl Acad Sci U S A 118

Bageshwar UK, Musser SM (2007) Two electrical potential-dependent steps are required for transport by the Escherichia coli Tat machinery. J Cell Biol 179: 87-99

Bageshwar UK, Whitaker N, Liang FC, Musser SM (2009) Interconvertibility of lipid- and translocon-bound forms of the bacterial Tat precursor pre-SufI. Mol Microbiol 74: 209-26

Bolhuis A, Mathers JE, Thomas JD, Barrett CM, Robinson C (2001) TatB and TatC Form a Functional and Structural Unit of the Twin-arginine Translocase from Escherichia coli. J Biol Chem 276: 20213-20219

Bruser T, Sanders C (2003) An alternative model of the twin arginine translocation system. Microbiol Res 158: 7-17

Celedon JM, Cline K (2012a) Intra-plastid protein trafficking; how plant cells adapted prokaryotic mechanisms to the eukaryotic condition. Biochim Biophys Acta

Celedon JM, Cline K (2012b) Stoichiometry for binding and transport by the twin arginine translocation system. J Cell Biol 197: 523-34

Chen FY, Lee MT, Huang HW (2003) Evidence for membrane thinning effect as the mechanism for peptide-induced pore formation. Biophys J 84: 3751-8

De Buck E, Lammertyn E, Anne J (2008) The importance of the twin-arginine translocation pathway for bacterial virulence. Trends Microbiol 16: 442-53

Di Cola A, Robinson C (2005) Large-scale translocation reversal within the thylakoid Tat system in vivo. J Cell Biol 171: 281-9

Endow JK, Singhal R, Fernandez DE, Inoue K (2015) Chaperone-assisted Post-translational Transport of Plastidic Type I Signal Peptidase 1. J Biol Chem 290: 28778-91

Frobel J, Rose P, Lausberg F, Blummel AS, Freudl R, Muller M (2012) Transmembrane insertion of twin-arginine signal peptides is driven by TatC and regulated by TatB. Nat Commun 3: 1311

Futaki S, Hirose H, Nakase I (2013) Arginine-rich peptides: methods of translocation through biological membranes. Curr Pharm Des 19: 2863-8

Futaki S, Suzuki T, Ohashi W, Yagami T, Tanaka S, Ueda K, Sugiura Y (2001) Arginine-rich peptides. An abundant source of membrane-permeable peptides having potential as carriers for intracellular protein delivery. J Biol Chem 276: 5836-40

Ganesan I, Shi LX, Labs M, Theg SM (2018) Evaluating the Functional Pore Size of Chloroplast TOC and TIC Protein Translocons: Import of Folded Proteins. Plant Cell 30: 2161-2173

Gouffi K, Santini CL, Wu LF (2002) Topology determination and functional analysis of the Escherichia coli TatC protein. FEBS Lett 525: 65-70

Hamsanathan S, Musser SM (2018) The Tat protein transport system: intriguing questions and

conundrums. FEMS Microbiol Lett 365
Hao B, Zhou W, Theg SM (2022a) Hydrophobic mismatch is a key factor in protein transport
across lipid bilayer membranes via the Tat pathway. J Biol Chem: 101991
Hao M, Zhang L, Chen P (2022b) Membrane Internalization Mechanisms and Design Strategies
of Arginine-Rich Cell-Penetrating Peptides. Int J Mol Sci 23
Hauer RS, Freudl R, Dittmar J, Jakob M, Klosgen RB (2017) How to achieve Tat transport with
alien TatA. Sci Rep 7: 8808
Hauer RS, Schlesier R, Heilmann K, Dittmar J, Jakob M, Klösgen RB (2013) Enough is enough:
TatA demand during Tat-dependent protein transport. Biochim Biophys Acta 1833: 957-65
Herce HD, Garcia AE (2007) Cell penetrating peptides: how do they do it? J Biol Phys 33: 345-
56
Hou B, Heidrich ES, Mehner-Breitfeld D, Bruser T (2018) The TatA component of the twin-
arginine translocation system locally weakens the cytoplasmic membrane of Escherichia coli
upon protein substrate binding. The Journal of biological chemistry
Huang X, Li G (2023) Antimicrobial Peptides and Cell-Penetrating Peptides: Non-Antibiotic
Membrane-Targeting Strategies Against Bacterial Infections. Infect Drug Resist 16: 1203-1219
Ize B, Gerard F, Zhang M, Chanal A, Voulhoux R, Palmer T, Filloux A, Wu LF (2002) In vivo
dissection of the tat translocation pathway in Escherichia coli. J Mol Biol 317: 327-335
Johnson MP, Brain AP, Ruban AV (2011a) Changes in thylakoid membrane thickness associated
with the reorganization of photosystem II light harvesting complexes during photoprotective
energy dissipation. Plant Signal Behav 6: 1386-90
Johnson MP, Goral TK, Duffy CD, Brain AP, Mullineaux CW, Ruban AV (2011b)
Photoprotective energy dissipation involves the reorganization of photosystem II light-harvesting
complexes in the grana membranes of spinach chloroplasts. Plant Cell 23: 1468-79
Kirchhoff H, Hall C, Wood M, Herbstova M, Tsabari O, Nevo R, Charuvi D, Shimoni E, Reich
Z (2011) Dynamic control of protein diffusion within the granal thylakoid lumen. Proc Natl
Acad Sci U S A 108: 20248-53
Koch S, Fritsch MJ, Buchanan G, Palmer T (2012) Escherichia coli TatA and TatB proteins have
N-out, C-in topology in intact cells. J Biol Chem 287: 14420-31
Lee M-T, Hung W-C, Chen F-Y, Huang HW (2005) Many-Body Effect of Antimicrobial
Peptides: On the Correlation Between Lipid's Spontaneous Curvature and Pore Formation.
Biophys J 89: 4006-4016
Leheny EA, Teter SA, Theg SM (1998) Identification of a role for an azide-sensitive factor in the
thylakoid transport of the 17-kilodalton subunit of the photosynthetic oxygen-eolving cmplex.
Plant Physiol 116: 805-14
Leveritt Iii John M, Pino-Angeles A, Lazaridis T (2015) The Structure of a Melittin-Stabilized
Pore. Biophys J 108: 2424-2426
Li L, Park E, Ling J, Ingram J, Ploegh H, Rapoport TA (2016) Crystal structure of a substrate-
engaged SecY protein-translocation channel. Nature 531: 395-399
Li M, Mukhopadhyay R, Svoboda V, Oung HMO, Mullendore DL, Kirchhoff H (2020)
Measuring the dynamic response of the thylakoid architecture in plant leaves by electron
microscopy. Plant Direct 4: e00280
Lo SM, Theg SM (2011) Protein targeting across and into chloroplast membranes. Methods Mol
Biol 684: 139-57
Lo SM, Theg SM (2012a) Role of vesicle-inducing protein in plastids 1 in cpTat transport at the
thylakoid. Plant J 71: 656-68

Lo SM, Theg SM (2012b) Role of Vesicle-Inducing Protein in Plastids 1 in cpTat transport at the thylakoid. Plant J

Louhivuori M, Risselada HJ, van der Giessen E, Marrink SJ (2010) Release of content through mechano-sensitive gates in pressurized liposomes. Proceedings of the National Academy of Sciences of the United States of America 107: 19856-60

Madani F, Lindberg S, Langel, #220, lo, Futaki S, Gr, #228, slund A (2011) Mechanisms of Cellular Uptake of Cell-Penetrating Peptides. Journal of biophysics 2011

Mecke A, Lee DK, Ramamoorthy A, Orr BG, Banaszak Holl MM (2005) Membrane thinning due to antimicrobial peptide binding: an atomic force microscopy study of MSI-78 in lipid bilayers. Biophys J 89: 4043-50

Mehner-Breitfeld D, Ringel MT, Tichy DA, Endter LJ, Stroh KS, Lünsdorf H, Risselada HJ, Brüser T (2022) TatA and TatB generate a hydrophobic mismatch important for the function and assembly of the Tat translocon in Escherichia coli. J Biol Chem 298

Mizusawa N, Wada H (2012) The role of lipids in photosystem II. Biochim Biophys Acta 1817: 194-208

Mori H, Cline K (2002) A twin arginine signal peptide and the pH gradient trigger reversible assembly of the thylakoid [Delta]pH/Tat translocase. J Cell Biol 157: 205-10

Murakami S, Packer L (1970a) Light-induced changes in the conformation and configuration of the thylakoid membrane of Ulva and Porphyra chloroplasts in vivo. Plant Physiol 45: 289-99

Murakami S, Packer L (1970b) Protonation and chloroplast membrane structure. J Cell Biol 47: 332-51

Musser SM, Theg SM (2000) Characterization of the early steps of OE17 precursor transport by the thylakoid DeltapH/Tat machinery. Eur J Biochem 267: 2588-2598

New CP, Ma Q, Dabney-Smith C (2018) Routing of thylakoid lumen proteins by the chloroplast twin arginine transport pathway. Photosynth Res 138: 289-301

Palmer T, Berks BC (2012) The twin-arginine translocation (Tat) protein export pathway. Nat Rev Microbiol 10: 483-96

Palmer T, Stansfeld PJ (2020) Targeting of proteins to the twin-arginine translocation pathway. Mol Microbiol 113: 861-871

Patel R, Smith SM, Robinson C (2014) Protein transport by the bacterial Tat pathway. Biochim Biophys Acta

Porcelli I, de Leeuw E, Wallis R, van den Brink-van der Laan E, de Kruijff B, Wallace BA, Palmer T, Berks BC (2002) Characterization and membrane assembly of the TatA component of the Escherichia coli twin-arginine protein transport system. Biochemistry 41: 13690-13697

Ramasamy S, Abrol R, Suloway CJ, Clemons WM, Jr. (2013) The glove-like structure of the conserved membrane protein TatC provides insight into signal sequence recognition in twin-arginine translocation. Structure 21: 777-88

Ramsey JD, Flynn NH (2015) Cell-penetrating peptides transport therapeutics into cells. Pharmacol Ther 154: 78-86

Rathmann C, Schlosser AS, Schiller J, Bogdanov M, Bruser T (2017) Tat transport in Escherichia coli requires zwitterionic phosphatidylethanolamine but no specific negatively charged phospholipid. FEBS Lett 591: 2848-2858

Rodriguez F, Rouse SL, Tait CE, Harmer J, De Riso A, Timmel CR, Sansom MS, Berks BC, Schnell JR (2013a) Structural model for the protein-translocating element of the twin-arginine transport system. Proc Natl Acad Sci U S A 110: E1092-101

Rodriguez F, Rouse SL, Tait CE, Harmer J, De Riso A, Timmel CR, Sansom MSP, Berks BC,

Schnell JR (2013b) Structural model for the protein-translocating element of the twin-arginine
transport system. Proceedings of the National Academy of Sciences 110: E1092-E1101
Rollauer SE, Tarry MJ, Graham JE, Jaaskelainen M, Jager F, Johnson S, Krehenbrink M, Liu
SM, Lukey MJ, Marcoux J, McDowell MA, Rodriguez F, Roversi P, Stansfeld PJ, Robinson CV,
Sansom MS, Palmer T, Hogbom M, Berks BC, Lea SM (2012) Structure of the TatC core of the
twin-arginine protein transport system. Nature 492: 210-4
Stanley NR, Palmer T, Berks BC (2000) The twin arginine consesus motif of Tat signal peptides
is in volved in Sec-independent protein targeting in E. coli. J Biol Chem 0000: 00-00
Stockwald ER, Steger LME, Vollmer S, Gottselig C, Grage SL, Bürck J, Afonin S, Fröbel J,
Blümmel AS, Setzler J, Wenzel W, Walther TH, Ulrich AS (2022) Length matters: functional
flip of the short TatA transmembrane helix. Biophys J
Teter SA, Theg SM (1998) Energy-transducing thylakoid membranes remain highly
impermeable to ions during protein translocation. Proc Natl Acad Sci U S A 95: 1590-4
Theg SM, Tom C (2011) Measurement of the DeltapH and electric field developed across
Arabidopsis thylakoids in the light. Methods Mol Biol 775: 327-41
van den Brink-van der Laan E, Chupin V, Killian JA, de Kruijff B (2004) Stability of KcsA
tetramer depends on membrane lateral pressure. Biochemistry 43: 4240-50
Walther TH, Gottselig C, Grage SL, Wolf M, Vargiu AV, Klein MJ, Vollmer S, Prock S,
Hartmann M, Afonin S, Stockwald E, Heinzmann H, Nolandt OV, Wenzel W, Ruggerone P,
Ulrich AS (2013) Folding and self-assembly of the TatA translocation pore based on a charge
zipper mechanism. Cell 152: 316-26
Yang L, Harroun TA, Weiss TM, Ding L, Huang HW Barrel-Stave Model or Toroidal Model? A
Case Study on Melittin Pores. Biophys J 81: 1475-1485
Yuan J, Jin F, Glatter T, Sourjik V (2017) Osmosensing by the bacterial PhoQ/PhoP two-
component system. Proc Natl Acad Sci U S A 114: E10792-e10798
Zhang Y, Wang L, Hu Y, Jin C (2014) Solution structure of the TatB component of the twin-
arginine translocation system. Biochim Biophys Acta 1838: 1881-8
Zhou W, Hao B, Bricker TM, Theg SM (2023) A Real-Time Analysis of Protein Transport via
the Twin Arginine Translocation Pathway in Response to Different Components of the
Protonmotive Force in vivo. bioRxiv: 2023.01.12.523868
Zoufaly S, Frobel J, Rose P, Flecken T, Maurer C, Moser M, Muller M (2012) Mapping
precursor-binding site on TatC subunit of twin arginine-specific protein translocase by site-
specific photo cross-linking. J Biol Chem 287: 13430-41
619

Figures

Figure 1. CPPs specifically stimulate transport on the Tat pathway. In A-D, thylakoids
were incubated with a radiolabeled substrate and CPPs at the indicated concentrations. A)
Thylakoids were treated with CPPs during a transport reaction of iOE17. Quantification of the
mature protein is shown. B) Thylakoids were incubated in the dark for 30 min (D) or light (L)
with iOE17 and with or without 5 μM penetratin. C) Thylakoids were treated with given HIVTat
concentrations and mixed with the transport incompetent precursor, iOE17KK. D) Thylakoids

were treated with HIVTat as indicated and assayed for transport of iOE33 through the Sec

pathway. Post-transport thylakoids were treated with thermolysin (therm) where indicated. E)

ECS measurements were made on 80 μg thylakoids with the indicated concentrations of melittin.

Two 9 ms-duration flashes were delivered 100 ms apart (arrows at the bottom of the panel); the

ECS signal was normalized to the first data point after the second flash. F) ΔpH measurements

were made by observing the quenching of NED fluorescence induced by application of actinic

light with indicated concentrations of melittin.

A

Control   1μM Melittin

20%   ΔpH

-i
-m

B

C

Control   2μM Penetrattin

ΔpH
20%   10%

-i
-m

D

637   **Figure 2. CPPs lower the ΔpH threshold for Tat transport**. ΔpH measurements were made

638   by observing the quenching of NED fluorescence by application of actinic light of varying

639   intensities. Transport reactions were stopped with 1mM nigericin and valinomycin immediately

640   followed by centrifugation and resuspension in SB. Protein transport was measured as in Fig. 1.

Figure 3. Thylakoids pellet more readily in the presence of CPPs. Thylakoids were

incubated with CPPs at indicated concentrations in the dark for 10 min. Thylakoids were then

centrifuged at 100 rcf for 1 min. The supernatant was then diluted tenfold into 89% Acetone.

Chlorophyll concentration was determined spectroscopically.

Figure 4. Stimulation of Tat transport in E. coli IMVs by CPPs. Biotinylated pre-SufI-IAC

substrate protein was transported into IMVs at 37 °C with indicated concentrations of CPPs.

Reactions were stopped after 8 min on ice. Samples were treated with Proteinase K and mature

protein was analyzed by SDS-PAGE and α-biotin Western blotting. Error bars indicate SD

(n≥4). Significance of CPP effect was determined by ANOVA post-hoc analysis (*$p<0.05$,

**$p<0.01$).

A

Transport in the presence
of Tha4 APH

B

Pelleting of TAT APHs

C

ΔpH of APHs

D

Normalized ECS
of TAT APHs

E

Transport in the
presence of Hcf106

F

Thermolysin digestion
of Hcf106 transport

Figure 5. Transport of iOE17 in the presence of Tat APHs. A) As in Fig. 1, thylakoids were

incubated with the Tha4 APH peptide and transport was performed. B) Pelleting assays were

performed as in Fig. 3. C) ΔpH analysis was performed as in Fig. 1. D) ECS measurements were

performed as in Fig. 1. E) As in Fig. 1, thylakoids were incubated with Hcf106 APH and

transport was performed. F) Transport reactions were thermolysin treated as indicated (+).

Protease was quenched with 25mM EDTA. Thylakoid reisolation and SDS-PAGE followed as

in Fig. 1.

Figure 6. Inhibition of Tat transport by TFE. Thylakoids were assayed for Tat (iOE17, blue)

or Sec (iOE33, orange) transport in the presence of indicated TFE concentrations. Thylakoids

were supplemented with ethanol such that the sum total concentration (%v/v) of TFE and ethanol

was equivalent in all reactions. Post-transport samples were treated with thermolysin (therm)

where indicated and analyzed by SDS PAGE and fluorography. Mature bands were quantified

(n≥3, SD). Thylakoid ΔpH was measured at indicated TFE concentrations (green, n=3, SD) as

described previously by 9AA fluorescence quenching (Theg & Tom, 2011).