Meta-analysis of gonadal transcriptomes in protogynous fishes provides novel perspective into

sex change machinery

Ryo Nozu[1,2], Mitsutaka Kadota[3], Masaru Nakamura[4], Shigehiro Kuraku[3,5,6], Hidemasa Bono[1,2*]

[1]Laboratory of Genome Informatics, Graduate School of Integrated Sciences for Life,

Hiroshima University, 3-10-23 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-0046, Japan

[2]Laboratory of BioDX, Genome Editing Innovation Center, Hiroshima University, Hiroshima,

Japan

[3]Laboratory for Phyloinformatics, RIKEN Center for Biosystems Dynamics Research (BDR),

Kobe, Japan

[4]Okinawa Churashima Research Center, Okinawa Churashima Foundation, Motobu-cho,

Okinawa, Japan

[5]Molecular Life History Laboratory, Department of Genomics and Evolutionary Biology,

National Institute of Genetics, Mishima, Japan

[6]Department of Genetics, Sokendai (Graduate University for Advanced Studies), Mishima,

Japan.

[*]To whom correspondence may be addressed: Hidemasa Bono

bonohu@hiroshima-u.ac.jp

Abstract

Protogynous fishes can change sex from female to male; concomitantly, their ovarian tissue

is completely replaced by testicular tissue. The high functional conservation of gonads across

organisms suggests the existence of as-yet-undefined common and essential factors in the

regulation of such gonadal transformation. To identify these factors, we conducted a

meta-analysis of public and newly derived gonadal transcriptome data from seven species.

Classical sex change-related genes were differentially expressed between the ovary and

transitional gonads, which corroborated previous reports on protogynous fishes. Moreover, we

identified novel factors not previously associated with gonadal sex change in fish. Notably, the

expression patterns of acetoacetyl-CoA synthetase (*aacs*) and apolipoprotein Eb (*apoeb*),

which are involved in cholesterol synthesis and catabolic processes, respectively, suggest that

the levels of cholesterol, a precursor of steroid hormones crucial for sex change, are decreased

upon sex change onset. Thus, cholesterol dynamics may influence gonadal transformation in

protogynous fishes.

Introduction

Sequential hermaphroditism, a phenomenon in which individuals undergo sex change during their lifetime, has been extensively recognized in teleosts over the past several decades[1]. Specifically, protogyny (i.e., the transformation from female to male) is prevalent in 20 families[2]. Phylogenetic analyses have shown that this characteristic has been independently acquired in various evolutionary lineages[3,4]. On the other hand, the complete replacement of ovarian tissue with testicular tissue is commonly observed during protogynous sex change[5-9], the intricate regulatory mechanisms responsible for inducing these remarkable transformations in gonadal tissue remain incompletely understood. Furthermore, considering the high functional conservation of gonads across organisms, the existence of a regulatory mechanism governing gonadal transformation comprising common and essential factors across protogynous fishes is plausible; however, a comprehensive perspective of this process remains elusive.

Accordingly, elucidation of the molecular mechanisms underlying the gametogenesis switch in sex-changing fish has subsequently become a focus of research. In particular, the expression patterns of multiple sex-related genes during sex change within the family Labridae, which serves as a model for protogynous sex change, have been explored[10-13]. Recently, next-generation sequencing technology has enabled researchers to conduct comprehensive analyses of gene expression, allowing the identification of global gene expression profiles during gonadal transformation in protogynous fish belonging to Labriformes, Perciformes, and Synbranchiformes[14-18]. Thus, these results suggest the involvement of several pivotal genes in the ovary-to-testis transformation in sex-changing fish.

For example, a repertoire of sex-related genes, including cytochrome P450, family 19, subfamily A, polypeptide 1a (*cyp19a1a*), forkhead box L2a (*foxl2a*), folliculogenesis specific bHLH transcription factor (*figla*), anti-Müllerian hormone (*amh*), gonadal somatic cell derived factor (*gsdf*), doublesex and mab-3 related transcription factor 1 (*dmrt1*) were identified as

differentially expressed genes (DEGs) during sex change across protogynous fishes. Notably,

the gonadal expression of *cyp19a1a* and *foxl2a*, which are key players in maintaining ovarian

status, is reduced upon the onset of sex change in multiple species[13,19,20]. *cyp19a1a* encodes

aromatase, an enzyme responsible for the conversion of androgens (e.g., testosterone) to

estrogens (e.g., estradiol-17ß; E2), whereas *foxl2a* encodes a transcription factor that targets

and up-regulates *cyp19a1a* in teleost fishes[21,22]. Furthermore, female-to-male sex change is

triggered by a rapid decline in E2 levels[6]. The expression profiles of these genes coordinate

well with the reduction in E2 levels upon sex change onset. Conversely, *amh*, *gsdf*, and *dmrt1*

are crucial for testicular differentiation and development in teleost fishes[23-29]. In several

sex-changing species, the upregulation of these genes coincides with the emergence of

testicular tissue during transition, suggesting that these genes are involved in testicular

formation during gonadal sex change[10-13,30].

Transcriptome analysis during gonadal transformation has also revealed differential

expression of several genes not previously implicated in sex determination or differentiation

pathways. In particular, upregulation of apolipoprotein Eb (*apoeb*), ectonucleotide

pyrophosphatase/phosphodiesterase 2 (*enpp2*), and l-rhamnose-binding lectin CSL2 (*csl2*)

expression, accompanied by downregulation of galectin-3 (*gal3*), was observed during the

early phase of gonadal sex change in the protogynous Asian swamp eel *Monopterus albus*.

These findings suggest that these factors are involved in gonadal transformation, particularly

during the initiation of sex changes[14,17]. However, verifying their direct involvement in the sex

change process remains challenging. Therefore, a novel approach is required to obtain

additional evidence.

The recent exponential growth of high-throughput sequencing data available in public

databases has opened new avenues for the effective utilization of meta-analysis. By

amalgamating transcriptome data from multiple studies, meta-analysis has attracted attention

as a powerful tool capable of providing novel insights overlooked by traditional

hypothesis-driven research methods. Indeed, several reports have described the discovery of

previously unrecognized gene expression profiles through meta-analysis of publicly available

transcriptome data[31-35]. Moreover, the commonalities revealed by meta-analytical approaches

not only provide more solid evidence than that from individual studies but also serve as

valuable launching points for data-driven research.

Therefore, the objective of the present study was to identify the shared factors involved in

gonadal transformation across species using a meta-analysis of publicly available

transcriptome data from multiple experiments and species. Moreover, to improve the

performance of the meta-analysis, we included data (*de novo* genome assembly data and RNA

sequencing (RNA-Seq) data from gonads undergoing sex change) generated in-house from an

additional well-studied species, the three-spot wrasse, *Halichoeres trimaculatus*, a

protogynous species belonging to the family Labridae[10-12]. Through meta-analysis of

RNA-Seq data encompassing seven protogynous species from three orders (Fig. 1), we

identified 168 DEGs between the transitional gonads and ovaries. Using bioinformatics and

gene set enrichment analyses, we reevaluated the relationship between these factors and

gonadal sex change, providing a novel perspective regarding the mechanisms underlying this

process.

Results

***De novo* genome assembly and transcriptome of gonadal tissue in the three-spot wrasse.**

High-throughput genome sequence data were obtained from a locally obtained female

three-spot wrasse from Okinawa, Japan. The obtained HiFi data, which comprised

approximately 32 Gb and encompassed approximately 1.9 million DNA sequences, were

assembled using the hifiasm program[36]. Subsequently, the acquired Hi-C data were used for

scaffolding, resulting in the generation of 227 scaffolds. The longest scaffold measured 44.2

Mb, with an N50 length (the shortest of the lengths covering 50% of the genome) of 37.9 Mb,

contributing to a total assembly length of approximately 849.9 Mb. Assessment using

BUSCO[37] revealed a completeness > 98%, indicating a high level of coverage and contiguity

in the *de novo* genome assembly of the three-spot wrasse (Table 1).

The RNA-Seq reads obtained from transitional gonads and ovaries of additional females

(n=2 each group) were mapped to the genome assembly, and the obtained mapping results

were used to assemble the transcriptome for the annotation file ("merged.gtf"). This process

yielded 27,573 transcript sequences (Table 2). Subsequently, 24,695 coding sequences (protein

sequences) were predicted from the set of transcript sequences using TransDecoder[38].

Data curation of public RNA-Seq data and genomic information. We downloaded

Sequence Read Archive (SRA) data from six fish species for which sex changes have been

reported:[2] bluehead *Thalassoma bifasciatum* (order Labriformes); orange-spotted grouper

Epinephelus coioides, Hong Kong grouper *E. akaara,* and black seabass *Centropristis striata*

(order Perciformes); and Asian swamp eel *Monopterus albus* and zig-zag eel *Mastacembelus*

armatus (order Synbranchiformes). Furthermore, using the metadata associated with the SRA

data, we curated the RNA-Seq data into comparable sample pairs of intersexual (transitional)

gonads and ovaries (a total of 31 paired data, including three-spot wrasse). The data pairs

analyzed in this study are listed in Supplementary Table 1.

The genome sequences of five of the seven species utilized in this study are publicly

available (Table 2) and served as reference genomes for read mapping. The genome sequences

of *T. bifasciatum* (GCA_008086565.1), *E. coioides* (GCA_900536245.1), *M. albus*

(GCF_001952655.1), and *M. armatus* (GCF_900324485.2) were retrieved from the National

Center for Biotechnology Information (NCBI) Genome Database (accessed on October 2022).

We also referenced the genome information of *E. akaara* reported by Ge et al.[39].

Transcriptome assembly and inference of ortholog relationships. For the five species with

public reference genomes, the number of transcript sequences obtained using StringTie[40] with

the "--merge" option is shown in Table 2. For black sea bass, (without a reference genome), the

number of transcriptome sequences assembled using Trinity[41] was 594,726. Among these,

95,941 transcript sequences corresponding to the predicted coding sequences (CDS) were

specified as references for expression quantification using Salmon[42]. The number of protein

sequences predicted using TransDecoder is shown in Table 2. Orthology inference and

consolidation of orthologs within the three-spot wrasse identified 7,289 orthologs that were

common to all species used in this study (Table 2), and thus were considered as candidate

sex-change factors.

Identification of differentially expressed transcripts. The total intersexual gonad and ovary

score (IO-score) was calculated for these 7,289 factors (Supplementary Table 2). Among these,

99 factors were upregulated (Fig. 2a, Fig. 3, Supplementary Table 3) and 69 factors were

downregulated (Fig. 2a, Fig. 4, Supplementary Table 4). Whereas the total IO-score of these

factors was occasionally derived from a single species (up: *cep97*, *zmat2*, *tmem237a*, *supt5h*;

down: *znf16l*, *lrrc8db*, *sh2b1*, *tmem263*, *cyldb*, *manea*, *nol11*), in most cases, the total

IO-scores were based on contributions from two or more species. Notably, the upregulated

genes included *amh* and *gsdf*, which have been previously reported to play important roles in

gonadal sex change, particularly in testicular formation and development, in some protogynous

fish. Conversely, the downregulated genes included *cyp19a1a*, *hsd17b1*, and *foxl2a*, which are

crucial for the onset of sex change (Table 3).

Enrichment analysis of DEGs. Sequence homology search of the three-spot wrasse proteins

using ggsearch36[43] against the reference zebrafish proteins (Ensembl Release 108)[44] resulted in

the identification of 90 homologs among the 99 upregulated factors (Supplementary Table 3).

No corresponding zebrafish orthologs were found for the remaining nine factors that were

common to all species analyzed in this study. The results of enrichment analysis using the 90

factors that could be converted to zebrafish orthologs are shown in Fig. 2b. The Gene Ontology

(GO) terms "homologous chromosome pairing at meiosis" and "germ cell development,"

which are related to reproduction, were significantly enriched among the upregulated factors.

The specific genes annotated with these GO terms included *hormad1*, *terb2*,

si:ch211-199g17.9, *gsdf*, and *sycp3* (Table 3).

We successfully obtained the corresponding zebrafish homologs for all of the 69

downregulated factors. The results of enrichment analysis are shown in Fig. 2c. Among the

downregulated factors, GO terms related to the regulation of steroid hormone levels, including

"regulation of hormone levels" and "steroid metabolic process," were significantly enriched.

Specific genes annotated with these enriched GO terms included *cyp19a1a*, *aacs*, *hsd17b1*,

ece1, *amacr*, and *sdr42e1* (Table 3).

Discussion

In this study, we conducted a meta-analysis of gonadal RNA-Seq data from seven protogynous

fish species using a unified analysis pipeline and threshold. The aim was to compare the

transcriptomes of ovaries and early sexual transition gonads, with the goal of identifying

common factors involved in protogynous gonadal sex change. Toward this end, we first

assembled the *de novo* genome of the three-spot wrasse. The genome assemblies of various

species from the family Labridae are listed in the NCBI genome database (accessed May 2023).

The assembly statistics of the three-spot wrasse (Table 1) were comparable to those of other

Labridae species, such as humphead wrasse (*Cheilinus undulatus*, GCF_018320785.1), ballan

wrasse (*Labrus bergylta*, GCF_900080235.1), and New Zealand spotty (*Notolabrus celidotus*,

GCF_009762535.1), previously deposited in the NCBI RefSeq database. These results

indicated that the *de novo* genome assembly of the three-spot wrasse exhibited high quality,

enabling us to perform expression analysis and orthology inference with confidence.

Through this meta-analysis, we successfully identified sex change-related factors including

cyp19a1a, *foxl2a*, *hsd17b1*, *amh*, and *gsdf*, that had been previously reported to play a role in

sex change in various protogynous fish species[10,13,15,17]. Furthermore, our enrichment analysis

of downregulated factors highlighted a term related to the "regulation of steroid hormone

production." This finding is consistent with those of previous reports, which indicate that a

rapid decrease in estrogen levels is crucial for female-to-male sex change[6,45,46]. Thus, by

pooling data across species, we were able to identify common factors involved in gonadal sex

change. Notably, our meta-analysis demonstrates the potential of utilizing transcriptome data

from multiple experiments to gain insights regarding the mechanisms underlying sex change in

fish.

Through the enrichment analysis, we identified several factors that have not been

previously associated with gonadal sex change. Among the up-regulated factors, we observed

the inclusion of "*hormad1*," "*terb2*," "*si:ch211-199g17.9*," and "*sycp3*," which are annotated

with a GO term related to "germ cells" (Table 3). Interestingly, *hormad1* and *sycp3* have been

reported to exhibit higher expression levels in the testis compared to the ovary in zebrafish[47].

Based on these facts, the increased expression of these factors may indicate the progression of

spermatogenesis during gonadal sex change. The timing of spermatogenesis during gonadal

sex change can vary among species. In Serranidae, spermatogenesis is observed early phase in

the sex change process, representing a histological characteristic at the onset of sex change

[9,20,48]. Conversely, in Labridae species such as the three-spot wrasse and bluehead, the

histological characteristic at the onset of sex change is the degeneration of ovarian tissue, and

spermatogenesis is histologically observed in the later stages of sex change [6,15,49,50]. In our

study, the RNA-seq data from Labridae species were obtained from gonads at an early stage of

sex change. Therefore, the unexpected increase in the expression of multiple factors involved

in spermatogenesis suggests that testicular formation including spermatogenesis may

commence early during gonadal transformation, even if it is not histologically recognizable.

This characteristic appears to be common to protogynous sex change.

Our findings demonstrate that *aacs*, annotated with the GO term "regulation of hormone

levels," is downregulated during sex change. Notably, no previous reports have suggested the

involvement of *aacs* in sex change, indicating that the encoded protein (acetoacetyl-CoA

synthetase; AACS) may represent a novel factor associated with this process. Typically, AACS

plays a critical role in cholesterol supply. Studies in mice have shown that AACS knockdown

decreases total serum cholesterol levels[51]. Additionally, in rat adrenal glands, inhibition of

AACS expression through hypermethylation suppresses cholesterol supply and steroid

synthesis[52]. Moreover, apolipoprotein Eb (encoded by *apoeb*, listed among the upregulated

factors), has been suggested to participate in cholesterol catabolism[53]. These observations

strongly support the hypothesis that the downregulation of *aacs* and the upregulation of *apoeb*

during sex change contribute to suppression of the cholesterol supply.

Cholesterol serves as a precursor for steroid hormones, including estrogens, androgens, and

glucocorticoids[54-56], which play crucial roles in protogynous sex change. Previous studies have

shown that a decline in endogenous estrogens, particularly E2, triggers the transition from

ovary to testis in protogynous fish[6,45,46]. E2 is synthesized from testosterone via an aromatase

enzyme, encoded by *cyp19a1a*. Downregulation of *cyp19a1a* expression at the onset of sex

change leads to reduced E2 production. Studies on *M. albus* and *T. bifasciatum* have revealed

higher DNA methylation levels of the *cyp19a1a* gene upstream region in the testes than those

in the ovaries, suggesting that epigenetic modifications may contribute to the suppression of

cyp19a1a expression during gonadal sex change[15,19]. Our present findings introduce an

alternative hypothesis, suggesting that the decreased E2 production at the initiation of sex

change may be attributed not only to the downregulation of aromatase activity but also to

suppression of the cholesterol supply, the precursor of steroid hormones. Although this

hypothesis requires further investigation and an understanding of cholesterol dynamics during

sex change, additional exploration is warranted. Notably, a study on medaka (*Oryzias latipes*)

subjected to starvation conditions implied a potential link between masculinization and lower

cholesterol levels[57]. Specifically, female-to-male sex reversal was observed in starved medaka

larvae, a gonochoristic fish, together with decreased cholesterol levels in the starved samples.

However, the precise pathways underlying masculinization and cholesterol dynamics remain

unknown. In general, cholesterol is synthesized not only in the liver and adrenal glands but

also in the gonads[58-60]. Therefore, gaining a comprehensive understanding of the lipid

metabolic mechanisms in fish gonads, including cholesterol synthesis and degradation during

sex change, may provide valuable insights regarding the mechanisms underlying gonadal

transformation. To clarify this perspective, the remaining challenge lies in providing

experimental evidence regarding whether it is possible to control gonadal transformation by

actually regulating cholesterol levels within the body.

In summary, we conducted a comprehensive meta-analysis by integrating transcriptome

data from multiple protogynous fishes spanning three orders. Our aim was to compare the

transcriptomes of sex-transitional gonads with those of ovaries using a unified pipeline to

identify the factors involved in gonadal sex change. Through this analysis, we identified 168

differentially expressed factors between the two gonadal states. These factors are likely to play

a universal role in gonadal sex change across diverse taxa of protogynous fishes. Notably, we

identified novel factors that were not previously associated with gonadal sex change. In

particular, we identified cholesterol dynamics as an important factor regulating the sex change

process. Moreover, this finding highlights the potential of transcriptomics meta-analysis as a

valuable approach to uncover overlooked mechanisms underlying conserved biological

processes including sex change. We anticipate that further data-driven research utilizing these

novel sex change-related factors will lead to new horizons in sex change research.

Methods

Animals. The adult female three-spot wrasse specimens collected in the field by local

fisherman in the northern region of Okinawa, Japan were purchased and employed for genome

and RNA sequencing. These specimens were subsequently cultivated in 1000 L tanks receiving

a continuous supply of unregulated seawater, and were subjected to natural photoperiod

conditions until sampling.

The maintenance and handling of fish in this study were conducted in accordance with the

"Guidelines for Animal Experiments of the Okinawa Churashima Foundation," with the same

consideration for animal care and welfare as that for higher vertebrates, including reptiles,

birds, and mammals. Animal ethical review and approval were not required for the present

study because the guidelines stipulated that approval from the Institutional Animal Care and

Use Committee of Okinawa Churashima Foundation is required for "higher" vertebrates, and is

waived for "lower" vertebrates including fish.

Genome sequencing and assembly for the three-spot wrasse. Muscle tissue from a mature

female individual was used for genome sequencing. The specimen was randomly selected from

cultivated tank. After anesthetizing the fish with 0.05% v/v 2-phenoxyethanol (FUJIFILM

Wako Pure Chemical), the muscle tissue was dissected, immediately frozen in liquid nitrogen,

and stored at −80 °C until genomic DNA extraction. High-molecular-weight DNA was

extracted from muscle using a NucleoBond AXG column (Macherey–Nagel), followed by

purification with phenol–chloroform. The concentration of the extracted DNA was measured

using Qubit (Thermo Fisher Scientific Inc.), and its size distribution was first analyzed with a

TapeStation 4200 (Agilent Technologies) to ensure high integrity, and subsequently analyzed

via pulse-field gel electrophoresis on a CHEF DR-II system (BioRad) to ensure a size range of

20–100 kb.

To obtain long-read sequence data, a SMRT sequence library was constructed using the

SMRTbell Express Template Prep Kit 2.0 (PacBio) and sequenced on a PacBio Sequel II

system. The sequencing output was processed to generate circular consensus sequences to

obtain HiFi sequence reads. Adapter sequences were removed from the HiFi reads using

HiFiAdapterFilt[61]. The obtained HiFi sequence reads were assembled using the hifiasm

(v0.16.1) assembler[36] with default parameters. The obtained contigs were first scaffolded with

the P_RNA_scaffolder[62] and further scaffolded using Hi-C data as follows. The Hi-C library

was prepared with the in situ method using the muscle tissue of the individual used for DNA

extraction according to the iconHi-C protocol[63], employing the restriction enzymes DpnII and

HinfI, and sequenced on a HiSeq X sequencing platform (Illumina). The obtained Hi-C read

pairs, amounting to 232 million read pairs, were aligned to the HiFi sequence contigs using the

Juicer program[64], and the HiFi sequence contigs were scaffolded via the 3d-dna to be

consistent with the chromatin contact profiles through manual curation on JuiceBox[65]. The

continuity and completeness of the resulting genome assembly were assessed using the

webserver gVolante[66], in which the pipeline BUSCO (v5.4, reference datasets:

vertebrata_odb10) was implemented[37].

RNA-Seq of gonadal tissue in the three-spot wrasse. Based on previous studies[10,45], we

triggered a sex change in three-spot wrasse 5 females (body length; 8.0–9.3 cm, body weight;

12.4–20.9 g) to obtain transitional gonads through treatment with an aromatase inhibitor (AI),

which exerts inhibitory effects on the enzyme responsible for estrogen synthesis. The collected

gonads were cut into small pieces, one piece was immersed in TRIzol reagent (Invitrogen) and

the rest was fixed in 4% paraformaldehyde/1× phosphate-buffered saline (PBS) for histological

observation. Total RNA in each sample was extracted promptly in accordance with the TRIzol

instruction manual. The extracted total RNA was dissolved in RNase-free water and stored at

−80 °C until RNA-Seq analysis. We selected two treated and two control samples for

RNA-Seq analysis on the basis of histological observation (Supplementary fig. 1).

Beads with oligo (dT) were used to isolate poly(A) mRNA after the total RNA was

collected from the samples. Fragmentation buffer was added to disrupt mRNA into short

fragments. Using these short fragments as templates, a random hexamer primer was used to

synthesize the first-strand cDNA. Second-strand cDNA was synthesized using buffer, dNTPs,

RHase H, and DNA polymerase I. Short fragments were purified using the QiaQuick PCR

extraction kit (Qiagen) and resolved with EB buffer for end reparation and adding an "A" base.

Subsequently, the short fragments were connected to sequencing adapters. Following agarose

gel electrophoresis, suitable fragments were selected as templates for PCR amplification.

Finally, the library was sequenced using an Illumina HiSeq 2000 as 2×90 bp paired-ends. The

acquired reads were subjected to the quality control procedures described below and used for

downstream expression analysis.

Retrieval and acquisition of RNA-Seq data in protogynous fishes from public resources.

To obtain RNA-Seq data from the gonads of protogynous fishes in public databases, we

accessed SRA[67] (accessed in October 2022) to search SRA data containing the following

keywords: "bony fishes (AND ovary OR gonad)." SRA data were converted into FASTQ files

using the Fasterq-dump program of the SRA Toolkit (v3.0.0).

Quality control of RNA-Seq data. To ensure the integrity of the RNA-Seq reads, trimming

and quality control procedures were performed. Specifically, we used Trim Galore (v0.6.7)[68]

with the option "--quality 20 --paired" to eliminate low-quality bases from paired-end reads.

Read mapping against the reference genome. For species with available genomic

information, RNA-Seq reads were mapped to the reference genome sequences using HISAT2

(v2.2.1)[69] with the parameter "-q -dta -x". The resulting SAM files were sorted and converted

into BAM files using SAMtools software (v1.15.1)[70].

Generating General Transfer Format (gtf) files based on genome sequence and RNA-Seq

reads for reference gene annotation when quantifying expression values. To ensure

consistency in the use of annotation information for expression quantification in species with

available reference genomes, we employed the merge function of StringTie (v2.1.7)[40] to

generate a gtf file that served as a guide for annotations. The transcriptome assembly for each

fish was conducted using StringTie with default settings based on the mapping results (BAM

file). The resulting gtf files were then merged using StringTie with the "--merge" option,

producing a unified file as the "merged.gtf". This "merged.gtf" file contained a non-redundant

set of assembled transcripts for each species. Generated "merged.gtf" files in each species were

designated as the reference annotation files for quantifying transcript expression.

Quantification of transcripts expression and detection of differentially expressed

transcripts. When reanalyzing high-throughput data from multiple experiments, it is crucial to

consider batch effects[71]. In the present study, we calculated the transcript per million (TPM)

value for each transcript and utilized the difference in TPM values between the paired data as

an indicator for identifying differentially expressed transcripts.

Transcript expression was quantified in species with available genome annotation using

StringTie, specifying the parameters as "-G merged.gtf -e". The resulting gtf file for each

sample provided the TPM values, which were used for downstream analysis.

TPM values were computed for species lacking reference genomes using Salmon

(v.1.9.0)[42] with the parameters "quant -i index -l A". To generate the transcriptome sequences

required for Salmon, we performed transcriptome assembly using the Trinity (v2.13.1)

assembler[41] with default settings. Open reading frames (ORFs) and CDS were predicted from

the transcriptome assembly using TransDecoder (v5.5.0)[38] with the same settings as described

below. The resulting gff3 file obtained from the TransDecoder run was used to extract

transcript sequences corresponding to the predicted CDS from the transcriptome assembly. The

extracted sequence set was then utilized as the reference transcriptome after eliminating

duplicate sequences using seqkit (v2.3.0)[72] with the "rmdup -s" command.

To assess changes in transcript expression, the expression ratio was determined using TPM

values obtained from intersexual gonad and ovary transcriptomes (referred to as the IO-ratio).

The IO-ratio for each transcript was calculated using Equation (1).

$$\text{IO ratio} = \log_2(\text{TPM}_{\text{intersexual}} + 0.01) - \log_2(\text{TPM}_{\text{ovary}} + 0.01), (1)$$

This calculation was performed for all pairs of intersexual gonad and ovary transcriptomes. To

convert zeros to logarithms, a value of "0.01" was added to the TPM of each transcript.

Transcripts with an IO-ratio $> \log_2(5)$ (indicating a 5-fold expression change) were classified

as "upregulated," whereas transcripts with an IO-ratio $< -\log_2(5)$ (representing a 0.2-fold

expression change) were considered "downregulated." Transcripts outside these thresholds

were deemed "unchanged." To determine differential expression between the intersexual gonad

and ovary, an IO-score was calculated for each transcript by subtracting the count of

"downregulated" pairs from that of "upregulated" pairs, as shown in Equation (2).

$$\text{IO-score} = (\text{count number}_{\text{upregulated}}) - (\text{count number}_{\text{downregulated}}), (2)$$

Using orthology inference as described below, the total IO-score for each transcript was

calculated by summing the IO-scores of all species. Total IO-score thresholds of 7 and −7 were

employed to identify transcripts that were differentially upregulated and downregulated,

respectively. All calculations for IO-ratios and IO-scores were performed using R (v 4.2.1)[73].

Orthology inference among species. GFFread (v0.12.1)[74] was utilized along with the

generated "merged.gtf" file to extract transcript sequences from the reference genomes. For

species without reference genomes, transcript sequences were obtained via assembly using

Trinity, as described previously. TransDecoder was used to identify the coding regions and

translate them into amino acid sequences for all species. ORFs were extracted from the

transcriptome sequences using TransDecoder Longorfs with the parameters "-m 100". CDS

were then predicted using TransDecoder Predict with default settings, and the protein

sequences converted from the predicted CDS were acquired. Subsequently, a reciprocal

homology search was conducted between the predicted protein sequences of the three-spot

wrasse and those of other species, and the reciprocal best hit (RBH) was identified as the

ortholog. The program Diamond (v2.0.15)[75] was used for the RBH search using the parameters

"--ultra-sensitive -e 1e-3". To extract the orthologs shared among all species, we consolidated

the RBH results with the three-spot wrasse-predicted protein IDs.

Gene set enrichment analysis. Differentially expressed gene sets were analyzed using the

web tool Metascape (v3.5)[76], which enables gene set enrichment analysis. However, Metascape

only supports the analysis of zebrafish among teleosts. Therefore, prior to analysis, the

predicted protein sequences of the three-spot wrasse were mapped to the corresponding

zebrafish Ensembl protein IDs. To obtain orthologous information, we utilized the ggsearch36

program from the FASTA package[43] to globally align the predicted protein sequences of the

three-spot wrasse with the proteome sequences of zebrafish (Ensembl Release 108)[44].

Data availability. All sequencing data, including assembled sequences and raw sequence reads,

were deposited in the DNA Data Bank of Japan (DDBJ) under the umbrella BioProject

accession number PRJDB15835. The genome assembly was deposited in the DDBJ under

accession numbers BSYP01000001–BSYP01000227. Additionally, raw sequence reads were

deposited in the DDBJ under the following accession numbers: DRR466686 (PacBio HiFi

reads), DRR486910 (Illumina HiC reads), and DRR483512-DRR483515 (Illumina RNA-Seq

reads).

References

1.   Atz JW. Intersexuality in fishes. In: *Intersexuality in Vertebrates Including Man* (eds Armstrong CN, Marshall AJ). Academic press (1964).

2.   Kuwamura T, Sunobe T, Sakai Y, Kadota T, Sawada K. Hermaphroditism in fishes: an annotated list of species, phylogeny, and mating system. *Ichthyol Res* **67**, 341–360 (2020).

3.   Avise JC, Mank JE. Evolutionary perspectives on hermaphroditism in fishes. *Sex Dev* **3**, 152–163 (2009).

4.   Erisman BE, Petersen CW, Hastings PA, Warner RR. Phylogenetic perspectives on the evolution of functional hermaphroditism in teleost fishes. *Integr Comp Biol* **53**, 736–754 (2013).

5.   Lo Nostro F, Grier H, Andreone L, Guerrero GA. Involvement of the gonadal germinal epithelium during sex reversal and seasonal testicular cycling in the protogynous swamp eel, *Synbranchus marmoratus* Bloch 1795 (Teleostei, Synbranchidae). *J Morphol* **257**, 107–126 (2003).

6.   Nakamura M, Hourigan TF, Yamauchi K, Nagahama Y, Grau EG. Histological and ultrastructural evidence for the role of gonadal steroid hormones in sex change in the protogynous wrasse *Thalassoma duperrey*. *Environ Biol Fishes* **24**, 117–136 (1989).

7.   Ohta K, Hirano M, Mine T, Mizutani H, Yamaguchi A, Matsuyama M. Body color change and serum steroid hormone levels throughout the process of sex change in the adult wrasse, *Pseudolabrus sieboldi*. *Mar Biol* **153**, 843–852 (2008).

8.   Muncaster S, Norberg B, Andersson E. Natural sex change in the temperate protogynous Ballan wrasse *Labrus bergylta*. *J Fish Biol* **82**, 1858–1870 (2013).

9.   Bhandari RK, Komuro H, Nakamura S, Higa M, Nakamura M. Gonadal restructuring and correlative steroid hormone profiles during natural sex change in protogynous honeycomb grouper (*Epinephelus merra*). *Zoolog Sci* **20**, 1399–1404 (2003).

10.   Horiguchi R, Nozu R, Hirai T, Kobayashi Y, Nagahama Y, Nakamura M. Characterization of gonadal soma-derived factor expression during sex change in the protogynous wrasse, *Halichoeres trimaculatus*. *Dev Dyn* **242**, 388–399 (2013).

11.   Nozu R, Horiguchi R, Kobayashi Y, Nakamura M. Expression profile of doublesex/male
abnormal-3-related transcription factor-1 during gonadal sex change in the protogynous wrasse,
Halichoeres trimaculatus. *Mol Reprod Dev* **82**, 859–866 (2015).

12.   Horiguchi R, Nozu R, Hirai T, Kobayashi Y, Nakamura M. Expression patterns of sex
differentiation-related genes during gonadal sex change in the protogynous wrasse, *Halichoeres*
trimaculatus. *Gen Comp Endocrinol* **257**, 67–73 (2018).

13.   Thomas JT*, et al.* Conservation and diversity in expression of candidate genes regulating
socially-induced female-male sex change in wrasses. *PeerJ* **7**, e7032 (2019).

14.   Senthilkumaran B, Chi W, Gao Y, Hu Q, Guo W, Li D. Genome-wide analysis of brain and
gonad transcripts reveals changes of key sex reversal-related genes expression and signaling
pathways in three stages of *Monopterus albus*. *PloS one* **12**, e0173974 (2017).

15.   Todd EV*, et al.* Stress, novel sex genes, and epigenetic reprogramming orchestrate socially
controlled sex change. *Sci Adv* **5**, eaaw7006 (2019).

16.   He Z*, et al.* Crosstalk between sex-related genes and apoptosis signaling reveals molecular
insights into sex change in a protogynous hermaphroditic teleost fish, ricefield eel *Monopterus*
albus. *Aquaculture* **552**, 737918 (2022).

17.   Fan M, Yang W, Zhang W, Zhang L. The ontogenic gonadal transcriptomes provide insights
into sex change in the ricefield eel *Monopterus albus*. *BMC Zool* **7**, 56 (2022).

18.   Wu X*, et al.* Integration of ATAC-seq and RNA-seq unravels chromatin accessibility during sex
reversal in Orange-spotted grouper (*Epinephelus coioides*). *Int J Mol Sci* **21**, 2800 (2020).

19.   Zhang Y, Zhang S, Liu Z, Zhang L, Zhang W. Epigenetic modifications during sex change
repress gonadotropin stimulation of cyp19a1a in a teleost ricefield eel (*Monopterus albus*).
Endocrinology **154**, 2881–2890 (2013).

20.   Li GL, Liu XC, Zhang Y, Lin HR. Gonadal development, aromatase activity and P450
aromatase gene expression during sex inversion of protogynous red-spotted grouper
Epinephelus akaara (Temminck and Schlegel) after implantation of the aromatase inhibitor,
fadrozole. *Aquac Res* **37**, 484–491 (2006).

21.   Wang DS*, et al.* Foxl2 up-regulates aromatase gene transcription in a female-specific manner
by binding to the promoter as well as interacting with ad4 binding protein/steroidogenic factor
1. *Mol Endocrinol* **21**, 712–725 (2007).

22.   Zhang C*, et al.* Transcription factors foxl2 and foxl3 regulate cyp19a1a and cyp11b in
orange-spotted grouper (*Epinephelus coioides*). *Aquac Rep* **25**, 101243 (2022).

23.   Shibata Y*, et al.* Expression of gonadal soma derived factor (GSDF) is spatially and temporally
correlated with early testicular differentiation in medaka. *Gene Expr Patterns* **10**, 283–289
(2010).

24.   Zhang X*, et al.* Autosomal gsdf acts as a male sex initiator in the fish medaka. *Sci Rep* **6**, 19738
19738 (2016).

25.   Kobayashi T, Kajiura-Kobayashi H, Guan G, Nagahama Y. Sexual dimorphic expression of
DMRT1 and Sox9a during gonadal differentiation and hormone-induced sex reversal in the
teleost fish Nile tilapia (*Oreochromis niloticus*). *Dev Dyn* **237**, 297–306 (2008).

26.   Rodríguez-Marí A, Yan Y-L, BreMiller RA, Wilson C, Cañestro C, Postlethwait JH.
Characterization and expression pattern of zebrafish anti-Müllerian hormone (amh) relative to
sox9a, sox9b, and cyp19a1a, during gonad development. *Gene Expr Patterns* **5**, 655–667
(2005).

27.   Wang XG, Orban L. Anti-Müllerian hormone and 11 β-hydroxylase show reciprocal expression
to that of aromatase in the transforming gonad of zebrafish males. *Dev Dyn* **236**, 1329–1338
(2007).

28.   Skaar KS, Nobrega RH, Magaraki A, Olsen LC, Schulz RW, Male R. Proteolytically activated,
recombinant anti-mullerian hormone inhibits androgen secretion, proliferation, and
differentiation of spermatogonia in adult zebrafish testis organ cultures. *Endocrinology* **152**,
3527–3540 (2011).

29.   Webster KA, Schach U, Ordaz A, Steinfeld JS, Draper BW, Siegfried KR. Dmrt1 is necessary
for male sexual development in zebrafish. *Dev Biol* **422**, 33–46 (2017).

30.   Zhu Y, Wang C, Chen X, Guan G. Identification of gonadal soma-derived factor involvement in
Monopterus albus (protogynous rice field eel) sex change. *Mol Biol Rep* **43**, 629–637 (2016).

31.   Bono H. Meta-analysis of oxidative transcriptomes in insects. *Antioxidants* **10**, 345 (2021).

32.   Ono Y, Bono H. Multi-omic meta-analysis of transcriptomes and the bibliome uncovers novel
hypoxia-inducible genes. *Biomedicines* **9**, 582 (2021).

33.   Toga K, Bono H. Meta-analysis of public RNA sequencing data revealed potential key genes
associated with reproductive division of labor in social hymenoptera and termites. *Int J Mol Sci*
24, 8353 (2023).

34.   Rahman MR*, et al.* Comprehensive analysis of RNA-Seq gene expression profiling of brain
transcriptomes reveals novel genes, regulators, and pathways in autism spectrum disorder.
Brain Sci **10**, 747 (2020).

35.   Shaar-Moshe L, Hübner S, Peleg Z. Identification of conserved drought-adaptive genes using a
cross-species meta-analysis approach. *BMC Plant Biol* **15**, 111 (2015).

36.   Cheng H, Concepcion GT, Feng X, Zhang H, Li H. Haplotype-resolved de novo assembly
using phased assembly graphs with hifiasm. *Nat Methods* **18**, 170–175 (2021).

37.   Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM, Kelley J. BUSCO update: novel
and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of
eukaryotic, prokaryotic, and viral genomes. *Mol Biol Evol* **38**, 4647–4654 (2021).

38.   Haas B. TransDecoder (v5.5.0). https://github.com/TransDecoder/TransDecoder

39.   Ge H*, et al.* De novo assembly of a chromosome-level reference genome of red-spotted grouper
(*Epinephelus akaara*) using nanopore sequencing and Hi-C. *Mol Ecol Resour* **19**, 1461–1469
(2019).

40.   Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. StringTie enables
improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol* **33**, 290–295
(2015).

41.   Grabherr MG*, et al.* Full-length transcriptome assembly from RNA-Seq data without a
reference genome. *Nat Biotechnol* **29**, 644–652 (2011).

42.   Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-aware
quantification of transcript expression. *Nat Methods* **14**, 417–419 (2017).

43.   Pearson WR, Lipman DJ. Improved tools for biological sequence comparison. *Proc Natl Acad*
Sci USA **85**, 2444–2448 (1988).

44.   Cunningham F*, et al.* Ensembl 2022. *Nucleic Acids Res* **50**, D988–D995 (2022).

45.   Nozu R, Kojima Y, Nakamura M. Short term treatment with aromatase inhibitor induces sex

change in the protogynous wrasse, *Halichoeres trimaculatus*. *Gen Comp Endocrinol* **161**,
360–364 (2009).
590

591 46. Kroon FJ, Liley NR. The role of steroid hormones in protogynous sex change in the Blackeye
592   goby, *Coryphopterus nicholsii* (Teleostei: Gobiidae). *Gen Comp Endocrinol* **118**, 273–283
593   (2000).
594

595 47. Yan Y-L*, et al.* A hormone that lost its receptor: anti-müllerian hormone (amh) in zebrafish
596   gonad development and sex determination. *Genetics* **213**, 529–553 (2019).
597

598 48. Peng C*, et al.* Natural sex change in mature protogynous orange☐spotted grouper (*Epinephelus
599   coioides*): gonadal restructuring, sex hormone shifts and gene profiles. *J Fish Biol* **97**, 785‒793
600   (2020).
601

602 49. Nozu R, Horiguchi R, Murata R, Kobayashi Y, Nakamura M. Survival of ovarian somatic cells
603   during sex change in the protogynous wrasse, *Halichoeres trimaculatus*. *Fish Physiol Biochem*
604   **39**, 47–51 (2013).
605

606 50. Nemtzov SC. Social control of sex change in the Red Sea razorfish *Xyrichtys pentadactylus*
607   (Teleostei, Labridae). *Environ Biol Fishes* **14**, 199–211 (1985).
608

609 51. Hasegawa S, Noda K, Maeda A, Matsuoka M, Yamasaki M, Fukui T. Acetoacetyl-CoA
610   synthetase, a ketone body-utilizing enzyme, is controlled by SREBP-2 and affects serum
611   cholesterol levels. *Mol Genet Metab* **107**, 553–560 (2012).
612

613 52. Wu D-M, He Z, Chen T, Liu Y, Ma L-P, Ping J. DNA hypermethylation of acetoacetyl-CoA
614   synthetase contributes to inhibited cholesterol supply and steroidogenesis in fetal rat adrenals
615   under prenatal nicotine exposure. *Toxicology* **340**, 43–52 (2016).
616

617 53. Gaudet P, Livstone MS, Lewis SE, Thomas PD. Phylogenetic-based propagation of functional
618   annotations within the Gene Ontology consortium. *Brief Bioinform* **12**, 449‒462 (2011).
619

620 54. Payne AH, Hales DB. Overview of steroidogenic enzymes in the pathway from cholesterol to
621   active steroid hormones. *Endocr Rev* **25**, 947–970 (2004).
622

623 55. Pikuleva IA. Cholesterol-metabolizing cytochromes P450. *Drug Metab Dispos* **34**, 513–520
624   (2006).
625

626 56. Miller WL. Steroidogenic enzymes. In: *Disorders of the Human Adrenal Cortex* (eds Flück CE,
627   Miller WL). S.Karger AG (2008).

57. Sakae Y*, et al.* Starvation causes female-to-male sex reversal through lipid metabolism in the teleost fish, medaka (*Olyzias latipes*). *Biol Open* **9**, bio050054 (2020).

58. Gwynne JT, Strauss JF. The role of lipoproteins in steroidogenesis and cholesterol metabolism in steroidogenic glands. *Endocr Rev* **3**, 299–329 (1982).

59. Sharpe RL, Drolet M, MacLatchy DL. Investigation of de novo cholesterol synthetic capacity in the gonads of goldfish (*Carassius auratus*) exposed to the phytosterol beta-sitosterol. *Reprod Biol Endocrinol* **4**, 60 (2006).

60. Bera A, Chadha NK, Dasgupta S, Chakravarty S, Sawant PB. Hypoxia-mediated inhibition of cholesterol synthesis leads to disruption of nocturnal sex steroidogenesis in the gonad of koi carp, *Cyprinus carpio*. *Fish Physiol Biochem* **46**, 2421–2435 (2020).

61. Sim SB, Corpuz RL, Simmonds TJ, Geib SM. HiFiAdapterFilt, a memory efficient read processing pipeline, prevents occurrence of adapter sequence in PacBio HiFi reads and their negative impacts on genome assembly. *BMC Genomics* **23**, 157 (2022).

62. Zhu B-H, Xiao J, Xue W, Xu G-C, Sun M-Y, Li J-T. P_RNA_scaffolder: a fast and accurate genome scaffolder using paired-end RNA-sequencing reads. *BMC Genomics* **19**, 175 (2018).

63. Kadota M, Nishimura O, Miura H, Tanaka K, Hiratani I, Kuraku S. Multifaceted Hi-C benchmarking: what makes a difference in chromosome-scale genome scaffolding? *Gigascience* **9**, giz158 (2020).

64. Durand NC*, et al.* Juicer provides a one-click system for analyzing loop-resolution Hi-C experiments. *Cell Syst* **3**, 95–98 (2016).

65. Dudchenko O*, et al.* The Juicebox assembly tools module facilitates *de novo* assembly of mammalian genomes with chromosome-length scaffolds for under $1000. *bioRxiv*, 254797 (2018).

66. Nishimura O, Hara Y, Kuraku S, Hancock J. gVolante for standardizing completeness assessment of genome and transcriptome assemblies. *Bioinformatics* **33**, 3635–3637 (2017).

67. Kodama Y, Shumway M, Leinonen R. The sequence read archive: explosive growth of sequencing data. *Nucleic Acids Res* **40**, D54–D56 (2011).

68. Trim Galore. https://github.com/FelixKrueger/TrimGalore

669   69.   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and
670   genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907–915 (2019).

672   70.   Danecek P*, et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008 (2021).

674   71.   Leek JT*, et al.* Tackling the widespread and critical impact of batch effects in high-throughput
675   data. *Nat Rev Genet* **11**, 733–739 (2010).

677   72.   Zou Q, Shen W, Le S, Li Y, Hu F. SeqKit: a cross-platform and ultrafast toolkit for FASTA/Q
678   file manipulation. *PloS one* **11**, e0163962 (2016).

680   73.   R Core Team. R: A language and environment for statistical computing. URL
681   https://www.R-project.org/

683   74.   Pertea G, Pertea M. GFF Utilities: GffRead and GffCompare. *F1000Res* **9**, 304 (2020).

685   75.   Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. *Nat
686   Methods* **12**, 59–60 (2014).

688   76.   Zhou Y*, et al.* Metascape provides a biologist-oriented resource for the analysis of
689   systems-level datasets. *Nat Commun* **10**, 1523 (2019).

Acknowledgements

All authors thank Dr. Osamu Nishimura at DNA Sequencing Facility operated by Laboratory

for Phyloinformatics in RIKEN BDR for assistance in Hi-C data assessment. We thank the

Sesoko Station, Tropical Biosphere Research Center, University of the Ryukyus for allowing

us to use their research facility. This research was supported by the Center of Innovation for

Bio-Digital Transformation (BioDX), a program on the open innovation platform for

industry-academia co-creation (COI-NEXT), and the Japan Science and Technology Agency

(JST, COI-NEXT, JPMJPF2010). This work was partially received financial support by

International Research Organization for Advanced Science and Technology, Kumamoto

University, Kumamoto, Japan.

Author contributions

RN and HB conceived and designed the study. MK and SK performed genome sequence and

de novo genome assembly of three-spot wrasse. RN & MN RNA-seq data production of

three-spot wrasse. RN performed RNA-seq data analysis and orthology inference. RN, MN,

SK and HB wrote a draft manuscript. All authors read and approved the final manuscript.

Competing interests

The authors have no conflicts of competing interest to declare that are relevant to the content

of this article.

Additional information

All supplementary information is accessible in figshare repository

(https://doi.org/10.6084/m9.figshare.c.6704160)

Supplementary figure 1. Histological sections of control ovaries and transitional gonads treated

AI for 7 days in the three-spot wrasse, *Halichoeres trimaculatus*.

Supplementary table 1

Supplementary table 2

Supplementary table 3

Supplementary table 4

Figure legends

Fig. 1. Schematic diagram of the analysis pipeline in this study.

Fig. 2 Scatter plot of the IO-score and gene set enrichment analysis of up- and

downregulated genes. a Scatterplot of total IO-scores for the 7,289 factors in descending

order. Blue circles indicate 99 upregulated factors; red circles indicate 69 downregulated

factors. Gray circles (which have the appearance of lines because they overlap each other)

indicate unchanged factors ($6 \leq$ total IO-score ≤ -6). **b,c** Results of enrichment analysis for 99

upregulated (**b**) and 69 downregulated (**c**) factors

Fig. 3 Composition of total IO-scores for upregulated factors (n=99). Numbers in

parentheses indicate the number of RNA-Seq data pairs for each species used in this study.

Asterisk indicates a gene for which the total IO-score is derived from a single species.

Fig. 4 Composition of total IO-scores for downregulated factors (n=69). Numbers in

parentheses indicate the number of RNA-Seq data pairs for each species used in this study.

Asterisk indicates a gene for which the total IO-score is derived from a single species.

Table 1. Statistics of genome assembly in Three-spot wrasse (fHalTri1.1) and other Labidae

fishes listed in NCBI Genomes.

	fHalTri1.1	*Cheilinus undulatus* (humphead wrasse)	*Notolabrus celidotus* (New Zealand spotty)	*Labrus bergylta* (ballan wrasse)	*Semicossyphus pulcher*	*Tautogolabrus adspersus* (cunner)
Assembly accession	BSYP01000001–BSYP01000227	GCF_018320785.1	GCF_009762535.1	GCF_900080235.1	GCA_022749685.1	GCA_020745685.1
Total seq length (Mb)	849.9	1173.4	846.7	805.5	794.1	723.6
Max length (Mb)	44.2	59.6	42.4	9.1	38.5	37.1
Num of scaffolds	227	45	467	13,466	178	347
Scaffold N50 (Mb)	37.9	51.5	37.1	0.8	32.1	31.7
Scaffold L50	11	11	11	252	12	11
Num of seqs > 1 Mb	25	24	24	197	35	24
% of seqs > 1 Mb	97.3	99.7	97.6	43.9	97.1	99.9

Num of gaps	117	284	1,133	257	8	283
Complete BUSCO (%)	98.7	99.0	97.1	98.9	98.9	98.1

755 **Table 2.** Statistics of orthology relationship among species.

Common name	Scientific name	Order	Reference genome	*Num of transcripts seqs obtained based on the RNA-seq data from gonadal tissue	Num of protein seqs predicted by TransDecoder	Num of RBH (vs Three-spot wrasse)
Three-spot wrasse	*Halichoeres trimaculatus*	Labriformes	Assembly in this study	27,573	24,965	24,965
Bluehead	*Thalassoma bifasciatum*	Labriformes	GCA_008086565.1	57,699	43,195	13,871
Hong Kong grouper	*Epinephelus akaara*	Perciformes	Ge et al., 2019	26,507	24,752	11,619
Orange-spotted grouper	*Epinephelus coioides*	Perciformes	GCA_900536245.1	30,867	34,191	12,507
Black seabass	*Centropristis striata*	Perciformes	NA	** 95,941	95,941	12,591
Asian swamp eel	*Monopterus albus*	Synbranchiformes	GCF_001952655.1	53,472	50,693	12,765
Zig-zag eel	*Mastacembelus armatus*	Synbranchiformes	GCF_900324485.2	48,724	40,908	13,095

756 * Based on the annotation file (merged gtf file), the extracted transcripts sequences from the

757 genome by gffread

758 ** Among transcriptome sequences (594,726 seqs) assembled by Trinity, transcripts sequences

759 corresponding to coding seqences predicted via TransDecoder

773 **Table 3.** List of genes that contributed to GO term enrichment.

Up/Down	GO ID	Category	GO term	*Genes
Up	GO:0007129	Biological Process	homologous chromosome pairing at meiosis	*hormad1, terb2, si:ch211-199g17.9*
Up	GO:0007369	Biological Process	gastrulation	*zeb1b, tmem237a, col12a1b, pfn1, enpp2*
Up	GO:1902531	Biological Process	regulation of intracellular signal transduction	*cep55l, psd3l, arhgef18b, ccl19a.1, sh3bp1, enpp2, pik3r6b*
Up	GO:0005929	Cellular Component	cilium	*septin6, prom1a, tmem237a, si:ch211-66i15.4, flr*
Up	GO:0007281	Biological Process	germ cell development	*hormad1, gsdf, sycp3*
Down	GO:0010817	Biological Process	regulation of hormone levels	*cyp19a1a, aacs, hsd17b1, ece1*
Down	GO:0006413	Biological Process	translational initiation	*eif4g2b, eif4eb, eif4e1c*
Down	GO:0019752	Biological Process	carboxylic acid metabolic process	*lpl, ldha, aacs, amacr, acaca, dars2*
Down	GO:0008202	Biological Process	steroid metabolic process	*hsd17b1, amacr, sdr42e1*
Down	GO:1904888	Biological Process	cranial skeletal system development	*nog3, cep55l, cyldb, ece1*
Down	GO:1901362	Biological Process	organic cyclic compound biosynthetic process	*tbx5a, hsd17b1, alad, acaca, sdr42e1, adcy6a*

774 * Gene symbols follow the zebrafish gene nomenclature in Ensembl.

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4.