Meta-analysis of gonadal transcriptome provides novel insights into sex change mechanism

across protogynous fishes

Short title: Meta-analysis of transcriptome in fishes

Keywords: Meta-analysis, Protogynous fishes, Gonad, Sex change, Genome, Transcriptome

Ryo Nozu[1,2], Mitsutaka Kadota[3,4], Masaru Nakamura[5], Shigehiro Kuraku[6,7], Hidemasa Bono[1,2*]

[1] Laboratory of Genome Informatics, Graduate School of Integrated Sciences for Life,

Hiroshima University, 3-10-23 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-0046, Japan

[2] Laboratory of BioDX, Genome Editing Innovation Center, Hiroshima University, Hiroshima,

Japan

[3] Laboratory for Phyloinformatics, RIKEN Center for Biosystems Dynamics Research (BDR),

Kobe, Japan

[4] Laboratory for Developmental Genome System, RIKEN Center for Biosystems Dynamics

Research (BDR), Kobe, Japan

[5] Okinawa Churashima Research Center, Okinawa Churashima Foundation, Motobu-cho,

Okinawa, Japan

[6] Molecular Life History Laboratory, Department of Genomics and Evolutionary Biology,

National Institute of Genetics, Mishima, Japan

[7] Department of Genetics, Sokendai (Graduate University for Advanced Studies), Mishima,

Japan.

[*]To whom correspondence may be addressed: Hidemasa Bono

bonohu@hiroshima-u.ac.jp

Abstract

Protogyny, being capable of changing from female to male during their lifetime, is prevalent in 20 families of teleosts but is believed to have evolved within specific evolutionary lineages. Therefore, shared regulatory factors governing the sex change process are expected to be conserved across protogynous fishes. However, a comprehensive understanding of this mechanism remains elusive. To identify these factors, we conducted a meta-analysis using gonadal transcriptome data from seven species. We curated data pairs of ovarian tissue and transitional gonad, and employed ratios of expression level as a unified criterion for differential expression, enabling a meta-analysis across species. Our approach revealed that classical sex change-related genes exhibited differential expression levels between the ovary and transitional gonads, consistent with previous reports. These results validate our methodology's robustness. Additionally, we identified novel genes not previously linked to gonadal sex change in fish. Notably, changes in the expression levels of acetoacetyl-CoA synthetase and apolipoprotein Eb, which are involved in cholesterol synthesis and transport, respectively, suggest that the levels of cholesterol, a precursor of steroid hormones crucial for sex change, are decreased upon sex change onset in the gonads. This implies a potential universal influence of cholesterol dynamics on gonadal transformation in protogyny.

Introduction

Sequential hermaphroditism, a phenomenon in which individuals undergo sex change during their lifetime, has been extensively recognized in teleosts over the past several decades (Atz, 1964). Sequential hermaphroditism has been categorized into three major types based on the direction of sex change (protogyny, where females change to males; protandry, where males change to females; and bidirectional sex change, where individuals can change both sexes serially). Phylogenetic analysis has shown that sequential hermaphroditism has been acquired in various evolutionary lineages (Mank et al., 2006) (Fig. 1). Interestingly, protogyny is prevalent in 20 families but is thought to have emerged once in evolutionally the Percomorpha (Kuwamura et al., 2020). Therefore, there are common features in the intricate regulatory mechanisms during protogynous sex change. Specifically, the replacement of ovarian tissue with testicular tissue is commonly observed during protogynous sex change (Bhandari et al., 2003; Lo Nostro et al., 2003; Muncaster et al., 2013; Nakamura et al., 1989; Ohta et al., 2008). Additionally, considering the high functional conservation of gonads across organisms, the existence of a regulatory mechanism governing gonadal transformation comprising common and essential factors across protogynous fishes is plausible; however, a comprehensive perspective of this process remains elusive.

Accordingly, elucidation of the molecular mechanisms underlying the gametogenesis switch in sex-changing fish has subsequently become a focus of research. In particular, the expression patterns of multiple sex-related genes during sex change within the family Labridae, which serves as a model for protogynous sex change, have been explored (Horiguchi et al., 2018; Miyake et al., 2012; Thomas et al., 2019). Recently, next-generation sequencing technology has enabled researchers to conduct comprehensive analyses of gene expression, allowing the identification of global gene expression profiles during gonadal transformation in protogynous fish belonging to Labriformes, Perciformes, and Synbranchiformes (Fan et al., 2022; He et al., 2022; Senthilkumaran et al., 2017; Todd et al., 2019; Wu et al., 2020). Thus,

these results suggest the involvement of several pivotal genes in the ovary-to-testis

transformation in sex-changing fish.

For example, a repertoire of sex-related genes, including cytochrome P450, family 19,

subfamily A, polypeptide 1a (*cyp19a1a*), forkhead box L2a (*foxl2a*), folliculogenesis specific

bHLH transcription factor (*figla*), anti-Müllerian hormone (*amh*), gonadal somatic cell derived

factor (*gsdf*), doublesex and mab-3 related transcription factor 1 (*dmrt1*) were identified as

differentially expressed genes (DEGs) during sex change across protogynous fishes. Notably,

the gonadal expression of *cyp19a1a* and *foxl2a*, which are key players in maintaining ovarian

status, is reduced upon the onset of sex change in multiple species (Li et al., 2006; Thomas et

al., 2019; Zhang et al., 2013). *cyp19a1a* encodes aromatase, an enzyme responsible for the

conversion of androgens (e.g., testosterone) to estrogens (e.g., estradiol-17ß; E2), whereas

foxl2a encodes a transcription factor that targets and up-regulates *cyp19a1a* in teleost fishes

(Wang et al., 2007; Zhang et al., 2022). Furthermore, the initiation of female-to-male sex

change, triggered by a rapid decline in E2 levels (Nakamura et al., 1989), is well-coordinated

with the expression profiles of these genes. Conversely, *amh*, *gsdf*, and *dmrt1* are crucial for

testicular differentiation and development in teleost fishes (Kobayashi et al., 2008;

Rodríguez-Marí et al., 2005; Shibata et al., 2010; Skaar et al., 2011; Wang & Orban, 2007;

Webster et al., 2017; Zhang et al., 2016). In several sex-changing species, the upregulation of

these genes coincides with the emergence of testicular tissue during transition, suggesting that

these genes are involved in testicular formation during gonadal sex change (Horiguchi et al.,

2013; Horiguchi et al., 2018; Nozu et al., 2015; Thomas et al., 2019; Zhu et al., 2016).

Transcriptome analysis during gonadal transformation has also revealed differential

expression of several genes not previously implicated in sex determination or differentiation

pathways. In particular, upregulation of apolipoprotein Eb (*apoeb*), ectonucleotide

pyrophosphatase/phosphodiesterase 2 (*enpp2*), and l-rhamnose-binding lectin CSL2 (*csl2*)

expression, accompanied by downregulation of galectin-3 (*gal3*), was observed during the

early phase of gonadal sex change in the protogynous Asian swamp eel *Monopterus albus*.

These findings suggest that these genes are involved in gonadal transformation, particularly

during the initiation of sex change (Fan et al., 2022; Senthilkumaran et al., 2017). However,

verifying their direct involvement and commonality in the sex change process remains

challenging.

Recently, by amalgamating transcriptome data from multiple studies, meta-analysis has

attracted attention as a powerful tool capable of providing novel insights overlooked by

traditional hypothesis-driven research methods (Bono, 2021; Rahman et al., 2020; Toga &

Bono, 2023). In other words, it is expected that by reanalyzing studies across multiple species

and studies, universal factors controlling gonadal transformation, which have not been

detected in individual studies, could be identified among protogynous fishes. Therefore, the

objective of the present study was to identify the shared factors involved in gonadal

transformation across species using a meta-analysis of publicly available transcriptome data

from multiple experiments and species. Furthermore, in this study, we focused on another

well-studied species, the three-spot wrasse, *Halichoeres trimaculatus*, a protogynous species

belonging to the family Labridae and produced *de novo* genome assembly and gonad

transcriptome data undergoing sex change. This new data consolidated our dataset

encompassing seven protogynous species from three orders allowing exhaustive

transcriptome analysis. Through this dataset and uniform pipeline, we identified genes with

differential expression levels between the transitional gonads and ovaries (Fig. S1). We

reevaluated the relationship between these genes and gonadal sex change, providing a novel

perspective regarding the mechanisms underlying this process.

Results

***De novo* genome assembly and transcriptome of gonadal tissue in the three-spot wrasse.**

The genome sequencing was required for the RNA-Seq analysis in this study, so the *de novo*

genome assembly of the three-spot wrasse, which had not been previously determined, was

constructed. High-throughput genome sequence data were obtained from a locally obtained

female three-spot wrasse from Okinawa, Japan. The obtained HiFi data, which comprised

approximately 32 Gbp and encompassed approximately 1.9 million DNA sequences, were

assembled using the hifiasm program (Cheng et al., 2021). Subsequently, the acquired Hi-C

data were used for scaffolding, resulting in the generation of 227 scaffolds. The longest

scaffold measured 44.2 Mbp, with an N50 length (the shortest of the lengths covering 50% of

the genome) of 37.9 Mbp, contributing to a total assembly length of approximately 849.9 Mbp.

Assessment using BUSCO v5.4 (Manni et al., 2021) revealed a completeness > 98%,

indicating a high level of coverage and contiguity in the *de novo* genome assembly of the

three-spot wrasse (Table 1).

The RNA-Seq reads obtained from transitional gonads and ovaries of additional females

were mapped to the genome assembly, and the obtained mapping results were used to assemble

the transcriptome for the annotation file ("merged.gtf"). This process yielded 27,573 transcript

sequences (Table 2). Subsequently, 24,695 coding sequences (protein sequences) were

predicted from the set of transcript sequences using TransDecoder (Haas, 2018).

Data curation of public RNA-Seq data and genomic information. We downloaded

Sequence Read Archive (SRA) data from six fish species for which sex changes have been

reported (Kuwamura et al., 2020): bluehead wrasse *Thalassoma bifasciatum* (Todd et al.,

2019) (order Labriformes), orange-spotted grouper *Epinephelus coioides* (Xiao et al., 2020)

(order Perciformes), Hong Kong grouper *E. akaara* (Ge et al., 2019) (order Perciformes)*,* black

seabass *Centropristis striata* (Breton et al., 2019) (order Perciformes); Asian swamp eel

Monopterus albus (Fan et al., 2022; Senthilkumaran et al., 2017; Zhao et al., 2018) (order

Synbranchiformes), zig-zag eel *Mastacembelus armatus* (Xue et al., 2021) (order

Synbranchiformes). Furthermore, using the metadata associated with the SRA data, we curated

the RNA-Seq data into comparable sample pairs of intersexual (transitional) gonads and

ovaries (a total of 31 paired data, including three-spot wrasse). Table S1 presents the metadata,

including Run accession numbers, for the RNA-Seq data analyzed in this study.

The genome sequences of five of the seven species utilized in this study are publicly

available (Table 2) and served as reference genomes for read mapping. The genome sequences

of *T. bifasciatum* (GCA_008086565.1), *E. coioides* (GCA_900536245.1), *M. albus*

(GCF_001952655.1), and *M. armatus* (GCF_900324485.2) were retrieved from the National

Center for Biotechnology Information (NCBI) Genome Database (accessed in October 2022).

We also referenced the genome information of *E. akaara* reported by Ge et al., 2019.

Transcriptome assembly and inference of ortholog relationships. For the five species with

public reference genomes, the number of transcript sequences obtained using StringTie (Pertea

et al., 2015) with the "--merge" option is shown in Table 2. For black sea bass, (without a

reference genome), the number of contiguous sequences assembled using Trinity (Grabherr et

al., 2011) was 594,726. Among these, 95,941 transcript sequences corresponding to the

predicted coding sequences (CDS) were specified as references for expression quantification

using Salmon (Patro et al., 2017). The number of protein sequences predicted using

TransDecoder is shown in Table 2. Orthology inference and consolidation of orthologs within

the three-spot wrasse identified 7,289 orthologs that were common to all species used in this

study (Table 2), and thus were considered as candidate sex-change genes.

Identification of differentially expressed genes. The total intersexual gonad and ovary score

(IO-score) was calculated for these 7,289 genes (Fig. 2a, Table S2, IO-score calculation; see

Methods). Among these, 99 genes were upregulated (Fig. 2b, Table S3) and 69 genes were

downregulated (Fig. 2c, Table S4). Whereas the total IO-score of these genes was occasionally

derived from a single species (up: *cep97*, *zmat2*, *tmem237a*, *supt5h*; down: *znf16l*, *lrrc8db*,

sh2b1, *tmem263*, *cyldb*, *manea*, *nol11*), in most cases, the total IO-scores were based on

contributions from two or more species. Notably, the upregulated genes included *amh* and *gsdf*,

which have been previously reported to play important roles in gonadal sex change,

particularly in testicular formation and development, in some protogynous fish. Conversely,

the downregulated genes included *cyp19a1a*, *hsd17b1*, and *foxl2a*, which are crucial for the

onset of sex change (Table 3).

Enrichment analysis of differentially expressed genes. Sequence homology search of the

three-spot wrasse proteins using ggsearch36 (Pearson & Lipman, 1988) against the reference

zebrafish proteins (Ensembl Release 108) (Cunningham et al., 2022) resulted in the

identification of 90 homologs among the 99 upregulated genes (Table S3). No corresponding

zebrafish orthologs were found for the remaining nine genes that were common to all species

analyzed in this study. The results of enrichment analysis using the 90 genes that could be

converted to zebrafish orthologs are shown in Fig. 3a. The Gene Ontology (GO) terms

"homologous chromosome pairing at meiosis" and "germ cell development," which are related

to reproduction, were significantly enriched among the upregulated genes. The specific genes

annotated with these GO terms included *hormad1*, *terb2*, *si:ch211-199g17.9*, *gsdf*, and *sycp3*

(Table 3).

We successfully obtained the corresponding zebrafish homologs for all of the 69

downregulated genes. The results of enrichment analysis are shown in Fig. 3b. Among the

downregulated genes, GO terms related to the regulation of steroid hormone levels, including

"regulation of hormone levels" and "steroid metabolic process," were significantly enriched.

Specific genes annotated with these enriched GO terms included *cyp19a1a*, *aacs*, *hsd17b1*,

ece1, *amacr*, and *sdr42e1* (Table 3).

Pathways mapping of candidate sex-change genes. Visualization of cholesterol and steroid

hormone biosynthesis pathways was conducted based on the GO terms obtained from the

enrichment analysis. By referencing pathway information for teleost fishes, pathway diagrams

were constructed and loaded with 7,289 genes converted to Ensembl zebrafish gene IDs (see

Methods). The results showed the mapping of most enzymes involved in the pathway from

acetyl-CoA to cholesterol synthesis (Fig. 4). Similarly, when mapping the steroid hormone

synthesis pathway, it was confirmed that enzymes involved in the synthesis pathway from

cholesterol to estradiol were included among the candidate sex-change genes (Fig. S2).

Discussion

In this study, we conducted a meta-analysis of gonadal RNA-Seq data from seven protogynous

fish species using a unified analysis pipeline and threshold. The aim was to compare the

transcriptomes of ovaries and sexual transition gonads, with the goal of identifying common

genes involved in protogynous gonadal sex change. Toward this end, we first assembled the *de*

novo genome of the three-spot wrasse. The assembly statistics of the three-spot wrasse (Table

1) were comparable to those of other Labridae species, such as humphead wrasse (*Cheilinus*

undulatus), ballan wrasse (*Labrus bergylta*), and New Zealand spotty (*Notolabrus celidotus*),

previously deposited in the NCBI RefSeq database. These results indicated that the *de novo*

genome assembly of the three-spot wrasse exhibited high quality, enabling us to perform

expression analysis and orthology inference with confidence.

In our meta-analysis, we broadly examined the gonadal sex change process in protogynous

fish, focusing on the replacement of ovarian tissue with testicular tissue. At a more detailed

level, histological characteristics during the sex change process vary among species. In

Labriformes, the early stages of sex change are predominantly characterized by ovarian tissue

regression (Nozu et al., 2009; Todd et al., 2019). In contrast, in Perciformes and

Synbranchiformes, testicular tissue emerges, and both ovarian and testicular tissues are present

simultaneously (Liu & de Mitcheson, 2009; Senthilkumaran et al., 2017; Xue et al., 2021).

Additionally, the methods used to obtain transitional gonad samples for RNA-seq also varied

in each project (Table S1). These differences are expected to impact gene expression levels. In

this context, we consider the differentially expressed genes identified through the

meta-analysis as crucial factors involved in gonadal transformation across species.

Through this meta-analysis, we successfully identified sex-related genes including

cyp19a1a, *foxl2a*, *hsd17b1*, *amh*, and *gsdf*, that had been previously reported to play a role in

sex change in various protogynous fish species (Fan et al., 2022; Horiguchi et al., 2013;

Thomas et al., 2019; Todd et al., 2019). Furthermore, our enrichment analysis of

downregulated genes highlighted a term related to the "regulation of steroid hormone

production." This finding connects to those of previous reports, which indicate that a rapid

decrease in estrogen levels is crucial for female-to-male sex change (Kroon & Liley, 2000;

Nakamura et al., 1989; Nozu et al., 2009). Thus, by pooling data across species, we were able

to identify common genes involved in gonadal sex change. Notably, our meta-analysis

demonstrates the potential of utilizing transcriptome data from multiple experiments to gain

insights regarding the mechanisms underlying sex change in fish.

On the other hand, some sex-related genes, generally identified as related to sex change,

showed no differential expression in this study, such as *figla*, *dmrt1*, *csl2*, and *gal3*. Particularly,

dmrt1, *csl2*, and *gal3* were not included in the set of 7289 orthologs common to all seven

species (Table S2). Among these genes, *dmrt1* is known as a key gene in male differentiation

(Herpin & Schartl, 2011), and its expression profiles have also been investigated in

sex-changing fish. In some protogynous fish, *dmrt1* expression increases during sex change,

coinciding with the appearance of testicular tissue (Nozu et al., 2015; Todd et al., 2019). Since

our dataset targets the early stages of sex change in gonads, it is suggested that the expression

levels of *dmrt1* were too low for the *dmrt1* transcripts to be assembled.

254   Our findings demonstrate that *aacs*, annotated with the GO term "regulation of hormone

255  levels," is downregulated during sex change. Notably, no previous reports have suggested the

256  involvement of *aacs* in sex change, indicating that the encoded protein (acetoacetyl-CoA

257  synthetase; Aacs) may represent a novel gene associated with this process. Typically, AACS

258  plays a critical role in cholesterol supply. Studies in mice have shown that *Aacs* knockdown

259  decreases total serum cholesterol levels (Hasegawa et al., 2012). Additionally, in rat adrenal

260  glands, inhibition of *AACS* expression through hypermethylation suppresses cholesterol supply

261  and steroid synthesis (Wu et al., 2016). Moreover, apolipoprotein Eb (encoded by *apoeb*, listed

262  among the upregulated genes), has been suggested to participate in cholesterol transportation in

263  zebrafish (Gaudet et al., 2011). Apolipoprotein E, encoded by *APOE* which is the human

264  ortholog of *apoeb*, serves as a crucial lipid transport protein involved in the modulation of

265  cholesterol metabolism and is also synthesized within peripheral tissues (Blue et al., 1983). In

266  addition, a previous study indicates that APOE gene expression exerts suppressive effects on

267  steroid production in adrenal cells (Reyland et al., 1991). Specifically, the high expression of

268  APOE in tissues with active steroid production, including the adrenal glands, appears to shift

269  cholesterol metabolism from delivery to the steroidogenic pathway to storage of esterified

270  cholesterol, resulting in a decrease in steroid production levels (Thorngate et al., 2002). These

271  observations strongly support the hypothesis that the downregulation of *aacs* and the

272  upregulation of *apoeb* during sex change contribute to suppression of the cholesterol supply

273  and entire steroidogenesis.

274   Cholesterol serves as a precursor for steroid hormones, including estrogens, androgens, and

275  glucocorticoids (Miller, 2008; Payne & Hales, 2004; Pikuleva, 2006), which play crucial roles

276  in protogynous sex change. Previous studies have shown that a decline in endogenous

277  estrogens, particularly E2, triggers the transition from ovary to testis in protogynous fish

278  (Kroon & Liley, 2000; Nakamura et al., 1989; Nozu et al., 2009). E2 is synthesized from

279  testosterone via an aromatase enzyme, encoded by *cyp19a1a*. Downregulation of *cyp19a1a*

expression at the onset of sex change leads to reduced E2 production. Studies on *M. albus* and

T. bifasciatum have revealed higher DNA methylation levels of the *cyp19a1a* gene upstream

region in the testes than those in the ovaries, suggesting that epigenetic modifications may

contribute to the suppression of *cyp19a1a* expression during gonadal sex change (Todd et al.,

2019; Zhang et al., 2013). Our present findings could contribute one more "piece of the

puzzle", suggesting that the trigger of gonadal transformation may be attributed not only to the

decreased E2 production by the downregulation of aromatase activity but also to the

suppression of the cholesterol supply, the precursor of steroid hormones, leading to the

inhibition of the entire steroid biosynthesis.

In fish, it is suggested that synthesized cholesterol affects the production of steroid

hormones (Bera et al., 2020; Sharpe et al., 2006). This is also supported by reports that the

exposure HMG-CoA reductase inhibitors in zebrafish leads to a decrease in steroid hormone

levels (Al-Habsi et al., 2016). The cholesterol synthesis pathway (mevalonate pathway)

primarily uses acetyl-CoA as a substrate, and *aacs* might be involved in the supply of this

acetyl-CoA (Fig. 4). Additionally, a decrease in the expression of lipoprotein lipase (*lpl*) has

been observed (Fig. 4, Table S4). Lipoprotein lipase is involved in the biosynthetic process of

fatty acid which serves as one of the sources of acetyl-CoA. This suggests the possibility of a

concurrent suppression of fatty acid supply to the gonads. Both results could lead to support

the hypothesis that the decrease in steroid hormone production is due to a reduced supply of

cholesterol caused by inhibition of cholesterol synthesis. Although this hypothesis requires

further investigation and an understanding of cholesterol dynamics during sex change,

additional exploration is warranted. Notably, a study on medaka (*Oryzias latipes*) subjected to

starvation conditions implied a potential link between masculinization and lower cholesterol

levels (Sakae et al., 2020). Specifically, female-to-male sex reversal was observed in starved

medaka larvae, a gonochoristic fish, together with decreased cholesterol levels in the starved

samples. However, the precise pathways underlying masculinization and cholesterol dynamics

remain unknown. In general, cholesterol is synthesized not only in the liver and adrenal glands

but also in the gonads (Bera et al., 2020; Gwynne & Strauss, 1982; Sharpe et al., 2006).

Therefore, gaining a comprehensive understanding of the lipid metabolic mechanisms in fish

gonads, including cholesterol synthesis and transportation during sex change, may provide

valuable insights regarding the mechanisms underlying gonadal transformation. To clarify this

perspective, the remaining challenge lies in providing experimental evidence regarding

whether it is possible to control gonadal transformation by actually regulating cholesterol

levels within the body.

Conclusions

In this study, our objective was to compare the transcriptomes of sex-transitional gonads with

those of ovaries using a unified pipeline to identify common genes involved in gonadal sex

change across protogynous fishes. We successfully constructed a high-quality *de novo* genome

assembly applicable for genome-wide analysis in the protogynous three-spot wrasse and

acquired newly generated gonadal transcriptome data. Leveraging this information, we

conducted a comprehensive meta-analysis by integrating both public and newly derived

gonadal transcriptome data from multiple protogynous fishes spanning three orders. Through

this analysis, we identified 168 differentially expressed genes between the two gonadal states,

suggesting their potential universal role in gonadal sex change across diverse taxa of

protogynous fishes. Notably, we identified novel genes that were not previously associated

with gonadal sex change. In particular, we identified cholesterol dynamics as an important

candidate factor regulating the sex change process. Moreover, this finding highlights the

potential of transcriptomics meta-analysis as a valuable approach to uncover overlooked

mechanisms underlying conserved biological processes including sex change. We anticipate

that further data-driven research utilizing these novel sex change-related genes will open new

horizons in sex change research.

Experimental procedures

Animals. The adult female three-spot wrasse specimens collected in the field by local

fisherman in the northern region of Okinawa, Japan were purchased and employed for genome

and RNA sequencing. These specimens were subsequently cultivated in 1000 L tanks receiving

a continuous supply of unregulated seawater, and were subjected to natural photoperiod

conditions until sampling.

The maintenance and handling of fish in this study were conducted in accordance with the

"Guidelines for Animal Experiments of the Okinawa Churashima Foundation," with the same

consideration for animal care and welfare as that for higher vertebrates, including reptiles,

birds, and mammals. Animal ethical review and approval were not required for the present

study because the guidelines stipulated that approval from the Institutional Animal Care and

Use Committee of Okinawa Churashima Foundation is required for "higher" vertebrates, and is

waived for "lower" vertebrates including fish.

Genome sequencing and assembly for the three-spot wrasse. Muscle tissue from a mature

female individual was used for genome sequencing. The specimen was randomly selected from

cultivated tank. After anesthetizing the fish with 0.05% v/v 2-phenoxyethanol (FUJIFILM

Wako Pure Chemical), the muscle tissue was dissected, immediately frozen in liquid nitrogen,

and stored at −80 °C until genomic DNA extraction. High-molecular-weight DNA was

extracted from muscle using a NucleoBond AXG column (Macherey–Nagel), followed by

purification with phenol–chloroform. The concentration of the extracted DNA was measured

using Qubit (Thermo Fisher Scientific Inc., Waltham, MA, USA), and its size distribution was

first analyzed with a TapeStation 4200 (Agilent Technologies Inc., Santa Clara, CA, USA) to

ensure high integrity, and subsequently analyzed via pulse-field gel electrophoresis on a CHEF

DR-II system (Bio-Rad Laboratories, Inc., Hercules, CA, USA) to ensure a size range of

20–100 kb.

To obtain long-read sequence data, a SMRT sequence library was constructed using the

SMRTbell Express Template Prep Kit 2.0 (PacBio, Menlo Park, CA, USA) and sequenced on a

PacBio Sequel II system. The sequencing output was processed to generate circular consensus

sequences to obtain HiFi sequence reads. Adapter sequences were removed from the HiFi

reads using HiFiAdapterFilt (Sim et al., 2022). The obtained HiFi sequence reads were

assembled using the hifiasm (v0.16.1) assembler (Cheng et al., 2021) with default parameters.

The obtained contigs were first scaffolded with the P_RNA_scaffolder (Zhu et al., 2018) and

further scaffolded using Hi-C data as follows. The Hi-C library was prepared with the *in situ*

method using the muscle tissue of the individual used for DNA extraction according to the

iconHi-C protocol (Kadota et al., 2020), employing the restriction enzymes DpnII and HinfI,

and sequenced on a HiSeq X sequencing platform (Illumina, Inc., San Diego, CA, USA). The

obtained Hi-C read pairs, amounting to 232 million read pairs, were aligned to the HiFi

sequence contigs using the Juicer program (Durand et al., 2016), and the HiFi sequence contigs

were scaffolded via the 3d-dna (version 201008) to be consistent with the chromatin contact

profiles through manual curation on JuiceBox v1.11.08 (Dudchenko et al., 2018). The

continuity and completeness of the resulting genome assembly were assessed using the

webserver gVolante (Nishimura et al., 2017), in which the pipeline BUSCO (v5.4, reference

datasets: vertebrata_odb10) was implemented (Manni et al., 2021).

RNA-Seq of gonadal tissue in the three-spot wrasse. Based on previous studies (Horiguchi

et al., 2013; Nozu et al., 2009), we triggered a sex change in three-spot wrasse n=5 females

(body length; 8.0–9.3 cm, body weight; 12.4–20.9 g) to obtain transitional gonads through

treatment with an aromatase inhibitor (AI), which exerts inhibitory effects on the enzyme

responsible for estrogen synthesis. The collected gonads were cut into small pieces, one piece

was immersed in TRIzol reagent (Invitrogen) and the rest was fixed in 4%

paraformaldehyde/1× phosphate-buffered saline (PBS) for histological observation. Total RNA

in each sample was extracted promptly in accordance with the TRIzol instruction manual. The

extracted total RNA was dissolved in RNase-free water and stored at −80 °C until RNA-Seq

analysis. We selected two treated and two control samples for RNA-Seq analysis on the basis

of histological observation (Fig. S3).

Beads with oligo (dT) were used to isolate poly(A) mRNA after the total RNA was

collected from the samples. Fragmentation buffer was added to disrupt mRNA into short

fragments. Using these short fragments as templates, a random hexamer primer was used to

synthesize the first-strand cDNA. Second-strand cDNA was synthesized using buffer, dNTPs,

Rhase H, and DNA polymerase I. Short fragments were purified using the QiaQuick PCR

extraction kit (Qiagen) and resolved with EB buffer for end reparation and adding an "A" base.

Subsequently, the short fragments were connected to sequencing adapters. Following agarose

gel electrophoresis, suitable fragments were selected as templates for PCR amplification.

Finally, the library was sequenced using an Illumina HiSeq 2000 as 2 × 90 bp paired-ends. The

acquired reads were subjected to the quality control procedures described below and used for

downstream expression analysis.

Retrieval and acquisition of RNA-Seq data in protogynous fishes from public resources.

To obtain RNA-Seq data from the gonads of protogynous fishes in public databases, we

accessed SRA (Kodama et al., 2011) (accessed in October 2022) to search SRA data containing

the following keywords: "bony fishes (AND ovary OR gonad)." SRA data were converted into

FASTQ files using the Fasterq-dump program of the SRA Toolkit (v3.0.0).

Quality control of RNA-Seq data. To ensure the integrity of the RNA-Seq reads, trimming

and quality control procedures were performed. Specifically, we used Trim Galore (v0.6.7)

with the option "--quality 20 --paired" to eliminate low-quality bases from paired-end reads.

Read mapping against the reference genome. For species with available genomic

information, RNA-Seq reads were mapped to the reference genome sequences using HISAT2

(v2.2.1) (Kim et al., 2019) with the parameter "-q -dta -x". The resulting SAM files were

sorted and converted into BAM files using SAMtools software (v1.15.1) (Danecek et al.,

2021).

Generating General Transfer Format (gtf) files based on genome sequence and RNA-Seq

reads for reference gene annotation when quantifying expression values. To ensure

consistency in the use of annotation information for expression quantification in species with

available reference genomes, we employed the merge function of StringTie (v2.1.7) (Pertea et

al., 2015) to generate a gtf file that served as a guide for annotations. The transcriptome

assembly for each fish was conducted using StringTie with default settings based on the

mapping results (BAM file). The resulting gtf files were then merged using StringTie with the

"--merge" option, producing a unified file as the "merged.gtf". This "merged.gtf" file

contained a non-redundant set of assembled transcripts for each species. Generated

"merged.gtf" files in each species were designated as the reference annotation files for

quantifying transcript expression.

Quantification of transcripts expression and detection of differentially expressed

transcripts. When reanalyzing high-throughput data from multiple experiments, it is crucial to

consider batch effects (Leek et al., 2010). In the present study, we, therefore, calculated the

transcript per million (TPM) value for each transcript and utilized the difference in TPM values

between the paired data as an indicator for identifying differentially expressed transcripts.

Prior to expression quantification, we performed a principal component analysis (PCA)

using count data from mapped reads to examine the expression profiles of ovaries and gonads

in each species (Fig. S4). The PCA and visualization were conducted using the 'pcaExplorer'

package (version 2.30.0) (Marini & Binder, 2019) in the R program.

Transcript expression was quantified in species with available genome annotation using

StringTie, specifying the parameters as "-G merged.gtf -e". The resulting gtf file for each

sample provided the TPM values, which were used for downstream analysis.

TPM values were computed for species lacking reference genomes using Salmon (v.1.9.0)

(Patro et al., 2017) with the parameters "quant -I index -l A". To generate the transcriptome

sequences required for Salmon, we performed transcriptome assembly using the Trinity

(v2.13.1) assembler (Grabherr et al., 2011) with default settings. Open reading frames (ORFs)

and CDS were predicted from the transcriptome assembly using TransDecoder (v5.5.0) (Haas,

2018) with the same settings as described below. The resulting gff3 file obtained from the

TransDecoder run was used to extract transcript sequences corresponding to the predicted CDS

from the transcriptome assembly. The extracted sequence set was then utilized as the reference

transcriptome after eliminating duplicate sequences using seqkit (v2.3.0) (Zou et al., 2016)

with the "rmdup -s" command.

To assess changes in transcript expression, the expression ratio was determined using TPM

values obtained from intersexual gonad and ovary transcriptomes (referred to as the IO-ratio).

The IO-ratio for each transcript was calculated using Equation (1).

$\text{IO ratio} = \log_2(\text{TPM}_{\text{intersexual}} + 0.01) - \log_2(\text{TPM}_{\text{ovary}} + 0.01)$, (1)

This calculation was performed for all pairs of intersexual gonad and ovary transcriptomes. To

convert zeros to logarithms, a value of "0.01" was added to the TPM of each transcript.

Transcripts with an IO-ratio $> \log_2(5)$ (indicating a 5-fold expression change) were classified

as "upregulated," whereas transcripts with an IO-ratio $< -\log_2(5)$ (representing a 0.2-fold

expression change) were considered "downregulated." Transcripts outside these thresholds

were deemed "unchanged." The threshold was set based on the ±2 SD (standard deviation)

range of expression ratios of transcripts in each species (Table S5). To determine differential

expression between the intersexual gonad and ovary, an IO-score was calculated for each

transcript by subtracting the count of "downregulated" pairs from that of "upregulated" pairs,

as shown in Equation (2).

IO-score = (count number $_{upregulated}$) – (count number $_{downregulated}$), (2)

Using orthology inference as described below, the total IO-score for each transcript was

calculated by summing the IO-scores of all species. Total IO-score thresholds of 7 and −7 were

employed to identify transcripts that were differentially upregulated and downregulated,

respectively. The thresholds were arbitrarily set based on 20% above the maximum or below

the minimum values (31 or -31). All calculations for IO-ratios and IO-scores were performed

using R (v 4.2.1) (R Core Team, 2022).

Orthology inference among species. GFFread (v0.12.1) (Pertea & Pertea, 2020) was utilized

along with the generated "merged.gtf" file to extract transcript sequences from the reference

genomes. For species without reference genomes, transcript sequences were obtained via

assembly using Trinity, as described previously. TransDecoder was used to identify the coding

regions and translate them into amino acid sequences for all species. ORFs were extracted

from the transcriptome sequences using TransDecoder Longorfs with the parameters "-m 100".

CDS were then predicted using TransDecoder Predict with default settings, and the protein

sequences converted from the predicted CDS were acquired. Subsequently, a reciprocal

homology search was conducted between the predicted protein sequences of the three-spot

wrasse and those of other species, and the reciprocal best hit (RBH) was identified as the

ortholog. The program Diamond (v2.0.15) (Buchfink et al., 2014) was used for the RBH search

using the parameters "--ultra-sensitive -e 1e-3". To extract the orthologs shared among all

species, we consolidated the RBH results with the three-spot wrasse-predicted protein IDs.

Gene set enrichment analysis. Differentially expressed gene sets were analyzed using the

web tool Metascape (v3.5) (Zhou et al., 2019), which enables gene set enrichment analysis.

However, Metascape only supports the analysis of zebrafish among teleosts. Therefore, prior to

analysis, the predicted protein sequences of the three-spot wrasse were mapped to the

corresponding zebrafish Ensembl protein IDs. To obtain orthologous information, we utilized

the ggsearch36 program from the FASTA package (Pearson & Lipman, 1988) to globally align

the predicted protein sequences of the three-spot wrasse with the proteome sequences of

zebrafish (Ensembl Release 108) (Cunningham et al., 2022).

Visualization of cholesterol and steroid hormone biosynthesis pathways and orthologs

mapping to the pathways. The hypothetical cholesterol biosynthesis and steroid hormone

biosynthesis pathway in teleost fish was illustrated using PathVisio (Murphy et al., 2015).

Modifying information from WikiPathways (Pico et al., 2008) intended for zebrafish (WP1387,

https://www.wikipathways.org/instance/WP1387) and data from the yellow perch, *Perca*

flavescens (Kemski et al., 2020), facilitated the depiction of the cholesterol biosynthesis

pathway. Additionally, the steroid hormone biosynthesis pathway was constructed by adjusting

information sourced from various fish models (Craft et al., 2015; Fernandino et al., 2013; Li et

al., 2006; Liu et al., 2022; Nagahama et al., 2021). Moreover, through PathVisio, 7,289

candidate sex-change genes, converted into Ensembl zebrafish gene IDs, were mapped onto

the established biosynthesis pathways.

Availability of data. All sequencing data, including assembled sequences and raw sequence

reads, were deposited in the DNA Data Bank of Japan (DDBJ) under the umbrella BioProject

accession number PRJDB15835. The genome assembly was deposited in the DDBJ under

accession numbers BSYP01000001–BSYP01000227. Additionally, raw sequence reads were

deposited in the DDBJ under the following accession numbers: DRR466686 (PacBio HiFi

reads), DRR486910 (Illumina HiC reads), and DRR483512–DRR483515 (Illumina RNA-Seq

reads).

Competing interests

The authors have no conflicts of competing interest to declare that are relevant to the content

of this article.

Author contributions

RN and HB conceived and designed the study. MK and SK performed genome sequence and

de novo genome assembly of three-spot wrasse. RN & MN RNA-seq data production of

three-spot wrasse. RN performed RNA-seq data analysis and orthology inference. RN, MN,

SK and HB wrote a draft manuscript. All authors read and approved the final manuscript.

Acknowledgements

All authors thank Dr. Osamu Nishimura at DNA Sequencing Facility operated by Laboratory

for Phyloinformatics in RIKEN BDR for assistance in Hi-C data assessment. We thank the

Sesoko Station, Tropical Biosphere Research Center, University of the Ryukyus for allowing

us to use their research facility. This research was supported by the Center of Innovation for

Bio-Digital Transformation (BioDX), a program on the open innovation platform for

industry-academia co-creation (COI-NEXT), and the Japan Science and Technology Agency

(JST, COI-NEXT, JPMJPF2010). This work was partially received financial support by

International Research Organization for Advanced Science and Technology, Kumamoto

University, Kumamoto, Japan.

References

Al-Habsi, A. A., Massarsky, A., & Moon, T. W. (2016). Exposure to gemfibrozil and atorvastatin affects
cholesterol metabolism and steroid production in zebrafish (*Danio rerio*). *Comparative*
Biochemistry and Physiology Part B: Biochemistry and Molecular Biology, 199, 87–96.
doi:10.1016/j.cbpb.2015.11.009

Atz, J. W. (1964). Intersexuality in fishes. In C. N. Armstrong & A. J. Marshall (Eds.), *Intersexuality in*
Vertebrates Including Man (pp. 145–232). London: Academic press.

Bera, A., Chadha, N. K., Dasgupta, S., Chakravarty, S., & Sawant, P. B. (2020). Hypoxia-mediated
inhibition of cholesterol synthesis leads to disruption of nocturnal sex steroidogenesis in the
gonad of koi carp, *Cyprinus carpio. Fish physiology and biochemistry,* 46, 2421–2435.
doi:10.1007/s10695-020-00887-5

Bhandari, R. K., Komuro, H., Nakamura, S., Higa, M., & Nakamura, M. (2003). Gonadal restructuring
and correlative steroid hormone profiles during natural sex change in protogynous honeycomb
grouper (*Epinephelus merra*). *Zoological Science,* 20, 1399–1404.

Blue, M. L., Williams, D. L., Zucker, S., Khan, S. A., & Blum, C. B. (1983). Apolipoprotein E synthesis
in human kidney, adrenal gland, and liver. *Proceedings of the National Academy of Sciences of*
the United States of America, 80, 283-287. doi:10.1073/pnas.80.1.283

Bono, H. (2021). Meta-analysis of oxidative transcriptomes in insects. *Antioxidants,* 10, 345.
doi:10.3390/antiox10030345

Breton, T. S., Kenter, L. W., Greenlaw, K., Montgomery, J., Goetz, G. W., Berlinsky, D. L., &
Luckenbach, J. A. (2019). Initiation of sex change and gonadal gene expression in black sea
bass (*Centropristis striata*) exposed to exemestane, an aromatase inhibitor. *Comparative*
Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 228, 51–61.
doi:10.1016/j.cbpa.2018.10.024

Buchfink, B., Xie, C., & Huson, D. H. (2014). Fast and sensitive protein alignment using DIAMOND.
Nature Methods, 12, 59–60. doi:10.1038/nmeth.3176

Cheng, H., Concepcion, G. T., Feng, X., Zhang, H., & Li, H. (2021). Haplotype-resolved de novo
assembly using phased assembly graphs with hifiasm. *Nature Methods,* 18, 170–175.
doi:10.1038/s41592-020-01056-5

Craft, J. A., Bain, P. A., Papanicolaou, A., & Kumar, A. (2015). Identification of putative nuclear
receptors and steroidogenic enzymes in Murray-Darling Rainbowfish (*Melanotaenia fluviatilis*)
using RNA-Seq and *de novo* transcriptome assembly. *PloS one,* 10, e0142636.
doi:10.1371/journal.pone.0142636

Cunningham, F., Allen, J. E., Allen, J., Alvarez-Jarreta, J., Amode, M R., Armean, Irina M., . . . Flicek, P.
(2022). Ensembl 2022. *Nucleic Acids Research,* 50, D988–D995. doi:10.1093/nar/gkab1049

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., . . . Li, H. (2021). Twelve
years of SAMtools and BCFtools. *Gigascience,* 10, giab008. doi:10.1093/gigascience/giab008
Dudchenko, O., Shamim, M. S., Batra, S. S., Durand, N. C., Musial, N. T., Mostofa, R., . . . Aiden, E. L.
(2018). The Juicebox assembly tools module facilitates *de novo* assembly of mammalian
genomes with chromosome-length scaffolds for under $1000. *bioRxiv*, 254797.
doi:10.1101/254797
Durand, N. C., Shamim, M. S., Machol, I., Rao, S. S. P., Huntley, M. H., Lander, E. S., & Aiden, E. L.
(2016). Juicer provides a one-click system for analyzing loop-resolution Hi-C experiments.
Cell Systems, 3, 95–98. doi:10.1016/j.cels.2016.07.002
Fan, M., Yang, W., Zhang, W., & Zhang, L. (2022). The ontogenic gonadal transcriptomes provide
insights into sex change in the ricefield eel *Monopterus albus*. *BMC Zoology,* 7, 56.
doi:10.1186/s40850-022-00155-4
Fernandino, J. I., Hattori, R. S., Moreno Acosta, O. D., Strussmann, C. A., & Somoza, G. M. (2013).
Environmental stress-induced testis differentiation: androgen as a by-product of cortisol
inactivation. *General and Comparative Endocrinology,* 192, 36–44.
doi:10.1016/j.ygcen.2013.05.024
Gaudet, P., Livstone, M. S., Lewis, S. E., & Thomas, P. D. (2011). Phylogenetic-based propagation of
functional annotations within the Gene Ontology consortium. *Briefings in Bioinformatics,* 12,
449–462. doi:10.1093/bib/bbr042
Ge, H., Lin, K., Shen, M., Wu, S., Wang, Y., Zhang, Z., . . . Zheng, L. (2019). De novo assembly of a
chromosome-level reference genome of red-spotted grouper (*Epinephelus akaara*) using
nanopore sequencing and Hi-C. *Molecular Ecology Resources,* 19, 1461–1469.
doi:10.1111/1755-0998.13064
Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., . . . Regev, A. (2011).
Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature*
Biotechnology, 29, 644–652. doi:10.1038/nbt.1883
Gwynne, J. T., & Strauss, J. F. (1982). The role of lipoproteins in steroidogenesis and cholesterol
metabolism in steroidogenic glands. *Endocrine reviews,* 3, 299–329. doi:10.1210/edrv-3-3-299
Haas, B. (2018, October 2022). TransDecoder (v5.5.0). Retrieved from
https://github.com/TransDecoder/TransDecoder
Hasegawa, S., Noda, K., Maeda, A., Matsuoka, M., Yamasaki, M., & Fukui, T. (2012). Acetoacetyl-CoA
synthetase, a ketone body-utilizing enzyme, is controlled by SREBP-2 and affects serum
cholesterol levels. *Molecular Genetics and Metabolism,* 107, 553–560.
doi:10.1016/j.ymgme.2012.08.017
He, Z., Deng, F., Yang, D., He, Z., Hu, J., Ma, Z., . . . Yan, T. (2022). Crosstalk between sex-related
genes and apoptosis signaling reveals molecular insights into sex change in a protogynous
hermaphroditic teleost fish, ricefield eel *Monopterus albus*. *Aquaculture,* 552, 737918.
doi:10.1016/j.aquaculture.2022.737918
Herpin, A., & Schartl, M. (2011). Dmrt1 genes at the crossroads: a widespread and central class of
sexual development factors in fish. *The FEBS Journal,* 278, 1010–1019.

doi:10.1111/j.1742-4658.2011.08030.x

Horiguchi, R., Nozu, R., Hirai, T., Kobayashi, Y., Nagahama, Y., & Nakamura, M. (2013).

Characterization of gonadal soma-derived factor expression during sex change in the

protogynous wrasse, *Halichoeres trimaculatus*. *Developmental Dynamics, 242*, 388–399.

doi:10.1002/dvdy.23929

Horiguchi, R., Nozu, R., Hirai, T., Kobayashi, Y., & Nakamura, M. (2018). Expression patterns of sex

differentiation-related genes during gonadal sex change in the protogynous wrasse, *Halichoeres*

trimaculatus. *General and Comparative Endocrinology,* 257, 67–73.

doi:10.1016/j.ygcen.2017.06.017

Kadota, M., Nishimura, O., Miura, H., Tanaka, K., Hiratani, I., & Kuraku, S. (2020). Multifaceted Hi-C

benchmarking: what makes a difference in chromosome-scale genome scaffolding?

Gigascience, 9, giz158. doi:10.1093/gigascience/giz158

Kemski, M. M., Rappleye, C. A., Dabrowski, K., Bruno, R. S., & Wick, M. (2020). Transcriptomic

response to soybean meal-based diets as the first formulated feed in juvenile yellow perch

(*Perca flavescens*). *Scientific Reports,* 10, 3998. doi:10.1038/s41598-020-59691-z

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome alignment

and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology,* 37, 907–915.

doi:10.1038/s41587-019-0201-4

Kobayashi, T., Kajiura-Kobayashi, H., Guan, G., & Nagahama, Y. (2008). Sexual dimorphic expression

of DMRT1 and Sox9a during gonadal differentiation and hormone-induced sex reversal in the

teleost fish Nile tilapia (*Oreochromis niloticus*). *Developmental Dynamics,* 237, 297–306.

doi:10.1002/dvdy.21409

Kodama, Y., Shumway, M., & Leinonen, R. (2011). The sequence read archive: explosive growth of

sequencing data. *Nucleic Acids Research,* 40, D54–D56. doi:10.1093/nar/gkr854

Kroon, F. J., & Liley, N. R. (2000). The role of steroid hormones in protogynous sex change in the

Blackeye goby, *Coryphopterus nicholsii* (Teleostei: Gobiidae). *General and Comparative*

Endocrinology, 118, 273–283. doi:10.1006/gcen.2000.7459

Kuwamura, T., Sunobe, T., Sakai, Y., Kadota, T., & Sawada, K. (2020). Hermaphroditism in fishes: an

annotated list of species, phylogeny, and mating system. *Ichthyological Research,* 67, 341–360.

doi:10.1007/s10228-020-00754-6

Leek, J. T., Scharpf, R. B., Bravo, H. C., Simcha, D., Langmead, B., Johnson, W. E., . . . Irizarry, R. A.

(2010). Tackling the widespread and critical impact of batch effects in high-throughput data.

Nature Reviews. Genetics, 11, 733–739. doi:10.1038/nrg2825

Li, G. L., Liu, X. C., Zhang, Y., & Lin, H. R. (2006). Gonadal development, aromatase activity and P450

aromatase gene expression during sex inversion of protogynous red-spotted grouper

Epinephelus akaara (Temminck and Schlegel) after implantation of the aromatase inhibitor,

fadrozole. *Aquaculture Research,* 37, 484–491.

Liu, M., & de Mitcheson, Y. S. (2009). Gonad development during sexual differentiation in

hatchery-produced orange-spotted grouper (*Epinephelus coioides*) and humpback grouper

(*Cromileptes altivelis*) (Pisces: Serranidae, Epinephelinae). *Aquaculture,* 287, 191–202.

doi:10.1016/j.aquaculture.2008.10.027

Liu, Y., Kossack, M. E., McFaul, M. E., Christensen, L. N., Siebert, S., Wyatt, S. R., . . . Draper, B. W. (2022). Single-cell transcriptome reveals insights into the development and function of the zebrafish ovary. *eLife,* 11. doi:10.7554/eLife.76014

Lo Nostro, F., Grier, H., Andreone, L., & Guerrero, G. A. (2003). Involvement of the gonadal germinal epithelium during sex reversal and seasonal testicular cycling in the protogynous swamp eel, *Synbranchus marmoratus* Bloch 1795 (Teleostei, Synbranchidae). *Journal of Morphology,* 257, 107–126. doi:10.1002/jmor.10105

Mank, J. E., Promislow, D. E. L., & Avise, J. C. (2006). Evolution of alternative sex-determining mechanisms in teleost fishes. *Biological Journal of the Linnean Society,* 87, 83–93. doi:10.1111/j.1095-8312.2006.00558.x

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., Zdobnov, E. M., & Kelley, J. (2021). BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Molecular Biology and Evolution,* 38, 4647–4654. doi:10.1093/molbev/msab199

Marini, F., & Binder, H. (2019). pcaExplorer: an R/Bioconductor package for interacting with RNA-seq principal components. *BMC Bioinformatics,* 20, 331. doi:10.1186/s12859-019-2879-1

Miller, W. L. (2008). Steroidogenic enzymes. In C. E. Flück & W. L. Miller (Eds.), *Disorders of the Human Adrenal Cortex* (pp. 1–18). Basel: S.Karger AG.

Miyake, Y., Sakai, Y., & Kuniyoshi, H. (2012). Molecular cloning and expression profile of sex-specific genes, Figla and Dmrt1, in the protogynous hermaphroditic fish, *Halichoeres poecilopterus*. *Zoological Science,* 29, 690–701. doi:10.2108/zsj.29.690

Muncaster, S., Norberg, B., & Andersson, E. (2013). Natural sex change in the temperate protogynous Ballan wrasse *Labrus bergylta*. *Journal of Fish Biology,* 82, 1858–1870. doi:10.1111/jfb.12113

Murphy, R. F., Kutmon, M., van Iersel, M. P., Bohler, A., Kelder, T., Nunes, N., . . . Evelo, C. T. (2015). PathVisio 3: an extendable pathway analysis toolbox. *PLoS Computational Biology,* 11, e1004085. doi:10.1371/journal.pcbi.1004085

Nagahama, Y., Chakraborty, T., Paul-Prasanth, B., Ohta, K., & Nakamura, M. (2021). Sex determination, gonadal sex differentiation, and plasticity in vertebrate species. *Physiological Reviews,* 101, 1237–1308. doi:10.1152/physrev.00044.2019

Nakamura, M., Hourigan, T. F., Yamauchi, K., Nagahama, Y., & Grau, E. G. (1989). Histological and ultrastructural evidence for the role of gonadal steroid hormones in sex change in the protogynous wrasse *Thalassoma duperrey*. *Environmental Biology of Fishes,* 24, 117–136. doi:10.1007/bf00001282

Nishimura, O., Hara, Y., Kuraku, S., & Hancock, J. (2017). gVolante for standardizing completeness assessment of genome and transcriptome assemblies. *Bioinformatics,* 33, 3635–3637. doi:10.1093/bioinformatics/btx445

Nozu, R., Horiguchi, R., Kobayashi, Y., & Nakamura, M. (2015). Expression profile of doublesex/male abnormal-3-related transcription factor-1 during gonadal sex change in the protogynous wrasse, *Halichoeres trimaculatus*. *Molecular Reproduction and Development,* 82, 859–866.

doi:10.1002/mrd.22527

Nozu, R., Kojima, Y., & Nakamura, M. (2009). Short term treatment with aromatase inhibitor induces
sex change in the protogynous wrasse, *Halichoeres trimaculatus*. *General and Comparative*
Endocrinology, 161, 360–364. doi:10.1016/j.ygcen.2009.01.024

Ohta, K., Hirano, M., Mine, T., Mizutani, H., Yamaguchi, A., & Matsuyama, M. (2008). Body color
change and serum steroid hormone levels throughout the process of sex change in the adult
wrasse, *Pseudolabrus sieboldi*. *Marine Biology,* 153, 843–852. doi:10.1007/s00227-007-0856-0

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon provides fast and
bias-aware quantification of transcript expression. *Nature Methods,* 14, 417–419.
doi:10.1038/nmeth.4197

Payne, A. H., & Hales, D. B. (2004). Overview of steroidogenic enzymes in the pathway from
cholesterol to active steroid hormones. *Endocrine reviews,* 25, 947–970.
doi:10.1210/er.2003-0030

Pearson, W. R., & Lipman, D. J. (1988). Improved tools for biological sequence comparison.
Proceedings of the National Academy of Sciences of the United States of America, 85,
2444–2448. doi:10.1073/pnas.85.8.2444

Pertea, G., & Pertea, M. (2020). GFF Utilities: GffRead and GffCompare. *F1000Research,* 9.
doi:10.12688/f1000research.23297.1

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T.-C., Mendell, J. T., & Salzberg, S. L. (2015).
StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature*
Biotechnology, 33, 290–295. doi:10.1038/nbt.3122

Pico, A. R., Kelder, T., van Iersel, M. P., Hanspers, K., Conklin, B. R., & Evelo, C. (2008).
WikiPathways: pathway editing for the people. *PLoS Biology,* 6, e184.
doi:10.1371/journal.pbio.0060184

Pikuleva, I. A. (2006). Cholesterol-metabolizing cytochromes P450. *Drug metabolism and disposition :*
the biological fate of chemicals, 34, 513–520. doi:10.1124/dmd.105.008789

R Core Team. (2022). R: A language and environment for statistical computing. Retrieved from URL
https://www.R-project.org/

Rabosky, D. L., Chang, J., Title, P. O., Cowman, P. F., Sallan, L., Friedman, M., . . . Alfaro, M. E. (2018).
An inverse latitudinal gradient in speciation rate for marine fishes. *Nature,* 559, 392–395.
doi:10.1038/s41586-018-0273-1

Rahman, M. R., Petralia, M. C., Ciurleo, R., Bramanti, A., Fagone, P., Shahjaman, M., . . . Nicoletti, F.
(2020). Comprehensive analysis of RNA-Seq gene expression profiling of brain transcriptomes
reveals novel genes, regulators, and pathways in autism spectrum disorder. *Brain Sciences,* 10.
doi:10.3390/brainsci10100747

Reyland, M. E., Gwynne, J. T., Forgez, P., Prack, M. M., & Williams, D. L. (1991). Expression of the
human apolipoprotein E gene suppresses steroidogenesis in mouse Y1 adrenal cells.
Proceedings of the National Academy of Sciences of the United States of America, 88,
2375-2379. doi:10.1073/pnas.88.6.2375

Rodríguez-Marí, A., Yan, Y.-L., BreMiller, R. A., Wilson, C., Cañestro, C., & Postlethwait, J. H. (2005).

Characterization and expression pattern of zebrafish anti-Müllerian hormone (amh) relative to sox9a, sox9b, and cyp19a1a, during gonad development. *Gene Expression Patterns,* 5, 655–667. doi:10.1016/j.modgep.2005.02.008

Sakae, Y., Oikawa, A., Sugiura, Y., Mita, M., Nakamura, S., Nishimura, T., . . . Tanaka, M. (2020). Starvation causes female-to-male sex reversal through lipid metabolism in the teleost fish, medaka (*Olyzias latipes*). *Biology Open,* 9, bio050054. doi:10.1242/bio.050054

Senthilkumaran, B., Chi, W., Gao, Y., Hu, Q., Guo, W., & Li, D. (2017). Genome-wide analysis of brain and gonad transcripts reveals changes of key sex reversal-related genes expression and signaling pathways in three stages of *Monopterus albus*. *PloS one,* 12, e0173974. doi:10.1371/journal.pone.0173974

Sharpe, R. L., Drolet, M., & MacLatchy, D. L. (2006). Investigation of de novo cholesterol synthetic capacity in the gonads of goldfish (*Carassius auratus*) exposed to the phytosterol beta-sitosterol. *Reproductive Biology and Endocrinology,* 4, 60. doi:10.1186/1477-7827-4-60

Shibata, Y., Paul-Prasanth, B., Suzuki, A., Usami, T., Nakamoto, M., Matsuda, M., & Nagahama, Y. (2010). Expression of gonadal soma derived factor (GSDF) is spatially and temporally correlated with early testicular differentiation in medaka. *Gene Expression Patterns,* 10, 283–289. doi:10.1016/j.gep.2010.06.005

Sim, S. B., Corpuz, R. L., Simmonds, T. J., & Geib, S. M. (2022). HiFiAdapterFilt, a memory efficient read processing pipeline, prevents occurrence of adapter sequence in PacBio HiFi reads and their negative impacts on genome assembly. *BMC Genomics,* 23, 157. doi:10.1186/s12864-022-08375-1

Skaar, K. S., Nobrega, R. H., Magaraki, A., Olsen, L. C., Schulz, R. W., & Male, R. (2011). Proteolytically activated, recombinant anti-mullerian hormone inhibits androgen secretion, proliferation, and differentiation of spermatogonia in adult zebrafish testis organ cultures. *Endocrinology,* 152, 3527–3540. doi:10.1210/en.2010-1469

Thomas, J. T., Todd, E. V., Muncaster, S., Lokman, P. M., Damsteegt, E. L., Liu, H., . . . Gemmell, N. J. (2019). Conservation and diversity in expression of candidate genes regulating socially-induced female-male sex change in wrasses. *PeerJ,* 7, e7032. doi:10.7717/peerj.7032

Thorngate, F. E., Strockbine, P. A., Erickson, S. K., & Williams, D. L. (2002). Altered adrenal gland cholesterol metabolism in the apoE-deficient mouse. *Journal of Lipid Research,* 43, 1920–1926. doi:10.1194/jlr.M200205-JLR200

Todd, E. V., Ortega-Recalde, O., Liu, H., Lamm, M. S., Rutherford, K. M., Cross, H., . . . Gemmell, N. J. (2019). Stress, novel sex genes, and epigenetic reprogramming orchestrate socially controlled sex change. *Science Advances,* 5, eaaw7006. doi:10.1126/sciadv.aaw7006

Toga, K., & Bono, H. (2023). Meta-analysis of public RNA sequencing data revealed potential key genes associated with reproductive division of labor in social hymenoptera and termites. *International Journal of Molecular Sciences,* 24, 8353. doi:10.3390/ijms24098353

Trim Galore. (October 2022). Retrieved from https://github.com/FelixKrueger/TrimGalore

Wang, D. S., Kobayashi, T., Zhou, L. Y., Paul-Prasanth, B., Ijiri, S., Sakai, F., . . . Nagahama, Y. (2007). Foxl2 up-regulates aromatase gene transcription in a female-specific manner by binding to the

promoter as well as interacting with ad4 binding protein/steroidogenic factor 1. *Molecular*
Endocrinology, 21, 712–725. doi:10.1210/me.2006-0248

Wang, X. G., & Orban, L. (2007). Anti-Müllerian hormone and 11 β-hydroxylase show reciprocal
expression to that of aromatase in the transforming gonad of zebrafish males. *Developmental*
Dynamics, 236, 1329–1338. doi:10.1002/dvdy.21129

Webster, K. A., Schach, U., Ordaz, A., Steinfeld, J. S., Draper, B. W., & Siegfried, K. R. (2017). Dmrt1
is necessary for male sexual development in zebrafish. *Developmental Biology,* 422, 33–46.
doi:10.1016/j.ydbio.2016.12.008

Wu, D.-M., He, Z., Chen, T., Liu, Y., Ma, L.-P., & Ping, J. (2016). DNA hypermethylation of
acetoacetyl-CoA synthetase contributes to inhibited cholesterol supply and steroidogenesis in
fetal rat adrenals under prenatal nicotine exposure. *Toxicology,* 340, 43–52.
doi:10.1016/j.tox.2016.01.002

Wu, X., Yang, Y., Zhong, C., Guo, Y., Wei, T., Li, S., . . . Liu, X. (2020). Integration of ATAC-seq and
RNA-seq unravels chromatin accessibility during sex reversal in orange-spotted grouper
(*Epinephelus coioides*). *International Journal of Molecular Sciences,* 21, 2800.
doi:10.3390/ijms21082800

Xiao, L., Guo, Y., Wang, D., Zhao, M., Hou, X., Li, S., . . . Zhang, Y. (2020). Beta-hydroxysteroid
dehydrogenase genes in orange-spotted grouper (*Epinephelus coioides*): genome-wide
identification and expression analysis during sex reversal. *Frontiers in Genetics,* 11, 161.
doi:10.3389/fgene.2020.00161

Xue, L., Jia, D., Xu, L., Huang, Z., Fan, H., Chen, B., . . . Gao, Y. (2021). Bulk and single-cell RNA-seq
reveal the sexually dimorphic expression pattern of dmrtb1 in zig-zag eel (*Mastacembelus*
armatus). *Aquaculture,* 545, 737194. doi:10.1016/j.aquaculture.2021.737194

Zhang, C., He, Q., Cheng, H., Li, L., Ruan, X., Duan, X., . . . Zhao, H. (2022). Transcription factors
foxl2 and foxl3 regulate cyp19a1a and cyp11b in orange-spotted grouper (*Epinephelus*
coioides). *Aquaculture Reports,* 25, 101243. doi:10.1016/j.aqrep.2022.101243

Zhang, X., Guan, G., Li, M., Zhu, F., Liu, Q., Naruse, K., . . . Hong, Y. (2016). Autosomal gsdf acts as a
male sex initiator in the fish medaka. *Scientific Reports,* 6, 19738. doi:10.1038/srep19738

Zhang, Y., Zhang, S., Liu, Z., Zhang, L., & Zhang, W. (2013). Epigenetic modifications during sex
change repress gonadotropin stimulation of cyp19a1a in a teleost ricefield eel (*Monopterus*
albus). *Endocrinology,* 154, 2881–2890. doi:10.1210/en.2012-2220

Zhao, X., Luo, M., Li, Z., Zhong, P., Cheng, Y., Lai, F., . . . Zhou, R. (2018). Chromosome-scale
assembly of the Monopterus genome. *Gigascience,* 7, giy046. doi:10.1093/gigascience/giy046

Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., . . . Chanda, S. K.
(2019). Metascape provides a biologist-oriented resource for the analysis of systems-level
datasets. *Nature Communications,* 10, 1523. doi:10.1038/s41467-019-09234-6

Zhu, B.-H., Xiao, J., Xue, W., Xu, G.-C., Sun, M.-Y., & Li, J.-T. (2018). P_RNA_scaffolder: a fast and
accurate genome scaffolder using paired-end RNA-sequencing reads. *BMC Genomics,* 19, 175.
doi:10.1186/s12864-018-4567-3

Zhu, Y., Wang, C., Chen, X., & Guan, G. (2016). Identification of gonadal soma-derived factor

involvement in *Monopterus albus* (protogynous rice field eel) sex change. *Molecular Biology*

Reports, 43, 629–637. doi:10.1007/s11033-016-3997-8

Zou, Q., Shen, W., Le, S., Li, Y., & Hu, F. (2016). SeqKit: a cross-platform and ultrafast toolkit for

FASTA/Q file manipulation. *PloS one,* 11, e0163962. doi:10.1371/journal.pone.0163962

Figure legends

Fig. 1. Summarized phylogenetic tree of Actinopterygii. The order-level phylogenetic tree is

shown. The phylogenetic tree was derived from Rabosky et al., (2018) and modified.

Diamonds indicate orders with sequential hermaphroditism. Numbers in parentheses indicate

the number of summarized orders. Orders containing the fish species used in this study are

shown in bold.

Fig. 2. Scatter plot of the total Intersexual gonad and Ovary (IO)-score and Composition

of total IO-scores for up- and downregulated genes. a Scatterplot of total IO-scores for the

7,289 genes in descending order. Red circles indicate 99 upregulated genes; blue circles

indicate 69 downregulated genes. Gray circles (which have the appearance of lines because

they overlap each other) indicate unchanged genes ($6 \leq$ total IO-score ≤ -6). Composition of

total IO-scores for up- (**b**, n=99) and downregulated genes (**c**, n=69). Numbers in parentheses

next to species name indicate the number of RNA-Seq data pairs for each species used in this

study. Asterisk indicates a gene for which the total IO-score is derived from a single species.

Fig. 3. Results of enrichment analysis for differentially expressed genes using Metascape.

a 99 upregulated genes. **b** 69 downregulated genes.

Fig. 4. Hypothetical ovarian cholesterol biosynthesis pathway in protogynous fishes. The

pathway was visualized using PathVisio (https://pathvisio.org), referencing several fish

cholesterol synthesis pathways. Gene products are represented in black text, while metabolites

are in blue text. Gene product annotations follow Ensembl Zebrafish gene symbols. Red-filled

boxes represent gene products with confirmed upregulated expression in this study, while

blue-filled boxes denote gene products with downregulated expression. Open boxes represent

gene products with no expression variation. Light green-filled boxes represent gene products

not included among the 7289 genes but confirmed to be present in at least the three-spot

wrasse transcriptome. Dark gray-filled boxes indicate gene products that could not be

identified within the three-spot wrasse transcriptome. Circles extending from gene products

represent catalytic reactions, while black arrows indicate the conversion of metabolites. The

dashed arrows indicate indirect pathways summarizing multi-step processes.

Table 1. Statistics of genome assembly in Three-spot wrasse (fHalTri1.1) and other Labidae

fishes listed in NCBI Genomes.

	fHalTri1.1	*Cheilinus undulatus* (humphead wrasse)	*Notolabrus celidotus* (New Zealand spotty)	*Labrus bergylta* (ballan wrasse)	*Semicossyphus pulcher* (California sheephead)	*Tautogolabrus adspersus* (cunner)
Assembly accession	BSYP01000001–BSYP01000227	GCF_018320785.1	GCF_009762535.1	GCF_900080235.1	GCA_022749685.1	GCA_020745685.1
Total seq length (Mb)	849.9	1173.4	846.7	805.5	794.1	723.6
Max length (Mb)	44.2	59.6	42.4	9.1	38.5	37.1
Num of scaffolds	227	45	467	13,466	178	347
Scaffold N50 (Mb)	37.9	51.5	37.1	0.8	32.1	31.7
Scaffold L50	11	11	11	252	12	11
Num of seqs > 1 Mb	25	24	24	197	35	24
% of seqs > 1 Mb	97.3	99.7	97.6	43.9	97.1	99.9
Num of gaps	117	284	1,133	257	8	283
Complete BUSCO (%)	98.7*	99.0	97.1	98.9	98.9	98.1

858 * BUSCO version 5.4, reference datasets: vertebrata_odb10

Table 2. Statistics of orthology relationship among species.

Common name	Scientific name	Order	Reference genome	*Num of transcripts seqs obtained based on the RNA-seq data from gonadal tissue	Num of protein seqs predicted by TransDecoder	Num of Reciprocal Best Hit (vs Three-spot wrasse)
Three-spot wrasse	*Halichoeres trimaculatus*	Labriformes	Assembly in this study	27,573	24,965	24,965
Bluehead wrasse	*Thalassoma bifasciatum*	Labriformes	GCA_008086565.1	57,699	43,195	13,871
Hong Kong grouper	*Epinephelus akaara*	Perciformes	Ge et al., 2019	26,507	24,752	11,619
Orange-spotted grouper	*Epinephelus coioides*	Perciformes	GCA_900536245.1	30,867	34,191	12,507
Black seabass	*Centropristis striata*	Perciformes	NA	** 95,941	95,941	12,591
Asian swamp eel	*Monopterus albus*	Synbranchiformes	GCF_001952655.1	53,472	50,693	12,765
Zig-zag eel	*Mastacembelus armatus*	Synbranchiformes	GCF_900324485.2	48,724	40,908	13,095

* Based on the annotation file (merged gtf file), the extracted transcripts sequences from the

genome by gffread

** Among transcriptome sequences (594,726 seqs) assembled by Trinity, transcripts sequences

corresponding to coding seqences predicted via TransDecoder

890   **Table 3.** List of genes that contributed to GO term enrichment.

Up/Down	GO ID	Category	GO term	*Genes
Up	GO:0007129	Biological Process	homologous chromosome pairing at meiosis	*hormad1, terb2, si:ch211-199g17.9*
Up	GO:0007369	Biological Process	gastrulation	*zeb1b, tmem237a, col12a1b, pfn1, enpp2*
Up	GO:1902531	Biological Process	regulation of intracellular signal transduction	*cep55l, psd3l, arhgef18b, ccl19a.1, sh3bp1, enpp2, pik3r6b*
Up	GO:0005929	Cellular Component	cilium	*septin6, prom1a, tmem237a, si:ch211-66i15.4, flr*
Up	GO:0007281	Biological Process	germ cell development	*hormad1, gsdf, sycp3*
Down	GO:0010817	Biological Process	regulation of hormone levels	*cyp19a1a, aacs, hsd17b1, ece1*
Down	GO:0006413	Biological Process	translational initiation	*eif4g2b, eif4eb, eif4e1c*
Down	GO:0019752	Biological Process	carboxylic acid metabolic process	*lpl, ldha, aacs, amacr, acaca, dars2*
Down	GO:0008202	Biological Process	steroid metabolic process	*hsd17b1, amacr, sdr42e1*
Down	GO:1904888	Biological Process	cranial skeletal system development	*nog3, cep55l, cyldb, ece1*
Down	GO:1901362	Biological Process	organic cyclic compound biosynthetic process	*tbx5a, hsd17b1, alad, acaca, sdr42e1, adcy6a*

891   * Gene symbols follow the zebrafish gene nomenclature in Ensembl.

Fig. 1.

Basal Actinopterygii
◇ Anguilliformes... (4 taxa)
Osteoglossiformes... (2 taxa)
Alepocephaliformes
◇ Clupeiformes
◇ Cypriniformes... (5 taxa)
Lepidogalaxiiformes
Argentiniformes... (3 taxa)
◇ Stomiatiformes... (2 taxa)
Galaxiiformes
Ateleopodiformes
Aulopiformes
Myctophiformes
Polymixiiformes... (5 taxa)
Lampridiformes
Beryciformes... (2 taxa)
Ophidiiformes
Batrachoidiformes
◇ **Labriformes**
◇ Tetraodontiformes... (2 taxa)
Acanthuriformes... (2 taxa)
◇ Spariformes
◇ **Perciformes... (5 taxa)**
◇ Gobiiformes... (4 taxa)
◇ **Synbranchiformes... (2 taxa)**
Pleuronectiformes... (4 taxa)
Mugiliformes
◇ Incertae sedis in Ovalentaria
◇ Cichliformes... (5 taxa)

Percomorpha

Fig. 2.

Fig. 3.

a Upregulated genes

GO:0007129: homologous chromosome pairing at meiosis
GO:0007369: gastrulation
GO:1902531: regulation of intracellular signal transduction
GO:0060271: cilium assembly
GO:0007281: germ cell development

-log10(P)

b Downregulated genes

GO:0010817: regulation of hormone levels
GO:0006413: translational initiation
GO:0019752: carboxylic acid metabolic process
GO:0008202: steroid metabolic process
GO:1904888: cranial skeletal system development
GO:1901362: organic cyclic compound biosynthetic process

-log10(P)

Fig. 4.