Gonadal transcriptome of sex change in fishes: quest for causal factors common to

phylogenetically distributed protogynous species

Ryo Nozu[1,2], Mitsutaka Kadota[3,4], Masaru Nakamura[5], Shigehiro Kuraku[6,7], Hidemasa Bono[1,2*]

[1] Laboratory of Genome Informatics, Graduate School of Integrated Sciences for Life,

Hiroshima University, 3-10-23 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-0046, Japan

[2] Laboratory of BioDX, Genome Editing Innovation Center, Hiroshima University, Hiroshima,

Japan

[3] Laboratory for Phyloinformatics, RIKEN Center for Biosystems Dynamics Research (BDR),

Kobe, Japan

[4] Laboratory for Developmental Genome System, RIKEN Center for Biosystems Dynamics

Research (BDR), Kobe, Japan

[5] Okinawa Churashima Research Center, Okinawa Churashima Foundation, Motobu-cho,

Okinawa, Japan

[6] Molecular Life History Laboratory, Department of Genomics and Evolutionary Biology,

National Institute of Genetics, Mishima, Japan

[7] Department of Genetics, Sokendai (Graduate University for Advanced Studies), Mishima,

Japan.

Email:

RN, rnozu22@hiroshima-u.ac.jp; MK, mitsutaka.kadota@riken.jp; MN,

masaru220101@gmail.com; SK, skuraku@nig.ac.jp

[*]To whom correspondence may be addressed: Hidemasa Bono

bonohu@hiroshima-u.ac.jp

Abstract

Protogyny, capable of changing from female to male during their lifetime, is prevalent in 20

families of teleosts but believed to have evolved within specific evolutionary lineages.

Therefore, shared regulatory factors governing this process are expected to be conserved across

protogynous fish. However, a comprehensive understanding of this mechanism remains elusive.

To identify these factors, we conducted a meta-analysis using gonadal transcriptome data from

seven species. We curated data pairs of ovarian tissue and transitional gonad, and employed

ratios of expression level as a unified criterion for differential expression, enabling a meta-

analysis across species. Our approach revealed that classical sex change-related genes exhibited

differential expression levels between the ovary and transitional gonads, consistent with

previous reports on protogynous fishes. These results validate our methodology's robustness.

Additionally, we identified novel factors not previously linked to gonadal sex change in fish.

Notably, changes in the expression levels of acetoacetyl-CoA synthetase (*aacs*) and

apolipoprotein Eb (*apoeb*), which are involved in cholesterol synthesis and transport,

respectively, suggest that the levels of cholesterol, a precursor of steroid hormones crucial for

sex change, are decreased upon sex change onset in the gonads. This implies a potential

universal influence of cholesterol dynamics on gonadal transformation in protogyny.

Keywords: Meta-analysis, Protogynous fishes, Gonad, Sex change, Genome, Transcriptome

Introduction

Sequential hermaphroditism, a phenomenon in which individuals undergo sex change during their lifetime, has been extensively recognized in teleosts over the past several decades[1]. Phylogenetic analysis has shown that sequential hermaphroditism has been acquired in various evolutionary lineages[2,3] (Fig. 1a). Especially, protogyny (i.e., the transformation from female to male) is prevalent in 20 families but is thought to have emerged once in evolutionally the Percomorpha (Fig. 1a)[4]. Therefore, there are common features in the intricate regulatory mechanisms during protogynous sex change. Specifically, the replacement of ovarian tissue with testicular tissue is commonly observed during protogynous sex change[5-9]. Additionally, considering the high functional conservation of gonads across organisms, the existence of a regulatory mechanism governing gonadal transformation comprising common and essential factors across protogynous fishes is plausible; however, a comprehensive perspective of this process remains elusive.

Accordingly, elucidation of the molecular mechanisms underlying the gametogenesis switch in sex-changing fish has subsequently become a focus of research. In particular, the expression patterns of multiple sex-related genes during sex change within the family Labridae, which serves as a model for protogynous sex change, have been explored[10-15]. Recently, next-generation sequencing technology has enabled researchers to conduct comprehensive analyses of gene expression, allowing the identification of global gene expression profiles during gonadal transformation in protogynous fish belonging to Labriformes, Perciformes, and Synbranchiformes[16-20]. Thus, these results suggest the involvement of several pivotal genes in the ovary-to-testis transformation in sex-changing fish.

For example, a repertoire of sex-related genes, including cytochrome P450, family 19, subfamily A, polypeptide 1a (*cyp19a1a*), forkhead box L2a (*foxl2a*), folliculogenesis specific bHLH transcription factor (*figla*), anti-Müllerian hormone (*amh*), gonadal somatic cell derived factor (*gsdf*), doublesex and mab-3 related transcription factor 1 (*dmrt1*) were identified as

differentially expressed genes (DEGs) during sex change across protogynous fishes. Notably,

the gonadal expression of *cyp19a1a* and *foxl2a*, which are key players in maintaining ovarian

status, is reduced upon the onset of sex change in multiple species[14,21,22]. *cyp19a1a* encodes

aromatase, an enzyme responsible for the conversion of androgens (e.g., testosterone) to

estrogens (e.g., estradiol-17ß; E2), whereas *foxl2a* encodes a transcription factor that targets and

up-regulates *cyp19a1a* in teleost fishes[23,24]. Furthermore, the initiation of female-to-male sex

change, triggered by a rapid decline in E2 levels[6], is well-coordinated with the expression

profiles of these genes. Conversely, *amh*, *gsdf*, and *dmrt1* are crucial for testicular

differentiation and development in teleost fishes[25-31]. In several sex-changing species, the

upregulation of these genes coincides with the emergence of testicular tissue during transition,

suggesting that these genes are involved in testicular formation during gonadal sex change[10-

12,14,32].

Transcriptome analysis during gonadal transformation has also revealed differential

expression of several genes not previously implicated in sex determination or differentiation

pathways. In particular, upregulation of apolipoprotein Eb (*apoeb*), ectonucleotide

pyrophosphatase/phosphodiesterase 2 (*enpp2*), and l-rhamnose-binding lectin CSL2 (*csl2*)

expression, accompanied by downregulation of galectin-3 (*gal3*), was observed during the early

phase of gonadal sex change in the protogynous Asian swamp eel *Monopterus albus*. These

findings suggest that these factors are involved in gonadal transformation, particularly during

the initiation of sex change[16,19]. However, verifying their direct involvement and commonality

in the sex change process remains challenging.

Recently, by amalgamating transcriptome data from multiple studies, meta-analysis has

attracted attention as a powerful tool capable of providing novel insights overlooked by

traditional hypothesis-driven research methods[33-35]. In other words, it is expected that by

reanalyzing studies across multiple species and studies, universal factors controlling gonadal

transformation, which have not been detected in individual studies, could be identified among

protogynous fishes. Therefore, the objective of the present study was to identify the shared

factors involved in gonadal transformation across species using a meta-analysis of publicly

available transcriptome data from multiple experiments and species. Furthermore, in this study,

we focused on another well-studied species, the three-spot wrasse, *Halichoeres trimaculatus*, a

protogynous species belonging to the family Labridae and produced *de novo* genome assembly

and gonad transcriptome data undergoing sex change. This new data consolidated our dataset

encompassing seven protogynous species from three orders allowing exhaustive genome-wide

investigation. Through this dataset and uniform pipeline, we identified factors with differential

expression levels between the transitional gonads and ovaries (Fig. 1b). We reevaluated the

relationship between these factors and gonadal sex change, providing a novel perspective

regarding the mechanisms underlying this process.

Results

***De novo* genome assembly and transcriptome of gonadal tissue in the three-spot wrasse.**

High-throughput genome sequence data were obtained from a locally obtained female three-spot

wrasse from Okinawa, Japan. The obtained HiFi data, which comprised approximately 32 Gbp

and encompassed approximately 1.9 million DNA sequences, were assembled using the hifiasm

program[36]. Subsequently, the acquired Hi-C data were used for scaffolding, resulting in the

generation of 227 scaffolds. The longest scaffold measured 44.2 Mbp, with an N50 length (the

shortest of the lengths covering 50% of the genome) of 37.9 Mbp, contributing to a total

assembly length of approximately 849.9 Mbp. Assessment using BUSCO [37] revealed a

completeness > 98%, indicating a high level of coverage and contiguity in the *de novo* genome

assembly of the three-spot wrasse (Table 1).

The RNA-Seq reads obtained from transitional gonads and ovaries of additional females

were mapped to the genome assembly, and the obtained mapping results were used to assemble

the transcriptome for the annotation file ("merged.gtf"). This process yielded 27,573 transcript

sequences (Table 2). Subsequently, 24,695 coding sequences (protein sequences) were predicted

from the set of transcript sequences using TransDecoder[38].

Data curation of public RNA-Seq data and genomic information. We downloaded Sequence

Read Archive (SRA) data from six fish species for which sex changes have been reported[4]:

bluehead wrasse *Thalassoma bifasciatum* (order Labriformes), orange-spotted grouper

Epinephelus coioides (order Perciformes), Hong Kong grouper *E. akaara* (order Perciformes)*,*

black seabass *Centropristis striata* (order Perciformes); Asian swamp eel *Monopterus albus*

(order Synbranchiformes), zig-zag eel *Mastacembelus armatus* (order Synbranchiformes).

Furthermore, using the metadata associated with the SRA data, we curated the RNA-Seq data

into comparable sample pairs of intersexual (transitional) gonads and ovaries (a total of 31

paired data, including three-spot wrasse). The data pairs analyzed in this study are listed in

Supplementary Table 1.

The genome sequences of five of the seven species utilized in this study are publicly

available (Table 2) and served as reference genomes for read mapping. The genome sequences

of *T. bifasciatum* (GCA_008086565.1), *E. coioides* (GCA_900536245.1), *M. albus*

(GCF_001952655.1), and *M. armatus* (GCF_900324485.2) were retrieved from the National

Center for Biotechnology Information (NCBI) Genome Database (accessed in October 2022).

We also referenced the genome information of *E. akaara* reported by Ge et al., 2019[39].

Transcriptome assembly and inference of ortholog relationships. For the five species with

public reference genomes, the number of transcript sequences obtained using StringTie [40] with

the "--merge" option is shown in Table 2. For black sea bass, (without a reference genome), the

number of contiguous sequences assembled using Trinity [41] was 594,726. Among these, 95,941

transcript sequences corresponding to the predicted coding sequences (CDS) were specified as

references for expression quantification using Salmon[42]. The number of protein sequences

predicted using TransDecoder is shown in Table 2. Orthology inference and consolidation of

orthologs within the three-spot wrasse identified 7,289 orthologs that were common to all

species used in this study (Table 2), and thus were considered as candidate sex-change factors.

Identification of differentially expressed factors. The total intersexual gonad and ovary score

(IO-score) was calculated for these 7,289 factors (Fig. 2a, Supplementary Table 2, IO-score

calculation; see Methods). Among these, 99 factors were upregulated (Fig. 2b, Supplementary

Table 3) and 69 factors were downregulated (Fig. 2c, Supplementary Table 4). Whereas the total

IO-score of these factors was occasionally derived from a single species (up: *cep97*, *zmat2*,

tmem237a, *supt5h*; down: *znf16l*, *lrrc8db*, *sh2b1*, *tmem263*, *cyldb*, *manea*, *nol11*), in most cases,

the total IO-scores were based on contributions from two or more species. Notably, the

upregulated genes included *amh* and *gsdf*, which have been previously reported to play

important roles in gonadal sex change, particularly in testicular formation and development, in

some protogynous fish. Conversely, the downregulated genes included *cyp19a1a*, *hsd17b1*, and

foxl2a, which are crucial for the onset of sex change (Table 3).

Enrichment analysis of differentially expressed factors. Sequence homology search of the

three-spot wrasse proteins using ggsearch36 [43] against the reference zebrafish proteins (Ensembl

Release 108) [44]resulted in the identification of 90 homologs among the 99 upregulated factors

(Supplementary Table 3). No corresponding zebrafish orthologs were found for the remaining

nine factors that were common to all species analyzed in this study. The results of enrichment

analysis using the 90 factors that could be converted to zebrafish orthologs are shown in Fig. 3a.

The Gene Ontology (GO) terms "homologous chromosome pairing at meiosis" and "germ cell

development," which are related to reproduction, were significantly enriched among the

upregulated factors. The specific genes annotated with these GO terms included *hormad1*,

terb2, *si:ch211-199g17.9*, *gsdf*, and *sycp3* (Table 3).

We successfully obtained the corresponding zebrafish homologs for all of the 69

downregulated factors. The results of enrichment analysis are shown in Fig. 3b. Among the

downregulated factors, GO terms related to the regulation of steroid hormone levels, including

"regulation of hormone levels" and "steroid metabolic process," were significantly enriched.

Specific genes annotated with these enriched GO terms included *cyp19a1a*, *aacs*, *hsd17b1*,

ece1, *amacr*, and *sdr42e1* (Table 3).

Pathways mapping of candidate sex-change factors. Visualization of cholesterol and steroid

hormone biosynthesis pathways was conducted based on the GO terms obtained from the

enrichment analysis. By referencing pathway information for teleost fishes, pathway diagrams

were constructed and loaded with 7,289 factors converted to Ensembl zebrafish gene IDs (see

Methods). The results showed the mapping of most enzymes involved in the pathway from

acetyl-CoA to cholesterol synthesis (Fig. 4). Similarly, when mapping the steroid hormone

synthesis pathway, it was confirmed that enzymes involved in the synthesis pathway from

cholesterol to estradiol were included among the candidate sex-change factors (supplementary

figure 2).

Discussion

In this study, we conducted a meta-analysis of gonadal RNA-Seq data from seven protogynous

fish species using a unified analysis pipeline and threshold. The aim was to compare the

transcriptomes of ovaries and sexual transition gonads, with the goal of identifying common

factors involved in protogynous gonadal sex change. Toward this end, we first assembled the *de*

novo genome of the three-spot wrasse. The assembly statistics of the three-spot wrasse (Table 1)

were comparable to those of other Labridae species, such as humphead wrasse (*Cheilinus*

undulatus), ballan wrasse (*Labrus bergylta*), and New Zealand spotty (*Notolabrus celidotus*),

previously deposited in the NCBI RefSeq database. These results indicated that the *de novo*

genome assembly of the three-spot wrasse exhibited high quality, enabling us to perform

expression analysis and orthology inference with confidence.

Through this meta-analysis, we successfully identified sex-related factors including

cyp19a1a, *foxl2a*, *hsd17b1*, *amh*, and *gsdf*, that had been previously reported to play a role in

sex change in various protogynous fish species[10,14,17,19]. Furthermore, our enrichment analysis of

downregulated factors highlighted a term related to the "regulation of steroid hormone

production." This finding connects to those of previous reports, which indicate that a rapid

decrease in estrogen levels is crucial for female-to-male sex change[6,45,46]. Thus, by pooling data

across species, we were able to identify common factors involved in gonadal sex change.

Notably, our meta-analysis demonstrates the potential of utilizing transcriptome data from

multiple experiments to gain insights regarding the mechanisms underlying sex change in fish.

Our findings demonstrate that *aacs*, annotated with the GO term "regulation of hormone

levels," is downregulated during sex change. Notably, no previous reports have suggested the

involvement of *aacs* in sex change, indicating that the encoded protein (acetoacetyl-CoA

synthetase; Aacs) may represent a novel factor associated with this process. Typically, AACS

plays a critical role in cholesterol supply. Studies in mice have shown that *Aacs* knockdown

decreases total serum cholesterol levels[47]. Additionally, in rat adrenal glands, inhibition of *AACS*

expression through hypermethylation suppresses cholesterol supply and steroid synthesis[48].

Moreover, apolipoprotein Eb (encoded by *apoeb*, listed among the upregulated factors), has

been suggested to participate in cholesterol transportation in zebrafish[49]. Apolipoprotein E,

encoded by *APOE* which is the human ortholog of *apoeb*, serves as a crucial lipid transport

protein involved in the modulation of cholesterol metabolism and is also synthesized within

peripheral tissues[50]. In addition, a previous study indicates that APOE gene expression exerts

suppressive effects on steroid production in adrenal cells[51]. Specifically, the high expression of

APOE in tissues with active steroid production, including the adrenal glands, appears to shift

cholesterol metabolism from delivery to the steroidogenic pathway to storage of esterified

cholesterol, resulting in a decrease in steroid production levels[52]. These observations strongly

support the hypothesis that the downregulation of *aacs* and the upregulation of *apoeb* during sex

change contribute to suppression of the cholesterol supply and entire steroidogenesis.

Cholesterol serves as a precursor for steroid hormones, including estrogens, androgens, and

glucocorticoids[53-55], which play crucial roles in protogynous sex change. Previous studies have

shown that a decline in endogenous estrogens, particularly E2, triggers the transition from ovary

to testis in protogynous fish[6,45,46]. E2 is synthesized from testosterone via an aromatase enzyme,

encoded by *cyp19a1a*. Downregulation of *cyp19a1a* expression at the onset of sex change leads

to reduced E2 production. Studies on *M. albus* and *T. bifasciatum* have revealed higher DNA

methylation levels of the *cyp19a1a* gene upstream region in the testes than those in the ovaries,

suggesting that epigenetic modifications may contribute to the suppression of *cyp19a1a*

expression during gonadal sex change[17,21]. Our present findings could contribute one more

"piece of the puzzle", suggesting that the trigger of gonadal transformation may be attributed

not only to the decreased E2 production by the downregulation of aromatase activity but also to

the suppression of the cholesterol supply, the precursor of steroid hormones, leading to the

inhibition of the entire steroid biosynthesis. Interestingly, a decrease in the expression of

lipoprotein lipase (*lpl*) has been observed (Fig 4, supplementary table 4). Lipoprotein lipase is

involved in the biosynthetic process of fatty acid which serves as one of the sources of acetyl-

CoA, a precursor for cholesterol synthesis. This suggests the possibility of a concurrent

suppression of fatty acid supply to the gonads, might be leading to support the hypothesis.

Although this hypothesis requires further investigation and an understanding of cholesterol

dynamics during sex change, additional exploration is warranted. Notably, a study on medaka

(*Oryzias latipes*) subjected to starvation conditions implied a potential link between

masculinization and lower cholesterol levels[56]. Specifically, female-to-male sex reversal was

observed in starved medaka larvae, a gonochoristic fish, together with decreased cholesterol

levels in the starved samples. However, the precise pathways underlying masculinization and

cholesterol dynamics remain unknown. In general, cholesterol is synthesized not only in the

liver and adrenal glands but also in the gonads[57-59]. Therefore, gaining a comprehensive

understanding of the lipid metabolic mechanisms in fish gonads, including cholesterol synthesis

and transportation during sex change, may provide valuable insights regarding the mechanisms

underlying gonadal transformation. To clarify this perspective, the remaining challenge lies in

providing experimental evidence regarding whether it is possible to control gonadal

transformation by actually regulating cholesterol levels within the body.

Conclusions

In this study, our objective was to compare the transcriptomes of sex-transitional gonads with

those of ovaries using a unified pipeline to identify common factors involved in gonadal sex

change across protogynous fishes. We successfully constructed a high-quality de novo genome

assembly applicable for genome-wide analysis in the protogynous three-spot wrasse and

acquired newly generated gonadal transcriptome data. Leveraging this information, we

conducted a comprehensive meta-analysis by integrating both public and newly derived gonadal

transcriptome data from multiple protogynous fishes spanning three orders. Through this

analysis, we identified 168 differentially expressed factors between the two gonadal states,

suggesting their potential universal role in gonadal sex change across diverse taxa of

protogynous fishes. Notably, we identified novel factors that were not previously associated

with gonadal sex change. In particular, we identified cholesterol dynamics as an important

factor regulating the sex change process. Moreover, this finding highlights the potential of

transcriptomics meta-analysis as a valuable approach to uncover overlooked mechanisms

underlying conserved biological processes including sex change. We anticipate that further data-

driven research utilizing these novel sex change-related factors will open new horizons in sex

change research.

Methods

Animals. The adult female three-spot wrasse specimens collected in the field by local fisherman

in the northern region of Okinawa, Japan were purchased and employed for genome and RNA

sequencing. These specimens were subsequently cultivated in 1000 L tanks receiving a

continuous supply of unregulated seawater, and were subjected to natural photoperiod

conditions until sampling.

The maintenance and handling of fish in this study were conducted in accordance with the

"Guidelines for Animal Experiments of the Okinawa Churashima Foundation," with the same

consideration for animal care and welfare as that for higher vertebrates, including reptiles, birds,

and mammals. Animal ethical review and approval were not required for the present study

because the guidelines stipulated that approval from the Institutional Animal Care and Use

Committee of Okinawa Churashima Foundation is required for "higher" vertebrates, and is

waived for "lower" vertebrates including fish.

Genome sequencing and assembly for the three-spot wrasse. Muscle tissue from a mature

female individual was used for genome sequencing. The specimen was randomly selected from

cultivated tank. After anesthetizing the fish with 0.05% v/v 2-phenoxyethanol (FUJIFILM Wako

Pure Chemical), the muscle tissue was dissected, immediately frozen in liquid nitrogen, and

stored at −80 °C until genomic DNA extraction. High-molecular-weight DNA was extracted

from muscle using a NucleoBond AXG column (Macherey–Nagel), followed by purification

with phenol–chloroform. The concentration of the extracted DNA was measured using Qubit

(Thermo Fisher Scientific Inc.), and its size distribution was first analyzed with a TapeStation

4200 (Agilent Technologies) to ensure high integrity, and subsequently analyzed via pulse-field

gel electrophoresis on a CHEF DR-II system (BioRad) to ensure a size range of 20–100 kb.

To obtain long-read sequence data, a SMRT sequence library was constructed using the

SMRTbell Express Template Prep Kit 2.0 (PacBio) and sequenced on a PacBio Sequel II

system. The sequencing output was processed to generate circular consensus sequences to

obtain HiFi sequence reads. Adapter sequences were removed from the HiFi reads using

HiFiAdapterFilt[60]. The obtained HiFi sequence reads were assembled using the hifiasm

(v0.16.1) assembler [36] with default parameters. The obtained contigs were first scaffolded with

the P_RNA_scaffolder [61] and further scaffolded using Hi-C data as follows. The Hi-C library

was prepared with the in situ method using the muscle tissue of the individual used for DNA

extraction according to the iconHi-C protocol[62], employing the restriction enzymes DpnII and

HinfI, and sequenced on a HiSeq X sequencing platform (Illumina). The obtained Hi-C read

pairs, amounting to 232 million read pairs, were aligned to the HiFi sequence contigs using the

Juicer program[63], and the HiFi sequence contigs were scaffolded via the 3d-dna (version

201008) to be consistent with the chromatin contact profiles through manual curation on

JuiceBox v1.11.08[64]. The continuity and completeness of the resulting genome assembly were

assessed using the webserver gVolante[65], in which the pipeline BUSCO (v5.4, reference

datasets: vertebrata_odb10) was implemented[37].

RNA-Seq of gonadal tissue in the three-spot wrasse. Based on previous studies[10,45], we

triggered a sex change in three-spot wrasse n=5 females (body length; 8.0–9.3 cm, body weight;

12.4–20.9 g) to obtain transitional gonads through treatment with an aromatase inhibitor (AI),

which exerts inhibitory effects on the enzyme responsible for estrogen synthesis. The collected

gonads were cut into small pieces, one piece was immersed in TRIzol reagent (Invitrogen) and

the rest was fixed in 4% paraformaldehyde/1× phosphate-buffered saline (PBS) for histological

observation. Total RNA in each sample was extracted promptly in accordance with the TRIzol

instruction manual. The extracted total RNA was dissolved in RNase-free water and stored at

−80 °C until RNA-Seq analysis. We selected two treated and two control samples for RNA-Seq

analysis on the basis of histological observation (Supplementary fig. 1).

Beads with oligo (dT) were used to isolate poly(A) mRNA after the total RNA was collected

from the samples. Fragmentation buffer was added to disrupt mRNA into short fragments.

Using these short fragments as templates, a random hexamer primer was used to synthesize the

first-strand cDNA. Second-strand cDNA was synthesized using buffer, dNTPs, Rhase H, and

DNA polymerase I. Short fragments were purified using the QiaQuick PCR extraction kit

(Qiagen) and resolved with EB buffer for end reparation and adding an "A" base. Subsequently,

the short fragments were connected to sequencing adapters. Following agarose gel

electrophoresis, suitable fragments were selected as templates for PCR amplification. Finally,

the library was sequenced using an Illumina HiSeq 2000 as 2 × 90 bp paired-ends. The acquired

reads were subjected to the quality control procedures described below and used for

downstream expression analysis.

Retrieval and acquisition of RNA-Seq data in protogynous fishes from public resources. To

obtain RNA-Seq data from the gonads of protogynous fishes in public databases, we accessed

SRA [66] (accessed in October 2022) to search SRA data containing the following keywords:

"bony fishes (AND ovary OR gonad)." SRA data were converted into FASTQ files using the

Fasterq-dump program of the SRA Toolkit (v3.0.0).

Quality control of RNA-Seq data. To ensure the integrity of the RNA-Seq reads, trimming and

quality control procedures were performed. Specifically, we used Trim Galore (v0.6.7) [67] with

the option "--quality 20 --paired" to eliminate low-quality bases from paired-end reads.

Read mapping against the reference genome. For species with available genomic

information, RNA-Seq reads were mapped to the reference genome sequences using HISAT2

(v2.2.1) [68]with the parameter "-q -dta -x". The resulting SAM files were sorted and converted

into BAM files using SAMtools software (v1.15.1)[69].

Generating General Transfer Format (gtf) files based on genome sequence and RNA-Seq

reads for reference gene annotation when quantifying expression values. To ensure

consistency in the use of annotation information for expression quantification in species with

available reference genomes, we employed the merge function of StringTie (v2.1.7) [40]to

generate a gtf file that served as a guide for annotations. The transcriptome assembly for each

fish was conducted using StringTie with default settings based on the mapping results (BAM

file). The resulting gtf files were then merged using StringTie with the "--merge" option,

producing a unified file as the "merged.gtf". This "merged.gtf" file contained a non-redundant

set of assembled transcripts for each species. Generated "merged.gtf" files in each species were

designated as the reference annotation files for quantifying transcript expression.

Quantification of transcripts expression and detection of differentially expressed

transcripts. When reanalyzing high-throughput data from multiple experiments, it is crucial to

consider batch effects[70]. In the present study, we therefore calculated the transcript per million

(TPM) value for each transcript and utilized the difference in TPM values between the paired

data as an indicator for identifying differentially expressed transcripts.

Transcript expression was quantified in species with available genome annotation using

StringTie, specifying the parameters as "-G merged.gtf -e". The resulting gtf file for each

sample provided the TPM values, which were used for downstream analysis.

TPM values were computed for species lacking reference genomes using Salmon (v.1.9.0)[42]

with the parameters "quant -I index -l A". To generate the transcriptome sequences required for

Salmon, we performed transcriptome assembly using the Trinity (v2.13.1) assembler [41] with

default settings. Open reading frames (ORFs) and CDS were predicted from the transcriptome

assembly using TransDecoder (v5.5.0) [38]with the same settings as described below. The

resulting gff3 file obtained from the TransDecoder run was used to extract transcript sequences

corresponding to the predicted CDS from the transcriptome assembly. The extracted sequence

set was then utilized as the reference transcriptome after eliminating duplicate sequences using

seqkit (v2.3.0) [71]with the "rmdup -s" command.

To assess changes in transcript expression, the expression ratio was determined using TPM

values obtained from intersexual gonad and ovary transcriptomes (referred to as the IO-ratio).

The IO-ratio for each transcript was calculated using Equation (1).

$\text{IO ratio} = \log_2(\text{TPM}_{\text{intersexual}} + 0.01) - \log_2(\text{TPM}_{\text{ovary}} + 0.01), (1)$

This calculation was performed for all pairs of intersexual gonad and ovary transcriptomes. To

convert zeros to logarithms, a value of "0.01" was added to the TPM of each transcript.

Transcripts with an IO-ratio $> \log_2(5)$ (indicating a 5-fold expression change) were classified as

"upregulated," whereas transcripts with an IO-ratio $< -\log_2(5)$ (representing a 0.2-fold

expression change) were considered "downregulated." Transcripts outside these thresholds were

deemed "unchanged." To determine differential expression between the intersexual gonad and

ovary, an IO-score was calculated for each transcript by subtracting the count of

"downregulated" pairs from that of "upregulated" pairs, as shown in Equation (2).

$\text{IO-score} = (\text{count number}_{\text{upregulated}}) - (\text{count number}_{\text{downregulated}}), (2)$

Using orthology inference as described below, the total IO-score for each transcript was

calculated by summing the IO-scores of all species. Total IO-score thresholds of 7 and −7 were

employed to identify transcripts that were differentially upregulated and downregulated,

respectively. The thresholds were arbitrarily set based on 20% above the maximum or below the

minimum values (31 or -31). All calculations for IO-ratios and IO-scores were performed using

R (v 4.2.1)[72].

Orthology inference among species. GFFread (v0.12.1) [73]was utilized along with the

generated "merged.gtf" file to extract transcript sequences from the reference genomes. For

species without reference genomes, transcript sequences were obtained via assembly using

Trinity, as described previously. TransDecoder was used to identify the coding regions and

translate them into amino acid sequences for all species. ORFs were extracted from the

transcriptome sequences using TransDecoder Longorfs with the parameters "-m 100". CDS

were then predicted using TransDecoder Predict with default settings, and the protein sequences

converted from the predicted CDS were acquired. Subsequently, a reciprocal homology search

was conducted between the predicted protein sequences of the three-spot wrasse and those of

other species, and the reciprocal best hit (RBH) was identified as the ortholog. The program

Diamond (v2.0.15) [74]was used for the RBH search using the parameters "--ultra-sensitive -e 1e-

3". To extract the orthologs shared among all species, we consolidated the RBH results with the

three-spot wrasse-predicted protein IDs.

Gene set enrichment analysis. Differentially expressed gene sets were analyzed using the web

tool Metascape (v3.5)[75], which enables gene set enrichment analysis. However, Metascape only

supports the analysis of zebrafish among teleosts. Therefore, prior to analysis, the predicted

protein sequences of the three-spot wrasse were mapped to the corresponding zebrafish Ensembl

protein IDs. To obtain orthologous information, we utilized the ggsearch36 program from the

FASTA package [43] to globally align the predicted protein sequences of the three-spot wrasse

with the proteome sequences of zebrafish (Ensembl Release 108)[44].

Visualization of cholesterol and steroid hormone biosynthesis pathways and orthologs

mapping to the pathways. The hypothetical cholesterol biosynthesis and steroid hormone

biosynthesis pathway in teleost fish was illustrated using PathVisio[76]. Modifying information

from WikiPathways [77] intended for zebrafish (WP1387,

https://www.wikipathways.org/instance/WP1387) and data from the yellow perch, *Perca*

flavescens[78], facilitated the depiction of the cholesterol biosynthesis pathway. Additionally, the

steroid hormone biosynthesis pathway was constructed by adjusting information sourced from

various fish models[22,79-82]. Moreover, through PathVisio, 7,289 candidate sex-change factors,

converted into Ensembl zebrafish gene IDs, were mapped onto the established biosynthesis

pathways.

References

1   Atz, J. W. in *Intersexuality in Vertebrates Including Man*   (eds C. N. Armstrong & A. J.
Marshall)   145–232 (Academic press, 1964).

2   Avise, J. C. & Mank, J. E. Evolutionary perspectives on hermaphroditism in fishes. *Sex. Dev.* **3**,
152–163, doi:10.1159/000223079 (2009).

3   Erisman, B. E., Petersen, C. W., Hastings, P. A. & Warner, R. R. Phylogenetic perspectives on the
evolution of functional hermaphroditism in teleost fishes. *Integr. Comp. Biol.* **53**, 736–754,
doi:10.1093/icb/ict077 (2013).

4   Kuwamura, T., Sunobe, T., Sakai, Y., Kadota, T. & Sawada, K. Hermaphroditism in fishes: an
annotated list of species, phylogeny, and mating system. *Ichthyological research* **67**, 341–360,
doi:10.1007/s10228-020-00754-6 (2020).

5   Lo Nostro, F., Grier, H., Andreone, L. & Guerrero, G. A. Involvement of the gonadal germinal
epithelium during sex reversal and seasonal testicular cycling in the protogynous swamp eel,
Synbranchus marmoratus Bloch 1795 (Teleostei, Synbranchidae). *Journal of morphology* **257**,
107–126, doi:10.1002/jmor.10105 (2003).

6   Nakamura, M., Hourigan, T. F., Yamauchi, K., Nagahama, Y. & Grau, E. G. Histological and
ultrastructural evidence for the role of gonadal steroid hormones in sex change in the protogynous
wrasse   *Thalassoma   duperrey*.   *Environmental   biology   of   fishes*   **24**,   117–136,
doi:10.1007/bf00001282 (1989).

7   Ohta, K. *et al.* Body color change and serum steroid hormone levels throughout the process of sex
change in the adult wrasse, *Pseudolabrus sieboldi*. *Marine Biology* **153**, 843–852,
doi:10.1007/s00227-007-0856-0 (2008).

8   Muncaster, S., Norberg, B. & Andersson, E. Natural sex change in the temperate protogynous
Ballan wrasse *Labrus bergylta*. *J Fish Biol* **82**, 1858–1870, doi:10.1111/jfb.12113 (2013).

9   Bhandari, R. K., Komuro, H., Nakamura, S., Higa, M. & Nakamura, M. Gonadal restructuring and
correlative steroid hormone profiles during natural sex change in protogynous honeycomb grouper
(*Epinephelus merra*). *Zoolog Sci* **20**, 1399–1404 (2003).

472	10	Horiguchi, R. *et al.* Characterization of gonadal soma-derived factor expression during sex change
473		in the protogynous wrasse, *Halichoeres trimaculatus*. *Dev Dyn* **242**, 388–399,
474		doi:10.1002/dvdy.23929 (2013).

472   10   Horiguchi, R. *et al.* Characterization of gonadal soma-derived factor expression during sex change
473   in the protogynous wrasse, *Halichoeres trimaculatus*. *Dev Dyn* **242**, 388–399,
474   doi:10.1002/dvdy.23929 (2013).

475   11   Nozu, R., Horiguchi, R., Kobayashi, Y. & Nakamura, M. Expression profile of doublesex/male
476   abnormal-3-related transcription factor-1 during gonadal sex change in the protogynous wrasse,
477   *Halichoeres trimaculatus*. *Mol Reprod Dev* **82**, 859–866, doi:10.1002/mrd.22527 (2015).

478   12   Horiguchi, R., Nozu, R., Hirai, T., Kobayashi, Y. & Nakamura, M. Expression patterns of sex
479   differentiation-related genes during gonadal sex change in the protogynous wrasse, *Halichoeres*
480   *trimaculatus*. *Gen Comp Endocrinol* **257**, 67–73, doi:10.1016/j.ygcen.2017.06.017 (2018).

481   13   Kobayashi, Y., Horiguchi, R., Nozu, R. & Nakamura, M. Expression and localization of forkhead
482   transcriptional factor 2 (Foxl2) in the gonads of protogynous wrasse, *Halichoeres trimaculatus*.
483   *Biology of sex differences* **1**, 3, doi:10.1186/2042-6410-1-3 (2010).

484   14   Thomas, J. T. *et al.* Conservation and diversity in expression of candidate genes regulating
485   socially-induced female-male sex change in wrasses. *PeerJ* **7**, doi:10.7717/peerj.7032 (2019).

486   15   Miyake, Y., Sakai, Y. & Kuniyoshi, H. Molecular cloning and expression profile of sex-specific
487   genes, Figla and Dmrt1, in the protogynous hermaphroditic fish, *Halichoeres poecilopterus*.
488   *Zoolog Sci* **29**, 690-701, doi:10.2108/zsj.29.690 (2012).

489   16   Senthilkumaran, B. *et al.* Genome-wide analysis of brain and gonad transcripts reveals changes of
490   key sex reversal-related genes expression and signaling pathways in three stages of *Monopterus*
491   *albus*. *PLoS One* **12**, doi:10.1371/journal.pone.0173974 (2017).

492   17   Todd, E. V. *et al.* Stress, novel sex genes, and epigenetic reprogramming orchestrate socially
493   controlled sex change. *Science Advances* **5**, doi:10.1126/sciadv.aaw7006 (2019).

494   18   He, Z. *et al.* Crosstalk between sex-related genes and apoptosis signaling reveals molecular
495   insights into sex change in a protogynous hermaphroditic teleost fish, ricefield eel *Monopterus*
496   *albus*. *Aquaculture* **552**, doi:10.1016/j.aquaculture.2022.737918 (2022).

497   19   Fan, M., Yang, W., Zhang, W. & Zhang, L. The ontogenic gonadal transcriptomes provide insights
498   into sex change in the ricefield eel *Monopterus albus*. *BMC zoology* **7**, doi:10.1186/s40850-022-
499   00155-4 (2022).

500   20   Wu, X. *et al.* Integration of ATAC-seq and RNA-seq unravels chromatin accessibility during sex
501   reversal in Orange-spotted grouper (*Epinephelus coioides*). *International Journal of Molecular*
502   *Sciences* **21**, doi:10.3390/ijms21082800 (2020).

503   21   Zhang, Y., Zhang, S., Liu, Z., Zhang, L. & Zhang, W. Epigenetic modifications during sex change
504   repress gonadotropin stimulation of cyp19a1a in a teleost ricefield eel (*Monopterus albus*).
505   *Endocrinology* **154**, 2881–2890, doi:10.1210/en.2012-2220 (2013).

506   22   Li, G. L., Liu, X. C., Zhang, Y. & Lin, H. R. Gonadal development, aromatase activity and P450
507   aromatase gene expression during sex inversion of protogynous red-spotted grouper *Epinephelus*
508   *akaara* (Temminck and Schlegel) after implantation of the aromatase inhibitor, fadrozole.
509   *Aquaculture research* **37**, 484–491 (2006).

510   23   Wang, D. S. *et al.* Foxl2 up-regulates aromatase gene transcription in a female-specific manner by
511   binding to the promoter as well as interacting with ad4 binding protein/steroidogenic factor 1. *Mol*

512 *Endocrinol* **21**, 712–725, doi:10.1210/me.2006-0248 (2007).

513 24 Zhang, C. *et al.* Transcription factors foxl2 and foxl3 regulate cyp19a1a and cyp11b in orange-
514   spotted grouper (*Epinephelus coioides*). *Aquaculture reports* **25**, doi:10.1016/j.aqrep.2022.101243
515   (2022).

516 25 Shibata, Y. *et al.* Expression of gonadal soma derived factor (GSDF) is spatially and temporally
517   correlated with early testicular differentiation in medaka. *Gene Expr Patterns* **10**, 283–289,
518   doi:10.1016/j.gep.2010.06.005 (2010).

519 26 Zhang, X. *et al.* Autosomal gsdf acts as a male sex initiator in the fish medaka. *Sci Rep* **6**,
520   doi:10.1038/srep19738 (2016).

521 27 Kobayashi, T., Kajiura-Kobayashi, H., Guan, G. & Nagahama, Y. Sexual dimorphic expression of
522   DMRT1 and Sox9a during gonadal differentiation and hormone-induced sex reversal in the teleost
523   fish Nile tilapia (*Oreochromis niloticus*). *Dev Dyn* **237**, 297–306, doi:10.1002/dvdy.21409 (2008).

524 28 Rodríguez-Marí, A. *et al.* Characterization and expression pattern of zebrafish anti-Müllerian
525   hormone (amh) relative to sox9a, sox9b, and cyp19a1a, during gonad development. *Gene
526   expression patterns* **5**, 655–667, doi:10.1016/j.modgep.2005.02.008 (2005).

527 29 Wang, X. G. & Orban, L. Anti-Müllerian hormone and 11 β-hydroxylase show reciprocal
528   expression to that of aromatase in the transforming gonad of zebrafish males. *Dev Dyn* **236**, 1329–
529   1338, doi:10.1002/dvdy.21129 (2007).

530 30 Skaar, K. S. *et al.* Proteolytically activated, recombinant anti-mullerian hormone inhibits androgen
531   secretion, proliferation, and differentiation of spermatogonia in adult zebrafish testis organ
532   cultures. *Endocrinology* **152**, 3527–3540, doi:10.1210/en.2010-1469 (2011).

533 31 Webster, K. A. *et al.* Dmrt1 is necessary for male sexual development in zebrafish. *Developmental
534   biology* **422**, 33–46, doi:10.1016/j.ydbio.2016.12.008 (2017).

535 32 Zhu, Y., Wang, C., Chen, X. & Guan, G. Identification of gonadal soma-derived factor involvement
536   in *Monopterus albus* (protogynous rice field eel) sex change. *Molecular biology reports* **43**, 629–
537   637, doi:10.1007/s11033-016-3997-8 (2016).

538 33 Bono, H. Meta-analysis of oxidative transcriptomes in insects. *Antioxidants* **10**,
539   doi:10.3390/antiox10030345 (2021).

540 34 Rahman, M. R. *et al.* Comprehensive analysis of RNA-Seq gene expression profiling of brain
541   transcriptomes reveals novel genes, regulators, and pathways in autism spectrum disorder. *Brain
542   sciences* **10**, doi:10.3390/brainsci10100747 (2020).

543 35 Toga, K. & Bono, H. Meta-analysis of public RNA sequencing data revealed potential key genes
544   associated with reproductive division of labor in social hymenoptera and termites. *International
545   Journal of Molecular Sciences* **24**, doi:10.3390/ijms24098353 (2023).

546 36 Cheng, H., Concepcion, G. T., Feng, X., Zhang, H. & Li, H. Haplotype-resolved de novo assembly
547   using phased assembly graphs with hifiasm. *Nature methods* **18**, 170–175, doi:10.1038/s41592-
548   020-01056-5 (2021).

549 37 Manni, M. *et al.* BUSCO update: novel and streamlined workflows along with broader and deeper
550   phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Molecular
551   Biology and Evolution* **38**, 4647–4654, doi:10.1093/molbev/msab199 (2021).

38 Haas, B. *TransDecoder (v5.5.0)*, <https://github.com/TransDecoder/TransDecoder> (

39 Ge, H. *et al.* De novo assembly of a chromosome-level reference genome of red-spotted grouper
(*Epinephelus akaara*) using nanopore sequencing and Hi-C. *Mol Ecol Resour* **19**, 1461–1469,
doi:10.1111/1755-0998.13064 (2019).

40 Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from RNA-seq
reads. *Nature biotechnology* **33**, 290–295, doi:10.1038/nbt.3122 (2015).

41 Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a reference
genome. *Nature biotechnology* **29**, 644–652, doi:10.1038/nbt.1883 (2011).

42 Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and bias-
aware quantification of transcript expression. *Nature methods* **14**, 417–419,
doi:10.1038/nmeth.4197 (2017).

43 Pearson, W. R. & Lipman, D. J. Improved tools for biological sequence comparison. *Proceedings*
of the National Academy of Sciences **85**, 2444–2448, doi:10.1073/pnas.85.8.2444 (1988).

44 Cunningham, F. *et al.* Ensembl 2022. *Nucleic Acids Res* **50**, D988–D995,
doi:10.1093/nar/gkab1049 (2022).

45 Nozu, R., Kojima, Y. & Nakamura, M. Short term treatment with aromatase inhibitor induces sex
change in the protogynous wrasse, *Halichoeres trimaculatus*. *Gen Comp Endocrinol* **161**, 360–
364, doi:10.1016/j.ygcen.2009.01.024 (2009).

46 Kroon, F. J. & Liley, N. R. The role of steroid hormones in protogynous sex change in the Blackeye
goby, *Coryphopterus nicholsii* (Teleostei: Gobiidae). *Gen Comp Endocrinol* **118**, 273–283,
doi:10.1006/gcen.2000.7459 (2000).

47 Hasegawa, S. *et al.* Acetoacetyl-CoA synthetase, a ketone body-utilizing enzyme, is controlled by
SREBP-2 and affects serum cholesterol levels. *Molecular Genetics and Metabolism* **107**, 553–560,
doi:10.1016/j.ymgme.2012.08.017 (2012).

48 Wu, D.-M. *et al.* DNA hypermethylation of acetoacetyl-CoA synthetase contributes to inhibited
cholesterol supply and steroidogenesis in fetal rat adrenals under prenatal nicotine exposure.
Toxicology **340**, 43–52, doi:10.1016/j.tox.2016.01.002 (2016).

49 Gaudet, P., Livstone, M. S., Lewis, S. E. & Thomas, P. D. Phylogenetic-based propagation of
functional annotations within the Gene Ontology consortium. *Briefings in Bioinformatics* **12**, 449–
462, doi:10.1093/bib/bbr042 (2011).

50 Blue, M. L., Williams, D. L., Zucker, S., Khan, S. A. & Blum, C. B. Apolipoprotein E synthesis in
human kidney, adrenal gland, and liver. *Proc Natl Acad Sci USA* **80**, 283–287,
doi:10.1073/pnas.80.1.283 (1983).

51 Reyland, M. E., Gwynne, J. T., Forgez, P., Prack, M. M. & Williams, D. L. Expression of the
human apolipoprotein E gene suppresses steroidogenesis in mouse Y1 adrenal cells. *Proc Natl*
Acad Sci USA **88**, 2375–2379, doi:10.1073/pnas.88.6.2375 (1991).

52 Thorngate, F. E., Strockbine, P. A., Erickson, S. K. & Williams, D. L. Altered adrenal gland
cholesterol metabolism in the apoE-deficient mouse. *J Lipid Res* **43**, 1920–1926,
doi:10.1194/jlr.M200205-JLR200 (2002).

53 Payne, A. H. & Hales, D. B. Overview of steroidogenic enzymes in the pathway from cholesterol

592   to active steroid hormones. *Endocr Rev* **25**, 947–970, doi:10.1210/er.2003-0030 (2004).

54   Pikuleva, I. A. Cholesterol-metabolizing cytochromes P450. *Drug metabolism and disposition :*
the biological fate of chemicals **34**, 513–520, doi:10.1124/dmd.105.008789 (2006).

55   Miller, W. L. in *Disorders of the Human Adrenal Cortex*   (eds C.E. Flück & Walter L. Miller)
1–18 (S.Karger AG, 2008).

56   Sakae, Y. *et al.* Starvation causes female-to-male sex reversal through lipid metabolism in the
teleost fish, medaka (*Olyzias latipes*). *Biol Open* **9**, doi:10.1242/bio.050054 (2020).

57   Gwynne, J. T. & Strauss, J. F. The role of lipoproteins in steroidogenesis and cholesterol
metabolism in steroidogenic glands. *Endocr Rev* **3**, 299–329, doi:10.1210/edrv-3-3-299 (1982).

58   Sharpe, R. L., Drolet, M. & MacLatchy, D. L. Investigation of de novo cholesterol synthetic
capacity in the gonads of goldfish (*Carassius auratus*) exposed to the phytosterol beta-sitosterol.
Reproductive biology and endocrinology **4**, doi:10.1186/1477-7827-4-60 (2006).

59   Bera, A., Chadha, N. K., Dasgupta, S., Chakravarty, S. & Sawant, P. B. Hypoxia-mediated
inhibition of cholesterol synthesis leads to disruption of nocturnal sex steroidogenesis in the gonad
of koi carp, *Cyprinus carpio*. *Fish Physiol Biochem* **46**, 2421–2435, doi:10.1007/s10695-020-
00887-5 (2020).

60   Sim, S. B., Corpuz, R. L., Simmonds, T. J. & Geib, S. M. HiFiAdapterFilt, a memory efficient
read processing pipeline, prevents occurrence of adapter sequence in PacBio HiFi reads and their
negative impacts on genome assembly. *BMC genomics* **23**, doi:10.1186/s12864-022-08375-1
(2022).

61   Zhu, B.-H. *et al.* P_RNA_scaffolder: a fast and accurate genome scaffolder using paired-end RNA-
sequencing reads. *BMC genomics* **19**, doi:10.1186/s12864-018-4567-3 (2018).

62   Kadota, M. *et al.* Multifaceted Hi-C benchmarking: what makes a difference in chromosome-scale
genome scaffolding? *GigaScience* **9**, doi:10.1093/gigascience/giz158 (2020).

63   Durand, N. C. *et al.* Juicer provides a one-click system for analyzing loop-resolution Hi-C
experiments. *Cell Systems* **3**, 95–98, doi:10.1016/j.cels.2016.07.002 (2016).

64   Dudchenko, O. *et al.* The Juicebox assembly tools module facilitates *de novo* assembly of
mammalian genomes with chromosome-length scaffolds for under $1000. *bioRxiv*,
doi:10.1101/254797 (2018).

65   Nishimura, O., Hara, Y., Kuraku, S. & Hancock, J. gVolante for standardizing completeness
assessment of genome and transcriptome assemblies. *Bioinformatics* **33**, 3635–3637,
doi:10.1093/bioinformatics/btx445 (2017).

66   Kodama, Y., Shumway, M. & Leinonen, R. The sequence read archive: explosive growth of
sequencing data. *Nucleic Acids Res* **40**, D54–D56, doi:10.1093/nar/gkr854 (2011).

67   *Trim Galore*, <https://github.com/FelixKrueger/TrimGalore> (

68   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nature biotechnology* **37**, 907–915,
doi:10.1038/s41587-019-0201-4 (2019).

69   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**,
doi:10.1093/gigascience/giab008 (2021).

70   Leek, J. T. *et al.* Tackling the widespread and critical impact of batch effects in high-throughput
data. *Nature reviews. Genetics* **11**, 733–739, doi:10.1038/nrg2825 (2010).

71   Zou, Q., Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: a cross-platform and ultrafast toolkit for
FASTA/Q file manipulation. *PLoS One* **11**, doi:10.1371/journal.pone.0163962 (2016).

72   R Core Team. *R: A language and environment for statistical computing.*, <URL https://www.R-
project.org/> (2022).

73   Pertea, G. & Pertea, M. GFF Utilities: GffRead and GffCompare. *F1000Research* **9**,
doi:10.12688/f1000research.23297.1 (2020).

74   Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND.
Nature methods **12**, 59–60, doi:10.1038/nmeth.3176 (2014).

75   Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of systems-level
datasets. *Nature communications* **10**, doi:10.1038/s41467-019-09234-6 (2019).

76   Murphy, R. F. *et al.* PathVisio 3: An Extendable Pathway Analysis Toolbox. *PLoS Comput Biol*
11, doi:10.1371/journal.pcbi.1004085 (2015).

77   Pico, A. R. *et al.* WikiPathways: Pathway Editing for the People. *PLoS Biol* **6**,
doi:10.1371/journal.pbio.0060184 (2008).

78   Kemski, M. M., Rappleye, C. A., Dabrowski, K., Bruno, R. S. & Wick, M. Transcriptomic
response to soybean meal-based diets as the first formulated feed in juvenile yellow perch (*Perca*
flavescens). *Sci Rep* **10**, doi:10.1038/s41598-020-59691-z (2020).

79   Liu, Y. *et al.* Single-cell transcriptome reveals insights into the development and function of the
zebrafish ovary. *Elife* **11**, doi:10.7554/eLife.76014 (2022).

80   Craft, J. A., Bain, P. A., Papanicolaou, A. & Kumar, A. Identification of Putative Nuclear Receptors
and Steroidogenic Enzymes in Murray-Darling Rainbowfish (*Melanotaenia fluviatilis*) Using
RNA-Seq and De Novo Transcriptome Assembly. *PloS one* **10**, doi:10.1371/journal.pone.0142636
(2015).

81   Nagahama, Y., Chakraborty, T., Paul-Prasanth, B., Ohta, K. & Nakamura, M. Sex determination,
gonadal sex differentiation, and plasticity in vertebrate species. *Physiol Rev* **101**, 1237–1308,
doi:10.1152/physrev.00044.2019 (2021).

82   Fernandino, J. I., Hattori, R. S., Moreno Acosta, O. D., Strussmann, C. A. & Somoza, G. M.
Environmental stress-induced testis differentiation: androgen as a by-product of cortisol
inactivation. *Gen Comp Endocrinol* **192**, 36-44, doi:10.1016/j.ygcen.2013.05.024 (2013).

83   Rabosky, D. L. *et al.* An inverse latitudinal gradient in speciation rate for marine fishes. *Nature*
559, 392-395, doi:10.1038/s41586-018-0273-1 (2018).

Acknowledgements

All authors thank Dr. Osamu Nishimura at DNA Sequencing Facility operated by Laboratory for

Phyloinformatics in RIKEN BDR for assistance in Hi-C data assessment. We thank the Sesoko

Station, Tropical Biosphere Research Center, University of the Ryukyus for allowing us to use

their research facility.

Funding

This research was supported by the Center of Innovation for Bio-Digital Transformation

(BioDX), a program on the open innovation platform for industry-academia co-creation (COI-

NEXT), and the Japan Science and Technology Agency (JST, COI-NEXT, JPMJPF2010). This

work was partially received financial support by International Research Organization for

Advanced Science and Technology, Kumamoto University, Kumamoto, Japan.

Authors' contributions

RN and HB conceived and designed the study. MK and SK performed genome sequence and de

novo genome assembly of three-spot wrasse. RN & MN RNA-seq data production of three-spot

wrasse. RN performed RNA-seq data analysis and orthology inference. RN, MN, SK and HB

wrote a draft manuscript. All authors read and approved the final manuscript.

Availability of data

All sequencing data, including assembled sequences and raw sequence reads, were deposited in

the DNA Data Bank of Japan (DDBJ) under the umbrella BioProject accession number

PRJDB15835. The genome assembly was deposited in the DDBJ under accession numbers

BSYP01000001–BSYP01000227. Additionally, raw sequence reads were deposited in the

DDBJ under the following accession numbers: DRR466686 (PacBio HiFi reads), DRR486910

(Illumina HiC reads), and DRR483512-DRR483515 (Illumina RNA-Seq reads).

Additional information

All supplementary information is accessible in figshare repository

(https://doi.org/10.6084/m9.figshare.c.6704160)

Supplementary figure 1. Histological sections of control ovaries and transitional gonads treated

AI for 7 days in the three-spot wrasse, *Halichoeres trimaculatus*.

Supplementary figure 2. Hypothetical ovarian steroid hormones biosynthesis pathway in

protogynous fishes. The pathway was visualized using PathVisio (https://pathvisio.org),

referencing several fish cholesterol synthesis pathways. Gene products are represented in black

text, while metabolites are in blue text. Gene product annotations follow Ensembl Zebrafish

gene symbols. Red-filled boxes represent gene products with confirmed upregulated expression

in this study, while blue-filled boxes denote gene products with downregulated expression.

Open boxes represent gene products with no expression variation. Light green-filled boxes

represent gene products not included among the 7289 factors but confirmed to be present in at

least the three-spot wrasse transcriptome. Dark gray-filled boxes indicate gene products that

could not be identified within the three-spot wrasse transcriptome. Circles extending from gene

products represent catalytic reactions, while black arrows indicate the conversion of

metabolites.

Supplementary table 1

Supplementary table 2

Supplementary table 3

Supplementary table 4

Competing interests

The authors have no conflicts of competing interest to declare that are relevant to the content of

this article.

 Figure legends

Fig. 1. Summarized phylogenetic tree of Actinopterygii and schematic diagram of the

analysis pipeline in this study. a The order-level phylogenetic tree is shown. The phylogenetic

tree was derived from Daniel et al., (2018)[83] and modified. Diamonds indicate orders with

sequential hermaphroditism. Numbers in parentheses indicate the number of summarized

orders. Orders containing the fish species used in this study are shown in bold. **b** The consistent

line style of the arrows indicates the utilization of common tools for conducting data processing.

The letters in parentheses next to the species names represent the following order names; L:

Labriformes, P: Perciformes, S: Synbranchiformes.

Fig. 2. Scatter plot of the total Intersexual gonad and Ovary (IO)-score and Composition

of total IO-scores for up- and downregulated factors. a Scatterplot of total IO-scores for the

7,289 factors in descending order. Red circles indicate 99 upregulated factors; blue circles

indicate 69 downregulated factors. Gray circles (which have the appearance of lines because

they overlap each other) indicate unchanged factors ($6 \leq$ total IO-score ≤ -6). Composition of

total IO-scores for up- (**b**, n=99) and downregulated factors (**c**, n=69). Numbers in parentheses

next to species name indicate the number of RNA-Seq data pairs for each species used in this

study. Asterisk indicates a factor for which the total IO-score is derived from a single species.

Fig. 3. Results of enrichment analysis for differentially expressed factors using Metascape.

a 99 upregulated factors. **b** 69 downregulated factors.

Fig. 4. Hypothetical ovarian cholesterol biosynthesis pathway in protogynous fishes. The

pathway was visualized using PathVisio (https://pathvisio.org), referencing several fish

cholesterol synthesis pathways. Gene products are represented in black text, while metabolites

are in blue text. Gene product annotations follow Ensembl Zebrafish gene symbols. Red-filled

boxes represent gene products with confirmed upregulated expression in this study, while blue-

filled boxes denote gene products with downregulated expression. Open boxes represent gene

products with no expression variation. Light green-filled boxes represent gene products not

included among the 7289 factors but confirmed to be present in at least the three-spot wrasse

transcriptome. Dark gray-filled boxes indicate gene products that could not be identified within

the three-spot wrasse transcriptome. Circles extending from gene products represent catalytic

reactions, while black arrows indicate the conversion of metabolites. The dashed arrows

indicate indirect pathways summarizing multi-step processes.

Table 1. Statistics of genome assembly in Three-spot wrasse (fHalTri1.1) and other Labidae

fishes listed in NCBI Genomes.

	fHalTri1.1	*Cheilinus undulatus* (humphead wrasse)	*Notolabrus celidotus* (New Zealand spotty)	*Labrus bergylta* (ballan wrasse)	*Semicossyphus pulcher* (California sheephead)	*Tautogolabrus adspersus* (cunner)
Assembly accession	BSYP01000001–BSYP01000227	GCF_018320785.1	GCF_009762535.1	GCF_900080235.1	GCA_022749685.1	GCA_020745685.1
Total seq length (Mb)	849.9	1173.4	846.7	805.5	794.1	723.6
Max length (Mb)	44.2	59.6	42.4	9.1	38.5	37.1
Num of scaffolds	227	45	467	13,466	178	347
Scaffold N50 (Mb)	37.9	51.5	37.1	0.8	32.1	31.7
Scaffold L50	11	11	11	252	12	11
Num of seqs > 1 Mb	25	24	24	197	35	24
% of seqs > 1 Mb	97.3	99.7	97.6	43.9	97.1	99.9
Num of gaps	117	284	1,133	257	8	283
Complete BUSCO (%)	98.7*	99.0	97.1	98.9	98.9	98.1

759   * BUSCO version 5.4, reference datasets: vertebrata_odb10

773 **Table 2.** Statistics of orthology relationship among species.

Common name	Scientific name	Order	Reference genome	*Num of transcripts seqs obtained based on the RNA-seq data from gonadal tissue	Num of protein seqs predicted by TransDecoder	Num of RBH (vs Three-spot wrasse)
Three-spot wrasse	*Halichoeres trimaculatus*	Labriformes	Assembly in this study	27,573	24,965	24,965
Bluehead wrasse	*Thalassoma bifasciatum*	Labriformes	GCA_008086565.1	57,699	43,195	13,871
Hong Kong grouper	*Epinephelus akaara*	Perciformes	Ge et al., 2019	26,507	24,752	11,619
Orange-spotted grouper	*Epinephelus coioides*	Perciformes	GCA_900536245.1	30,867	34,191	12,507
Black seabass	*Centropristis striata*	Perciformes	NA	** 95,941	95,941	12,591
Asian swamp eel	*Monopterus albus*	Synbranchiformes	GCF_001952655.1	53,472	50,693	12,765
Zig-zag eel	*Mastacembelus armatus*	Synbranchiformes	GCF_900324485.2	48,724	40,908	13,095

774   * Based on the annotation file (merged gtf file), the extracted transcripts sequences from the

775   genome by gffread

776   ** Among transcriptome sequences (594,726 seqs) assembled by Trinity, transcripts sequences

777   corresponding to coding seqences predicted via TransDecoder

791 **Table 3.** List of genes that contributed to GO term enrichment.

Up/Down	GO ID	Category	GO term	*Genes
Up	GO:0007129	Biological Process	homologous chromosome pairing at meiosis	*hormad1, terb2, si:ch211-199g17.9*
Up	GO:0007369	Biological Process	gastrulation	*zeb1b, tmem237a, col12a1b, pfn1, enpp2*
Up	GO:1902531	Biological Process	regulation of intracellular signal transduction	*cep55l, psd3l, arhgef18b, ccl19a.1, sh3bp1, enpp2, pik3r6b*
Up	GO:0005929	Cellular Component	cilium	*septin6, prom1a, tmem237a, si:ch211-66i15.4, flr*
Up	GO:0007281	Biological Process	germ cell development	*hormad1, gsdf, sycp3*
Down	GO:0010817	Biological Process	regulation of hormone levels	*cyp19a1a, aacs, hsd17b1, ece1*
Down	GO:0006413	Biological Process	translational initiation	*eif4g2b, eif4eb, eif4e1c*
Down	GO:0019752	Biological Process	carboxylic acid metabolic process	*lpl, ldha, aacs, amacr, acaca, dars2*
Down	GO:0008202	Biological Process	steroid metabolic process	*hsd17b1, amacr, sdr42e1*
Down	GO:1904888	Biological Process	cranial skeletal system development	*nog3, cep55l, cyldb, ece1*
Down	GO:1901362	Biological Process	organic cyclic compound biosynthetic process	*tbx5a, hsd17b1, alad, acaca, sdr42e1, adcy6a*

792   * Gene symbols follow the zebrafish gene nomenclature in Ensembl.

Fig. 1.

a

Basal Actinopterygii
◇ Anguilliformes... (4 taxa)
Osteoglossiformes... (2 taxa)
Alepocephaliformes
◇ Clupeiformes
◇ Cypriniformes... (5 taxa)
Lepidogalaxiiformes
Argentiniformes... (3 taxa)
◇ Stomiatiformes... (2 taxa)
Galaxiiformes
Ateleopodiformes
Aulopiformes
Myctophiformes
Polymixiiformes... (5 taxa)
Lampridiformes
Beryciformes... (2 taxa)
Ophidiiformes
Batrachoidiformes
◇ **Labriformes**
◇ Tetraodontiformes... (2 taxa)
Acanthuriformes... (2 taxa)
◇ Spariformes
◇ **Perciformes... (5 taxa)**
◇ Gobiiformes... (4 taxa)
◇ **Synbranchiformes... (2 taxa)**
Pleuronectiformes... (4 taxa)
Mugiliformes
◇ Incertae sedis in Ovalentaria
◇ Cichliformes... (5 taxa)

Percomorpha

b

7 species
(3 orders)

Black seabass (P)

Bluehead wrasse (L)
Hong Kong grouper (P)
Orange spotted grouper (P)
Asian swamp eel (S)
Zig-zag eel (S)

Threespot wrasse (L)

RNA-seq data   Public data   Public data   In-house data

Genome   NA   Public data   *De novo* assembly

Quality Control

Transcriptome assembly

Read Mapping

Creation of reference annotation

Quantification of transcripts expression
Calculation of IO-ratio/-score in each species

Prediction of CDS
(protein sequences)

Ortholog inference & Identification of
proteins common to all species

Calculation of total IO-score

Enrichment analysis

Fig. 2.

Fig. 3.

a

Upregulated factors

GO:0007129: homologous chromosome pairing at meiosis
GO:0007369: gastrulation
GO:1902531: regulation of intracellular signal transduction
GO:0060271: cilium assembly
GO:0007281: germ cell development

b

Downregulated factors

GO:0010817: regulation of hormone levels
GO:0006413: translational initiation
GO:0019752: carboxylic acid metabolic process
GO:0008202: steroid metabolic process
GO:1904888: cranial skeletal system development
GO:1901362: organic cyclic compound biosynthetic process

Fig. 4.