Meta-analysis of gonadal transcriptomes in protogynous fishes provides novel perspective into

sex change machinery

Ryo Nozu[1,2], Mitsutaka Kadota[3,4], Masaru Nakamura[5], Shigehiro Kuraku[6,7], Hidemasa Bono[1,2#]

[1] Laboratory of Genome Informatics, Graduate School of Integrated Sciences for Life,

Hiroshima University, 3-10-23 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-0046, Japan

[2] Laboratory of BioDX, Genome Editing Innovation Center, Hiroshima University, Hiroshima,

Japan

[3] Laboratory for Phyloinformatics, RIKEN Center for Biosystems Dynamics Research (BDR),

Kobe, Japan

[4] Laboratory for Developmental Genome System, RIKEN Center for Biosystems Dynamics

Research (BDR), Kobe, Japan

[5] Okinawa Churashima Research Center, Okinawa Churashima Foundation, Motobu-cho,

Okinawa, Japan

[6] Molecular Life History Laboratory, Department of Genomics and Evolutionary Biology,

National Institute of Genetics, Mishima, Japan

[7] Department of Genetics, Sokendai (Graduate University for Advanced Studies), Mishima,

Japan.

Email:

RN, rnozu22@hiroshima-u.ac.jp; MK, mitsutaka.kadota@riken.jp; MN,

m-nakamura@okichura.jp; SK, skuraku@nig.ac.jp

[#]To whom correspondence may be addressed: Hidemasa Bono

bonohu@hiroshima-u.ac.jp

Abstract

Background: Protogynous fishes can change sex from female to male; concomitantly, their ovarian tissue is completely replaced by testicular tissue. The high functional conservation of gonads across organisms suggests the existence of as-yet-undefined common and essential factors in the regulation of such gonadal transformation. To identify these factors, we conducted a meta-analysis of public and newly derived gonadal transcriptome data from seven species.

Results: Classical sex change-related genes, including *cyp19a1a*, *foxl2a*, *hsd17b1*, *amh*, and *gsdf*, were differentially expressed between the ovary and transitional gonads, which corroborated previous reports on protogynous fishes. Moreover, we identified novel factors not previously associated with gonadal sex change in fish. Notably, the expression patterns of acetoacetyl-CoA synthetase (*aacs*) and apolipoprotein Eb (*apoeb*), which are involved in cholesterol synthesis and transport, respectively, suggest that the levels of cholesterol, a precursor of steroid hormones crucial for sex change, are decreased upon sex change onset in the gonads. Thus, cholesterol dynamics may influence gonadal transformation in protogynous fishes.

Conclusion: This finding indicates the potential of transcriptome meta-analysis as a valuable approach to uncover previously overlooked mechanisms underlying sex change. We anticipate that further data-driven research utilizing these novel sex change-related factors will open new horizons in sex change research.

Keywords: Meta-analysis, Protogynous fishes, Gonad, Sex change, Genome, Transcriptome

Background

Sequential hermaphroditism, a phenomenon in which individuals undergo sex change during

their lifetime, has been extensively recognized in teleosts over the past several decades [1].

Specifically, protogyny (i.e., the transformation from female to male) is prevalent in 20

families [2]. Phylogenetic analyses have shown that this characteristic has been independently

acquired in various evolutionary lineages [3, 4]. On the other hand, the complete replacement

of ovarian tissue with testicular tissue is commonly observed during protogynous sex change

[5-9], the intricate regulatory mechanisms responsible for inducing these remarkable

transformations in gonadal tissue remain incompletely understood. Furthermore, considering

the high functional conservation of gonads across organisms, the existence of a regulatory

mechanism governing gonadal transformation comprising common and essential factors across

protogynous fishes is plausible; however, a comprehensive perspective of this process remains

elusive.

Accordingly, elucidation of the molecular mechanisms underlying the gametogenesis

switch in sex-changing fish has subsequently become a focus of research. In particular, the

expression patterns of multiple sex-related genes during sex change within the family Labridae,

which serves as a model for protogynous sex change, have been explored [10-15]. Recently,

next-generation sequencing technology has enabled researchers to conduct comprehensive

analyses of gene expression, allowing the identification of global gene expression profiles

during gonadal transformation in protogynous fish belonging to Labriformes, Perciformes, and

Synbranchiformes [16-20]. Thus, these results suggest the involvement of several pivotal genes

in the ovary-to-testis transformation in sex-changing fish.

For example, a repertoire of sex-related genes, including cytochrome P450, family 19,

subfamily A, polypeptide 1a (*cyp19a1a*), forkhead box L2a (*foxl2a*), folliculogenesis specific

bHLH transcription factor (*figla*), anti-Müllerian hormone (*amh*), gonadal somatic cell derived

factor (*gsdf*), doublesex and mab-3 related transcription factor 1 (*dmrt1*) were identified as

differentially expressed genes (DEGs) during sex change across protogynous fishes. Notably,

the gonadal expression of *cyp19a1a* and *foxl2a*, which are key players in maintaining ovarian

status, is reduced upon the onset of sex change in multiple species [14, 21, 22]. *cyp19a1a*

encodes aromatase, an enzyme responsible for the conversion of androgens (e.g., testosterone)

to estrogens (e.g., estradiol-17ß; E2), whereas *foxl2a* encodes a transcription factor that targets

and up-regulates *cyp19a1a* in teleost fishes [23, 24]. Furthermore, the initiation of

female-to-male sex change, triggered by a rapid decline in E2 levels [6], is well-coordinated

with the expression profiles of these genes. Conversely, *amh*, *gsdf*, and *dmrt1* are crucial for

testicular differentiation and development in teleost fishes [25-31]. In several sex-changing

species, the upregulation of these genes coincides with the emergence of testicular tissue

during transition, suggesting that these genes are involved in testicular formation during

gonadal sex change [10-12, 14, 32].

Transcriptome analysis during gonadal transformation has also revealed differential

expression of several genes not previously implicated in sex determination or differentiation

pathways. In particular, upregulation of apolipoprotein Eb (*apoeb*), ectonucleotide

pyrophosphatase/phosphodiesterase 2 (*enpp2*), and l-rhamnose-binding lectin CSL2 (*csl2*)

expression, accompanied by downregulation of galectin-3 (*gal3*), was observed during the

early phase of gonadal sex change in the protogynous Asian swamp eel *Monopterus albus*.

These findings suggest that these factors are involved in gonadal transformation, particularly

during the initiation of sex change [16, 19]. However, verifying their direct involvement in the

sex change process remains challenging.

Recently, by amalgamating transcriptome data from multiple studies, meta-analysis has

attracted attention as a powerful tool capable of providing novel insights overlooked by

traditional hypothesis-driven research methods[33-35]. Therefore, the objective of the present

study was to identify the shared factors involved in gonadal transformation across species

using a meta-analysis of publicly available transcriptome data from multiple experiments and

species. To improve the performance of the meta-analysis, we included data (*de novo* genome

assembly data and RNA sequencing (RNA-Seq) data from gonads undergoing sex change)

generated in-house from an additional well-studied species, the three-spot wrasse,

Halichoeres trimaculatus, a protogynous species belonging to the family Labridae [10-13].

Through meta-analysis of RNA-Seq data encompassing seven protogynous species from three

orders, we identified 168 DEGs between the transitional gonads and ovaries (Fig. 1).

Subsequently, we reevaluated the relationship between these factors and gonadal sex change,

providing a novel perspective regarding the mechanisms underlying this process.

Results

***De novo* genome assembly and transcriptome of gonadal tissue in the three-spot wrasse.**

High-throughput genome sequence data were obtained from a locally obtained female

three-spot wrasse from Okinawa, Japan. The obtained HiFi data, which comprised

approximately 32 Gbp and encompassed approximately 1.9 million DNA sequences, were

assembled using the hifiasm program [36]. Subsequently, the acquired Hi-C data were used for

scaffolding, resulting in the generation of 227 scaffolds. The longest scaffold measured 44.2

Mbp, with an N50 length (the shortest of the lengths covering 50% of the genome) of 37.9

Mbp, contributing to a total assembly length of approximately 849.9 Mbp. Assessment using

BUSCO [37] revealed a completeness > 98%, indicating a high level of coverage and

contiguity in the *de novo* genome assembly of the three-spot wrasse (Table 1).

The RNA-Seq reads obtained from transitional gonads and ovaries of additional females

were mapped to the genome assembly, and the obtained mapping results were used to assemble

the transcriptome for the annotation file ("merged.gtf"). This process yielded 27,573 transcript

sequences (Table 2). Subsequently, 24,695 coding sequences (protein sequences) were

predicted from the set of transcript sequences using TransDecoder [38].

Data curation of public RNA-Seq data and genomic information. We downloaded

Sequence Read Archive (SRA) data from six fish species for which sex changes have been

reported [2]: bluehead wrasse *Thalassoma bifasciatum* (order Labriformes), orange-spotted

grouper *Epinephelus coioides* (order Perciformes), Hong Kong grouper *E. akaara* (order

Perciformes)*, black seabass *Centropristis striata* (order Perciformes); Asian swamp eel

Monopterus albus (order Synbranchiformes), zig-zag eel *Mastacembelus armatus* (order

Synbranchiformes). Furthermore, using the metadata associated with the SRA data, we curated

the RNA-Seq data into comparable sample pairs of intersexual (transitional) gonads and

ovaries (a total of 31 paired data, including three-spot wrasse). The data pairs analyzed in this

study are listed in Supplementary Table 1.

The genome sequences of five of the seven species utilized in this study are publicly

available (Table 2) and served as reference genomes for read mapping. The genome sequences

of *T. bifasciatum* (GCA_008086565.1), *E. coioides* (GCA_900536245.1), *M. albus*

(GCF_001952655.1), and *M. armatus* (GCF_900324485.2) were retrieved from the National

Center for Biotechnology Information (NCBI) Genome Database (accessed in October 2022).

We also referenced the genome information of *E. akaara* reported by Ge et al., 2019 [39].

Transcriptome assembly and inference of ortholog relationships. For the five species with

public reference genomes, the number of transcript sequences obtained using StringTie [40]

with the "--merge" option is shown in Table 2. For black sea bass, (without a reference

genome), the number of contiguous sequences assembled using Trinity [41] was 594,726.

Among these, 95,941 transcript sequences corresponding to the predicted coding sequences

(CDS) were specified as references for expression quantification using Salmon [42]. The

number of protein sequences predicted using TransDecoder is shown in Table 2. Orthology

inference and consolidation of orthologs within the three-spot wrasse identified 7,289

orthologs that were common to all species used in this study (Table 2), and thus were

considered as candidate sex-change factors.

Identification of differentially expressed transcripts. The total intersexual gonad and ovary

score (IO-score) was calculated for these 7,289 factors (Supplementary Table 2, IO-score

calculation; see Methods). Among these, 99 factors were upregulated (Fig. 2a, Fig. 3,

Supplementary Table 3) and 69 factors were downregulated (Fig. 2a, Fig. 4, Supplementary

Table 4). Whereas the total IO-score of these factors was occasionally derived from a single

species (up: *cep97*, *zmat2*, *tmem237a*, *supt5h*; down: *znf16l*, *lrrc8db*, *sh2b1*, *tmem263*, *cyldb*,

manea, *nol11*), in most cases, the total IO-scores were based on contributions from two or

more species. Notably, the upregulated genes included *amh* and *gsdf*, which have been

previously reported to play important roles in gonadal sex change, particularly in testicular

formation and development, in some protogynous fish. Conversely, the downregulated genes

included *cyp19a1a*, *hsd17b1*, and *foxl2a*, which are crucial for the onset of sex change (Table

3).

Enrichment analysis of DEGs. Sequence homology search of the three-spot wrasse proteins

using ggsearch36 [43] against the reference zebrafish proteins (Ensembl Release 108) [44]

resulted in the identification of 90 homologs among the 99 upregulated factors (Supplementary

Table 3). No corresponding zebrafish orthologs were found for the remaining nine factors that

were common to all species analyzed in this study. The results of enrichment analysis using the

90 factors that could be converted to zebrafish orthologs are shown in Fig. 2b. The Gene

Ontology (GO) terms "homologous chromosome pairing at meiosis" and "germ cell

development," which are related to reproduction, were significantly enriched among the

upregulated factors. The specific genes annotated with these GO terms included *hormad1*,

terb2, *si:ch211-199g17.9*, *gsdf*, and *sycp3* (Table 3).

180   We successfully obtained the corresponding zebrafish homologs for all of the 69

181 downregulated factors. The results of enrichment analysis are shown in Fig. 2c. Among the

182 downregulated factors, GO terms related to the regulation of steroid hormone levels, including

183 "regulation of hormone levels" and "steroid metabolic process," were significantly enriched.

184 Specific genes annotated with these enriched GO terms included *cyp19a1a*, *aacs*, *hsd17b1*,

185 *ece1*, *amacr*, and *sdr42e1* (Table 3).

187 **Pathways mapping of candidate sex-change factors.** Visualization of cholesterol and steroid

188 hormone biosynthesis pathways was conducted based on the GO terms obtained from the

189 enrichment analysis. By referencing pathway information for teleost fishes, pathway diagrams

190 were constructed and loaded with 7,289 factors converted to Ensembl zebrafish gene IDs (see

191 Methods). The results showed the mapping of most enzymes involved in the pathway from

192 acetyl-CoA to cholesterol synthesis (Fig. 5). Similarly, when mapping the steroid hormone

193 synthesis pathway, it was confirmed that enzymes involved in the synthesis pathway from

194 cholesterol to estradiol were included among the candidate sex-change factors (supplementary

195 figure 2).

198 **Discussion**

199 In this study, we conducted a meta-analysis of gonadal RNA-Seq data from seven protogynous

200 fish species using a unified analysis pipeline and threshold. The aim was to compare the

201 transcriptomes of ovaries and sexual transition gonads, with the goal of identifying common

202 factors involved in protogynous gonadal sex change. Toward this end, we first assembled the

203 *de novo* genome of the three-spot wrasse. The assembly statistics of the three-spot wrasse

204 (Table 1) were comparable to those of other Labridae species, such as humphead wrasse

205 (*Cheilinus undulatus*), ballan wrasse (*Labrus bergylta*), and New Zealand spotty (*Notolabrus*

celidotus), previously deposited in the NCBI RefSeq database. These results indicated that the

de novo genome assembly of the three-spot wrasse exhibited high quality, enabling us to

perform expression analysis and orthology inference with confidence.

Through this meta-analysis, we successfully identified sex-related factors including

cyp19a1a, *foxl2a*, *hsd17b1*, *amh*, and *gsdf*, that had been previously reported to play a role in

sex change in various protogynous fish species[10, 14, 17, 19]. Furthermore, our enrichment

analysis of downregulated factors highlighted a term related to the "regulation of steroid

hormone production." This finding connects to those of previous reports, which indicate that a

rapid decrease in estrogen levels is crucial for female-to-male sex change [6, 45, 46]. Thus, by

pooling data across species, we were able to identify common factors involved in gonadal sex

change. Notably, our meta-analysis demonstrates the potential of utilizing transcriptome data

from multiple experiments to gain insights regarding the mechanisms underlying sex change in

fish.

Our findings demonstrate that *aacs*, annotated with the GO term "regulation of hormone

levels," is downregulated during sex change. Notably, no previous reports have suggested the

involvement of *aacs* in sex change, indicating that the encoded protein (acetoacetyl-CoA

synthetase; Aacs) may represent a novel factor associated with this process. Typically, AACS

plays a critical role in cholesterol supply. Studies in mice have shown that *Aacs* knockdown

decreases total serum cholesterol levels [47]. Additionally, in rat adrenal glands, inhibition of

AACS expression through hypermethylation suppresses cholesterol supply and steroid

synthesis [48]. Moreover, apolipoprotein Eb (encoded by *apoeb*, listed among the upregulated

factors), has been suggested to participate in cholesterol transportation in zebrafish [49].

Apolipoprotein E, encoded by *APOE* which is the human ortholog of *apoeb*, serves as a crucial

lipid transport protein involved in the modulation of cholesterol metabolism and is also

synthesized within peripheral tissues [50]. In addition, a previous study indicates that APOE

gene expression exerts suppressive effects on steroid production in adrenal cells [51].

Specifically, the high expression of APOE in tissues with active steroid production, including

the adrenal glands, appears to shift cholesterol metabolism from delivery to the steroidogenic

pathway to storage of esterified cholesterol, resulting in a decrease in steroid production levels

[52]. These observations strongly support the hypothesis that the downregulation of *aacs* and

the upregulation of *apoeb* during sex change contribute to suppression of the cholesterol

supply and entire steroidogenesis.

Cholesterol serves as a precursor for steroid hormones, including estrogens, androgens, and

glucocorticoids [53-55], which play crucial roles in protogynous sex change. Previous studies

have shown that a decline in endogenous estrogens, particularly E2, triggers the transition from

ovary to testis in protogynous fish [6, 45, 46]. E2 is synthesized from testosterone via an

aromatase enzyme, encoded by *cyp19a1a*. Downregulation of *cyp19a1a* expression at the onset

of sex change leads to reduced E2 production. Studies on *M. albus* and *T. bifasciatum* have

revealed higher DNA methylation levels of the *cyp19a1a* gene upstream region in the testes

than those in the ovaries, suggesting that epigenetic modifications may contribute to the

suppression of *cyp19a1a* expression during gonadal sex change [17, 21]. Our present findings

could contribute one more "piece of the puzzle", suggesting that the trigger of gonadal

transformation may be attributed not only to the decreased E2 production by the

downregulation of aromatase activity but also to the suppression of the cholesterol supply, the

precursor of steroid hormones, leading to the inhibition of the entire steroid biosynthesis.

Interestingly, a decrease in the expression of lipoprotein lipase (*lpl*) has been observed (Fig 5,

supplementary table 4). Lipoprotein lipase is involved in the biosynthetic process of fatty acid

which serves as one of the sources of acetyl-CoA, a precursor for cholesterol synthesis. This

suggests the possibility of a concurrent suppression of fatty acid supply to the gonads, might be

leading to support the hypothesis. Although this hypothesis requires further investigation and

an understanding of cholesterol dynamics during sex change, additional exploration is

warranted. Notably, a study on medaka (*Oryzias latipes*) subjected to starvation conditions

implied a potential link between masculinization and lower cholesterol levels [56]. Specifically,

female-to-male sex reversal was observed in starved medaka larvae, a gonochoristic fish,

together with decreased cholesterol levels in the starved samples. However, the precise

pathways underlying masculinization and cholesterol dynamics remain unknown. In general,

cholesterol is synthesized not only in the liver and adrenal glands but also in the gonads

[57-59]. Therefore, gaining a comprehensive understanding of the lipid metabolic mechanisms

in fish gonads, including cholesterol synthesis and transportation during sex change, may

provide valuable insights regarding the mechanisms underlying gonadal transformation. To

clarify this perspective, the remaining challenge lies in providing experimental evidence

regarding whether it is possible to control gonadal transformation by actually regulating

cholesterol levels within the body.

Conclusions

In this study, our objective was to compare the transcriptomes of sex-transitional gonads with

those of ovaries using a unified pipeline to identify common factors involved in gonadal sex

change across protogynous fishes. We successfully constructed a high-quality de novo genome

assembly applicable for genome-wide analysis in the protogynous three-spot wrasse and

acquired newly generated gonadal transcriptome data. Leveraging this information, we

conducted a comprehensive meta-analysis by integrating both public and newly derived

gonadal transcriptome data from multiple protogynous fishes spanning three orders. Through

this analysis, we identified 168 differentially expressed factors between the two gonadal states,

suggesting their potential universal role in gonadal sex change across diverse taxa of

protogynous fishes. Notably, we identified novel factors that were not previously associated

with gonadal sex change. In particular, we identified cholesterol dynamics as an important

factor regulating the sex change process. Moreover, this finding highlights the potential of

transcriptomics meta-analysis as a valuable approach to uncover overlooked mechanisms

underlying conserved biological processes including sex change. We anticipate that further

data-driven research utilizing these novel sex change-related factors will open new horizons in

sex change research.

Methods

Animals. The adult female three-spot wrasse specimens collected in the field by local

fisherman in the northern region of Okinawa, Japan were purchased and employed for genome

and RNA sequencing. These specimens were subsequently cultivated in 1000 L tanks receiving

a continuous supply of unregulated seawater, and were subjected to natural photoperiod

conditions until sampling.

The maintenance and handling of fish in this study were conducted in accordance with the

"Guidelines for Animal Experiments of the Okinawa Churashima Foundation," with the same

consideration for animal care and welfare as that for higher vertebrates, including reptiles,

birds, and mammals. Animal ethical review and approval were not required for the present

study because the guidelines stipulated that approval from the Institutional Animal Care and

Use Committee of Okinawa Churashima Foundation is required for "higher" vertebrates, and is

waived for "lower" vertebrates including fish.

Genome sequencing and assembly for the three-spot wrasse. Muscle tissue from a mature

female individual was used for genome sequencing. The specimen was randomly selected from

cultivated tank. After anesthetizing the fish with 0.05% v/v 2-phenoxyethanol (FUJIFILM

Wako Pure Chemical), the muscle tissue was dissected, immediately frozen in liquid nitrogen,

and stored at −80 °C until genomic DNA extraction. High-molecular-weight DNA was

extracted from muscle using a NucleoBond AXG column (Macherey–Nagel), followed by

purification with phenol–chloroform. The concentration of the extracted DNA was measured

using Qubit (Thermo Fisher Scientific Inc.), and its size distribution was first analyzed with a

TapeStation 4200 (Agilent Technologies) to ensure high integrity, and subsequently analyzed

via pulse-field gel electrophoresis on a CHEF DR-II system (BioRad) to ensure a size range of

20–100 kb.

To obtain long-read sequence data, a SMRT sequence library was constructed using the

SMRTbell Express Template Prep Kit 2.0 (PacBio) and sequenced on a PacBio Sequel II

system. The sequencing output was processed to generate circular consensus sequences to

obtain HiFi sequence reads. Adapter sequences were removed from the HiFi reads using

HiFiAdapterFilt [60]. The obtained HiFi sequence reads were assembled using the hifiasm

(v0.16.1) assembler [36] with default parameters. The obtained contigs were first scaffolded

with the P_RNA_scaffolder [61] and further scaffolded using Hi-C data as follows. The Hi-C

library was prepared with the in situ method using the muscle tissue of the individual used for

DNA extraction according to the iconHi-C protocol [62], employing the restriction enzymes

DpnII and HinfI, and sequenced on a HiSeq X sequencing platform (Illumina). The obtained

Hi-C read pairs, amounting to 232 million read pairs, were aligned to the HiFi sequence

contigs using the Juicer program [63], and the HiFi sequence contigs were scaffolded via the

3d-dna (version 201008) to be consistent with the chromatin contact profiles through manual

curation on JuiceBox v1.11.08 [64]. The continuity and completeness of the resulting genome

assembly were assessed using the webserver gVolante [65], in which the pipeline BUSCO

(v5.4, reference datasets: vertebrata_odb10) was implemented [37].

RNA-Seq of gonadal tissue in the three-spot wrasse. Based on previous studies [10, 45], we

triggered a sex change in three-spot wrasse n=5 females (body length; 8.0–9.3 cm, body

weight; 12.4–20.9 g) to obtain transitional gonads through treatment with an aromatase

inhibitor (AI), which exerts inhibitory effects on the enzyme responsible for estrogen synthesis.

The collected gonads were cut into small pieces, one piece was immersed in TRIzol reagent

(Invitrogen) and the rest was fixed in 4% paraformaldehyde/1× phosphate-buffered saline

(PBS) for histological observation. Total RNA in each sample was extracted promptly in

accordance with the TRIzol instruction manual. The extracted total RNA was dissolved in

RNase-free water and stored at −80 °C until RNA-Seq analysis. We selected two treated and

two control samples for RNA-Seq analysis on the basis of histological observation

(Supplementary fig. 1).

Beads with oligo (dT) were used to isolate poly(A) mRNA after the total RNA was

collected from the samples. Fragmentation buffer was added to disrupt mRNA into short

fragments. Using these short fragments as templates, a random hexamer primer was used to

synthesize the first-strand cDNA. Second-strand cDNA was synthesized using buffer, dNTPs,

Rhase H, and DNA polymerase I. Short fragments were purified using the QiaQuick PCR

extraction kit (Qiagen) and resolved with EB buffer for end reparation and adding an "A" base.

Subsequently, the short fragments were connected to sequencing adapters. Following agarose

gel electrophoresis, suitable fragments were selected as templates for PCR amplification.

Finally, the library was sequenced using an Illumina HiSeq 2000 as 2×90 bp paired-ends. The

acquired reads were subjected to the quality control procedures described below and used for

downstream expression analysis.

Retrieval and acquisition of RNA-Seq data in protogynous fishes from public resources.

To obtain RNA-Seq data from the gonads of protogynous fishes in public databases, we

accessed SRA [66] (accessed in October 2022) to search SRA data containing the following

keywords: "bony fishes (AND ovary OR gonad)." SRA data were converted into FASTQ files

using the Fasterq-dump program of the SRA Toolkit (v3.0.0).

Quality control of RNA-Seq data. To ensure the integrity of the RNA-Seq reads, trimming

and quality control procedures were performed. Specifically, we used Trim Galore (v0.6.7)

[67] with the option "--quality 20 --paired" to eliminate low-quality bases from paired-end

reads.

Read mapping against the reference genome. For species with available genomic

information, RNA-Seq reads were mapped to the reference genome sequences using HISAT2

(v2.2.1) [68] with the parameter "-q -dta -x". The resulting SAM files were sorted and

converted into BAM files using SAMtools software (v1.15.1) [69].

Generating General Transfer Format (gtf) files based on genome sequence and RNA-Seq

reads for reference gene annotation when quantifying expression values. To ensure

consistency in the use of annotation information for expression quantification in species with

available reference genomes, we employed the merge function of StringTie (v2.1.7) [40] to

generate a gtf file that served as a guide for annotations. The transcriptome assembly for each

fish was conducted using StringTie with default settings based on the mapping results (BAM

file). The resulting gtf files were then merged using StringTie with the "--merge" option,

producing a unified file as the "merged.gtf". This "merged.gtf" file contained a non-redundant

set of assembled transcripts for each species. Generated "merged.gtf" files in each species were

designated as the reference annotation files for quantifying transcript expression.

Quantification of transcripts expression and detection of differentially expressed

transcripts. When reanalyzing high-throughput data from multiple experiments, it is crucial to

consider batch effects [70]. In the present study, we therefore calculated the transcript per

million (TPM) value for each transcript and utilized the difference in TPM values between the

paired data as an indicator for identifying differentially expressed transcripts.

Transcript expression was quantified in species with available genome annotation using

StringTie, specifying the parameters as "-G merged.gtf -e". The resulting gtf file for each

sample provided the TPM values, which were used for downstream analysis.

TPM values were computed for species lacking reference genomes using Salmon

(v.1.9.0)[42] with the parameters "quant -I index -l A". To generate the transcriptome

sequences required for Salmon, we performed transcriptome assembly using the Trinity

(v2.13.1) assembler [41] with default settings. Open reading frames (ORFs) and CDS were

predicted from the transcriptome assembly using TransDecoder (v5.5.0) [38] with the same

settings as described below. The resulting gff3 file obtained from the TransDecoder run was

used to extract transcript sequences corresponding to the predicted CDS from the

transcriptome assembly. The extracted sequence set was then utilized as the reference

transcriptome after eliminating duplicate sequences using seqkit (v2.3.0) [71] with the "rmdup

-s" command.

To assess changes in transcript expression, the expression ratio was determined using TPM

values obtained from intersexual gonad and ovary transcriptomes (referred to as the IO-ratio).

The IO-ratio for each transcript was calculated using Equation (1).

$\text{IO ratio} = \log_2(\text{TPM}_{\text{intersexual}} + 0.01) - \log_2(\text{TPM}_{\text{ovary}} + 0.01)$, (1)

This calculation was performed for all pairs of intersexual gonad and ovary transcriptomes. To

convert zeros to logarithms, a value of "0.01" was added to the TPM of each transcript.

Transcripts with an IO-ratio $> \log_2(5)$ (indicating a 5-fold expression change) were classified

as "upregulated," whereas transcripts with an IO-ratio $< -\log_2(5)$ (representing a 0.2-fold

expression change) were considered "downregulated." Transcripts outside these thresholds

were deemed "unchanged." To determine differential expression between the intersexual gonad

and ovary, an IO-score was calculated for each transcript by subtracting the count of

"downregulated" pairs from that of "upregulated" pairs, as shown in Equation (2).

$\text{IO-score} = (\text{count number}_{\text{upregulated}}) - (\text{count number}_{\text{downregulated}})$, (2)

Using orthology inference as described below, the total IO-score for each transcript was

calculated by summing the IO-scores of all species. Total IO-score thresholds of 7 and −7 were

employed to identify transcripts that were differentially upregulated and downregulated,

respectively. The thresholds were arbitrarily set based on 20% above the maximum or below

the minimum values (31 or -31). All calculations for IO-ratios and IO-scores were performed

using R (v 4.2.1) [72].

Orthology inference among species. GFFread (v0.12.1) [73] was utilized along with the

generated "merged.gtf" file to extract transcript sequences from the reference genomes. For

species without reference genomes, transcript sequences were obtained via assembly using

Trinity, as described previously. TransDecoder was used to identify the coding regions and

translate them into amino acid sequences for all species. ORFs were extracted from the

transcriptome sequences using TransDecoder Longorfs with the parameters "-m 100". CDS

were then predicted using TransDecoder Predict with default settings, and the protein

sequences converted from the predicted CDS were acquired. Subsequently, a reciprocal

homology search was conducted between the predicted protein sequences of the three-spot

wrasse and those of other species, and the reciprocal best hit (RBH) was identified as the

ortholog. The program Diamond (v2.0.15) [74] was used for the RBH search using the

parameters "--ultra-sensitive -e 1e-3". To extract the orthologs shared among all species, we

consolidated the RBH results with the three-spot wrasse-predicted protein IDs.

Gene set enrichment analysis. Differentially expressed gene sets were analyzed using the

web tool Metascape (v3.5) [75], which enables gene set enrichment analysis. However,

Metascape only supports the analysis of zebrafish among teleosts. Therefore, prior to analysis,

the predicted protein sequences of the three-spot wrasse were mapped to the corresponding

zebrafish Ensembl protein IDs. To obtain orthologous information, we utilized the ggsearch36

program from the FASTA package [43] to globally align the predicted protein sequences of the

three-spot wrasse with the proteome sequences of zebrafish (Ensembl Release 108) [44].

Visualization of cholesterol and steroid hormone biosynthesis pathways and orthologs

mapping to the pathways. The hypothetical cholesterol biosynthesis and steroid hormone

biosynthesis pathway in teleost fish was illustrated using PathVisio [76]. Modifying

information from WikiPathways [77] intended for zebrafish (WP1389,

https://www.wikipathways.org/instance/WP1387) and data from the yellow perch, *Perca*

flavescens [78], facilitated the depiction of the cholesterol biosynthesis pathway. Additionally,

the steroid hormone biosynthesis pathway was constructed by adjusting information sourced

from various fish models [22, 79-82]. Moreover, through PathVisio, 7,289 candidate

sex-change factors, converted into Ensembl zebrafish gene IDs, were mapped onto the

established biosynthesis pathways.

Declarations

Ethics approval

The maintenance and handling of fish in this study were conducted in accordance with the

"Guidelines for Animal Experiments of the Okinawa Churashima Foundation," with the same

consideration for animal care and welfare as that for higher vertebrates, including reptiles,

birds, and mammals. Animal ethical review and approval were not required for the present

study because the guidelines stipulated that approval from the Institutional Animal Care and

Use Committee of Okinawa Churashima Foundation is required for "higher" vertebrates, and is

waived for "lower" vertebrates including fish.

Consent for publication

Not applicable

Availability of data

All sequencing data, including assembled sequences and raw sequence reads, were deposited

in the DNA Data Bank of Japan (DDBJ) under the umbrella BioProject accession number

PRJDB15835. The genome assembly was deposited in the DDBJ under accession numbers

BSYP01000001–BSYP01000227. Additionally, raw sequence reads were deposited in the

DDBJ under the following accession numbers: DRR466686 (PacBio HiFi reads), DRR486910

(Illumina HiC reads), and DRR483512-DRR483515 (Illumina RNA-Seq reads).

Competing interests

The authors have no conflicts of competing interest to declare that are relevant to the content

of this article.

Funding

This research was supported by the Center of Innovation for Bio-Digital Transformation

(BioDX), a program on the open innovation platform for industry-academia co-creation

(COI-NEXT), and the Japan Science and Technology Agency (JST, COI-NEXT, JPMJPF2010).

This work was partially received financial support by International Research Organization for

Advanced Science and Technology, Kumamoto University, Kumamoto, Japan.

Authors' contributions

RN and HB conceived and designed the study. MK and SK performed genome sequence and

de novo genome assembly of three-spot wrasse. RN & MN RNA-seq data production of

three-spot wrasse. RN performed RNA-seq data analysis and orthology inference. RN, MN,

SK and HB wrote a draft manuscript. All authors read and approved the final manuscript.

Acknowledgements

All authors thank Dr. Osamu Nishimura at DNA Sequencing Facility operated by Laboratory

for Phyloinformatics in RIKEN BDR for assistance in Hi-C data assessment. We thank the

Sesoko Station, Tropical Biosphere Research Center, University of the Ryukyus for allowing

us to use their research facility.

Additional information

All supplementary information is accessible in figshare repository

(https://doi.org/10.6084/m9.figshare.c.6704160)

Supplementary figure 1. Histological sections of control ovaries and transitional gonads treated

AI for 7 days in the three-spot wrasse, *Halichoeres trimaculatus*.

Supplementary figure 2. Hypothetical ovarian steroid hormones biosynthesis pathway in

protogynous fishes. The pathway was visualized using PathVisio (https://pathvisio.org),

referencing several fish cholesterol synthesis pathways. Gene products are represented in black

text, while metabolites are in blue text. Gene product annotations follow Ensembl Zebrafish

gene symbols. Red-filled boxes represent gene products with confirmed upregulated

expression in this study, while blue-filled boxes denote gene products with downregulated

expression. Open boxes represent gene products with no expression variation. Light

green-filled boxes represent gene products not included among the 7289 factors but confirmed

to be present in at least the three-spot wrasse transcriptome. Dark gray-filled boxes indicate

gene products that could not be identified within the three-spot wrasse transcriptome. Circles

extending from gene products represent catalytic reactions, while black arrows indicate the

conversion of metabolites.

Supplementary table 1

Supplementary table 2

Supplementary table 3

Supplementary table 4

References

1.   Atz JW: **Intersexuality in fishes**. In: *Intersexuality in Vertebrates Including Man*. Edited by
Armstrong CN, Marshall AJ. London: Academic press; 1964: 145–232.

2.   Kuwamura T, Sunobe T, Sakai Y, Kadota T, Sawada K: **Hermaphroditism in fishes: an**
annotated list of species, phylogeny, and mating system. *Ichthyol Res* 2020, **67**: 341–360.

3.   Avise JC, Mank JE: **Evolutionary perspectives on hermaphroditism in fishes**. *Sex Dev* 2009,
3: 152–163.

4.   Erisman BE, Petersen CW, Hastings PA, Warner RR: **Phylogenetic perspectives on the**
evolution of functional hermaphroditism in teleost fishes. *Integr Comp Biol* 2013, **53**:
736–754.

5.   Lo Nostro F, Grier H, Andreone L, Guerrero GA: **Involvement of the gonadal germinal**
epithelium during sex reversal and seasonal testicular cycling in the protogynous swamp
eel, *Synbranchus marmoratus* **Bloch 1795 (Teleostei, Synbranchidae)**. *J Morphol* 2003, **257**:
107–126.

6.   Nakamura M, Hourigan TF, Yamauchi K, Nagahama Y, Grau EG: **Histological and**
ultrastructural evidence for the role of gonadal steroid hormones in sex change in the
protogynous wrasse *Thalassoma duperrey*. *Environ Biol Fishes* 1989, **24**: 117–136.

7.   Ohta K, Hirano M, Mine T, Mizutani H, Yamaguchi A, Matsuyama M: **Body color change and**
serum steroid hormone levels throughout the process of sex change in the adult wrasse,
Pseudolabrus sieboldi. *Mar Biol* 2008, **153**: 843–852.

8.   Muncaster S, Norberg B, Andersson E: **Natural sex change in the temperate protogynous**
Ballan wrasse *Labrus bergylta*. *J Fish Biol* 2013, **82**: 1858–1870.

9.   Bhandari RK, Komuro H, Nakamura S, Higa M, Nakamura M: **Gonadal restructuring and**
correlative steroid hormone profiles during natural sex change in protogynous
honeycomb grouper (*Epinephelus merra* **)**. *Zoolog Sci* 2003, **20**: 1399–1404.

10.   Horiguchi R, Nozu R, Hirai T, Kobayashi Y, Nagahama Y, Nakamura M: **Characterization of**
gonadal soma-derived factor expression during sex change in the protogynous wrasse,
Halichoeres trimaculatus. *Dev Dyn* 2013, **242**: 388–399.

11.   Nozu R, Horiguchi R, Kobayashi Y, Nakamura M: **Expression profile of doublesex/male**
abnormal-3-related transcription factor-1 during gonadal sex change in the protogynous
wrasse, *Halichoeres trimaculatus*. *Mol Reprod Dev* 2015, **82**: 859–866.

12.   Horiguchi R, Nozu R, Hirai T, Kobayashi Y, Nakamura M: **Expression patterns of sex**
differentiation-related genes during gonadal sex change in the protogynous wrasse,

Halichoeres trimaculatus. *Gen Comp Endocrinol* 2018, **257**: 67–73.

13. Kobayashi Y, Horiguchi R, Nozu R, Nakamura M: **Expression and localization of forkhead transcriptional factor 2 (Foxl2) in the gonads of protogynous wrasse, *Halichoeres trimaculatus*. *Biol Sex Differ* 2010, **1**: 3.

14. Thomas JT, Todd EV, Muncaster S, Lokman PM, Damsteegt EL, Liu H, Soyano K, Gléonnec F, Lamm MS, Godwin JR *et al*: **Conservation and diversity in expression of candidate genes regulating socially-induced female-male sex change in wrasses**. *PeerJ* 2019, **7**: e7032.

15. Miyake Y, Sakai Y, Kuniyoshi H: **Molecular cloning and expression profile of sex-specific genes, Figla and Dmrt1, in the protogynous hermaphroditic fish, *Halichoeres poecilopterus*. *Zoolog Sci* 2012, **29**: 690–701.

16. Senthilkumaran B, Chi W, Gao Y, Hu Q, Guo W, Li D: **Genome-wide analysis of brain and gonad transcripts reveals changes of key sex reversal-related genes expression and signaling pathways in three stages of *Monopterus albus***. *PloS one* 2017, **12**: e0173974.

17. Todd EV, Ortega-Recalde O, Liu H, Lamm MS, Rutherford KM, Cross H, Black MA, Kardailsky O, Marshall Graves JA, Hore TA *et al*: **Stress, novel sex genes, and epigenetic reprogramming orchestrate socially controlled sex change**. *Sci Adv* 2019, **5**: eaaw7006.

18. He Z, Deng F, Yang D, He Z, Hu J, Ma Z, Zhang Q, He J, Ye L, Chen H *et al*: **Crosstalk between sex-related genes and apoptosis signaling reveals molecular insights into sex change in a protogynous hermaphroditic teleost fish, ricefield eel *Monopterus albus***. *Aquaculture* 2022, **552**: 737918.

19. Fan M, Yang W, Zhang W, Zhang L: **The ontogenic gonadal transcriptomes provide insights into sex change in the ricefield eel *Monopterus albus***. *BMC Zool* 2022, **7**: 56.

20. Wu X, Yang Y, Zhong C, Guo Y, Wei T, Li S, Lin H, Liu X: **Integration of ATAC-seq and RNA-seq unravels chromatin accessibility during sex reversal in Orange-spotted grouper (*Epinephelus coioides*)**. *Int J Mol Sci* 2020, **21**: 2800.

21. Zhang Y, Zhang S, Liu Z, Zhang L, Zhang W: **Epigenetic modifications during sex change repress gonadotropin stimulation of cyp19a1a in a teleost ricefield eel (*Monopterus albus*)**. *Endocrinology* 2013, **154**: 2881–2890.

22. Li GL, Liu XC, Zhang Y, Lin HR: **Gonadal development, aromatase activity and P450 aromatase gene expression during sex inversion of protogynous red-spotted grouper *Epinephelus akaara* (Temminck and Schlegel) after implantation of the aromatase inhibitor, fadrozole**. *Aquac Res* 2006, **37**: 484–491.

23. Wang DS, Kobayashi T, Zhou LY, Paul-Prasanth B, Ijiri S, Sakai F, Okubo K, Morohashi K, Nagahama Y: **Foxl2 up-regulates aromatase gene transcription in a female-specific manner by binding to the promoter as well as interacting with ad4 binding protein/steroidogenic factor 1**. *Mol Endocrinol* 2007, **21**: 712–725.

24. Zhang C, He Q, Cheng H, Li L, Ruan X, Duan X, Huang F, Yang H, Zhang H, Shi H *et al*: **Transcription factors foxl2 and foxl3 regulate cyp19a1a and cyp11b in orange-spotted grouper (*Epinephelus coioides*)**. *Aquac Rep* 2022, **25**: 101243.

25. Shibata Y, Paul-Prasanth B, Suzuki A, Usami T, Nakamoto M, Matsuda M, Nagahama Y:

595   **Expression of gonadal soma derived factor (GSDF) is spatially and temporally correlated**

596   **with early testicular differentiation in medaka**. *Gene Expr Patterns* 2010, **10**: 283–289.

597   26.   Zhang X, Guan G, Li M, Zhu F, Liu Q, Naruse K, Herpin A, Nagahama Y, Li J, Hong Y:

598   **Autosomal gsdf acts as a male sex initiator in the fish medaka**. *Sci Rep* 2016, **6**: 19738.

599   27.   Kobayashi T, Kajiura-Kobayashi H, Guan G, Nagahama Y: **Sexual dimorphic expression of**

600   **DMRT1 and Sox9a during gonadal differentiation and hormone-induced sex reversal in**

601   **the teleost fish Nile tilapia (*Oreochromis niloticus*)**. *Dev Dyn* 2008, **237**: 297–306.

602   28.   Rodríguez-Marí A, Yan Y-L, BreMiller RA, Wilson C, Cañestro C, Postlethwait JH:

603   **Characterization and expression pattern of zebrafish anti-Müllerian hormone (amh)**

604   **relative to sox9a, sox9b, and cyp19a1a, during gonad development**. *Gene Expr Patterns*

605   2005, **5**: 655–667.

606   29.   Wang XG, Orban L: **Anti-Müllerian hormone and 11 β-hydroxylase show reciprocal**

607   **expression to that of aromatase in the transforming gonad of zebrafish males**. *Dev Dyn*

608   2007, **236**: 1329–1338.

609   30.   Skaar KS, Nobrega RH, Magaraki A, Olsen LC, Schulz RW, Male R: **Proteolytically activated,**

610   **recombinant anti-mullerian hormone inhibits androgen secretion, proliferation, and**

611   **differentiation of spermatogonia in adult zebrafish testis organ cultures**. *Endocrinology*

612   2011, **152**: 3527–3540.

613   31.   Webster KA, Schach U, Ordaz A, Steinfeld JS, Draper BW, Siegfried KR: **Dmrt1 is necessary**

614   **for male sexual development in zebrafish**. *Dev Biol* 2017, **422**: 33–46.

615   32.   Zhu Y, Wang C, Chen X, Guan G: **Identification of gonadal soma-derived factor**

616   **involvement in *Monopterus albus* (protogynous rice field eel) sex change**. *Mol Biol Rep*

617   2016, **43**: 629–637.

618   33.   Bono H: **Meta-analysis of oxidative transcriptomes in insects**. *Antioxidants* 2021, **10**: 345.

619   34.   Rahman MR, Petralia MC, Ciurleo R, Bramanti A, Fagone P, Shahjaman M, Wu L, Sun Y,

620   Turanli B, Arga KY *et al*: **Comprehensive analysis of RNA-Seq gene expression profiling of**

621   **brain transcriptomes reveals novel genes, regulators, and pathways in autism spectrum**

622   **disorder**. *Brain Sci* 2020, **10**: 747.

623   35.   Toga K, Bono H: **Meta-analysis of public RNA sequencing data revealed potential key**

624   **genes associated with reproductive division of labor in social hymenoptera and termites**.

625   *Int J Mol Sci* 2023, **24**: 8353.

626   36.   Cheng H, Concepcion GT, Feng X, Zhang H, Li H: **Haplotype-resolved de novo assembly**

627   **using phased assembly graphs with hifiasm**. *Nat Methods* 2021, **18**: 170–175.

628   37.   Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM, Kelley J: **BUSCO update:**

629   **novel and streamlined workflows along with broader and deeper phylogenetic coverage**

630   **for scoring of eukaryotic, prokaryotic, and viral genomes**. *Mol Biol Evol* 2021, **38**:

631   4647–4654.

632   38.   **TransDecoder (v5.5.0)** [https://github.com/TransDecoder/TransDecoder]

633   39.   Ge H, Lin K, Shen M, Wu S, Wang Y, Zhang Z, Wang Z, Zhang Y, Huang Z, Zhou C *et al*: ***De***

634   ***novo* assembly of a chromosome-level reference genome of red-spotted grouper**

(*Epinephelus akaara*) **using nanopore sequencing and Hi-C**. *Mol Ecol Resour* 2019, **19**:
1461–1469.

40. Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL: **StringTie enables**
improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol* 2015, **33**:
290–295.

41. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,
Raychowdhury R, Zeng Q *et al*: **Full-length transcriptome assembly from RNA-Seq data**
without a reference genome. *Nat Biotechnol* 2011, **29**: 644–652.

42. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C: **Salmon provides fast and bias-aware**
quantification of transcript expression. *Nat Methods* 2017, **14**: 417–419.

43. Pearson WR, Lipman DJ: **Improved tools for biological sequence comparison**. *Proc Natl*
Acad Sci USA 1988, **85**: 2444–2448.

44. Cunningham F, Allen JE, Allen J, Alvarez-Jarreta J, Amode M R, Armean Irina M,
Austine-Orimoloye O, Azov Andrey G, Barnes I, Bennett R *et al*: **Ensembl 2022**. *Nucleic*
Acids Res 2022, **50**: D988–D995.

45. Nozu R, Kojima Y, Nakamura M: **Short term treatment with aromatase inhibitor induces**
sex change in the protogynous wrasse, *Halichoeres trimaculatus*. *Gen Comp Endocrinol*
2009, **161**: 360–364.

46. Kroon FJ, Liley NR: **The role of steroid hormones in protogynous sex change in the**
Blackeye goby, *Coryphopterus nicholsii* **(Teleostei: Gobiidae)**. *Gen Comp Endocrinol* 2000,
118: 273–283.

47. Hasegawa S, Noda K, Maeda A, Matsuoka M, Yamasaki M, Fukui T: **Acetoacetyl-CoA**
synthetase, a ketone body-utilizing enzyme, is controlled by SREBP-2 and affects serum
cholesterol levels. *Mol Genet Metab* 2012, **107**: 553–560.

48. Wu D-M, He Z, Chen T, Liu Y, Ma L-P, Ping J: **DNA hypermethylation of acetoacetyl-CoA**
synthetase contributes to inhibited cholesterol supply and steroidogenesis in fetal rat
adrenals under prenatal nicotine exposure. *Toxicology* 2016, **340**: 43–52.

49. Gaudet P, Livstone MS, Lewis SE, Thomas PD: **Phylogenetic-based propagation of**
functional annotations within the Gene Ontology consortium. *Brief Bioinform* 2011, **12**:
449–462.

50. Blue ML, Williams DL, Zucker S, Khan SA, Blum CB: **Apolipoprotein E synthesis in human**
kidney, adrenal gland, and liver. *Proc Natl Acad Sci USA* 1983, **80**: 283–287.

51. Reyland ME, Gwynne JT, Forgez P, Prack MM, Williams DL: **Expression of the human**
apolipoprotein E gene suppresses steroidogenesis in mouse Y1 adrenal cells. *Proc Natl*
Acad Sci USA 1991, **88**: 2375–2379.

52. Thorngate FE, Strockbine PA, Erickson SK, Williams DL: **Altered adrenal gland cholesterol**
metabolism in the apoE-deficient mouse. *J Lipid Res* 2002, **43**: 1920–1926.

53. Payne AH, Hales DB: **Overview of steroidogenic enzymes in the pathway from cholesterol**
to active steroid hormones. *Endocr Rev* 2004, **25**: 947–970.

54. Pikuleva IA: **Cholesterol-metabolizing cytochromes P450**. *Drug Metab Dispos* 2006, **34**:

513–520.

55.   Miller WL: **Steroidogenic enzymes**. In: *Disorders of the Human Adrenal Cortex*. Edited by
Flück CE, Miller WL. Basel: S.Karger AG; 2008: 1–18.

56.   Sakae Y, Oikawa A, Sugiura Y, Mita M, Nakamura S, Nishimura T, Suematsu M, Tanaka M:
Starvation causes female-to-male sex reversal through lipid metabolism in the teleost fish,
medaka (*Oryzias latipes*). *Biol Open* 2020, **9**: bio050054.

57.   Gwynne JT, Strauss JF: **The role of lipoproteins in steroidogenesis and cholesterol**
metabolism in steroidogenic glands. *Endocr Rev* 1982, **3**: 299–329.

58.   Sharpe RL, Drolet M, MacLatchy DL: **Investigation of de novo cholesterol synthetic**
capacity in the gonads of goldfish (*Carassius auratus*) exposed to the phytosterol
beta-sitosterol. *Reprod Biol Endocrinol* 2006, **4**: 60.

59.   Bera A, Chadha NK, Dasgupta S, Chakravarty S, Sawant PB: **Hypoxia-mediated inhibition of**
cholesterol synthesis leads to disruption of nocturnal sex steroidogenesis in the gonad of
koi carp, *Cyprinus carpio*. *Fish Physiol Biochem* 2020, **46**: 2421–2435.

60.   Sim SB, Corpuz RL, Simmonds TJ, Geib SM: **HiFiAdapterFilt, a memory efficient read**
processing pipeline, prevents occurrence of adapter sequence in PacBio HiFi reads and
their negative impacts on genome assembly. *BMC Genomics* 2022, **23**: 157.

61.   Zhu B-H, Xiao J, Xue W, Xu G-C, Sun M-Y, Li J-T: **P_RNA_scaffolder: a fast and accurate**
genome scaffolder using paired-end RNA-sequencing reads. *BMC Genomics* 2018, **19**: 175.

62.   Kadota M, Nishimura O, Miura H, Tanaka K, Hiratani I, Kuraku S: **Multifaceted Hi-C**
benchmarking: what makes a difference in chromosome-scale genome scaffolding?
Gigascience 2020, **9**: giz158.

63.   Durand NC, Shamim MS, Machol I, Rao SSP, Huntley MH, Lander ES, Aiden EL: **Juicer**
provides a one-click system for analyzing loop-resolution Hi-C experiments. *Cell Syst* 2016,
3: 95–98.

64.   Dudchenko O, Shamim MS, Batra SS, Durand NC, Musial NT, Mostofa R, Pham M, Hilaire
BGS, Yao W, Stamenova E *et al*: **The Juicebox assembly tools module facilitates *de novo***
assembly of mammalian genomes with chromosome-length scaffolds for under $1000.
bioRxiv 2018: 254797.

65.   Nishimura O, Hara Y, Kuraku S, Hancock J: **gVolante for standardizing completeness**
assessment of genome and transcriptome assemblies. *Bioinformatics* 2017, **33**: 3635–3637.

66.   Kodama Y, Shumway M, Leinonen R: **The sequence read archive: explosive growth of**
sequencing data. *Nucleic Acids Res* 2011, **40**: D54–D56.

67.   **Trim Galore** [https://github.com/FelixKrueger/TrimGalore]

68.   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL: **Graph-based genome alignment and**
genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 2019, **37**: 907–915.

69.   Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,
McCarthy SA, Davies RM *et al*: **Twelve years of SAMtools and BCFtools**. *Gigascience* 2021,
10: giab008.

70.   Leek JT, Scharpf RB, Bravo HC, Simcha D, Langmead B, Johnson WE, Geman D, Baggerly K,

Irizarry RA: **Tackling the widespread and critical impact of batch effects in high-throughput data**. *Nat Rev Genet* 2010, **11**: 733–739.

71. Zou Q, Shen W, Le S, Li Y, Hu F: **SeqKit: a cross-platform and ultrafast toolkit for FASTA/Q file manipulation**. *PloS one* 2016, **11**: e0163962.

72. **R: A language and environment for statistical computing.** [URL https://www.R-project.org/]

73. Pertea G, Pertea M: **GFF Utilities: GffRead and GffCompare**. *F1000Res* 2020, **9**: 304.

74. Buchfink B, Xie C, Huson DH: **Fast and sensitive protein alignment using DIAMOND**. *Nat Methods* 2014, **12**: 59–60.

75. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, Benner C, Chanda SK: **Metascape provides a biologist-oriented resource for the analysis of systems-level datasets**. *Nat Commun* 2019, **10**: 1523.

76. Murphy RF, Kutmon M, van Iersel MP, Bohler A, Kelder T, Nunes N, Pico AR, Evelo CT: **PathVisio 3: An Extendable Pathway Analysis Toolbox**. *PLoS Comput Biol* 2015, **11**: e1004085.

77. Pico AR, Kelder T, van Iersel MP, Hanspers K, Conklin BR, Evelo C: **WikiPathways: Pathway Editing for the People**. *PLoS Biol* 2008, **6**: e184.

78. Kemski MM, Rappleye CA, Dabrowski K, Bruno RS, Wick M: **Transcriptomic response to soybean meal-based diets as the first formulated feed in juvenile yellow perch (*Perca flavescens*)**. *Sci Rep* 2020, **10**: 3998.

79. Liu Y, Kossack ME, McFaul ME, Christensen LN, Siebert S, Wyatt SR, Kamei CN, Horst S, Arroyo N, Drummond IA *et al*: **Single-cell transcriptome reveals insights into the development and function of the zebrafish ovary**. *Elife* 2022, **11**: e76014.

80. Craft JA, Bain PA, Papanicolaou A, Kumar A: **Identification of Putative Nuclear Receptors and Steroidogenic Enzymes in Murray-Darling Rainbowfish (*Melanotaenia fluviatilis*) Using RNA-Seq and De Novo Transcriptome Assembly**. *PloS one* 2015, **10**: e0142636.

81. Nagahama Y, Chakraborty T, Paul-Prasanth B, Ohta K, Nakamura M: **Sex determination, gonadal sex differentiation, and plasticity in vertebrate species**. *Physiol Rev* 2021, **101**: 1237–1308.

82. Fernandino JI, Hattori RS, Moreno Acosta OD, Strussmann CA, Somoza GM: **Environmental stress-induced testis differentiation: androgen as a by-product of cortisol inactivation**. *Gen Comp Endocrinol* 2013, **192**: 36–44.

Figure legends

Fig. 1. Schematic diagram of the analysis pipeline in this study. The consistent line style of the arrows indicates the utilization of common tools for conducting data processing.

Fig. 2. Scatter plot of the IO-score and gene set enrichment analysis of up- and downregulated genes. a Scatterplot of total IO-scores for the 7,289 factors in descending order. Red circles indicate 99 upregulated factors; blue circles indicate 69 downregulated factors. Gray circles (which have the appearance of lines because they overlap each other) indicate unchanged factors ($6 \leq$ total IO-score ≤ -6). **b,c** Results of enrichment analysis for 99 upregulated (**b**) and 69 downregulated (**c**) factors

Fig. 3. Composition of total IO-scores for upregulated factors (n=99). Numbers in parentheses indicate the number of RNA-Seq data pairs for each species used in this study. Asterisk indicates a gene for which the total IO-score is derived from a single species.

Fig. 4. Composition of total IO-scores for downregulated factors (n=69). Numbers in parentheses indicate the number of RNA-Seq data pairs for each species used in this study. Asterisk indicates a gene for which the total IO-score is derived from a single species.

Fig. 5. Hypothetical ovarian cholesterol biosynthesis pathway in protogynous fishes. The pathway was visualized using PathVisio (https://pathvisio.org), referencing several fish cholesterol synthesis pathways. Gene products are represented in black text, while metabolites are in blue text. Gene product annotations follow Ensembl Zebrafish gene symbols. Red-filled boxes represent gene products with confirmed upregulated expression in this study, while blue-filled boxes denote gene products with downregulated expression. Open boxes represent gene products with no expression variation. Light green-filled boxes represent gene products not included among the 7289 factors but confirmed to be present in at least the three-spot wrasse transcriptome. Dark gray-filled boxes indicate gene products that could not be identified within the three-spot wrasse transcriptome. Circles extending from gene products represent catalytic reactions, while black arrows indicate the conversion of metabolites. The

778   dashed arrows indicate indirect pathways summarizing multi-step processes. The dashed

779   arrows indicate indirect pathways summarizing multi-step processes.

783   **Table 1.** Statistics of genome assembly in Three-spot wrasse (fHalTri1.1) and other Labidae

784   fishes listed in NCBI Genomes.

	fHalTri1.1	*Cheilinus undulatus* (humphead wrasse)	*Notolabrus celidotus* (New Zealand spotty)	*Labrus bergylta* (ballan wrasse)	*Semicossyphus pulcher* (California sheephead)	*Tautogolabrus adspersus* (cunner)
Assembly accession	BSYP01000001–BSYP01000227	GCF_018320785.1	GCF_009762535.1	GCF_900080235.1	GCA_022749685.1	GCA_020745685.1
Total seq length (Mb)	849.9	1173.4	846.7	805.5	794.1	723.6
Max length (Mb)	44.2	59.6	42.4	9.1	38.5	37.1
Num of scaffolds	227	45	467	13,466	178	347
Scaffold N50 (Mb)	37.9	51.5	37.1	0.8	32.1	31.7
Scaffold L50	11	11	11	252	12	11
Num of seqs > 1 Mb	25	24	24	197	35	24
% of seqs > 1 Mb	97.3	99.7	97.6	43.9	97.1	99.9
Num of gaps	117	284	1,133	257	8	283
Complete BUSCO (%)	98.7	99.0	97.1	98.9	98.9	98.1

799   **Table 2.** Statistics of orthology relationship among species.

Common name	Scientific name	Order	Reference genome	*Num of transcripts seqs obtained based on the RNA-seq data from gonadal tissue	Num of protein seqs predicted by TransDecoder	Num of RBH (vs Three-spot wrasse)
Three-spot wrasse	*Halichoeres trimaculatus*	Labriformes	Assembly in this study	27,573	24,965	24,965
Bluehead wrasse	*Thalassoma bifasciatum*	Labriformes	GCA_008086565.1	57,699	43,195	13,871
Hong Kong grouper	*Epinephelus akaara*	Perciformes	Ge et al., 2019	26,507	24,752	11,619
Orange-spotted grouper	*Epinephelus coioides*	Perciformes	GCA_900536245.1	30,867	34,191	12,507
Black seabass	*Centropristis striata*	Perciformes	NA	** 95,941	95,941	12,591
Asian swamp eel	*Monopterus albus*	Synbranchiformes	GCF_001952655.1	53,472	50,693	12,765
Zig-zag eel	*Mastacembelus armatus*	Synbranchiformes	GCF_900324485.2	48,724	40,908	13,095

800   * Based on the annotation file (merged gtf file), the extracted transcripts sequences from the

801   genome by gffread

802   ** Among transcriptome sequences (594,726 seqs) assembled by Trinity, transcripts sequences

803   corresponding to coding seqences predicted via TransDecoder

817   **Table 3.** List of genes that contributed to GO term enrichment.

Up/Down	GO ID	Category	GO term	*Genes
Up	GO:0007129	Biological Process	homologous chromosome pairing at meiosis	*hormad1, terb2, si:ch211-199g17.9*
Up	GO:0007369	Biological Process	gastrulation	*zeb1b, tmem237a, col12a1b, pfn1, enpp2*
Up	GO:1902531	Biological Process	regulation of intracellular signal transduction	*cep55l, psd3l, arhgef18b, ccl19a.1, sh3bp1, enpp2, pik3r6b*
Up	GO:0005929	Cellular Component	cilium	*septin6, prom1a, tmem237a, si:ch211-66i15.4, flr*
Up	GO:0007281	Biological Process	germ cell development	*hormad1, gsdf, sycp3*
Down	GO:0010817	Biological Process	regulation of hormone levels	*cyp19a1a, aacs, hsd17b1, ece1*
Down	GO:0006413	Biological Process	translational initiation	*eif4g2b, eif4eb, eif4e1c*
Down	GO:0019752	Biological Process	carboxylic acid metabolic process	*lpl, ldha, aacs, amacr, acaca, dars2*
Down	GO:0008202	Biological Process	steroid metabolic process	*hsd17b1, amacr, sdr42e1*
Down	GO:1904888	Biological Process	cranial skeletal system development	*nog3, cep55l, cyldb, ece1*
Down	GO:1901362	Biological Process	organic cyclic compound biosynthetic process	*tbx5a, hsd17b1, alad, acaca, sdr42e1, adcy6a*

818   * Gene symbols follow the zebrafish gene nomenclature in Ensembl.

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4.

Fig. 5.