Quantifying mutational synergy using computational models predicts survival in

haematological cancers

Richard Norris [1], John Jones[1], Erika Mancini[2], Timothy Chevassut[1], Chris Pepper[1], Andrea
Pepper[1], Simon Mitchell[1*]

[1]Brighton and Sussex Medical School, Department of Clinical and Experimental Medicine,
Brighton, United Kingdom

[2]School of Life Sciences, University of Sussex, Brighton, United Kingdom

* Correspondence: S.A.Mitchell@bsms.ac.uk

Running Title: Mutational synergy predicts survival

Abstract

Genetic heterogeneity and co-occurring driver mutations contribute to poor clinical outcomes in

cancer. However, the impact of multiple mutations on complex signalling networks is not easily

predicted. We found that, by placing mutations into their cellular context, multi-scale agent-

based mathematical models could predict how genetic events combine in haematological

malignancies. Simulations of lymphoma and myeloma predicted co-occurring mutations

synergised to increase tumour cell expansion beyond what would be expected from the impact

of the individual mutations alone. Mutational synergy between *MYC* and *BCL2* was consistent

with the more aggressive disease course of patients with double-hit lymphoma, and mutational

synergy between MCL1 and CKS1B was predictive of outcome in patients with gain 1q multiple

myeloma. Incorporating patient-specific mutational profiles into personalised models of

lymphoma patients with the worst clinical outcomes revealed a correlation between simulated

mutational synergy and overall survival, which outperformed widely used classifications of

lymphoma informed by gene expression or mutational data alone. Our results demonstrated that

mutational synergy scores enabled prediction of the impact of co-occurring mutations and may

improve personalised prognostic predictions.

Keywords: Cancer / Computational Biology / DLBCL / Myeloma / Systems Biology

Introduction

Mutational heterogeneity represents a major barrier to understanding the molecular

mechanisms responsible for tumour progression, and to the development of novel treatment

strategies. With the advent of whole exome sequencing (WES) many malignancies, including B-

cell malignancies such as Diffuse Large B-Cell Lymphoma (DLBCL) and Multiple Myeloma

(MM), have become characterised by interpatient mutational heterogeneity and within-patient

mutational complexity (1, 2). This factor contributes to variation in response to current

treatments such that first line therapy for DLBCL is ineffective for around 40% of patients, while

almost all MM patients will eventually relapse. The most aggressive malignancies often contain

genetic aberrations affecting multiple genes. 'Double hit' (DH) DLBCL, featuring overexpression

of *MYC* and *BCL2* (or *BCL6*)*,* is among the most aggressive lymphoid malignancies, with very

poor outcomes for these patients (3, 4). The prognostic implications of DH lymphomas were the

basis for the 2017 update on DLBCL by the World Health Organization. Increased expression of

MYC, which codes for the transcription factor Myc (also known as c-Myc), is a significant factor

in the growth of multiple tumour types (5), and mutations affecting *MYC* occur in approximately

33% of DLBCL patients (6). Aberrant expression of *BCL2* is anti-apoptotic (7), and as such the

genes defining DH lymphoma represent archetypal examples of two of the hallmarks of cancer:

sustained proliferation and resistance to cell death (8). *BCL6* encodes a transcriptional

repressor expressed in healthy germinal centre B-cells. Like *BCL2*, over-expression of *BCL6*

can prevent apoptosis in DLBCL (9).

Initial attempts to characterise the heterogeneity of DLBCL utilised gene expression profiling

that split cases into subgroups based on their putative cell of origin (germinal centre – GC, or

activated B cell – ABC) (10). Subsequent studies leveraged WES to identify 5 or more

prognostically informative clusters of patients based on their mutational profiles (11-13).

LymphGen identified a DH gene expression signature in the *EZB-MYC*[+] genetic cluster,

characterised by *BCL2* translocation in 72% of cases, but only bearing genetic abnormalities in

MYC in 38% of cases (13). Double hit gene expression signatures have also been identified in

tumours lacking rearrangements of *MYC* and/or *BCL2* (14). Overall, fewer than 10% of DLBCL

cases are true DH lymphoma, yet 30-40% of DLBCL patients relapse following frontline

treatment and have a poor prognosis (15). Comparisons can be made in MM, where specific

genetic events are known to confer a more aggressive phenotype. An example is gain of

chromosome 1q, which occurs in 30-50% of tumours and is associated with poorer outcome

(16, 17). However, even in the absence of known high-risk genetic changes, around 70% of

patients will succumb to the disease within 10 years. Taken together these data demonstrate

limits to our ability to identify combinations of mutations that confer aggressive disease and poor

prognosis. Importantly, clustering patients by mutational burden ignores the context of the

complex gene regulatory signalling networks involving mutated genes and the gene products

they encode.

Computational models of B-cell fates have provided insight into healthy immune responses, with

predictions of cell proliferation and survival validated by laboratory experiments (18-21). Here,

we used mechanistic models to simulate how mutations combine in B-cell malignancies. We

quantified the emergent behaviour resulting from combinations of mutations impacting multiple

signalling networks by simulating individual mutations and co-occurring mutations in the same

cell. We developed a novel method to quantify the extent to which mutations combine to confer

a worse prognosis than expected based on the impact of each individual mutation in isolation.

Using WES data to create individual virtual DLBCL patients, we found that quantification of

mutational synergy predicted overall survival better than clustering (of WES data) or gene

expression-based classification. To further validate our model in an alternative B cell malignancy

we modelled the impact of gain(1q) MM. Chromosome 1q is commonly amplified in MM, but the

impact of putative driver genes in this chromosomal region on prognosis is not well understood.

We computationally identify a gene-dosage effect, with increasing copies of chromosome 1q

correlating with increased mutational synergy, a prediction that aligns with outcomes in clinical

trials. Taken together, we demonstrated that "multi-hit" haematological malignancies, with poor

prognosis resulting from co-occurring driver mutations, can be identified by incorporating

mutational data into systems biology modelling to quantify mutational synergy.

Results

Simulating individual mutations is insufficient to explain aggressive disease.

Computational systems biology provides a unique opportunity to predict the impact of mutations

on individual molecular networks in isolation, as it is challenging to isolate the molecular

machinery controlling proliferation and apoptosis in cell-based systems. Initially, we sought to

model the effect of gain-of-function mutations in *MYC* and *BCL2* on cell cycle and apoptosis

signalling pathways, prior to investigating the effect of combining the mutations. Given the broad

impact of c-Myc, modelling its actions is challenging. However, previous modelling work was

able to recapitulate both single cell and bulk experimental measurements by encoding c-Myc as

acting on B-cell growth in cell cycle models (18, 19). Modelling the function of the Bcl2 family of

proteins meanwhile was centred around programmed cell death, and Bcl2 itself an inhibitor of

apoptosis (22).

Simulating the impact of overexpression of *MYC* predicted a modest impact on the timing of cell

cycle phases and overall time to divide (Fig 1A). A sensitivity analysis also predicted that

increased *MYC* expression has a small effect on cell cycle time relative to expression changes

of other genes in the pathway (Fig S1). This result recapitulated experimental tracking of

division times in B-cells transfected with c-Myc and cellular proliferation measurement in

lymphocytes from Eµ-Myc mice (23). In simulations of increasing levels of *MYC* expression, the

decrease in total cell cycle time was primarily due to a shortening of G1 phase. A dose-

dependent effect was seen with increasing *MYC* expression up to the equivalent of 5 extra

copies, after which no further effect is seen (Fig 1A, Fig S2). Simulating the impact of *BCL2*

overexpression, in a computational model of apoptosis (24), predicted a dose-dependent delay

of poly ADP-ribose polymerase (PARP) cleavage up to the equivalent of 5 extra copies of *BCL2*,

beyond which the cells survive the full length of the simulation (Fig 1B). These simulations

represented an average B-cell using published parameters, but it is not clear that mutations will

affect each cell equally within a heterogeneous population of individual B-cells.

Simulating a population of molecularly heterogeneous individual B-cells, by incorporating cell-to-

cell heterogeneity as described previously (18), resulted in cell cycle timings that recapitulated

the range and distribution of cell cycle duration and cell death timings measured by time-lapse

microscopy in murine B lymphocytes (25) (Fig 1C and D). Increased *MYC* expression was

predicted to have a minor effect (decreasing total cell cycle time) in all cells (Fig 1C). These

simulations also predicted a small population of cells (~80/1000) to be in cell cycle arrest in the

WT simulation. Simulated cells in this state did not progress towards a constant cell cycle time

(limit cycle), but rather their cell mass fell due to attempts to proliferate faster than they could

gain mass, which lead to cell cycle arrest. Concentrations of the cell cycle inhibitor p27, and

complexes containing p27, were predicted to be upregulated in these arrested cells (Fig S3),

which corresponds with higher p27 levels in experimental measurements of cell cycle arrest

(24). *MYC* overexpression was predicted to reverse the cell cycle arrest in >80% of these cells,

and strikingly they become the most rapidly proliferating cells (Fig 1C). The number of cells

switched was proportional to the level of *MYC* expression (Fig S4). *MYC* is amongst the most

differentially expressed genes in quiescence and reverses quiescence in haematopoietic stem

cells (26, 27), therefore it is plausible that this effect may also be present in B-cells. The impact

of *BCL2* over-expression was predicted to proportionally increase with the survival time of the

wild-type cell, with the largest Impact in cells molecularly predisposed for long survival (Fig 1E

and F).

As the fates of cell division and apoptosis are mutually exclusive and competing fates in

individual cells (dead cells can't divide, while cells that survive may divide again) (18, 28), how

these two fates combine in individual cells with DH lymphoma is unclear. To gain a better

understanding of this, simulating these signalling networks simultaneously in individual cells is

required. If within a DH cell population there exists individual cells that concurrently have the

shortest cell cycle times (Fig 1C) and the longest survival times (Fig 1E), then the mutations are

likely to combine deleteriously. However, if the most rapidly cycling cells are also the most

rapidly dying cells, then the combined effect of the mutations is unlikely to be greater than

expected from their individual impact.

Multi-scale modelling enables quantification of mutational synergy.

To predict how the DH mutations combine in single cells, we employed multi-scale, agent-based

models of B-cells, which enable simulation of multiple signalling networks simultaneously in a

heterogeneous population of individual cells. We used an established multi-scale model of B-

cell fates that enables simulation of proliferation, apoptosis and terminal B-cell differentiation

(Fig 2A). The input to this simulation was mitogenic signalling through NF-κB essential

modulator (NEMO)-IκB kinase (IKK) control over the Nuclear Factor κB (NF-κB) transcription

factor. In this modelling framework NF-κB induced pro-proliferation and anti-apoptotic target

genes within the cell cycle and apoptosis modules of the model. In this way we could simulate a

cell population in which cells that undergo apoptosis were removed from the cell population

(determined by an increase in cleaved PARP), while cells that divide introduced new simulations

of daughter cells (determined by an increase in Cdh1, which increases upon completion of

mitosis) (Fig 2A). These models have been validated against cell-population and single-cell data

from primary B-ells, and allowed us to simulate large heterogeneous populations of virtual B-

cells (18, 19). We sought to use this modelling approach to quantify the predicted effect of

mutations impacting multiple signalling pathways. We developed a method of visualising and

quantifying mutational synergy to evaluate the degree to which multiple simulated mutations

confer increased cell population size compared to what would be expected from the sum of the

effects of the individual mutations (Fig 2B). Here, synergy was assumed to be deleterious, as

more synergy results in more proliferative disease.

Mutational synergy in simulations aligns with poor prognosis in patients.

The results of simulations of individual mutations in individual cells (Fig 1) indicated that

overexpression of *MYC* would result in a small increase in cell population size, while the

overexpression of *BCL2* would have a larger effect, and this was indeed seen in multi-scale

modelling results (Fig 2C, red region). Interestingly, the effect of *BCL6* and *MYC* overexpression

on cell number is predicted to be similar, but less than *BCL2* overexpression (Fig 2D). We found

that the *MYC*+*BCL2* double hit simulations, with each gene upregulated 1.5-fold (approximating

one additional copy of each gene), showed a small amount of synergy, producing larger cell

populations than the additive effects of individual mutations (Fig 2C, red region). This synergistic

effect was predicted to be stronger for the DH featuring *MYC*+*BCL2* than for *MYC*+*BCL6* (Fig

2C and 2D). This difference was found to be more pronounced when DH genes were

overexpressed 5 times (vs x1.5), which is consistent with the 4-6 fold change in gene

expression seen in lymphoma patient samples with genomic events affecting *BCL2* and *MYC*

(29)(Fig S5 and S6). Interestingly, this difference in effect size on mutations synergy when

MYC+BCL6 are overexpressed x1.5 to x5 is under half the size seen with increased expression

of *MYC+BCL2* (Fig 2F compared to Fig 2E and Fig S5). If mutational synergy correlates with

prognosis, these simulations predicted that double hits involving *BCL2* are more likely to confer

aggressive disease than those involving *BCL6* (Fig 2C-F). Indeed, this was consistent with

results from a multi-centre study that found the prognosis for *MYC+BCL2* double hits was

significantly worse than for *MYC+BCL6* (30). The dose-dependent increase in synergy with

extra copies of *BCL2* and *MYC* also aligns with clinical outcome as *MYC* rearrangements, in

which the gene is highly expressed, confer worse prognosis than copy number alterations (31).

Given the alignment between mutational synergy and outcomes we found for lymphoma, and to

establish whether the utility of simulated mutational synergy is restricted to lymphoma, we

simulated the impact of genetic events that commonly occur in cases of MM with poor

prognosis. Gain 1q MM is a commonly occurring cytogenetic event in MM associated with

therapeutic failure and shortened survival (32). Simulating chromosomal gains with the multi-

scale modelling framework required mapping genes on the chromosome to model parameters.

Genes encoding the Bcl2-family protein Mcl1 and cell cycle regulator Cks1b both reside on

chromosome 1q21 and have been implicated in the pathogenesis of gain1q MM (33). We

mapped these genes to modelled parameters (see methods) and found a similar pattern to DH

lymphoma, i.e., a small amount of synergy for x1.5-fold increase (equivalent to one extra copy)

and increasing synergy proportional to chromosome 1q copy number (Fig 3A-C). Strikingly, with

5-fold increases in expression of *MCL1* and *CKS1b* the synergistic effect is such that the cell

population continues to increase in size throughout the simulation, even beyond the point at

which the input signal (IKK activity) is removed (Fig 3C, and IKK input curve; Fig 2A). The dose-

dependent increase in mutational synergy aligns with MM patient outcome data, which shows

that prognosis worsens with increasing copies of chromosome 1q (Fig 3D and Fig S7). Taken

together we found that by mapping mutational events and chromosomal abnormalities to model

parameters, then simulating the impact of individual mutations and the combined impact of

multiple mutations, the resultant mutational synergy correlated with outcomes in patients with

poor prognosis lymphoma and MM.

Disease severity can be predicted from multi-hit mutational synergy in patients with poor

prognosis.

Only a small subgroup of lymphoma patients (~5%) have well-defined DH lymphoma. Despite

this, over 20% of patients will not survive 3 years after diagnosis. While co-morbidities and non-

genetic changes likely contribute to this variability in prognosis, we wondered whether

substantially more patients exist with complex mutational combinations that confer poor

prognosis. Given the observations from simulating DH lymphoma and MM we hypothesised that

calculating per-patient mutational synergy using the mutational profile of individual patients may

reveal a correlation with overall survival. To test this hypothesis, we leveraged comprehensive

WES data from patients with long-term follow-up (Fig 4A)(11). To create virtual patients, we

mapped mutated genes and chromosomal abnormalities to modelled parameters (see

Methods). Initial analysis, in which we visualised peak cell population size and mutational

synergy in a small subset of patients, indicated that patients predicted to have high mutational

synergy had relatively poor prognoses (Fig 4B & 4C).

To test whether simulated mutational synergy can predict overall survival we simulated both

individual mutations and the combined mutational profile of all deceased DLBCL patients in the

WES cohort with OS <36 months (11); the majority of events in trials that stratify DH lymphoma

patients occur in the first 3 years (11, 34, 35). This analysis required simulating 100 founder

cells for each mutation which, following cell proliferation within the agent-based modelling

framework, resulted in simulating 100-893 cells per condition. Performing this analysis for these

patients required solving 15783 individual cells, and a total of 306902 ordinary differential

equations. This analysis resulted in simulated cell numbers for each mutation reported in the

WES data for this cohort, and cell numbers for the combination of mutations found in each

patient. We assumed the effect size of each mutation was 1.5-fold for upregulation or 0.5 for

downregulation. Simulations for every patient are provided in the supplementary material. To

quantify the magnitude of synergy we created a mutational synergy score (MSS), defined as the

difference in cell number between the sum of individual mutations and combinations of all

mutations, normalised to the cell population size for each patient. MSS can be visualised as the

largest height of the red region in Fig 2, 3 and 4B-C, normalised to the peak red line in each

plot.

Comparing MSS to OS revealed a significant inverse correlation between the MSS and survival

(Fig 4D R^2 = 21.2%, p = 0.0156 using a generalised linear regression analysis). Simulated cell

number alone (peak simulated cell number rather than MSS) did not correlate as strongly with

OS (R^2=0.1, Fig S8). Interestingly, when we compared MMS in the DLBCL cohort to those

calculated for the classic DH simulations, we identified two patients with higher synergy scores

than we found for 5-fold overexpression of both *MYC* and *BCL2* 5x (Fig 2E, Fig S6, MSS=

0.76). Both patients have OS <7 months and co-occurring mutations impacting NF-κB (REL

copy number increase), apoptosis (*BCL2* or *MCL1*), and blocking B-cell differentiation (*BLIMP1*

loss). While one patient also has a cell cycle mutation (*RB1*), the other lacks cell cycle activating

mutations but may be driven into a highly proliferative regime via NF-κB due to a mutation in

CARD11. Stratifying patients by MSS revealed that patients with low synergy survived

significantly longer than patients with high mutational synergy (Fig 4E, p = 0.02). By

comparison, stratification of the same patient cohort by widely used gene expression-based cell-

of-origin classifications and more recent mutational-based clustering did not reach significance

for association with survival (Fig 4E). We thus demonstrated that using WES data to create

computational models of individual patients, and quantifying mutational synergy, enables

prediction of OS in a cohort of patients with poor prognosis.

Discussion

Many types of cancer are clinically and genetically heterogeneous. Developments in sequencing

technologies have led to the discovery of distinct subgroups of cancer based on mutational

profiles. Often, as is the case in B-cell lymphoma, these disease subtypes can be differentiated

by the signalling pathways that are most dysregulated (11-13). Despite this, knowledge of the

signalling pathways and their interactions are not routinely used to classify disease or predict

prognosis. Computational models of signalling networks can encode decades of knowledge

(20), and many of these models were originally parameterised or validated in wet-lab

experiments where genes are mutated (36). Here we demonstrated that by mapping genetic

data onto computational models we could quantitatively predict the degree to which

combinations of mutations combine deleteriously. We simulated mutations and chromosomal

abnormalities by adjusting appropriate model parameters and thereby placed the genes effected

in the context of molecular signalling networks. We also developed a novel metric: "mutational

synergy score", or MSS, to quantify how genomic events combine to contribute to disease

severity. We demonstrated the ability of mutational synergy to identify established subgroups of

lymphoma and myeloma that have the worst prognosis. This approach enabled us to create

virtual patients using WES data and predict prognosis in patients by identifying the most

deleterious combinations of mutations, even when clustering patients by their gene expression

or mutational burden was not prognostically informative.

We used established multi-scale, agent-based models of B cells (18, 19), in which cellular

signalling networks were encoded in continuous ODE models. Logical modelling is frequently

used to simulate networks of this size and complexity, and indeed has been previously applied

to B-cell lymphoma (37)**.** Here, through leveraging a continuous modelling framework, we were

able to identify many dose-dependent effects, including how chromosomal gain and

amplification confer distinct prognoses. Given the size of the ODE systems used here (194

equations, and 563 parameters) a large amount of information is required to parameterise these

ODE models. Here we performed no parameter fitting, as these parameters have been

accumulated and validated across multiple cellular contexts and cell types (18, 20, 21, 36). We

assume that parameters, other than those effected by mutations, remain consistent between

healthy B-cells and cancerous B-cells. Indeed, many of these parameters have been shown to

be consistent in multiple cell types including T-cells, dendritic cells and fibroblasts (20).

Embedding Large ODE systems in agent-based models, that result in simulations of thousands

of cells, presents a substantial computational challenge. Here, these challenges were overcome

through high performance computing and by leveraging results that demonstrate that B-cell

fates are reliably inherited (18), enabling simulation of a single branch down each lineage tree

for each founder cell and interference of the fate of all remaining daughter cells. We also utilised

the fact that individual mutations are present in multiple patients to reduce the need to simulate

each individual mutation in each individual patient. We benefited from the performance of the

Julia programming language (38), the DifferentialEquations package (39), and used Docker

containers to ensure reproducibility.

Whilst we find computational modelling aligns with and exceeds clinically utilised disease

classifications for these poor prognosis patients, further work is required to develop this

approach into a clinical tool. The correlation between mutational synergy and overall survival is

not strong enough alone to inform clinical prognostic decisions. This may result from some of

the assumptions we needed to make, including assuming that all mutations have the same

effect size, equivalent to one extra copy or loss of copy of each gene. We expect that identifying

and quantifying the impact of each mutation on each gene would improve the correlation

between mutational synergy and OS. Given that only WES data were used to inform the model,

and the assumptions made to map mutations to model parameters, the correlation between

mutational synergy and outcome is perhaps surprising. Simulating the effect of targeted

inhibitors in the patient-specific simulations presented here may provide insights into which

patients will respond to these drugs, identify new treatment strategies, and shed light on

treatment resistance.

Our analysis is limited to poor prognosis patients who succumb to the disease within

approximately 3 years. Further work is required to simulate how treatment may perturb

biological networks harbouring mutations across long time scales, by simulating the effects of R-

CHOP chemotherapy, and polatuzumab- or etoposide-containing regimes. This work may

predict and provide mechanistic insight into the development of relapsed DLBCL. Modelling

long-term responses will require consideration of the causes of relapse. Previous modelling

studies have addressed this challenge by simultaneously simulating populations of malignant

cells, unmutated cells, and premalignant cells to predict the time taken for malignant cells to re-

emerge is predicted (40). Clonal expansion dynamics and cellular plasticity, which are often

associated with increased aggressiveness and treatment resistance, are likely to contribute to

relapse over longer time scales than considered here (41). Expression of epithelial-

mesenchymal transition transcription factors can drive indolent type lymphoma to take on an

aggressive phenotype (42). Regulation of these gene expression programs has been the

subject of multiple mathematical modelling approaches (43). Another model, which simulates

the transition of cancer cells between stem and non-stem states over time, predicted that

plasticity increased tumour growth rate and invasion (44). Creating models that can predict long-

term outcomes and the emergence of treatment resistance from diagnostically-available data

will require reconciling the timescales between the work we present here and models of

treatment resistance/cellular plasticity. Focusing on the poor-prognosis patients in DH

lymphoma enabled us to predict prognosis using computational models focused on molecular

signalling, while also focusing this study on the largest unmet need in this disease. Similarly,

gain(1q) MM represents a substantial unmet need, and our findings that computational

modelling can also predict the correlation between outcome and copies of chromosome 1q seen

in clinical trials indicates that this approach will have utility in multiple malignancies.

Beyond DLBCL and MM, the modelling methodology and synergy calculation described here

can be applied to other tumour types when mutational data and comprehensive, experimentally

validated simulations are available. Many of the genes dysregulated in DLBCL and MM harbour

common cancer-associated mutations. It is well-established that c-Myc deregulation is involved

in the development of a wide variety of cancers (45), Bcl2 is dysregulated in many malignancies

including breast cancer and gastric carcinoma (46, 47), while NF-κB is implicated in numerous

cancers (48). A mathematical model, simulating 17 cancer types, estimated that the number of

carcinogenic mutations (hits) varies from two to eight depending on the cancer type (49). With

the increasing availability of cancer genome data and the evolution of computational methods to

identify the mutational burden of cancer patients, the approach described here is rapidly

becoming feasible in numerous cancers. Mutational synergy metrics are likely to be most

impactful in mutationally heterogeneous cancers, such as breast cancer, where heterogeneity

challenges early diagnosis, treatment selection and prognosis prediction (50).

The goal of cancer research, beyond understanding how the disease develops, is to find

effective treatments for all patients. As knowledge of human cancers has developed it has

become clear that, although patterns may emerge, every patient's cancer is different. We

therefore need better tools, not only to diagnose cancer, but to predict survival and response to

treatment. Modern sequencing techniques have provided ever more data, but we have yet to

fully utilise it. Computational models with the power to translate patient-specific mutation data

into personalised treatment plans will be at the forefront of patient care. The work presented

here is a step towards this, and in future we hope to develop it further and demonstrate its

potential in a range of malignancies.

Methods

Model generation

Ordinary differential equation (ODE) model files were generated from individual reaction,

parameter and rate law tables using CSV2Julia, a Python script developed in-house (see

https://github.com/SiFTW/CSV2JuliaDiffEq). Jupyter Notebooks with Julia code used to

generate and run all computational models and plot output are available at

https://github.com/SiFTW/norrisEtAl. Ordinary differential equation models were solved using

DifferentialEquations.jl (38, 39). Full description of the models reactions, parameters, and rate

laws are defined in the GitHub repository.

Cell cycle model

Each cell was simulated until either a limit cycle was hit, or the maximum time allowed for the

simulation ($1x10^6$ minutes). We defined the limit cycle as when the maximum value of cell mass

is the same, within 3 decimal places, for 5 consecutive cycles

(https://github.com/SiFTW/norrisEtAl/CellCycle_WTvsMycOE.ipynb). Simulations of cell

populations were performed using distributions of parameters representing expression and

degradation of modelled molecular species as described previously with 11% coefficient of

variance sampled from a normal distribution around the published parameter value truncated at

0 (18) (https://github.com/SiFTW/norrisEtAl/CellCycleParamDist.ipynb).

Apoptosis model

The model was initially run to steady state. The final values for each molecular species were

then used (unmutated) or modified (to reflect mutations) as input to a time course phase. Values

for PARP and cleaved PARP were extracted and the ratio of PARP to cleaved PARP was

calculated for each time point in the simulation. Time to death was defined as the point at which

this ratio was >0.1 (https://github.com/SiFTW/norrisEtAl/Apoptosis_Bcl2.ipynb,

https://github.com/SiFTW/norrisEtAl/ApoptosisDist.ipynb).

Multi-scale model

The multi-scale model was generated using module definition files (reactions, rate laws and

parameters) for individual modules (NF-κB signalling, cell cycle, apoptosis, and differentiation)

as well as a set of linking reactions. Any reactions or parameters appearing in individual

modules that overlap with linking reactions were replaced by the corresponding information from

the linking files. Cell populations were simulated according to a set of pre-defined conditions

including condition name, parameter(s) to modify and amount to modify the parameter(s) by. All

these conditions were kept constant for all simulations presented here except for individual

parameters changed to individual mutations.

Callbacks were used to modify the values of molecular species during the simulation upon

reaching a set of pre-defined conditions. In the multi-scale model this allows cells to divide or

die, and in the case of division introduces new simulations. The two callbacks used in multi-

scale modelling were maintained from previous work (cleaved PARP >2500 = apoptosis, CDH1

>0.1 = cell division) (18, 19).

The introduction of two daughter cells with each cell division resulted in an infeasible simulation

size in highly proliferative simulations, and cell fates have been shown to be reliably inherited in

B-cells such that both daughter cells will achieve similar fates at similar times (16). Therefore, at

each cell division we introduced a single daughter representing both daughter cells, with a

variable to track the number of cells represented by each daughter, using the formula: [cells

represented] = $2^{(generation-1)}$.

Files that define the multi-scale model along with Jupyter notebooks to run the simulations

described here are available in the GitHub repository.

Modelling patients

We restricted our analysis to mutations that impact the core simulated network of the multi-scale

model using a gene list of recurrent mutations that are known to impact well-defined nodes of

the model (gene list: https://github.com/SiFTW/norrisEtAl/geneList.txt). Querying this gene list

through the cBioPortal we converted mutations to parameter value changes based on literature

regarding the effect of each mutation. For gain-of-function mutations we multiplied the

appropriate parameter value by 1.5 and for loss-of-function we multiplied by 0.5. For copy

number alterations (CNA) we found each gene on the respective chromosome section and

multiplied by 1.5 for an amplification and 0.5 for heterozygous loss. If multiple mutations

affected the same parameter, then the individual effects of those mutations were multiplied

together. For mutations in genes not modelled directly we assigned an indirect effect. For

example, the effect of mutations in p53 is modelled as overexpression of *BCL2* as it is known to

inhibit Bcl2, and *CARD11* mutations are modelled as increasing NEMO-IKK activity (51).

Calculating Metrics

For each simulation the number of cells alive in the population was exported, and we then

calculated a series of metrics (see

https://github.com/SiFTW/norrisEtAl/calculateCellNumbersForAllConditions.ipynb):

PeakCN = maximum number of cells at any time point in the simulation of a patients' full

genotype

PeakEIA = maximum number of cells at any time point calculated from the sum of all individual

mutations minus WT.

PeakSyn = maximum value of peakCN minus peakEIA

NormalisedSyn = peakSyn/peakEIA

Acknowledgements

The authors would like to thank Ulf Klein, Jana Wolf, and Richard Norris for critical discussions and

Eleanor Jayawant for providing critical input into the manuscript. Funding for SM: Leukaemia UK John

Goldman Fellowship (2020/JGF/003) and UKRI Future Leaders Fellowship (MR/T041889/1). Funding for

AP: MRC Research Grant (MR/V009095/1). Funding for CP: Blood Cancer UK (23004).

Author contributions

RN and SM performed computational analysis and modelling. RN, CP, AP and SM

conceptualized the study, and EM, JJ and TC conceptualized the Figure 3. AP, CP and SM

supervised the project. All authors contributed to the article, critically revised the manuscript,

and approved the submitted version.

Disclosure and competing interests statement

The authors state they have no competing interests or disclosures.

Data Availability

The datasets and computer code produced in this study are available in the following

databases:

Modelling computer scripts: GitHub (https://github.com/SiFTW/norrisEtAl).

References

1.	Maura F, Bolli N, Angelopoulos N, Dawson KJ, Leongamornlert D, Martincorena I, et al. Genomic landscape and chronological reconstruction of driver events in multiple myeloma. Nat Commun. 2019;10(1):3835.

2.	Mosquera Orgueira A, Ferreiro Ferro R, Díaz Arias J, Aliste Santos C, Antelo Rodríguez B, Bao Pérez L, et al. Detection of new drivers of frequent B-cell lymphoid neoplasms using an integrated analysis of whole genomes. PLoS One. 2021;16(5):e0248886.

3.	Cheah C, Herbert K, O'rourke K, Kennedy GA, George A, Fedele P, et al. A multicentre retrospective comparison of central nervous system prophylaxis strategies among patients with high-risk diffuse large B-cell lymphoma. British journal of cancer. 2014;111(6):1072-9.

4.	Hartert KT, Wenzl K, Krull JE, Manske M, Sarangi V, Asmann Y, et al. Targeting of inflammatory pathways with R2CHOP in high-risk DLBCL. Leukemia. 2021;35(2):522-33.

5.	Walz S, Lorenzin F, Morton J, Wiese KE, von Eyss B, Herold S, et al. Activation and repression by oncogenic MYC shape tumour-specific gene expression profiles. Nature. 2014;511(7510):483-7.

6.	Xu-Monette ZY, Deng Q, Manyam GC, Tzankov A, Li L, Xia Y, et al. MYC mutation profiling and prognostic significance in de novo diffuse large B-cell lymphoma. Clinical cancer research: an official journal of the American Association for Cancer Research. 2016;22(14):3593.

7.	Yip K, Reed J. Bcl-2 family proteins and cancer. Oncogene. 2008;27(50):6398-406.

8.	Hanahan D, Weinberg RA. The hallmarks of cancer. cell. 2000;100(1):57-70.

9.	Ci W, Polo JM, Cerchietti L, Shaknovich R, Wang L, Yang SN, et al. The BCL6 transcriptional program features repression of multiple oncogenes in primary B cells and is deregulated in DLBCL. Blood, The Journal of the American Society of Hematology. 2009;113(22):5536-48.

10.	Rosenwald A, Wright G, Chan WC, Connors JM, Campo E, Fisher RI, et al. The use of molecular profiling to predict survival after chemotherapy for diffuse large-B-cell lymphoma. New England Journal of Medicine. 2002;346(25):1937-47.

11.	Chapuy B, Stewart C, Dunford AJ, Kim J, Kamburov A, Redd RA, et al. Molecular subtypes of diffuse large B cell lymphoma are associated with distinct pathogenic mechanisms and outcomes. Nat Med. 2018;24(5):679-90.

12.	Lacy SE, Barrans SL, Beer PA, Painter D, Smith AG, Roman E, et al. Targeted sequencing in DLBCL, molecular subtypes, and outcomes: a Haematological Malignancy Research Network report. Blood. 2020;135(20):1759-71.

13.	Schmitz R, Wright GW, Huang DW, Johnson CA, Phelan JD, Wang JQ, et al. Genetics and Pathogenesis of Diffuse Large B-Cell Lymphoma. N Engl J Med. 2018;378(15):1396-407.

14.	Ennishi D, Jiang A, Boyle M, Collinge B, Grande BM, Ben-Neriah S, et al. Double-hit gene expression signature defines a distinct subgroup of germinal center B-cell-like diffuse large B-cell lymphoma. Journal of Clinical Oncology. 2019;37(3):190.

15.	Sehn LH, Salles G. Diffuse Large B-Cell Lymphoma. N Engl J Med. 2021;384(9):842-58.

16.	Liu X, Jia S, Chu Y, Tian B, Gao Y, Zhang C, et al. Chromosome 1q21 gain is an adverse prognostic factor for newly diagnosed multiple myeloma patients treated with bortezomib-based regimens. Front Oncol. 2022;12:938550.

17.	Abdallah N, Greipp P, Kapoor P, Gertz MA, Dispenzieri A, Baughn LB, et al. Clinical characteristics and treatment outcomes of newly diagnosed multiple myeloma with chromosome 1q abnormalities. Blood Adv. 2020;4(15):3509-19.

18.	Mitchell S, Roy K, Zangle TA, Hoffmann A. Nongenetic origins of cell-to-cell variability in B lymphocyte proliferation. Proceedings of the National Academy of Sciences. 2018;115(12):E2888-E97.

19.	Roy K, Mitchell S, Liu Y, Ohta S, Lin Y-s, Metzig MO, et al. A regulatory circuit controlling the dynamics of NFκB cRel transitions B cells from proliferation to plasma cell differentiation. Immunity. 2019;50(3):616-28. e6.

20.   Mitchell S, Tsui R, Tan ZC, Pack A, Hoffmann A. The NF-κB multidimer system model: A
knowledge base to explore diverse biological contexts. Science Signaling. 2023;16(776):eabo2838.
21.   Shokhirev MN, Almaden J, Davis-Turak J, Birnbaum HA, Russell TM, Vargas JA, et al. A multi-scale
approach reveals that NF-κB cRel enforces a B-cell decision to divide. Mol Syst Biol. 2015;11(1):783.
22.   Youle RJ, Strasser A. The BCL-2 protein family: opposing activities that mediate cell death. Nat
Rev Mol Cell Biol. 2008;9(1):47-59.
23.   Heinzel S, Binh Giang T, Kan A, Marchingo JM, Lye BK, Corcoran LM, et al. A Myc-dependent
division timer complements a cell-death timer to regulate T cell and B cell responses. Nat Immunol.
2017;18(1):96-103.
24.   Loriaux PM, Tesler G, Hoffmann A. Characterizing the relationship between steady state and
response using analytical expressions for the steady states of mass action models. PLoS computational
biology. 2013;9(2):e1002901.
25.   Dowling MR, Kan A, Heinzel S, Zhou JH, Marchingo JM, Wellard CJ, et al. Stretched cell cycle
model for proliferating lymphocytes. Proc Natl Acad Sci U S A. 2014;111(17):6377-82.
26.   Cai C, Hu X, Dai P, Zhang T, Jiang M, Wang L, et al. c-Myc regulates neural stem cell quiescence
and activation by coordinating the cell cycle and mitochondrial remodeling. Signal Transduct Target
Ther. 2021;6(1):306.
27.   Cheng Y, Luo H, Izzo F, Pickering BF, Nguyen D, Myers R, et al. m(6)A RNA Methylation Maintains
Hematopoietic Stem Cell Identity and Symmetric Commitment. Cell Rep. 2019;28(7):1703-16.e6.
28.   Duffy KR, Hodgkin PD. Intracellular competition for fates in the immune system. Trends Cell Biol.
2012;22(9):457-64.
29.   Meriranta L, Pasanen A, Alkodsi A, Haukka J, Karjalainen-Lindsberg M-L, Leppä S. Molecular
background delineates outcome of double protein expressor diffuse large B-cell lymphoma. Blood
advances. 2020;4(15):3742-53.
30.   El-Sharkawi D, Sud A, Prodger C, Khwaja J, Shotton R, Hanley B, et al. A retrospective study of
MYC rearranged diffuse large B-cell lymphoma in the context of the new WHO and ICC classifications.
Blood Cancer J. 2023;13(1):54.
31.   Quesada AE, Medeiros LJ, Desai PA, Lin P, Westin JR, Hawsawi HM, et al. Increased MYC copy
number is an independent prognostic factor in patients with diffuse large B-cell lymphoma. Mod Pathol.
2017;30(12):1688-97.
32.   Schmidt TM, Fonseca R, Usmani SZ. Chromosome 1q21 abnormalities in multiple myeloma.
Blood Cancer J. 2021;11(4):83.
33.   Hanamura I. Gain/Amplification of Chromosome Arm 1q21 in Multiple Myeloma. Cancers
(Basel). 2021;13(2).
34.   Sarkozy C, Traverse-Glehen A, Coiffier B. Double-hit and double-protein-expression lymphomas:
aggressive and refractory lymphomas. The Lancet Oncology. 2015;16(15):e555-e67.
35.   Akyurek N, Uner A, Benekli M, Barista I. Prognostic significance of MYC, BCL2, and BCL6
rearrangements in patients with diffuse large B-cell lymphoma treated with cyclophosphamide,
doxorubicin, vincristine, and prednisone plus rituximab. Cancer. 2012;118(17):4173-83.
36.   Hoffmann A, Levchenko A, Scott ML, Baltimore D. The IkappaB-NF-kappaB signaling module:
temporal control and selective gene activation. Science. 2002;298(5596):1241-5.
37.   Thobe K, Konrath F, Chapuy B, Wolf J. Patient-Specific Modeling of Diffuse Large B-Cell
Lymphoma. Biomedicines. 2021;9(11).
38.   Bezanson J, Edelman A, Karpinski S, Shah VB. Julia: A fresh approach to numerical computing.
SIAM review. 2017;59(1):65-98.
39.   Rackauckas C, Nie Q. Differentialequations. jl–a performant and feature-rich ecosystem for
solving differential equations in julia. Journal of open research software. 2017;5(1).

40.   Abubakar SD, Takaki M, Haeno H. Computational modeling of locoregional recurrence with
spatial structure identifies tissue-specific carcinogenic profiles. Front Oncol. 2023;13:1116210.
41.   Qin S, Jiang J, Lu Y, Nice EC, Huang C, Zhang J, et al. Emerging role of tumor cell plasticity in
modifying therapeutic response. Signal Transduct Target Ther. 2020;5(1):228.
42.   Chen SC, Liao TT, Yang MH. Emerging roles of epithelial-mesenchymal transition in
hematological malignancies. J Biomed Sci. 2018;25(1):37.
43.   Tripathi S, Xing J, Levine H, Jolly MK. Mathematical Modeling of Plasticity and Heterogeneity in
EMT. Methods Mol Biol. 2021;2179:385-413.
44.   Poleszczuk J, Enderling H. Cancer Stem Cell Plasticity as Tumor Growth Promoter and Catalyst of
Population Collapse. Stem Cells Int. 2016;2016:3923527.
45.   Kalkat M, De Melo J, Hickman KA, Lourenco C, Redel C, Resetca D, et al. MYC Deregulation in
Primary Human Cancers. Genes (Basel). 2017;8(6).
46.   Joensuu H, Pylkkänen L, Toikkanen S. Bcl-2 protein expression and long-term survival in breast
cancer. Am J Pathol. 1994;145(5):1191-8.
47.   Inada T, Kikuyama S, Ichikawa A, Igarashi S, Ogata Y. Bcl-2 expression as a prognostic factor of
survival of gastric carcinoma. Anticancer Res. 1998;18(3b):2003-10.
48.   Luo JL, Kamata H, Karin M. IKK/NF-kappaB signaling: balancing life and death--a new approach
to cancer therapy. J Clin Invest. 2005;115(10):2625-32.
49.   Anandakrishnan R, Varghese RT, Kinney NA, Garner HR. Estimating the number of genetic
mutations (hits) required for carcinogenesis based on the distribution of somatic mutations. PLoS
Comput Biol. 2019;15(3):e1006881.
50.   Guo L, Kong D, Liu J, Zhan L, Luo L, Zheng W, et al. Breast cancer heterogeneity and its
implication in personalized precision therapy. Exp Hematol Oncol. 2023;12(1):3.
51.   Hemann MT, Lowe SW. The p53-Bcl-2 connection. Cell Death Differ. 2006;13(8):1256-9.

Figure 1. Computational modelling of the cell cycle and apoptosis reveal limited impact of archetypal double hit mutations on their respective molecular networks.

A. (top) Output of the cell cycle model run to a limit cycle for a WT cell and a cell in which *MYC* is overexpressed (x1.5). Lines indicate the values of molecular species (used to calculate phase transition times) over time from the first growth phase (G1) to cell division (M). Cyclin is abbreviated to "Cyc".

A. (bottom) Cell cycle phase timing for cells with varying *MYC* expression.

B. (top) The percentage of cleaved PARP over time for two simulations (WT and *BCL2* overexpressed) using the apoptosis model. B (bottom). Time to death for cells with varying *BCL2* expression.

C, D. Comparison of total cell cycle times from limit cycles of simulations of heterogeneous unmutated (c-MYC x1.0) cells and cells with a *MYC* mutation (c-MYC x1.5). C. Total cell cycle times in c-MYC x1.0 simulations are indicated with a black line, while c-MYC x1.5 is indicated with green bars. Each unmutated cell is plotted alongside the corresponding *MYC* mutant (by cell ID). The only difference between matched cells is *MYC* expression level but each cell pair has distributed parameters compared to every other pair. The inset bar plot indicates the number of cells (in c-MYC x1.0 & x1.5 simulations) in which cell cycle arrest is predicted. D. Cell cycle times for the same simulations represented as panel C but displayed density plots. c-MYC x1.0 is shown in black and c-MYC x1.5 in green.

E, F. Comparisons of time to death for simulations of heterogeneous populations of matched cells differing only in expression level of *BCL2*. Simulations are shown as both a horizontal bar plot (E) with BCL2 x1.0 shown in black and BCL2 x1.5 shown in green, and a density plot (F) with the same colour scheme.

Figure 2. Multi-scale modelling of heterogeneous cell populations reveals that mutations combine with oncogenic synergy.

A) Schematics of the multi-scale model, including the signalling networks within the model and the cell fates they control. The full reaction scheme of the model is defined in the Github repository associated with this study: https://github.com/SiFTW/norrisEtAl. Simulation input, NEMO and IKK activity curves, are indicated. A schematic tree structures of cell division and death in simulations of cell populations with and without mutations is provided (right).

B) Explanation of the method for calculating a comparison of individual mutations and combinations of mutations (mutational synergy). Mutational synergy is defined as the difference in cell number between the sum of cell numbers from all simulations containing individual mutations, and the cells added in a simulation that combines all mutations. This difference defines the red region in this schematic and future plots.

C-F) Cell population size (cell count) over time for wild type/unmutated (blue), individual mutations (orange and red) and double hits (green). Each simulation started with 100 heterogeneous B-cells stimulated by the indicated IKK input curve (panel A). C & E) Cell numbers and synergistic regions for *MYC* and *BCL2* overexpression x1.5 (C) and x5 (E). D & F) *MYC* and *BCL6* overexpression x1.5 (D) and x5 (F).

D) Multiple Myeloma Clinical Trial Data

1q copies	PFS (95% CI, months)
2	65.1 (53.4-NR)
3	55.9 (32.6–78.9)
≥4	34.6 (15.6–61.1)

Schmidt *et al.* (2019) Blood Cancer Journal

Figure 3. Mutations associated with aggressive multiple myeloma are also predicted to create mutational synergy.

A-C) Cell population size (cell count) over time for wild type/unmutated (blue), individual mutations (orange and green) and double hits (pink). *CKS1B* and *MCL1* overexpression x1.5 (A), x2 (B) and x5 (C).

D) Data from Schmidt et al., (32) showing the average progression free survival (PFS) for patients in a Multiple Myeloma cohort with varying chromosome 1q copy number.

Figure 4. Mutational synergy is correlated with overall survival in patients with the worst outcomes.

A) OncoPrint output showing mutations and clinical data for deceased patients from Chapuy et al., (11) with overall survival <36 months. All data is from Chapuy et al., (11) except mutational synergy which is calculated here.

B, C) Plots of cell number vs time for simulations of individual mutations and all mutations combined for patients with low (B) and high (C) synergy.

D) Plot of normalised synergy score vs overall survival for deceased patients with OS <36 months. The blue line shows best fit for synergy vs OS correlation; blue shadowing shows 95% confidence interval; orange and blue bars (right) delineate groups of patients with low and high synergy.

E) Box plots showing mean survival (months) for groups of patients with high (blue) and low (orange) synergy. Adjacent box plots show how stratifying patients by cell of origin and mutational clusters compares to the normalised synergy metric when plotted against overall survival (OS). P> 0.1 for all pairwise comparisons between mutational clusters for these patients from Chapuy et al. (11).