Glutamatergic neuronal activity regulates angiogenesis and blood-retinal barrier maturation via Norrin/β-catenin signaling

Saptarshi Biswas[1,*], Sanjid Shahriar[2, 3], Galina Bachay[4], Panos Arvanitis[5], Danny Jamoul[1, 6], William J. Brunken[4] and Dritan Agalliu[1, 2, *]

[1]Department of Neurology, Columbia University Irving Medical Center, New York, NY 10032, USA.

[2]Department of Pathology and Cell Biology, Columbia University Irving Medical Center, New York, NY 10032, USA.

[3]Wyss Institute for Biologically Inspired Engineering, Boston, MA 02115, USA.

[4]Department of Ophthalmology and Visual Sciences, SUNY Upstate Medical University, Syracuse, NY 13210, USA.

[5]Warren Alpert Medical School, Brown University, Providence, RI 02903, USA.

[6]John Jay College of Criminal Justice, City University of New York, New York, NY 10019, USA.

*Corresponding authors:

Saptarshi Biswas Ph.D.   Email – sb4082@cumc.columbia.edu

Dritan Agalliu Ph.D.   Email – da191@cumc.columbia.edu

Lead contact:

Dritan Agalliu Ph.D.   Email – da191@cumc.columbia.edu

Keywords:

Glutamate, neuronal activity, angiogenesis, blood-retinal barrier, tight junctions, Norrin/β-catenin signaling

SUMMARY

Interactions among neuronal, glial and vascular components are crucial for retinal angiogenesis and blood-retinal barrier (BRB) maturation. Although synaptic dysfunction precedes vascular abnormalities in many retinal pathologies, how neuronal activity, specifically glutamatergic activity, regulates retinal angiogenesis and BRB maturation remains unclear. Using *in vivo* genetic studies in mice, single-cell RNA-sequencing and functional validation, we show that deep plexus angiogenesis and paracellular BRB maturation are delayed in *Vglut1*$^{-/-}$ retinas where neurons fail to release glutamate. In contrast, deep plexus angiogenesis and paracellular BRB maturation are accelerated in *Gnat1*$^{-/-}$ retinas where constitutively depolarized rods release excessive glutamate. Norrin expression and endothelial Norrin/β-catenin signaling are downregulated in *Vglut1*$^{-/-}$ retinas, and upregulated in *Gnat1*$^{-/-}$ retinas. Pharmacological activation of endothelial Norrin/β-catenin signaling in *Vglut1*$^{-/-}$ retinas rescued defects in deep plexus angiogenesis and paracellular BRB maturation. Our findings demonstrate that glutamatergic neuronal activity regulates retinal angiogenesis and BRB maturation by modulating endothelial Norrin/β-catenin signaling.

INTRODUCTION

The crosstalk among neuronal, glial, and vascular components is essential, not only for proper angiogenesis, but also for the establishment of the inner blood-retinal barrier (BRB) in the developing retina [1]. Retinal angiogenesis is tightly coordinated with the maturation of the BRB properties by endothelial cells (ECs), through the presence of tight-junctions (TJ) that limit paracellular permeability between ECs, specialized transporters to facilitate bidirectional transport of nutrients, and suppression of transcytosis that reduces transcellular EC permeability [2,3]. Although multiple signaling pathways operating in ECs, pericytes (PCs), neuronal or glial cells have emerged as critical regulators of retinal angiogenesis and BRB maturation [reviewed in [4]], it is unclear how distinct types of neuronal synaptic activities influence these processes, and whether neurotransmitters act directly on ECs to mediate their effects. This is highly significant since early loss of neurovascular coupling, impaired response of photoreceptors to light, gradual neurodegeneration, gliosis and neuroinflammation are found prior to any obvious vascular pathologies in diabetic retinopathy and other retinal disease [5-7]. Moreover, it has been postulated that diabetes-induced changes in neurons and glial cells lead to vascular defects [8], yet the underlying mechanisms remain elusive.

Distinct types of neuronal activities are present in the postnatal mouse retina during angiogenesis and BRB maturation **(Figure 1A)**. From birth to postnatal day (P) 10, starburst amacrine cells propagate spontaneous cholinergic waves [9-11]. These cholinergic waves gradually subside and spontaneous glutamatergic waves emerge from P10 to P14 [11-14]. The majority of neuronal activities in the mouse retina from P14 onwards are mediated by light-dependent glutamatergic synaptic activity of photoreceptors, and photoactivation of intrinsically-photosensitive retinal ganglion cells (ipRGCs) [reviewed in [4]]. However, light activation of photoreceptors has been observed as early as P8 [13,15]. The superficial vascular plexus in the retina develops between P0 - P8 [16] over the ganglion cell layer (GCL) synchronous with neuronal activities from the cholinergic waves and photoactivation of ipRGCs. In contrast, the deep plexus angiogenesis (P8 - P12) in the outer plexiform layer (OPL) [16] and BRB maturation (P6 - P18) span both

cholinergic and glutamatergic spontaneous waves, glutamatergic synaptic activity of photoreceptors as well

as photoactivation of ipRGCs [reviewed in [4]]. Few studies have addressed how neuronal activity regulates

retinal angiogenesis and BRB maturation in the developing retina. Photoactivation of both Opn4[+ 17] and

Opn5[+ 18] RGCs regulates angiogenesis in the retina. Pharmacological blockade of cholinergic waves before

P8 delays deep plexus angiogenesis and BRB maturation, suggesting an important role for cholinergic

activity in these processes [14]. However, the role of glutamatergic neuronal activities in retinal angiogenesis

and BRB maturation remains unclear, although it is the predominant type of neuronal activity when most

of the vascular development and BRB maturation occurs in the retina.

The Norrin/β-catenin signaling is critical for retinal angiogenesis and BRB maturation by inducing

stabilization and nuclear translocation of β-catenin. Genetic elimination of Norrin (ligand), Frizzled-4,

Lrp5/6, Tspan12 (receptors) and β-catenin in the mouse retina delays superficial vascular plexus

angiogenesis, induces loss of the deep plexus [19-23] and BRB integrity [24,25]. How Norrin expression and

endothelial Norrin/β-catenin signaling are related to the glutamatergic activities in the developing retina is

also unknown.

Using *in vivo* anatomical and functional characterizations of the vasculature, as well as single-cell

transcriptomic analyses and validation studies, we show that deep plexus angiogenesis and paracellular

BRB maturation are delayed in *Vglut1*[-/-] retinas that have reduced synaptic release of glutamate from

photoreceptors and bipolar cells [12]. In contrast, deep plexus angiogenesis is accelerated and paracellular

BRB matures precociously in *Gnat1*[-/-] retinas, where rod photoreceptors remain constitutively depolarized

due to the absence of α-Transducin, a Gα protein critical for phototransduction [26,27]. These phenotypes are

not due to abnormalities in neurovascular unit (NVU) assembly, aberrant specification of neuronal or glial

subtypes, or changes in the neuronal metabolic demand. Importantly, *Norrin* mRNA expression in the inner

nuclear layer (INL), and as a consequence endothelial Norrin/β-catenin signaling, are downregulated in

Vglut1^{-/-} retinas, and upregulated in *Gnat1^{-/-}* retinas. Pharmacological activation of endothelial Norrin/β-

catenin signaling in *Vglut1^{-/-}* retinas rescues defects in both deep plexus angiogenesis and paracellular BRB

integrity. Our study elucidates a critical mechanism by which glutamatergic neuronal activities regulate

retinal angiogenesis and BRB maturation by modulating endothelial Norrin/β-catenin signaling in the

developing retina. These findings have important implications for development of potential interventions

in various neurovascular pathologies.

RESULTS

Deep plexus angiogenesis is delayed in the *Vglut1^{-/-}* retina.

Photoreceptors release glutamate in the OPL in the dark and suppress glutamate release upon photon

capture [28]. The synaptic release of glutamate is dependent on vGluT1, a vesicular glutamate transporter [29].

Consequently, *Vglut1^{-/-}* mice have reduced synaptic release of glutamate from photoreceptors and bipolar

cells **(Figure 1B)**[12]. To understand if extracellular glutamate levels affect retinal angiogenesis, we compared

the vascular growth in the superficial and deep plexuses between wild-type (WT) and *Vglut1^{-/-}* retinas. Since

the central retina is developmentally more mature than the peripheral retina, we analyzed these two regions

separately **(Figure 1C)**. Although the vascular coverage of the retina in the superficial plexus was similar

in both genotypes at P8 and P10 **(Figure 1D, E, H, I)**, the vascular coverage of the retina in the deep plexus

and its branch-point density were reduced significantly in the P10 and P12 *Vglut1^{-/-}* retinas **(Figure 1E-G,**

J-M). By P14, however, the vascular coverage of the retina in the deep plexus was similar between the two

genotypes, although the branch-point density of the *Vglut1^{-/-}* deep plexus was still significantly lower than

the WT **(Figure 1F, G, J-M)**. Importantly, these vascular abnormalities were consistent in both the central

and peripheral *Vglut1^{-/-}* retinas (**Figure 1J-M**).

To understand the molecular mechanisms underlying these vascular phenotypes, we performed

single-cell RNA-sequencing analyses (scRNA-seq) of ECs that were dissociated from P10 WT and *Vglut1⁻*

/- retinas and sorted for CD31 using fluorescence-activated cell sorting (FACS; **Figure S1A, B**). The purity

of ECs was confirmed by uniform expression of EC markers [*Pecam1* (CD31) and *Cdh5* (VE-Cadherin)]

in the entire population, and absence of mural cell [*Pdgfrb* (PDGFRβ) and *Des* (Desmin)] and astrocyte

[*Gfap* and *Pdgfra*] markers **(Figure S1C)**. The majority of ECs isolated from either genotype were capillary

ECs (Mfsd2a+), as opposed to arterial or venous ECs **(Figure S1D)**, consistent with the predominant

distribution of this EC subtype in both the superficial and deep plexuses of the retina (the deep plexus has

only capillaries). ECs isolated from P10 WT and *Vglut1*$^{-/-}$ retinas clustered separately in the Uniform

Manifold Approximation and Projection (UMAP) plot based on differences in their transcriptomic

signatures **(Figure 2A; Tables S1, S2)**. Gene ontology (GO) analyses of the differentially expressed genes

(DEGs) between the two genotypes revealed that pathways related to "angiogenesis", "blood vessel

morphogenesis" and "cell proliferation" were downregulated in P10 *Vglut1*$^{-/-}$ compared to WT retinal ECs

(Figure 2B; Tables S3, S5), consistent with the phenotypic analyses (**Figure 1**). In contrast, pathways

related to "cellular and protein metabolic processes" were upregulated in P10 *Vglut1*$^{-/-}$ compared to WT

retinal ECs **(Figure 2C; Tables S4)**. Expression levels of several pro-angiogenic genes [e.g. *Ptprb*, *Aplnr*,

Unc5b etc.], including two effectors of Norrin/β-catenin signaling [*Ctnnb1* (β-catenin) and *Lef1*], were

downregulated significantly in P10 *Vglut1*$^{-/-}$ compared to WT retinal ECs **(Figure 2D, F; Table S5)**. In

contrast, the majority of genes enriched in the deep plexus tip cells (D-tip cells) compared to the superficial

plexus tip cells (S-tip cells) [e.g. *Bsg*, *Rgcc*, *Plxnd1*, *Cdkn1a*, *Pde4b*, *Prcp*], as well as those enriched in D-

tip cells compared to all ECs [e.g. *Lamb1*, *Pcdh17*, *Adm*] [30] were upregulated in P10 *Vglut1*$^{-/-}$ compared to

WT retinal ECs **(Figure 2E, G; Table S6)**, indicative of delayed deep plexus angiogenesis at the

transcriptome level. To validate these molecular findings, we labeled proliferating cells with Ki67 in both

WT and *Vglut1*$^{-/-}$ retinas and assessed whether endothelial stalk cell proliferation was reduced in *Vglut1*$^{-/-}$

retinal ECs (**Figure 2H**) [31]. Consistent with the scRNA-seq data, there was a significant reduction in the

number of proliferating Ki67$^+$ ECs in P12 *Vglut1*$^{-/-}$ retinal deep plexus compared to the WT **(Figure 2I, J)**,

while the number of apoptotic ECs (cleaved Caspase-3$^+$) was not significantly different **(Figure 2J)**.

Therefore, the delayed angiogenesis in *Vglut1*[-/-] retinal deep plexus is due, in part, to reduced EC

proliferation. Consistent with the scRNA-seq data, we also observed numerous Dll4[+] tip cells [32,33] in the

leading edge of the *Vglut1*[-/-] deep plexus at P12, while the WT deep plexus was completely vascularized at

this stage **(Figure 2K, L)**. Thus, angiogenesis (EC proliferation) is delayed and the number of endothelial

tip cells is increased in the deep, but not superficial, plexus of *Vglut1*[-/-] retinas lacking extracellular

glutamate release by neurons.

Deep plexus vessels are poorly perfused in the *Vglut1*[-/-] retina.

Next, we asked if the deep plexus vessels are polarized and function normally in the *Vglut1*[-/-] retinas.

We stained blood vessels with Podocalyxin, which labels the apical (luminal) side of ECs, and is a marker

for endothelial apical-basal polarity **(Figure 3A)** [34]. Compared to the WT, Podocalyxin coverage of the

deep plexus vessels was significantly reduced in *Vglut1*[-/-] retinas at both P10 and P14 **(Figure 3B, C-E)**,

suggesting that EC polarity is abnormal in the deep vascular plexus. Next, we examined whether deep

plexus vessels are perfused normally in *Vglut1*[-/-] retinas. We performed intracardiac injection of

fluorescently-labeled Lectin and analyzed its binding to the retinal vasculature as a readout of perfusion.

By P14, the entire OPL in *Vglut1*[-/-] retinas is vascularized **(Figure 1)**. However, the fraction of Lectin[+]

perfused blood vessels was significantly lower in the deep plexus of P14 *Vglut1*[-/-] compared to the WT

retinas **(Figure 3C, F)**. These data demonstrate that, although the deep plexus is fully formed by P14 in the

Vglut1[-/-] retinas (**Figure 1F, G**), EC polarity and vessel perfusion remain abnormal.

Paracellular BRB maturation is impaired in the *Vglut1*[-/-] retina.

Next, we asked if diminished glutamate release from photoreceptors and bipolar cells affects BRB

maturation. During retinal development, the transcellular (caveolae-mediated) BRB matures by P10 [35]

whereas the paracellular (tight junction-mediated) BRB does not mature until P18 [36] **(Figure 4A)**. We

therefore injected intravenously biocytin-tetramethylrhodamine (biocytin-TMR, 870 Da) tracer in WT and

Vglut1⁻ᐟ⁻ mice at P10 and P14 and quantified its leakage in the retina as a functional readout of paracellular

BRB maturation **(Figure 4B)** [36,37]. Biocytin-TMR leakage (area and intensity) was higher in the *Vglut1⁻ᐟ⁻*

compared to the WT retinas at both time points **(Figure 4C-I)**. Importantly, these patterns were consistent

in both the central and peripheral retina, and the superficial and deep plexuses. Consistent with a delay in

BRB maturation, scRNA-seq analyses revealed reduced transcript levels for most "BRB maturation genes"

in P10 *Vglut1⁻ᐟ⁻* compared to WT retinal ECs. These include tight junction-associated genes [*Cldn5*

(Claudin-5) and *Tjp1* (ZO-1)], transporters [*Slc7a5* (Lat1), *Slc2a1* (Glut1) etc.,] and effectors of Norrin/β-

catenin signaling [*Ctnnb1* and *Lef1*] **(Figure 4J, K; Table S7)**. Surprisingly, transcript levels for *Ocln*

(Occludin; tight junction protein), vBM proteins (Collagen IV, Laminin), BBB transcription factors (e.g.

Foxq1, *Foxf2*, *Zic3* etc.), or caveolar-related transport (*Cav-1*, *Cav-2*, *Cavin-1*, *Mfsd2a*) genes were

unaffected in P10 *Vglut1⁻ᐟ⁻* retinal ECs **(Tables S2, S7)**.

We validated the scRNA-seq findings for several tight junction proteins in the *Vglut1⁻ᐟ⁻* retina.

Consistent with the scRNA-seq data, there was a significant reduction in Claudin-5 and ZO-1 protein

localization to EC junctions at P14 *Vglut1⁻ᐟ⁻* compared to the WT retina in both the superficial and deep

plexuses by immunofluorescence (**Figure 4L-O**). Importantly, these phenotypes were not caused by

abnormalities in the NVU assembly, as mural cell (PDGFRβ⁺), glial endfeet (Aquaporin-4; Aqp-4⁺) and

vascular basement membrane (vBM; Laminin) coverage of retinal blood vessels were comparable between

the two genotypes **(Figure S2)**.

We next asked whether the lack of glutamate release by neurons affects the maturation of the

transcellular BRB properties. We injected intravenously albumin-Alexa594 (MW= 60kDa), a protein that

crosses the BRB primarily through a transcellular route [36] into P10 WT and *Vglut1⁻ᐟ⁻* mice and assessed its

extravasation into the retina. The tracer was largely confined within the blood vessels and its intensity in

the parenchyma was very low and identical between the two genotypes **(Figure S3A, B)**. In addition, there

was no difference in endogenous serum IgG leakage between P10 WT and *Vglut1*[-/-] retinas **(Figure S3C)**.

These functional studies are consistent with the scRNA-seq data showing no change in *Mfsd2a* (a gene

required for suppression of transcellular BRB permeability [35]) transcript levels between the two genotypes

(Tables S2, S7). In addition, PLVAP expression, a component of the diaphragms of EC fenestrae normally

found in the immature BRB and peripheral ECs [38], was absent in retinal ECs from both genotypes **(Figure**

S3D) despite a slight downregulation in *Plvap* mRNA in P10 *Vglut1*[-/-] retinal ECs by scRNA-seq **(Figure**

4J; Table S7). Similarly, immunofluorescence analysis showed no difference in retinal EC expression of

Caveolin-1 and Mfsd2a, two critical proteins responsible for regulating EC transcellular permeability

[4,35,39,40], between the two genotypes **(Figure S3E, F).** Thus, paracellular, but not transcellular, BRB

maturation is delayed in both the superficial and deep plexuses of *Vglut1*[-/-] retinas with decreased

extracellular glutamate **(Figure S3G)**.

BRB maturation is accelerated in the absence of cholinergic waves due to an early onset of

glutamatergic activities.

During development, the cholinergic waves persist beyond P10 in the *Vglut1*[-/-] retina lacking

glutamatergic neuronal activities [12]. Conversely, the glutamatergic waves begin earlier (P8) than usual (P10)

in the *Chrnb2*[-/-] retina where the cholinergic waves are abolished **(Figure 5A)** [11]. Recently it was reported

that pharmacological blockade of cholinergic waves before P8 delays deep plexus angiogenesis and BRB

maturation, suggesting a critical role for the cholinergic waves in promoting these processes in the

developing retina [14]. In contrast, our phenotypic analysis of the *Vglut1*[-/-] retinal vasculature implies that the

cholinergic waves are not sufficient to induce paracellular BRB maturation since it is very delayed in the

Vglut1[-/-] retina throughout P14 (**Figure 4**). To test this hypothesis, we examined angiogenesis and the

paracellular BRB integrity in the *Chrnb2*[-/-] retina where cholinergic waves are abolished and glutamatergic

waves begin earlier (P8) than usual [11] **(Figure 5A)**. At P7, before the onset of glutamatergic waves in the

Chrnb2[-/-] retina, there was no difference in the vascularized area of the superficial plexus **(Figure 5B, C),**

and biocytin-TMR leakage **(Figure 5D-F)** between WT and *Chrnb2*[-/-] retinas. In contrast, there was less biocytin-TMR leakage in the parenchyma **(Figure 5J-L),** and increased Claudin-5 coverage of blood vessels **(Figure 5M, N)** in the P10 *Chrnb2*[-/-] retina compared to the WT. However, vascular coverage in either the superficial or deep plexus was not significantly different between P10 WT and *Chrnb2*[-/-] retinas **(Figure 5G-I)**. These data are consistent with the idea that the earlier onset of glutamatergic waves in the *Chrnb2*[-/-] retina is sufficient to promote a precocious BRB maturation, but does not affect deep plexus angiogenesis. Moreover, they demonstrate that glutamatergic neuronal activities, but not the cholinergic waves, are the primary drivers of paracellular BRB maturation in the developing retina.

Deep plexus angiogenesis and paracellular BRB maturation are accelerated in the *Gnat1*[-/-] retina.

To further evaluate the effect of excess glutamate release by retinal neurons, we examined the retinal vascular phenotypes in mice lacking α−Transducin. Phototransduction by rod photoreceptors is a complex cascade, dependent in part on the Gα subunit, Transducin 1 (*Gnat1*). *Gnat1*[-/-] rod photoreceptors remain constitutively depolarized [26,27] leading to increased glutamate release from rods **(Figure 6A, B)**. To understand how excess glutamate release by rod photoreceptors affects retinal angiogenesis, we compared the vascular growth of the superficial and deep plexuses between the WT and *Gnat1*[-/-] retinas. While the vascular coverage of the retina in the superficial plexus was similar between the two genotypes at P10 **(Figure 6C, D)**, both the vascular coverage of the retina in the deep plexus, and its branch-point density were significantly increased in the *Gnat1*[-/-] retina **(Figure 6E-J)**. However, the angiogenesis phenotype was transient, and by P14 there was no difference between the two genotypes **(Figure 6F, G, I, J; Figure S4A)**. Importantly, these patterns were similar in both central and peripheral retina. Consistent with an accelerated deep plexus angiogenesis in the P10 *Gnat1*[-/-], there was also increased EC proliferation (Ki67[+] ECs) in mutant deep vascular plexus compared to the WT **(Figure 6K, L)**.

Next, we asked whether excess glutamate release by rod photoreceptors affects paracellular BRB

maturation. Biocytin-TMR leakage area and fluorescence intensity were reduced in *Gnat1*[-/-] compared to

the WT retinas in both central and peripheral retinas and superficial and deep plexuses at P10 and P14

(Figures 6M-O; S4B-D). Consistent with the functional data, there was a significant increase in Claudin-5

protein localization at BRB tight junctions in the P10 *Gnat1*[-/-] retina compared to the WT **(Figure 6P-R)**.

In summary, retinal deep plexus angiogenesis is accelerated and paracellular BRB integrity is precociously

mature when there is excess glutamate release by the rods.

To address whether aberrant specification of neuronal subtypes or Müller glia in either *Vglut1*[-/-] or

Gnat1[-/-] retinas may be responsible for the vascular phenotypes seen in these mutants, we immunolabelled

and quantified the number of neuronal subtypes and Müller glia. There was no significant difference in the

number of ganglion cells [NeuN[+] [41,42]], amacrine (except for a slight reduction in the *Vglut1*[-/-] retina) and

horizontal cells [Calbindin[+] [43]], photoreceptors [measured by outer nuclear layer (ONL) thickness] among

the three genotypes **(Figure S5A-I)**. Similarly, the number of Sox9[+] Müller glia [44,45] was similar among

WT, *Vglut1*[-/-] and *Gnat1*[-/-] retinas **(Figure S5J, K)**. Importantly, altered levels of extracellular glutamate

did not trigger Müller cell gliosis in *Vglut1*[-/-] and *Gnat1*[-/-] retinas, as there was no upregulation of GFAP

expression [a marker of gliosis [46,47]] in Müller glia processes spanning multiple retinal layers (**Figure S5L**).

GFAP expression was only detected in retinal astrocytes over the GCL (colocalized with PDGFRα) **(Figure**

S5L, M). Therefore, the angiogenesis and BRB phenotypes seen in *Vglut1*[-/-] and *Gnat1*[-/-] retinas are not

caused by defects in neuronal and glial cell type specification.

Finally, we examined whether cholinergic, GABAergic and dopaminergic neuronal populations are

altered in mutant retinas with abnormal glutamatergic activities which may indirectly affect vascular

development. There was no difference in the number of ChAT[+] (cholinergic), GAD67[+] (GABAergic) and

Tyrosine Hydroxylase (TH)[+] (dopaminergic) neurons between WT, *Vglut1*[-/-] and *Gnat1*[-/-] retinas **(Figure**

S6A, B). Overall, our data show that changes in extracellular glutamate does not affect neuronal or glial

subtypes specification or neurons involved in other types of neurotransmission.

Neurons and Müller glia in the inner nuclear layer of the retina respond to altered levels of

extracellular glutamate.

To understand how extracellular glutamate affects retinal angiogenesis and paracellular BRB

maturation, we examined several parameters that could influence these processes. First, we asked whether

glutamate acts directly on retinal ECs. Our scRNA-seq analyses of P10 and P14 WT retinal ECs showed

that neither ionotropic (*Grin1*, *Grin2b*, *Grin2d*, *Gria1*), nor metabotropic (*Grm3*, *Grm5*, *Grm6*, *Grm7*,

Grm8) glutamate receptor subunit transcripts are expressed in developing retinal ECs **(Figure S6C).**

Analysis of a previously published scRNA-seq database of all cell types from the P14 retina [48] also

confirmed our findings that glutamate receptor subunit transcripts are not expressed in P14 retinal ECs.

Thus, glutamate cannot act directly on developing retinal ECs. In contrast, ionotropic and metabotropic

glutamate receptor subunit mRNAs are expressed by the inner nuclear layer (INL) neurons and Müller glia

(Figure S6D), suggesting that extracellular glutamate could act on INL neurons and Müller glia.

To examine whether altered levels of extracellular glutamate are sensed by Müller glia in *Vglut1⁻/⁻*

and *Gnat1⁻/⁻* retinas, we initially measured mean fluorescence intensity of glutamine synthetase (GS), an

enzyme that coverts glutamate to glutamine, and the Glutamate Aspartate Transporter [GLAST, excitatory

aminoacid transporter 1 (EAAT-1)] from immunofluorescence staining. These proteins are known to be

upregulated in Müller glia exposed to excess extracellular glutamate [49,50]. GS and GLAST protein levels

were downregulated in *Vglut1⁻/⁻*, and upregulated in *Gnat1⁻/⁻* retinas, compared to the WT retinas **(Figure**

7A-D). To determine whether altered levels of extracellular glutamate are sensed by neurons in *Vglut1⁻/⁻*

and *Gnat1⁻/⁻* retinas, we measured Glutamate Transporter 1 protein levels [GLT-1, or excitatory aminoacid

transporter 1 (EAAT-2)]. This transporter is expressed primarily in retinal neurons [51], and its expression is

regulated by extracellular glutamate [52]. GLT-1 expression in INL neurons was downregulated in the *Vglut1⁻*

⁻, and upregulated in the *Gnat1⁻ᐟ⁻* retinas, compared to the WT **(Figure 7E-F)**. These data confirm that

Müller glia and INL neurons respond to abnormal extracellular glutamate levels in *Vglut1⁻ᐟ⁻* or *Gnat1⁻ᐟ⁻*

retinas.

Retinal metabolic demand, hypoxia and VEGF-A signaling are not affected in the *Vglut1⁻ᐟ⁻* retina.

Since Müller glia and INL neurons respond to changes in extracellular glutamate levels, we asked

whether neuronal metabolic demand is also affected in *Vglut1⁻ᐟ⁻* and *Gnat1⁻ᐟ⁻* retinas, as alterations in

neuronal metabolic demand trigger vascular abnormalities [reviewed in [4]]. We examined Hexokinase-2

(HK2) and Pyruvate Kinase M-2 (PKM2) protein levels by Western blotting since they are key glycolytic

enzymes of neuronal metabolism readout [53]. The levels of these enzymes were similar among P10 WT,

Gnat1⁻ᐟ⁻ and *Vglut1⁻ᐟ⁻* retinas **(Figure S7A-L)**. Thus, alterations in extracellular glutamate levels caused by

genetic perturbations in glutamate release from neurons do not affect expression of key enzymes for the

neuronal metabolism in the retina.

Neuronal-derived VEGF-A is critical for nervous system function [54] and VEGF-A signaling is a

major regulator of retinal angiogenesis [14,55], prompting us to examine whether VEGF-A levels change in

Vglut1⁻ᐟ⁻ and *Gnat1⁻ᐟ⁻* retinas. VEGF-A protein levels measured with Western blotting were not different

among WT, *Vglut1⁻ᐟ⁻* and *Gnat1⁻ᐟ⁻* retinas **(Figure S7M-P)**. These results are also consistent with our

scRNA-seq data showing no changes in transcripts for VEGF-A signaling components [*Nr4a2*, *Egr3*,

Igfbp3, *Hlx1*, *Crem*, *Per1*, *Mef2c*, *Thbd*, *Ndrg1*, *Dnajb9*, *Mycn* [56]] in *Vglut1⁻ᐟ⁻* retinal ECs **(Figure S7Q;**

Table S2). Since VEGF-A expression depends on tissue hypoxia [57], we measured retinal hypoxia with a

hypoxyprobe, Pimonidazole [58,59]. Consistent with no change in VEGF-A protein and downstream target

transcript levels, there was no difference in tissue hypoxia between *Vglut1⁻ᐟ⁻* and WT retinas **(Figure S7R,**

S). Therefore, the vascular phenotypes observed in mutants with altered extracellular glutamate are not due

to changes in VEGF-A signaling and hypoxia.

Extracellular glutamate levels affect Norrin expression in retinal neurons and Müller glia and as a

consequence endothelial Norrin/β-catenin signaling.

Our scRNA-seq analyses showed that transcript levels for two Norrin/β-catenin signaling effectors

(*Ctnnb1* and *Lef1*), essential for angiogenesis and BRB maturation, were downregulated in P10 *Vglut1*[-/-]

compared to WT retinal ECs **(Figures 2D, F; 4J)**. Among seven key transcription factors (*Foxf2*, *Foxl2*,

Foxq1, *Lef1*, *Ppard*, *Zfp551* and *Zic3*) required for BBB maturation [60], only *Lef1* was downregulated

significantly in P10 *Vglut1*[-/-] retinal ECs, suggesting a specific effect on Norrin/β-catenin signaling. To

address changes in Norrin/β-catenin signaling, we examined *Norrin* (*Ndp*) mRNA expression in *Vglut1*[-/-]

and *Gnat1*[-/-] retinas by fluorescent *in situ* hybridization (FISH). *Norrin* mRNA expression was reduced in

the INL of P10 *Vglut1*[-/-] retinas, and upregulated in the INL of P10 *Gnat1*[-/-] retinas, compared to the WT

(Figure 7G, H). The upregulation of *Norrin* mRNA persisted in the INL of *Gnat1*[-/-] retinas through P14

(Figure S4E). Consistent with our data, analysis of a previously published scRNA-seq database [48]

demonstrated that INL neurons (horizontal cells and a subset of amacrine and bipolar cells) and Müller glia

express high levels of *Norrin* mRNA and glutamate transporters [*Slc1a2* (GLT-1) and *Slc1a3* (GLAST)]

respectively [48] **(Figure 7I)**. These transcriptional data and the response of the INL neurons and Müller glia

to extracellular glutamate in the *Vglut1*[-/-] and *Gnat1*[-/-] retinas (**Figure 7A-F**) suggest that glutamate may act

directly on Müller glia and INL neurons to induce *Norrin* mRNA expression.

To test whether glutamate can induce *Norrin* mRNA expression in Müller glia and INL neurons, we

performed *ex vivo* organotypic cultures of P10 WT retinas in the presence or absence of glutamate

(0.01mM) and measured *Norrin* mRNA expression in the INL by FISH (fluorescence intensity) after 6 and

12 hours in culture (**Figure 7J**). *Norrin* mRNA expression in the INL was upregulated after 6 hours of

glutamate exposure, although it was not statistically significant (p=0.16; **Figure 7K, L**). By 12 hours,

Norrin mRNA levels in the INL were upregulated significantly with glutamate compared to media alone

(Figure 7K, M). Therefore, glutamate induces *Norrin* mRNA expression in Müller glia and INL neurons.

Glutamate-mediated induction of *Norrin* mRNA expression in Müller glia and INL neurons may be

regulated through glutamate uptake via GLAST and GLT-1. To test this directly, we performed *ex vivo*

organotypic cultures of P10 WT retinas with or without glutamate and TFB-TBOA, a pharmacological

inhibitor of GLAST and GLT-1[61-63] **(Figure 7J)**. *Norrin* mRNA expression in the INL was downregulated

significantly with glutamate and TFB-TBOA compared to glutamate alone conditions after 6 and 12 hours

in culture **(Figure 7K-M)**. Importantly, TFB-TBOA alone did not alter *Norrin* mRNA levels in the INL

compared to the media alone **(Figure 7K, M)**, suggesting that TFB-TBOA did not have any non-specific

effect on *Norrin* mRNA expression. Based on these pharmacological data, we conclude that glutamate

uptake by GLAST and GLT-1 likely induces *Norrin* mRNA expression in both Müller glia and INL

neurons.

Next, we examined Norrin/β-catenin signaling activation in *Vglut1*$^{-/-}$ and *Gnat1*$^{-/-}$ retinal ECs. We

immunolabeled P14 WT, *Vglut1*$^{-/-}$ and *Gnat1*$^{-/-}$ retinal sections with Lef1 and quantified the number of

Lef1$^+$ ECs in both the superficial and deep plexuses. There were significantly fewer Lef1$^+$ ECs in the *Vglut1*$^-$

$^{/-}$ retina and significantly higher in the *Gnat1*$^{-/-}$ retina compared to the WT **(Figure 8A, B)**. These data

suggest that angiogenesis and BRB phenotypes in *Vglut1*$^{-/-}$ and *Gnat1*$^{-/-}$ retinas are likely driven by altered

Norrin/β-catenin signaling.

Activation of endothelial Norrin/β-catenin signaling restores retinal angiogenesis and BRB deficits in

***Vglut1*$^{-/-}$ mice.**

If angiogenesis and BRB phenotypes observed in *Vglut1*[-/-] and *Gnat1*[-/-] retinas are driven by altered

endothelial Norrin/β-catenin activity, then activation of Norrin/β-catenin signaling in retinal ECs should

rescue angiogenesis and BRB phenotypes in *Vglut1*[-/-] retinas. We injected intraperitoneally either BIO, a

compound that activates Wnt (Norrin)/β-catenin signaling by inhibiting GSK-3β, or an inactive analog

MeBIO [64], in WT and *Vglut1*[-/-] mice between P6 - P8 and analyzed the vascular and BRB phenotypes in the

retina at P10. BIO-treatment significantly increased the number of Lef1[+] ECs in the *Vglut1*[-/-] retina

compared to the MeBIO-treated *Vglut1*[-/-] retina, whereas MeBIO administration had no effect on endothelial

Lef1 [there were significantly fewer Lef1[+] ECs in the MeBIO-treated *Vglut1*[-/-] compared to the MeBIO-

treated WT retina (**Figure 8C, D**)]. Importantly, there were no Lef1[+] nuclei outside the vasculature with

either treatment **(Figure 8C, D)**. Therefore, BIO-treatment activated Norrin/β-catenin signaling specifically

in *Vglut1*[-/-] retinal ECs.

BIO-treatment significantly increased vascular coverage by the deep plexus in *Vglut1*[-/-] retina to

levels comparable to MeBIO-treated WT retinas **(Figure 8E-G)**. However, the vascular coverage of the

deep plexus was significantly lower in the MeBIO-treated *Vglut1*[-/-] retina compared to the MeBIO-treated

WT retina, consistent with no effect of MeBIO on endothelial Norrin/β-catenin activation (**Figure 8C, D**).

Similarly, BIO-treatment significantly decreased biocytin-TMR leakage area and intensity in the *Vglut1*[-/-]

retina similar to those observed in MeBIO-treated WT levels **(Figure 8H-K),** whereas Me-BIO treatment

had no effect [biocytin-TMR leakage area and intensity were significantly higher in the MeBIO-treated

Vglut1[-/-] retina compared to the MeBIO-treated WT retina (**Figure 8H-K**)]. Interestingly, although BIO

administration in WT mice decreased modestly biocytin-TMR leakage in the retina compared to the MeBIO

administration **(Figure S8C-E)**, the vascular coverage of the deep plexus was reduced significantly in the

BIO-treated compared to the MeBIO-treated WT retina **(Figure S8A, B)**. These data are consistent with

our previous findings that endothelial Norrin/β-catenin activation needs to be finely tuned for proper deep

plexus angiogenesis and BRB maturation in the developing retina [36]. In summary, activation of endothelial

Norrin/β-catenin signaling was sufficient to rescue angiogenesis and BRB phenotypes in the *Vglut1⁻/⁻* retina.

Taken together, our data demonstrate that glutamate released by neurons regulates retinal

angiogenesis and paracellular BRB integrity by modulating Norrin expression in retinal neurons and Müller

glia and endothelial Norrin/β-catenin signaling activation.

DISCUSSION

It is well established that neuronal and vascular development in the CNS, including the retina, are

interdependent. Axonal growth and sprouting angiogenesis use multiple similar guidance cues [65,66]. CNS

vascular growth is also regulated by the neuronal metabolism [43,67,68], as CNS is the most metabolically

active organ system in the body. CNS neurons use a multitude of neurotransmitters such as glutamate, acetyl

choline, dopamine and GABA. However, it is unclear whether specific types of neuronal activities and

neurotransmitters affect CNS vascular growth. A recent study suggested that dopaminergic activities of

ganglion cells regulate retinal superficial plexus angiogenesis [69]. Cholinergic waves propagated by starburst

amacrine cells have also been implicated in the development of retinal deep plexus vessels and BRB

maturation [14]. While glutamate is the predominant neurotransmitter in the retina during vascular

development and in adulthood [70], nothing is known about whether glutamatergic neuronal activities regulate

retinal angiogenesis and BRB maturation. Our study elucidates this critical missing step that links

glutamatergic neuronal activities with vascular development and maturation in the retina - a process that

may also be relevant for several retinal disease. Below, we discuss: a) the role of glutamate in angiogenesis

and barriergenesis, b) the signaling pathways that function downstream of glutamate to promote these

processes, and c) the relevance for the disease.

408    In our study, we used three mouse models to address the role of glutamate in angiogenesis and

409 barriergenesis: a) *Vglut1*[-/-], where retinal photoreceptors and bipolar cells fail to release glutamate into the

410 synaptic cleft, b) *Gnat1*[-/-], where depolarized rod photoreceptors release excess glutamate into the synaptic

411 cleft, and c) *Chrnb2*[-/-], where cholinergic waves are abolished but glutamatergic waves begin at P8 instead

412 of P10. We found that deep plexus angiogenesis and paracellular BRB maturation in the retina are critically

413 linked to the extracellular glutamate levels. These processes are delayed in *Vglut1*[-/-] retinas, where neurons

414 fail to release glutamate. In contrast, deep plexus angiogenesis and paracellular BRB maturation are

415 accelerated in *Gnat1*[-/-] retinas, where constitutively depolarized rods release excessive glutamate. Although

416 the deep plexus angiogenesis, which occurs from P8-P12, was affected in all mutant mouse models, the

417 superficial plexus angiogenesis (P0-P8) was unperturbed. This is likely because the superficial plexus is

418 fully formed by P8, before the onset of glutamatergic neuronal activities in the mouse retina. This

419 developmental time course can also explain why treatment with 2-amino-4-phosphonobutyrate, a

420 compound that blocks glutamatergic waves, from P3- P7 did not affect retinal angiogenesis in a previous

421 study [14], because the treatment was performed before the onset of glutamatergic waves in the developing

422 mouse retina. Similarly, we found that the transcellular, but not paracellular BRB, was also not affected in

423 the *Vglut1*[-/-] retina. This is consistent with previous studies showing that transcellular BRB in the mouse

424 retina is fully mature by P10 [3], whereas paracellular BRB does not fully mature until P18 [36]. Thus, it is

425 conceivable that glutamatergic neuronal activities kick in too late to affect the maturation of the transcellular

426 BRB, but have enough time to influence the maturation of the paracellular BRB.

428    Interestingly, *Vglut1*[-/-] retinas lacking glutamatergic neuronal activities show persistent cholinergic

429 waves beyond P10 [12]. Since deficits in the deep plexus angiogenesis and paracellular BRB integrity persist

430 in the *Vglut1*[-/-] retina throughout P14, our data suggest that the cholinergic waves are not sufficient to

431 promote deep plexus angiogenesis and paracellular BRB maturation. In contrast, paracellular BRB is

432 precociously mature in the *Chrnb2*[-/-] retina where cholinergic waves are abolished and glutamatergic waves

begin earlier (P8) than usual [11]. Overall these data indicate that glutamatergic neuronal activities are the

primary driver of paracellular BRB maturation. We were not able to extend our studies beyond P14 in

Vglut1[-/-] mice, as the pups die around P15 [71]. Notably, the vascular phenotypes, albeit opposite, were more

severe in the *Vglut1*[-/-] retina compared to the *Gnat1*[-/-] retina. This is likely because glutamate release is

abolished from both bipolar cells and photoreceptors (rods and cones) in the *Vglut1*[-/-] retina [12]. In contrast,

only depolarized rod photoreceptors release excess glutamate in the *Gnat1*[-/-] retina [26,27]. This may likely

explain why the vascular phenotypes are less prominent in the P10 *Gnat1*[-/-] retina, and possibly why the

angiogenic phenotype (but not BRB phenotype) in the *Gnat1*[-/-] retina normalizes to wild-type levels by P14.

How does extracellular glutamate trigger angiogenesis and BRB maturation in the developing

retina? It is possible that the vascular phenotypes in *Vglut1*[-/-] and *Gnat1*[-/-] retinas could be due to alterations

in either neuronal numbers which have been shown to affect retinal angiogenesis by titrating the level of

available pro-angiogenic factor VEGF-A [55]. However, we did not observe any significant differences in the

numbers of neuronal subtypes between WT, *Vglut1*[-/-] and *Gnat1*[-/-] retinas. While there was a slight decrease

in calbindin-labelled amacrine cells in the *Vglut1*[-/-] retina, identified classes of amacrine cells, including

cholinergic, GABAergic and dopaminergic neurons, were unaffected in these mutant retinas. These data

suggest that angiogenesis and BRB phenotypes in *Vglut1*[-/-] and *Gnat1*[-/-] retinas are not caused by alterations

in neuronal numbers. Metabolic demand of retinal neurons also modulates angiogenesis [43,67,68]. However,

the expression levels of HK2 and PKM2, two key glycolytic enzymes that are readouts of photoreceptor

cell metabolism [53], were unaffected in both P10 *Vglut1*[-/-] and *Gnat1*[-/-] retinas compared to the WT, in spite

of the prominent vascular phenotypes in these mutant retinas at this developmental stage. Similarly, retinal

hypoxia as well as VEGF-A expression levels were also unaffected in both P10 *Vglut1*[-/-] and *Gnat1*[-/-] retinas

compared to the WT. These data suggest that angiogenesis and BRB maturation abnormalities seen in these

mutant retinas are not due to alterations in neuronal numbers and neuronal metabolic demand or VEGF-A

signaling.

Our data demonstrate that *Norrin* mRNA expression and endothelial Norrin/β-catenin activity are

downregulated in *Vglut1⁻ᐟ⁻* retinas and upregulated in *Gnat1⁻ᐟ⁻* retinas between P10-P14, and therefore are

likely regulated by the extracellular glutamate concentration. Consistent with these findings our scRNA-

seq analyses showed that expression levels of effectors of Norrin/β-catenin pathway [*Ctnnb1* and *Lef1*] are

downregulated in *Vglut1⁻ᐟ⁻* retinal ECs. However, *Norrin* mRNA is expressed in the retina before P10 [20],

and perturbation of Norrin/β-catenin signaling delays superficial plexus angiogenesis as early as P5 [20,25],

well before the onset of glutamatergic neuronal activity. Therefore, early (P5-P8) *Norrin* mRNA expression

in the retina, which is critical for superficial plexus angiogenesis, is likely regulated by a glutamate-

independent mechanism [48]. Our data in conjunction with a previous study [48] also suggest that *Norrin* mRNA

is expressed by several cell types in the developing retina including Müller glia and INL neurons (horizontal

cells and subsets of amacrine and bipolar cells). Using *ex vivo* organotypic cultures, we also show that

glutamate can directly induce *Norrin* mRNA expression in the Müller glia and INL neurons, and this occurs

via glutamate uptake by GLAST and GLT-1, two glutamate transporters expressed in Müller glia and INL

neurons. We used a high concentration of TFB-TBOA (GLAST and GLT-1 inhibitor) in our study in order

to ensure that glia- and neuron-mediated glutamate uptake is completely inhibited in our *ex vivo* culture

system. Importantly, the presence of TFB-TBOA in the media in the absence of glutamate did not

significantly alter *Norrin* mRNA expression compared to the retinas with neither glutamate nor TFB-TBOA

in the media, confirming that TFB-TBOA does not affect *Norrin* mRNA expression in the retina. Future

studies will elucidate the relative contribution of Müller glia- or INL neurons-derived Norrin, regulated by

glutamate, for the deep plexus angiogenesis and paracellular BRB maturation. Importantly,

pharmacological activation of endothelial Norrin/β-catenin signaling was sufficient to rescue both deep

plexus angiogenesis and paracellular BRB integrity in the *Vglut1⁻ᐟ⁻* retina, consistent with the idea that

Norrin/β-catenin signaling functions downstream of extracellular glutamate to promote these processes.

Our scRNA-seq data also show a slight downregulation of *Unc5b* in *Vglut1⁻ᐟ⁻* retinal ECs. Unc5b has been

shown to modulate Wnt/β-catenin signaling in CNS ECs [72], suggesting the possibility that the cross-talk

between these two pathways may be affected in *Vglut1*$^{-/-}$ retinal ECs. Interestingly, none of the downstream

targets of VEGF-A signaling was affected in *Vglut1*$^{-/-}$ retinal ECs at the transcriptome level. Thus, VEGF-

A signaling pathway does not appear to be regulated by glutamate in the developing retina.

Taken together our data suggest a new mechanism by which glutamatergic neuronal activity regulate

retinal angiogenesis and BRB maturation in the developing retina (**Figure 8L**). We propose that

extracellular glutamate, released either during spontaneous glutamatergic waves or light-dependent

glutamatergic synaptic activity of photoreceptors from P8 onwards, signals to Müller glia and INL neurons.

Glutamate signaling induces Norrin mRNA and protein expression and secretion by Müller glia and INL

neurons. Norrin, in turn, activates endothelial Norrin/β-catenin signaling in ECs which promotes

angiogenesis in the deep plexus and paracellular BRB maturation in both superficial and deep plexuses of

the retina (**Figure 8L**). Our genetic data in *Vglut1*$^{-/-}$ (no glutamate release into the synaptic cleft), *Gnat1*$^{-/-}$

(excess glutamate release into the synaptic cleft) and *Chrnb2*$^{-/-}$ (early onset of the glutamatergic waves),

and the pharmacological rescue of *Vglut1*$^{-/-}$ phenotypes by activation of endothelial Norrin/β-catenin

signaling, are consistent with the proposed model that extracellular glutamate levels control deep plexus

angiogenesis and barriergenesis. Although, the exact mechanisms by which glutamate mediates these

effects are not fully clear, our single-cell RNA-seq data together with the published datasets [48] suggest that

glutamate cannot act directly on ECs as they do not express either ionotropic or metabotropic glutamate

receptor subunits or glutamate transporters. The published transcriptional data of the glutamate receptor

and transporter expression [48], and our analysis of the response of the INL cells (neurons and Müller glia) to

extracellular glutamate (i.e, changes in GS, GLAST, GLT-1 protein levels) in the *Vglut1*$^{-/-}$ and *Gnat1*$^{-/-}$

retinas, suggest that glutamate could directly act on Müller glia and INL neurons to induce *Norrin* mRNA

expression. These cells express subsets of ionotropic or metabotropic glutamate receptors and glutamate

transporters that make them exquisite sensors of extracellular glutamate levels. In addition, *Norrin* mRNA

is predominantly expressed by the INL cells during vascular development and maturation. Future studies will address in more detail the molecular mechanisms by which glutamate affects Norrin expression in the INL cells and specific contributions of INL cell-derived Norrin to angiogenesis and barriergenesis.

Structural and functional abnormalities in neurons or glial cells precede vascular pathologies in multiple CNS neurological diseases with prominent vascular involvement, including those in the retina. For example, in diabetic retinopathy there is an early loss of neurovascular coupling, impaired response of photoreceptors to light, gradual neurodegeneration, gliosis and neuroinflammation before any observable vascular pathologies [5-7]. Moreover, the early phases of either age-related macular degeneration, or retinitis pigmentosa, characterized by changes in synaptic activity and metabolic deficits in Müller glia [73-75], may also affect blood vessel function leading to neurodegeneration. Although vascular pathologies, including aberrant angiogenesis and BRB breakdown, are found in diabetic retinopathy and proliferative vitreoretinopathy that cause acquired adult blindness, as well as retinopathy of prematurity which leads to blindness in premature infants [1,76-78], the molecular mechanisms that link neuronal or glial abnormalities with vascular deficits remain elusive. Our findings not only extend our understanding of the molecular mechanisms by which vascular development and homeostasis occurs in the CNS [reviewed in [4]] including the role that neuronal activity plays in these processes both in the cortex [79] and retina [14,17,18], but also have important implications to identify putative interventions to treat CNS neurovascular pathologies, including those in the retina from diabetic retinopathy, retinitis pigmentosa to age-related macular degeneration.

ACKNOWLEDGEMENTS

We thank Dr. Vilas Menon (Columbia University Irving Medical Center) for his consultation on computational analyses and Dr. Eduardo Solessio (Upstate Medical University, State University of New York) for his consultation about *Gnat1*[-/-] retinal physiology. S.B., S.S., and D.A. are supported by grants from the National Eye Institute (R01EY033994 and K99EY033909), National Heart Lung and Blood

Institute (R61HL159949, R33HL159949), National Institute of Aging (RF1AG078352), National Institute

of Neurological Disorders and Stroke (R21 NS130265), the National Multiple Sclerosis Society (RG-1901-

33218) and in part by a gift donation. W.J.B. and G.B. are supported by grants from the National Eye

Institute (R01 EY033994 and R21 EY034696) and in part by funds from an unrestricted grant from

Research to Prevent Blindness to the Department of Ophthalmology and Visual Sciences, Upstate Medical

University.

AUTHOR CONTRIBUTIONS

Conceptualization: S.B., W.J.B., D.A.; Methodology: S.B., D.A., G.B.; Software: S.S.; Validation: S.B.,

G.B; Formal analysis: S.B., S.S., G.B., P.A., D.J.; Investigation: S.B., S.S., G.B.; Resources: W.J.B., D.A.;

Data curation: S.B., S.S., G.B.; Writing - original draft: S.B., D.A.; Writing - review & editing: S.B., S.S.,

G.B., W.J.B., D.A.; Visualization: S.B.; Supervision: W.J.B., D.A., Project administration: D.A., Funding

acquisition: W.J.B., D.A.

DECLARATION OF INTEREST

The authors declare no competing or financial interests.

STAR ★ METHODS

Detailed methods are provided in the online version of this paper and include the following:

KEY RESOURCES TABLE

RESOURCE AVAILABILITY

o   Lead contact

o   Materials availability

o   Data and code availability

EXPERIMENTAL MODEL AND SUBJECT DETAILS

o   Mice

METHOD DETAILS

o   Immunohistochemistry

o   Western blot

o   Antibodies

o   *In situ* hybridization

o   Barrier permeability assays

o   Vessel perfusion assay

o   Hypoxia measurement

o   Retinal endothelial cell isolation

o   Retinal organotypic culture

o   Single-cell RNA-sequencing analysis

o   BIO and MeBIO treatment

QUANTIFICATION AND STATISTICAL ANALYSIS

o   Image quantification

o   Statistical analysis

 STAR ★ METHODS

 KEY RESOURCES TABLE

Reagent or resource	Source	Identifier
Primary antibodies		
Rat monoclonal anti-CD31	BD Biosciences	Cat #553370, RRID:AB_394816
FITC rat monoclonal anti-mouse CD31	BD Pharmingen	Cat #553372, RRID:AB_394818
APC rat monoclonal anti-mouse CD31	BD Pharmingen	Cat #551262, RRID:AB_398497
Rabbit monoclonal anti-Ki67	Abcam	Cat #ab16667, RRID:AB_302459
Rabbit monoclonal anti-Cleaved Caspase-3	Cell Signaling Technology	Cat #9664S, RRID:AB_2070042
Goat polyclonal anti-DLL4	R & D Systems	Cat #AF1389, RRID:AB_354770
Rabbit polyclonal anti-Claudin-5	ThermoFisher Scientific	Cat #34-1600, RRID:AB_2533157
Mouse monoclonal anti-ZO-1	ThermoFisher Scientific	Cat #33-9100, RRID:AB_2533147
Rabbit polyclonal anti-SOX9	Millipore Sigma	Cat #ABE571, Lot #2535307
Mouse monoclonal anti-Glutamine synthetase	Millipore Sigma	Cat #MAB302, RRID:AB_2110656

Rabbit monoclonal anti-EAAT1 (GLAST)	Abcam	Cat #ab181036, RRID:AB_2885103
Rabbit polyclonal anti-EAAT2 (GLT-1)	Abcam	Cat #ab41621, RRID:AB_941782
Rabbit polyclonal anti-GFAP	Millipore Sigma	Cat #AB5804, RRID:AB_2109645
Rat monoclonal anti-CD140a (PDGFRα)	BD Biosciences	Cat #558774, RRID:AB_397117
Rabbit monoclonal anti-LEF1	Cell Signaling Technology	Cat# 2230s, RRID:AB_823558
Goat polyclonal anti-PDGFRβ	R & D Systems	Cat #AF1042, RRID:AB_2162633
Rabbit polyclonal anti-Laminin	Millipore Sigma	Cat #L9393, RRID:AB_477163
Rabbit polyclonal anti-Collagen IV	Abcam	Cat #ab6586, RRID:AB_305584
Rabbit polyclonal anti-Aquaporin-4	Millipore Sigma	Cat #A5971, RRID:AB_258270
Rabbit polyclonal anti-Caveolin-1	Abcam	Cat #ab18199, RRID:AB_444314
Rabbit polyclonal anti-PLVAP	Proteintech	Cat #14452-1-AP, RRID:AB_2166100
Mouse monoclonal anti-NeuN	Millipore Sigma	Cat #MAB377, RRID:AB_2298772

Rabbit polyclonal anti-Calbindin	Millipore Sigma	Cat #AB1778, RRID:AB_2068336
Mouse monoclonal anti-PKCα	Santa Cruz Biotechnology	Cat #sc-8393, RRID:AB_628142
Goat polyclonal anti-Podocalyxin	R & D Systems	Cat #AF1556-SP, RRID:AB_354858
Goat polyclonal anti-Choline Acetyltransferase (ChAT)	Novus Biologicals	Cat #NBP1-30052, RRID:AB_1968484
Mouse monoclonal anti-GAD67	Millipore Sigma	Cat #MAB5406, RRID:AB_2278725
Rabbit polyclonal anti-Tyrosine Hydroxylase (TH)	Abcam	Cat #ab6211, RRID:AB_2240393
Rabbit polyclonal anti-Mfsd2a	Gift from Dr. Chenghua Gu (Cell Signaling Technology)	Cat #14-87 Mfsd2a, RRID:AB_2617168
Rabbit monoclonal anti-Hexokinase II	Abcam	Cat #ab209847, RRID:AB_2904621
Rabbit monoclonal anti-Pyruvate kinase M2	Cell Signaling Technology	Cat #4053, RRID:AB_1904096
Rabbit polyclonal anti-VEGF-A	Millipore Sigma	Cat #SAB5700629
Mouse monoclonal anti-GAPDH	Proteintech	Cat #60004-1-Ig, RRID:AB_2107436
Chemicals, peptides, and recombinant proteins		

Biocytin-TMR (5-(and-6)-tetramethylrhodamine biocytin)	ThermoFisher Scientific	Cat #T12921
Bovine Serum Albumin (BSA), AlexaFluor 594 conjugate	ThermoFisher Scientific	Cat #A13101
Fluorescein-labeled *Griffonia* (*Bandeiraea*) *Simplicifolia* Lectin I (GSL I, BSL I)	Vector Laboratories	Cat #FL-1101, RRID:AB_2336490
Rhodamine-labeled *Griffonia* (*Bandeiraea*) *Simplicifolia* Lectin I (GSL I, BSL I)	Vector Laboratories	Cat #RL-1102, RRID:AB_2336492
Hypoxyprobe™-1 Kit (100 mg of Pimonidazole HCl plus 1 ml of 11.23.22.r rat monoclonal antibody	Hypoxyprobe	Cat #HP1-100Kit
6-BIO ((2*Z*,3*E*)-6-bromoindirubin- 3-acetoxime, BIO-Acetoxime)	Calbiochem	Cat #361551
MeBIO (1-methyl-BIO)	Calbiochem	Cat #361556
Propidium iodide	ThermoFisher Scientific	Cat #P1304MP
TFB-TBOA	TOCRIS	Cat #2532
CD16/CD32 Fc block	BD Pharmingen	Cat #553141
Experimental Models: Organisms/Strains		
Vglut1[+/-] (C57/BL6J background)	Gift from Dr. Robert H. Edwards [71]	N/A

Gnat1^{+/-} (C57/BL6J background)	Gift from Dr. William J. Brunken [26]	N/A
Chrnb2^{+/-} (C57/BL6J background)	Gift from Dr. Marina Picciotto [11]	N/A
Oligonucleotides		
Vglut1-WT forward: 5'CCAAGCAAGGTTAAGCCTAG3'	IDT	N/A
Vglut1-WT reverse: 5'GGTGAATTTGGAAAAGAGC3'	IDT	N/A
Vglut1-KO forward: 5'TCGCAGCGCATCGCCTTCTA3'	IDT	N/A
Vglut1-KO reverse: 5'GCCACACGTTTCTTCTGAGG3'	IDT	N/A
Gnat1-WT forward: 5'GCCATCATCTACGGCAACACTCTGC3'	IDT	N/A
Gnat1-WT reverse: 5'GCCGGCGGAATTGAGCTGGTA3'	IDT	N/A
Gnat1-KO forward: 5'GCCGGTGAATCCGGGAAGAGC3'	IDT	N/A
Gnat1-KO reverse:	IDT	N/A

5'GGGCTGCTAAAGCGCATGCTC 3'		
Chrnb2-WT forward: *5*'CGGAGCATTTGAACTCTG AGCAGTGGGGTCGC3'	IDT	N/A
Chrnb2-WT reverse: 5'CTCGCTGACACAAGGGCT GCGGAC3'	IDT	N/A
Chrnb2-KO forward: *5*'CACTACGTCTGAACG TCGAAAACCCG3'	IDT	N/A
Chrnb2-KO reverse: 5'CGGGCAAATAATATC GGTGGCCGTGG3'	IDT	N/A
Softwares and algorithms		
ImageJ/Fiji	NIH	https://imagej.net/software/fiji/downloads
Zen microscopy software	ZEISS	https://www.zeiss.com/microscopy/en/products/software/zeiss-zen.html#packages
Seurat	Seurat	https://satijalab.org/seurat/
Tidyverse	Tidyverse	https://www.tidyverse.org

DAVID	DAVID bioinformatics resources	https://david.ncifcrf.gov
Prism 9	GraphPad	https://www.graphpad.com/ scientific-software/prism/
Other		
RNA *in situ* probe for mouse *Ndp* (Norrin)	[36]	N/A
C57B6 wild-type P14 retina by drop-seq (single-cell RNA-sequencing database)	[48]	https://singlecell.broadinstitute.org/single_cell/study/SCP301/c57b6-wild-type-p14-retina-by-drop-seq

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to the lead contact, Dritan

Agalliu (da191@cumc.columbia.edu).

Materials availability

This study did not generate new unique reagents.

Data and code availability

All data are available in the manuscript and the supplementary information. The codes used in this study

have been previously published and are freely available (see Key Resources Table). Any additional

information required to reanalyze the data reported in this study will be available upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mice

All animal usage and protocols were approved by the Institutional Animal Care and Use Committee (IACUC) at Columbia University Irving Medical Center. Experimental procedures were optimized to minimize animal stress and the number of animals used for experiments. Mice were maintained on 12 light/12 dark cycle. Both sexes (males and females) were used for experiments and analyses. All mice were maintained in C57/BL6J background. The *in vivo* BRB permeability assays were performed either at P10 or P14. Generation of *Vglut1*[-/-] mice was described before [71]. These mice were bred as heterozygous and gave birth in Mendelian ratios. Age-matched littermate WT and *Vglut1*[-/-] mice were used for experiments (*Vglut1*[-/-] mice die around P15). Generation of *Gnat1*[-/-] mice was described before [26]. These mice were bred as heterozygous and gave birth in Mendelian ratios. Age-matched littermate WT and *Gnat1*[-/-] mice were used for experiments. *Gnat1*[-/-] mice are viable and fertile. Generation of *Chrnb2*[-/-] mice was described before [11]. These mice were bred as *Chrnb2*[+/-] and gave birth in Mendelian ratios. Age-matched littermate WT and *Chrnb2*[-/-] mice were used for experiments. *Chrnb2*[-/-] mice are viable and fertile.

METHOD DETAILS

Immunofluorescence

Unless otherwise specified, isolated tissues (retina, brain and liver) were fixed in 4% PFA for 2 hours for immunostaining. The exceptions were retinal flat-mount staining for Laminin, in which cases retinas were fixed for 10 minutes in 2% PFA. For retinal flat-mounts, tissues were washed with 1X PBS following PFA fixation and blocked overnight in blocking buffer (5% goat or donkey serum, 0.3% Triton X-100 in 1× PBS) at 4°C. Samples were then incubated with primary antibodies in antibody-diluting solution (5% goat or donkey serum; 0.01% Triton X-100 in 1X PBS) for 48h at 4°C, washed, and incubated with secondary antibodies for 24 hours. For radial section staining, samples were washed in 1 × PBS,

cryoprotected in 30% sucrose overnight, embedded in optimal cutting temperature compound (OCT;

Tissue-Tek; Torrance, CA, USA), and sectioned at a thickness of 12 μm. Tissue sections were blocked for

2h at room temperature followed by primary antibody incubation overnight at 4°C. Following 1X PBS

washes, sections were incubated with secondary antibodies for 4h at room temperature. Samples were

imaged in room temperature using a Zeiss LSM700 confocal microscope [ZEISS 20X air (Objective Plan-

Apochromat 20x/0.8 M27; numerical aperture: 0.8) or 40X water (Objective C-Apochromat 40x/1.2 W

autocorr M27; numerical aperture: 1.2) objectives; ZEN image acquisition software] and analyzed using

Fiji software.

Western blotting

Retinas were collected from mice after perfusion with PBS. Six retinas from three mice were

combined for each sample. Collected tissue was homogenized in lysis buffer containing protease and

phosphatase inhibitors using a Dounce homogenizer followed by sonication. Protein concentration was

assayed by BCA (Bio-Rad) and resolved on SDS-PAGE. Protein levels in P10 retinas were measured by

fluorescent western blot analysis [4-15% or 7.5% Mini-PROTEAN TGX gels (Bio-Rad)] and quantitation

was performed using the LICOR system, as previously described. Fluorescent quantification of protein

levels was performed using the Odyssey SA infrared imaging system. Values are displayed as protein levels

corrected with GAPDH (loading control) and normalized to the WT values.

Antibodies

Primary antibodies used were rat anti-CD31 (BD Biosciences, Cat #553370, 1:250), rabbit anti-

Ki67 (Abcam, Cat #ab16667, 1:500), rabbit anti-cleaved Caspase-3 (Cell Signaling Technology, Cat

#9664S, 1:500), Goat anti-DLL4 (R & D Systems, Cat #AF1389, 1:200), rabbit anti-Claudin-5

(ThermoFisher Scientific, Cat #34-1600, 1:500), rabbit anti-SOX9 (Millipore Sigma, Cat #ABE571, 1:500),

mouse anti-Glutamine Synthetase (Millipore Sigma, Cat #MAB302, 1:500), rabbit anti-EAAT1 (Abcam,

Cat #ab181036, 1:250), rabbit anti-EAAT2 (Abcam, Cat #ab41621, 1:250), rabbit anti-GFAP (Millipore

Sigma, Cat #AB5804, 1:500), rat anti-CD140a (BD Biosciences, Cat #558774, 1:100), rabbit anti-LEF1

(Cell Signaling Technology, Cat #2230s, 1:100), goat anti-PDGFRβ (R & D Systems, Cat #AF1042, 1:250),

rabbit anti-Laminin (Millipore Sigma, Cat #L9393, 1:500), rabbit anti-Aquaporin-4 (Millipore Sigma, Cat

#A5971, 1:250), rabbit anti-Caveolin-1 (Abcam, Cat #ab18199, 1:500), rabbit anti-PLVAP (Proteintech,

Cat #14452-1-AP, 1:250), mouse anti-NeuN (Millipore Sigma, Cat #MAB377, 1:500), rabbit anti-

Calbindin (Millipore Sigma, Cat #AB1778, 1:250), mouse anti-PKCα (Santa Cruz Biotechnology, Cat #sc-

8393, 1:1000), rabbit anti-Mfsd2a (a gift from Dr. Chenghua Gu, 1:300), goat anti-Podocalyxin (R & D

Systems, Cat #AF1556-SP, 1:500), goat anti-ChAT (Novus Biologicals, Cat #NBP1-30052, 1:100), mouse

anti-GAD67 (Millipore Sigma, Cat #MAB5406, 1:2000), rabbit anti-Tyrosine Hydroxylase (Abcam, Cat

#ab6211, 1:1000), rabbit anti-Hexokinase II (Abcam, Cat #ab209847, 1:1000), rabbit anti-Pyruvate Kinase

M2 (Cell Signaling Technology, Cat #4053, 1:1000), rabbit anti-VEGF-A (Millipore Sigma, Cat

#SAB5700629, 1:2000), mouse anti-GAPDH (Proteintech, Cat #60004-1-Ig, 1:50000), FITC rat anti-

mouse CD31 (BD Pharmingen, Cat #553372, 1:200), and APC rat anti-mouse CD31 (BD Pharmingen, Cat

#551262, 1:200). Secondary antibodies used were: goat anti-rat Alexa Fluor 594 (A11007), goat anti-rabbit

Alexa Fluor 594 (A11012), goat anti-rat Alexa Fluor 488 (A11006), goat anti-rabbit Alexa Fluor 488

(A11034), donkey anti-rat Alexa Fluor 594 (A21209), donkey anti-rabbit Alexa Fluor 594 (A32754),

donkey anti-rat Alexa Fluor 488 (A21208), donkey anti-rabbit Alexa Fluor 488 (A21206), donkey anti-rat

Alexa Fluor 647 (A48272), donkey anti-rabbit Alexa Fluor 647 (A32795), donkey anti-goat Alexa Fluor

594 (A32758) and donkey anti-goat Alexa Fluor 488 (A11055) (all 1:250; ThermoFisher

Scientific). IRDyes 680 and 800 (1:20000; LI-COR) were used as secondary antibodies for Western blot.

***In situ* hybridization**

Both alkaline phosphatase and fluorescent *in situ* hybridization of fresh-frozen (in OCT) retinal

sections using antisense mRNA probe for full-length mouse *Ndp* (Norrin) were performed as described

before [36]. The samples were imaged using either a ZEISS Axio Imager (alkaline phosphatase) or a ZEISS

LSM700 confocal (fluorescent) microscope and analyzed using Fiji software.

Barrier permeability assays

Mice were anesthetized using isoflurane and injected with either 1% Biocytin-5(and-6-)-

Tetramethylrhodamine (biocytin-TMR: ThermoFisher Scientific, Cat #T12921) or 1% Bovine Serum

Albumin (BSA) - AlexaFluor 594 conjugate (albumin-Alexa594: ThermoFisher Scientific, Cat #A13101)

into the tail veins. After 40 minutes in circulation, mice were re-anesthetized and perfused with 1X PBS via

the cardiac route. Retinas and livers were isolated and fixed in 4% PFA for 2 hours prior to immunostaining

with FITC-conjugated Lectin. Since blood vessels in the liver do not have a barrier, Biocytin-TMR or

Albumin-594 fluorescence levels in the livers of corresponding animals were used as internal controls for

measurement of biocytin-TMR and albumin-Alexa594 fluorescence levels in the retinas as described [36,37].

Vessel perfusion assay

P14 pups were anesthetized using isoflurane and injected with 75 µl of Rhodamine-labeled Lectin

(Vector Laboratories, Cat #RL-1102) into the left ventricles of the heart. Lectin was allowed to circulate

for 5 minutes under anesthesia. After that, retinas were isolated and fixed in 4% PFA for 2 hours prior to

immunostaining with vascular markers.

Hypoxia measurement

Hypoxyprobe staining was performed as per the vendor's (Hypoxyprobe, Cat #HP1-100Kit)

instructions. Briefly, the probe was dissolved in saline and administered into P10 pups by intraperitoneal

injection at a concentration of 60 mg/kg body weight. After 90 minutes, the pups were anesthetized using

isoflurane and perfused for 4 minutes with ice-cold sterile 1X PBS. After this, retinas were harvested,

washed with 1X PBS following PFA fixation (4% PFA for 2 hours) and blocked overnight in blocking

buffer at 4°C. Samples were then incubated with primary antibody (11.23.22.r) provided with the kit in

antibody-diluting solution (1:50) for 48h at 4°C. Retinas were then washed with 1X PBS, and incubated

with secondary donkey anti-rat Alexa594 antibody and Fluorescein-labeled Lectin for 24h.

Retinal organotypic culture

P10 WT pups were perfused with sterile 1X PBS under isoflurane-induced anesthesia, and eyeballs

were enucleated. Eyeballs were dissected in sterile neurobasal media (ThermoFisher Scientific, Cat

#21103049) supplemented with serum free B-27 (ThermoFisher Scientific, Cat #17504044). Corneas and

lenses were removed from the eyeballs, and eyecups were placed in the upper chambers of sterile 12 mm

Transwell with 0.4 μm Pore Polycarbonate Membrane Insert (Corning Life Sciences, Cat #3401). Eyecups

were divided into four groups based on the media (neurobasal media + B-27) composition: 1) without

glutamate, 2) with 0.1 mM glutamate, 3) without glutamate + 100 μm TFB-TBOA, 4) with 0.1 mM

glutamate + 100 μm TFB-TBOA. The upper chambers of the transwell received 350 μl of the corresponding

media, and the lower chambers of the transwell received 650 μl of the corresponding media. Under these

conditions, the eyecups were placed in the sterile incubator for either 6 hours or 12 hours at 37° C and with

5% CO_2. After 6 or 12 hours, the eyecups were removed from trans-well and fresh-frozen in OCT for future

sectioning and fluorescent *in situ* hybridization.

Retinal endothelial cell isolation

P10 WT and *Vglut1*[-/-] as well as P14 WT mice were anesthetized using isoflurane and perfused for

4 minutes with sterile 1X PBS. The eyes were enucleated and retinas were dissected in Earl's balanced salt

solution (EBSS). Retinas were dissociated in a single-cell suspension using a published protocol [80]. Once

the retinas were dissociated into single cells, it was resuspended in 200 ml of CD16/CD32 Fc block (BD

Pharmingen, Cat #553141, 1:200 in FACS buffer) and incubated at room temperature for 15 minutes. The

samples were washed with 2 ml of FACS buffer, with 100 ml kept aside for use as unstained and

fluorescence minus one (FMO) controls. The cells were then resuspended in 200 ml of anti-mouse CD31

[either FITC rat anti-mouse CD31 (BD Pharmingen, Cat #553372, 1:200) or and APC rat anti-mouse CD31

(BD Pharmingen, Cat #551262), 1:200 in FACS buffer)] antibody to label ECs, covered in aluminum foil

and incubated on ice for 60 mins. Following antibody labeling, the samples were washed twice with 2 ml

of FACS buffer and resuspended in 400 ml of propidium iodide (Thermofisher, Cat. #P1304MP; 1/10,000

in FACS buffer). Gates for the surface stain were set using the unstained and FMO controls, and either

CD31-APC$^+$ or CD31-FITC$^+$ ECs were sorted out using the BD FACSAria Flow Cytometer (Columbia

Stem Cell Initiative Flow Cytometry Core) for single-cell RNA-sequencing.

Single-cell RNA-sequencing (scRNA-seq) analysis

Total of 1104 ECs for P10 WT retinas and 1046 ECs for P10 *Vglut1*$^{-/-}$ retinas were analyzed for

differential gene expression analyses (**Table S1**). The 10x Genomics Chromium platform [81,82] was used to

generate low-depth data (~2000 genes/cell), and was processed using the Cell Ranger analysis pipeline to

align reads and generate feature-barcode matrices. The Seurat R package [83] was implemented to read the

output of the Cell Ranger pipeline and merge cells from all the samples (WT and mutant) into a single R

object. The standard pre-processing workflow for scRNAseq data was carried out in Seurat, which consisted

of the selection and filtration of cells based on quality control metrics (e.g., number of unique genes detected

in each cell > 200, total number of molecules detected within a cell > 1,000 and < 50,000, percentage of

reads within a cell that map to the mitochondrial genome < 20, etc.), data normalization and scaling, and

the detection of highly variable features. Next, linear dimensionality reduction (PCA) was performed on

the scaled data using the previously determined highly variable features, an alternative heuristic method

('Elbow plot') was implemented to determine the 'dimensionality' of the dataset, and cells were clustered by

applying the weighted shared nearest-neighbor graph-based clustering method using the first 20 PCs [84].

Lastly, the 20 PCA dimensions were further reduced into two-dimensional space using the non-linear

dimensional reduction technique, uniform manifold approximation and projection for dimension reduction

(UMAP), to visualize and explore the scRNA-seq dataset and each of its unique EC clusters. All the ECs

isolated from either genotype were plotted in the UMAP. Differential gene expression analyses were

performed for all ECs from either genotype, and not on specific clusters. From the gene signatures that are

uniquely present in mutant ECs compared to WT ECs, and validated by fluorescent in situ hybridization

and/or immunofluorescence. Gene ontology (GO) enrichment analysis was performed using DAVID

bioinformatics database to elucidate significantly up- and downregulated pathways in mutant ECs.

BIO and MeBIO treatment

The *in vivo* injections of 6-BIO ((2Z,3E)-6-bromoindirubin- 3-acetoxime, BIO-Acetoxime;

Calbiochem, Cat #361551) and control MeBIO (1-methyl-BIO; Calbiochem, Cat #361556) were performed

as previously published [64]. Pups were anesthetized with 1% isoflurane. Intraperitoneal injections of BIO

and MeBIO were performed at a concentration of 5 mg/kg/day in sterile PBS from P6 to P8. Injected pups

were re-anesthetized and perfused with 1X PBS to harvest tissues for analyses at P10.

QUANTIFICATION AND STATISTICAL ANALYSIS

Image quantification

For retinal flat-mounts, each measurement was made in at least 3 quadrants per sample. The

measurements were done separately in the central (near the optic nerve-head) and peripheral (near the edge)

retina. For sections, at least three sections were imaged per sample. In each case, multiple representative

images were acquired and averaged. To measure vascular area and branch point density, area positive for

CD31 or Lectin was measured from each image in Fiji software (National Institutes of Health) using signal

masking. Branch point numbers were normalized to unit area ($1 mm^2$). To measure number of proliferative

or apoptotic ECs, the number of $Ki67^+$ or cleaved Caspase-3^+ ECs, respectively, were normalized to unit

$CD31^+$ vascular area ($1 mm^2$). The area of Biocytin-TMR leakage was done separately in the central and

peripheral retina. The intensity of biocytin-TMR and albumin-Alexa594 outside of retinal vessels was

measured as described [36]. Biocytin-TMR and albumin-Alexa594 signal intensities in the retina and cerebella

were normalized to those of livers. For vascular area positive for Claudin-5 and Caveolin-1, area positive

for Lectin was selected from each image in Fiji software using signal masking, followed by measuring the

percentage of Lectin$^+$/Claudin-5$^+$ or Lectin$^+$/Caveolin-1$^+$ in corresponding images. To measure the number

of ganglion (NeuN$^+$), amacrine, horizontal (Calbindin$^+$) and bipolar cells (PKCα^+), the number of cells were

normalized to the length of either ganglion cell layer (GCL; for ganglion cells) or inner nuclear layer (INL;

for amacrine, horizontal and bipolar cells). To measure the number of photoreceptors, the thickness of the

outer nuclear layer (ONL) was quantified as a function of photoreceptor number. To measure the number

of Müller glia, the number of Sox9$^+$ cells were normalized to the length of INL. To measure the relative

expression of GS and GLAST, the M.F.I were calculated for each field and data was presented as values

normalized to the number of Sox9$^+$ Müller glia. To measure *Ndp* expression, mean fluorescence intensity

(M.F.I) of *Ndp* in the INL was calculated with histogram analysis in Fiji software and normalized to the

WT values. For Lef1$^+$ EC quantification, the number of Lef1$^+$ ECs was normalized to unit CD31$^+$ or Lectin$^+$

vascular area (1 mm^2). For western blots, values are displayed as target protein levels corrected with

GAPDH (loading control) and normalized to the WT values using Odyssey SA infrared imaging system.

Statistical analysis

Retinas from 3-5 different animals (for *in vivo* experiments) were used for statistical analysis. For

western blots, retinas from 3 animals/genotype were combined into one sample, and 4 samples/genotype

was used for quantification. To test for statistical significance of any differences, unpaired Student's *t* test

or one-way ANOVA was performed using GraphPad Prism. A value of *P* < 0.05 was considered

statistically significant. For scRNA-seq analysis, we considered transcripts with adjusted p value < 0.05,

and >1.2-fold ($\log_2 fc > 0.25$ and $\log_2 fc < -0.25$) difference in expression as significant. For GO analysis,

Gene sets with false discovery rate (FDR) < 0.01 were considered significant.

REFERENCES

Klaassen, I., Van Noorden, C. J. & Schlingemann, R. O. Molecular basis of the inner blood-retinal barrier and its breakdown in diabetic macular edema and other pathological conditions. *Prog Retin Eye Res* **34**, 19-48, doi:10.1016/j.preteyeres.2013.02.001 (2013).

Zhao, Z., Nelson, A. R., Betsholtz, C. & Zlokovic, B. V. Establishment and Dysfunction of the Blood-Brain Barrier. *Cell* **163**, 1064-1078, doi:10.1016/j.cell.2015.10.067 (2015).

Chow, B. W. & Gu, C. The molecular constituents of the blood-brain barrier. *Trends Neurosci* **38**, 598-608, doi:10.1016/j.tins.2015.08.003 (2015).

Biswas, S., Cottarelli, A. & Agalliu, D. Neuronal and glial regulation of CNS angiogenesis and barriergenesis. *Development* **147**, doi:10.1242/dev.182279 (2020).

Umino, Y. & Solessio, E. Loss of scotopic contrast sensitivity in the optomotor response of diabetic mice. *Invest Ophthalmol Vis Sci* **54**, 1536-1543, doi:10.1167/iovs.12-10825 (2013).

Pardue, M. T. *et al.* Rodent Hyperglycemia-Induced Inner Retinal Deficits are Mirrored in Human Diabetes. *Transl Vis Sci Technol* **3**, 6, doi:10.1167/tvst.3.3.6 (2014).

Abcouwer, S. F. & Gardner, T. W. Diabetic retinopathy: loss of neuroretinal adaptation to the diabetic metabolic environment. *Ann N Y Acad Sci* **1311**, 174-190, doi:10.1111/nyas.12412 (2014).

Feit-Leichman, R. A. *et al.* Vascular damage in a mouse model of diabetic retinopathy: relation to neuronal and glial changes. *Invest Ophthalmol Vis Sci* **46**, 4281-4287, doi:10.1167/iovs.04-1361 (2005).

Feller, M. B., Wellis, D. P., Stellwagen, D., Werblin, F. S. & Shatz, C. J. Requirement for cholinergic synaptic transmission in the propagation of spontaneous retinal waves. *Science* **272**, 1182-1187, doi:10.1126/science.272.5265.1182 (1996).

Zhou, Z. J. Direct participation of starburst amacrine cells in spontaneous rhythmic activities in the developing mammalian retina. *J Neurosci* **18**, 4155-4165, doi:10.1523/JNEUROSCI.18-11-04155.1998 (1998).

Bansal, A. *et al.* Mice lacking specific nicotinic acetylcholine receptor subunits exhibit dramatically altered spontaneous activity patterns and reveal a limited role for retinal waves in forming ON and OFF circuits in the inner retina. *J Neurosci* **20**, 7672-7681, doi:10.1523/JNEUROSCI.20-20-07672.2000 (2000).

Blankenship, A. G. *et al.* Synaptic and extrasynaptic factors governing glutamatergic retinal waves. *Neuron* **62**, 230-241, doi:10.1016/j.neuron.2009.03.015 (2009).

Tiriac, A., Smith, B. E. & Feller, M. B. Light Prior to Eye Opening Promotes Retinal Waves and Eye-Specific Segregation. *Neuron* **100**, 1059-1065 e1054, doi:10.1016/j.neuron.2018.10.011 (2018).

Weiner, G. A. *et al.* Cholinergic neural activity directs retinal layer-specific angiogenesis and blood retinal barrier formation. *Nat Commun* **10**, 2477, doi:10.1038/s41467-019-10219-8 (2019).

Bonezzi, P. J., Stabio, M. E. & Renna, J. M. The Development of Mid-Wavelength Photoresponsivity in the Mouse Retina. *Curr Eye Res* **43**, 666-673, doi:10.1080/02713683.2018.1433859 (2018).

Stahl, A. *et al.* The mouse retina as an angiogenesis model. *Invest Ophthalmol Vis Sci* **51**, 2813-2826, doi:10.1167/iovs.10-5176 (2010).

Rao, S. *et al.* A direct and melanopsin-dependent fetal light response regulates mouse eye development. *Nature* **494**, 243-246, doi:10.1038/nature11823 (2013).

Nguyen, M. T. *et al.* An opsin 5-dopamine pathway mediates light-dependent vascular development in the eye. *Nat Cell Biol* **21**, 420-429, doi:10.1038/s41556-019-0301-x (2019).

Luhmann, U. F. *et al.* Role of the Norrie disease pseudoglioma gene in sprouting angiogenesis during development of the retinal vasculature. *Invest Ophthalmol Vis Sci* **46**, 3372-3382, doi:10.1167/iovs.05-0174 (2005).

20 Ye, X. *et al.* Norrin, frizzled-4, and Lrp5 signaling in endothelial cells controls a genetic program
for retinal vascularization. *Cell* **139**, 285-298, doi:10.1016/j.cell.2009.07.047 (2009).
21 Junge, H. J. *et al.* TSPAN12 regulates retinal vascular development by promoting Norrin- but not
Wnt-induced FZD4/beta-catenin signaling. *Cell* **139**, 299-311, doi:10.1016/j.cell.2009.07.048
(2009).
22 Xia, C. H., Yablonka-Reuveni, Z. & Gong, X. LRP5 is required for vascular development in deeper
layers of the retina. *PLoS One* **5**, e11676, doi:10.1371/journal.pone.0011676 (2010).
23 Zuercher, J., Fritzsche, M., Feil, S., Mohn, L. & Berger, W. Norrin stimulates cell proliferation in
the superficial retinal vascular plexus and is pivotal for the recruitment of mural cells. *Hum Mol*
Genet **21**, 2619-2630, doi:10.1093/hmg/dds087 (2012).
24 Paes, K. T. *et al.* Frizzled 4 is required for retinal angiogenesis and maintenance of the blood-retina
barrier. *Invest Ophthalmol Vis Sci* **52**, 6452-6461, doi:10.1167/iovs.10-7146 (2011).
25 Wang, Y. *et al.* Norrin/Frizzled4 signaling in retinal vascular development and blood brain barrier
plasticity. *Cell* **151**, 1332-1344, doi:10.1016/j.cell.2012.10.042 (2012).
26 Calvert, P. D. *et al.* Phototransduction in transgenic mice after targeted deletion of the rod transducin
alpha -subunit. *Proc Natl Acad Sci U S A* **97**, 13913-13918, doi:10.1073/pnas.250478897 (2000).
27 Schroeder, M. M. *et al.* The Roles of Rods, Cones, and Melanopsin in Photoresponses of M4
Intrinsically Photosensitive Retinal Ganglion Cells (ipRGCs) and Optokinetic Visual Behavior.
Front Cell Neurosci **12**, 203, doi:10.3389/fncel.2018.00203 (2018).
28 Okawa, H., Sampath, A. P., Laughlin, S. B. & Fain, G. L. ATP consumption by mammalian rod
photoreceptors in darkness and in light. *Curr Biol* **18**, 1917-1921, doi:10.1016/j.cub.2008.10.029
(2008).
29 Sherry, D. M., Wang, M. M., Bates, J. & Frishman, L. J. Expression of vesicular glutamate
transporter 1 in the mouse retina reveals temporal ordering in development of rod vs. cone and ON
vs. OFF circuits. *J Comp Neurol* **465**, 480-498, doi:10.1002/cne.10838 (2003).
30 Zarkada, G. *et al.* Specialized endothelial tip cells guide neuroretina vascularization and blood-
retina-barrier formation. *Dev Cell* **56**, 2237-2251 e2236, doi:10.1016/j.devcel.2021.06.021 (2021).
31 Siemerink, M. J., Klaassen, I., Van Noorden, C. J. & Schlingemann, R. O. Endothelial tip cells in
ocular angiogenesis: potential target for anti-angiogenesis therapy. *J Histochem Cytochem* **61**, 101-
115, doi:10.1369/0022155412467635 (2013).
32 Hellstrom, M. *et al.* Dll4 signalling through Notch1 regulates formation of tip cells during
angiogenesis. *Nature* **445**, 776-780, doi:10.1038/nature05571 (2007).
33 Suchting, S. *et al.* The Notch ligand Delta-like 4 negatively regulates endothelial tip cell formation
and vessel branching. *Proc Natl Acad Sci U S A* **104**, 3225-3230, doi:10.1073/pnas.0611177104
(2007).
34 Pelton, J. C., Wright, C. E., Leitges, M. & Bautch, V. L. Multiple endothelial cells constitute the tip
of developing blood vessels and polarize to promote lumen formation. *Development* **141**, 4121-
4126, doi:10.1242/dev.110296 (2014).
35 Chow, B. W. & Gu, C. Gradual Suppression of Transcytosis Governs Functional Blood-Retinal
Barrier Formation. *Neuron* **93**, 1325-1333 e1323, doi:10.1016/j.neuron.2017.02.043 (2017).
36 Mazzoni, J. *et al.* The Wnt Inhibitor Apcdd1 Coordinates Vascular Remodeling and Barrier
Maturation of Retinal Blood Vessels. *Neuron* **96**, 1055-1069 e1056,
doi:10.1016/j.neuron.2017.10.025 (2017).
37 Biswas, S. *et al.* Mural Wnt/beta-catenin signaling regulates Lama2 expression to promote
neurovascular unit maturation. *Development* **149**, doi:10.1242/dev.200610 (2022).
38 Bosma, E. K., van Noorden, C. J. F., Schlingemann, R. O. & Klaassen, I. The role of plasmalemma
vesicle-associated protein in pathological breakdown of blood-brain and blood-retinal barriers:
potential novel therapeutic target for cerebral edema and diabetic macular edema. *Fluids Barriers*
CNS **15**, 24, doi:10.1186/s12987-018-0109-2 (2018).

39 Gu, X. *et al.* Spatial and temporal localization of caveolin-1 protein in the developing retina. *Adv*
Exp Med Biol **801**, 15-21, doi:10.1007/978-1-4614-3209-8_3 (2014).
40 Gu, X. *et al.* Loss of caveolin-1 causes blood-retinal barrier breakdown, venous enlargement, and
mural cell alteration. *Am J Pathol* **184**, 541-555, doi:10.1016/j.ajpath.2013.10.022 (2014).
41 Buckingham, B. P. *et al.* Progressive ganglion cell degeneration precedes neuronal loss in a mouse
model of glaucoma. *J Neurosci* **28**, 2735-2744, doi:10.1523/JNEUROSCI.4443-07.2008 (2008).
42 Hajdu, R. I. *et al.* Detailed Evaluation of Possible Ganglion Cell Loss in the Retina of Zucker
Diabetic Fatty (ZDF) Rats. *Sci Rep* **9**, 10463, doi:10.1038/s41598-019-46879-1 (2019).
43 Usui, Y. *et al.* Neurovascular crosstalk between interneurons and capillaries is required for vision.
J Clin Invest **125**, 2335-2346, doi:10.1172/JCI80297 (2015).
44 Poche, R. A., Furuta, Y., Chaboissier, M. C., Schedl, A. & Behringer, R. R. Sox9 is expressed in
mouse multipotent retinal progenitor cells and functions in Muller glial cell development. *J Comp*
Neurol **510**, 237-250, doi:10.1002/cne.21746 (2008).
45 Serjanov, D., Bachay, G., Hunter, D. D. & Brunken, W. J. Laminin beta2 Chain Regulates Retinal
Progenitor Cell Mitotic Spindle Orientation via Dystroglycan. *J Neurosci* **38**, 5996-6010,
doi:10.1523/JNEUROSCI.0551-18.2018 (2018).
46 Hippert, C. *et al.* Muller glia activation in response to inherited retinal degeneration is highly varied
and disease-specific. *PLoS One* **10**, e0120415, doi:10.1371/journal.pone.0120415 (2015).
47 Lee, S. H., Park, Y. S., Paik, S. S. & Kim, I. B. Differential Response of Muller Cells and Microglia
in a Mouse Retinal Detachment Model and Its Implications in Detached and Non-Detached Regions.
Cells **10**, doi:10.3390/cells10081972 (2021).
48 Macosko, E. Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells Using
Nanoliter Droplets. *Cell* **161**, 1202-1214, doi:10.1016/j.cell.2015.05.002 (2015).
49 Taylor, S., Srinivasan, B., Wordinger, R. J. & Roque, R. S. Glutamate stimulates neurotrophin
expression in cultured Muller cells. *Brain Res Mol Brain Res* **111**, 189-197, doi:10.1016/s0169-
328x(03)00030-5 (2003).
50 Shen, F. *et al.* Glutamate-induced glutamine synthetase expression in retinal Muller cells after short-
term ocular hypertension in the rat. *Invest Ophthalmol Vis Sci* **45**, 3107-3112, doi:10.1167/iovs.03-
0948 (2004).
51 Harada, T. *et al.* Functions of the two glutamate transporters GLAST and GLT-1 in the retina. *Proc*
Natl Acad Sci U S A **95**, 4663-4666, doi:10.1073/pnas.95.8.4663 (1998).
52 Russo, R. *et al.* Impairment of neuronal glutamate uptake and modulation of the glutamate
transporter GLT-1 induced by retinal ischemia. *PLoS One* **8**, e69250,
doi:10.1371/journal.pone.0069250 (2013).
53 Cheng, S. Y. *et al.* Altered photoreceptor metabolism in mouse causes late stage age-related macular
degeneration-like pathologies. *Proc Natl Acad Sci U S A* **117**, 13094-13104,
doi:10.1073/pnas.2000339117 (2020).
54 Mackenzie, F. & Ruhrberg, C. Diverse roles for VEGF-A in the nervous system. *Development* **139**,
1371-1380, doi:10.1242/dev.072348 (2012).
55 Okabe, K. *et al.* Neurons limit angiogenesis by titrating VEGF in retina. *Cell* **159**, 584-596,
doi:10.1016/j.cell.2014.09.025 (2014).
56 Schweighofer, B. *et al.* The VEGF-induced transcriptional response comprises gene clusters at the
crossroad of angiogenesis and inflammation. *Thromb Haemost* **102**, 544-554, doi:10.1160/TH08-
12-0830 (2009).
57 Liu, Y., Cox, S. R., Morita, T. & Kourembanas, S. Hypoxia regulates vascular endothelial growth
factor gene expression in endothelial cells. Identification of a 5' enhancer. *Circ Res* **77**, 638-643,
doi:10.1161/01.res.77.3.638 (1995).
58 Lee, M., Leskova, W., Eshaq, R. S. & Harris, N. R. Retinal hypoxia and angiogenesis with
methamphetamine. *Exp Eye Res* **206**, 108540, doi:10.1016/j.exer.2021.108540 (2021).

Morita, A. *et al.* Transient phenotypic changes in endothelial cells and pericytes in neonatal mouse retina following short-term blockade of vascular endothelial growth factor receptors. *Dev Dyn* **247**, 699-711, doi:10.1002/dvdy.24614 (2018).

Hupe, M. *et al.* Gene expression profiles of brain endothelial cells during embryonic development at bulk and single-cell levels. *Sci Signal* **10**, doi:10.1126/scisignal.aag2476 (2017).

Jiang, E., Yan, X. & Weng, H. R. Glial glutamate transporter and glutamine synthetase regulate GABAergic synaptic strength in the spinal dorsal horn. *J Neurochem* **121**, 526-536, doi:10.1111/j.1471-4159.2012.07694.x (2012).

Sullivan, S. J. & Miller, R. F. AMPA receptor mediated D-serine release from retinal glial cells. *J Neurochem* **115**, 1681-1689, doi:10.1111/j.1471-4159.2010.07077.x (2010).

Shimamoto, K. *et al.* Characterization of novel L-threo-beta-benzyloxyaspartate derivatives, potent blockers of the glutamate transporters. *Mol Pharmacol* **65**, 1008-1015, doi:10.1124/mol.65.4.1008 (2004).

Corada, M. *et al.* Fine-Tuning of Sox17 and Canonical Wnt Coordinates the Permeability Properties of the Blood-Brain Barrier. *Circ Res* **124**, 511-525, doi:10.1161/CIRCRESAHA.118.313316 (2019).

Carmeliet, P. & Tessier-Lavigne, M. Common mechanisms of nerve and blood vessel wiring. *Nature* **436**, 193-200, doi:10.1038/nature03875 (2005).

Sapieha, P. Eyeing central neurons in vascular growth and reparative angiogenesis. *Blood* **120**, 2182-2194, doi:10.1182/blood-2012-04-396846 (2012).

Joyal, J. S. *et al.* Retinal lipid and glucose metabolism dictates angiogenesis through the lipid sensor Ffar1. *Nat Med* **22**, 439-445, doi:10.1038/nm.4059 (2016).

Sapieha, P. *et al.* The succinate receptor GPR91 in neurons has a major role in retinal angiogenesis. *Nat Med* **14**, 1067-1076, doi:10.1038/nm.1873 (2008).

Liang, J. H. *et al.* Dopamine signaling from ganglion cells directs layer-specific angiogenesis in the retina. *Curr Biol* **33**, 3821-3834 e3825, doi:10.1016/j.cub.2023.07.040 (2023).

Boccuni, I. & Fairless, R. Retinal Glutamate Neurotransmission: From Physiology to Pathophysiological Mechanisms of Retinal Ganglion Cell Degeneration. *Life (Basel)* **12**, doi:10.3390/life12050638 (2022).

Fremeau, R. T., Jr. *et al.* Vesicular glutamate transporters 1 and 2 target to functionally distinct synaptic release sites. *Science* **304**, 1815-1819, doi:10.1126/science.1097468 (2004).

Boye, K. *et al.* Endothelial Unc5B controls blood-brain barrier integrity. *Nat Commun* **13**, 1169, doi:10.1038/s41467-022-28785-9 (2022).

Jones, B. W. *et al.* Retinal remodeling in human retinitis pigmentosa. *Exp Eye Res* **150**, 149-165, doi:10.1016/j.exer.2016.03.018 (2016).

Jones, B. W. *et al.* Retinal Remodeling and Metabolic Alterations in Human AMD. *Front Cell Neurosci* **10**, 103, doi:10.3389/fncel.2016.00103 (2016).

Pfeiffer, R. L., Marc, R. E., Kondo, M., Terasaki, H. & Jones, B. W. Muller cell metabolic chaos during retinal degeneration. *Exp Eye Res* **150**, 62-70, doi:10.1016/j.exer.2016.04.022 (2016).

Chan-Ling, T. & Stone, J. Degeneration of astrocytes in feline retinopathy of prematurity causes failure of the blood-retinal barrier. *Invest Ophthalmol Vis Sci* **33**, 2148-2159 (1992).

Prokofyeva, E. & Zrenner, E. Epidemiology of major eye diseases leading to blindness in Europe: a literature review. *Ophthalmic Res* **47**, 171-188, doi:10.1159/000329603 (2012).

Liegl, R., Hellstrom, A. & Smith, L. E. Retinopathy of prematurity: the need for prevention. *Eye Brain* **8**, 91-102, doi:10.2147/EB.S99038 (2016).

Lacoste, B. *et al.* Sensory-related neural activity regulates the structure of vascular networks in the cerebral cortex. *Neuron* **83**, 1117-1130, doi:10.1016/j.neuron.2014.07.034 (2014).

Vanlandewijck, M. *et al.* A molecular atlas of cell types and zonation in the brain vasculature. *Nature* **554**, 475-480, doi:10.1038/nature25739 (2018).

81   Yuan, J. & Sims, P. A. An Automated Microwell Platform for Large-Scale Single Cell RNA-Seq.
Sci Rep **6**, 33883, doi:10.1038/srep33883 (2016).
82   Yuan, J. *et al.* Single-cell transcriptome analysis of lineage diversity in high-grade glioma. *Genome*
Med **10**, 57, doi:10.1186/s13073-018-0567-9 (2018).
83   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 e3529,
doi:10.1016/j.cell.2021.04.048 (2021).
84   Menon, V. Clustering single cells: a review of approaches on high-and low-depth single-cell RNA-
seq data. *Brief Funct Genomics* **17**, 240-245, doi:10.1093/bfgp/elx044 (2018).
991

FIGURES AND FIGURE LEGENDS

Figure 1. Deep plexus angiogenesis is delayed in the *Vglut1⁻ᐟ⁻* retina. A) Timeline of the neuronal activity

[cholinergic and glutamatergic waves and glutamatergic synaptic activity of photoreceptors

(phototransduction)] and angiogenesis (superficial, deep and intermediate plexuses) in the postnatal wild-

type (WT) retina (P1-P22). **B)** Schematic diagram of glutamate release from bipolar cells and

photoreceptors in WT and *Vglut1*$^{-/-}$ retinas. *Vglut1*$^{-/-}$ photoreceptors and bipolar cells fail to release

glutamate into the synaptic cleft. **C)** Schematic diagram of distinct developmental regions in the retina.

Central retina is developmentally more mature than peripheral retina. **D)** Representative images of the

superficial plexus from P8 WT and *Vglut1*$^{-/-}$ retinal flat mounts stained for CD31 (green; EC marker). **E)**

Representative images of the retinas from P10 WT and *Vglut1*$^{-/-}$ retinal flat mounts stained for CD31.

Superficial plexus is color-coded in purple and deep plexus is color-coded in yellow. Dashed white line

shows the extent of deep plexus vascularization. **F, G)** Representative images of the deep plexus from P10

and P14 WT and *Vglut1*$^{-/-}$ retinal flat mounts visualized with CD31 staining (green). **H, I)** Dotted bar graphs

of the vascular coverage of the superficial plexus in P8 and P10 WT and *Vglut1*$^{-/-}$ central (**H**) and peripheral

(**I**) retina (n = 3 mice/genotype). **J-M)** Dotted bar graphs of the vascular coverage (**J, L**) and branch point

density (**K, M**) of the deep plexus in P10, P12 and P14 WT and *Vglut1*$^{-/-}$ central (**J, K**) and peripheral (**L,**

M) retina (n = 4 mice/genotype). Scale bars: E = 250 µm; D, F, G = 62 µm. Statistical analyses were done

with Students t-test: *p<0.05, **p<0.02, NS = not significant. Error bars: Mean ± S.E.M.

1012

Figure 2. Aberrant angiogenic transcriptomic signatures are present in *Vglut1⁻/⁻* ECs from single cell RNA-sequencing (scRNA-seq). A) A Uniform Manifold Approximation and Projection (UMAP) for

dimension reduction projection shows separation of P10 WT and *Vglut1*[-/-] retinal ECs based on their transcriptomic signatures. **B, C)** Gene ontology (GO) analyses of P10 WT and *Vglut1*[-/-] retinal ECs show top downregulated (**B**) and upregulated (**C**) pathways. **D, E)** Dot plot of expression levels for angiogenesis (**D**) and tip cell (**E**) genes in P10 WT and *Vglut1*[-/-] retinal ECs. The dot size indicates the percent of the population expressing each marker, the color scale indicates the average level of gene expression. Genes enriched in deep plexus tip cells (D-tip cells) compared to superficial plexus tip cells (S-tip cells) are shown in red. Genes enriched in deep plexus tip cells (D-tip cells) compared to all ECs are shown in purple. **F, G)** Violin plots of representative angiogenesis (**F**) and tip cells (**G**) genes in P10 WT and *Vglut1*[-/-] retinal ECs. **H)** Schematic of a vascular sprout consisting of leading tip cells and proliferative stalk cells. **I)** Representative images of the deep plexus from P12 WT and *Vglut1*[-/-] retinal flat-mounts stained for Ki67 (green) and CD31 (purple). **J)** Measurements of Ki67[+] (proliferative), or cleaved Caspase3[+] (apoptotic) ECs in P12 WT and *Vglut1*[-/-] retinal deep plexus (n = 4). **K)** Representative images of the deep plexus from P12 WT and *Vglut1*[-/-] retinal flat-mounts stained for CD31 (green). The dotted line shows the border of the vascular front in the *Vglut1*[-/-] retina. **L)** Representative images of the deep plexus from P12 WT and *Vglut1*[-/-] retinal flat-mounts stained for CD31 and Dll4. Arrowheads point to Dll4[+] tip cells in the *Vglut1*[-/-] retina. Scale bars = 41 μm. Students t-test: **<0.02, NS = not significant. Error bars: Mean ± S.E.M.

.036   **Figure 3. Deep plexus vessels in the *Vglut1^{-/-}* retina are poorly perfused. A)** Schematic of Podocalyxin

.037   localization on the luminal surface of ECs. BM: basement membrane. **B)** Representative images of the deep

.038   plexus of P10 WT and *Vglut1^{-/-}* retinal flat mounts labeled with CD31 (red) and Podocalyxin (grey). **C)**

.039   Representative images of the deep plexus of P14 WT and *Vglut1^{-/-}* retinal flat mounts labeled with CD31

040   (red) and Podocalyxin (grey) 5 minutes after intracardiac injection of Lectin (blue). Empty arrowheads

041   show CD31$^+$ vessels that are not perfused with Lectin. **D, E)** Dotted bar graphs of Podocalyxin coverage of

042   the deep plexus in P10 (**D**) and P14 (**E**) WT and *Vglut1$^{-/-}$* retinas (n = 3-4 mice/genotype). **F)** Dotted bar

043   graphs of Lectin perfusion of the deep plexus in P14 WT and *Vglut1$^{-/-}$* retinas (n=3 mice/genotype). Scale

044   bars = 50 μm. Students t-test: *<0.05, **<0.02, NS = not significant. Error bars: Mean ± S.E.M.

Figure 4. Paracellular BRB maturation is impaired in *Vglut1⁻/⁻* retina. A) Timeline of the paracellular

and transcellular BRB maturation in the postnatal WT retina (P1-P22). **B)** Schematic of biocytin-TMR

paracellular leakage through impaired tight junctions at the BRB; the leaked biocytin-TMR is taken up by

neurons. vBM = vascular basement membrane. **C)** Representative low magnification images of P10 WT

and *Vglut1*[-/-] retinal flat mounts 40 minutes after intravenous injection of biocytin-TMR, followed by PBS

perfusion. **D-F)** Representative high magnification images of the superficial **(D, E)** and deep **(F)** plexuses

of P10 and P14 WT and *Vglut1*[-/-] retinal flat mounts labeled with Lectin (green) 40 minutes after intravenous

injection of biocytin-TMR, followed by PBS perfusion. **G-I)** Measurements of the area **(G, H)** and

fluorescence intensity (**I**) of leaked biocytin-TMR in P10 and P14 WT and *Vglut1*[-/-] retinal flat mounts (n =

4). BRB formation is impaired in *Vglut1*[-/-] retina. **J)** Dot plot of expression levels for BRB genes in P10 WT

and *Vglut1*[-/-] retinal ECs. The dot size indicates the percent of the population expressing each marker; the

color scale indicates the average level of gene expression. **K)** Violin plots of representative BRB genes in

P10 WT and *Vglut1*[-/-] retinal ECs. **L)** Representative images of the superficial and deep plexuses of P14

WT and *Vglut1*[-/-] retinal flat mounts stained for Claudin-5 (Cldn5; purple) and Lectin (green). Empty

arrowheads point to Lectin[+] blood vessels devoid of Claudin-5 at tight junctions. **M)** Representative images

of the superficial and deep plexuses in P14 WT and *Vglut1*[-/-] retinal flat-mounts stained for ZO-1 (*Tjp1*,

purple) and Lectin (green). Empty arrowheads point at Lectin[+] blood vessels devoid of ZO-1 localization

to tight junctions. **N, O)** Measurement of Claudin-5 (**N**) and ZO-1 (**O**) coverage of blood vessels in P14

WT and *Vglut1*[-/-] retinas (n=4 mice / genotype). Scale bars: C = 150 μm; D, L, M = 41 μm. Students t-test:

*<0.05, **<0.02. Error bars: Mean ± S.E.M.

Figure 5. Paracellular BRB matures precociously in the *Chrnb2*-/- retina. A) Schematic diagram

illustrating that glutamatergic wave begins earlier (P8) than usual (P10) in the *Chrnb2*-/- retina where

cholinergic wave is abolished. **B)** Representative images of the superficial plexus of P7 WT and *Chrnb2$^{-/-}$*

retinal flat-mounts labeled with Lectin (green). **C)** Quantification of vascular coverage in the superficial

plexus of P7 WT and *Chrnb2$^{-/-}$* retinas. **D)** Representative images of superficial and deep plexuses of P7

WT and *Chrnb2$^{-/-}$* retinal flat-mounts labeled with Lectin (green) 40 minutes after intravenous biocytin-

TMR injection, followed by PBS perfusion. **E, F)** Measurements of the area (**E**) and fluorescence intensity

(**F**) of leaked biocytin-TMR in P7 WT and *Chrnb2$^{-/-}$* retinal flat-mounts (n = 4 mice / genotype). **G)**

Representative images of superficial and deep plexuses of P10 WT and *Chrnb2$^{-/-}$* retinal flat-mounts labeled

with CD31 (green). **H, I)** Quantification of vascular coverage in the superficial (**H**) and deep (**I**) plexuses

of P10 WT and *Chrnb2$^{-/-}$* retinas. **J)** Representative images of superficial and deep plexuses of P10 WT and

Chrnb2$^{-/-}$ retinal flat-mounts labeled with Lectin (green) 40 minutes after intravenous biocytin-TMR

injection, followed by PBS perfusion. **K, L)** Measurements of the area (**K**) and fluorescence intensity (**L**)

of leaked biocytin-TMR in P10 WT and *Chrnb2$^{-/-}$* retinal flat-mounts (n = 4 mice / genotype). **M)**

Representative images of the superficial and deep plexuses of P10 WT and *Chrnb2$^{-/-}$* retinal flat-mounts

stained for Claudin-5 (purple) and Lectin (green). Empty arrowheads point at Lectin^{+} blood vessels devoid

of Claudin-5 localization to tight junctions. **N)** Measurement of Claudin-5 coverage of blood vessels in P10

WT and *Chrnb2$^{-/-}$* retinas (n = 3 mice / genotype). Scale bars: B, G = 62 μm, D, J, M = 41 μm. Students t-

test: **<0.02. Error bars: Mean ± S.E.M.

Figure 6. Deep plexus angiogenesis and paracellular BRB maturation are accelerated in the *Gnat1*⁻/⁻ retina. A, B) Schematic diagrams of glutamate release from rod photoreceptors in WT (**A**) and *Gnat1*⁻/⁻ (**B**) retinas. *Gnat1*⁻/⁻ rod photoreceptors (depolarized) release excess glutamate into the synaptic cleft. **C)** Representative images of the superficial plexus in P10 WT and *Gnat1*⁻/⁻ retinal flat-mounts stained for Lectin (green). **D)** Measurements of vascular coverage of the superficial plexus in P10 WT and *Vglut1*⁻/⁻

103 central and peripheral retina (n = 3 mice / genotype). **E, H)** Representative images of the deep plexus in

104 P10 WT and *Gnat1*$^{-/-}$ retinal flat-mounts stained for Lectin (green). **F, G, I, J)** Measurements of vascular

105 coverage (**F, I**) and branch point density (**G, J**) of the deep plexus in P10 and P14 WT and *Gnat1*$^{-/-}$ central

106 (**F, G**) and peripheral (**I, J**) retina (n = 4 mice / genotype). **K)** Representative images of the deep plexus in

107 P10 WT and *Gnat1*$^{-/-}$ retinal flat-mounts stained for Ki67 (purple) and CD31 (green). **L)** Measurements of

108 Ki67$^+$ ECs in P10 WT and *Gnat1*$^{-/-}$ retinal deep plexus (n = 4 mice / genotype). **M)** Representative images

109 of superficial and deep plexuses of P10 WT and *Gnat1*$^{-/-}$ retinal flat-mounts labeled with Lectin (green) 40

110 minutes after intravenous injection of biocytin-TMR (purple), followed by PBS perfusion. **N, O)**

111 Measurements of the area (**N**) and fluorescence intensity (**O**) of leaked biocytin-TMR in P10 WT and *Gnat1*$^{-}$

112 $^{/-}$ retinal flat-mounts (n = 4 mice / genotype). **P, Q)** Representative images of the superficial (**P**) and deep

113 (**Q**) plexuses in P10 WT and *Gnat1*$^{-/-}$ retinal flat-mounts stained for Claudin-5 (purple) and Lectin (green).

114 P' and Q' are magnified images of the corresponding boxed areas. Empty arrowheads point at Lectin$^+$ blood

115 vessels devoid of Claudin-5 localization to tight junctions. **R)** Measurement of Claudin-5 coverage of blood

116 vessels in P10 WT and *Gnat1*$^{-/-}$ retinas (n = 4 mice / genotype). Scale bars: **C, E, H** = 62 μm and **K, M, P,**

117 **Q** = 41 μm. Students t-test: *<0.05, **<0.02, NS = not significant. Error bars: Mean ± S.E.M.

Figure 7. Expression of glutamine synthetase, glutamate transporters and Norrin are downregulated in *Vglut1*-/- retinas and upregulated in *Gnat1*-/- retinas. A, B) P14 WT, *Vglut1*-/- and *Gnat1*-/- retinal sections were stained for glutamine synthetase (GS) (**A**), and the mean fluorescence intensity (M.F.I) in the INL was quantified (**B**) for each genotype (n = 4 mice / genotype). **C, D)** P14 WT, *Vglut1*-/- and *Gnat1*-/- retinal sections were stained for GLAST (**C**), and the M.F.I in the INL was quantified (**D**) for each genotype (n = 4 mice / genotype). **E, F)** P14 WT, *Vglut1*-/- and *Gnat1*-/- retinal sections were stained for GLT-1 (**E**),

and the M.F.I in the INL was quantified (**F**) for each genotype (n=4 mice / genotype). **G)** Fluorescent *in*

situ hybridization of P10 WT, *Vglut1*[-/-] and *Gnat1*[-/-] retinal sections with an antisense probe against *Ndp*

(Norrin), followed by staining with DAPI. Norrin is expressed throughout the INL. **H)** Quantification of

the M.F.I of *Ndp* in the INL of P10 WT, *Vglut1*[-/-] and *Gnat1*[-/-] retinal sections (n = 4 mice / genotype). **I)**

Expression profiles of *Ndp, Slc1a3* (GLAST) and *Slc1a2* (GLT-1) in the P14 WT retina from an existing

scRNA-seq database [48]. *Ndp* mRNA is expressed predominantly by the horizontal cells and Müller glia in

the INL (small subsets of amacrine and bipolar cells also express *Ndp*). In the INL, *Slc1a3* is primarily

expressed by Müller glia and astrocytes, whereas *Slc1a2* is expressed in retinal neurons as well as Müller

glia. **J)** Schematic of *ex vivo* organotypic culture experiments. **K)** Fluorescent *in situ* hybridization of P10

WT retinal sections, cultured *ex vivo* for 12 hours with or without glutamate or TFB-TBOA in the media,

with an antisense probe against *Ndp* (*Norrin*) mRNA, followed by staining with DAPI. *Norrin* mRNA is

expressed throughout the INL. **L, M)** Quantification of the M.F.I of *Ndp* mRNA in the INL of P10 WT

retinal sections, cultured *ex vivo* for either 6 **(L)** or 12 **(M)** hours with or without glutamate or TFB-TBOA

in the media (n = 4 retinas / group). GCL: ganglion cell layer, INL: inner nuclear layer, ONL: outer nuclear

layer. Scale bars = 41 μm. One-way ANOVA: *<0.05, **<0.02, NS = not significant. Error bars: Mean ±

S.E.M.

Figure 8. Upregulation of Norrin/β-catenin signaling rescues vascular deficits of the *Vglut1-/-* retina.

A, B) P14 WT, *Vglut1-/-* and *Gnat1-/-* retinal sections were stained (**A**) for Lef1 (green), CD31 (purple) and

DAPI (blue), and the number of Lef1+ ECs was quantified (**B**) for each genotype (n = 4 mice / genotype).

C, D) MeBIO (control)- or BIO (activates Norrin/β-catenin signaling)-treated P10 WT and *Vglut1-/-* retinal

flat-mounts were stained (**C**) with Lectin (purple) and Lef1 (green), and the number of Lef1+ ECs / vessel

area (mm^2) for each treatment was quantified (**D**) for each treatment (n = 4 mice / group). BIO activates

Norrin/β-catenin signaling in ECs of *Vglut1-/-* retinas. **E-G)** MeBIO- or BIO-treated P10 WT and *Vglut1-/-*

retinas were stained with Lectin [low (grey; **E**) and high (green; **F**) magnifications], and vascular coverage

of the peripheral deep plexus was quantified (**G**) for each treatment (n = 4 mice / group). Red dotted lines

depict the extent of deep plexus growth. **H, I)** Representative images of flat-mount MeBIO- or BIO-treated

P10 WT and *Vglut1⁻ᐟ⁻* retinas 40 minutes after intravenous injection of biocytin-TMR [low (grey; **H**) and

high (purple; **I**) magnifications] and stained with Lectin (green). P10 *Vglut1⁻ᐟ⁻* retinas treated with MeBIO

have high biocytin-TMR leakage. P10 *Vglut1⁻ᐟ⁻* retinas treated with BIO have low biocytin-TMR leakage

similar to P10 WT retinas. **J, K)** Dotted bar graphs show quantification of biocytin-TMR fluorescence

intensity (**J**) and leakage area (**K**) (n = 5 mice / group). **L)** According to our working model, glutamate is

taken up into synaptic vesicles by vGluT1 (*Vglut1*), and is released into the synaptic cleft by neurons, either

during spontaneous glutamatergic waves (from bipolar cells) or light-dependent glutamatergic synaptic

activities of photoreceptors in the retina. The inhibition of glutamate release from rods after light exposure

is regulated by Transducin (*Gnat1*). The released glutamate signals to Müller glia and inner nuclear layer

(INL) neurons including horizontal, amacrine and bipolar cells. As a result, Müller glia and INL neurons

upregulate Norrin (*Ndp*) expression and likely its secretion; however, the relative contribution from Müller

glia and INL neurons on Norrin levels in this process is unclear. Glutamate-dependent secretion of Norrin

activates endothelial Norrin/β-catenin signaling, which in turn promotes deep plexus angiogenesis and the

paracellular BRB maturation in both the superficial and deep plexuses of the retina. GCL: ganglion cell

layer, INL: inner nuclear layer. Scale bars **A, D, G** = 41 μm, **C, F** = 200 μm. One-way ANOVA: ****<0.02**.

Error bars: Mean ± S.E.M.