Expression status transition of *NOTCH1* accompanies chromatin remodeling in human early

retinal progenitor cells

Yoshitoku Watabe[1,2], Sakurako Kobayashi[1,2], Takahiro Nakayama[1,2], Satoru Takahashi[2,3], Masaharu

Yoshihara[2,4,†]

[1]College of Medicine, School of Medicine and Health Sciences, University of Tsukuba, Tsukuba,

Japan

[2]Department of Anatomy and Embryology, Institute of Medicine, University of Tsukuba, Tsukuba,

Japan

[3]Transborder Medical Center, Institute of Medicine, University of Tsukuba, Tsukuba, Japan

[4]Department of Primary Care and Medical Education, Institute of Medicine, University of Tsukuba,

Tsukuba, Japan

[†]**Corresponding author:**

Masaharu Yoshihara

Laboratory Animal Resource Center

1-1-1 Tennodai, Tsukuba, 305-8575 Ibaraki, Japan

Tel.: 81-29-853-7516

Fax: 81-29-853-6965

Email: yoshihara.masahar.ly@alumni.tsukuba.ac.jp

Abstract

Objective: The regulation of receptor expression is crucial for fine-tuned signal transduction. Notch

signaling is a key signaling pathway involved in retinal development. Although the involvement of

this signaling pathway in the differentiation of retinal ganglion cells has been documented, less is

known about its involvement in earlier stages of retinal progenitor cell differentiation. To clarify the

timing of Notch receptor expression in undifferentiated retinal progenitor cells and elucidate the

possible involvement of chromatin remodeling in the regulation of Notch receptor expressions, we re-

analyzed publicly available human fetal retina single-cell RNA-seq and ATAC-seq data.

Results: On days 59, 74, and 78, we observed *NOTCH1* mRNA expression in early retinal progenitor

cells, which diminished at later stages of differentiation. Integration of single-cell RNA-seq and

ATAC-seq revealed that chromatin remodeling in part of the *NOTCH1* locus was accompanied by

transitions in its mRNA expression. Importantly, chromatin accessibility in the region upstream of

NOTCH1 depended on the differentiated cell types. These results suggest that chromatin remodeling

may be involved in the differential expression of *NOTCH1*, although another type of Notch mRNA

expression regulation may exist.

Keywords: differentiation, epigenetics, eye development, single-cell ATAC-seq, single-cell RNA-seq

Introduction

Signal transduction depends on the expression of receptors regulated at multiple levels. These

regulations include chromatin remodeling and DNA-binding proteins, such as transcription factors

and transcriptional repressors. Because fine-tuned signal transduction is necessary for development, it

is important to clarify the regulatory mechanisms of receptor expression.

Notch signaling in mammals is dependent on the binding of five canonical DSL ligands and four Notch

receptors [1]. It has been suggested that Notch loci are subject to chromatin remodeling under both

normal [2-7] and pathological [8-16] conditions. In terms of developmental biology, the retina is a

good model for investigating cellular differentiation involving Notch signaling [17]. The developed

retina is composed of multiple cell types, including retinal ganglion cells, bipolar cells, photoreceptors,

amacrine cells, horizontal cells, and Müllar glial cells, all of which originate from the retinal progenitor

cells (RPCs). RPC is characterized by the expression of *Lhx2*, *Pax6*, *Rax*, and *Vsx2* while there are

some additional marker genes for developing horizontal cells/retinal ganglion cells (*Onecut1/2*) [18],

developing amacrine cells (*Elavl* genes) [19], developing photoreceptors/amacrine cells/Müllar glial

cells (*Eef1a1*) [20], developing retinal ganglion cells (*Meis2*) [21], developing horizontal cells (*Lhx1*,

Ptf1a) [22], and glial cells (*Pax2*) [23]. Of the four Notch receptors in mammals, Notch1, has been

suggested to be involved in the maintenance of RPCs and differentiation into retinal ganglion cells

[24]. However, it is unclear when and how Notch receptor expression is switched on and off in RPCs

during early differentiation. Here, we re-analyzed a public multi-omics dataset of single-cell RNA-seq

and single-cell ATAC-seq from three human fetal retinas [25] to clarify the timing of Notch receptor

expression and examine the involvement of chromatin remodeling in this receptor expression switch.

Methods

A single-cell multi-omics dataset (GSE183684) [25] was downloaded from the Gene Expression

Omnibus (https://www.ncbi.nlm.nih.gov/geo/); data from days 59, 74, and 78 in the dataset were

selectively used because they contained many undifferentiated retinal progenitor cells. The data were

processed in Seurat version 5.1.0 [26] and Signac version 1.14.0 [27] pipelines in R version 4.4.1 on

the Ubuntu 22.04.4 LTS environment. Pseudotime analysis and integration of the single-cell RNA-seq

data and the single-cell ATAC-seq data was conducted by using the "FindTransferAnchors" function

of Signac and Monocle3 version 1.3.7 [28], respectively. The conditions used in the analysis are

provided in the GitHub repository.

Results

Re-analysis day 59 human fetal retina

First, early gestational stage samples were characterized (day 59). UMAP analysis of single-cell RNA-

seq identified 13 clusters (Figure 1A), which were further characterized by marker gene expression

(Figure 1B). This sample primarily contained RPCs with various differentiation statuses except for

PAX2-expressing glial cells. Early RPCs (RPC1-3 and *MKI67*-expressing Proliferating RPC) were

characterized by *LHX2*, *PAX6*, *RAX*, and *VSX2*. In addition, pseudotime analysis suggested that these

RPCs differentiated into either *ONECUT1/2*-expressing, *ELAVL2/4*-expressing, or *EEF1A1*-

expressing RPCs (Figure 1C). Next, we examined Notch mRNA expression and found that *NOTCH1-*

3 was expressed primarily in early RPCs, with *NOTCH1* and *NOTCH3* being the most prominent

genes (Figure 1D). Then, we re-analyzed single-cell ATAC-seq data for this day 59 sample, which was

mathematically integrated with its single-cell RNA-seq data by using the "FindTransferAnchors"

function of Signac (Figure 1E). In the early RPCs (RPC1/2 and *MKI67*-expressing Proliferating RPC),

we observed peaks between Chr9 136510000-136520000 of the *NOTCH1* gene, which diminished in

the other RPC clusters. These results imply that Notch expression transition during RPC differentiation

might be associated with the chromatin remodeling of *NOTCH1* (Figure 1F). However, we noted that

chromatin accessibility of the upstream region of the *NOTCH1* locus remained high in *ONECUT-*

expressing RPCs. In contrast to these populations, chromatin accessibility of the *NOTCH1* locus in

ELAVL2/4-expressing RPCs was very low, which was consistent with the low mRNA expression in

these RPC populations. In addition, the chromatin accessibility of the *NOTCH3* locus remained

unchanged during RPC differentiation, although its mRNA expression diminished (Figure 1G). Since

NOTCH2 and *NOTCH4* were less prominent, the feature plots of the marker genes and the dot plots

of the Notch genes from single-cell RNA-seq along with coverage plots of *NOTCH2* and *NOTCH4*

from single-cell ATAC-seq are available in "Additional_file_1."

Figure 1 *NOTCH1* mRNA expression decrease and concomitant chromatin remodeling in the

day 59 sample. (A) UMAP analysis of the single-cell RNA-seq data identified 13 clusters. (B) Dot

plot of marker genes. Note that early RPC clusters expressed *LHX2*, *PAX6*, *RAX*, and *VSX2*. (C)

Monocle3 pseudotime analysis for clarifying the differentiation status. (D) Feature plot of *NOTCH1-*

4. Note that *NOTCH1* and *NOTCH3* expression was prominent in the early RPC clusters. (E) UMAP

analysis of the single-cell ATAC-seq data which was integrated with the single-cell RNA-seq data. (F)

Coverage plot of the *NOTCH1* locus showing the chromatin accessibility. Note that the upstream of

the gene is on the right. (G) Coverage plot of the *NOTCH3* locus. Note that the upstream of the gene

is on the right.

Re-analysis day 74 human fetal retina

To examine whether the changes in Notch mRNA expression and chromatin accessibility of the Notch

loci in the day 59 sample were representative, we investigated another early gestational stage sample

(day 74). UMAP analysis identified 15 clusters (Figure 2A) that were further characterized by marker

gene expression (Figure 2B). This sample primarily contained RPCs with various differentiation

statuses. Early RPCs (RPC1-6) were characterized by *LHX2*, *PAX6*, *RAX*, and *VSX2*. In addition,

pseudotime analysis suggested that these RPCs differentiated into either *ELAVL2/4*-expressing,

ONECUT1/MEIS2-expressing, or *ONECUT2*-expressing RPCs (Figure 2C). *ONECUT1/MEIS2*-

expressing RPCs differentiated into RPCs that markedly expressed *VSX2*. Next, we examined Notch

mRNA expression and found that *NOTCH1-3* was expressed primarily in early RPCs, with *NOTCH1*

and *NOTCH3* being the most prominent genes, similar to the day 59 sample (Figure 2D). We then re-

analyzed the single-cell ATAC-seq data for this sample, which were integrated with the single-cell

RNA-seq data (Figure 2E). In the early RPCs (RPC1-3), we observed peaks between Chr9 136510000-

136520000 of the *NOTCH1* gene, which diminished in the other RPC clusters. These results confirmed

that Notch expression transition during RPC differentiation might be associated with the chromatin

remodeling of *NOTCH1* (Figure 2F). However, we noted that the chromatin accessibility of the

upstream region of the *NOTCH1* locus remained high in *ONECUT*-expressing RPCs. In contrast to

these populations, the chromatin accessibility of the *NOTCH1* locus in *ELAVL2/4*-expressing RPCs

was very low, which was consistent with the low mRNA expression in these RPC populations. In

addition, the chromatin accessibility of the *NOTCH3* locus remained unchanged during RPC

differentiation (Figure 2G). In summary, similar to the day 59 sample, in the day 74 sample, a

concomitant mRNA decrease and chromatin remodeling in *NOTCH1* was observed, while chromatin

accessibility in the upstream region of the *NOTCH1* locus remained high in *ONECUT*-expressing

RPCs. The feature plots of the marker genes and dot plots of the Notch genes from single-cell RNA-

seq along with coverage plots of *NOTCH2* and *NOTCH4* from single-cell ATAC-seq are available in

"Additional_file_2."

Figure 2 *NOTCH1* mRNA expression decrease and concomitant chromatin remodeling of the

day 74 sample. (A) UMAP analysis of the single-cell RNA-seq data identified 15 clusters. (B) Dot

plot of marker genes. Note that early RPC clusters expressed *LHX2*, *PAX6*, *RAX,* and *VSX2*. (C)

Monocle3 pseudotime analysis. (D) Feature plot of *NOTCH1-4*. Note that *NOTCH1* and *NOTCH3*

expression was prominent in the early RPC clusters. (E) UMAP analysis of the single-cell ATAC-seq

data which was integrated with the single-cell RNA-seq data. (F) Coverage plot of the *NOTCH1* locus.

(G) Coverage plot of the *NOTCH3* locus.

Re-analysis day 78 human fetal retina

To further confirm the developmental changes in Notch mRNA expression and chromatin accessibility

at the Notch loci, we investigated an additional early gestational stage sample (day 78). UMAP

analysis identified 17 clusters (Figure 3A) that were further characterized by marker gene expression

(Figure 3B). The sample primarily contained RPCs with various differentiation statuses. Early RPCs

(RPC1-3 and *MKI67*-expressing Proliferating RPC1-2 cells) were characterized by *LHX2*, *PAX6*, *RAX*,

and *VSX2*. In addition, pseudotime analysis suggested that these RPCs differentiated into

ONECUT1/2-expressing RPCs, which later differentiated into *PTF1A* and *LHX1*-expressing RPCs,

MEIS2-expressing RPCs, or *ELAVL2/4*-expressing RPCs (Figure 3C). Next, we examined Notch

mRNA expression and found that *NOTCH1-3* was expressed primarily in early RPCs, with *NOTCH1*

and *NOTCH3* being the most prominent genes (Figure 3D). We then re-analyzed the single-cell ATAC-

seq data for the day 78 sample, which was integrated with the single-cell RNA-seq data (Figure 3E).

In the early RPCs (RPC1-3), we observed peaks between Chr9 136510000-136520000 of the

NOTCH1 gene, which diminished in the other RPC clusters (Figure 3F). However, we noted that the

chromatin accessibility of the upstream region of the *NOTCH1* locus remained high in *ONECUT*-

expressing RPCs. In contrast to these populations, the chromatin accessibility of the *NOTCH1* locus

in *ELAVL2/4*-expressing RPCs was very low, which was consistent with the low mRNA expression in

these RPC populations. In addition, the chromatin accessibility of the *NOTCH3* locus remained

unchanged during RPC differentiation (Figure 3G). In summary, examination of all three independent

samples suggested that *NOTCH1* mRNA expression decreased, which was concomitant with

chromatin remodeling in Chr9 136510000-136520000 of the *NOTCH1* locus. The feature plots of the

marker genes and the dot plots of the *NOTCH* genes from single-cell RNA-seq along with coverage

plots of *NOTCH2* and *NOTCH4* from single-cell ATAC-seq are available in "Additional_file_3"

Figure 3 *NOTCH1* mRNA expression decrease and concomitant chromatin remodeling of the

day 78 sample. (A) UMAP analysis of the single-cell RNA-seq data identified 17 clusters. (B) Dot

plot of marker genes. Note that early RPC clusters expressed *LHX2*, *PAX6*, *RAX*, and *VSX2*. (C)

Monocle3 pseudotime analysis. (D) Feature plot of *NOTCH1-4*. Note that *NOTCH1* and *NOTCH3*

expression was prominent in early RPC clusters. (E) UMAP analysis of the single-cell ATAC-seq data

which was integrated with the single-cell RNA-seq data. (F) Coverage plot of the *NOTCH1* locus. (G)

Coverage plot of the *NOTCH3* locus.

Discussion

The involvement of Notch signaling in cell fate choices is well documented, including in *Drosophila*

neurogenesis [29] and mammalian biliary development [30]. Although the regulation of Notch

receptor expression is necessary for these processes, to the best of our knowledge, few studies have

used genome-wide investigations of the underlying molecular mechanisms. To examine chromatin

remodeling in such regulatory mechanisms, we re-analyzed a single-cell RNA-seq and ATAC-seq

dataset from developing retinas in which differentiation trajectories were well characterized. By re-

analyzing three independent samples, we observed chromatin remodeling in part of the *NOTCH1* locus,

concomitant with changes in its mRNA expression during RPC differentiation.

Transcriptional regulation occurs at many levels, including DNA-binding proteins and miRNAs, as

well as chromatin remodeling. Importantly, the chromatin accessibility of the upstream regions of the

NOTCH1 locus was unaffected in *ONECUT*-expressing RPCs. Therefore, the observed mRNA

changes might be driven by pathways other than chromatin remodeling. Indeed, *NOTCH3* mRNA

expression also diminished during differentiation, although we observed no chromatin remodeling in

the *NOTCH3* locus. Because chromatin accessibility in *ELAVL2/4*-expressing RPCs was very low, this

ensured low mRNA expression of *NOTCH1*.

An ophthalmological study revealed that the epigenetic landscape of cell type-specific enhancers

shifted during differentiation of RPCs [31]. For example, in the single-cell ATAC-seq data from

embryonic day 14.5 mouse retina, motif enrichment for *Lhx2*, *Rax* and *Pax6* in the early RPCs were

observed, and footprinting analysis validated binding of those transcription factors to their motifs.

These high chromatin accessibilities decreased as they differentiated into retinal ganglion cells and

non-retinal ganglion cells. Although that study is excellent in providing comprehensive and in-depth

insights, ours is unique in focusing on the Notch loci for clarifying the regulatory mechanisms in view

of Notch signaling biology.

Limitations

This study elucidated the chromatin regions in which chromatin accessibility differed among RPC

subsets. We identified the transition of Notch receptor expression and the accompanying chromatin

remodeling of Notch loci in the early stages of RPC differentiation. Further investigations, such as a

large deletion of these regions, will be needed to evaluate the contribution of these regions to the

differential expression of Notch receptors in RPC subsets.

Abbreviations

RPC: Retinal progenitor cell

UMAP: Uniform manifold approximation and projection

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

A single-cell multiomics dataset (GSE183684) [18] was downloaded from the Gene Expression

Omnibus database (https://www.ncbi.nlm.nih.gov/geo/). The code used in the present study is

available   at   our   GitHub   repository

(https://github.com/Yoshitokky/Notch_multiomics_retina/tree/v1.0) that was archived using Zenodo

(https://doi.org/10.5281/zenodo.14058572).

Competing interests

The authors declare that they have no competing interests.

Funding

This study was supported by the JSPS KAKENHI Grant-in-Aid for Early-Career Scientists (grant no.

JP23K14429).

Authors' contributions

Y. W. performed the analyses and wrote the original manuscript. S.K. assisted in writing the manuscript.

T.N. designed the study. S.T. revised and supervised the study. M.Y. designed the study, performed

analyses, wrote the original manuscript, and acquired funding.

Acknowledgements

The authors thank Editage (https://www.editage.jp) for their support with English language editing.

References

1.   Kopan R, Ilagan MX. The canonical Notch signaling pathway: unfolding the activation

mechanism. Cell. 2009;137:216-33.

2.   He T, Fan Y, Du J, Yi M, Li Y, Liu M, et al. Stuxnet fine-tunes Notch dose during development

using a functional Polycomb response element. Development. 2023;150:dev201297.

3.   Schlereth K, Weichenhan D, Bauer T, Heumann T, Giannakouri E, Lipka D, et al. The

transcriptomic and epigenetic map of vascular quiescence in the continuous lung endothelium.

eLife. 2018;7:e34423.

4.   Lussi YC, Mariani L, Friis C, Peltonen J, Myers TR, Krag C, et al. Impaired removal of H3K4

methylation affects cell fate determination and gene transcription. Development. 2016;143:3751-

62.

5.   Marttila S, Kananen L, Jylhävä J, Nevalainen T, Hervonen A, Jylhä M, et al. Length of paternal

lifespan is manifested in the DNA methylome of their nonagenarian progeny. Oncotarget.

2015;6:30557-67.

6.   Gu B, Watanabe K, Sun P, Fallahi M, Dai X. Chromatin effector Pygo2 mediates Wnt-notch

crosstalk to suppress luminal/alveolar potential of mammary stem and basal cells. Cell Stem Cell.

2013;13:48-61.

7.   Rincon-Arano H, Halow J, Delrow JJ, Parkhurst SM, Groudine M. UpSET recruits HDAC

complexes and restricts chromatin accessibility and acetylation at promoter regions. Cell.

2012;151:1214-28.

8. Hussan SS, Ali MS, Fatima M, Altaf M, Sadaf S. Epigenetically dysregulated NOTCH-Delta-

HES signaling cascade can serve as a subtype classifier for acute lymphoblastic leukemia. Ann

Hematol. 2024;103:511-23.

9. Zhu Y, Ye M, Xu H, Gu R, Ma X, Chen M, et al. Methylation status of CpG sites in the NOTCH4

promoter region regulates NOTCH4 expression in patients with tetralogy of Fallot. Mol Med Rep.

2020;22:4412-22.

10. Jeong GY, Park MK, Choi HJ, An HW, Park YU, Choi HJ, et al. NSD3-induced methylation of

H3K36 activates NOTCH signaling to drive breast tumor initiation and metastatic progression.

Cancer Res. 2021;81:77-90.

11. Huang YC, Lin SJ, Shih HY, Chou CH, Chu HH, Chiu CC, et al. Epigenetic regulation of

NOTCH1 and NOTCH3 by KMT2A inhibits glioma proliferation. Oncotarget. 2017;8:63110-20.

12. Liu M, Liang K, Zhen J, Zhou M, Wang X, Wang Z, et al. Sirt6 deficiency exacerbates podocyte

injury and proteinuria through targeting Notch signaling. Nat Commun. 2017;8:413.

13. Liu S, Liu D, Chen C, Hamamura K, Moshaverinia A, Yang R, et al. MSC transplantation

improves osteopenia via epigenetic regulation of notch signaling in lupus. Cell Metab.

2015;22:606-18.

14. Kuang SQ, Fang Z, Zweidler-McKay PA, Yang H, Wei Y, Gonzalez-Cervantes EA, et al.

Epigenetic inactivation of Notch-Hes pathway in human B-cell acute lymphoblastic leukemia.

PLOS ONE. 2013;8:e61807.

15. Jithesh PV, Risk JM, Schache AG, Dhanda J, Lane B, Liloglou T, et al. The epigenetic landscape

of oral squamous cell carcinoma. Br J Cancer. 2013;108:370-9.

16. Jeannet R, Mastio J, Macias-Garcia A, Oravecz A, Ashworth T, Geimer Le Lay AS, et al.

Oncogenic activation of the Notch1 gene by deletion of its promoter in Ikaros-deficient T-ALL.

Blood. 2010;116:5443-54.

17. Mills EA, Goldman D. The regulation of notch signaling in retinal development and regeneration.

Curr Pathobiol Rep. 2017;5:323-31.

18. Sapkota D, Chintala H, Wu F, Fliesler SJ, Hu Z, Mu X. Onecut1 and Onecut2 redundantly

regulate early retinal cell fates during development. Proc Natl Acad Sci U S A. 2014;111:E4086-

95.

19. Wutikeli H, Yu Y, Zhang T, Cao J, Nawy S, Shen Y. Role of Elavl-like RNA-binding protein in

retinal development and signal transduction. Biochim Biophys Acta Mol Basis Dis.

2024;1871:167518.

20. Roesch K, Jadhav AP, Trimarchi JM, Stadler MB, Roska B, Sun BB, et al. The transcriptome of

retinal Müller glial cells. J Comp Neurol. 2008;509:225-38.

21. Rheaume BA, Jereen A, Bolisetty M, Sajid MS, Yang Y, Renna K, et al. Single cell transcriptome

profiling of retinal ganglion cells identifies cellular subtypes. Nat Commun. 2018;9:2759.

22. Boije H, Shirazi Fard S, Edqvist PH, Hallböök F. Horizontal cells, the odd ones out in the retina,

give insights into development and disease. Front Neuroanat. 2016;10:77.

23. Stanke J, Moose HE, El-Hodiri HM, Fischer AJ. Comparative study of Pax2 expression in glial

cells in the retina and optic nerve of birds and mammals. J Comp Neurol. 2010;518:2316-33.

24. Dvoriantchikova G, Perea-Martinez I, Pappas S, Barry AF, Danek D, Dvoriantchikova X, et al.

Molecular characterization of Notch1 positive progenitor cells in the developing retina. PLOS

ONE. 2015;10:e0131054.

25. Lyu P, Hoang T, Santiago CP, Thomas ED, Timms AE, Appel H, et al. Gene regulatory networks

controlling temporal patterning, neurogenesis, and cell-fate specification in mammalian retina.

Cell Rep. 2021;37:109994.

26. Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, et al. Dictionary learning

for integrative, multimodal and scalable single-cell analysis. Nat Biotechnol. 2024;42:293-304.

27. Stuart T, Srivastava A, Madad S, Lareau CA, Satija R. Single-cell chromatin state analysis with

Signac. Nat Methods. 2021;18:1333-41.

28. Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The dynamics and

regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. Nat

Biotechnol. 2014;32:381-6.

29. Heitzler P, Simpson P. The choice of cell fate in the epidermis of Drosophila. Cell. 1991;64:1083-

92.

30. Tanimizu N, Miyajima A. Notch signaling controls hepatoblast differentiation by altering the

expression of liver-enriched transcription factors. J Cell Sci. 2004;117:3165-74.

31. Ge Y, Chen X, Nan N, Bard J, Wu F, Yergeau D, et al. Key transcription factors influence the

epigenetic landscape to regulate retinal cell differentiation. Nucleic Acids Res. 2023;51:2151-

76.

Supplementary file

File name: Additional_file_1

File format: JPG

Title of data: Supplementary Figure for day 59 data

Description of data: (A) Feature plot of the marker gene mRNA expression. (B) Dot plot showing

Notch mRNA expression. (C) Chromatin accessibility at the *NOTCH2* locus as revealed by coverage

plots. (D) Coverage plot of the *NOTCH4* locus.

File name: Additional_file_2

File format: JPG

Title of data: Supplementary Figure for day 74 data

Description of data: (A) Feature plot of the marker gene mRNA expression. (B) Dot plot showing

Notch mRNA expression. (C) Chromatin accessibility at the *NOTCH2* locus as revealed by coverage

plots. (D) Coverage plot of the *NOTCH4* locus.

File name: Additional_file_3

File format: JPG

Title of data: Supplementary Figure for day 78 data

Description of data: (A) Feature plot of the marker gene mRNA expression. (B) Dot plot showing

Notch mRNA expression. (C) Chromatin accessibility at the *NOTCH2* locus as revealed by coverage

plots. (D) Coverage plot of the *NOTCH4* locus.

Figure legends

Figure 1 *NOTCH1* mRNA expression decrease and concomitant chromatin remodeling of the day 59 sample

(A) UMAP analysis of the single-cell RNA-seq data identified 13 clusters. (B) Dot plot of marker genes. Note that early RPC clusters expressed *LHX2*, *PAX6*, *RAX*, and *VSX2*. (C) Monocle3 pseudotime analysis for clarifying the differentiation status. (D) Feature plot of *NOTCH1-4*. Note that *NOTCH1* and *NOTCH3* expression was prominent in the early RPC clusters. (E) UMAP analysis of the single-cell ATAC-seq data, which was integrated with the single-cell RNA-seq data. (F) The coverage plot of the *NOTCH1* locus showing the chromatin accessibility. Note that the upstream of the gene is on the right. (G) The coverage plot of the *NOTCH3* locus. Note that the upstream of the gene is on the right.

Figure 2 *NOTCH1* mRNA expression decrease and concomitant chromatin remodeling of the day 74 sample

(A) UMAP analysis of the single-cell RNA-seq data identified 15 clusters. (B) Dot plot of marker genes. Note that early RPC clusters expressed *LHX2*, *PAX6*, *RAX*, and *VSX2*. (C) Monocle3 pseudotime analysis. (D) Feature plot of *NOTCH1-4*. Note that *NOTCH1* and *NOTCH3* expression

were prominent in the early RPC clusters. (E) UMAP analysis of the single-cell ATAC-seq data which

was integrated with the single-cell RNA-seq data. (F) The coverage plot of the *NOTCH1* locus. (G)

The coverage plot of the *NOTCH3* locus.

Figure 3 *NOTCH1* mRNA expression decrease and concomitant chromatin remodeling of the

day 78 sample

(A) UMAP analysis of the single-cell RNA-seq data identified 17 clusters. (B) Dot plot of marker

genes. Note that early RPC clusters expressed *LHX2*, *PAX6*, *RAX*, and *VSX2*. (C) Monocle3

pseudotime analysis. (D) Feature plot of *NOTCH1-4*. Note that *NOTCH1* and *NOTCH3* expression

were prominent in early RPC clusters. (E) UMAP analysis of the single-cell ATAC-seq data which

was integrated with the single-cell RNA-seq data. (F) The coverage plot of the *NOTCH1* locus. (G)

The coverage plot of the *NOTCH3* locus.