Direct measurements of luminal Ca^{2+} with endo-lysosomal GFP-aequorin reveal functional IP_3 receptors

Calvo B[1*], Torres-Vidal P[1*], Delrio-Lorenzo A[2], Rodriguez C[1], Aulestia FJ[3], Rojo-Ruiz J[1], McVeigh BM[4], Moiseenkova-Bell V[4], Yule DI[5], Garcia-Sancho J[1], Patel S[6**] and Alonso MT[1**]

[1]Unidad de Excelencia, Instituto de Biomedicina y Genética Molecular de Valladolid, Universidad de Valladolid y Consejo Superior de Investigaciones Científicas (CSIC), C/ Sanz y Forés 3, 47003 Valladolid, Spain

[2]Present address: Heart Failure and Inherited Cardiac Diseases Unit, Department of Cardiology, Hospital Universitario Puerta de Hierro, Madrid; Instituto de Salud Carlos III, Madrid; Centro Nacional de Investigaciones Cardiovasculares, CIBER Cardiovascular; C/ Melchor Fernandez Almagro 3, 28029 Madrid. Spain

[3]Present address: Mabxience, C/ Julia Morros 1, 24009, León, Spain

[4]Department of Systems Pharmacology and Translational Therapeutics, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, 19104 USA

[5]Department of Pharmacology and Physiology, University of Rochester, 601 Elmwood Avenue, Rochester, NY, 14642, USA.

[6]Department of Cell and Developmental Biology, University College London, Gower Street, London WC1E 6BT, UK.

* These authors contributed equally to this work

** Corresponding authors: talonso@uva.es and patel.s@ucl.ac.uk

25   **KEYWORDS:**

27   Endo-lysosome, Ca^{2+}, Ca^{2+} signalling, aequorin, genetically encoded Ca^{2+}

28   indicator, acidic organelle, IP_3R, Ca^{2+} release

ABSTRACT

Endo-lysosomes are considered acidic Ca^{2+} stores but direct measurements of luminal Ca^{2+} within them are limited. Here we report that the Ca^{2+}-sensitive luminescent protein aequorin does not reconstitute with its cofactor at highly acidic pH but that a significant fraction of the probe is functional within a mildly acidic compartment when targeted to the endo-lysosomal system. We leveraged this probe (ELGA) to report Ca^{2+} dynamics in this compartment. We show that Ca^{2+} uptake is ATP-dependent and sensitive to blockers of endoplasmic reticulum Ca^{2+} pumps. We find that the Ca^{2+} mobilizing messenger IP_3 which typically targets the endoplasmic reticulum evokes robust luminal responses in wild type cells, but not in IP_3 receptor knock-out cells. Responses were comparable to those evoked by activation of the endo-lysosomal ion channel TRPML1. Stimulation with IP_3-forming agonists also mobilized the store in intact cells. Super-resolution microscopy analysis confirmed the presence of IP_3 receptors within the endo-lysosomal system, both in live and fixed cells. Our data reveal a physiologically-relevant, IP_3-sensitive store of Ca^{2+} within the endo-lysosomal system.

INTRODUCTION

The endo-lysosomal (EL) system has emerged as a relevant player in subcellular Ca^{2+} signalling alongside the better characterised endoplasmic reticulum (ER) (Morgan et al., 2011; Patel and Docampo, 2010; Yang et al., 2019). EL Ca^{2+} has been implicated in most of the organelle's functions, such as late endosome-lysosome fusion, lysosomal exocytosis, and membrane repair. Impaired Ca^{2+} signalling can cause defects in EL trafficking and eventually can provoke lysosome storage diseases (LSDs). These so called acidic Ca^{2+} stores are thought to accumulate high amounts of Ca^{2+} (in the range of 0.4-0.6 mM), similar to the ER (Christensen et al., 2002; Lloyd-Evans et al., 2008). Akin to ER-localised receptor/Ca^{2+} channels for the second messenger IP_3 which mediate Ca^{2+} release in response to IP_3-forming hormones, various Ca^{2+}-permeable channels have been shown to localize to membranes of the EL system. These include Mucolipin transient receptor potential (TRPML) channels, two pore channels (TPCs) and P_2X_4 channels. TRPML1 is an ubiquitously expressed channel specifically activated by phosphatidylinositol 3,5-bisphosphate ($PI(3,5)P_2$), a low abundant phosphoinositide enriched in endomembranes, and which is mutated in type IV mucolipidosis (Dong et al., 2010).

The ER accumulates Ca^{2+} via SERCA. In contrast, the mechanism responsible for EL Ca^{2+} uptake has not been yet fully elucidated, mainly because of the technical difficulties associated to measure intraluminal Ca^{2+} in an acidic compartment. It is assumed that the vacuolar-type ATPase (V-ATPase) which is present in the EL membrane and maintains the proton gradient, drives Ca^{2+} into the lysosomal lumen via a type of transporter of H^+ for Ca^{2+} similar to the Ca^{2+}/H^+ exchanger present in yeast and in some animals (Melchionda et al., 2016; Mindell, 2012). This view has been recently challenged by the proposal that the ER, not the pH gradient, through IP_3R tunnels Ca^{2+} into the endo-lysosomes (Atakpa et al., 2019; Garrity et al., 2016). But these conclusions were drawn from indirect monitoring of cytosolic Ca^{2+} using conventional organic Ca^{2+} dyes or genetically-encoded Ca^{2+} indicators targeted to the cytosolic site of the EL system (Kilpatrick et al., 2016; Shen et al., 2012) thereby providing only qualitative insights into EL Ca^{2+} uptake.

Direct measurements of luminal Ca^{2+} within the EL system have been challenging due to the highly acidic pH which interferes with Ca^{2+} binding and thus requires

careful calibration. In this study, we report the use of the Ca^{2+}-sensitive

luminescent protein aequorin targeted to the endo-lysosome. In general, aequorin

is considered less sensitive to acidic pH than fluorescent genetically encoded Ca^{2+}

indicators (Johnson and Shimomura, 1978; Shimomura et al., 1974). But

surprisingly, we find that aequorin does not reconstitute well at extreme acidic pH

and therefore reports Ca^{2+} in a mildly acidic compartment of the EL system. We

show that this compartment accumulates Ca^{2+} via active uptake and releases it via

endogenous IP_3 receptors. Our data extend the actions of IP_3 to the endo-

lysosomal system.

RESULTS

ELGA as a novel endo-lysosomal Ca^{2+} indicator

We recently introduced genetically encoded Ca^{2+} indicators based on aequorin

fused to GFP (Rodriguez-Garcia et al., 2014; Rodriguez-Prados et al., 2015).

Here, we targeted them to the EL system by fusing the N-terminal of GFP-

aequorin (GA) to the C-terminal of the tetanus-insensitive vesicle-associated

membrane protein (VAMP7), a marker of the late endosomes and secretory

lysosome (Pryor et al., 2004; Vats and Galli, 2022) (***Figure 1A***). VAMP-7 is

composed of a cytosolic region, an EL transmembrane domain and a luminal C-

terminal domain. Therefore, the Ca^{2+} indicators are expected to be confined to the

EL lumen (***Figure 1B***). Due to the expected high endo-lysosomal luminal Ca^{2+}, we

introduced the D119A mutation in aequorin to lower the affinity for Ca^{2+}. This

probe was dubbed ELGA (for Endo-Lysosomal GFP-Aequorin). We generated a

second variant (ELGA1) with additional substitutions (D117A, and D163A) to

further reduce the affinity, lowering rate of aequorin consumption and thereby

allowing longer recordings (up to ~1h). Note that GA1 was a variant previously

published as GAP1 (Rodriguez-Garcia et al., 2014; Rodriguez-Prados et al., 2015)

but renamed here to maintain a comprehensive nomenclature. Both probes can

measure Ca^{2+} in the range of 100-1000 µM (***Figure 1 – figure supplement 1A***).

The Ca^{2+} sensors were expressed (either transiently or stably) in mammalian cells

using a relatively weak herpes simplex virus type 1 (HSV-1) promoter (Geller,

1991), to limit the amount of expressed protein and, hence, to minimize
mistargeting.

As shown in **Figure 1C- supplement movie 1**, ELGA and ELGA1 both localized
to punctate structures. To characterize the subcellular distribution of the ELGAs,
we performed colocalization analysis with various endogenous organellar markers.
Figure 1D shows representative confocal images of HEK293T cells transfected
with ELGA and fixed one day later. The endogenous lysosome-associated
membrane protein 2 (LAMP-2) was assayed by immunodetection with a specific
antibody (**Figure 1D**). There was colocalization of ELGA with LAMP2, whereas
there was no significant overlap with markers for other organelles such as ER
(Farrell et al., 2024), early endosome, or mitochondria (**Figure 1D- figure**
supplement 2 and table supplement 1). Altogether, these results indicate that
the majority of the ELGA indicator is localized to a subset of endo-lysosomes.

ELGA reports Ca^{2+} in mildly acidic compartments

To analyse the ability of aequorin to report Ca^{2+} at acidic pHs, we compared the
maximal luminescence signal triggered by Ca^{2+} in the pH range from 4.5 up to 7.2.
Recombinant GA regenerated with coelenterazine showed maximal luminescence
activity at pH 7.2 that decreased proportionally at lower pHs (6.5 and 6.0 pH) up to
a threshold of 5.5 pH, below which the Ca^{2+} sensor was not functional (**Figure 1E).**
Similar results were obtained with GA1 or with the original aequorin, either
reconstituted with native coelenterazine or with its lower affinity *n* analogue
(**Figure supplement 3**), indicating that the effect of acidic pH on the luminescent
function is independent on the affinity of aequorin for Ca^{2+}. This was surprising,
given previous calibration/performance studies at acidic pH (Mitchell et al., 2001;
Ronco et al., 2015; SantoDomingo et al., 2010). If the aequorin-based Ca^{2+} sensor
was first reconstituted at near-neutral pH (7.2) and then, shifted to pH 4.5 for the
luminescence assay, the luminescence activity was recovered, suggesting that the
lack of light emission is due to the binding of the cofactor to apoaequorin (**Figure**
1F).

The above findings precluded measurement of Ca^{2+} at highly acidic pH such as
that found in lysosomes. But because the pH of endocytic organelles is diverse
(e.g., early endosome, pH 6.5; late endosomes, pH 6.0; lysosome, pH 4.5), we

reasoned that ELGAs could be used to report Ca^{2+} in less acidic compartments within the EL system. To determine whether ELGAs reside in such compartments, we leveraged the pH sensitivity of the GFP moiety in ELGA. GFP fluorescence is quenched at acidic pH (pKa ~6). Thus, any detectable GFP fluorescence in live cells should reflect targeting to a compartment with a pH>5 within the range where aequorin can be reconstituted. As shown in **Figure 1G**, punctate GFP fluorescence was readily detectable in live HeLa and HEK293 cells, both transfected with ELGA.

To assess the properties of the probes at less acidic pH, we performed Ca^{2+} calibrations at pH 7.2 and 6. Essentially similar values were obtained at both pH values, indicating no major effect within this range. As shown in **Figure 1H-supplement 1A**, the affinity of GA1 for Ca^{2+} was ~350 µM at pH 7.2.

Taken together, these results identify ELGAs as reporters of Ca^{2+} within mildly acidic compartments.

Uptake of Ca^{2+} is ATP-dependent and sensitive to blockers of SERCA

In order to directly monitor uptake and release of Ca^{2+} within the EL compartment, we used digitonin-permeabilized HEK293T cells expressing ELGA1 perfused with a cytosol-like solution. As we expected the EL system to be a high Ca^{2+} content store, it needed to be depleted of Ca^{2+} before aequorin was regenerated with its cofactor to avoid its premature consumption. But the mechanisms of store filling are unclear. We first tested Bafilomycin A1 prior to adding the coelenterazine: this compound inhibits the V-type ATPase and has been used widely to disrupt the activity of putative $Ca^{2+}-H^+$ exchangers. Bafilomycin A1 treatment did not produce a measurable luminescent signal (data not shown), indicating that aequorin had been consumed likely because the target store was not emptied of Ca^{2+}. These data are consistent with functional targeting of ELGA to a non-highly acidic compartment.

By contrast, incubation with the reversible SERCA inhibitor tert-butylhydroquinone (tBHQ), produced a robust luminescence with a good signal to noise ratio (**Figure 2A**). Under these conditions perfusion with cytosol-like medium (free $[Ca^{2+}]$ = 200 nM) initiated Ca^{2+} uptake at a rate of 8 ± 0.6 µM/s (n=36), reaching a value of $[Ca^{2+}]_{EL}$ at the steady-state of 365 ± 21 µM (n=54) in 3 ± 0.6 min (n=7) (**Figure 2A**

and B). Other cell types tested, e.g., HeLa, CHO, MDCK or cortical astrocytes
displayed similar responses (*Figure 2B- figure supplement 4).* We tested the
effect of Bafilomycin A1. Bafilomycin A1 (1 µM) added acutely or following
incubation for 1 h had little effect on the $[Ca^{2+}]_{EL}$ (*Figure supplement 5*),
consistent with the failure to reconstitute the probe in its presence.

Ca^{2+} uptake was ATP-dependent because there was little change in luminescence
in the absence of ATP (90 ± 4% inhibition, n=7; *Figure 2C and F - figure*
supplement 1B-C). Similarly, the SERCA specific inhibitors thapsigargin or tBHQ
provoked a complete inhibition of the EL Ca^{2+} uptake (98 ± 1%, n=7; and 97 ± 2%,
n=8; *Figure 2D and E,* respectively). In this last case, washing out of the
reversible inhibitor rapidly restored the Ca^{2+} uptake up to normal rate (*Figure 2E*).
Once the endo-lysosomal lumen was fully replenished with Ca^{2+}, its removal
(addition of 0.1 mM EGTA) caused a progressive and complete decline in the
$[Ca^{2+}]_{EL}$ at a rate of ~2.0 µM/s (*Figure 2G)*. Likewise, removal of ATP or
application of tBHQ, either in the presence or absence of Ca^{2+}, emptied the EL-
Ca^{2+} store. Statistical quantifications of the EL-Ca^{2+}-uptake and Ca^{2+}-exit rate are
shown in *Figure 2F and H.*

Collectively, these results indicate that the endo-lysosome can accumulate high
amounts of Ca^{2+} through a SERCA mechanism.

IP_3 mobilizes Ca^{2+} from endo-lysosomal Ca^{2+} stores

To explore endogenous Ca^{2+} release pathways, we examined the effect of
NAADP, which activates endo-lysosomal TPCs. NAADP evoked a fast reversible
response that released 30% of the EL-Ca^{2+} content, both in HEK and HeLa cells.
But the responses were inconsistent and observable in around ~ 50% of trials
(*Figure supplement 6*). Surprisingly, addition of IP_3 (2 µM, 30s) reliably produced
a fast and strong Ca^{2+} release from the endo-lysosomal store that reduced the
endo-lysosomal luminal $[Ca^{2+}]_{EL}$ by 72 ± 1% (n= 18) (*Figure 3A and 3F*). The IP_3-
evoked Ca^{2+} release was very robust and consistent in all cell types tested (HeLa,
CHO or MDCK cells; representative results are shown in *figure supplement 7*).
This effect was concentration dependent, from 0.01 µM, that produced a small but
detectable effect, up to 10 µM, a maximal effect, comparable to that with 30 µM,
the highest IP_3 concentration tested (*Figure 3B,* and summary data in *Figure 3C*).

To test specificity of the IP$_3$ response, we used IP$_3$R-null HEK293 (HEK-3KO) cell

line generated by CRISPR/Cas9 technology lacking all three IP$_3$ receptor subtypes

(Alzayady et al., 2016). As shown in **Figure 3D**, IP$_3$-induced Ca^{2+} release was

abrogated (1.36 ± 0.97 % (n= 12). Interestingly, in these cells both the EL-Ca^{2+}

refilling rate (6.7 ± 0.7 µM/s) and the [Ca^{2+}]$_{EL}$ reached at the steady-state (369 ±

32 µM; n= 12) coincided with those values obtained in IP$_3$ receptor replete control

cells (8.0 ± 0.6 µM/s; n= 36 and 365 ± 21 µM; n= 54) (**figure supplement 8**). This

suggests that, unless there are compensatory effects, IP$_3$R is not essential for

refilling the EL Ca^{2+} store. We also examined the effect of IP$_3$ in HEK-3KO cells

upon re-expression of IP$_3$R1. As shown in **Figure 3E and F**, the IP$_3$ response was

partially rescued (54 ± 1 %; n= 7)**.**

In an independent approach, we used heparin, an inhibitor of the IP$_3$R. Heparin

completely blocked the IP$_3$ response (92 ± 2 % inhibition; n= 5; **Figure 3G and**

3H). Collectively, both genetic and pharmacological approaches reveal the

presence of functional IP$_3$Rs within the EL Ca^{2+} store that support robust luminal

Ca^{2+} responses.

ELGA responses are independent of the ER

Because of the similarities in the properties of the Ca^{2+} stores reported by ELGA

and the ER with respect to both Ca^{2+} uptake and Ca^{2+} release, we considered the

possibility that a fraction of the probe may be mistargeted to the ER.

To address this, we expressed TRPML1, a major endo-lysosomal Ca^{2+} channel.

Colocalization analysis of ELGA with TRPML in fixed cells is shown in **Figure 4A**.

Both probes were targeted to the same compartment. Colocalization significantly

decreased in live cells (**Figures 4A and B**) indicating that a proportion of ELGA is

located to acidic, non-fluorescent vesicles in live cells that become fluorescent

when the pH increases upon fixation. These results provide further evidence that

ELGA is a Ca^{2+} reporter in a fraction of the EL system.

Interestingly, TRPML1 expression drastically reduced the levels of Ca^{2+} reported

by ELGA1 (**Figure 4C**). In contrast, Ca^{2+} levels in the ER measured using the

same reporter targeted to the ER were unchanged upon TRPML1 expression,

indicating that the channel is specifically leaky in the EL store. These functional

data distinguish EL and ER Ca^{2+} stores. In accord, there was no correlation

between EL and ER Ca^{2+} levels reported by the corresponding GA1 in the cell
types used (**Figure 4D).**

To further explore the properties of TRPML1, we activated it with the specific
membrane-permeable synthetic agonist ML-SA1. In transiently expressing
TRPML1 HEK293T cells, ML-SA1 (20 µM, 30s) produced a robust EL-Ca^{2+}
release of 38 ± 4% (n= 15) of the total EL luminal Ca^{2+}content, as reported by
ELGA1 at the steady-state (**Figure 4E and J**). Upon application, ML-SA1 exhibited
a lag time of ~20 s and it continued releasing Ca^{2+} once the agonist was washed
out. Consecutive applications of ML-SA1 evoked responses of similar amplitude,
indicating no desensitization of the channel. The time to refill the store after the
washing out was ~5 min. Full concentration-effect relationship for the Ca^{2+} release
response showed an EC_{50} value of 7 ± 1 µM (**Figures 4F** and **G**). Comparable
responses were observed in TRPML1-expressing HeLa cells (**figure supplement**
9). In accord, the response to ML-SA1 was selectively blocked by the specific
TRPML-antagonists ML-SI1 **(Figure 4H -figure supplement 9B and 9C)** or ML-
SI3 (**figure supplement 9C**) and expression of TRPML1 did not affect the
response to IP_3 (**figure supplement 9D).**

We also activated TRPML1 with its endogenous activator PI(3,5)P_2. As shown in
Figure 4I, PI(3,5)P_2 evoked EL-Ca^{2+} release in cells expressing TRPML1 albeit
modestly compared to ML-SA1. Statistical quantification is summarized in **Figure**
4J. The EL-Ca^{2+} release provoked by PI(3,5)P_2 was concentration-dependent
(**Figure 4K and L).**

These data show that ELGA accurately reports the properties of a *bona fide* EL
Ca^{2+} channel.

A subset of IP_3 receptors localize to endo-lysosomes

We applied several high resolution microscopy techniques to assess whether IP_3
receptors localized to the EL system. We used HeLa cells in which endogenous
IP_3R1 is tagged with EGFP by gene editing (Thillaiappan et al., 2017). Results
from stimulated emission depletion (STED) super-resolution microscopy showed
that IP_3R1-GFP displayed a well resolved punctate distribution in fixed cells
(**Figure 5A**). This distribution was compared to expressed EL channels. We
detected a number of IP_3R1 puncta that colocalized with TRPML1 (**Figures 5Aa-**

c). Essentially, similar results were obtained with TPC2 (***Figures 5Ad-f).*** In

addition, we found part colocalization of IP_3R1 (***Figures 5Ag-i***) and SERCA2

(***figure supplement 10***) with endogenous LAMP2 although overlap was modest.

We also used STED microscopy to localize endogenous IP_3R3 in HEK cells. We

found evidence for part colocalization of IP_3R3 with endogenous LAMP1 (***Figure***

5B).

To seek evidence for IP_3R localization with EL system in live cells, we loaded

IP_3R1-GFP HeLa cells with fluorescent acidotropes (LysoView650 and Lysotracker

red). Some IP_3R1 clearly formed puncta with endo-lysosomes using either super-

resolution microscopy (STED; ***Figure 5C)*** or total internal reflection fluorescence

microscopy (TIRFM; ***Figure 5D).***

In sum, these data support the view that a subpopulation of IP_3Rs are physically

localized to the endo-lysosomal system.

Physiological IP_3-forming stimuli mobilize Ca^{2+} from endo-lysosomal Ca^{2+}

stores

To further characterize endo-lysosomal Ca^{2+} dynamics, we assessed luminal Ca^{2+}

levels in intact cells. Addition of 1 mM $CaCl_2$ refilled the EL Ca^{2+}store up to a

steady-state of 234 ± 29 µM (n= 12) (***Figure 6A***).

To probe the functionality of IP_3 receptors, we stimulated live cells with ATP which

activates endogenous P2Y receptors or carbachol which activates muscarinic

receptors. ATP and carbachol provoked fast and reversible EL Ca^{2+}responses of

35.81 ± 2.34 % (n= 7) and 19.88 ± 1.87 % (n= 5), respectively (***Figure 6A***)***.*** ML-

SA1 also produced reproducible responses in cells expressing TRPML1 although

it was ineffective in wild type cells (***Figure 6A and B***). Similar experiments were

performed in polarised MDCK cells ***(Figure 6C).*** This cell type displayed a

comparable steady-state EL-Ca^{2+} that was almost completely discharged upon cell

activation with ATP (73.72 ± 16.85 %; n= 4).

Finally, we examined EL responses in a primary cell type. In mouse cortical

astrocytes ***(Figure 6D)***, ML-SA1 application elicited a negligible effect whereas a

subsequent pulse of ATP almost completely discharged the EL-Ca^{2+} store (87.68 ±

5.90 %; n= 6). A summary of the data is shown in ***Figure 6E***.

In sum, these data identify a widespread physiologically-relevant IP$_3$-sensitive

Ca^{2+} store within the EL system.

DISCUSSION

Using a new genetically-encoded Ca^{2+} indicator based on aequorin, we demonstrate here the existence of a *bona-fide* endo-lysosomal Ca^{2+} store, able to accumulate a high $[Ca^{2+}]$ and to release it upon physiological activation (***Figure 6F***).

Our results are best explained if this endo-lysosome is functionally coupled with the ER to accumulate Ca^{2+} via active transport such as via SERCA pumps. Although the two compartments work in parallel, they are distinct in the amount of Ca^{2+} that they can store, with the luminal Ca^{2+} concentration being significantly lower in the EL system (~200-400 µM) than the ER (~400-500 µM) and in their sensitivity to expression of TRPML1, which selectively lowers the EL Ca^{2+} store content. The EL Ca^{2+} values are in agreement with previous reports (Narayanaswamy et al., 2019; Ronco et al., 2015) but contrast with others (Albrecht et al., 2015). The SERCA might be located at the EL membrane where it directly pumps Ca^{2+} into its lumen, or, alternatively, Ca^{2+} could be indirectly fuelled from the ER into the EL lumen, through an undetermined channel connecting the two organelles. The ER directly contacts the EL system through membrane contact sites (MCS), especially abundant in late endosomes, where they allow exchange of lipids and other molecules, including Ca^{2+} (Kilpatrick et al., 2013; Lee and Blackstone, 2020; Raiborg and Stenmark, 2015). Close association of ELGA-positive compartments with ER reported here might provide a route for uptake.

We show that activation of TRPML1 with ML-SA1/PI(3,5)P_2 and TPCs with NAADP produces luminal responses, but the latter were not consistent. Recent studies have shown that NAADP binds to soluble binding proteins to activate TPCs (Gunaratne et al., 2023; Saito et al., 2023). It is possible that loss of these proteins upon permeabilization may account for NAADP insensitivity. Surprisingly, IP$_3$ consistently produced robust Ca^{2+} release responses comparable to those evoked by bonafide EL channels. In accord with these functional data, our results with super resolution microscopy and TIRFM provide direct evidence for the presence of IP$_3$R on the endo-lysosome vesicles. Importantly, proteomic analysis of endo-lysosomal vesicles isolated from HEK293 cells by multiple different protocols, has also identified the two isoforms of IP$_3$R and SERCA in this organelle

(Singh et al., 2020). Moreover, IP$_3$Rs are detected in numerous additional cell

lines, both of mouse and human origin (Akter et al., 2023). Finally, we have also

confirmed the presence of IP$_3$Rs and SERCA in endo-lysosomes using LysoIP

(unpublished results). Functional IP$_3$ receptors have been documented in other

acidic organelles such as secretory granules in professional secretory cell types

(Gerasimenko et al., 2006; Santodomingo et al., 2008) and the acidocalcisome in

Trypanosomes (Huang et al., 2013). But we are not aware of their presence within

the EL system, which extends IP$_3$ action to potentially all cells.

Previous work suggested that IP$_3$ fills the EL system with Ca^{2+} (Atakpa et al.,

2018; Garrity et al., 2016) as opposed to emptying it as reported here. But direct

measurements of luminal Ca^{2+} in response to IP$_3$ were not reported previously.

Instead, these conclusions were based almost exclusively on indirect

measurements of Ca^{2+} in the cytosol. Additionally, because reconstitution of ELGA

is pH sensitive, it reports Ca^{2+} in only a sub-compartment of the EL system. The

EL system is a hybrid compartment, very heterogeneous in morphology,

composition, and function. Although this organelle is globally considered acidic, its

luminal pH varies from 4.5 up to 6.5 in different vesicles. The acidity of the vesicles

has been correlated with subcellular location (Bright et al., 2016; Johnson et al.,

2016) likely indicative of functional diversity. Thus, Ca^{2+} handling, might also show

heterogeneity among the various endo-lysosomal vesicles not least because of the

relationship between luminal Ca^{2+} and pH (Christensen et al., 2002). Probes that

allow spatial mapping of Ca^{2+} are required to pinpoint the presence of endosomal

IP$_3$ receptors that might release Ca^{2+} and ER IP$_3$ receptors that might drive Ca^{2+}

uptake.

Our intact cell studies provide evidence that IP$_3$-sensitive stores can be tapped

during physiological stimulation with IP$_3$-forming stimuli. The fast exit of Ca^{2+} from

the EL into the cytosol through IP$_3$Rs would generate local cytosolic Ca^{2+} signals.

These, in turn, could be amplified either via Ca^{2+}-induced Ca^{2+} release (CICR)

through IP$_3$Rs (or RyRs in other cell types) present in the ER membrane to

generate global signals. Indeed, IP$_3$Rs have been shown to be preferentially

associate with the lysosomes at contact sites (Atakpa et al., 2019). Such coupling

would be analogous to that triggering of Ca^{2+} signals by TRPML (Kilpatrick et al.,

2016) or TPCs (Xu and Ren, 2015; Yuan et al., 2024; Zhu et al., 2010). Our results

are in line with the '*chatter'* between the two organelles (Patel et al., 2010) and the

essential role of the ER in the EL signal in various cell types (Garrity et al., 2016;

López-Sanjurjo et al., 2013; Morgan et al., 2013; Yang et al., 2019). Alternatively,

local IP_3R-dependent Ca^{2+} signals may regulate endo-lysosomal function at basal

(unstimulated) levels of IP_3, again in analogy with TPCs and TRPML, which are

well established in regulating membrane traffic (Pryor et al., 2000; Ruas et al.,

2010; Vassileva et al., 2020).

In conclusion, our direct luminal Ca^{2+} measurements help to fill the gaps in our

knowledge on the Ca^{2+} signalling toolkit of the endo-lysosome system and provide

new insight into the dynamics of these small acidic Ca^{2+} stores and their functional

relationship with the ER Ca^{2+} stores.

 FIGURE LEGENDS

Figure 1. Design, characterization and calibration of endo-lysosomal Ca^{2+}

indicators. (A*)* Domain structure and mutations of the endo-lysosomal indicators

drawn approximately at scale. VAMP7, vesicle-associated membrane protein 7;

AEQ, aequorin; ELGA, endo-lysosomal GFP-Aequorin; ELGA1, endo-lysosomal

GAP1. **(B)** Predicted topology of ELGA and ELGA1 Ca^{2+} probes within the endo-

lysosome. **(C)** Representative confocal fluorescent images of equatorial plane of

fixed MDCK cells transiently transfected with ELGA or ELGA1. Scale bar for both

images, 10 µm. **(D)** *Upper row:* ELGA expressing HEK293T cells (*Left*, green)

were immunostained with a specific anti-LAMP2 antibody (*Center*, magenta).

Green and magenta images are merged (*Right, merge*). ***Lower row:***

Colocalization of ELGA expressing HEK293T cells with the following organellar

markers (magenta): ER, er-Ruby-GCaMP210 (ER); early endosome,

immunostained with the anti-early endosome antigen 1 (EE); mitochondria,

immunostained with anti-TOM20 antibody (mt). Scale bar for all images, 5 µm. **(E)**

Effect of acidic pH on GA luminescence emission. *E.coli* recombinant GA

apoprotein was reconstituted with native (*left*) or *n* (*right*) coelenterazine (CTZ) for

20 min at the indicated pH, and then subjected to luminescence emission assay

triggered with 10 mM CaCl$_2$ (arrow) and measured at the same pH. Buffers used

were: acetate for range 4.5-5.5 pH; Na-MOPS for 6 and 6.5 pH; and Na-Hepes for

7.2 pH. Each trace is the value normalized with respect to its background emission

before addition of Ca^{2+}. Each trace is the average of 3-4 independent experiments.

(F) Luminescence activity of GA1 after regeneration with *n* coelenterazine at 7.2 or

4.5 pH. Peak value (mean ± S.E.M; n=3) obtained from an experiment similar to

(E) performed at pHs 7.2 *(blue bar)* and 4.5 *(black bar)* or reconstituted with

coelenterazine at pH 7.2 and switched to pH 4.5 to measure luminescence (*pink*

bar). **(G)** Expression of ELGA in live HeLa and HEK293T cells. Scale bar, 10 µm.

(H) Ca^{2+} titration curves of GA1 at pH 6.0 and 7.2. Each value is the mean ± S.E.M

of 3-6 measurements obtained from two independent protein preparations.

Figure 2. The endo-lysosomal Ca^{2+} store accumulates Ca^{2+} via SERCA. (A)

Representative EL-Ca^{2+} signals recorded in HEK293T cells transiently transfected

with ELGA1. After Ca^{2+} depletion prior to reconstitution with coelenterazine *n*, cells

were permeabilized in Ca^{2+}-free intracellular medium (containing 1 mM ATP) and

then perfused with 200 nM free [Ca^{2+}], as indicated. **(B)** Summary data (mean ± S.E.M.) of [Ca^{2+}]$_{EL}$ at steady-state obtained in populations of HEK293T, HeLa and MDCK cells with the number of independent transfections in parenthesis. **(C)** Cells were perfused as in (A) with an intracellular medium without ATP (-ATP) and reintroduced (1 mM) when indicated. **(D)** Cells were incubated with thapsigargin (1 µM) during the final 10 min of reconstitution followed by the application of the identical protocol described in (A). **(E)** Cells were perfused in an intracellular medium containing 2,5-di-tert-butylhydroquinone (tBHQ, 10 µM) followed by washout of the inhibitor. **(F)** Summary results (mean ± S.E.M.) with statistical analysis. ANOVA with Tukey's post hoc test was applied. **(G)** Cells were perfused as in (A) and, once the [Ca^{2+}]$_{EL}$ steady-state was reached, different conditions were tested (arrow): none (200 nM Ca^{2+}; Control), EGTA (0.1 mM)**,** tBHQ (10 µM), EGTA+ tBHQ, or removal of ATP (-ATP)**. (H)** Summary results of EL-Ca^{2+} exit responses shown in (G), obtained from at least 3 independent days. ANOVA with Tukey's post hoc test was applied. *p < 0.05, ***p < 0.001, and ****p < 0.0001.

Figure 3. IP$_3$ releases Ca^{2+} from the endo-lysosomal Ca^{2+} store. (A) Representative response to IP$_3$ (2 µM, 30 s) in wt HEK293T cells expressing ELGA1. **(B)** Representative EL-Ca^{2+} release (in L/Lt ratio) upon application of four consecutive pulses (30s) of increasing concentrations of IP$_3$ (in µM). **(C)** Concentration-dependent response of EL-Ca^{2+} release (expressed as % of maximal response obtained with 30 µM IP$_3$) upon addition of increasing concentrations of IP$_3$ (0.01, 0.1, 1 and 10 µM). Each point is the mean ± S.E.M. of 3-4 values obtained from independent transfections. **(D)** Representative response to IP$_3$ (2 µM, 30 s) in IP$_3$R-null HEK293 (IP3R-3KO) cell line expressing ELGA1. **(E)** Representative response to IP$_3$ in IP3R-3KO cells transiently expressing IP$_3$R1. **(F)** Statistical quantification of IP$_3$ induced-EL-Ca^{2+} release (mean ± S.E.M.) with the number of independent transfections in parenthesis. ANOVA with Tukey's post hoc test was applied. **(G)** Inhibition of the IP$_3$ EL-Ca^{2+} release by heparin (100 µg/mL) in ELGA1 expressing HEK293T cells. **(H)** Statistical quantification of heparin inhibition of IP$_3$ response. Data are expressed as mean ± S.E.M. In all bar graphs, numbers in parenthesis correspond to the number of independent transfections from at least 2-3 experiments. A Student´s *t* test was applied. ****p < 0.0001.

Figure 4. ELGA is functionally located to the endo-lysosome. (A) Colocalization of ELGA fluorescence with TRPML1-cherry (magenta) in either fixed or live HEK293T cells expressing both proteins. Scale bar in all images, 10 µm. **(B)** Quantitative colocalization analysis of confocal fluorescence microscopy images were expressed as Pearson's (*P*) correlation coefficient shown in (A). Each point represents an individual cell from at least 2 independent experiments (mean ± S.E.M). A Student's *t* test was applied. **(C)** Effect of expression of TRPML1 on the [Ca^{2+}] at steady-state recorded from HEK293T expressing either ELGA1 (EL) or ERGA1 (ER). Data are expressed as mean ± S.E.M. In all bar graphs, numbers in parenthesis correspond to the independent transfections. A Student's *t* test was applied. **(D)** Lack of correlation between resting [Ca^{2+}] in the ER and the EL recorded with ERGA1 or ELGA1, respectively, in six cell types: HEK, TRPML1-expressing HEK, 3KO HEK, HeLa, CHO cells and cultured murine cortical astrocytes. Data are expressed as mean ± S.E.M. ER-Ca^{2+} in astrocytes taken from (Rodríguez-Prados et al., 2020). **(E)** Representative ML-SA1 (20 µM, 30 s) responses inducing EL-Ca^{2+} releases (expressed as L/Lt) from HEK293T cells expressing TRPML1 measured with ELGA1. **(F)** Representative Ca^{2+} responses (expressed as L/Lt) to application of four consecutive pulses (30s) of increasing concentrations of ML-SA1. **(G)** Concentration-dependent EL-Ca^{2+} release (expressed as % of the maximal response) in response to increasing concentrations of ML-SA1. Each point is the mean ± S.E.M. of 3-4 values obtained from 3 independent experiments. **(H)** Effect of the TRPML1 inhibitor ML-SI1 (10 µM) on the ML-SA1 (20 µM) response (expressed as L/Lt) in TRPML1 transfected HEK293T cells. **(I)** Effect of PI(3,5)P$_2$ (1 µM) on the EL-Ca^{2+} release in TRPML1 expressing HEK293T cells. **(J)** Summary results and statistical quantification of EL-Ca^{2+} release responses under various experimental conditions. Control (non-transfected) cells and TRPML1 transfected HEK293T cells were stimulated with ML-SA1 or PI(3,5)P$_2$. Percentage of response related to total [Ca^{2+}]$_{EL}$ at the steady-state is indicated. Data are expressed as mean ± S.E.M. and numbers in parenthesis indicate total *n* of independent transfections. One-way ANOVA with a Tukey's post hoc test (MLSA1 data) or a Student's *t* test (PI(3,5)P$_2$) were applied. **(K)** Concentration-dependent response (expressed as L/Lt) for EL-Ca^{2+} release upon increasing concentrations of PI(3,5)P$_2$. **(L)** Summary results and statistical significance of the responses determined from three independent transfection are

shown in (K)**.** Results are expressed as mean ± S.E.M. An ANOVA with a Tukey's

post hoc test was applied. ****p < 0.0001 and ***p < 0.001.

Figure 5. ELGA is physically located to the endo-lysosome. (A) Colocalization

of endogenous IP$_3$R1 with various endo-lysosome markers in EGFP-IP$_3$R1 HeLa

cells using STED microscopy. Cells were transiently transfected with cherry-

TRPML1 (a-c) or mRFP-TPC2 (d-f). EGFP-IP$_3$R1 and LAMP2 were detected using

specific antibodies against EGFP or LAMP2. **(B)** Association (arrows) of

endogenous IP$_3$R3 with LAMP1 detected with specific antibodies anti-IP$_3$R3 and

anti-LAMP1 in fixed wild type HEK293 cells using STED Super-resolution

microscopy. **(C)** Colocalization (arrows) of endo-lysosomes stained with

LysoView650 (pink) and endogenous IP$_3$R1-EGFP in live Hela cells using live cell

STED microscopy. **(D)** Colocalization (arrows) of endo-lysosomes stained with

lysotracker red (pink) and endogenous IP$_3$R1-EGFP in HeLa cells using TIRFM.

Figure 6. Physiological endo-lysosomal Ca^{2+} release. (A) Representative EL-

Ca^{2+} responses (in L/Lt) to agonists (ML-SA1, 20 µM; ATP, 100 µM; Carbachol,

(CCh), 100 µM) in intact HEK293T cells expressing ELGA1. **(B)** Similar to (A), in

transiently expressing TRPML1 HEK293T cells. **(C)** Protocol similar to (A) applied

in MDCK cells. **(D)** Protocol similar to (A) applied in astrocytes. **(E)** Summary

results of EL-Ca^{2+} release induced by ML-SA1 and ATP in different cell types or

conditions. Data are expressed as mean ± S.E.M. and numbers in brackets

correspond to the total number of independent transfections. Responses are in

percentages related to total [Ca^{2+}]$_{EL}$ at the steady-state. **(F)** A proposed model of

Ca^{2+} dynamics in the endo-lysosome (see text for explanation).

METHODS

Gene construction

H. sapiens VAMP7 cDNA was amplified by PCR using specific primers (#324; forward: 5'-CCAaagcttGCCACCATGGCGATTCTTTTT-3'; #325; reverse: 5'-ATaagcttTTTCTTCACACAGCTTGGCCATG-3'; *HindIII* sites are in lowercase). This fragment was digested with *HindIII* and inserted upstream of the *HindIII* fragment encoding the GA or the GA1 cDNAs, in the pHSV plasmid leading to pHSV-ELGA or pHSV-ELGA1 plasmids, respectively. The GA gene is a fusion of EGFP and aequorin (Manjarres et al., 2008) containing the D119A mutation (Kendall et al., 1992). The GA1 (previously described as GAP1) gene is a fusion of GFP (C3) and aequorin containing the mutations D117A, D119A and D163A (Rodriguez-Garcia et al., 2014). For expression in bacteria, GA or GA1 cDNAs were cloned into the pET28a vector. For measurements of ER-Ca^{2+}, the pHSV-ERGA1 (previously described as erGAP1) was used (Rodriguez-Prados et al., 2015).

Virus production

pHSV-ELGA or pHSV-ELGA1 were packaged into HSV-1 amplicon particles using a deletion mutant packaging system. Briefly, these plasmids were transfected into 2-2 cells and then, infected with a defective 5dl 1.2 helper virus. Titration was performed as previously described (Alonso et al., 1996; Lim et al., 1996).

Bacterial expression of GA and GA1 proteins

GA or GA1 cloned in pET28a were transformed in *E. coli* BL21 (Stratagene). Bacteria were grown at 37º C to A$_{600}$ >0.6 in LB containing 40 mg/L kanamycin and induced with 0.05 mM isopropyl β-D-thiogalactoside (IPTG) for 5h at 30º. Cells were then pelleted by centrifugation at 6000 g for 30 min and sonicated in a buffer containing (in mM): NaCl, 50; DTT, 0.5; Tris-HCl 20, pH 7.5; and Roche protein inhibitors. The bacterial lysate was centrifuged at 30,000 g for 10 min. Protein was purified by using a HisPur Ni-NTA resin (ThermoFisher, 88222). Protein quantification was performed by Bradford (range 20-60 µg/µL).

Bioluminescence activity at various pHs

Measurements were performed in the Tecan Genios Pro 96-well plate reader.

Recombinant *E.coli* GA or GA1 protein was reconstituted with coelenterazine *n* (1

µM) for 20 min and then 5-10 µL were aliquoted into each well to a final volume of

100 µL with a buffer containing (in mM): KCl, 140; $MgCl_2$, 1; MOPS/Tris, 20, pH

7.2. The effects of pH were monitored using different buffers for each pH range:

acetate for pH in the range 4.5-5.5; MOPS/Na for pH range 6-6.5; and Hepes/Na

for pH 7.2. Maximal luminescence was acquired by injecting 10-25 µL of saturating

(50 mM) Ca^{2+}. Calibrations were performed in Ca^{2+}-free medium (0.1 mM EGTA

added), or in the same solutions containing increasing Ca^{2+} concentrations

between 50 µM and 50 mM. This Ca^{2+} solution (10 µL) was injected with a first

injector, read for 15 seconds and then a saturating Ca^{2+} solution (10 mM) was

injected with a second injector.

Cell culture and gene expression

HeLa (CCL-2), HEK 293-T (CLR-1573), CHO- K1 (CCL-61) or MDCK cells (CCL-

34), were all purchased from ATCC. EGFP-IP_3R1 HeLa cells were earlier

described (Thillaiappan et al., 2017). All cells were maintained in Dulbecco's

Modified Eagle Medium (DMEM, Invitrogen) supplemented with 10% fetal bovine

serum (FBS, Sigma), 2 mM L-glutamine, 100 µg/ml streptomycin and 100 U/ml

penicillin. IP_3R-null HEK293 (HEK-3KO) cells (Alzayady et al., 2016) were

maintained in DMEM/F-12 with GlutaMAX (ThermoFisher) supplemented with

10%, FBS. Periodic PCR screening confirmed that all cells were free of

mycoplasma infection.

Mouse cortical astrocytes were isolated using the protocol described previously

(Rodríguez-Prados et al., 2020) with the approval of the Institutional Animal Care

and Use Committee at the University of Valladolid (ethics approval reference

number: 10207729). Briefly, cortices were dissected from three P2-3 C57BL/6J Crl

(Charles Rivers) mice in cold HBSS and digested with 0.5 mg/mL papain and 0.04

mg/mL DNase dissolved in a Ca^{2+}- and Mg-free HBSS with 10 mM glucose and

0.1% bovine serum albumin (BSA) at 37ºC for 20 min. Cells were resuspended

with Minimal Essential Medium (MEM) supplemented with 2 mM glutamax, 27 mM

glucose, 100 U/mL penicillin, 100 µg/mL streptomycin and 10% (v/v) horse serum,

and plated on a poly-L-lysine coated 75 cm^2 flask. After 2 and 7 days, flasks were

slapped to remove neurons and other loosely attached cells such as microglia and oligodendrocytes. After one week, cells were trypsinized and seeded in the same medium onto poly-L-lysine coated 12 mm-diameter glass coverslips at a density of 2.5×10^4 cells/coverslip. The cultures were used within 1-2 weeks.

Cells were transiently transfected with 0.2 µg (HEK293T) or 0.2-0.5 µg (HeLa) pHSV-ELGA or pHSV-ELGA1 using Lipofectamine 2000 (Invitrogen), according to the manufacturer's instructions. In some experiments, this plasmid was cotransfected with other plasmids encoding: mRFP-TPC2 (0.1 µg); EGFP-trpML1 (0.02 µg); mCherry-trpML1 (0.05 µg; Addgene #135189); er-Ruby-GCaMP210 (0.5 µg); or IP$_3$R1 (0.02 µg). Stably transfected HeLa clones expressing ELGA were generated by cotransfection of pHSV-ELGA and pcDNA3 (carrying the *neomycin* gene). GFP positive cells were enriched by sorting in a FACSAria II followed by antibiotic selection with 800 µg/mL G418 and single cell cloning was performed by limited dilution. ELGA-HeLa clones (No. 3 and 6) were maintained with 200 µg/mL G418.

ELGAs expression in mouse cortical astrocyte cultures was achieved by infection with herpes HSV-ELGA, HSV-ELGA1 or HSV-ERGA1 (previously named erGAP1) amplicons at a multiplicity of infection ranging between 0.01 and 0.1, one day before use.

Ca^{2+} bioluminescence measurements

Cells expressing ELGA1 (or ELGA in some experiments) were seeded on 4-well plates at a density of 8×10^4 (HeLa, CHO, or MDCK cells) or 2×10^5 (HEK-293T and HEK-3KO cells). Cells were Ca^{2+} depleted by incubation with the reversible sarco-endoplasmic reticulum Ca^{2+}-ATPase (SERCA) inhibitor 2,5-di-tert-butylhydroquinone (TBH; 10 µM) in the extracellular buffer (145 mM NaCl; 5 mM KCl; 1 mM MgCl$_2$; 10 mM glucose; 10 mM Na-HEPES, pH 7.4) supplemented with 0.5 mM EGTA for 10 min at 22ºC. Apo-aequorin was then reconstituted by incubation with 1 µM *n* coelenterazine in the same medium during 1 hour prior to measurements. Cells were then placed in the luminometer (Cairn Research, UK). Light acquisition was recorded each second as cps.

Intact cells were perfused (5 mL/min) with the extracellular buffer containing 1 mM CaCl$_2$ plus additions as stated. In the experiments with permeabilized cells,

perfusion was performed with an intracellular-like (IL) medium with the following
composition (in mM): KCl, 140; KH_2PO_4, 1; $MgCl_2$ 1; Mg-ATP, 1; sodium
succinate, 2; sodium-pyruvate, 1; Na-HEPES, 20, pH 7.2. The cells were
permeabilized by perfusion of digitonin (50 µM) in the IL medium containing 0.5
mM EGTA during 1 min. The digitonin was washed out and the cells were then
switched to IL medium containing 200 nM Ca^{2+} made by blending titrated solutions
of EGTA and EGTA-Ca^{2+} in the required amounts, calculated using the program
MaxChelator (Patton et al., 2004). If not stated otherwise, all stimuli or drugs were
dissolved and applied in the IL-buffer containing 200 nM Ca^{2+} during 30s, as
indicated in the graphs. At the end of each run, cells were lysed with 100 µM
digitonin in a 10 mM $CaCl_2$ solution to release all the remaining aequorin
luminescence. The EL-Ca^{2+} was expressed as luminescence at a certain time
point (L) divided by total luminescence till that moment (L/Lt). All measurements
were performed at 22ºC. Calibration curve allowed to transform L/Lt into [Ca^{2+}]
(µM) (Rodriguez-Prados et al., 2015).

Immunofluorescence

2-5 x 10^4 HEK cells were seeded onto 12 mm-glass coverslips and fixed, either
with cold 100% methanol or with 4% PFA in PBS for 20 min, followed by
incubation with 10% goat serum, 0.2% saponin (or 1% goat serum with 0.1%
Tween-20 in some experiments) in PBS for 20 min at 22 ºC. The incubation with
the primary antibody was performed overnight at 4ºC. The following primary
antibodies were used: anti-IP3R3 (1:100; BD biosciences); anti-LAMP1 (1:500;
Cell Signaling); anti-LAMP2 (H4B4) antibody (1:100; DSHB / University of Iowa);
anti-EEA1 (1:200; BD Transduction Laboratories); anti-TOM20 (1:100; Santa
Cruz); anti-GFP (1:500; Invitrogen). The secondary antibodies (anti-mouse Alexa
Fluor 568-conjugated antibodies) were incubated for 1h at 22 ºC. Nuclei were
stained with 4',6-diamidino-2-phenylindole (DAPI). Cells were mounted with
VectashieldR (Vector).

Confocal microscopy

624 Intact or fixed cells were imaged under a 63X Plan-Apochromat (NA 1.4) oil
625 immersion objective using a confocal microscope (SP5 Leica) equipped with a

626 white laser. Images were analyzed with ImageJ software using the JACoP plugin
627 (Bolte and Cordelières, 2006).

STED microscopy

3D STED microscopy was performed using a Leica SP8-STED microscope
equipped with an HC PL APO CS2 x100/1.4NA oil immersion objective.
Fluorescence depletion was made by a white laser and using a LAS-X
deconvolution software. Some experiments were performed with an Abberior
Instruments Expert Line STED microscope equipped with an Olympus UPLSAPO
×100/1.4NA oil immersion objective. Sequential confocal and STED images were
obtained following excitation of Alexa Fluor 594 and STAR RED by 594 and 640
nm lasers, respectively. Both fluorophores were depleted in three dimensions with
a 775 nm pulsed STED laser. Z-stacks were obtained by collecting images at 50
nm intervals using the 3D STED mode. Rescue STED was employed to minimize
the light dosage.

TIRFM

Images were acquired using an Olympus IX81 inverted Total Internal Reflection
Fluorescence Microscopy (TIRFM) equipped with oil-immersion PLAPO OTIRFM
60x objective lens/1.45 numerical aperture. Olympus CellSens Dimensions 2.3
(Build 189987) software was used for imaging. The cells were illuminated using
488 or 568 nm laser and the emitted fluorescence was collected through a band-
pass filter by a Hamamatsu ORCA-Fusion CMOS camera. The angle of the
excitation beam was adjusted to achieve TIRF with a penetration depth of ~140
nm.

Statistical analysis

Results are expressed as mean ± S.E.M., as indicated. Data were analyzed with
the GraphPad Prism 9 software (San Diego, CA, USA). Statistical significance
between two groups was analyzed by the unpaired Student's t test, whereas more
than two groups were compared using one-way analysis of variance (ANOVA) and
Tukey post-test analysis. Differences were considered statistically significant when
probability (p) values were less or equal than 0.05 (*), 0.01 (**), 0.001 (***) or
0.0001(****). Sample sizes (n) refer to independent cell wells from, at least, 3
different days. Individual traces show representative experiments.

Funding

This work was supported by grants from the Ministerio de Ciencia e Innovación de España (PID2020-116086RB-I00 to M.T.A.), by the Programa Estratégico del IBGM, Escalera de Excelencia, Junta de Castilla y León (Ref. CLU-2019-02 to M.T.A.), from the BBSRC, UK (BB/T015853/1 and BB/W01551X/1 to S.P.), from the National Institute of Health (RO1DE019245 to D.Y.) and Maximizing Investigator's Research Award grant (R35GM144120 to V.M.B.).

Credit authorship contribution statement

FJA made the original observations; BC, PT-V, AD-L, CR, FJA, JR-R, BMM and DIY performed investigation, formal analysis, and interpretation of data; DIY, VM, JG-S, SP and MTA provided conceptualization and funding acquisition; MTA and SP wrote original draft; all authors reviewed and edited the manuscript.

Declaration of Competing interest

The authors declare no competing financial interests.

Acknowledgments

We thank Dr. H. Xu for the VAMP7 plasmid; Dr. De Juan-Sanz for erRuby-GCaMP210; and, C. Taylor for IP$_3$R1 cDNA and EGFP-IP$_3$R1 HeLa cells. We also thank Dr. T. Schimmang for his valuable comments and suggestions. We also thank I. López and J. Fernández, as well as the Servicio de Microscopía, (C.I.C., Salamanca), for technical assistance.

REFERENCES

Akter, F., Bonini, S., Ponnaiyan, S., Kögler-Mohrbacher, B., Bleibaum, F., Damme, M., Renard, B. Y. and Winter, D. (2023). Multi-Cell Line Analysis of Lysosomal Proteomes Reveals Unique Features and Novel Lysosomal Proteins. *Mol Cell Proteomics* **22**, 100509.

Albrecht, T., Zhao, Y., Nguyen, T. H., Campbell, R. E. and Johnson, J. D. (2015). Fluorescent biosensors illuminate calcium levels within defined beta-cell endosome subpopulations. *Cell Calcium* **57**, 263-274.

Alonso, M. T., Lim, F., Nunez, L., Represa, J., Giraldez, F. and Schimmang, T. (1996). HSV-1 vector mediated transfer of BDNF into cerebellar granule cells. *Neuroreport* **7**, 3105-3108.

Alzayady, K. J., Wang, L., Chandrasekhar, R., Wagner, L. E., 2nd, Van Petegem, F. and Yule, D. I. (2016). Defining the stoichiometry of inositol 1,4,5-trisphosphate binding required to initiate Ca2+ release. *Sci Signal* **9**, ra35.

Atakpa, P., Thillaiappan, N. B., Mataragka, S., Prole, D. L. and Taylor, C. W. (2018). IP(3) Receptors Preferentially Associate with ER-Lysosome Contact Sites and Selectively Deliver Ca(2+) to Lysosomes. *Cell Rep* **25**, 3180-3193.e3187.

Atakpa, P., van Marrewijk, L. M., Apta-Smith, M., Chakraborty, S. and Taylor, C. W. (2019). GPN does not release lysosomal Ca(2+) but evokes Ca(2+) release from the ER by increasing the cytosolic pH independently of cathepsin C. *J Cell Sci* **132**.

Bolte, S. and Cordelières, F. P. (2006). A guided tour into subcellular colocalization analysis in light microscopy. *J Microsc* **224**, 213-232.

Bright, N. A., Davis, L. J. and Luzio, J. P. (2016). Endolysosomes Are the Principal Intracellular Sites of Acid Hydrolase Activity. *Current biology : CB* **26**, 2233-2245.

Christensen, K. A., Myers, J. T. and Swanson, J. A. (2002). pH-dependent regulation of lysosomal calcium in macrophages. *J Cell Sci* **115**, 599-607.

Dong, X. P., Shen, D., Wang, X., Dawson, T., Li, X., Zhang, Q., Cheng, X., Zhang, Y., Weisman, L. S., Delling, M., et al. (2010). PI(3,5)P(2) controls membrane trafficking by direct activation of mucolipin Ca(2+) release channels in the endolysosome. *Nat Commun* **1**, 38.

Farrell, R. J., Bredvik, K. G., Hoppa, M. B., Hennigan, S. T., Brown, T. A. and Ryan, T. A. (2024). A ratiometric ER calcium sensor for quantitative comparisons across cell types and subcellular regions. *bioRxiv*.

Garrity, A. G., Wang, W., Collier, C. M., Levey, S. A., Gao, Q. and Xu, H. (2016). The endoplasmic reticulum, not the pH gradient, drives calcium refilling of lysosomes. *Elife* **5**.

Geller, A. I. (1991). A system, using neural cell lines, to characterize HSV-1 vectors containing genes which affect neuronal physiology, or neuronal promoters. *J Neurosci Methods* **36**, 91-103.

Gerasimenko, J. V., Sherwood, M., Tepikin, A. V., Petersen, O. H. and Gerasimenko,
O. V. (2006). NAADP, cADPR and IP3 all release Ca2+ from the endoplasmic
reticulum and an acidic store in the secretory granule area. *J Cell Sci* **119**, 226-
238.

Gunaratne, G. S., Brailoiu, E., Kumar, S., Yuan, Y., Slama, J. T., Walseth, T. F., Patel,
S. and Marchant, J. S. (2023). Convergent activation of two-pore channels
mediated by the NAADP-binding proteins JPT2 and LSM12. *Sci Signal* **16**,
eadg0485.

Huang, G., Bartlett, P. J., Thomas, A. P., Moreno, S. N. and Docampo, R. (2013).
Acidocalcisomes of Trypanosoma brucei have an inositol 1,4,5-trisphosphate
receptor that is required for growth and infectivity. *Proc Natl Acad Sci U S A* **110**,
1887-1892.

Johnson, D. E., Ostrowski, P., Jaumouillé, V. and Grinstein, S. (2016). The position of
lysosomes within the cell determines their luminal pH. *J Cell Biol* **212**, 677-692.

Johnson, F. H. and Shimomura, O. (1978). [30] Introduction to the bioluminescence of
medusae, with special reference to the photoprotein aequorin. In *Methods in*
Enzymology, pp. 271-291: Academic Press.

Kendall, J. M., Sala-Newby, G., Ghalaut, V., Dormer, R. L. and Campbell, A. K. (1992).
Engineering the CA(2+)-activated photoprotein aequorin with reduced affinity for
calcium. *Biochem Biophys Res Commun* **187**, 1091-1097.

Kilpatrick, B. S., Eden, E. R., Schapira, A. H., Futter, C. E. and Patel, S. (2013). Direct
mobilisation of lysosomal Ca2+ triggers complex Ca2+ signals. *J Cell Sci* **126**, 60-
66.

Kilpatrick, B. S., Yates, E., Grimm, C., Schapira, A. H. and Patel, S. (2016). Endo-
lysosomal TRP mucolipin-1 channels trigger global ER Ca2+ release and Ca2+
influx. *Journal of Cell Science* **129**, 3859-3867.

Lee, C. A. and Blackstone, C. (2020). ER morphology and endo-lysosomal crosstalk:
Functions and disease implications. *Biochim Biophys Acta Mol Cell Biol Lipids*
1865, 158544.

Lim, F., Hartley, D., Starr, P., Lang, P., Song, S., Yu, L., Wang, Y. and Geller, A. I.
(1996). Generation of high-titer defective HSV-1 vectors using an IE 2 deletion
mutant and quantitative study of expression in cultured cortical cells.
Biotechniques **20**, 460-469.

Lloyd-Evans, E., Morgan, A. J., He, X., Smith, D. A., Elliot-Smith, E., Sillence, D. J.,
Churchill, G. C., Schuchman, E. H., Galione, A. and Platt, F. M. (2008).
Niemann-Pick disease type C1 is a sphingosine storage disease that causes
deregulation of lysosomal calcium. *Nat Med* **14**, 1247-1255.

López-Sanjurjo, C. I., Tovey, S. C., Prole, D. L. and Taylor, C. W. (2013). Lysosomes
shape Ins(1,4,5)P3-evoked Ca2+ signals by selectively sequestering Ca2+
released from the endoplasmic reticulum. *J Cell Sci* **126**, 289-300.

Manjarres, I. M., Chamero, P., Domingo, B., Molina, F., Llopis, J., Alonso, M. T. and
Garcia-Sancho, J. (2008). Red and green aequorins for simultaneous monitoring
of Ca2+ signals from two different organelles. *Pflugers Arch* **455**, 961-970.

Melchionda, M., Pittman, J. K., Mayor, R. and Patel, S. (2016). Ca2+/H+ exchange by
acidic organelles regulates cell migration in vivo. *J Cell Biol* **212**, 803-813.

Mindell, J. A. (2012). Lysosomal acidification mechanisms. *Annu Rev Physiol* **74**, 69-86.

Mitchell, K. J., Pinton, P., Varadi, A., Tacchetti, C., Ainscow, E. K., Pozzan, T.,
Rizzuto, R. and Rutter, G. A. (2001). Dense core secretory vesicles revealed as a
dynamic Ca(2+) store in neuroendocrine cells with a vesicle-associated membrane
protein aequorin chimaera. *J Cell Biol* **155**, 41-51.

Morgan, A. J., Davis, L. C., Wagner, S. K., Lewis, A. M., Parrington, J., Churchill, G.
C. and Galione, A. (2013). Bidirectional Ca²⁺ signaling occurs between the
endoplasmic reticulum and acidic organelles. *J Cell Biol* **200**, 789-805.

Morgan, A. J., Platt, F. M., Lloyd-Evans, E. and Galione, A. (2011). Molecular
mechanisms of endolysosomal Ca2+ signalling in health and disease. *Biochem J*
439, 349-374.

Narayanaswamy, N., Chakraborty, K., Saminathan, A., Zeichner, E., Leung, K.,
Devany, J. and Krishnan, Y. (2019). A pH-correctable, DNA-based fluorescent
reporter for organellar calcium. *Nat Methods* **16**, 95-102.

Patel, S. and Docampo, R. (2010). Acidic calcium stores open for business: expanding
the potential for intracellular Ca2+ signaling. *Trends Cell Biol* **20**, 277-286.

Patel, S., Marchant, J. S. and Brailoiu, E. (2010). Two-pore channels: Regulation by
NAADP and customized roles in triggering calcium signals. *Cell Calcium* **47**, 480-
490.

Patton, C., Thompson, S. and Epel, D. (2004). Some precautions in using chelators to
buffer metals in biological solutions. *Cell Calcium* **35**, 427-431.

Pryor, P. R., Mullock, B. M., Bright, N. A., Gray, S. R. and Luzio, J. P. (2000). The role
of intraorganellar Ca(2+) in late endosome-lysosome heterotypic fusion and in the
reformation of lysosomes from hybrid organelles. *J Cell Biol* **149**, 1053-1062.

Pryor, P. R., Mullock, B. M., Bright, N. A., Lindsay, M. R., Gray, S. R., Richardson, S.
C., Stewart, A., James, D. E., Piper, R. C. and Luzio, J. P. (2004). Combinatorial
SNARE complexes with VAMP7 or VAMP8 define different late endocytic fusion
events. *EMBO Rep* **5**, 590-595.

Raiborg, C. and Stenmark, H. (2015). An ER clamp for endosome fission. *Embo j* **34**,
136-137.

Rodriguez-Garcia, A., Rojo-Ruiz, J., Navas-Navarro, P., Aulestia, F. J., Gallego-
Sandin, S., Garcia-Sancho, J. and Alonso, M. T. (2014). GAP, an aequorin-
based fluorescent indicator for imaging Ca2+ in organelles. *Proc Natl Acad Sci U S*
A **111**, 2584-2589.

Rodriguez-Prados, M., Rojo-Ruiz, J., Aulestia, F. J., Garcia-Sancho, J. and Alonso,
M. T. (2015). A new low-Ca(2)(+) affinity GAP indicator to monitor high Ca(2)(+) in
organelles by luminescence. *Cell Calcium* **58**, 558-564.

Rodríguez-Prados, M., Rojo-Ruiz, J., García-Sancho, J. and Alonso, M. T. (2020).
Direct monitoring of ER Ca(2+) dynamics reveals that Ca(2+) entry induces ER-
Ca(2+) release in astrocytes. *Pflugers Arch* **472**, 439-448.

Ronco, V., Potenza, D. M., Denti, F., Vullo, S., Gagliano, G., Tognolina, M., Guerra,
G., Pinton, P., Genazzani, A. A., Mapelli, L., et al. (2015). A novel Ca²⁺-mediated
cross-talk between endoplasmic reticulum and acidic organelles: implications for
NAADP-dependent Ca²⁺ signalling. *Cell Calcium* **57**, 89-100.

Ruas, M., Rietdorf, K., Arredouani, A., Davis, L. C., Lloyd-Evans, E., Koegel, H.,
Funnell, T. M., Morgan, A. J., Ward, J. A., Watanabe, K., et al. (2010). Purified
TPC isoforms form NAADP receptors with distinct roles for Ca(2+) signaling and
endolysosomal trafficking. *Curr Biol* **20**, 703-709.

Saito, R., Mu, Q., Yuan, Y., Rubio-Alarcón, M., Eznarriaga, M., Zhao, P., Gunaratne,
G., Kumar, S., Keller, M., Bracher, F., et al. (2023). Convergent activation of
Ca(2+) permeability in two-pore channel 2 through distinct molecular routes. *Sci*
Signal **16**, eadg0661.

SantoDomingo, J., Fonteriz, R. I., Lobaton, C. D., Montero, M., Moreno, A. and
Alvarez, J. (2010). Ca2+ dynamics in the secretory vesicles of neurosecretory
PC12 and INS1 cells. *Cell Mol Neurobiol* **30**, 1267-1274.

Santodomingo, J., Vay, L., Camacho, M., Hernández-Sanmiguel, E., Fonteriz, R. I.,
Lobatón, C. D., Montero, M., Moreno, A. and Alvarez, J. (2008). Calcium
dynamics in bovine adrenal medulla chromaffin cell secretory granules. *Eur J*
Neurosci **28**, 1265-1274.

Shen, D., Wang, X., Li, X., Zhang, X., Yao, Z., Dibble, S., Dong, X. P., Yu, T.,
Lieberman, A. P., Showalter, H. D., et al. (2012). Lipid storage disorders block
lysosomal trafficking by inhibiting a TRP channel and lysosomal calcium release.
Nat Commun **3**, 731.

Shimomura, O., Johnson, F. H. and Morise, H. (1974). Mechanism of the luminescent
intramolecular reaction of aequorin. *Biochemistry* **13**, 3278-3286.

Singh, J., Kaade, E., Muntel, J., Bruderer, R., Reiter, L., Thelen, M. and Winter, D.
(2020). Systematic Comparison of Strategies for the Enrichment of Lysosomes by
Data Independent Acquisition. *J Proteome Res* **19**, 371-381.

Thillaiappan, N. B., Chavda, A. P., Tovey, S. C., Prole, D. L. and Taylor, C. W. (2017).
Ca(2+) signals initiate at immobile IP(3) receptors adjacent to ER-plasma
membrane junctions. *Nat Commun* **8**, 1505.

Vassileva, K., Marsh, M. and Patel, S. (2020). Two-pore channels as master regulators
of membrane trafficking and endocytic well-being. *Curr Opin Physiol* **17**, 163-168.

Vats, S. and Galli, T. (2022). Role of SNAREs in Unconventional Secretion-Focus on the
VAMP7-Dependent Secretion. *Front Cell Dev Biol* **10**, 884020.

Xu, H. and Ren, D. (2015). Lysosomal physiology. *Annu Rev Physiol* **77**, 57-80.

Yang, J., Zhao, Z., Gu, M., Feng, X. and Xu, H. (2019). Release and uptake
mechanisms of vesicular Ca(2+) stores. *Protein Cell* **10**, 8-19.

Yuan, Y., Arige, V., Saito, R., Mu, Q., Brailoiu, G. C., Pereira, G. J. S., Bolsover, S. R.,
Keller, M., Bracher, F., Grimm, C., et al. (2024). Two-pore channel-2 and inositol
trisphosphate receptors coordinate Ca(2+) signals between lysosomes and the
endoplasmic reticulum. *Cell Rep* **43**, 113628.

Zhu, M. X., Ma, J., Parrington, J., Calcraft, P. J., Galione, A. and Evans, A. M. (2010).
Calcium signaling via two-pore channels: local or global, that is the question. *Am J*
Physiol Cell Physiol **298**, C430-441.

Fig.1

Fig.2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

A *wt HEK293*

1 mM Ca²⁺

MLSA1 ATP CCh

EL-Ca²⁺ (L/Lt × 10⁻⁴)

2 min

B *TRPML1 HEK293*

1 mM Ca²⁺

MLSA1 ATP

EL-Ca²⁺ (L/Lt × 10⁻⁴)

2 min

C MDCK

1 mM Ca²⁺

MLSA1 ATP

EL-Ca²⁺ (L/Lt × 10⁻⁴)

1 min

D *Cortical astrocytes*

1 mM Ca²⁺

MLSA1 ATP

EL-Ca²⁺ (L/Lt × 10⁻⁴)

1 min

E

■ MLSA 1
■ ATP

% EL-Ca²⁺ release

HEKT TRPML1-HEKT HeLa MDCK Astrocyte

F

SERCA
?
EL
TRPML1
IP₃R IP₃R IP₃R
ER
Local signal Global signal