Title: Treatment with furosemide indirectly increases inhibitory transmission in the developing hippocampus

Authors: C. Peerboom[1], T. Wijne[1] and C.J. Wierenga[1,2]

Corresponding author: Corette J. Wierenga

Affiliations:

[1] Cell Biology, Neurobiology and Biophysics, Biology department, Utrecht University, 3584 CH, Utrecht, the Netherlands,

[2] Donders Institute for Brain, Cognition and Behaviour, Radboud University, 6525 AJ, Nijmegen, the Netherlands

Author contributions

C.P., and C.J.W. designed research; C.P and T.W., performed research and analyzed data; C.P., and C.J.W. wrote the paper.

Declaration of Interests

The authors declare no competing interests.

Acknowledgements

This research was supported by a TOP grant from ZonMW (#91216021) and by the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO; #OCENW.KLEIN.150). We thank Prof. Thomas Kuner for providing the Cre-dependent SClm construct (Boffi et al., 2018). We thank Prof. Kevin Staley for providing the SuperClomeleon[lox/-] mouse line, Dr. Stefan Berger for the CamKIIα[Cre/-] mice and Dr. Henk Karst for sharing these mice with us. We thank Dunya Selemangel for help with SuperClomeleon data analysis and René van Dorland for excellent technical support.

Abstract

During the first two postnatal weeks intraneuronal chloride concentrations in rodents

gradually decrease, causing a shift from depolarizing to hyperpolarizing γ-aminobutyric acid

(GABA) responses. GABAergic depolarization in the immature brain is crucial for the formation

and maturation of excitatory synapses, but when GABAergic signaling becomes inhibitory it

no longer promotes synapse formation. Here we examined the role of chloride transporters

in developing postnatal hippocampal neurons using furosemide, an inhibitor of the chloride

importer NKCC1 and chloride exporter KCC2 with reported anticonvulsant effects. We treated

organotypic hippocampal cultures made from 6 to 7-day old mice with 200 μM furosemide

from DIV1 to DIV8. Using perforated patch clamp recordings we observed that the GABA

reversal potential was depolarized after acute furosemide application, but after a week of

furosemide treatment the GABA reversal potential but was more hyperpolarized compared

to control. Expression levels of the chloride cotransporters were unaffected after one week

furosemide treatment. This suggests that furosemide inhibited KCC2 acutely, while prolonged

treatment resulted in (additional) inhibition of NKCC1, but we cannot exclude changes in

HCO_3^-. We assessed the effects of accelerating the GABA shift by furosemide treatment on

inhibitory synapses onto CA1 pyramidal cells. Directly after cessation of furosemide treatment

at DIV9, inhibitory synapses were not affected. However at DIV21, two weeks after ending

the treatment, we found that the frequency of inhibitory currents was increased, and VGAT

puncta density in *stratum Radiatum* was increased. In addition, cell capacitance of CA1

pyramidal neurons was reduced in furosemide-treated slices at DIV21 in an activity-

dependent manner. Our results suggest that furosemide indirectly promotes inhibitory

transmission, but the underlying mechanism remains unresolved. The furosemide-induced

increase in inhibitory transmission might constitute an additional mechanism via which

furosemide reduces seizure susceptibility in the epileptic brain.

Introduction

γ-Aminobutyric acid (GABA) is the main inhibitory, hyperpolarizing neurotransmitter in the adult brain, but during early development GABA actually depolarizes neurons. Ionotropic GABA$_A$ receptors primarily conduct chloride (~70%), but also HCO$_3^-$ ions (~30%) (Kaila et al., 2014). While HCO$_3^-$ levels remain constant, intracellular chloride levels decrease during brain development (Kaila et al., 2014). This developmental decrease in chloride causes the reversal potential (E$_{GABA}$) to gradually drop below resting membrane potential. As a result, the GABAergic driving force shifts from depolarizing in immature neurons to hyperpolarizing in the mature brain (Rivera et al., 1999; Hübner et al., 2001; Tyzio et al., 2007; Romo-Parra et al., 2008; Kirmse et al., 2015; Tsukahara et al., 2015; Sulis Sato et al., 2017; Murata and Colonnese, 2020). The decrease in chloride levels is the direct result of an increased function of the chloride exporter KCC2 (K-Cl cotransporter 2) relative to the chloride importer NKCC1 (Na-K-2Cl cotransporter 1) (Rivera et al., 1999; Gulyás et al., 2001; Yamada et al., 2004; Dzhala et al., 2005; Otsu et al., 2020). In humans, the shift in chloride transporter expression occurs during the first year after birth (Dzhala et al., 2005; Sedmak et al., 2016). In rodents, the GABA shifts during the first two postnatal weeks (Rivera et al., 1999; Stein et al., 2004; Tyzio et al., 2007; Romo-Parra et al., 2008; Kirmse et al., 2015; Sulis Sato et al., 2017; Murata and Colonnese, 2020).

Depolarizing GABA signaling plays a critical role in the formation and maturation of excitatory synapses in the developing brain (Leinekugel et al., 1997; Akerman and Cline, 2006; Wang and Kriegstein, 2008, 2011; Chancey et al., 2013; van Rheede et al., 2015; Oh et al., 2016). However, after the first postnatal week, GABA becomes inhibitory and it no longer promotes synapse formation (Salmon et al., 2020; Peerboom et al., 2023). Here we examined the role of chloride transporters in developing postnatal hippocampal neurons using furosemide, a well-known inhibitor of chloride transport. Low furosemide concentrations (100-1000 μM) have been employed previously to block KCC2 and elevate chloride concentrations in (slice) cultures (Thompson and Gahwiler, 1989; Jarolimek et al., 1999; Deeb et al., 2013; Wright et al., 2017), as furosemide has a preference for inhibiting KCC2 over NKCC1 (IC$_{50}$ ~30 μM for rat NKCC1 and IC$_{50}$~10 μM for rat KCC2, expressed in Xenopus oocytes (Orlov et al., 2015)). In addition, furosemide can inhibit α6- and α4-subunit containing GABA$_A$ receptors (Thompson

and Gahwiler, 1989; Pearce, 1993; Korpi et al., 1995), and proteins that regulate neuronal HCO_3^- levels (Halligan et al., 1991; Temperini et al., 2009; Ruusuvuori and Kaila, 2014; Uwera et al., 2015). Moreover, furosemide was shown to inhibit activity-induced swelling of epileptic brain tissue (Hochman et al., 1995; Gutschmidt et al., 1999; Hochman, 2012).

Furosemide is commonly used as a diuretic, but has anti-epileptic actions as well. A case-control study on hypertension and epilepsy showed that diuretics including furosemide protected patients with hypertension against an increased risk on seizures in later life (Hesdorffer et al., 1996). Subsequently, the antiepileptic effects of furosemide were demonstrated in several animal models (Hochman et al., 1995; Gutschmidt et al., 1999; Holtkamp et al., 2003; Barbaro et al., 2004; Viitanen et al., 2010; Hochman, 2012; Uwera et al., 2015; Chen et al., 2022). Since then, several epidemiological and experimental studies have reported anticonvulsant actions of furosemide, but because of its wide range of actions, the mechanism behind furosemide's anti-epileptic effects remains mostly unclear (Hesdorffer et al., 1996; Staley, 2002; Maa et al., 2011).

We examined the effects of 200 µM furosemide in organotypic hippocampal cultures. We observed that acute furosemide application depolarized GABA reversal potential, in line with a preferential inhibition of KCC2. In contrast, furosemide treatment from DIV1 to DIV8 resulted in hyperpolarization of the GABA reversal potential. Expression levels of the chloride-cotransporters were unaffected after furosemide-treatment. This suggests that sustained application of furosemide resulted in inhibition of NKCC1, although we cannot exclude changes in HCO_3^- levels. We assessed the consequences of the accelerated GABA shift during furosemide treatment on inhibitory transmission. We found that inhibitory synapses were not affected directly after furosemide treatment at DIV9. However at DIV21, when E_{GABA} had normalized, inhibitory currents were strongly increased in furosemide-treated slices, possibly via an increased number of inhibitory synapses in the *stratum Radiatum*. In addition, we found evidence for shrinkage of CA1 pyramidal neurons in furosemide-treated slices at DIV21. Our results suggest that furosemide may indirectly promote inhibitory transmission, which could constitute a new mechanism via which furosemide can reduce seizures in the epileptic brain.

Methods

Animals

All animal experiments were performed in compliance with the guidelines for the welfare of experimental animals issued by the Federal Government of the Netherlands and were approved by the Dutch Central Committee Animal experiments (CCD), project AVD1080020173847 and project AVD1150020184927. In this study we used male and female transgenic mice: GAD65-GFP mice (López-Bendito et al., 2004) (bred as a heterozygous line in a C57BL/6JRj background), C57BL/6JRj littermates, and SuperClomeleon (SClm) mice. The SClm mice (Herstel et al., 2022) are SuperClomeleon$^{lox/-}$ mice (Rahmati et al., 2021), a gift from Kevin Staley (Massachusetts General Hospital, Boston, MA) crossed with CamKIIα$^{Cre/-}$ mice (Tsien et al., 1996; Casanova et al., 2001a). SClm mice express the chloride sensor SClm in up to 70% of the pyramidal neurons in the hippocampus (Casanova et al., 2001b; Wang et al., 2013). GAD65-GFP mice express GFP in ~20% of GABAergic interneurons in the CA1 region of the hippocampus (Wierenga et al., 2010). As we did not detect any differences between slices from male and female mice as well as between GAD65-GFP mice and C57BL/6JRj littermates, all data were pooled.

Organotypic hippocampal culture preparation and furosemide-treatment

Organotypic hippocampal cultures were made from P6-7 mice as described before (Hu et al., 2019; Herstel et al., 2022), based on the *Stoppini method* (Stoppini et al., 1991). Mice were decapitated and their brain was rapidly placed in ice-cold Grey's Balanced Salt Solution (GBSS; containing (in mM): 137 NaCl, 5 KCl, 1.5 CaCl$_2$, 1 MgCl$_2$, 0.3 MgSO$_4$, 0.2 KH$_2$PO$_4$ and 0.85 Na$_2$HPO$_4$) with 25 mM glucose, 12.5 mM HEPES and 1 mM kynurenic acid (pH set at 7.2, osmolarity set at ~320 mOsm, sterile filtered). 400 μm thick transverse hippocampal slices were cut with a McIlwain tissue chopper. Slices were placed on Millicell membrane inserts (Millipore) in wells containing 1 ml culture medium (consisting of 48% MEM, 25% HBSS, 25% horse serum, 25 mM glucose, and 12.5 mM HEPES, with an osmolarity of ~325 mOsm and a pH of 7.3 – 7.4). Slices were stored in an incubator (35°C with 5% CO$_2$). Culture medium was replaced by culture medium supplemented with 0.1% dimethyl sulfoxide (DMSO, Sigma-Aldrich) or 200 μM furosemide (furosemide, Merck, in 0.1 % DMSO). In addition, a small drop

of medium supplemented with DMSO or furosemide was carefully placed on top of the slices.

In this way, cultures were treated 3 times per week until DIV8. From DIV8 onwards, cultures

received normal culture medium 3 times per week. Experiments were performed at day in

vitro 1-3 (DIV2), 8-10 (DIV9) or 20-22 (DIV21). Please note that perforated patch experiments

at DIV9 were performed 1 to 55 hr after cessation of the furosemide treatment. E_{GABA} did not

correlate with hours after treatment and was not affected by addition of furosemide to the

ACSF during the perforated patch recording.

Electrophysiology and analysis

Organotypic cultures were transferred to a recording chamber and continuously perfused

with carbonated (95% O_2, 5% CO_2) artificial cerebrospinal fluid (ACSF, in mM: 126 NaCl, 3 KCl,

2.5 $CaCl_2$, 1.3 $MgCl_2$, 26 $NaHCO_3$, 1.25 NaH_2PO_4, 20 glucose; with an osmolarity of ~310 mOsm)

at a rate of approximately 1 ml/min. Bath temperature was maintained at 29-32°C.

Recordings were acquired using an MultiClamp 700B amplifier (Molecular Devices), filtered

with a 3 kHz Bessel filter and stored using pClamp10 software. For acute application, ACSF

containing 0.1% DMSO or 200 µM furosemide (dissolved in 0.1% DMSO) was bath applied for

5 minutes.

Perforated patch clamp recordings were performed in CA1 pyramidal neurons. Recording

pipettes with resistances of 2-4 MΩ were pulled from borosilicate glass capillaries (World

Precision Instruments). The pipette tip was filled with gramicidin-free KCl solution (140 mM

KCl and 10 mM HEPES, pH adjusted to 7.2 with KOH, and osmolarity 285 mOsm/l) and then

backfilled with solution containing gramicidin (60 µg/ml, Sigma). Neurons were held at -65

mV and the access resistance of the perforated cells was monitored constantly before and

during recordings by a 5 mV seal test. An access resistance of 50 MΩ was considered

acceptable to start recording. Recordings were excluded if the resting membrane potential

exceeded -50 mV or if the series resistance after the recording deviated more than 30% from

its original value. $GABA_A$ currents were induced by upon local application of 50 µM muscimol

dissolved in HEPES-ACSF (containing in mM: 135 NaCl, 3 KCl, 2.5 $CaCl_2$, 1.3 $MgCl_2$, 1.25

$Na_2H_2PO_4$, 20 Glucose, and 10 HEPES) every 30 sec using a Picospritzer II. Muscimol responses

were recorded in the presence of 1 µM TTX (Abcam) at holding potentials between -100 mV

and -30 mV in 10 mV steps. We plotted response amplitude as a function of holding potential

and calculated the chloride reversal potential from the intersection of the linear current-

voltage curve with the x-axis. To assess the size of GABA$_A$ currents independent of holding

and reversal potential, we calculated the slope of the current-voltage-plot.

For recordings of spontaneous inhibitory postsynaptic currents (sIPSCs) and miniature

inhibitory currents (mIPSCs) we performed whole-cell patch clamp recordings from CA1

pyramidal neurons using borosilicate glass pipettes with resistances of 3-6 MΩ. Pipettes were

filled with a KCl-based internal solution (containing in mM: 70 K-gluconate, 70 KCl, 0.5

ethyleneglycol- bis(β-aminoethyl)-N,N,N',N'-tetraacetic Acid (EGTA), 4 Na$_2$phosphocreatine,

4 MgATP, 0.4 NaGTP and 10 HEPES; pH adjusted to 7.2 with KOH) and ACSF was supplemented

with 20 µM DNQX (Tocris) and 50 µM DL-APV (Tocris). For recordings of miniature IPSCs

(mIPSCs), 1 µM TTX (Abcam) was also added to the ACSF. sIPSCs and mIPSCs were recorded

for 6 minutes. For wash in experiments, an sIPSC baseline was recorded for 6 minutes and

compared to sIPSCs after 20 minutes wash in of 1 µM TTX (Abcam), 0.2 µM K252a (Tocris), 5

µM AM251 (Sigma) or 1 µM agatoxin-IVA (smartox-biotech), again for 6 minutes. Membrane

capacitance and series resistance were monitored during the recordings by a 5 mV seal test.

Cells were discarded if series resistance was above 35 MΩ or if the resting membrane

potential exceeded -50 mV or if the series resistance after the recording deviated more than

30% from its original value.

All data were blinded before analysis. Events were selected using a template in Clampfit10

software. Further analysis was performed using custom-written MATLAB scripts. Rise time of

sIPSCs was determined as the time between 10% and 90% of the peak value. The decay tau

was fitted with a single exponential function. Only events with a goodness of fit $R^2 \geq 0.75$ were

included.

Two-photon SuperClomeleon imaging and analysis

We performed two-photon chloride imaging in CA1 pyramidal cells using the SuperClomeleon

sensor (Grimley et al., 2013) in cultured slices from SClm mice (Herstel et al., 2022). The SClm

sensor consists of Cerulean (a CFP mutant) and Topaz (a YFP mutant) joined by a flexible

linker. In the absence of chloride, Fluorescence Resonance Energy Transfer (FRET) occurs from

the CFP donor to the YFP acceptor. Chloride binding to YFP increases the distance between

the fluorophores and decreases FRET (Grimley et al., 2013).

At the recording day, slices were transferred to a recording chamber and continuously perfused with carbonated (95% O_2, 5% CO_2) ACSF supplemented with 1 mM Trolox, at a rate of approximately 1 ml/min. Bath temperature was monitored and maintained at 30-32 °C. Two-photon imaging of CA1 pyramidal neurons or VGAT-positive interneurons in the CA1 area of the hippocampus was performed using a customized two-photon laser scanning microscope (Femto3D-RC, Femtonics, Budapest, Hungary) with a Ti-Sapphire femtosecond pulsed laser (MaiTai HP, Spectra-Physics) and a 60x water immersion objective (Nikon NIR Apochromat; NA 1.0) The CFP donor was excited at 840 nm. The emission light was split using a dichroic beam splitter at 505 nm and detected using two GaAsP photomultiplier tubes. We collected fluorescence emission of Cerulean/CFP (485 ± 15 nm) and Topaz/YFP (535 ± 15 nm) in parallel. Of each slice from SClm mice, 2-5 image stacks were acquired in different fields of view. The resolution was 8.1 pixels/µm (1024x1024 pixels, 126x126 µm) with 1 µm z-steps.

Image analysis was performed using ImageJ software, as described before (Herstel et al., 2022). We manually determined regions of interest (ROIs) around individual neuron somata. To select a representative cell population in SClm slices, in each image z-stack we selected four z-planes at comparable depths in which three pyramidal cells were identified that varied in CFP brightness (bright, middle and dark). We subtracted the mean fluorescence intensity of the background in the same image plane before calculating the fluorescence ratio of CFP and YFP. We limited our analysis to cells which were located within 450 pixels from the center of the image, as FRET ratios showed slight aberrations at the edge of our images. We excluded cells with a FRET ratio < 0.5 or > 1.6 to avoid unhealthy cells.

Protein extraction and Western Blot analysis

Organotypic hippocampal cultures were washed in cold PBS and subsequently lysed in cold protein extraction buffer containing: 150 mM NaCl, 10 mM EDTA 10 mM HEPES, 1% Triton-X100 and 1x protease and 1x phosphatase inhibitors cocktails (Complete Mini EDTA-free and phosSTOP, Roche). Lysates were cleared of debris by centrifugation (14,000 rpm, 1 min, 4°C) and measured for protein concentration before storage at −20°C until use. Lysates were denatured by adding loading buffer and heating to 95°C for 5 min. For each sample an equal mass of proteins was resolved on 4–15 % polyacrylamide gel (Bio-Rad). The proteins were

then transferred (300 mA, 3h) onto ethanol-activated Immobilon-P PVDF membrane

(Millipore) before blocking with 3% Bovine Serum Albumin in Tris- buffered saline-Tween

(TBST, 20 mM Tris, 150 mM NaCl, 0.1% Tween-20) for 1 h. Primary antibodies used in this

study were: mouse anti-NKCC1 (T4, Developmental Studies Hybridoma Bank, 1:1000), rabbit

anti-KCC2 (07-432, Merck, 1:1000) and s940-pKCC2 (p1551-940, LuBioSciences, 1:1000),

mouse anti-tubulin (T-5168, Sigma, 1:10000). Primary antibodies were diluted in blocking

buffer and incubated with the blots overnight at 4 °C under gentle rotation. The membrane

was washed 3 times 15 minutes with TBST before a 1h incubation of horseradish peroxidase

(HRP)-conjugated antibodies (P0447 goat anti-mouse IgG HRP, Dako, 1:2500 or P0399 swine

anti-rabbit IgG HRP, Dako, 1:2500), and washed 3 times 15 minutes in TBST again before

chemiluminescence detection. For chemiluminescence detection, blots were incubated with

Enhanced luminol-based Chemiluminescence substrate (Promega), and the exposure was

captured using the ImageQuant 800 system (Amersham). Images were analyzed in ImageJ, by

drawing rectangular boxes around each band and measuring average intensities. Protein

levels were normalized to the Tubulin loading controls.

Immunohistochemistry, confocal microscopy and analysis

Slices were fixed 4 % paraformaldehyde solution in PBS for 30 minutes at room temperature.

After washing slices in phosphate buffered saline (PBS), they were permeabilized for 15

minutes in 0.5 % Triton X-100 in PBS for 15 minutes, followed by 1 hour in a blocking solution

consisting of 10 % normal goat serum and 0.2 % Triton X-100 in PBS. Slices were incubated in

primary antibody solution at 4 °C overnight. The following primary antibodies were used:

rabbit polyclonal anti-KCC2 (07-432, Merck, 1:1000), rabbit anti-VGAT (131003, Synaptic

Systems, 1:1000), guinea pig anti-VGLUT (AB5905, Merck, 1:1000), mouse anti-NeuN

(MAB377, Millipore, 1:500), guinea pig anti-NeuN (ABN90, Merck, 1:1000). Slices were

washed in PBS and incubated in secondary antibody solution for four hours at room

temperature. Secondary antibodies used were: goat anti-mouse Alexa Fluor 467 (Life

Technologies, A21236, 1:500), goat anti-rabbit Alexa Fluor 405 (A31556, Life Technologies,

1:500), goat anti-guinea pig Alexa Fluor 488 (A11073, Life Technologies, 1:1000) and Goat

anti-guineapig Alexa Fluor 568 (A11075, Life Technologies, 1:500) Goat anti-rabbit Alexa fluor

647 (A21245, Life Technologies, 1:500). After another PBS wash, slices were mounted in

Vectashield mounting medium (Vector labs).

Confocal images were taken on a Zeiss LSM-700 confocal laser scanning microscopy system with a Plan-Apochromat 40x 1.3 NA immersion objective for KCC2 staining and 63x 1.4 NA oil immersion objective for inhibitory synapse staining. KCC2 staining was imaged in two fields of view in the CA1 *sPyr* of each slice. Inhibitory synapses were imaged in the CA1 *sPyr* and *sRad* of each slice. Image stacks (1024x1024 pixels) were acquired at 6.55 pixels/µm for KCC2 staining, 10.08 pixels/µm for VGLUT/VGAT staining. Step size in z was 0.5 µm for KCC2 staining and 0.4 µm for inhibitory synapse staining.

Confocal images were blinded before analysis. For the analysis of KCC2 levels in control and furosemide-treated slices, ROIs were drawn around 5 NeuN positive cell bodies and this was repeated in 1-3 z-planes (depending on the depth of KCC2 signal) at comparable depths, to measure average KCC2 fluorescence of 5-15 neurons per image stack. For analysis of inhibitory synapses a custom-made macro was used (Ruiter et al., 2020). Briefly, three Z-planes were averaged and background was subtracted with rolling ball radius of 10 pixels. VGAT puncta were identified using watershed segmentation. PV and CB1R channels were thresholded and colocalization with VGAT puncta was analyzed.

Statistical Analyses

Statistical analysis was performed in Prism (Graphpad). Normality was tested using the D'Agostino & Pearson test. For the comparison of two groups we either used an unpaired Student's t test (UT; parametric), a Mann-Whitney test (MW; non-parametric), a paired Student's t test (PT; parametric) or a Wilcoxon signed-rank test (WSR; non-parametric). For comparison of multiple groups, a Kruskal Wallis test (KW; nonparametric) was used, followed by a Dunn's Multiple Comparison posthoc test (DMC) or a two way ANOVA (2W ANOVA; parametric), followed by a by a Sidak's Multiple Comparisons posthoc test (SMC). Data are presented as mean ± standard error of the mean. Significance is reported as *$p < 0.05$; **$p \leq 0.01$; ***$p \leq 0.001$.

Results

Furosemide depolarizes GABA responses acutely, but hyperpolarizes GABA responses after one week treatment.

We used perforated patch recordings upon muscimol application in CA1 pyramidal neurons of cultured hippocampal slices to record the reversal potential of $GABA_A$ receptor currents. Acute administration of 200 µM furosemide in DIV2 slices did not affect the GABA reversal potential (E_{GABA}) (Fig. 1A) or GABA driving force (GABA DF) (Fig. 1B), in agreement with a low expression of KCC2 at this age (Salmon et al., 2020; Peerboom et al., 2023). At DIV21, when KCC2 expression is high (Salmon et al., 2020; Peerboom et al., 2023), acute furosemide application elevated E_{GABA} and GABA DF (Fig. 1C,D). This is in agreement with previous reports (Thompson and Gahwiler, 1989; Jarolimek et al., 1999; Deeb et al., 2013; Wright et al., 2017) and comparable with acute inhibition of KCC2 by the specific blocker VU0463271 (VU) in slices at the same developmental stage (Peerboom et al., 2023).

We then treated cultured hippocampal slices from DIV2 to DIV8 with 200 µM furosemide. Surprisingly, we observed a strong hyperpolarization of E_{GABA} and a negative shift of the GABA DF at DIV9 after one week of furosemide treatment (Fig. 1E,F). Since furosemide has been shown to inhibit α6- and α4-subunit containing $GABA_A$ receptors (Thompson and Gahwiler, 1989; Pearce, 1993; Korpi et al., 1995), we checked if $GABA_A$ receptor inhibition may have interfered with our measurements of E_{GABA} and GABA DF. As a rough indication, we determined the slope of the current-voltage (IV) plot to assess GABA response amplitudes independently of holding potential and reversal potential. However, IV-slopes were not different (MW, p=0.46), suggesting that furosemide did not have a major effect on $GABA_A$ receptors.

 A hyperpolarizing shift in E_{GABA} suggest that furosemide treatment resulted in a reduction of intraneuronal chloride or HCO_3^- levels. To directly assess intracellular chloride levels after furosemide treatment, we employed the chloride sensor SuperClomeleon (SClm) (Grimley et al., 2013; Boffi et al., 2018; Rahmati et al., 2021). We used slices from SClm mice in which CA1 pyramidal cells express the SClm sensor and we recorded FRET ratios in control and furosemide-treated slices using two-photon microscopy (Herstel et al., 2022). Remarkably,

we did not observe a difference in SClm FRET ratios after one week of furosemide treatment

(Fig. 1G,H). The discrepancy between our results from perforated patch recordings and SClm

imaging may be due to the decreased sensitivity of the SClm sensor for low chloride levels

(Grimley et al., 2013; Herstel et al., 2022). Alternatively, it may reflect concomitant changes

in chloride and HCO_3^-, as SClm displays substantial pH-sensitivity (Grimley et al., 2013; Boffi

et al., 2018; Lodovichi et al., 2022). These possibilities are discussed in more detail in the

discussion section.

To examine if the hyperpolarizing GABA signaling after furosemide treatment was due to

changes in chloride cotransporter expression, we assessed total NKCC1, KCC2 or s940-pKCC2

protein levels in control and furosemide-treated slices using Western blots. No significant

changes in the levels of NKCC1, KCC2 or S940-pKCC2 were observed (Fig. 1I,J). In addition, we

examined KCC2 localization in treated and control slices using immunohistochemistry. We

used NeuN to identify individual cell bodies and we estimated KCC2 levels in somata and

membranes. We previously showed that we can detect a reduction in KCC2 levels after shRNA

using these methods (Peerboom et al., 2023). Furosemide treatment did not significantly

affect KCC2 levels in somata or membranes of CA1 pyramidal neurons (Fig. 1K,L). Together,

this indicates that furosemide treatment during DIV2 to DIV8 hyperpolarized GABA signaling

without changing the expression or surface levels of KCC2, nor by changing expression levels

of NKCC1 and S940-pKCC2. These results suggest that furosemide treatment hyperpolarized

E_{GABA} via inhibition of chloride transporters KCC2 and NKCC1 without altering their expression.

Furosemide treatment does not affect inhibitory synapses at DIV9.

The results above show that one week of furosemide treatment hyperpolarized E_{GABA}, thereby

accelerating the GABA shift. Next, we assessed the consequences of this accelerated GABA

shift on inhibitory transmission. We recorded spontaneous and miniature inhibitory currents

(sIPSCs and mIPSCs) using whole-cell voltage clamp recordings in CA1 pyramidal cells in

control and furosemide-treated slices at DIV9. We found that the frequency and amplitude of

sIPSCs were similar in control and furosemide-treated slices at DIV9 (Fig. 2A,B). There were

also no differences in rise and decay kinetics and the membrane capacitance (C_m) of CA1

pyramidal neurons (Fig. 2C-E). This indicates that GABA signaling does not directly affect the

development of inhibitory synapses at this developmental stage (Peerboom et al., 2023).

When recording sIPSCs, we noticed that the sIPSC frequency slightly increased by ~20% during

the 20 minutes of recording in both control and furosemide-treated slices (Fig. 2F,G). When

we washed in TTX to block neuronal activity in control slices, this 'run-up' was prevented (Fig.

2H), indicating that the run-up is activity dependent. However, TTX did not prevent the run-

up in furosemide-treated slices, suggesting that the run up had different mechanisms in

control and furosemide-treated slices (Fig. 2I). We considered the possibility that the activity-

dependent run up is due to release of activity-dependent factors, such as endocannabinoids

or BDNF. We assessed the contribution of the BDNF receptor TrkB by washing in TrkB

antagonist K252a during sIPSC recordings in control and furosemide-treated slices. We found

that sIPSCs were enhanced after wash in K252a, suggesting that there is ongoing suppression

of GABA release by BDNF in our slices at DIV9. However, there was no differential contribution

of BDNF between control and furosemide-treated slices (Fig. 2J). We also washed in AM251,

a specific antagonist of the endocannabinoid receptor CB1. We found that endocannabinoid

signaling does not have a prominent influence on sIPSCs, and there was no difference

between control and furosemide-treated slices (Fig. 2J). Blocking presynaptic P/Q calcium

channels with agatoxin-IVA (aga) (Goswami et al., 2012) also did not significantly alter sIPSC

frequency in control and furosemide-treated slices (Fig. 2J). It therefore remains unclear why

activity-dependent run up of sIPSCs was different in furosemide-treated slices.

Inhibitory transmission is increased in furosemide-treated slices at DIV21.

Two weeks after ending the treatment at DIV21, E_{GABA} and GABA DF in pyramidal neurons had

normalized and reached control levels (Fig. 3A,B). We previously found that a transient

elevation of chloride using VU treatment from DIV2 to DIV8 resulted in indirect changes in

inhibitory transmission at DIV21 (Peerboom et al., 2023). We therefore assessed if furosemide

treatment from DIV2 to DIV8 affected sIPSCs at DIV21 (Fig. 3C). There was large variability in

sIPSC frequency between cells, but we did not detect any significant difference between

furosemide-treated and control slices (Fig. 3D). We did observe a small increase in sIPSC

amplitude in furosemide-treated slices (Fig. 3E), while the rise and decay time of the sIPSCs

were not different (Fig. 3F, G). In addition, we observed that cell capacitance C_m of CA1

pyramidal cells was decreased in furosemide-treated slices compared to control slices (Fig.

3H). To assess whether the changes in inhibitory transmission were activity-dependent, we

also recorded mIPSCs (Fig. 3I). We observed an increased mIPSCs frequency in furosemide-

treated slices at DIV21, while amplitude, rise and decay time were not different from control

slices (Fig. 3J-M). Interestingly, a difference in membrane capacitance C_m was not detected

when recording mIPSCs in the presence of TTX (Fig. 3N). When we compared C_m values during

sIPSC and mIPSC recording we found a significant interaction (p=0.02 ; 2w ANOVA), suggesting

that furosemide treatment affected cell capacitance in an activity-dependent manner.

A change in mIPSC frequency while amplitude remain unaffected could have resulted from an

increase in the number of inhibitory synapses in furosemide-treated slices compared to

control slices at DIV21. To assess synapse numbers, we used immunohistochemistry to

visualize inhibitory synapses in *stratum Pyramidale (sPyr)* and *stratum Radiatum* (*sRad*) using

the vesicular GABA transporter VGAT. In addition, we stained for CB1 receptors (CB1R) and

parvalbumin (PV), to identify inhibitory synapses from specific interneuron subtypes (Fig.

4A,B). The density and size of VGAT puncta in the CA1 area were similar in control and

furosemide-treated slices at DIV9 (Fig. 4C-F) and we did not observe any differences in the

density of CB1R+ and PV+ VGAT puncta (Fig. 4G-I). At DIV21 however (Fig. 5A,B), we observed

an increased density and reduced size of VGAT puncta in the *sRad* in furosemide-treated slices

compared to controls (Fig. 5E,F), but not *sPyr* (Fig. 5C,D)*.* This suggests that the number of

inhibitory synapses in the dendritic region was increased at DIV21 in furosemide-treated

slices. The density of PV- and CB1R-positive puncta were not different in furosemide-treated

and control slices (Fig. 5G-I), suggesting that the increase in VGAT puncta is due to an increase

in synapses from another interneuron subtype.

Discussion

In this study, we used furosemide to manipulate the GABA shift in organotypic hippocampal

slices. Unexpectedly, we observed that treatment with furosemide from DIV1 to DIV8 had an

opposite effect compared to acute furosemide application. While acute application

depolarized the reversal potential for GABA signaling (E_{GABA}), one week furosemide treatment

resulted in a strong hyperpolarizing shift of E_{GABA}. This shift was not accompanied by a change

in SuperClomeleon (SClm) FRET ratios or chloride co-transporter expression. We observed

that sIPSCs were not affected by the hyperpolarization of E_{GABA} after furosemide treatment at

DIV9, consistent with the notion that GABA signaling in the second postnatal week no longer

promotes synapse formation (Wang and Kriegstein, 2011; Salmon et al., 2020; Peerboom et

al., 2023). However, we found that at DIV21, two weeks after ending furosemide treatment

and when E_{GABA} had normalized, inhibitory currents were increased. This was accompanied

by an increase of the density of VGAT puncta in *sRad*. In addition, we found that cell

capacitance was affected in furosemide-treated slices at DIV21 in an activity-dependent

manner. Together, these results show that accelerating the GABA shift with furosemide does

not directly affect the development of inhibitory synapses. However, transient furosemide

treatment appeared to indirectly alter cell volume regulation and increase inhibitory

transmission two weeks later.

Effect of furosemide on neuronal chloride levels

The depolarizing effect of acute furosemide application at DIV21 is comparable to that of

acute inhibition of KCC2 by specific KCC2 blocker VU0463271 (VU) in slices at the same

developmental stage (Peerboom et al., 2023), suggesting that acutely applied furosemide

merely inhibited KCC2. At the concentration used here (200 µM), furosemide is expected to

block both KCC2 and NKCC1 (Orlov et al., 2015). However, NKCC1 maintains chloride levels

only until approximately P15-21 (Romo-Parra et al., 2008; Sulis Sato et al., 2017; Salmon et

al., 2020). Previous studies reporting a depolarizing effect of furosemide acutely applied the

compound to (slice) cultures at a developmental stage equal to P14 or later (Thompson and

Gahwiler, 1989; Jarolimek et al., 1999; Deeb et al., 2013; Wright et al., 2017). In our slices at

DIV21 (prepared from P7 mice), NKCC1 may no longer directly regulate chloride levels, which

explains why acute furosemide application only acted via KCC2 at DIV21.

In stark contrast, furosemide treatment from DIV1 to DIV8 resulted in hyperpolarizing E_{GABA},

suggestive of NKCC1 block. We checked that furosemide treatment did not affect expression

levels of chloride transporters. If furosemide treatment indeed hyperpolarized E_{GABA} by

blocking NKCC1, we would expect to observe an increase in SClm FRET ratios (reflecting a

decrease in chloride levels) after furosemide treatment, which was not the case. However,

the sensitivity of the SClm sensor is reduced at chloride concentrations below ~5 mM (Grimley

et al., 2013; Boffi et al., 2018; Herstel et al., 2022). This raises the possibility that chloride

levels in our DIV9 slices are too low to detect a further decrease by furosemide treatment

with the SClm sensor.

Possible effect of furosemide on pH

In addition to blocking chloride transporters, furosemide may also interfere with enzymes

that are involved in HCO_3^- regulation (Halligan et al., 1991; Temperini et al., 2009; Uwera et

al., 2015). $GABA_A$ receptors are permeable for chloride and HCO_3^- ions with a permeability of

around 70% and 30% respectively (Kaila et al., 2014). The reversal potential of HCO_3^- (E_{HCO3})

is maintained by pH-regulatory proteins at around -10 mV (Kaila et al., 2014), which is much

more positive than the chloride reversal potential (E_{Cl}). As a result, E_{GABA} is slightly more

positive than E_{Cl} and the flow of chloride through $GABA_A$ receptors is accompanied by an

outflow of HCO_3^- (Kaila et al., 2014). The lack of change in SClm FRET ratios after furosemide

treatment raises the possibility that the observed hyperpolarization of E_{GABA} reflect changes

in intracellular HCO_3^- levels, rather than chloride. If furosemide would decrease intraneuronal

HCO_3^- levels, the outflow of HCO_3^- upon $GABA_A$ receptor activation would be reduced, and

E_{GABA} would become more negative. Previous studies have suggested that furosemide can

inhibit the Cl^-/HCO_3^- anion exchanger AE3 (Halligan et al., 1991; Uwera et al., 2015). AE3

exchanges intracellular HCO_3^- for extracellular chloride, thereby lowering intracellular pH and

increasing intracellular chloride concentrations in response to intracellular alkali loads

(Gonzalez-Islas et al., 2009; Pfeffer et al., 2009; Romero et al., 2013). Furosemide can also

block carbonic anhydrase (CA) (Temperini et al., 2009), which is a cytosolic enzyme that

mediates the conversion of HCO_3^- ($CO_2 + H_2O \rightleftarrows HCO_3^- + H^+$) to buffer intraneuronal pH after

alkali or acid loads (Ruusuvuori and Kaila, 2014). To our knowledge there are no studies

directly examining the contribution of AE3 or CA to E_{GABA}. Theoretically, inhibition of AE3 by

furosemide would decrease intracellular chloride levels while increasing HCO_3^-, which have

opposing effects on E_{GABA}. Inhibition of CA would in theory reduce the contribution of E_{HCO3} to

E_{GABA} and result in a negative shift of E_{GABA} towards E_{Cl}. If the hyperpolarization of E_{GABA} was

solely due to changing HCO_3^- levels without changing chloride, pH values would be

unphysiologically low. A strong decrease in pH would decrease IPSC frequency and amplitude

and increase rise and decay times (Mozrzymas et al., 2003; Dietrich and Morad, 2010), while

we observed that IPSCs were unaffected at DIV9. We therefore deem it unlikely that

furosemide induced large pH changes. However, a small decrease in intracellular pH would

decrease SClm FRET ratios (Grimley et al., 2013; Boffi et al., 2018; Lodovichi et al., 2022),

which may have obscured an increase in FRET ratios due to a decrease in chloride levels. We

therefore conclude that furosemide treatment most likely resulted in a decrease in chloride

levels, but our data do not exclude a subtle change in pH regulation.

Possible effect of furosemide on cell volume regulation

We observed that the membrane capacitance of CA1 pyramidal neurons was reduced in

furosemide-treated slices compared to control slices at DIV21 (Fig. 3H), but not at DIV9 (Fig.

2E). A reduction in whole cell membrane capacitance may reflect a reduction in cell size

(Taylor, 2012). Remarkably, neuronal capacitance was not affected when neuronal activity

was blocked during mIPSC recordings (compare Fig. 3H and 3N), raising the possibility that

the difference in membrane capacitance reflects activity-dependent cell shrinkage. Previous

studies have shown that furosemide inhibits activity-induced changes in the volume of

neuronal tissue during epilepsy (Hochman et al., 1995; Gutschmidt et al., 1999; Hochman,

2012). It has been shown that this effect is mainly mediated by inhibition of NKCC1 on glia

(Ransom et al., 1985; Hochman, 2012). Glia disperse local, activity-induced elevations in

potassium and chloride ions which are followed by water and result in glial swelling (Ransom

et al., 1985; Hochman et al., 1995; Gutschmidt et al., 1999; Hochman, 2012). Our results

suggest that one week treatment with furosemide indirectly affect activity-induced swelling

of neurons, but the mechanism remains unresolved. In interpreting our results and that of

previous studies, it is hard to disentangle the versatile pharmacological profile of furosemide

and the many roles for neuronal chloride in regulating cell volume (Jentsch and Pusch, 2018)

and possibly cellular excitability (Huang et al., 2012; Seja et al., 2012; Jentsch and Pusch, 2018;

Goutierre et al., 2019; Sinha et al., 2022; Peerboom et al., 2023).

Furosemide-treatment indirectly increased inhibitory transmission

We observed that sIPSCs were not affected by the hyperpolarization of E_{GABA} at DIV9, immediately after the furosemide treatment (Fig. 2A-E). These results confirm previous studies reporting that synapse formation is mostly independent from GABA signaling after the first postnatal week (Wang and Kriegstein, 2011; Peerboom et al., 2023). More surprisingly, we found that furosemide treatment resulted in an increase in inhibitory transmission at DIV21. The amplitude of sIPSCs and the frequency of mIPSCs were strongly increased, and this was associated with an increased density of inhibitory synapses in *sRad* in furosemide-treated slices compared to control slices. However, one should be cautious in interpreting synaptic density in the two conditions, as furosemide treatment may have induced overall cell shrinkage, which would increase apparent synaptic density. To assess synaptic densities independently of cell shrinkage, the density of synapses along the dendrites (Feng et al., 2021) or per soma (Bijlsma et al., 2022) should be assessed.

In a recent study we treated slice cultures with VU, a specific blocker of the chloride exporter KCC2, to delay the postnatal GABA shift and maintain a depolarizing E_{GABA} until DIV8. Similar to the current results, we also observed an indirect increase in sIPSC frequency at DIV21 cells (Peerboom et al., 2023). However, the mechanism seems different as mIPSC frequency and synapse numbers were not affected in VU-treated slices (Peerboom et al., 2023). These observations underscore the need for a better understanding of the role of chloride as an intracellular messenger in developing neurons. Although the precise mechanism remains obscure, our current results suggest that furosemide treatment indirectly increases inhibitory transmission, which adds to its previously described anticonvulsant action.

References

Akerman CJ, Cline HT (2006) Depolarizing GABAergic conductances regulate the balance of excitation to inhibition in the developing retinotectal circuit in vivo. J Neurosci 26:5117–5130.

Barbaro NM, Takahashi DK, Baraban SC (2004) A potential role for astrocytes in mediating the antiepileptic actions of furosemide in vitro. Neuroscience 128:655–663.

Bijlsma A, Omrani A, Spoelder M, Verharen JPH, Bauer L, Cornelis C, de Zwart B, van Dorland R, Vanderschuren LJMJ,
Wierenga CJ (2022) Social Play Behavior Is Critical for the Development of Prefrontal Inhibitory Synapses and
Cognitive Flexibility in Rats. J Neurosci 42:8716–8728.

Boffi JC, Knabbe J, Kaiser M, Kuner T (2018) KCC2-dependent steady-state intracellular chloride concentration and ph in
cortical layer 2/3 neurons of anesthetized and awake mice. Front Cell Neurosci 12:1–14.

Casanova E, Fehsenfeld S, Mantamadiotis T, Lemberger T, Greiner E, Stewart AF, Schtz G (2001a) A CamKIIα iCre BAC allows
brain-specific gene inactivation. Genesis 31:37–42.

Casanova E, Fehsenfeld S, Mantamadiotis T, Lemberger T, Greiner E, Stewart AF, Schtz G (2001b) A CamKIIα iCre BAC allows
brain-specific gene inactivation. Genesis 31:37–42.

Chancey JH, Adlaf EW, Sapp MC, Pugh PC, Wadiche JI, Overstreet-Wadiche LS (2013) GABA Depolarization Is Required for
Experience-Dependent Synapse Unsilencing in Adult-Born Neurons. J Neurosci 33:6614–6622.

Chen L, Yu J, Wan L, Wu Z, Wang G, Hu Z, Ren L, Zhou J, Qian B, Zhao X, Zhang J, Liu X (2022) Furosemide prevents
membrane KCC2 downregulation during convulsant stimulation in the hippocampus. IBRO Neurosci Rep 12:355–
365.

Deeb TZ, Nakamura Y, Frost GD, Davies PA, Moss SJ (2013) Disrupted Cl-homeostasis contributes to reductions in the
inhibitory efficacy of diazepam during hyperexcited states. Eur J Neurosci 38:2453–2467.

Dietrich CJ, Morad M (2010) Synaptic acidification enhances GABAA signaling. J Neurosci 30:16044–16052.

Dzhala VI, Talos DM, Sdrulla DA, Brumback AC, Mathews GC, Benke TA, Delpire E, Jensen FE, Staley KJ (2005) NKCC1
transporter facilitates seizures in the developing brain. Nat Med 11:1205–1213.

Feng T, Alicea C, Pham V, Kirk A, Pieraut S (2021) Experience-dependent inhibitory plasticity is mediated by CCK1 basket
cells in the developing dentate gyrus. J Neurosci 41:4607–4619.

Gonzalez-Islas C, Chub N, Wenner P (2009) NKCC1 and AE3 appear to accumulate chloride in embryonic motoneurons. J
Neurophysiol 101:507–518.

Goswami SP, Bucurenciu I, Jonas P (2012) Miniature IPSCs in hippocampal granule cells are triggered by voltage-gated Ca2+
channels via microdomain coupling. J Neurosci 32:14294–14304.

Goutierre M, Al Awabdh S, Donneger F, François E, Gomez-Dominguez D, Irinopoulou T, Menendez de la Prida L, Poncer JC
(2019) KCC2 Regulates Neuronal Excitability and Hippocampal Activity via Interaction with Task-3 Channels. Cell
Rep 28:91–103.

Grimley JS, Li L, Wang W, Wen L, Beese LS, Hellinga HW, Augustine GJ (2013) Visualization of Synaptic Inhibition with an
Optogenetic Sensor Developed by Cell-Free Protein Engineering Automation. J Neurosci 33:16297–16309.

Gulyás AI, Sík A, Payne JA, Kaila K, Freund TF (2001) The KCl cotransporter, KCC2, is highly expressed in the vicinity of
excitatory synapses in the rat hippocampus. Eur J Neurosci 13:2205–2217.

Gutschmidt KU, Stenkamp K, Buchheim K, Heinemann U, Meierkord H (1999) Anticonvulsant actions of furosemide in vitro.
Neuroscience 91:1471–1481.

Halligan RD, Shelat H, Kahn AM (1991) Na+-independent Cl--HCO : exchange in sarcolemmal vesicles from vascular smooth
muscle. Am J Physiol 260:C347-54.

Herstel LJ, Peerboom C, Uijtewaal S, Selemangel D, Karst H, Wierenga CJ (2022) Using SuperClomeleon to measure changes
in intracellular chloride during development and after early life stress. Eneuro November/D.

Hesdorffer DC, Hauser WA, Annegers JF, Rocca WA (1996) Severe, uncontrolled hypertension and adult-onset seizures: A
case- control study in Rochester, Minnesota. Epilepsia 37:736–741.

Hochman DW (2012) The extracellular space and epileptic activity in the adult brain: Explaining the antiepileptic effects of
furosemide and bumetanide. Epilepsia 53:18–25.

Hochman DW, Baraban SC, Owens JWM, Schwartzkroin PA (1995) Dissociation of synchronization and excitability in furosemide blockade of epileptiform activity. Science 270:99–102.

Holtkamp M, Matzen J, Buchheim K, Walker MC, Meierkord H (2003) Furosemide terminates limbic status epilepticus in freely moving rats. Epilepsia 44:1141–1144.

Hu HY, Kruijssen DLH, Frias CP, Rózsa B, Hoogenraad CC, Wierenga CJ (2019) Endocannabinoid Signaling Mediates Local Dendritic Coordination between Excitatory and Inhibitory Synapses. Cell Rep 27:666-675.e5.

Huang WC, Xiao S, Huang F, Harfe BD, Jan YN, Jan LY (2012) Calcium-Activated Chloride Channels (CaCCs) Regulate Action Potential and Synaptic Response in Hippocampal Neurons. Neuron 74:179–192.

Hübner C a, Stein V, Hermans-Borgmeyer I, Meyer T, Ballanyi K, Jentsch TJ (2001) Disruption of KCC2 reveals an essential role of K-Cl cotransport already in early synaptic inhibition. Neuron 30:515–524.

Jarolimek W, Lewen A, Misgeld U (1999) A furosemide-sensitive K+ -Cl- cotransporter counteracts intracellular Cl- accumulation and depletion in cultured rat midbrain neurons. J Neurosci 19:4695–4704.

Jentsch TJ, Pusch M (2018) CLC chloride channels and transporters: Structure, function, physiology, and disease. Physiol Rev 98:1493–1590.

Kaila K, Price TJ, Payne JA, Puskarjov M, Voipio J (2014) Cation-chloride cotransporters in neuronal development, plasticity and disease. Nat Rev Neurosci 15:637–654.

Kirmse K, Kummer M, Kovalchuk Y, Witte OW, Garaschuk O, Holthoff K (2015) GABA depolarizes immature neurons and inhibits network activity in the neonatal neocortex in vivo. Nat Commun 6:1–13.

Korpi ER, Kuner T, Seeburg PH, Luddens H (1995) Selective antagonist for the cerebellar granule cell-specific g-aminobutyric acid type A receptor. MolPharmacol 47:283–289.

Leinekugel X, Medina I, Khalilov I, Ben Ari Y, Khazipov R (1997) Ca 2+ Oscillations Mediated by the Synergistic Excitatory Actions of GABA. Neuron 18:243–255.

Lodovichi C, Ratto GM, Trevelyan AJ, Arosio D (2022) Genetically encoded sensors for Chloride concentration. J Neurosci Methods 368:109455.

Maa EH, Kahle KT, Walcott BP, Spitz MC, Staley KJ (2011) Diuretics and epilepsy: Will the past and present meet? Epilepsia 52:1559–1569.

Mozrzymas J, Zarnowska E, Pytel M, Mercik K (2003) Modulation of GABAA receptors by hydrogen ions reveals synaptic GABA transient and a crucial role of the desensitization process. J Neurosci 23:7981–7992.

Murata Y, Colonnese MT (2020) GABAergic interneurons excite neonatal hippocampus in vivo. Sci Adv 6:1–10.

Oh WC, Lutzu S, Castillo PE, Kwon HB (2016) De novo synaptogenesis induced by GABA in the developing mouse cortex - supplemental. Science 353:1037–1040.

Orlov SN, Koltsova S V, Kapilevich L V, Gusakova S V, Dulin NO (2015) NKCC1 and NKCC2: The pathogenetic role of cation-chloride cotransporters in hypertension. 2:186–196.

Otsu Y, Donneger F, Schwartz EJ, Poncer JC (2020) Cation–chloride cotransporters and the polarity of GABA signalling in mouse hippocampal parvalbumin interneurons. J Physiol 598:1–16.

Pearce RA (1993) Physiological evidence for two distinct GABAA responses in rat hippocampus. Neuron 10:189–200.

Peerboom C, Kater S De, Jonker N, Rieter M, Wijne T, Wierenga CJ (2023) Delaying the GABA shift indirectly affects membrane properties in the developing hippocampus. bioRxiv.

Pfeffer CK, Stein V, Keating DJ, Maier H, Rinke I, Rudhard Y, Hentschke M, Rune GM, Jentsch TJ, Hubner CA (2009) NKCC1-Dependent GABAergic Excitation Drives Synaptic Network Maturation during Early Hippocampal Development. J Neurosci 29:3419–3430.

Rahmati N, Normoyle KP, Glykys J, Dzhala VI, Lillis KP, Kahle KT, Raiyyani R, Jacob T, Staley KJ (2021) Unique actions of gaba
arising from cytoplasmic chloride microdomains. J Neurosci 41:4967–4975.

Ransom B, Yamate C, Connors B (1985) Activity-dependent in rat optic nerve: a developmental study. J Neurosci 5:532–
535.

Rivera C, Voipio J, Payne J a, Ruusuvuori E, Lahtinen H, Lamsa K, Pirvola U, Saarma M, Kaila K (1999) The K+/Cl- co-
transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. Nature 397:251–255.

Romero MF, Chen AP, Parker MD, Boron WF (2013) The SLC4 family of bicarbonate (HCO3-) transporters. Mol Aspects Med
34:159–182.

Romo-Parra H, Treviño M, Heinemann U, Gutiérrez R (2008) GABA actions in hippocampal area CA3 during postnatal
development: Differential shift from depolarizing to hyperpolarizing in somatic and dendritic compartments. J
Neurophysiol 99:1523–1534.

Ruiter M, Herstel LJ, Wierenga CJ, Ma T (2020) Reduction of Dendritic Inhibition in CA1 Pyramidal Neurons in Amyloidosis
Models of Early Alzheimer's Disease. J Alzheimers Dis 78:951–964.

Ruusuvuori E, Kaila K (2014) Carbonic Anhydrases and Brain pH in the Control of Neuronal Excitability. In: Carbonic
Anhydrase: Mechanism, Regulation, Links to Disease, and Industrial Applications, pp 271–290 Subcellular
Biochemistry. Springer International Publishing. Available at: https://doi.org/10.1007/978-94-007-7359-2_14.

Salmon CK, Pribiag H, Gizowski C, Farmer WT, Cameron S, Jones E V., Mahadevan V, Bourque CW, Stellwagen D, Woodin
MA, Murai KK (2020) Depolarizing GABA Transmission Restrains Activity-Dependent Glutamatergic Synapse
Formation in the Developing Hippocampal Circuit. Front Cell Neurosci 14:1–16.

Sedmak G, Puskarjov NJ-MM, Ulamec M, Krušlin B, Kaila K, Judaš M (2016) Developmental Expression Patterns of KCC2 and
Functionally Associated Molecules in the Human Brain. Cereb Cortex 27:4060–4072.

Seja P, Schonewille M, Spitzmaul G, Badura A, Klein I, Rudhard Y, Wisden W, Hübner CA, De Zeeuw CI, Jentsch TJ (2012)
Raising cytosolic Cl - in cerebellar granule cells affects their excitability and vestibulo-ocular learning. EMBO J
31:1217–1230.

Sinha AS, Wang T, Watanabe M, Hosoi Y, Sohara E, Akita T, Uchida S, Fukuda A (2022) WNK3 kinase maintains neuronal
excitability by reducing inwardly rectifying K+ conductance in layer V pyramidal neurons of mouse medial
prefrontal cortex. Front Mol Neurosci 15.

Staley KJ (2002) Diuretics as Antiepileptic Drugs: Should We Go with the Flow? Epilepsy Curr 2:39–40.

Stein V, Hermans-Borgmeyer I, Jentsch TJ, Hübner CA (2004) Expression of the KCl Cotransporter KCC2 Parallels Neuronal
Maturation and the Emergence of Low Intracellular Chloride. J Comp Neurol 468:57–64.

Stoppini L, Buchs PA, Muller D (1991) A simple method for organotypic cultures of nervous tissue. J Neurosci Methods
37:173–182.

Sulis Sato S, Artoni P, Landi S, Cozzolino O, Parra R, Pracucci E, Trovato F, Szczurkowska J, Luin S, Arosio D, Beltram F,
Cancedda L, Kaila K, Ratto GM (2017) Simultaneous two-photon imaging of intracellular chloride concentration
and pH in mouse pyramidal neurons in vivo. Proc Natl Acad Sci 114:E8770–E8779.

Taylor AL (2012) What we talk about when we talk about capacitance measured with the voltage-clamp step method. J
Comput Neurosci 32:167–175.

Temperini C, Cecchi A, Scozzafava A, Supuran CT (2009) Carbonic anhydrase inhibitors. Comparison of chlorthalidone,
indapamide, trichloromethiazide, and furosemide X-ray crystal structures in adducts with isozyme II, when
several water molecules make the difference. Bioorg Med Chem 17:1214–1221.

Thompson SM, Gahwiler BH (1989) Activity-dependent disinhibition. II. Effects of extracellular potassium, furosemide, and
membrane potential on E(Cl)- in hippocampal CA3 neurons. J Neurophysiol 61:512–523.

Tsien JZ, Chen DF, Gerber D, Tom C, Mercer EH, Anderson DJ, Mayford M, Kandel ER, Tonegawa S (1996) Subregion- and
cell type-restricted gene knockout in mouse brain. Cell 87:1317–1326.

Tsukahara T, Masuhara M, Iwai H, Sonomura T, Sato T (2015) Repeated stress-induced expression pattern alterations of the
hippocampal chloride transporters KCC2 and NKCC1 associated with behavioral abnormalities in female mice.
Biochem Biophys Res Commun 465:145–151.

Tyzio R, Holmes GL, Ben-Ari Y, Khazipov R (2007) Timing of the developmental switch in GABAA mediated signaling from
excitation to inhibition in CA3 rat hippocampus using gramicidin perforated patch and extracellular recordings.
Epilepsia 48:96–105.

Uwera J, Nedergaard S, Andreasen M (2015) A novel mechanism for the anticonvulsant effect of furosemide in rat
hippocampus in vitro. Brain Res 1625:1–8.

van Rheede JJ, Richards BA, Akerman CJ (2015) Sensory-Evoked Spiking Behavior Emerges via an Experience-Dependent
Plasticity Mechanism. Neuron 87:1050–1062.

Viitanen T, Ruusuvuori E, Kaila K, Voipio J (2010) The K+-Cl- cotransporter KCC2 promotes GABAergic excitation in the
mature rat hippocampus. J Physiol 588:1527–1540.

Wang BS, Feng L, Liu M, Liu X, Cang J (2013) Environmental Enrichment Rescues Binocular Matching of Orientation
Preference in Mice that Have a Precocious Critical Period. Neuron 80:198–209.

Wang DD, Kriegstein AR (2008) GABA Regulates Excitatory Synapse Formation in the Neocortex via NMDA Receptor
Activation. J Neurosci 28:5547–5558.

Wang DD, Kriegstein AR (2011) Blocking early GABA depolarization with bumetanide results in permanent alterations in
cortical circuits and sensorimotor gating deficits. Cereb Cortex 21:574–587.

Wierenga CJ, Müllner FE, Rinke I, Keck T, Stein V, Bonhoeffer T (2010) Molecular and electrophysiological characterization
of GFP-expressing ca1 interneurons in GAD65-GFP mice. PLoS ONE 5:1–11.

Wright R, Newey SE, Ilie A, Wefelmeyer W, Raimondo J V., Ginham R, Mcllhinney RAJ, Akerman CJ (2017) Neuronal
Chloride Regulation via KCC2 Is Modulated through a GABA $_B$ Receptor Protein Complex. J Neurosci 37:5447–
5462.

Yamada J, Okabe A, Toyoda H, Kilb W, Luhmann H, Fukuda A (2004) Cl– uptake promoting depolarizing GABA actions in
immature rat neocortical neurones is mediated by NKCC1. J Physiol 557:829–841.

Figure legends

Figure 1. *Furosemide acutely depolarizes the GABAergic driving force, but results in hyperpolarization after*
one week treatment.

A, B) GABA reversal potential (E_{GABA}) (PT, p=0.33) and GABA Driving Force (GABA DF) (PT, p=0.19) values recorded
in CA1 pyramidal cells at DIV2 before and after acute application of furosemide. Data from 4 cells, 4 slices and 4
mice.

C, D) E_{GABA} (MW, p=0.004) and GABA DF (MW, p=0.002) values recorded in CA1 pyramidal cells at DIV21 before
and after acute application of furosemide. Data from 6 cells, 5 slices and 5 mice.

E, F) E_{GABA} (MW, p=0.001) and GABA DF (MW, p=0.005) recorded in CA1 pyramidal cells at DIV9 in control (CTL)
and furosemide-treated slices. Data from 7-9 cells, 5-8 slices and 4-5 mice per group.

G) Average SClm FRET ratios in CTL and furosemide-treated slice cultures (UT, p=0.75). Data from 10-14 slices
and 6-10 mice per group.

H) Cumulative distribution of FRET ratios in individual cells in CTL and furosemide -treated cultures. (KS, p=0.23).
Data from 15 randomly selected cells per slice.

I) Western blots of NKCC1, KCC2 and S940-pKCC2 (pKCC2) protein levels in CTL and furosemide-treated slice
cultures. Tubulin (Tub) was used as loading control.

J) Summary of data for NKCC1, KCC2 and for s940-pKCC2 (pKCC2) protein levels. Values were normalized to the
protein level in CTL cultures (NKCC1: MW, p>0.99; KCC2: MW, p=0.70; pKCC2: MW, p=0.48). Data from 3-5
experiments and 3 mice per group.

K) Confocal images of NeuN and KCC2 staining in CTL and furosemide-treated cultures. Yellow lines indicate the
outlines of individual cells. Scale bar: 10 µm.

L) Total KCC2 levels (UT, p=0.55) and KCC2 levels in membrane (UT, p=0.37) in CTL and furosemide-treated
cultures. Each datapoint represents the mean KCC2 intensity of all neurons in one image. Data from 8-17 images,
4-8 slices and 3 mice per group.

Figure 2. *Furosemide does not affect inhibitory transmission at DIV9.*
A-E) sIPSC frequency (MW, p=0.084), amplitude (UT, p=0.83), risetime (MW, p=0.80) and decay tau (MW,
p=0.57) and membrane capacitance (Cm) (UT, p=0.63) in control (CTL) and furosemide-treated slice cultures at
DIV9. Data from 14-17 cells, 9 slices and 9 mice per group.

F-I) sIPSC frequency after ACSF wash in as a control CTL (WSR, p=0.008) and furosemide-treated (WSR, p=0.002)
slice cultures at DIV9. sIPSC frequency after TTX wash in of in CTL (WSR, p=0.80) and furosemide-treated WSR,
p=0.002) slice cultures at DIV9. Data from 8-15 cells, 8-15 slices and 7-10 mice per group.

J) Change in sIPSC frequency after ACSF wash in as a control, versus after wash in of TTX, K252a, AM251 and
agatoxin-IVA (aga) to block neuronal activity, BDNF, endocannabinoids and P/Q calcium channels respectively in
control and furosemide-treated slices at DIV9 (2W ANOVA, treatment p=0.18, blocker p=0.002; SMC, control
versus TTX, p=0.95; control versus K252a, p=0.020; control versus AM251, p=0.99; control versus agatoxin-IVA
(aga), p=0.82). Data from 7-15 cells, 7-15 slices and 4-10 mice per group.

Figure 3. *Furosemide increases inhibitory transmission at DIV21.*

A,B) GABA reversal potential (E_{GABA}) (MW, p=0.56) and GABA Driving Force (GABA DF) (MW, p=0.76) in control

(CTL) and furosemide-treated slice cultures at DIV21. Data from 9-13 cells, 9 slices and 9 mice per group.

C) sIPSC recording from CA1 pyramidal cells in CTL and furosemide-treated organotypic cultures at DIV21. sIPSCs

are indicated with *.

D-G) sIPSC frequency (UT, p=0.10), amplitude (UT, p=0.034), risetime (UT, p=0.54) and decay tau (UT, p=0.34) in

CTL and furosemide-treated slice cultures at DIV21.

I) Membrane capacitance (Cm) of CA1 pyramidal cells in CTL and furosemide-treated slice cultures at DIV21 (UT,

p=0.035).

I) mIPSC recording from CA1 pyramidal cells in control and furosemide-treated organotypic cultures at DIV21.

mIPSCs are indicated with *.

J-M) mIPSC frequency (UT, p=0.0002), amplitude (UT, p=0.34), risetime (MW, p=0.73) and decay tau (UT, p=0.06)

in CTL and furosemide-treated organotypic cultures at DIV21.

N) Membrane capacitance (Cm) of CA1 pyramidal cells in the presence of TTX in CTL and furosemide-treated

slice cultures at DIV21 (UT, p=0.35). Data in D-H from 11-12 cells, 5-6 slices and 3-5 mice per group; data in J-N

from 12-13 cells, 5 slices and 4 mice per group.

Figure 4. *Furosemide does not affect inhibitory synapses at DIV9.*

A,B) VGAT, PV and CB1R immunofluorescence in the CA1 *stratum Pyramidale* (*sPyr)* and *stratum Radiatum*

(*sRad)* of control (CTL) and furosemide-treated organotypic cultures at DIV9. Scale bar=10 µm.

C,D) The density (UT, p=0.50) and size (UT, p=0.47) of VGAT puncta in the *sPyr* of CTL and furosemide-treated

organotypic cultures at DIV9.

E,F) The density (UT, p=0.59) and size (UT, p=0.46) of VGAT puncta in *sRad* of CTL and furosemide-treated

organotypic cultures at DIV9.

G,H) The density of CB1R-positive VGAT puncta (UT, p=0.43) and PV positive VGAT puncta (UT, p=0.82) in *sPyr*

of CTL and furosemide-treated organotypic cultures at DIV9.

I) The density of CB1R-positive VGAT puncta (UT, p=0.68) in *sRad* of CTL and furosemide-treated organotypic

cultures at DIV9.

Data in C-I from 11-12 images, 6 slices and 2 mice per group.

Figure 5. *Furosemide increases the number of VGAT puncta at DIV21 in sRad.*

A,B) VGAT, PV and CB1R immunofluorescence in the CA1 *stratum Pyramidale* (*sPyr)* and *stratum Radiatum*

(*sRad)* of control (CTL) and furosemide-treated organotypic cultures at DIV21. Scale bar=10 µm.

C,D) The density (UT, p=0.09) and size (UT, p=0.14) of VGAT puncta in *sPyr* of CTL and furosemide-treated

organotypic cultures at DIV21. Data from 16 images, 8 slices and 3 mice per group.

E,F) The density (MW, p=0.019) and size (MW, p=0.015) of VGAT puncta in *SRad* of CTL and furosemide-treated

organotypic cultures at DIV21. Data from 13 images, 7 slices and 3 mice per group.

G,H) The density of CB1R (UT, p=0.11) and PV positive VGAT puncta (UT, p=0.89) in *sPyr* of CTL and furosemide-

treated organotypic cultures at DIV21. Data from 16 images, 8 slices and 3 mice per group.

I) The density of CB1R-positive VGAT puncta (MW, p=0.30) in *SRad* of CTL and furosemide-treated organotypic

cultures at DIV21. Data from 13 images, 7 slices and 3 mice per group.

Peerboom et al. - Figure 1

Peerboom et al. - Figure 2

Peerboom et al. - Figure 3

Peerboom et al. - Figure 4

Peerboom et al. - Figure 5