High quality genome assembly and annotation (v1) of the eukaryotic terrestrial microalga *Coccomyxa viridis* SAG 216-4

Anton Kraege[1], Edgar A. Chavarro-Carrero[1], Nadège Guiglielmoni[2], Eva Schnell[1], Joseph
Kirangwa[2], Stefanie Heilmann-Heimbach[3,4], Kerstin Becker[5,6], Karl Köhrer[6], WGGC Team,
DeRGA Community, Philipp Schiffer[2], Bart P. H. J. Thomma[1,7] and Hanna Rovenich[1]

[1] Institute for Plant Sciences, University of Cologne, Germany; [2] Institute of Zoology, University
of Cologne, Germany; [3] Institute of Human Genetics, University Hospital of Bonn & University
of Bonn, Bonn, Germany; [4] NGS Core Facility, Medical Faculty of the University of Bonn, Bonn,
Germany; [5] Cologne Center for Genomics (CCG), Medical Faculty, University of Cologne,
50931 Cologne, Germany; [6] Biological and Medical Research Centre (BMFZ), Genomics &
Transcriptomics Laboratory, Heinrich-Heine-University Düsseldorf, Germany; [7] Cluster of
Excellence on Plant Sciences (CEPLAS), Germany

corresponding author: hanna.roevenich@uni-koeln.de

Abstract

Unicellular green algae of the genus *Coccomyxa* are recognized for their worldwide distribution and ecological versatility. Most species described to date live in close association with various host species, such as in lichen associations. However, little is known about the molecular mechanisms that drive such symbiotic lifestyles. We generated a high-quality genome assembly for the lichen photobiont *Coccomyxa viridis* SAG 216-4 (formerly *C. mucigena*). Using long-read PacBio HiFi and Oxford Nanopore Technologies in combination with chromatin conformation capture (Hi-C) sequencing, we assembled the genome into 21 scaffolds with a total length of 50.9 Mb, an N50 of 2.7 Mb and a BUSCO score of 98.6%. While 19 scaffolds represent full-length nuclear chromosomes, two additional scaffolds represent the mitochondrial and plastid genomes. Transcriptome-guided gene annotation resulted in the identification of 13,557 protein-coding genes, of which 68% have annotated PFAM domains and 962 are predicted to be secreted.

Keywords: *Coccomyxa viridis*, EBP, ERGA, long-read sequencing, genome assembly, genome annotation, Trebouxiophyceae

Introduction

Green algae are photosynthesizing eukaryotic organisms that differ greatly in terms of morphology and colonize a large variety of aquatic and terrestrial habitats. Phylogenetically, green algae form a paraphyletic group that has recently been proposed to comprise three lineages including the Prasinodermophyta in addition to the Chlorophyta and Streptophyta (Li et al., 2020). This new phylum diverged before the split of the Chlorophyta and Streptophyta that occurred between 1,000 and 700 million years ago (Morris et al., 2018). While the streptophyte lineage encompasses charophyte green algae as well as land plants, the chlorophyte lineage consists of 7 prasinophyte classes, which gave rise to 4 phycoplast-containing core chlorophyte classes (Chlorodendrophyceae, Trebouxiophyceae, Ulvophyceae, Chlorophyceae) with one independent sister class (Pedinophyceae) (Leliaert et al., 2012; Marin, 2012).

The *Coccomyxa* genus is represented by coccoid unicellular green algae that belong to the class of Trebouxiophyceae. Morphologically, *Coccomyxa* spp. are characterized by irregular elliptical to globular cells that range from 6–14 x 3–6 μm in size, with a single parietal chloroplast lacking pyrenoids and the absence of flagellate stages (Schmidle, 1901). Members of this genus are found in freshwater, marine, and various terrestrial habitats where they occur free-living or in symbioses with diverse hosts (Darienko et al., 2015; Gustavs et al., 2017; Malavasi et al., 2016). Several *Coccomyxa* species establish stable, mutualistic associations with fungi that result in the formation of complex three-dimensional architectures, known as lichens (Faluaburu et al., 2019; Gustavs et al., 2017; Jaag, 1933; Yahr et al., 2015; Zoller and Lutzoni, 2003). Others associate with vascular plants or lichens as endo- or epiphytes, respectively (Cao et al., 2018a; Cao et al., 2018b; Tagirdzhanova et al., 2023; Trémouillaux-Guiller et al., 2002), and frequently occur on the bark of trees (Kulichovà et al., 2014; Štifterovà and Neustupa, 2015) where they may interact with other microbes. One novel species was recently found in association with carnivorous plants, even though the nature of this relationship remains unclear (Sciuto et al., 2019). Besides, *Coccomyxa* also establishes parasitic interactions with different mollusk species affecting their filtration ability and reproduction (Gray et al., 1999; Sokolnikova et al., 2016; Sokolnikova et al., 2022; Vaschenko et al., 2013).

Despite this ecological versatility, little is known about the molecular mechanisms that determine the various symbiotic lifestyles in *Coccomyxa*. One short read-based genome is available for *C. subellipsoidea* C-169 that was isolated on Antarctica where it occurred on dried algal peat (Blanc et al., 2012), whereas another high-quality genome has recently been made available for a non-symbiotic strain of *C. viridis* that was isolated from a lichen thallus (Tagirdzhanova et al., 2023). For *Coccomyxa* sp. Obi, LA000219 and SUA001 chromosome-

, scaffold- and contig-level assemblies are available on NCBI, respectively, as well as two
metagenome-assembled genomes of *C. subellipsoidea*. To facilitate the study of *Coccomyxa*
symbiont-associated traits and their evolutionary origin, we here present the generation of a
high-quality chromosome-scale assembly of the phycobiont *C. mucigena* SAG 216-4 using
long-read PacBio HiFi and Oxford Nanopore Technology (ONT) combined with Hi-C and RNA
sequencing. Recent SSU and ITS rDNA sequencing-based re-evaluations of the *Coccomyxa*
phylogeny placed the SAG 216-4 isolate in the clade of *C. viridis* (Darienko et al., 2015;
Malavasi et al., 2016). Hence, this isolate will be referred to as *C. viridis* here and data have
been deposited under the corresponding Taxonomy ID.
83
Materials & Methods
Sample information
Coccomyxa viridis (formerly *Coccomyxa mucigena*) SAG 216-4 was ordered from the Culture
Collection of Algae at the Georg-August-University Göttingen (*Sammlung von Algenkulturen*
der Universität Göttingen, international acronym SAG), Germany. The stock culture was
reactivated in liquid modified Waris-H growth medium (McFadden and Melkonian, 1986) with
soil extract and 3x vitamins (0.15 nM vitamin B12, 4.1 nM biotin, 0.3 µM thiamine-HCl, 0.8 nM
niacinamide), and maintained through regular medium replacement. Cultures were grown at
~ 15 µmol photons m^{-2} s^{-1} (fluorescent light tubes: L36W/640i energy saver cool white and
L58W/956 BioLux, Osram, Munich, Germany) in a 14/10 h light/dark cycle at 20°C.
94
DNA and *RNA extraction*
Cells of a 7-week-old *C. viridis* culture were harvested over 0.8 µm cellulose nitrate filters
(Sartorius, Göttingen, Germany) using a vacuum pump. Material was collected with a spatula,
snap-frozen and ground in liquid nitrogen using mortar and pestle. The ground material was
used for genomic DNA extraction with the RSC Plant DNA Kit (Promega, Madison, WI, USA)
using the Maxwell® RSC device according to manufacturer's instructions. To prevent shearing
of long DNA fragments, centrifugation was carried out at 10,000 *g* during sample preparation.
Following DNA extraction, DNA fragments <10,000 bp were removed using the SRE XS kit
(Circulomics, Baltimore, MD, USA) according to manufacturer's instructions. DNA quantity and
quality were assessed using the Nanodrop 2000 spectrometer and Qubit 4 fluorometer with
the dsDNA BR assay kit (Invitrogen, Carlsbad, CA, USA), and integrity was confirmed by gel
electrophoresis. High-molecular weight DNA was stored at 4°C.
For total RNA extraction, algal cells were collected from a dense nine-day-old culture
and ground in liquid nitrogen using mortar and pestle. RNA was extracted with the Maxwell
RSC Plant RNA kit (Promega, Madison, WI, USA) using the Maxwell® RSC device according

to manufacturer's instructions. RNA quality and quantity was determined using the Nanodrop

2000 and stored at -80°C.

Pacific Biosciences High-Fidelity (PacBio HiFi) sequencing

114   HiFi libraries were prepared with the Express 2.0 Template kit (Pacific Biosciences, Menlo

115   Park, CA, USA) and sequenced on a Sequel II/Sequel IIe instrument with 30h movie time. HiFi

116   reads were generated using SMRT Link (v10; (Pacific Biosciences, Menlo Park, CA, USA)

117   with default parameters.

Oxford Nanopore Technologies (ONT) sequencing

Library preparation with the Rapid Sequencing Kit (SQK-626 RAD004) was performed with

~400 ng HMW DNA according to manufacturer's instructions (Oxford Nanopore Technologies,

Oxford, UK). The sample was loaded onto an R9.4.1 flow cell in a miniION Mk1B device

(Oxford Nanopore Technologies, Oxford, UK), which was run for 24 h. Subsequent base

calling was performed using Guppy (version 630 3.1.3; Oxford Nanopore Technologies,

Oxford, UK). Adapter sequences were removed using Porechop (version 0.2.4 with default

settings) (Wick, 2018), and the reads were self-corrected and trimmed using Canu (version

1.8) (Koren et al., 2017).

Chromosome conformation capture (Hi-C) and sequencing

C. viridis cells were cross-linked in 3% formaldehyde for 1 hour at room temperature. The

reaction was quenched with glycine at a final concentration of 250 mM. Cells were collected

by centrifugation at 16,000 g for 10 min. Pellets were flash-frozen in liquid nitrogen and ground

using mortar and pestle. Hi-C libraries were prepared using the Arima-HiC+ kit (Arima

Genomics, Carlsbad, CA, USA) according to manufacturer's instructions, and subsequently

paired-end (2x150 bp) sequenced on a NovaSeq 6000 instrument (Illumina, San Diego, CA,

USA).

RNA sequencing

139   Library preparation for full-length mRNASeq was performed using the NEB Ultra II Directional

140   RNA Library Prep with NEBNext Poly(A) mRNA Magenetic Isolation Module and 500 ng total

141   RNA as starting material, except for W-RNA Lplaty, where library prep was based on 100 ng

142   total RNA as starting material. Sequencing was performed on an Illumina NovaSeq 6000

143   device with 2x150 bp paired-end sequencing protocol and >50 M reads per sample.

Genome assembly

PacBio HiFi reads were assembled using Raven (v1.8.1) (Vaser and Šikić, 2021) with default
settings. Hi-C reads were mapped onto this assembly with Juicer (v2.0) using the "assembly"
option to skip the post-processing steps and generate the merged_nodups.txt file (Durand et
al., 2016b). For the juicer pipeline, restriction site maps were generated using the *Dpn*II
(GATC) and *Hinf*I (GANTC) restriction site profile and the assembly was indexed with BWA
index (v0.7.17-r1188) (Li and Durbin, 2009), and used to polish the assembly using 3d-dna
(v180922) (Dudchenko et al., 2017). Afterwards, Juicebox (v1.11.08) was used to manually
curate the genome assembly by splitting contigs and rearranging them according to the Hi-C
pattern (Durand et al., 2016a). Contigs were merged to scaffolds according to the Hi-C map
and Ns were introduced between contigs within scaffolds, gaps between contigs were
removed and contigs were merged. Subsequently, ONT reads were mapped to the assembly
using Minimap2 (v2.24-r1122) and Samtools (v1.10) and mapped reads were visualized in
Integrative Genome Viewer (v2.11.2) (Danecek et al., 2021; Li, 2021; Robinson et al., 2011).
Whenever gaps between contigs were spanned by at least five reads with a mapping quality
of 30, the contigs were fused in the assembly.

Potential telomeres were identified using tapestry (v1.0.0) with "AACCCT" as telomere
sequence (Davey et al., 2020). To check for potential contaminations, Blobtools (v1.1.1) and
BLAST (v2.13.0+) were used to create a Blobplot including taxonomic annotation at genus
level (Camacho et al., 2009; Laetsch and Blaxter, 2017). To check completeness of the
assembly and retrieve ploidy information, kat comp from the Kmer Analysis Toolkit (v2.4.2)
was used, and results were visualized using the kat plot spectra-cn function with the -x 800
option to extend the x-axis (Mapleson et al., 2016). Genome synteny to the closest sequenced
relative *C. subellipsoidea* C-169 was determined using Mummer3 (Blanc et al., 2012; Kurtz et
al., 2004). In detail, the two assemblies were first aligned using Nucmer, followed by a filtering
step with Delta-filter using the many-to-many option (-m). Finally, the alignment was visualized
with Mummerplot.

Annotation

To annotate repetitive elements in the nuclear genome, a database of simple repeats was
created with RepeatModeler (v2.0.3) that was expanded with transposable elements (TE) from
the TransposonUltimate resonaTE (v1.0) pipeline (Flynn et al., 2020; Riehl et al., 2022). This
pipeline uses multiple tools for TE prediction and combines the prediction output. For the
prediction of TEs in *Coccomyxa viridis* helitronScanner, ltrHarvest, mitefind, mitetracker,
RepeatModeler, RepeatMasker, sinefind, tirvish, transposonPSI and NCBICDD1000 were
used within TransposonUltimate resonaTE and TEs that were predicted by at least two tools
were added to the database. TEclass (v2.1.3) was used for classification (Abrusán et al.,
2009). To softmask the genome and obtain statistics on the total TE and repetitive element

content in the genome, RepeatMasker (v4.1.2-p1)(Smit et al., 2012) was used with excln

option to exclude Ns in the masking.

Gene annotation in the nuclear genome was performed making use of RNA

sequencing data. To this end, the genome was indexed, and reads were mapped with HiSat2

(v2.2.1) using default settings (Kim et al., 2019). Afterwards, BRAKER1 (v2.1.6) was used for

transcriptome-guided gene prediction based on the RNA sequencing data with default settings

(Hoff et al., 2016). To generate protein and coding sequence files the Braker output was

transformed with Gffread (v0.12.7) (Pertea and Pertea, 2020). PFAM domain annotation was

performed with InterProScan (v5.61) (Paysan-Lafosse et al., 2023). To estimate the number

of secreted proteins, SignalP (v6.0) was run in the slow-sequential mode on the annotated

proteins (Teufel et al., 2022). Finally, BUSCO (v5.3.2) was run with the Chlorophyta database

(chlorophyta_odb10) to estimate the completeness of the gene annotation (Manni et al.,

2021). The circos plot visualization of the annotation was created with R (v4.2.0) and Circlize

(v0.4.14) (Gu et al., 2014). All software and tools used for the genome assembly and

annotation are summarized in Table S1.

Organelle genomes were annotated separately. Scaffolds were identified as organelle

genomes based on their lower GC content and smaller size. The mitochondrial genome was

annotated using MFannot (Lang et al., 2023) as well as GeSeq (Tillich et al., 2017) and the

annotation was combined within the GeSeq platform. The plastid genome was annotated

using GeSeq alone. The annotations were visualized using the OGDraw webserver (Greiner

et al., 2019).

206 Results

The version 1 genome of *C. viridis* was assembled from 32.2 Gbp of PacBio HiFi reads with

a mean read length of 15 kb, 0.95 Gbp Nanopore reads with a mean read length of 8.8 kb and

15 million pairs of Hi-C seq data. The PacBio HiFi reads were first assembled using Raven

(Vaser and Šikić, 2021), yielding 27 contigs. These contigs were scaffolded and manually

curated using Hi-C data (Dudchenko et al., 2017; Durand et al., 2016a; Durand et al., 2016b;

Li and Durbin, 2009). To close the remaining gaps between contigs within scaffolds, ONT

reads were mapped onto the assembly (Danecek et al., 2021; Li, 2021) and gaps that were

spanned by at least 5 ONT reads with a mapping quality >30 were manually closed, finally

resulting in 21 scaffolds consisting of 26 contigs with a total length of 50.9 Mb and an N50 of

2.7 Mb (Figure 1, Table 1). Using Tapestry (Davey et al., 2020), telomeric regions

([AACCCT]*n*) were identified at both ends of nine of the 21 scaffolds (≥5 repeats) (Figure 1a),

suggesting that these represent full-length chromosomes, which was confirmed by Hi-C

analysis (Figure 1b). Additionally, the Hi-C contact map indicated centromeres for some of the chromosomes. However, the determination of exact centromere locations on all chromosomes will require ChIP-seq analysis and CenH3 mapping. While Tapestry detected telomeric sequences at only one end of eight other scaffolds and none for scaffold 18 and 19, the Hi-C map points towards the presence of telomeric repeats at both ends of all scaffolds 1-19 (Figure 1b), suggesting that the v1 assembly contains 19 full-length chromosomes that compose the nuclear genome. Scaffolds 20 and 21 were considerably shorter with ~162 kb and ~70 kb and displayed a markedly lower GC content at 41-42% (Figure 1a), suggesting that these scaffolds represent the chloroplast and mitochondrial genomes, respectively. BLAST analyses confirmed the presence of plastid and mitochondrial genes on the respective scaffolds, and the overall scaffold lengths corresponded with the sizes of the plastid and mitochondrial genomes of *Coccomyxa subellipsoidea* C-169 with 175 kb and 65 kb, respectively (Blanc et al., 2012). Full annotation of scaffolds 20 and 21 showed that they indeed represent chloroplast and mitochondrial genomes, respectively (Figure 2).

To rule out the presence of contaminants, the assembly and PacBio HiFi raw reads were used to produce a Blobplot (Camacho et al., 2009; Laetsch and Blaxter, 2017), which indicates that 98.76% of the reads match only the *Coccomyxa* genus (Figure 3) and, consequently, that the original sample was free of contaminating organisms. Finally, a KAT analysis showed a single peak of k-mer multiplicity based on HiFi reads that were represented once in the assembly (Figure 4) (Mapleson et al., 2016), indicative of a high-quality, haploid genome.

To annotate the nuclear genome, we first assessed the presence of repetitive elements. In total, we found 8.9% of the genome to be repetitive (Table 2), comparable to the 7.2% of repetitive sequences found in the genome of *C. supellipsoidea* C-169 (Blanc et al., 2012). These 8.9% repetitive elements were annotated as either simple repeats (2.3%) or transposable elements (6.6%). Of the transposable elements, 36% were annotated as retrotransposons and 64% as DNA transposons. The distribution of the repetitive elements was even across the genome with only a few repeat-rich regions (Figure 5). Next, we aimed to produce a high-quality genome annotation using RNA sequencing data. In total 13,557 genes were annotated with an average length of 3.1 kb (Table 2). The amount of alternative splicing in the genome is predicted to be very low, given the average of one transcript per gene model. To confirm the actual amount of alternative splicing, however, further analyses will be required. Of the 13,557 genes, 68% have annotated PFAM domains and 962 are predicted to carry a signal peptide for secretion. A total of 1,489 (98.6 %) complete gene models among 1,519 conserved Benchmarking Universal Single-Copy Orthologs (BUSCO) (Manni et al., 2021) in the chlorophyta_odb10 database were identified (Table 2), suggesting a highly complete genome annotation.

Until recently, the taxonomic classification and definition of *Coccomyxa* species was based on environmentally variable morphological and cytological characteristics. This classification was reviewed based on the phylogenetic analyses of nuclear SSU and ITS rDNA sequences, which resulted in the definition of 27 currently recognized *Coccomyxa* species (Darienko et al., 2015; Malavasi et al., 2016). Dot plot analysis of the high-quality genome assembly of *C. viridis* SAG216-4 with the assembly of the most closely related sequenced relative *C. subellipsoidea* C-169 revealed a lack of synteny since the few identified orthologous sequences were < 1 kb and, therefore, do not represent full-length genes (Figure 6a, Table 2). This lack of synteny was no technical artifact since the *C. viridis* assembly could be fully aligned to itself (Figure 6b), and BLAST analyses with five out of six non-identical ITS sequences identified in the *C. viridis* SAG 216-4 assembly confirmed its species identity. A comparison of the assembly of *C. subellipsoidea* C-169 to that of *Chlorella variabilis* (Chlorophyte, *Trebouxiophyceae*) has previously identified few syntenic regions which displayed poor gene collinearity (Blanc et al., 2012). Future studies will help to clarify whether the absence of synteny between *C. viridis* and *C. subellipsoidea* is due to the quality of the available assemblies or whether it has biological implications.

Data availability

Data for *C. viridis* SAG 216-4 with the ToLID ucCocViri1 is available via the European Nucleotide Archive (ENA) under the study accession number PRJNA1054215. Fastqc reports of raw data can be found in (Kraege et al., 2023).

Acknowledgements

This genome project is part of the DeRGA pilot study (https://www.erga-biodiversity.eu/pilot-project) and we acknowledge coordination and support by Dr. Astrid Böhne and Prof. Dr. Ann-Marie Waldvogel. Development of the pilot ERGA Data Portal (https://portal.erga-biodiversity.eu/) was funded by the European Molecular Biology Laboratory. We thank the West German Genome Center (WGGC) and the Cologne Center for Genomics (CCG) for library generation and quality control as well as their support for PacBio Hifi, Hi-C and RNA sequencing.

Conflict of interest

The authors declare no conflict of interest.

Funding information

BPHJT acknowledges funding by the Alexander von Humboldt Foundation in the framework of an Alexander von Humboldt Professorship endowed by the German Federal Ministry of Education and Research and is furthermore supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC 2048/1 – Project ID: 390686111. This research was also funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – SFB1535 - Project ID 458090666. PHS was funded by a DFG ENP grant (grant number: 434028868), which also funded JK's position. NG was first funded through a DFG grant to PHS (458953049) and subsequently through the European Union's Horizon Europe Research and Innovation program under the Marie Skłodowska-Curie grant agreement No. 101110569.

References

Abrusán, G., Grundmann, N., DeMester, L., and Makalowski, W. (2009). TEclass--a tool for automated classification of unknown eukaryotic transposable elements. Bioinformatics *25*, 1329-1330.

Blanc, G., Agarkova, I., Grimwood, J., Kuo, A., Brueggeman, A., Dunigan, D.D., Gurnon, J., Ladunga, I., Lindquist, E., Lucas, S.*, et al.* (2012). The genome of the polar eukaryotic microalga *Coccomyxa subellipsoidea* reveals traits of cold adaptation. Genome Biol *13*, R39.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden, T.L. (2009). BLAST+: architecture and applications. BMC Bioinformatics *10*, 421.

Cao, S., Zhang, F., Zheng, H., Liu, C., Peng, F., and Zhou, Q. (2018a). *Coccomyxa antarctica* sp. nov. from the Antarctic lichen *Usnea aurantiacoatra*. PhytoKeys, 107-115.

Cao, S., Zhang, F., Zheng, H., Peng, F., Liu, C., and Zhou, Q. (2018b). *Coccomyxa greatwallensis* sp. nov. (Trebouxiophyceae, Chlorophyta), a lichen epiphytic alga from Fildes Peninsula, Antarctica. PhytoKeys, 39-50.

Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A., Keane, T., McCarthy, S.A., Davies, R.M.*, et al.* (2021). Twelve years of SAMtools and BCFtools. Gigascience *10*.

Darienko, T., Gustavs, L., Eggert, A., Wolf, W., and Proschold, T. (2015). Evaluating the species boundaries of green microalgae (Coccomyxa, Trebouxiophyceae, Chlorophyta) using integrative taxonomy and DNA barcoding with further implications for the species identification in environmental samples. PLoS One *10*, e0127838.

Davey, J.W., Davis, S.J., Mottram, C., and Ashton, P.D. (2020). Tapestry: validate and edit small eukaryotic genome assemblies with long reads. bioRxiv.

Dudchenko, O., Batra, S.S., Omer, A.D., Nyquist, S.K., Hoeger, M., Durand, N.C., Shamim, M.S., Machol, I., Lander, E.S., Aiden, A.P.*, et al.* (2017). *De novo* assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds. Science *356*, 92-95.

Durand, N.C., Robinson, J.T., Shamim, M.S., Machol, I., Mesirov, J.P., Lander, E.S., and Aiden, E.L. (2016a). Juicebox provides a visualization system for Hi-C contact maps with unlimited zoom. Cell Syst *3*, 99-101.

Durand, N.C., Shamim, M.S., Machol, I., Rao, S.S.P., Huntley, M.H., Lander, E.S., and Aiden, E.L. (2016b). Juicer provides a one-click system for analyzing loop-resolution Hi-C experiments. Cell Systems *3*, 95-98.

Faluaburu, M.S., Nakai, R., Imura, S., and Naganuma, T. (2019). Phylotypic characterization of
mycobionts and photobionts of rock tripe lichen in East Antarctica. Microorganisms *7*.

Flynn, J.M., Hubley, R., Goubert, C., Rosen, J., Clark, A.G., Feschotte, C., and Smit, A.F. (2020).
RepeatModeler2 for automated genomic discovery of transposable element families. Proc Natl Acad
Sci U S A *117*, 9451-9457.

Gray, A.P., Lucas, I.A.N., Seed, R., and Richardson, C.A. (1999). *Mytilus edulis chilensis* infested with
Coccomyxa parasitica (Chlorococcales, Coccomyxaceae). Journal of Molluscan Studies *65*, 289-294.

Greiner, S., Lehwark, P., and Bock, R. (2019). OrganellarGenomeDRAW (OGDRAW) version 1.3.1:
expanded toolkit for the graphical visualization of organellar genomes. Nucleic Acids Res *47*, W59-
W64.

Gu, Z., Gu, L., Eils, R., Schlesner, M., and Brors, B. (2014). Circlize implements and enhances
circular visualization in R. Bioinformatics *30*, 2811-2812.

Gustavs, L., Schiefelbein, U., Darienko, T., and Pröschold, T. (2017). Symbioses of the green algal
genera *Coccomyxa* and *Elliptochloris* (Trebouxiophyceae, Chlorophyta). In Algal and Cyanobacteria
Symbioses, M. Grube, J. Seckbach, and L. Muggia, eds. (Europe: World Scientific), pp. 169-208.

Hoff, K.J., Lange, S., Lomsadze, A., Borodovsky, M., and Stanke, M. (2016). BRAKER1:
Unsupervised RNA-Seq-based genome annotation with GeneMark-ET and AUGUSTUS.
Bioinformatics *32*, 767-769.

Jaag, O. (1933). *Coccomyxa* Schmidle- Monographie einer Algengattung (Bern: Gebrüder Fretz).

Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol *37*, 907-915.

Koren, S., Walenz, B.P., Berlin, K., Miller, J.R., Bergman, N.H., and Phillippy, A.M. (2017). Canu:
scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation.
Genome Res *27*, 722-736.

Kraege, A., Thomma, B.P.H.J., and Rovenich, H. (2023). Fastqc reports of sequencing data from
Coccomyxa viridis SAG 216-4.

Kulichovà, J., Škaloud, P., and Neustupa, J. (2014). Molecular diveristy of green corticolous
microalgae from two sub-Mediterranean European localities. European Journal of Phycology *49*, 345-
355.

Kurtz, S., Phillippy, A., Delcher, A.L., Smoot, M., Shumway, M., Antonescu, C., and Salzberg, S.L.
(2004). Versatile and open software for comparing large genomes. Genome Biol *5*, R12.

Laetsch, D.R., and Blaxter, M.L. (2017). BlobTools: Interrogation of genome assemblies. F1000
Research *6*, 1287.

Lang, B.F., Beck, N., Prince, S., Sarrasin, M., Rioux, P., and Burger, G. (2023). Mitochondrial genome
annotation with MFannot: a critical analysis of gene identification and gene model prediction. Front
Plant Sci *14*, 1222186.

Leliaert, F., Smith, D.R., Moreau, H., Herron, M.D., Verbruggen, H., Delwiche, C.F., and De Clerck, O.
(2012). Phylogeny and molecular evolution of the green algae. Crit Rev Plant Sci *31*, 1-46.

Li, H. (2021). New strategies to improve minimap2 alignment accuracy. Bioinformatics *37*, 4572-4574.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics *25*, 1754-1760.

Li, L., Wang, S., Wang, H., Sahu, S.K., Marin, B., Li, H., Xu, Y., Liang, H., Li, Z., Cheng, S.*, et al.* (2020). The genome of *Prasinoderma coloniale* unveils the existence of a third phylum within green plants. Nat Ecol Evol *4*, 1220-1231.

Malavasi, V., Skaloud, P., Rindi, F., Tempesta, S., Paoletti, M., and Pasqualetti, M. (2016). DNA-Based taxonomy in ccologically versatile microalgae: A re-evaluation of the species concept within the coccoid green algal genus *Coccomyxa* (Trebouxiophyceae, Chlorophyta). PLoS One *11*, e0151137.

Manni, M., Berkeley, M.R., Seppey, M., Simao, F.A., and Zdobnov, E.M. (2021). BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes. Mol Biol Evol *38*, 4647-4654.

Mapleson, D., Accinelli, G.G., Kettleborough, G., Wright, J., and Clavijo, B.J. (2016). KAT: A K-mer Analysis Toolkit to quality control NGS datasets and genome assemblies. Bioinformatics *33*, 574-576.

Marin, B. (2012). Nested in the Chlorellales or independent class? Phylogeny and classification of the Pedinophyceae (Viridiplantae) revealed by molecular phylogenetic analyses of complete nuclear and plastid-encoded rRNA operons. Protist *163*, 778-805.

McFadden, G.I., and Melkonian, M. (1986). Use of Hepes buffer for microalgal culture media and fixation for electron microscopy. Phycologia *25*, 551-557.

Morris, J.L., Puttick, M.N., Clark, J.W., Edwards, D., Kenrick, P., Pressel, S., Wellman, C.H., Yang, Z., Schneider, H., and Donoghue, P.C.J. (2018). The timescale of early land plant evolution. Proc Natl Acad Sci U S A *115*, E2274-E2283.

Paysan-Lafosse, T., Blum, M., Chuguransky, S., Grego, T., Pinto, B.L., Salazar, G.A., Bileschi, M.L., Bork, P., Bridge, A., Colwell, L.*, et al.* (2023). InterPro in 2022. Nucleic Acids Res *51*, D418-D427.

Pertea, G., and Pertea, M. (2020). GFF utilities: GffRead and GffCompare. F1000 Research *9*, 304.

Riehl, K., Riccio, C., Miska, E.A., and Hemberg, M. (2022). TransposonUltimate: software for transposon classification, annotation and detection. Nucleic Acids Res *50*, e64.

Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and Mesirov, J.P. (2011). Integrative genomics viewer. Nat Biotechnol *29*, 24-26.

Schmidle, W. (1901). Über drei Algengenera. Berichte der deutschen botanischen Gesellschaft *19*, 10-24.

Sciuto, K., Baldan, B., Maracto, S., and Moro, I. (2019). *Coccomyxa cimbria* sp. nov., a green microalga found in association with carnivorous plants of the genus *Drosera* L. European Journal of Phycology *54*, 531-547.

Smit, A.F., Hubley, R., and Green, P. (2012). RepeatMasker (Retrieved from https://repeatmasker.org).

Sokolnikova, Y., Magarlamov, T., Stenkova, A., and Kumeiko, V. (2016). Permanent culture and parasitic impact of the microalga *Coccomyxa parasitica*, isolated from horse mussel *Modiolus kurilensis*. J Invertebr Pathol *140*, 25-34.

Sokolnikova, Y., Tumas, A., Stenkova, A., Slatvinskaya, V., Magarlamov, T., and Smagina, E. (2022). Novel species of parasitic green microalgae *Coccomyxa veronica* sp. nov. infects *Anadara broughtonii* from Sea of Japan. Symbiosis *87*, 293-305.

Štifterovà, A., and Neustupa, J. (2015). Community structure of corticolous microalgae within a single forest stand: evaluating the effects of bark surface pH and tree species. Fottea Olomouc *15*, 113-122.

Tagirdzhanova, G., Scharnagl, K., Yan, X., and Talbot, N.J. (2023). Genomic analysis of *Coccomyxa*
viridis, a common low-abundance alga associated with lichen symbioses. Sci Rep *13*, 21285.

Teufel, F., Almagro Armenteros, J.J., Johansen, A.R., Gislason, M.H., Pihl, S.I., Tsirigos, K.D.,
Winther, O., Brunak, S., von Heijne, G., and Nielsen, H. (2022). SignalP 6.0 predicts all five types of
signal peptides using protein language models. Nat Biotechnol *40*, 1023-1025.

Tillich, M., Lehwark, P., Pellizzer, T., Ulbricht-Jones, E.S., Fischer, A., Bock, R., and Greiner, S.
(2017). GeSeq - versatile and accurate annotation of organelle genomes. Nucleic Acids Res *45*, W6-
W11.

Trémouillaux-Guiller, J., Rohr, T., Rohr, R., and Huss, V.A.R. (2002). Discovery of an endophytic alga
in *Ginkgo biloba*. Am J Bot *89*, 727-733.

Vaschenko, M.A., Kovaleva, A.L., Syasina, I.G., and Kukhlevsky, A.D. (2013). Reproduction-related
effects of green alga *Coccomyxa* sp. infestation in the horse mussel *Modiolus modiolus*. J Invertebr
Pathol *113*, 86-95.

Vaser, R., and Šikić, M. (2021). Time- and memory-efficient genome assembly with Raven. Nature
Computational Science *1*, 332-336.

Wick, R. (2018). Porechop (Retrieved from https://github.com/rrwick/Porechop).

Yahr, R., Florence, A., Škaloud, P., and Voytsekhovich, A. (2015). Molecular and morphological
diversity in photobionts associated with *Micarea* s. str. (*Lecanorales*, Ascomycota). The Lichenologist
47, 403-414.

Zoller, S., and Lutzoni, F. (2003). Slow algae, fast fungi: exceptionally high nucleotide substitution rate
differences between lichenized fungi Omphalina and their symbiotic green algae Coccomyxa. Mol
Phylogenet Evol *29*, 629-640.
436
437

Figure legends

Figure 1. Genome assembly of *Coccomyxa viridis* SAG 216-4. (a) An overview of the *C. viridis* genome assembly depicts chromosome-scale scaffolds. Green bars indicate scaffold sizes and red bars represent telomeres. Variations in color intensities correlate with read coverage. Read coverage per scaffold is determined by mapping PacBio HiFi reads onto the assembly. Scaffolds 20 and 21 were identified as chloroplast and mitochondrial genomes based on size and low GC contents, and BLAST analyses. (b) Hi-C contact map showing interaction frequencies between regions in the nuclear genome of *Coccomyxa viridis*. Scaffolds are framed by blue lines while contigs within scaffolds are depicted in green.

Figure 2 Scaffolds 20 and 21 represent the plastid and mitochondrial genomes of *C. viridis* SAG 216-4. Gene maps of the chloroplast (a) and mitochondrial (b) genomes. The inner circles indicate the GC content and mapped genes are shown on the outer circles. Genes that are transcribed clockwise are placed inside the outer circles, and genes that are transcribed counterclockwise at the outside of the outer circles.

Figure 3. Taxonomic annotation indicates absence of contaminations in the genome assembly. (b) Taxon-annotated GC coverage scatter plot (Blobplot) of the contigs from the genome assembly shows that all scaffolds are taxon-annotated as *Coccomyxa* and all scaffolds that belong to the nuclear genome have similar GC contents (~54%). The GC content of the mitochondrial and plastid genomes are considerably lower (~41%). (b) In total 98.76% of the reads can be mapped onto the assembly and are therefore classified as *Coccomyxa* reads.

Figure 4. The *Coccomyxa viridis* SAG 216-4 genome is haploid. The KAT specra-cn plot depicts the 27-mer multiplicity of the PacBio HiFi reads against the genome assembly. Black areas under the peaks represent k-mers present in the reads but absent from the assembly, colored peaks indicate k-mers that are present once to multiple times in the assembly. The single red peak in the KAT specra-cn plot suggests that *Coccomyxa viridis* has a haploid genome, while the black peak at low multiplicity shows that the assembly is highly complete and that all reads are represented in the assembly.

Figure 5. Circos plot summarizing the nuclear genome annotation of *Coccomyxa viridis* SAG 216-4. From outside to inside the tracks display: GC content (over 1-kb windows), gene density (blue) and repetitive element density (red).

Figure 6. No synteny detected between related *Coccomyxa* species. (a) Dot plot of
orthologous sequences in the genome assemblies of *C. viridis* SAG 216-4 and *C.*
subellipsoidea C-169. Violet and blue dots represent orthologous sequences on same and
opposite strands, respectively. Dot sizes does not correlate with the length of the sequences
they represent, which were all < 1 kb. The width of each box corresponds to the length (bp) of
the respective scaffold. (b) Dot plot of the genome assembly of *C. viridis* SAG216-4 against
itself.
481
482

Figure 1. Genome assembly of *Coccomyxa viridis* SAG 216-4. (a) An overview of the *C.*
viridis genome assembly depicts chromosome-scale scaffolds. Green bars indicate scaffold
sizes and red bars represent telomeres. Variations in color intensities correlate with read

487 coverage. Read coverage per scaffold is determined by mapping PacBio HiFi reads onto the
488 assembly. Scaffolds 20 and 21 were identified as chloroplast and mitochondrial genomes
489 based on size and low GC contents, and BLAST analyses. (b) Hi-C contact map showing
490 interaction frequencies between regions in the nuclear genome of *Coccomyxa viridis*.
491 Scaffolds are framed by blue lines while contigs within scaffolds are depicted in green.
492
493

495 **Figure 2 Scaffolds 20 and 21 represent the plastid and mitochondrial genomes of *C.***
496 ***viridis* SAG 216-4.** Gene maps of the chloroplast (a) and mitochondrial (b) genomes. The
497 inner circles indicate the GC content and mapped genes are shown on the outer circles. Genes
498 that are transcribed clockwise are placed inside the outer circles, and genes that are
499 transcribed counterclockwise at the outside of the outer circles.

Figure 3. Taxonomic annotation indicates absence of contaminations in the genome assembly. (b) Taxon-annotated GC coverage scatter plot (Blobplot) of the contigs from the genome assembly shows that all scaffolds are taxon-annotated as *Coccomyxa* and all scaffolds that belong to the nuclear genome (N) have similar GC contents (~54%). The GC

506　content of the mitochondrial (M) and plastid (P) genomes are considerably lower (~41%). (b)

507　In total 98.76% of the reads can be mapped onto the assembly and are therefore classified

508　as *Coccomyxa* reads.

Figure 4. The *Coccomyxa viridis* SAG 216-4 nuclear genome is haploid. The KAT specra-cn plot depicts the 27-mer multiplicity of the PacBio HiFi reads against the nuclear genome assembly. Black areas under the peaks represent k-mers present in the reads but absent from the assembly, colored peaks indicate k-mers that are present once to multiple times in the assembly. The single red peak in the KAT specra-cn plot suggests that *Coccomyxa viridis* has a haploid genome, while the black peak at low multiplicity shows that the assembly is highly complete and that all reads are represented in the assembly.

Figure 5. Circos plot summarizing the nuclear genome annotation of *Coccomyxa viridis*

SAG 216-4. From outside to inside the tracks display: GC content (over 1-kb windows), gene

density (blue) and repetitive element density (red).

Figure 6. No synteny detected between related _Coccomyxa_ species. (a) Dot plot of

orthologous sequences in the genome assemblies of _C. viridis_ SAG 216-4 and _C._

subellipsoidea C-169. Violet and blue dots represent orthologous sequences on same and

opposite strands, respectively. Dot sizes does not correlate with the length of the sequences

they represent, which were all < 1 kb. The width of each box corresponds to the length (bp) of

the respective scaffold. (b) Dot plot of the genome assembly of _C. viridis_ SAG216-4 against

itself.

Table 1. Genome features of *C. viridis* SAG 216-4 including the mitochondrial and plastid genomes.

Assembly ID	*C. viridis* SAG 216-4 genomes
Total length (bp)	50,911,578
No. of contigs	27
No. of scaffolds	21
Longest scaffold (bp)	4,477,725
N50 (bp)	2,669,017
L50	8
GC content (%)	54.5

541   **Table 2** Annotation features of the *C. viridis* SAG 216-4 nuclear genome.

Genome annotation	
Repeat content (%)	8.85
Retrotransposons	2.4
DNA transposons	4.2
Simple repeats	2.25
No. gene models	13,557
Average gene length (bp)	3146
No. exons	122,978
Average no. exons per gene model	9
Average exon length (bp)	158
No. transcripts	14,024
Average no. transcripts/gene model	1
No. gene models <200 bp length	0
No. proteins with ≥1 PFAM domain	9205
No. proteins with signal peptide	962
BUSCO (chlorophyta_odb10)	**C: 98.6% [S: 82.5%, D: 16.1%], F: 0.1%, M: 1.3%, N: 1519**

544 **Table S1.** Summary of bioinformatics tools used for genome assembly and annotation.

Assembly		Annotation	
Tool	Version	Tool	Version
Raven	v1.8.1	RepeatModeler	v2.0.3
Juicer	v2.0	TransposonUltimate	v1.0
BWA	v0.7.17-r1188	TEclass	v2.1.3
3d-dna	v180922	RepeatMasker	v4.1.2-p1
Juicebox	v1.11.08	HiSat2	v2.2.1
Minimap2	v2.24-r1122	Braker	v2.1.6
Samtools	v1.10	Gffread	v0.12.7
Integrative Genome Viewer	v2.11.2	SignalP	v6.0
Tapestry	v1.0.0	BUSCO	v5.3.2
Blobtools	v1.1.1	R	v4.2.0
BLAST	2.13.0+	Circilize	v0.4.14
Kmer Analysis Toolkit	V2.4.2	InterProScan	v5.61
Mummer	C3.23		