High quality genome assembly and annotation (v1) of the eukaryotic terrestrial microalga *Coccomyxa viridis* SAG 216-4

Anton Kraege[1], Edgar Chavarro-Carrero[1], Nadège Guiglielmoni[2], Eva Schnell[1], Joseph Kirangwa[2], Stefanie Heilmann-Heimbach[3,4], Kerstin Becker[5,6], Karl Köhrer[6], WGGC Team, DeRGA Community, Philipp Schiffer[2], Bart P. H. J. Thomma[1,7] and Hanna Rovenich[1]

[1] Institute of Plant Sciences, University of Cologne, Germany; [2] Institute of Zoology, University of Cologne, Germany; [3] Institute of Human Genetics, University Hospital of Bonn & University of Bonn, Bonn, Germany; [4] NGS Core Facility, Medical Faculty of the University of Bonn, Bonn, Germany; [5] Cologne Center for Genomics (CCG), Medical Faculty, University of Cologne, 50931 Cologne, Germany; [6] Biological and Medical Research Centre (BMFZ), Genomics & Transcriptomics Laboratory, Heinrich-Heine-University Düsseldorf, Germany; [7] Cluster of Excellence on Plant Sciences (CEPLAS), Germany

corresponding author: hanna.roevenich@uni-koeln.de

Abstract

Unicellular green algae of the genus *Coccomyxa* are recognized for their worldwide distribution and ecological versatility. Most species described to date live in close association with various host species, such as in lichen associations. However, little is known about the molecular mechanisms that drive such symbiotic lifestyles. We generated a high-quality genome assembly for the lichen photobiont *Coccomyxa viridis* SAG 216-4 (formerly *C. mucigena*). Using long-read PacBio HiFi and Oxford Nanopore Technologies in combination with chromatin conformation capture (Hi-C) sequencing, we assembled the genome into 21 scaffolds with a total length of 50.9 Mb, an N50 of 2.7 Mb and a BUSCO score of 98.6%. While nineteen scaffolds represent full-length nuclear chromosomes, two additional scaffolds represent the mitochondrial and plastid genomes. Transcriptome-guided gene annotation resulted in the identification of 13,557 protein-coding genes, of which 68% have annotated PFAM domains and 962 are predicted to be secreted.

Keywords: *Coccomyxa viridis*, EBP, ERGA, long-read sequencing, genome assembly, genome annotation, Trebouxiophyceae

Introduction

Green algae are photosynthesizing eukaryotic organisms that differ greatly in terms of morphology and colonize a large variety of aquatic and terrestrial habitats. Phylogenetically, green algae form a paraphyletic group that comprises the chlorophyte and streptophyte lineages, which diverged between 1,000 and 700 million years ago (Morris et al., 2018). While the streptophyte lineage encompasses charophyte green algae as well as land plants, the chlorophyte lineage consists of 7 early diverging prasinophyte classes, which gave rise to 4 phycoplast-containing core chlorophyte classes (Chlorodendrophyceae, Trebouxiophyceae, Ulvophyceae, Chlorophyceae) with one independent sister class (Pedinophyceae) (Leliaert et al., 2012; Marin, 2012).

The *Coccomyxa* genus is represented by coccoid unicellular green algae that belong to the class of Trebouxiophyceae. Morphologically, *Coccomyxa* spp. are characterized by irregular elliptical to globular cells that range from 6–14 x 3–6 μm in size, with a single parietal chloroplast lacking pyrenoids and the absence of flagellate stages (Schmidle, 1901). Members of this genus are found in freshwater, marine, and various terrestrial habitats where they occur free-living or in symbioses with diverse hosts (Darienko et al., 2015; Gustavs et al., 2017; Malavasi et al., 2016). Several *Coccomyxa* species establish stable, mutualistic associations with fungi that result in the formation of complex three-dimensional architectures, known as lichens (Faluaburu et al., 2019; Gustavs et al., 2017; Jaag, 1933; Yahr et al., 2015; Zoller and Lutzoni, 2003). Others associate with vascular plants or lichens as endo- or epiphytes, respectively (Cao et al., 2018a; Cao et al., 2018b; Trémouillaux-Guiller et al., 2002), and frequently occur on the bark of trees (Kulichovà et al., 2014; Štifterovà and Neustupa, 2015) where they may interact with other microbes. One novel species was recently found in association with carnivorous plants, even though the nature of this relationship remains unclear (Sciuto et al., 2019). Besides, *Coccomyxa* also establishes parasitic interactions with different mollusk species affecting their filtration ability and reproduction (Gray et al., 1999; Sokolnikova et al., 2016; Sokolnikova et al., 2022; Vaschenko et al., 2013).

Despite this ecological versatility, little is known about the molecular mechanisms that determine the various symbiotic lifestyles in *Coccomyxa* and a genome sequence is available only for a single *Coccomyxa* species, namely *Coccomyxa subellipsoidea* C-169 that was isolated on Antarctica where it occurred on dried algal peat (Blanc et al., 2012). Here, we combined long-read PacBio HiFi and Oxford Nanopore Technology (ONT) sequencing with chromatin conformation capture (Hi-C) sequencing to generate a high-quality chromosome-scale assembly of *C. mucigena* SAG 216-4. Recent SSU and ITS rDNA sequencing-based re-evaluations of the *Coccomyxa* phylogeny placed the SAG 216-4 isolate in the clade of *C.*

72 *viridis* (Darienko et al., 2015; Malavasi et al., 2016). Hence, this isolate will be referred to as

73 *C. viridis* here and data have been deposited under the corresponding Taxonomy ID.

Materials & Methods

Sample information

Coccomyxa viridis (formerly *Coccomyxa mucigena*) SAG 216-4 was ordered from the Culture

Collection of Algae at the Georg-August-University Göttingen (*Sammlung von Algenkulturen*

der Universität Göttingen, international acronym SAG), Germany. The stock culture was

reactivated in liquid modified Waris-H growth medium (McFadden and Melkonian, 1986) with

soil extract and 3x vitamins, and maintained through regular medium replacement. Cultures

were grown at ~ 15 µmol photons m^{-2} s^{-1} (fluorescent light tubes: L36W/640i energy saver cool

white and L58W/956 BioLux, Osram, Munich, Germany) in a 14/10 hr light/dark cycle at 20°C.

DNA and *RNA extraction*

Cells of a 7-week-old *C. viridis* culture were harvested over 0.8 µm cellulose nitrate filters

(Sartorius, Göttingen, Germany) using a vacuum pump. Material was collected with a spatula,

snap-frozen and ground in liquid nitrogen using mortar and pestle. The ground material was

used for genomic DNA extraction with the RSC Plant DNA Kit (Promega, Madison, WI, USA)

using the Maxwell® RSC device according to manufacturer's instructions. To prevent shearing

of long DNA fragments, centrifugation was carried out at 10,000 *g* during sample preparation.

Following DNA extraction, DNA fragments <10,000 bp were removed using the SRE XS kit

(Circulomics, Baltimore, MD, USA) according to manufacturer's instructions. DNA quality and

quality were assessed using the Nanodrop 2000 spectrometer and Qubit 4 fluorometer with

the dsDNA BR assay kit (Invitrogen, Carlsbad, CA, USA), and integrity was confirmed by gel

electrophoresis. High-molecular weight DNA was stored at -20°C.

For total RNA extraction, algal cells were collected from a dense 9-day-old culture and

ground in liquid nitrogen using mortar and pestle. RNA was extracted with the Maxwell RSC

Plant RNA kit (Promega, Madison, WI, USA) using the Maxwell® RSC device according to

manufacturer's instructions. RNA quality and quantity was determined using the Nanodrop

2000 and stored at -80°C.

Pacific Biosciences High-Fidelity (PacBio HiFi) sequencing

104 HiFi libraries were prepared with the Express 2.0 Template kit (Pacific Biosciences, Menlo

105 Park, CA, USA) and sequenced on a Sequel II/Sequel IIe instrument with 30h movie time. HiFi

106 reads were generated using SMRT Link (v10; (Pacific Biosciences, Menlo Park, CA, USA)

107 with default parameters.

Oxford Nanopore Technologies (ONT) sequencing

Library preparation with the Rapid Sequencing Kit (SQK-626 RAD004) was performed with ~400 ng HMW DNA according to manufacturer's instructions (Oxford Nanopore Technologies, Oxford, UK). The sample was loaded onto an R9.4.1 flow cell (Oxford Nanopore Technologies, Oxford, UK), which was run for 24 h. Subsequent base calling was performed using Guppy (version 630 3.1.3; Oxford Nanopore Technologies, Oxford, UK). Adapter sequences were removed using Porechop (version 0.2.4 with default settings) (Wick, 2018), and the reads were self-corrected and trimmed using Canu (version 1.8) (Koren et al., 2017).

Chromosome conformation capture (Hi-C) and sequencing

C. viridis cells were cross-linked in 3% formaldehyde for 1 hour at room temperature. The reaction was quenched with glycine at a final concentration of 250 mM. Cells were collected by centrifugation at 16,000 *g* for 10 min. Pellets were flash-frozen in liquid nitrogen and ground using mortar and pestle. Hi-C libraries were prepared using the Arima-HiC+ kit (Arima Genomics, Carlsbad, CA, USA) according to manufacturer's instructions, and subsequently paired-end (2x150 bp) sequenced on a NovaSeq 6000 instrument (Illumina, San Diego, CA, USA).

RNA sequencing

Library preparation for full-length mRNASeq was performed using the NEB Ultra II Directional RNA Library Prep with NEBNext Poly(A) mRNA Magenetic Isolation Module and 500ng total RNA as starting material, except for W-RNA Lplaty, where library prep was based on 100ng total RNA as starting material. Sequencing was performed on an Illumina NovaSeq 6000 device with 2x150bp paired-end sequencing protocol and >50M reads per sample.

Genome assembly

PacBio HiFi reads were assembled using Raven (v1.8.1) (Vaser and Šikić, 2021) with default settings. Hi-C reads were mapped onto this assembly with Juicer (v2.0) using the "assembly" option to skip the post-processing steps and generate the merged_nodups.txt file (Durand et al., 2016b). For the juicer pipeline, restriction site maps were generated using the *Dpn*II (GATC) and *Hin*fI (GANTC) restriction site profile and the assembly was indexed with BWA index (v0.7.17-r1188) (Li and Durbin, 2009), and used to polish the assembly using 3d-dna (v180922) (Dudchenko et al., 2017). Afterwards, Juicebox (v1.11.08) was used to manually curate the genome assembly by splitting contigs and rearranging them according to the Hi-C pattern (Durand et al., 2016a). Contigs were merged to scaffolds according to the Hi-C map and Ns were introduced between contigs within scaffolds, gaps between contigs were removed and contigs were merged. Subsequently, ONT reads were mapped to the assembly

using Minimap2 (v2.24-r1122) and Samtools (v1.10) and mapped reads were visualized in
Integrative Genome Viewer (v2.11.2) (Danecek et al., 2021; Li, 2021; Robinson et al., 2011).
Whenever gaps between contigs were spanned by at least five reads with a mapping quality
of 30, the contigs were fused in the assembly.

Potential telomeres were identified using tapestry (v1.0.0) with "AACCCT" as telomere
sequence (Davey et al., 2020). To check for potential contaminations, Blobtools (v1.1.1) and
BLAST (v2.13.0+) were used to create a Blobplot including taxonomic annotation at genus
level (Camacho et al., 2009; Laetsch and Blaxter, 2017). To check completeness of the
assembly and retrieve ploidy information, kat comp from the Kmer Analysis Toolkit (v2.4.2)
was used, and results were visualized using the kat plot spectra-cn function with the -x 800
option to extend the x-axis (Mapleson et al., 2016). Genome synteny to the closest sequenced
relative *C. subellipsoidea* C-169 was determined using Mummer3 (Blanc et al., 2012; Kurtz et
al., 2004). In detail, the two assemblies were first aligned using Nucmer, followed by a filtering
step with Delta-filter using the many-to-many option (-m). Finally, the alignment was visualized
with Mummerplot.

Annotation

To annotate repetitive elements, a database of simple repeats was created with
RepeatModeler (v2.0.3) that was expanded with transposable elements (TE) from the
TransposonUltimate resonaTE (v1.0) pipeline (Flynn et al., 2020; Riehl et al., 2022). This
pipeline uses multiple tools for TE prediction and combines the prediction output. For the
prediction of TEs in *Coccomyxa viridis* helitronScanner, ltrHarvest, mitefind, mitetracker,
RepeatModeler, RepeatMasker, sinefind, tirvish, transposonPSI and NCBICDD1000 were
used within TransposonUltimate resonaTE and TEs that were predicted by at least two tools
were added to the database. TEclass (v2.1.3) was used for classification (Abrusán et al.,
2009). To softmask the genome and obtain statistics on the total TE and repetitive element
content in the genome, RepeatMasker (v4.1.2-p1)(Smit et al., 2012) was used with excln
option to exclude Ns in the masking.

Gene annotation was performed making use of RNA sequencing data. To this end,
the genome was indexed, and reads were mapped with HiSat2 (v2.2.1) using default settings
(Kim et al., 2019). Afterwards, Braker (v2.1.6) was used for transcriptome-guided gene
prediction with default settings (Bruna et al., 2021). To generate protein and coding sequence
files the Braker output was transformed with Gffread (v0.12.7) (Pertea and Pertea, 2020).
PFAM domain annotation was performed with InterProScan (v5.61) (Paysan-Lafosse et al.,
2023). To estimate the number of secreted proteins, SignalP (v6.0) was run in the slow-
sequential mode on the annotated proteins (Teufel et al., 2022). Finally, BUSCO (v5.3.2) was
run with the Chlorophyta database (chlorophyta_odb10) to estimate the completeness of the

gene annotation (Manni et al., 2021). The circos plot visualization of the annotation was
created with R (v4.2.0) and Circilize (v0.4.14) (Gu et al., 2014). All software and tools used for
the genome assembly and annotation are summarized in Table 1.

Results

The version 1 genome of *C. viridis* was assembled from 32.2 Gbp of PacBio HiFi reads with
a mean read length of 15 kb, 0.95 Gbp Nanopore reads with a mean read length of 8.8 kb and
15 million pairs of Hi-C seq data. The PacBio HiFi reads were first assembled using Raven
(Vaser and Šikić, 2021), yielding 27 contigs. These contigs were scaffolded and manually
curated using Hi-C data (Dudchenko et al., 2017; Durand et al., 2016a; Durand et al., 2016b;
Li and Durbin, 2009). To close the remaining gaps between contigs within scaffolds, ONT
reads were mapped onto the assembly (Danecek et al., 2021; Li, 2021) and gaps that were
spanned by at least 5 ONT reads with a mapping quality >30 were manually closed, finally
resulting in 21 scaffolds consisting of 26 contigs with a total length of 50.9 Mb and an N50 of
2.7 Mb (Figure 1, Table 2). Using Tapestry (Davey et al., 2020), telomeric regions
([AACCCT]*n*) were identified at both ends of nine of the 21 scaffolds (≥5 repeats) (Figure 1a),
suggesting that these represent full-length chromosomes, which was confirmed by Hi-C
analysis (Figure 1b). While Tapestry detected telomeric sequences at only one end of eight
other scaffolds and none for scaffold 18 and 19, the Hi-C map points towards the presence of
telomeric repeats at both ends of all scaffolds 1-19 (Figure 1b), suggesting that the v1
assembly contains 19 full-length chromosomes that compose the nuclear genome. Scaffolds
20 and 21 were considerably shorter with ∼162 kb and ∼70 kb and displayed a markedly lower
GC content (Figure 1a), suggesting that these scaffolds represent the chloroplast and
mitochondrial genomes, respectively. BLAST analyses confirmed the presence of plastid and
mitochondrial genes on the respective scaffolds, and the overall scaffold lengths
corresponded with the sizes of the plastid and mitochondrial genomes of *Coccomyxa*
subellipsoidea C-169 (Smith et al., 2011). To rule out the presence of contaminants, the
assembly and PacBio HiFi raw reads were used to produce a Blobplot (Camacho et al., 2009;
Laetsch and Blaxter, 2017), which indicates that 98.76% of the reads match only the
Coccomyxa genus (Figure 2) and, consequently, that the original sample was free of
contaminating organisms. Finally, a KAT analysis showed a single peak of k-mer multiplicity
based on HiFi reads that were represented once in the assembly (Figure 3) (Mapleson et al.,
2016), indicative of a high-quality, haploid genome.

To annotate the genome, we first assessed the presence of repetitive elements. In
total, we found 8.9% of the genome to be repetitive (Table 3), comparable to the 7.2% of
repetitive sequences found in the genome of *C. supellipsoidea* C-169 (Blanc et al., 2012).

These 8.9% repetitive elements were annotated as either simple repeats (2.3%) or transposable elements (6.6%). Of the transposable elements, 36% were annotated as retrotransposons and 64% as DNA transposons. The distribution of the repetitive elements was even across the genome with only a few repeat-rich regions (Figure 4). Next, we aimed to produce a high-quality genome annotation using RNA sequencing data. In total 13,557 genes were annotated with an average length of 3.1 kb (Table 3). The amount of alternative splicing in the genome is predicted to be very low, given the average of one transcript per gene model. Of the 13,557 genes, 68% have annotated PFAM domains and 962 are predicted to carry a signal peptide for secretion. A total of 1,489 (98.6 %) complete gene models among 1,519 conserved Benchmarking Universal Single-Copy Orthologs (BUSCO) (Manni et al., 2021) in the chlorophyta_odb10 database were identified (Table 3), suggesting a highly complete genome annotation.

Until recently, the taxonomic classification and definition of *Coccomyxa* species was based on environmentally variable morphological and cytological characteristics. This classification was reviewed based on the phylogenetic analyses of nuclear SSU and ITS rDNA sequences, which resulted in the definition of 27 currently recognized *Coccomyxa* species (Darienko et al., 2015; Malavasi et al., 2016). Dot plot analysis of the high-quality genome assembly of *C. viridis* SAG216-4 with the assembly of the most closely related sequenced relative *C. subellipsoidea* C-169 revealed a lack of synteny since the few identified orthologous sequences were < 1 kb and, therefore, do not represent full-length genes (Figure 5a, Table 3). This lack of synteny was no technical artifact since the *C. viridis* assembly could be fully aligned to itself (Figure 5b), and BLAST analyses of six identified ITS sequences in the *C. viridis* SAG 216-4 assembly confirmed its species identity. A comparison of the assembly of *C. subellipsoidea* C-169 to that of *Chlorella variabilis* (Chlorophyte, *Trebouxiophyceae*) has previously identified few syntenic regions which displayed poor gene collinearity (Blanc et al., 2012). Future studies will help to clarify whether the absence of synteny between *C. viridis* and *C. subellipsoidea* is due to the quality of the available assemblies or whether it has biological implications.

Data availability

Data submission for *C. viridis* SAG 216-4 with the ToLID ucCocViri1 to the European Nucleotide Archive (ENA) is in progress, accession numbers are pending. Fastqc reports of raw data can be found in (Kraege et al., 2023).

Acknowledgements

This genome project is part of the DeRGA pilot study (https://www.erga-biodiversity.eu/pilot-
project) and we acknowledge coordination and support by Dr. Astrid Böhne and Prof. Dr. Ann-
Marie Waldvogel. Development of the pilot ERGA Data Portal (https://portal.erga-
biodiversity.eu/) was funded by the European Molecular Biology Laboratory. We thank the
West German Genome Center (WGGC) and the Cologne Center for Genomics (CCG) for
library generation and quality control as well as their support for PacBio Hifi, Hi-C and RNA
sequencing.
260

Conflict of interest

262 The authors declare no conflict of interest.

Funding information

BPHJT acknowledges funding by the Alexander von Humboldt Foundation in the framework
of an Alexander von Humboldt Professorship endowed by the German Federal Ministry of
Educaion and Research and is furthermore supported by the Deutsche
Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence
Strategy – EXC 2048/1 – Project ID: 390686111. PHS was funded by a DFG ENP grant (grant
number: 434028868), which also funded JK's position. NG was first funded through a DFG
grant to PHS (458953049) and subsequently through the European Union's Horizon Europe
Research and Innovation program under the Marie Skłodowska-Curie grant agreement No.
101110569.

References

Abrusán, G., Grundmann, N., DeMester, L., and Makalowski, W. (2009). TEclass--a tool for
automated classification of unknown eukaryotic transposable elements. Bioinformatics *25*, 1329-1330.
doi:10.1093/bioinformatics/btp084.

Blanc, G., Agarkova, I., Grimwood, J., Kuo, A., Brueggeman, A., Dunigan, D.D., Gurnon, J., Ladunga,
I., Lindquist, E., Lucas, S*., et al.* (2012). The genome of the polar eukaryotic microalga *Coccomyxa*
subellipsoidea reveals traits of cold adaptation. Genome Biol *13*, R39. doi:10.1186/gb-2012-13-5-r39.

Bruna, T., Hoff, K.J., Lomsadze, A., Stanke, M., and Borodovsky, M. (2021). BRAKER2: automatic
eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein
database. NAR Genom Bioinform *3*, lqaa108. doi:10.1093/nargab/lqaa108.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden, T.L.
(2009). BLAST+: architecture and applications. BMC Bioinformatics *10*, 421. doi:10.1186/1471-2105-
10-421.

Cao, S., Zhang, F., Zheng, H., Liu, C., Peng, F., and Zhou, Q. (2018a). *Coccomyxa antarctica* sp.
nov. from the Antarctic lichen *Usnea aurantiacoatra*. PhytoKeys, 107-115.
doi:10.3897/phytokeys.98.25360.

Cao, S., Zhang, F., Zheng, H., Peng, F., Liu, C., and Zhou, Q. (2018b). *Coccomyxa greatwallensis* sp. nov. (Trebouxiophyceae, Chlorophyta), a lichen epiphytic alga from Fildes Peninsula, Antarctica. PhytoKeys, 39-50. doi:10.3897/phytokeys.110.26961.

Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A., Keane, T., McCarthy, S.A., Davies, R.M.*, et al.* (2021). Twelve years of SAMtools and BCFtools. Gigascience *10*. doi:10.1093/gigascience/giab008.

Darienko, T., Gustavs, L., Eggert, A., Wolf, W., and Proschold, T. (2015). Evaluating the species boundaries of green microalgae (Coccomyxa, Trebouxiophyceae, Chlorophyta) using integrative taxonomy and DNA barcoding with further implications for the species identification in environmental samples. PLoS One *10*, e0127838. doi:10.1371/journal.pone.0127838.

Davey, J.W., Davis, S.J., Mottram, C., and Ashton, P.D. (2020). Tapestry: validate and edit small eukaryotic genome assemblies with long reads. bioRxiv. doi:10.1101/2020.04.24.059402.

Dudchenko, O., Batra, S.S., Omer, A.D., Nyquist, S.K., Hoeger, M., Durand, N.C., Shamim, M.S., Machol, I., Lander, E.S., Aiden, A.P.*, et al.* (2017). *De novo* assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds. Science *356*, 92-95. doi:10.1126/science.aal3327.

Durand, N.C., Robinson, J.T., Shamim, M.S., Machol, I., Mesirov, J.P., Lander, E.S., and Aiden, E.L. (2016a). Juicebox provides a visualization system for Hi-C contact maps with unlimited zoom. Cell Syst *3*, 99-101. doi:10.1016/j.cels.2015.07.012.

Durand, N.C., Shamim, M.S., Machol, I., Rao, S.S.P., Huntley, M.H., Lander, E.S., and Aiden, E.L. (2016b). Juicer provides a one-click system for analyzing loop-resolution Hi-C experiments. Cell Systems *3*, 95-98. doi:10.1016/j.cels.2016.07.002.

Faluaburu, M.S., Nakai, R., Imura, S., and Naganuma, T. (2019). Phylotypic characterization of mycobionts and photobionts of rock tripe lichen in East Antarctica. Microorganisms *7*. doi:10.3390/microorganisms7070203.

Flynn, J.M., Hubley, R., Goubert, C., Rosen, J., Clark, A.G., Feschotte, C., and Smit, A.F. (2020). RepeatModeler2 for automated genomic discovery of transposable element families. Proc Natl Acad Sci U S A *117*, 9451-9457. doi:10.1073/pnas.1921046117.

Gray, A.P., Lucas, I.A.N., Seed, R., and Richardson, C.A. (1999). *Mytilus edulis chilensis* infested with *Coccomyxa parasitica* (Chlorococcales, Coccomyxaceae). Journal of Molluscan Studies *65*, 289-294. doi:10. 1093/ mollus/ 65.3. 289.

Gu, Z., Gu, L., Eils, R., Schlesner, M., and Brors, B. (2014). Circlize implements and enhances circular visualization in R. Bioinformatics *30*, 2811-2812. doi:10.1093/bioinformatics/btu393.

Gustavs, L., Schiefelbein, U., Darienko, T., and Pröschold, T. (2017). Symbioses of the green algal genera *Coccomyxa* and *Elliptochloris* (Trebouxiophyceae, Chlorophyta). In Algal and Cyanobacteria Symbioses, M. Grube, J. Seckbach, and L. Muggia, eds. (Europe: World Scientific), pp. 169-208.

Jaag, O. (1933). *Coccomyxa* Schmidle- Monographie einer Algengattung (Bern: Gebrüder Fretz).

Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol *37*, 907-915. doi:10.1038/s41587-019-0201-4.

Koren, S., Walenz, B.P., Berlin, K., Miller, J.R., Bergman, N.H., and Phillippy, A.M. (2017). Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res *27*, 722-736. doi:10.1101/gr.215087.116.

Kraege, A., Thomma, B.P.H.J., and Rovenich, H. (2023). Fastqc reports of sequencing data from *Coccomyxa viridis* SAG 216-4. doi:10.5281/zenodo.8084901.

Kulichovà, J., Škaloud, P., and Neustupa, J. (2014). Molecular diveristy of green corticolous
microalgae from two sub-Mediterranean European localities. European Journal of Phycology *49*, 345-
355. doi:10.1080/09670262.2014.945190.

Kurtz, S., Phillippy, A., Delcher, A.L., Smoot, M., Shumway, M., Antonescu, C., and Salzberg, S.L.
(2004). Versatile and open software for comparing large genomes. Genome Biol *5*, R12.
doi:10.1186/gb-2004-5-2-r12.

Laetsch, D.R., and Blaxter, M.L. (2017). BlobTools: Interrogation of genome assemblies. F1000
Research *6*, 1287. doi:10.12688/f1000research.12232.1.

Leliaert, F., Smith, D.R., Moreau, H., Herron, M.D., Verbruggen, H., Delwiche, C.F., and De Clerck, O.
(2012). Phylogeny and molecular evolution of the green algae. Crit Rev Plant Sci *31*, 1-46.
doi:10.1080/07352689.2011.615705.

Li, H. (2021). New strategies to improve minimap2 alignment accuracy. Bioinformatics *37*, 4572-4574.
doi:10.1093/bioinformatics/btab705.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics *25*, 1754-1760. doi:10.1093/bioinformatics/btp324.

Malavasi, V., Skaloud, P., Rindi, F., Tempesta, S., Paoletti, M., and Pasqualetti, M. (2016). DNA-
Based taxonomy in ccologically versatile microalgae: A re-evaluation of the species concept within the
coccoid green algal genus *Coccomyxa* (Trebouxiophyceae, Chlorophyta). PLoS One *11*, e0151137.
doi:10.1371/journal.pone.0151137.

Manni, M., Berkeley, M.R., Seppey, M., Simao, F.A., and Zdobnov, E.M. (2021). BUSCO update:
novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of
eukaryotic, prokaryotic, and viral genomes. Mol Biol Evol *38*, 4647-4654.
doi:10.1093/molbev/msab199.

Mapleson, D., Accinelli, G.G., Kettleborough, G., Wright, J., and Clavijo, B.J. (2016). KAT: A K-mer
Analysis Toolkit to quality control NGS datasets and genome assemblies. Bioinformatics *33*, 574-576.
doi:10.1093/bioinformatics/btw663.

Marin, B. (2012). Nested in the Chlorellales or independent class? Phylogeny and classification of the
Pedinophyceae (Viridiplantae) revealed by molecular phylogenetic analyses of complete nuclear and
plastid-encoded rRNA operons. Protist *163*, 778-805. doi:10.1016/j.protis.2011.11.004.

McFadden, G.I., and Melkonian, M. (1986). Use of Hepes buffer for microalgal culture media and
fixation for electron microscopy. Phycologia *25*, 551-557. doi:10.2216/i0031-8884-25-4-551.1.

Morris, J.L., Puttick, M.N., Clark, J.W., Edwards, D., Kenrick, P., Pressel, S., Wellman, C.H., Yang, Z.,
Schneider, H., and Donoghue, P.C.J. (2018). The timescale of early land plant evolution. Proc Natl
Acad Sci U S A *115*, E2274-E2283. doi:10.1073/pnas.1719588115.

Paysan-Lafosse, T., Blum, M., Chuguransky, S., Grego, T., Pinto, B.L., Salazar, G.A., Bileschi, M.L.,
Bork, P., Bridge, A., Colwell, L.*, et al.* (2023). InterPro in 2022. Nucleic Acids Res *51*, D418-D427.
doi:10.1093/nar/gkac993.

Pertea, G., and Pertea, M. (2020). GFF utilities: GffRead and GffCompare. F1000 Research *9*, 304.
doi:10.12688/f1000research.23297.1.

Riehl, K., Riccio, C., Miska, E.A., and Hemberg, M. (2022). TransposonUltimate: software for
transposon classification, annotation and detection. Nucleic Acids Res *50*, e64.
doi:10.1093/nar/gkac136.

Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and Mesirov,
J.P. (2011). Integrative genomics viewer. Nat Biotechnol *29*, 24-26. doi:10.1038/nbt.1754.

Schmidle, W. (1901). Über drei Algengenera. Berichte der deutschen botanischen Gesellschaft *19*,
10-24.

Sciuto, K., Baldan, B., Maracto, S., and Moro, I. (2019). *Coccomyxa cimbria* sp. nov., a green
microalga found in association with carnivorous plants of the genus *Drosera* L. European Journal of
Phycology *54*, 531-547. doi:10.1080/09670262.2019.1618920.

Smit, A.F., Hubley, R., and Green, P. (2012). RepeatMasker (Retrieved from
https://repeatmasker.org).

Smith, D.R., Burki, F., Yamada, T., Grimwood, J., Grigoriev, I.V., Van Etten, J.L., and Keeling, P.J.
(2011). The GC-rich mitochondrial and plastid genomes of the green alga *Coccomyxa* give insight into
the evolution of organelle DNA nucleotide landscape. PLoS One *6*, e23624.
doi:10.1371/journal.pone.0023624.

Sokolnikova, Y., Magarlamov, T., Stenkova, A., and Kumeiko, V. (2016). Permanent culture and
parasitic impact of the microalga *Coccomyxa parasitica*, isolated from horse mussel *Modiolus*
kurilensis. J Invertebr Pathol *140*, 25-34. doi:10.1016/j.jip.2016.07.012.

Sokolnikova, Y., Tumas, A., Stenkova, A., Slatvinskaya, V., Magarlamov, T., and Smagina, E. (2022).
Novel species of parasitic green microalgae *Coccomyxa veronica* sp. nov. infects *Anadara*
broughtonii from Sea of Japan. Symbiosis *87*, 293-305. doi:10.1007/s13199-022-00877-6.

Štifterovà, A., and Neustupa, J. (2015). Community structure of corticolous microalgae within a single
forest stand: evaluating the effects of bark surface pH and tree species. Fottea Olomouc *15*, 113-122.
doi:10.5507/fot.2015.013.

Teufel, F., Almagro Armenteros, J.J., Johansen, A.R., Gislason, M.H., Pihl, S.I., Tsirigos, K.D.,
Winther, O., Brunak, S., von Heijne, G., and Nielsen, H. (2022). SignalP 6.0 predicts all five types of
signal peptides using protein language models. Nat Biotechnol *40*, 1023-1025. doi:10.1038/s41587-
021-01156-3.

Trémouillaux-Guiller, J., Rohr, T., Rohr, R., and Huss, V.A.R. (2002). Discovery of an endophytic alga
in *Ginkgo biloba*. Am J Bot *89*, 727-733. doi:10.3732/ajb.89.5.727.

Vaschenko, M.A., Kovaleva, A.L., Syasina, I.G., and Kukhlevsky, A.D. (2013). Reproduction-related
effects of green alga *Coccomyxa* sp. infestation in the horse mussel *Modiolus modiolus*. J Invertebr
Pathol *113*, 86-95. doi:10.1016/j.jip.2013.02.003.

Vaser, R., and Šikić, M. (2021). Time- and memory-efficient genome assembly with Raven. Nature
Computational Science *1*, 332-336. doi:10.1038/s43588-021-00073-4.

Wick, R. (2018). Porechop (Retrieved from https://github.com/rrwick/Porechop).

Yahr, R., Florence, A., Škaloud, P., and Voytsekhovich, A. (2015). Molecular and morphological
diversity in photobionts associated with *Micarea* s. str. (*Lecanorales*, Ascomycota). The Lichenologist
47, 403-414. doi:10.1017/S0024282915000341.

Zoller, S., and Lutzoni, F. (2003). Slow algae, fast fungi: exceptionally high nucleotide substitution rate
differences between lichenized fungi Omphalina and their symbiotic green algae Coccomyxa. Mol
Phylogenet Evol *29*, 629-640. doi:10.1016/s1055-7903(03)00215-x.
417
418

Figure legends

Figure 1. Genome assembly of *Coccomyxa viridis* SAG 216-4. (a) An overview of the *C. viridis* genome assembly depicts chromosome-scale scaffolds. Green bars indicate scaffold sizes and red bars represent telomeres. Variations in color intensities correlate with read coverage. Read coverage per scaffold is determined by mapping PacBio HiFi reads onto the assembly. Scaffolds 20 and 21 were identified as chloroplast and mitochondrial genomes based on size and low GC contents, and BLAST analyses. (b) Hi-C contact map showing interaction frequencies between regions in the nuclear genome of *Coccomyxa viridis*. Scaffolds are framed by blue lines while contigs within scaffolds are depicted in green.

Figure 2. Taxonomic annotation indicates absence of contaminations in the genome assembly. (b) Taxon-annotated GC coverage scatter plot (Blobplot) of the contigs from the genome assembly shows that all scaffolds are taxon-annotated as *Coccomyxa* and all scaffolds that belong to the nuclear genome have similar GC contents (~54%). The GC content of the mitochondrial and plastid genomes are considerably lower (~41%). (b) In total 98.76% of the reads can be mapped onto the assembly and are therefore classified as *Coccomyxa* reads.

Figure 3. The *Coccomyxa viridis* SAG 216-4 genome is haploid. The KAT specra-cn plot depicts the 27-mer multiplicity of the PacBio HiFi reads against the genome assembly. Black areas under the peaks represent k-mers present in the reads but absent from the assembly, colored peaks indicate k-mers that are present once to multiple times in the assembly. The single red peak in the KAT specra-cn plot suggests that *Coccomyxa viridis* has a haploid genome, while the black peak at low multiplicity shows that the assembly is highly complete and that all reads are represented in the assembly.

Figure 4. Circos plot summarizing the nuclear genome annotation of *Coccomyxa viridis* SAG 216-4. From outside to inside the tracks display: GC content (over 1-kb windows), gene density (blue) and repetitive element density (red).

Figure 5. No synteny detected between related *Coccomyxa* species. (a) Dot plot of orthologous sequences in the genome assemblies of *C. viridis* SAG 216-4 and *C. subellipsoidea* C-169. Violet and blue dots represent orthologous sequences on same and opposite strands, respectively. Dot sizes does not correlate with the length of the sequences they represent, which were all < 1 kb. The width of each box corresponds to the length (bp) of

454   the respective scaffold. (b) Dot plot of the genome assembly of *C. viridis* SAG216-4 against

455   itself.

Figure 1. Genome assembly of *Coccomyxa viridis* SAG 216-4. (a) An overview of the *C. viridis* genome assembly depicts chromosome-scale scaffolds. Green bars indicate scaffold sizes and red bars represent telomeres. Variations in color intensities correlate with read

462 coverage. Read coverage per scaffold is determined by mapping PacBio HiFi reads onto the
463 assembly. Scaffolds 20 and 21 were identified as chloroplast and mitochondrial genomes
464 based on size and low GC contents, and BLAST analyses. (b) Hi-C contact map showing
465 interaction frequencies between regions in the nuclear genome of *Coccomyxa viridis*.
466 Scaffolds are framed by blue lines while contigs within scaffolds are depicted in green.
467
468

Figure 2. Taxonomic annotation indicates absence of contaminations in the genome assembly. (b) Taxon-annotated GC coverage scatter plot (Blobplot) of the contigs from the genome assembly shows that all scaffolds are taxon-annotated as *Coccomyxa* and all scaffolds that belong to the nuclear genome have similar GC contents (~54%). The GC

content of the mitochondrial and plastid genomes are considerably lower (~41%). (b) In total

98.76% of the reads can be mapped onto the assembly and are therefore classified as

Coccomyxa reads.

Figure 3. The *Coccomyxa viridis* SAG 216-4 genome is haploid. The KAT specra-cn plot

depicts the 27-mer multiplicity of the PacBio HiFi reads against the genome assembly. Black

areas under the peaks represent k-mers present in the reads but absent from the assembly,

colored peaks indicate k-mers that are present once to multiple times in the assembly. The

single red peak in the KAT specra-cn plot suggests that *Coccomyxa viridis* has a haploid

genome, while the black peak at low multiplicity shows that the assembly is highly complete

and that all reads are represented in the assembly.

Figure 4. Circos plot summarizing the nuclear genome annotation of *Coccomyxa viridis* SAG 216-4. From outside to inside the tracks display: GC content (over 1-kb windows), gene density (blue) and repetitive element density (red).

Figure 5. No synteny detected between related *Coccomyxa* species. (a) Dot plot of orthologous sequences in the genome assemblies of *C. viridis* SAG 216-4 and *C. subellipsoidea* C-169. Violet and blue dots represent orthologous sequences on same and opposite strands, respectively. Dot sizes does not correlate with the length of the sequences they represent, which were all < 1 kb. The width of each box corresponds to the length (bp) of the respective scaffold. (b) Dot plot of the genome assembly of *C. viridis* SAG216-4 against itself.

506 **Table 1.** Summary of bioinformatics tools used for genome assembly and annotation.

Assembly		Annotation	
Tool	Version	Tool	Version
Raven	v1.8.1	RepeatModeler	v2.0.3
Juicer	v2.0	TransposonUltimate	v1.0
BWA	v0.7.17-r1188	TEclass	v2.1.3
3d-dna	v180922	RepeatMasker	v4.1.2-p1
Juicebox	v1.11.08	HiSat2	v2.2.1
Minimap2	v2.24-r1122	Braker	v2.1.6
Samtools	v1.10	Gffread	v0.12.7
Integrative Genome Viewer	v2.11.2	SignalP	v6.0
Tapestry	v1.0.0	BUSCO	v5.3.2
Blobtools	v1.1.1	R	v4.2.0
BLAST	2.13.0+	Circilize	v0.4.14
Kmer Analysis Toolkit	V2.4.2	InterProScan	v5.61
Mummer	C3.23		

510 **Table 2.** Genome features of *C. viridis* SAG 216-4 including the mitochondrial and plastid genomes.

Assembly ID	*C. viridis* SAG 216-4 genome (including mitochondrial genome)
Total length (bp)	50,911,578
No. of contigs	27
No. of scaffolds	21
Longest scaffold (bp)	4,477,725
N50 (bp)	2,669,017
L50	8
GC content (%)	54.5

513 **Table 3** Annotation features of the *C. viridis* SAG 216-4 genome.

Genome annotation	
Repeat content (%)	8.85
Retrotransposons	2.4
DNA transposons	4.2
Simple repeats	2.25
No. gene models	13,557
Average gene length (bp)	3146
No. exons	122,978
Average no. exons per gene model	9
Average exon length (bp)	158
No. transcripts	14,024
Average no. transcripts/gene model	1
No. gene models <200 bp length	0
No. proteins with ≥1 PFAM domain	9205
No. proteins with signal peptide	962
BUSCO (chlorophyta_odb10)	**C: 98.6% [S: 82.5%, D: 16.1%], F: 0.1%, M: 1.3%, N: 1519**