Ankyrin B Promotes Developmental Spine Regulation in the Mouse Prefrontal Cortex

Authors: Kelsey E. Murphy[1,2], Bryce W. Duncan[1,2], Justin E. Sperringer[2], Erin Y. Zhang[2], Victoria A. Haberman[2], Elliott V. Wyatt[2], and Patricia F. Maness*[2]

Department of Biochemistry and Biophysics

University of North Carolina School of Medicine at Chapel Hill

Campus Box 7260

Chapel Hill, NC 27599

*Submitting and corresponding author Patricia F. Maness srclab@med.unc.edu

Orcid ID 0000-0003-248-5610

Phone 919-428-0942

Fax 919-966-3532

[1] co-first authors

Running Title: Developmental Spine Pruning through Ankyrin B

Keywords: Ankyrin, dendritic spines, cerebral cortex, cell adhesion molecules

Abstract

Postnatal regulation of dendritic spine formation and refinement in cortical pyramidal neurons is critical for excitatory/inhibitory balance in neocortical networks. Recent studies have identified a selective spine pruning mechanism in the mouse prefrontal cortex (PFC) mediated by class 3 Semaphorins and the L1-CAM cell adhesion molecules Neuron-glia related CAM (NrCAM), Close Homolog of L1 (CHL1), and L1. L1-CAMs bind Ankyrin B (AnkB), an actin-spectrin adaptor encoded by *Ankyrin2* (*ANK2*), a high confidence gene for autism spectrum disorder (ASD). In a new inducible mouse model (Nex1Cre-ERT2: *Ank2*flox: RCE), *Ank2* deletion in early postnatal pyramidal neurons increased spine density on apical dendrites in PFC layer 2/3 of homozygous and heterozygous *Ank2*-deficient mice. In contrast, *Ank2* deletion in adulthood had no effect on spine density. Sema3F-induced spine pruning was impaired in cortical neuron cultures from AnkB-null mice and was rescued by re-expression of the 220 kDa AnkB isoform but not 440 kDa AnkB. AnkB bound to NrCAM at a cytoplasmic domain motif (FIGQY[1231]), and mutation to FIGQH inhibited binding, impairing Sema3F-induced spine pruning in neuronal cultures. Identification of a novel function for AnkB in dendritic spine regulation provides insight into cortical circuit development, as well as potential molecular deficiencies in ASD.

Introduction

Developmental regulation of dendritic spine formation and remodeling in pyramidal neurons is necessary to achieve excitatory/inhibitory balance in neocortical circuits. During early stages of development, dendritic spines and excitatory synapses on

cortical pyramidal neurons are initially overproduced, then selectively remodeled by elimination mechanisms that are incompletely understood (Huttenlocher 1979; Alvarez *et al.* 2007; Petanjek *et al.* 2011). In the prefrontal cortex (PFC) appropriately balanced circuits are important for such vital functions as working memory, cognitive flexibility, and sociability. In Autism Spectrum Disorders (ASD) these behaviors are impaired, and elevated spine density has been reported in frontal cortical areas, which could potentially contribute to network hyper-excitability (Hutsler *et al.* 2010; Tang *et al.* 2014; Forrest *et al.* 2018).

Recent investigation into molecular mechanisms of spine elimination in the mouse neocortex demonstrated that secreted Semaphorins of class 3 (Sema3) prune distinct populations of dendritic spines during development and homeostatic scaling (Tran *et al.* 2009; Wang *et al.* 2017; Mohan *et al.* 2019a; Mohan *et al.* 2019b; Duncan *et al.* 2021b). As shown in Figure 1A, Sema3 ligands bind to heterotrimeric receptors comprising L1 cell adhesion molecules (L1-CAMs), Neuropilins (Npn1/2), and PlexinAs (PlexA1-4), to activate intracellular signaling resulting in spine pruning (Duncan *et al.* 2021a). Neuron-glial related CAM (NrCAM), Npn2, and PlexA3 constitute a receptor complex for Sema3F (Demyanenko *et al.* 2014; Mohan *et al.* 2019a), while Close Homolog of L1 (CHL1), Npn2, and PlexA4 form a receptor for Sema3B (Mohan *et al.* 2019b). Genetic knockout of NrCAM, CHL1 or L1 (Murphy *et al.* 2023a) in mice increased spine density on apical dendrites of cortical pyramidal neurons. Sema3F and Sema3B are secreted by neurons in an activity-dependent manner, consistent with the notion that less active or immature spines may be pruned by Sema3s released from active synaptic neighbors to refine cortical circuits in development (Wang *et al.* 2017; Mohan *et al.* 2019a).

All L1-CAMs bind the actin-spectrin adaptor protein Ankyrin B (AnkB) at a conserved cytoplasmic domain motif (FIGQY) (Bennett *et al.* 2009). Mice with a point mutation in the L1 Ankyrin binding site (FIGQH) display increased spine density in the prefrontal cortex (Murphy *et al.* 2023a) suggesting that AnkB may be required for spine pruning *in vivo*. This L1 mutation is a human pathological mutation associated with intellectual disability (Hortsch *et al.* 2014). AnkB is encoded by the high confidence ASD gene *ANK2*. Multiple *de novo* ASD variants in *ANK2* have been identified in the Simon Simplex Collection and Autism Sequencing Consortium (Iossifov *et al.* 2012; Willsey *et al.* 2013; De Rubeis *et al.* 2014; Iossifov *et al.* 2015; Wang *et al.* 2016; Satterstrom *et al.* 2019; Wang *et al.* 2020; Zhou *et al.* 2022). *ANK2* is subject to alternative splicing resulting in a number of AnkB isoforms including a principal isoform of 220 kDa, expressed in brain and other tissues, and a neuron-specific 440 kDa isoform (Jenkins *et al.* 2015). Over half of the 23 distinct variants in *ANK2* are predicted to be loss-of-function mutations affecting both AnkB isoforms.

Knowledge of AnkB function in developing brain is limited, because germline deletion of AnkB in mouse models is lethal in early postnatal life (Scotland *et al.* 1998). To interrogate a potential role for AnkB in developmental spine regulation and to provide a model for loss-of-function ASD mutations, we generated a tamoxifen-inducible *Ank2* mouse line (Nex1Cre-ERT2: *Ank2^flox^*: RCE), which can delete all AnkB isoforms from postmitotic pyramidal neurons postnatally, circumventing lethality associated with germline deletion. Here we show that homozygous or heterozygous *Ank2* deletion in Nex1Cre-ERT2: *Ank2^flox^*: RCE mice during active stages of spine remodeling increased

spine density and excitatory synaptic puncta on apical dendrites of prefrontal cortical

pyramidal neurons, whereas *Ank2* deletion in adulthood had no effect.

AnkB localized to spines and dendrites of cortical pyramidal cells and was

enriched in synaptoneurosomes from mouse brain. The Ankyrin binding motif in NrCAM

was required for association with AnkB-220, as shown by mutation of tyrosine to

histidine mutation in the NrCAM FIGQY[1231] sequence. Notably, cortical neuron cultures

from AnkB-null or NrCAM-null mice were deficient in Sema3F-induced spine pruning,

and re-expression of AnkB-220 but not AnkB-440 in *Ank2*-null neurons restored spine

pruning *in vitro*. These results suggest that AnkB loss or haploinsufficiency could alter

developmental spine regulation and have consequences on cortical circuits relevant to

ASD.

Materials and Methods

Generation and Tamoxifen Induction of Ank2 Conditional Mutant Mice

We generated an inducible *Ank2* mouse line Nex1Cre-ERT2: *Ank2^flox^*: EGFP^flox^

(C57Bl/6) to delete *Ank2* from post-mitotic, postmigratory cortical pyramidal neurons.

Nex1Cre-ERT2 mice express a modified Cre recombinase fused to the human estrogen

receptor, which is under control of endogenous Nex1 regulatory sequences and

activated by tamoxifen (TMX) (Agarwal *et al.* 2012). The Nex1 promoter drives the

TMX-inducible Cre-ERT2 recombinase only in postmitotic pyramidal neurons and not

interneurons, oligodendroglia, astrocytes, or non-neural cells (Agarwal *et al.* 2012).

Nex1-Cre mice have been extensively employed to target postmitotic pyramidal neurons

with high specificity (Golonzhka *et al.* 2015; Fame *et al.* 2016; Mohan *et al.* 2019a;

Miller *et al.* 2021). For our approach Nex1Cre-ERT2 mice were intercrossed with RCE:

loxP reporter mice (from Gordon Fishell), in which induction of Cre-ERT2 recombines a "floxed stop cassette" enabling expression of EGFP in Cre-induced neurons (Sousa *et al.* 2009). The resulting mice were intercrossed with $Ank2^{flox}$ mice (from Peter Mohler), in which recombination deletes exon 24, which has 73 bp of coding sequence, resulting in a frameshift that places a stop codon in exon 25 (Smith *et al.* 2015; Roberts *et al.* 2019).

For *in vivo* induction, TMX (Sigma-Aldrich, #10540-29-1) was dissolved (10 mg/ml) in sunflower seed oil and administered by intraperitoneal injection at 100 mg/kg body weight every 24 hours for 4 (P10-P13) or 10 (P50-P60) consecutive days. Mice were analyzed at P50 (young adult) or P80 (older adult), respectively. Postnatal TMX induction in Nex1-CreERT2 mice has been shown to achieve cell-specific targeting of postmitotic cortical and hippocampal pyramidal neurons (Agarwal *et al.* 2012). For breeding Nex1Cre-ERT2$^{+/+}$: $Ank2^{flox/+}$: RCE$^{+/+}$ mice were crossed with $Ank2^{flox}$ mice, yielding expected ratios of ~50% $Ank2^{flox}$ homozygotes (Nex1Cre-ERT2$^{+/-}$: $Ank2^{flox/flox}$: RCE$^{+/-}$) and 50% $Ank2^{flox/+}$ heterozygotes (Nex1Cre-ERT2$^{+/-}$: $Ank2^{flox/+}$: RCE$^{+/-}$). Mice with wild type (WT) $Ank2$ alleles (Nex1Cre-ERT2$^{+/-}$: $Ank2^{+/+}$: RCE$^{+/-}$) were produced by intercrossing heterozygotes. Each allele was genotyped by PCR from genomic DNA.

Other mouse strains included wild type (WT) mice (JAX:000664) and *NrCAM*-null mice on the C57Bl/6 background (Sakurai *et al.* 2001; Sakurai *et al.* 2006). *Ank2*-null mice on a hybrid background Sv129/C57Bl were produced by intercrossing heterozygotes of each strain as described (Scotland *et al.* 1998). Mice were maintained according to policies of the University of North Carolina Institutional Animal Care and

Use Committee (IACUC; AAALAC Institutional Number: #329; ID# 18-073, 21-039) in accordance with NIH guidelines.

<u>Immunoreagents and Immunolabeling</u>

Polyclonal antibodies were directed against the following proteins: NrCAM (Abcam #24344, RRID:AB_448024 or R&D Systems AF8538), Npn2 (R&D Systems, AF567), and GFP (Abcam #13970, RRID:AB_300798). Monoclonal antibodies were directed against AnkB (ThermoFisher, #33-3700, RRID:AB_2533115; Biolegend, #821403, RRID:AB_2728536), vinculin (ThermoFisher, #MA5-11690), α-tubulin (Abcam, ab7291), actin (Sigma-Aldrich, MAB1501), and parvalbumin (Neuromab L11413). Sheep anti-AnkB antibodies were a gift from Paul Jenkins (University of Michigan) (Qu *et al.* 2016). Also used were antibodies to vesicular glutamate transporter-1 (vGlut1, #821301, Biolegend) and Homer-1 (#160002, Synaptic Systems). Non-immune rabbit IgG (NIg), HRP- and AlexaFluor 488, 555, and 594-conjugated secondary antibodies were from Jackson Immunoresearch.

<u>Brain Fixation, Immunostaining, and Imaging</u>

Mice at indicated postnatal ages were anesthetized with 2.5% Avertin, perfused transcardially with 4% paraformaldehyde (PFA)/PBS and processed for staining as described (Demyanenko *et al.* 1999). Brains were postfixed in 4% PFA overnight at 4°C, followed by 0.02% PBS-azide, then sectioned coronally on a vibratome (60 μm) and mounted on glass slides. Sections were permeabilized in 0.3% Triton X-100 and blocked in 10% normal donkey serum in PBS for 3 hours at room temperature, then incubated with chicken anti-GFP antibody (1:250) and/or mouse monoclonal anti-AnkB antibody 33-3700 (1:250) for 48 hours at 4°C. After washing, sections were incubated

with anti-chicken Alexa Fluor 488 and/or anti-mouse Alexa Fluor 555 secondary antibodies (1:250) for 2 hours before mounting with Prolong Glass (Thermofisher). Confocal z-stacks were obtained by imaging on a Zeiss LSM 700 microscope in the UNC Microscopy Services Laboratory with an EC Plan Neofluar 40x objective using a 1.3 N.A. oil lens (0.3 µm optical sections). Images were acquired using a pinhole size of 1 AU. Zoom was adjusted to obtain pixel sizes of 0.13-0.14 µm. For AnkB localization in immunostained brain sections, pixel intensities were quantified from confocal maximum intensity projections in FIJI. Regions of interest (ROIs) were selected using a grid, and mean grey values obtained using the FIJI analysis menu to yield pixel intensities per unit area. Pixel intensities of negative control staining with secondary antibodies alone were obtained similarly to determine background fluorescence, which was subtracted from mean pixel intensities of AnkB immunofluorescence. The resulting mean pixel intensities per unit area for each genotype were compared by the Mann-Whitney test (two tailed, unequal variance) for significant differences ($p < 0.05$). AnkB localization in cortical neurons in culture (DIV14) was analyzed by indirect immunofluorescence staining using sheep anti-AnkB (Qu *et al.* 2016) and AlexaFluor secondary antibodies (Thermofisher #A-11039, RRID: AB-2534096). Stimulated emission depletion microscopy (STED) was used for AnkB localization in cortical neurons in culture by immunofluorescence staining with mouse monoclonal anti-AnkB antibody 33-3700 and AF-594-conjugated secondary antibodies (Abcam, #ab245992). STED images were acquired using a Leica SP8 STED confocal microscope with Lightning deconvolution in the UNC Neuroscience Research Center.

Excitatory synaptic puncta on spines of EGFP-expressing pyramidal neurons in Nex1Cre-ERT2:RCE and Nex1Cre-ERT2: *Ank2* F/F: RCE mice were analyzed confocally in coronal brain sections through the PFC in layer 2/3 after immunofluorescence staining of presynaptic vesicular glutamate transporter-1 (vGlut1) and postsynaptic Homer-1 with AlexaFluor-conjugated secondary antibodies. Quantification of excitatory synaptic puncta was performed blind to observer in single optical confocal sections (0.3 µm) using Imaris software. The mean number of puncta per 10 µm dendrite length was calculated. Puncta densities were compared between genotypes by the Mann-Whitney test (two tailed, unequal variance; $p < 0.05$).

Analysis of Dendritic Spine Density and Morphology

Spine densities of layer 2/3 pyramidal neurons in the PFC (primary cingulate area) were measured using Neurolucida software (MBF Bioscience) as described (Demyanenko *et al.* 2014; Mohan *et al.* 2019a; Mohan *et al.* 2019b). Briefly, EGFP-labeled spines were traced and quantified blind to the observer on 30 µm segments of the first branch of apical or basal dendrites from maximum intensity projections. Images were compared to confocal z-stacks to verify that spines emerged from dendritic segments. Mean spine number per 10 µm of dendritic length (density) was calculated. Neurons (n = 10-21) were analyzed for each genotype on apical or basal dendrites. For production of 3D reconstructions dendritic z-stacks were deconvolved using AutoQuant 3 software (Media Cybernetics) with default deconvolution settings in Imaris (Bitplane). Mean spine densities/10 µM dendrite length ± SEM were analyzed by 2-factor ANOVA with Tukey's posthoc testing ($p<0.05$).

Once spines densities were determined, spine morphologies of the same layer 2/3 pyramidal neurons in the PFC (primary cingulate area, P50) were manually scored on constructed Z-projections as mushroom, stubby, or thin (filopodial) on apical dendrites. Spines were measured in three categories using the straight-line tool in FIJI: Spine length, head width, and neck width. Spine length was defined as the total length of the spine from its base at the dendrite to its terminus. Head width was defined as the width of a distinct head, or, when lacking a distinct head, the measurement of the widest point in the terminal third of the spine. Neck width was defined as the width of the distinct neck of the spine, or, when a distinct neck was not discernible, the widest point in the basal third of the spine. Based on these three measurements, mushroom spines were defined by a spine length >0.5 μm, and a head width to neck width ratio >1.5. Stubby spines were defined by a length <1.0 μm and a ratio <1.5, while thin spines were defined by a length >1.0 μm and a ratio >1.0. To compare spine morphologies between genotypes of mice in PFC (layer 2/3), spine morphological types were calculated as a percentage of the total spine number. P-values were computed as indicated in the legend to Figure 2.

Dendritic arborization was measured by Sholl analysis of processes crossing concentric rings centered on the soma of EGFP-labeled pyramidal neurons in confocal z-stacks (20x). The center was defined as the middle of the cell body at a soma detector sensitivity of 1.5 μm, and the automatic tracing mode of Neurolucida was used to seed and trace dendritic arbors. Images in DAT format were subjected to Sholl analysis using Neurolucida Explorer (MBF Bioscience) with a starting radius of 10 μm and radius increments of 10 μm ending at 300 μm. Apical and basal dendrites were not able to be

scored separately due to overlap at increasing distances from soma. Data were analyzed using two-factor ANOVA (p<0.05).

The midline widths of the corpus callosum and anterior commissure in TMX-induced Nex1Cre-ERT2:RCE and Nex1Cre-ERT2: *Ank2 flox*: RCE mouse brains were measured in serial coronal brain sections (60 μm) after immunostaining with antibodies against GFP. Confocal images were captured and analyzed in FIJI. The dorsoventral width of each tract was measured at the midline of each section using the straight-line tool in FIJI, and mean widths calculated. Means (± SEM) were compared by the Mann-Whitney test (two-tailed, unequal variance).

Cortical Neuron Cultures and Sema3F-Induced Spine Retraction

Cortical neurons were isolated from brains of WT, *NrCAM*-null, or *Ank2*-null embryos (E15.5) and plated onto Lab-Tek II chamber slides (1.5 x 10^6 cells/well) coated with poly-D-lysine and laminin as described (Mohan *et al.* 2019a). For *Ank2*-null primary cultures, rapid genotyping of embryos was done while cortices of embryos remained on ice in Hibernate-E media containing B27 Plus supplement. AraC was added at 5 days *in vitro* (DIV5) to limit the growth of glia and fibroblasts, and media was changed on DIV7. At DIV11, cells were transfected with pCAG-IRES-mEGFP with or without additional plasmids (AnkB-220 (pEGFP-N1-ΔEGFP) or AnkB-440 (pAB)) using Lipofectamine 2000 (Mohan *et al.* 2019a; Duncan *et al.* 2021a). At DIV 14, cultures were treated with purified Fc from human IgG (Abcam #ab90285) or recombinant mouse Sema3F-Fc fusion protein (R&D Systems #3237-S3) at 5 nM for 30 minutes. Cultures were fixed with 4% PFA, quenched with 0.1M glycine, permeabilized with 0.1% Triton X-100, and blocked with 10% donkey serum. Cells were incubated with chicken

anti-GFP primary antibody and AlexaFluor AF488-conjugated goat anti-chicken secondary antibody (1:500), washed, and mounted. At least 10 images of apical dendrites of labeled pyramidal neurons were captured per condition. Apical dendrites were differentiated from basal dendrites by cell morphology. Pyramidal neurons extend a single prominent, thicker apical dendrite from the apex of the pyramidal-shaped soma, and multiple, thinner basal dendrites from the base of the soma (Spruston 2008). Any ambiguous dendrites were not analyzed. Confocal z-stacks were obtained using 0.2 μm optical sections of field size 64.02 x 64.02 μm, using a 40x oil objective and 2.4x digital zoom, and subjected to deconvolution. Spines from maximum intensity projections were traced and scored using Neurolucida. Mean spine densities (number/10 μm ± SEM) were calculated and compared by two-factor ANOVA with Tukey's posthoc testing (p<0.05).

Immunoprecipitation and immunoblotting

Protein-protein interactions were assessed by co-immunoprecipitation and immunoblotted from cortical lysates of mouse forebrain, synaptoneurosomes, and transfected HEK293T cells. For preparation of mouse cortical lysates, forebrains were dissociated and subjected to Dounce homogenization for 20 strokes in RIPA buffer. Homogenates were incubated for 15 minutes on ice, then centrifuged at 16,000 x g for 10 minutes. The supernatant was retained, and protein concentration determined by BCA. Synaptoneurosomes were isolated as described (Villasana *et al.* 2006). Briefly, WT mice (P32) were anesthetized, decapitated and cortices were isolated. Following Dounce homogenization in Triton lysis buffer (20 mM Tris, pH 7.0, 150mM NaCl, 5 mM EDTA, 1 mM EGTA, 1% Triton X-100, 10 mM NaF, 1x Protease Inhibitor Cocktail),

homogenates were sonicated, filtered, and centrifuged at 1000 x g for 10 minutes at 4°C. Pellets were resuspended in Triton lysis buffer, nutated, and centrifuged at 16,000 x g for 10 minutes at 4°C. Supernatants were retained as the synaptoneurosome fraction. Protein concentrations were determined by BCA. Postsynaptic density protein 95 (PSD95), AnkB, α-tubulin, and actin levels were quantified by western blotting in the synaptoneurosome fraction compared to the filtered homogenate and 1000xg supernatant fraction as described previously(Villasana *et al.* 2006; Demyanenko *et al.* 2014; Murphy *et al.* 2023b). HEK293T cells were grown in DMEM/ gentamicin/kanamycin/10% FBS in a humidified incubator with 5% CO_2. Cells were seeded at 2×10^6 cells/100mm dish the day before transfection. Plasmids were transfected with Lipofectamine 2000 in Opti-MEM. Media was changed to DMEM after 18 h, and cells were lysed and collected 48 h post-transfection. Cells were harvested in Brij98 lysis buffer (1% Brij98, 10 mM Tris-Cl pH 7.0, 150 mM NaCl, 1mM EDTA, 1mM EGTA, 10 mM NaF, protease inhibitors (SigmaAldrich #P8340).

For immunoprecipitation, lysates of mouse forebrain (1 mg) or HEK293T cells (0.5 mg) were precleared for 30 minutes at 4°C using Protein A/G Sepharose beads (ThermoFisher). Precleared lysates (equal amounts of protein) were incubated with 3 µg rabbit polyclonal antibody to NrCAM (Abcam #24344) or nonimmune IgG (NIg) for 2 hr on ice. Protein A/G Sepharose beads were added for an additional 30 min with nutation at 4°C before washing with RIPA or Triton lysis buffer (synaptoneurosomes). Beads were washed 4 times, then immunoprecipitated proteins were eluted from the beads by boiling in SDS-PAGE sample buffer. Samples (50 µg) were subjected to SDS-PAGE (6%) and transferred to nitrocellulose. Membranes were blocked in TBST containing 5%

nonfat dried milk and incubated overnight with primary antibodies (1:1000), washed, and incubated with HRP-secondary antibodies (1:5000) for 1 h. Antibodies were diluted in 5% milk/Tris buffered saline/0.1% Tween-20 (TBST). For detection of AnkB-440, a 3.5-17.5% gradient gel was used, and AnkB antibodies were diluted in 5% BSA/TBST. Blots were developed using Western Bright ECL Substrate (Advansta) and exposed to film for times yielding a linear response of signal. Membranes were stripped and reprobed with rabbit anti-NrCAM antibodies (R&D Systems AF8538). Bands were quantified on densitometric scans in FIJI. The ratio of AnkB to NrCAM was determined from 3 replicate experiments to obtain mean and SEM values.

All experiments were designed to provide sufficient power (80–90%) to discriminate significant differences ($p<0.05$) in means (± SEM) between independent controls and experimental subjects as described (Dupont and Plummer, 1990). The type I error probability associated with tests of the null hypothesis was set at 0.05. Full details of the experimental design, sample sizes, and number of repetitions for each experiment are reported in the figure legends.

Results

Postnatal Deletion of Ankyrin B from Prefrontal Pyramidal Neurons Increases Spine Density and Immature Spine Morphology

To identify consequences of postnatal AnkB deletion on dendritic spine remodeling in cortical pyramidal neurons, we generated a TMX-inducible mouse line, Nex1Cre-ERT2: $Ank2^{flox}$: RCE (C57Bl/6). In Nex1Cre-ERT2:RCE mice, loxP recombination is activated by Cre-ERT2 recombinase under control of the Nex1 promoter in postmitotic pyramidal neurons treated with TMX (Agarwal *et al.* 2012;

Mohan *et al.* 2019a). Reporter EGFP expression is induced by loxP recombination in the same cells (Sousa *et al.* 2009). To achieve *Ank2* deletion during the major postnatal period of spine formation and remodeling (Culotta *et al.* 2020), mice were given daily TMX injections intraperitoneally from P10 to P13 as described (Agarwal *et al.* 2012). During the P14-P21 time period dendritic spines are overproduced and pruned to appropriate levels, and spine remodeling decreases substantially from P19-P34 as mature circuits are established (Trachtenberg *et al.* 2002; Holtmaat *et al.* 2005). *Ank2* deletion at these postnatal stages were chosen to overcome early postnatal lethality of germline *Ank2* deletion (Scotland *et al.* 1998), thus circumventing effects on premigratory pyramidal cells and early neuronal precursors.

We focused on layer 2/3 pyramidal neurons of the PFC, because of their importance in social and cognitive circuits, which are affected in ASD (Yizhar 2012; Kroon *et al.* 2019). Treatment with TMX at P10-P13 efficiently upregulated EGFP expression in pyramidal neurons of young adult Nex1Cre-ERT2: *Ank2$^{flox/flox}$*: RCE mice (termed Ank2 F/F) in the primary cingulate area (Cg1) of the PFC at P50 (Fig. 1 B, C). EGFP-expressing neurons were distributed across the cortical laminae in induced but not uninduced Ank2 F/F mice, indicating that Cre recombinase expression was tightly regulated. Prefrontal cortical regions in mice lack a canonical layer 4 (L4) as well as gene expression characterizing L4 neurons (Tasic *et al.* 2016; Wang *et al.* 2018; Anastasiades *et al.* 2021).

Figure 1

Induced *Ank2* F/F mice were healthy and viable even as older adults (P150).

Decreased expression of AnkB protein was also evident in the PFC of TMX-induced

Ank2 F/F mice, as shown by diminished AnkB immunofluorescence staining compared

to Nex1Cre: *Ank2*[+/+]: RCE mice (termed *Ank2* +/+), using pan-AnkB antibodies that

recognized both AnkB-220 and -440 principal isoforms (Fig. 1D). Immunofluorescence

was quantified by measuring mean pixel intensities per unit area in single optical

confocal sections and found to be significantly lower in PFC layer 2/3 of *Ank2* F/F

compared to *Ank2* +/+ mice (Fig. 1D, p = 0.03). Negative control staining with

secondary AlexaFluor secondary antibodies alone indicated that AnkB immunostaining

in the *Ank2* F/F cortex was substantially greater than background (Fig. 1D, control).

Residual AnkB immunofluorescence in the *Ank2* F/F cortex was seen adjacent to

EGFP-negative soma (arrows), and in the neuropil between soma. This

immunoreactivity likely derived from interneuron processes, glia, or axons from neuronal

cells whose soma are located in subcortical areas, such as the amygdala and ventral

tegmental area, which do not express Nex1. This is supported by the expression of

Ank2 transcripts in pyramidal cells and interneurons, as well as in astrocytes and

oligodendroglia of the adult mouse cortex (Allen Brain Map Transcriptomics Explorer

(https://celltypes/brain-map.org/maseq).

Western blotting of cortical lysates from *Ank2* F/F mice (P50) confirmed that

there was decreased protein expression of both AnkB-220 and AnkB-440 isoforms in

Ank2 F/F forebrain compared to uninduced *Ank2* F/F following TMX induction (Fig. 1E).

Densitometric measurements of replicate blots showed that both isoforms decreased

significantly upon induction relative to the vinculin control, but that AnkB-440 decreased

to a greater extent than AnkB-220. This was probably because neuron-specific AnkB-

440 is largely under Nex1 control, whereas AnkB-220 expression in interneurons and

glia is Nex1-independent. Note also that AnkB-220 and -440 required different

electrophoresis and transfer conditions for Western blotting due to their large difference

in molecular weight.

Basket interneurons account for ~ 50% of cortical interneurons, of which

parvalbumin-expressing basket cells constitute the majority (Markram *et al.* 2004). To

validate the specificity of Nex1Cre-ERT2 induced recombination, we carried out triple

immunostaining for parvalbumin, AnkB, and EGFP in PFC layer 2/3 of TMX induced

Ank2 F/F and *Ank2* +/+ mice (P50) (Fig. 1F). Merged confocal images (a,e) showed

overall decreased AnkB staining in *Ank2* F/F compared to *Ank2* +/+ cortex in agreement

with Fig. 1D. Parvalbumin+ interneurons in *Ank2* +/+ and *Ank2* F/F cortex (blue; b,f,

arrows) were distinct from EGFP-expressing pyramidal neurons (green; c,g), but did

exhibit AnkB immunoreactivity (d,h, arrows). These results indicated that *Ank2* was not

deleted in parvalbumin+ basket cells upon TMX induction of *Ank2* F/F mice.

There was no major alteration in the size of the corpus callosum or anterior

commissure, major tracts comprising contralaterally crossing pyramidal cell axons. The

mean width of EGFP-labeled axonal tracts were not different between genotypes as

measured in serial coronal sections of *Ank2* F/F and *Ank2* +/+ brains (Fig. 1G). Gross

neuroanatomical defects including the size of brain ventricles (lateral and third) and

cortical thickness were also not observed in *Ank2* F/F mice (not shown).

To investigate an *in vivo* role of AnkB in dendritic spine regulation in the postnatally

developing PFC and effect of gene dosage, recombination was induced in juvenile *Ank2*

F/F, *Ank2* F/+, and *Ank2* +/+ mice at P10-P13 and analyzed in young adults at P50

(Laviola *et al.* 2003). During this time frame dendritic spines are actively overproduced

and spine turnover occurs concurrently, followed by a decrease in spine number from

P19-P34 as mature circuits are stabilized (Trachtenberg *et al.* 2002; Holtmaat *et al.*

2005). A significant increase in the density of EGFP-labeled spines on apical dendrites

of pyramidal neurons in PFC layer 2/3 was observed in homozygous *Ank2* F/F mice

compared to *Ank2* +/+ mice (Fig. 2A, B; 88.2% increase, *$p<0.0001$). Homozygous

Ank2 mice reflect AnkB loss-of-function, whereas heterozygotes may reflect haplo-

insufficiency associated with neurological diseases (Lim *et al.* 2013). Accordingly, we

compared spine density in layer 2/3 pyramidal neurons of *Ank2* heterozygotes (P50)

induced at P10-P13. *Ank2* heterozygotes displayed a significant increase in spine

density on apical dendrites compared to *Ank2* +/+ mice (36.5% increase, *$p=0.02$) and a

significant decrease in spine density compared to *Ank2* F/F homozygotes (*$p=0.001$,

Fig. 2A, B), consistent with an intermediate phenotype. In contrast spine densities on

basal dendrites of layer 2/3 pyramidal cells were not significantly different across any of

the genotypes (*Ank2* +/+, F/+, F/F mice; Fig. 2B). Because *Ank2* remains deleted after

recombination, effects seen at P50 could result from changes at any point during this

postnatal period. It is worth noting that the presence of spines in *Ank2*-deficient mice

indicates that AnkB is not required during this period to maintain spine integrity. This

may be mediated by AnkG, a different gene product localized to adult spine

nanodomains, where it promotes spine maintenance and plasticity (Smith *et al.* 2014;

Yoon *et al.* 2020).

     To determine if deletion of *Ank2* in adulthood alters spine regulation that is

associated with homeostasis, *Ank2* F/F, *Ank2* F/+, and *Ank2* +/+ mice were TMX-

treated in young adults at P50-P60 and analyzed for spine density in older adults at

P80. There were no significant differences in spine density on apical dendrites of layer

2/3 pyramidal neurons in *Ank2* +/+, *Ank2* F/+, *Ank2* F/F PFC at P80 (Fig. 2A, C). These

results suggested that spine regulation in the adult PFC is not substantially altered by

depletion of AnkB in cortical pyramidal neurons.

Figure 2

Dendritic spines acquire diverse morphologies (mushroom, stubby,

thin/filopodial), which have different functional properties and are classified based on

size and shape (Peters *et al.* 1970). Spine morphologies are dynamically

interchangeable, comprising a continuum from immature spines (thin and stubby), which

can have immature synaptic contacts associated with juvenile plasticity (Bourne *et al.*

2007), to mushroom spines with mature synapses (Bhatt *et al.* 2009; Holtmaat *et al.*

2009; Berry *et al.* 2017). Several studies have shown that immature spine morphology

is associated with Fragile X syndrome (Phillips *et al.* 2015) and intellectual disability

(Forrest *et al.* 2018). In *Ank2* F/F mice TMX-induced deletion at P10-13 increased the percent of stubby spines and decreased the percent of mushroom spines on apical dendrites of layer 2/3 pyramidal neurons in the PFC at P53 (Fig. 2D). These differences between *Ank2* +/+ and *Ank2* F/F were statistically significant by ANOVA multiple comparisons (p < 0.01). The percent of thin spines was not significantly different between genotypes. These findings were consistent with the hypothesis of preferential elimination of immature spines mediated by AnkB in normal postnatal development. However, due to the dynamic transformation among morphologies it is also possible that AnkB also promotes elimination of other spine types. Arborization of neuronal processes (primarily dendrites) was then evaluated by Sholl analysis of PFC layer 2/3 pyramidal cells at P50 following homozygous *Ank2* deletion at P10-P13. No differences were observed in dendritic branching of EGF-labeled *Ank2* F/F neurons compared to *Ank2* +/+ cells (Fig. 2E). Taken together, these results support a role for AnkB in constraining spine density on apical dendrites of layer 2/3 pyramidal neurons in the developing PFC.

To assess whether the observed increase in spine density of *Ank2* F/F pyramidal cells resulted in a corresponding increase in excitatory synapses, synaptic puncta were identified on spines by juxtaposed immunofluorescence staining of the presynaptic marker vesicular glutamate transporter 1 (vGlut1) and postsynaptic marker Homer-1 on EGFP-expressing apical dendrites in PFC layer 2/3 in brain sections of *Ank2* F/F and *Ank2* +/+ mice (P50). Juxtaposed VGlut1 and Homer-1 immunofluorescence was observed on EGFP-labeled spines of both genotypes (Fig. 2F). Quantification of dual-labeled puncta on spines revealed a significant increase in *Ank2* F/F compared to *Ank2*

+/+ neurons (Fig. 2G). These results were consistent with an increase in excitatory synapses, rather than orphan synapses lacking pre- or postsynaptic specializations. Although we have not evaluated synaptic function, immature spines with stubby or thin, but not filopodial morphology, have been shown to contain functional synapses, albeit with a low AMPA to NMDA ratio (Berry *et al.* 2017).

AnkyrinB-220 Mediates Sema3F-Induced Spine Pruning

To assess a potential role for AnkB in Sema3F-induced spine pruning, and to evaluate AnkB isoform specificity in the response, we developed a spine retraction assay using *Ank2*-null cortical neuron cultures in which all *Ank2* isoforms are deleted. Most *Ank2*-null mice on a C57Bl/6 background die shortly after birth (Scotland *et al.* 1998), and cortical neuron cultures from these embryos (E15.5) degenerated at approximately DIV6, prior to significant spine elaboration. We found that *Ank2*-null mice on a first-generation hybrid background C57Bl/6/SV129 survived to at least P5, and cortical neuron cultures from *Ank2*-null embryos (E15.5) were viable through DIV14, when spines were abundant. Intercrossing *Ank2* heterozygotes from each background strain produced WT and *Ank2*-null hybrid embryos in the same litters. Embryos (E15.5) were subjected to rapid genotyping then cortical neuron cultures were plated. Cortical cultures from WT and *Ank2*-null embryos were transfected on DIV11 with reporter plasmid pCAG-IRES-EGFP, then treated with Sema3F-Fc or Fc (5 nM, 30 min) on DIV14. After fixation and immunostaining for EGFP, spine density was quantified on apical dendrites of EGFP-expressing neurons. WT and Ank2-null neurons in control cultures treated with Fc exhibited equivalent spine densities, consistent with the interpretation that AnkB was not required for spine maintenance (Fig. 3A, B). Notably,

Sema3F-Fc decreased spine density in WT but not Ank2-null neurons, indicating that AnkB was required for the Sema3F pruning response (Fig. 3A, B). Spine retraction in response to Sema3F-Fc in WT neurons was incomplete, since only the subpopulation of spines expressing NrCAM responds to Sema3F, while CHL1-expressing neurons respond to Sema3B (Mohan *et al.* 2019b). Additionally, there was no difference in spine density in control Fc-treated WT and *Ank2*-null neurons (Fig. 3B), since the cultures *in vitro* are not exposed to semaphorins, or other factors present *in vivo*.

To determine if AnkB-220 or -440 isoforms mediated Sema3F-induced spine retraction, the ability of each isoform to rescue spine pruning was assayed after transfection of the relevant cDNAs into AnkB-null neurons. Plasmids expressing AnkB-220 or AnkB-440 under control of the CMV enhancer/promoter were co-transfected on DIV11 into *Ank2*-null neurons with pCAG-IRES-EGFP, then treated on DIV14 with Sema3F-Fc or Fc. Expression of AnkB-220 effectively rescued Sema3F-induced spine retraction to the same extent as in WT neurons (Fig. 3B). In contrast, expression of AnkB-440 in AnkB-null neurons did not rescue Sema3F-induced spine retraction (Fig. 3B). These results supported the conclusion that AnkB-220 specifically mediates Sema3F-induced spine retraction *in vitro*.

Figure 3

To examine AnkB localization in neurons, immunostaining was carried out on WT

cortical neuron cultures (DIV14) transfected for EGFP expression. Immunolabeling with

AnkB antibodies recognizing both isoforms (Qu *et al.* 2016) showed AnkB localized to

apical and basal dendrites as well as axons and soma in confocal images (Fig. 3C).

AnkB immunostaining was evident on spines of mushroom (mr), filopodial (fp), and

stubby (st) morphology (Fig. 3C, d arrows). Higher magnification confocal (g) and super-

resolution STED microscopy (h) showed AnkB immunofluorescence in spine head

domains (arrowheads) and adjacent dendritic segments (Fig. 3C). The presence of

AnkB staining in diverse cellular compartments of neuronal cultures is consistent with previous work using pan-AnkB antibodies and AnkB-440 specific antibodies concluding that AnkB-220 preferentially localizes to the somato-dendritic domain of neurons (Stevens *et al.* 2021), whereas AnkB-440 is enriched in premyelinated axons (Kunimoto *et al.* 1991; Kunimoto 1995; Yang *et al.* 2019). NrCAM is also present in all subcellular compartments of cortical neurons in culture (Demyanenko *et al.* 2014), although Npn2 is specifically localized to apical dendrites (Tran *et al.* 2009).

<u>The NrCAM Ankyrin Binding Motif is Required for Sema3F-Induced Spine Pruning</u>

L1-CAMs bear a conserved cytoplasmic domain containing a FIG(Q/A)Y motif that reversibly binds Ankyrin (Bennett *et al.* 2009). Phosphorylation of the tyrosine residue in the FIGQY motif can be achieved by receptor tyrosine kinases that are activated by NGF, bFGF (Garver *et al.* 1997) or EphrinB (Dai *et al.* 2013), and this phosphorylation reverses Ankyrin binding to L1-CAMs (Bennett *et al.* 2009). A charge reversal mutation of L1-FIGQY to FIGQH prevents Ankyrin association, and is a pathological mutation in the human L1 syndrome of intellectual disability/aphasia (Hortsch *et al.* 2014). In a knock-in mouse mutant expressing L1-FIGQH axonal connectivity is altered, as demonstrated by aberrant topographic mapping of retinocollicular axons (Buhusi *et al.* 2008; Dai *et al.* 2012). The L1-FIGQH mutation also suppresses axon branching (Yang *et al.* 2019) and perisomatic innervation by interneurons (Tai *et al.* 2019). However, a role for Ankyrin association with L1-CAMs in spine elimination has not been demonstrated.

Structure-function studies showed that NrCAM binding to Npn2 increases binding between Npn2 and PlexA3, necessary for Sema3F-induced spine pruning (Duncan *et*

al. 2021a). To investigate whether the Ankyrin binding motif FIGQY in the NrCAM

cytoplasmic domain also promotes Sema3F-induced spine pruning, we analyzed the

effect of the NrCAM FIGQY[1231]H mutation on Sema3F-induced spine retraction in

neuronal cultures. NrCAM-null neurons in culture are refractory to Sema3F-induced

spine retraction but re-expression of WT NrCAM rescues the response (Mohan *et al.*

2019a). In the present study NrCAM-null neurons were transfected with pCAG-IRES-

EGFP plasmids encoding WT NrCAM or NrCAM-FIGQY[1231]H. Neuronal cultures were

then treated with Sema3F-Fc or control Fc proteins (5 nM, 30 min) as described (Mohan

et al. 2019a). Spine density on apical dendrites of EGFP-expressing neurons was

measured after immunostaining for EGFP. Sema3F-Fc promoted spine retraction in WT

neurons but not in NrCAM-null cortical neurons, and re-expression of WT NrCAM

restored the responsiveness to Sema3F-Fc, as reported (Mohan *et al.* 2019a) (Fig. 3D,

E). In contrast, NrCAM-null neurons transfected with the NrCAM FIGQY[1231]H mutant

failed to respond to Sema3F-Fc and did not show a decrease in spine density (Fig. 3D,

E). These results indicated that the Ankyrin binding site in the NrCAM cytoplasmic

domain is a determinant of Sema3F-induced spine retraction.

Ankyrin B Association with NrCAM

AnkB-220 and -440 isoforms are both expressed in mouse brain (Jenkins *et al.*

2015; Stevens *et al.* 2021), and share a membrane binding domain that binds L1-CAMs

and certain ion channels (Chen *et al.* 2017). Because of the selective ability of AnkB-

220 to rescue Sema3F-induced spine pruning in AnkB-null cortical neurons, we

examined the expression of AnkB-220 at different postnatal and adult stages in mouse

cortex by Western blotting of forebrain lysates (equal protein). AnkB-220 was expressed

at approximately equivalent levels in postnatal (P22, P34), young adult (P50) and older

adult (P105) stages (Fig. 4A). NrCAM (130 kDa) also showed a relatively uniform

expression pattern at these stages. (Fig. 4A). To examine the association of AnkB-220

with NrCAM, NrCAM was immunoprecipitated from cortex lysates (P22-P105, equal

protein), and immune complexes immunoblotted for AnkB-220 and NrCAM. AnkB-220

co-immunoprecipitated with NrCAM from cortex lysates at postnatal (P22, P34) and

adult (P50, P105) stages (Fig. 4B). The relative amounts of associated AnkB-220 and

NrCAM remained approximately equivalent from P22 to P50. The association of AnkB-

220 with NrCAM was further assessed in synaptoneurosomes, a fraction enriched in

pre- and postsynaptic terminals from mouse forebrain (P28) (Villasana *et al.* 2006).

AnkB-220 co-immunoprecipitated with NrCAM from synaptoneurosome lysates (Fig.

4C). AnkB shows enrichment in synaptoneurosome samples over filtered homogenate

and first supernatant samples (equal protein), similar to postsynaptic density protein 95

(PSD95, Fig. 4D), a known synaptic protein, which had been observed previously

(Murphy *et al.* 2023b). Conversely, tubulin, a non-synaptic protein, shows a decrease in

expression in synaptoneurosome samples when compared to filtered homogenate and

first supernatant (Villasana *et al.* 2006; Demyanenko *et al.* 2014)(Fig. 4D).

To probe the role of the NrCAM FIGQY motif in AnkB-220 binding, HEK293 cells

were transfected with plasmids expressing AnkB-220 and either WT NrCAM or NrCAM

FIGQY[1231]H. NrCAM was immunoprecipitated from equal amounts of HEK293 cell

lysates then immunoblotted for AnkB. The relative amounts of AnkB/NrCAM in the

immune complexes were quantified by densitometry. Results showed that AnkB-220 co-

immunoprecipitated efficiently with WT NrCAM but much less with NrCAM FIGQY[1231]H

(Fig. 4E). Input blots verified equivalent levels of AnkB-220, NrCAM, and NrCAM

FIGQY^{1231}H in cell lysates.

Figure 4

Discussion

To investigate a role for AnkB in regulating dendritic spine regulation, we generated a conditional mouse model (Nex1Cre-ERT2: $Ank2^{F/F}$: RCE) to inducibly delete *Ank2* from postmitotic, postmigratory pyramidal neurons at postnatal and adult stages in the PFC. Homozygous deletion of *Ank2* during early postnatal development when spines are most actively remodeled, resulted in elevated spine density on apical but not basal dendrites of layer 2/3 pyramidal neurons persisting in adulthood. Heterozygous deletion of *Ank2* postnatally increased spine density to an intermediate extent only on apical dendrites. In contrast, *Ank2* deletion in young adults did not alter spine density in older adult stages suggesting that mature functions of plasticity and homeostasis may be minimally if at all impacted. Because apical dendrites have different synaptic inputs than basal dendrites (Brzdak *et al.* 2019), it is possible that distinct synaptic connections are remodeled through AnkB interaction with L1 family adhesion molecules in response to different Semaphorins. However, the identity of such inputs has not been determined.

AnkB was found to be essential for Sema3F-induced spine retraction, as demonstrated by a lack of response to Sema3F-Fc in *Ank2*-null neurons *in vitro*. Molecular replacement experiments in *Ank2*-null cortical neurons further revealed AnkB-220 was the isoform responsible for the Sema3F pruning response, as opposed to the neuron-specific isoform AnkB-440. This result was in accord with the preferential localization of AnkB-220 on neuronal dendrites and AnkB-440 on axons (Kunimoto *et al.* 1991; Kunimoto 1995; Chen *et al.* 2020). Our localization studies, which were limited to the use of pan-AnkB antibodies, demonstrated the presence of AnkB within spines in addition to other neuronal compartments, but could not differentiate between AnkB-220 and Ankb-440.

The phenotype of elevated spine density on apical dendrites in Nex1Cre-ERT2:

Ank2^{F/F}: RCE mice is also observed in *NrCAM* conditional mutant mice (Nex1Cre-ERT2:

NrCAM^{F/F}: RCE) (Mohan *et al.* 2019b), as well as in Sema3F-null, Npn2-null, and

PlexA3-null mice (Tran *et al.* 2009; Mohan *et al.* 2019b). This elevated spine phenotype

on apical dendrites is consistent with AnkB-mediated spine elimination through the

Sema3F holoreceptor complex and was also observed in the PFC of CHL1-null and L1-

null mutant mice (Murphy *et al.* 2023a). Since CHL1 shares a homologous Ankyrin

binding site (FIGAY) and binds Npn2, Sema3B-induced retraction of a distinct spine

subpopulation likely also depends on AnkB. Both Sema3F (Wang *et al.* 2017; Mohan *et*

al. 2019a) and Sema3B (Mohan *et al.* 2019b) are expressed in an activity-dependent

manner in culture, suggesting that these ligands may be released at active synaptic

contacts to locally prune weak or inactive spines as cortical circuits mature. In line with

this notion, we found that deletion of *Ank2* increased the relative proportion of stubby

spines compared to mature mushroom spines, suggesting that immature spines may be

selectively eliminated through AnkB function, or that AnkB promotes spine maturation.

NrCAM engaged AnkB-220 at a conserved FIGQY motif in the NrCAM cytoplasmic

domain. This motif was required for spine elimination induced by Sema3F, as mutation

of FIGQY to FIGQH disrupted AnkB-220 binding to NrCAM and blocked spine retraction

in cortical neurons, similar to a previous study in L1 knock-in mice with a substitution in

the FIGQY motif (FIGQH) (Murphy *et al.* 2023a). As shown in Figure 4F, NrCAM binds

Npn2 through extracellular domain interactions which increase Npn2/PlexA3 affinity

(Mohan *et al.* 2019a), activating the intrinsic Rap-GAP activity of PlexA3 (Wang *et al.*

2013). Downstream signal transduction ensues via pathways governed by Rac1 and

RhoA GTPases leading to spine elimination on apical dendrites (Duncan *et al.* 2021a)(Fig. 4F). The recruitment of AnkB to the NrCAM cytoplasmic tail may further stabilize the Sema3F holoreceptor complex, prolonging or enhancing signaling leading to spine retraction. Doublecortin-like kinase 1 (DCLK1) also binds to the Ankyrin binding motif FIGQY in NrCAM (Murphy *et al.* 2023b). However, DCLK1 deletion in postnatal pyramidal neurons of Nex1Cre-ERT2: *DCLK1*$^{flox/flox}$: RCE mice decreases spine density on apical dendrites in contrast to *Ank2* deletion, which increases spine density. Thus, NrCAM binding to Ankyrin, rather than DCLK1, most likely is responsible for constraining spine density.

The Nex1Cre-ERT2: *Ank2*$^{F/F}$: RCE mouse reported here will be useful as a new animal model for *in vivo* studies relevant to ASD that involve prefrontal pyramidal circuits, such as social and cognitive behaviors. Our model most closely resembles ASD *de novo* truncating and frameshift (fs) mutations (R895, R990, L1448, Q3683fs), in which both AnkB-220 and -440 isoforms are deficient. Our model also mimics phenotypes seen in ASD, such as elevated spine density and increased excitatory synaptic puncta. It differs from a frameshift mouse mutant (ABe37fs/fs), in which a stop codon in exon 37 interrupts the neuron-specific sequence unique to AnkB-440 (Yang *et al.* 2019). These juvenile ABe37fs/fs mice displayed transiently increased spines in the somatosensory cortex, and more prominent axonal branching in hippocampal cultures, however these effects were normalized by adulthood. Axons were not extensively analyzed in our *Ank2* F/F model but 2 major axon tracts, the corpus callosum and anterior commissure, were of normal size. Nex1Cre-induced deletion of *Ank2* at P10-P13 may minimize axon defects as deletion is restricted to postmitotic, postmigratory

pyramidal neurons after many axons have reached their targets. Spine defects were not reported in a Nestin-Cre: *Ank2* exon37 F/F mutant in which 90% of AnkB-440 is lost from both neuronal and glial progenitors, or in a heterozygous truncation mutant (ABe22) targeting both AnkB-220 and -440 isoforms (homozygotes were lethal) (Yang *et al.* 2019). All these mouse genetic models will be important in future studies to determine the relative roles of AnkB isoforms at different stages of dendritic and axonal development.

In conclusion, these results illuminate a new molecular function for AnkB in Sema3F-induced spine elimination in postnatally developing pyramidal neurons of the PFC. The findings contribute to our understanding of how cortical pyramidal cells may protect active spines while eliminating others in developing prefrontal circuits. This work also may provide molecular insight into pathology associated with ASD, which is associated with elevated spine density, hyper-excitability and over-connectivity in cortical circuitry (Hinton *et al.* 1991; Irwin *et al.* 2001; Hutsler *et al.* 2010; Tang *et al.* 2014; Sellier *et al.* 2017).

Figure Legends

Figure 1: Conditional Deletion of AnkB in Postnatal Pyramidal Neurons of Nex1Cre-ERT2: _Ank2_$^{F/F}$: RCE Mice

(A) Schematic of the binding of class 3 semaphorins to heterotrimeric receptors consisting of L1 cell adhesion molecules (L1-CAMs), Neuropilins (Npn1/2), and PlexinAs (PlexA1-4). This binding event activates intracellular signaling pathways that induce spine elimination.

(B) Recombination in _Ank2_ F/F primary cingulate cortex (Cg1) at P50 indicated by EGFP immunofluorescence (Alexafluor-488) in pyramidal neurons after TMX induction (P10-13). Cortical layers 1-3, 5, and 6 are labeled. Bar = 100 μm.

(C) EGFP-labeled neurons are not evident in Cg1 of uninduced _Ank2_ F/F mice (P50). Bar = 100 μm.

(D) Representative confocal images showing EGFP-labeled pyramidal neurons (green, Alexafluor-488) and reduced immunofluorescence of AnkB protein (red, Alexafluor-555) in Cg1 layer 2/3 following TMX induction of _Ank2_ F/F mutant mice compared to _Ank2_ +/+ mice. Single optical sections are shown. Mean pixel intensity (± SEM) of AnkB immunofluorescence/unit area (below) was significantly lower in _Ank2_ F/F compared to _Ank2_ +/+ mice (n= 3 mice/genotype; Mann-Whitney 2-tailed test, p=0.03). Arrows in _Ank2_ F/F cortex point to residual AnkB immunofluorescence at the perisomatic region of EGFP-nonexpressing cells. Negative control with both AlexaFluor-conjugated secondary antibodies without primary antibodies showed no detectable fluorescence. All images were captured at same confocal settings.

(E) Representative AnkB immunoblots of cortical lysates (5 µg) from uninduced and induced *Ank2* F/F mice (P50, n= 3 mice/condition). Densitometric scanning of protein bands showed that the ratios of AnkB-220 or AnkB-440 to vinculin as control decreased significantly following TMX induction in *Ank2* F/F cortex compared to uninduced *Ank2* F/F cortex (mean ratio ± SEM; Mann-Whitney 2-tailed test, p=0.01 for each isoform). Ratios in uninduced cortex were set at 100. Different SDS-PAGE and transfer conditions were required to resolve AnkB-220 and AnkB-440 isoforms. (Right) Immunoblotting of cortical lysates indicate AnkB antibody is specific to AnkB and AnkB is not labeled by control antibodies (Actin).

(F) *Ank2* +/+ and *Ank2* F/F mice were induced with TMX and immunostained at P50 for parvalbumin (blue), EGFP (green), and AnkB (red). Panels a, e show 3-channel merged confocal images. Insert in panel e (lower left) shows control staining with secondary AlexaFluor antibodies alone. Single channels show immunostaining of (b,f) parvalbumin (blue channel), (c,g) EGFP (green channel), and (d,h) AnkB (red channel). Parvalbumin+ basket interneurons are EGFP-negative but express AnkB at the soma (arrows). Images are single optical sections captured at the same confocal settings (scale bar = 10 μm).

(G) Representative confocal images of the corpus callosum and anterior commissure at the midline (white brackets) in coronal brain sections of *Ank2* +/+ mice (n=4) and *Ank2* F/F mice (n=3, P50) expressing EGFP after TMX induction (P10-P13). Measurements of mean width (± SEM) were made using the straight-line tool in FIJI. Results indicated no significant difference in mean width of either tract for *Ank2* F/F compared to *Ank2* +/+

mice at the corpus callosum (Mann-Whitney test, 2- tailed, p= 0.39; scale bar = 50 µm) or anterior commissure (p=0.72; scale bar = 25 µm).

Figure 2: Increased Spine Density and Immature Synapses on Apical Dendrites in *Ank2* F/F Pyramidal Neurons in PFC Layer 2/3

(A) Representative images of EGFP-labeled apical dendrites and spines on pyramidal neurons in layer 2/3 of the medial PFC were subjected to 3D reconstruction of deconvolved dendritic z-stacks in Imaris (scale bar = 7 µm). P50 panels show that deletion of *Ank2* at P10-P13 by TMX induction increased spine density on apical dendrites in *Ank2* F/F and *Ank2* F/+ mice compared to *Ank2* +/+ mice at P50. P80 panels show that deletion of *Ank2* by TMX induction at P50 did not increase spine density on apical dendrites in older adults at P80. Three panels at right are representative low magnification confocal images of cortical neurons from P50 mice, with arrows indicating apical dendrites (A). Scale bar = 20 µm.

(B) Quantification of mean spine density per 10 µm dendritic length (± SEM) on apical and basal dendrites of pyramidal neurons in layer 2/3 of PFC in *Ank2* +/+, F/+, and F/F mice induced at P10-P13 and analyzed at P50. Each point represents the mean spine intensity per mouse analyzed. One-factor ANOVA with Tukey's posthoc testing showed significantly increased spine density on apical dendrites of *Ank2* F/F pyramidal neurons (9.2 spines/10 µm ± 0.5, n=11 mice) compared to *Ank2* +/+ (4.9 spines/10 µm ± 0.2, n=13 mice, *p<0.0001). *Ank2* F/+ mice exhibited intermediate spine density on apical dendrites (6.7 spines/10 µm ± 0.4, n=6 mice), which was significantly greater than *Ank2* +/+ mice (*p=0.02), and significantly less than *Ank2* F/F (*p=0.001). Spine density on basal dendrites of *Ank2* +/+ pyramidal neurons in layer 2/3 PFC (4.2 spines/10 µm ±

0.2; n=10 mice) was not significantly different from heterozygous *Ank2* F/+ (4.4

spines/10 µm ± 0.3; n=4 mice, p = 0.91) or *Ank2* F/F mice (3.8 ± 0.2; n=6 mice, p =

0.39). *Ank2* F/+ mice were not significantly different from *Ank2* F/F mice (p = 0.33). The

number of neurons analyzed per mouse ranged from 10-21.

(C) Quantification of mean spine density per 10 µm dendritic length (± SEM) on apical

dendrites of pyramidal neurons in layer 2/3 of PFC in *Ank2* +/+, F/+, and F/F mice

induced at P50 and analyzed at P80. Each point represents the mean spine intensity

per mouse. One-factor ANOVA with Tukey's posthoc testing showed no difference in

apical dendritic spine density in *Ank2* +/+ mice (5.4 ± 0.5, n=6 mice) compared to *Ank2*

F/+ mice (5.4 ± 0.3, n=6 mice, p = 0.99) or *Ank2* F/F mice (6.0 spines/10 µm ± 0.5; n=9

mice, p = 0.72). *Ank2* F/+ mice were also not significantly different from *Ank2* F/F mice

(p = 0.64).

(D) Spine morphologies were quantified in *Ank2* +/+, *Ank2* F/+, and *Ank2* F/F mice

induced at P10-P13 and analyzed at P50. Quantities of each morphological type are

reported as percentages of total number of spines (± SEM). Spines were identified as

one of three morphologies based on spine length and the ratio of head width to neck

width. One-factor ANOVA with Tukey's posthoc testing showed a significantly increased

percentage of stubby spines in *Ank2* F/F (34.5 % of total spines ± 0.9) compared to

Ank2 +/+ mice (27.7% ± 1.5, *p = 0.005), but no significant difference between *Ank2* F/+

stubby spines (31.4% ± 1.4) and either *Ank2* F/F stubby spines (p = 0.82) or *Ank2* +/+

stubby spines (p = 0.67). Analysis also showed a significant decrease in mushroom

spines in *Ank2* F/F (49.7% ± 1.1) compared to *Ank2* +/+ mice (56.6% ± 1.3, *p = 0.004),

but no significant difference between *Ank2* F/+ mushroom spines (55.1% ± 2.1) and

Ank2 F/F mushroom spines (p = 0.16) or *Ank2* +/+ mushroom spines (p = 0.99). The percent of thin spines was not significantly different between genotypes (p > 0.05). The total number of neurons analyzed per genotype was between 9-15.

(E) Sholl analysis of neuronal process branching of EGFP-expressing pyramidal neurons in PFC layer 2/3 showed no significant differences (n.s.) in the number of intersections of branches (primarily dendrites) with concentric circles at various distances from the soma of *Ank2* +/+ and *Ank2* F/F mice (P50) following TMX induction at P10-P13 (n =15 cells/genotype, 3 mice/genotype, two-factor ANOVA, p = 0.85).

(F) Excitatory synaptic puncta were visualized by immunofluorescence staining of juxtaposed vGlut1 with Alexafluor-555 (red) and Homer-1 with Alexafluor-647(blue) in layer 2/3 of the PFC on apical dendritic spines of EGFP-expressing pyramidal neurons in *Ank2* +/+ and *Ank2* F/F mice (P50) after TMX induction at P10-P13. Representative confocal images (merged) show synaptic puncta of adjacent vGlut1 and Homer-1 labeling (arrows), scale bar = 1 μm.

(G) Quantification of vGlut1/Homer-1 double-labeled puncta on spines per 10 μm dendritic length in single optical sections. *Ank2* F/F neurons exhibited an increased mean density of these puncta compared to *Ank2* +/+ neurons (Mann-Whitney 2-tailed test, *p=0.004; n= 4 mice/genotype, 22-24 neurons/genotype). Points represent puncta density per neuron.

Figure 3: Sema3F-Induced Spine Retraction is Impaired in Cortical Neuron Cultures from *Ank2*-null, NrCAM-null, and NrCAM Y^{1231}H Mutant Mice

(A) WT and *Ank2*-null cortical neuronal cultures were transfected with pCAG-IRES-EGFP, treated with 5 nM Fc or Sema3F-Fc for 30 min at DIV14, immunostained for

EGFP, and apical dendrites imaged confocally (scale bar = 5 μm). *Ank2*-null cultures were co-transfected with Ank2-220 or Ank2-440 cDNAs. Representative images show EGFP-labeled apical dendrites with spines in WT and *Ank2*-null cortical neuronal cultures treated with Fc or Sema3F-Fc.

(B) Quantification of mean spine intensity ± SEM on apical dendrites of neuronal cultures described in (A). Each point represents the mean spine density per neuron per 10 μm dendrite length. Two-factor ANOVA with Tukey post hoc testing of spine density on *Ank2*-null neurons demonstrated that *Ank2* deletion inhibited Sema3F-induced spine retraction (Fc-treated, 3.23 spines/10 μm ± 0.18; Sema3F-Fc treated, 3.01 spines/10 μm ± 0.15, p = 0.99). In contrast WT neurons exhibited spine retraction in response to Sema3F (Fc-treated, 3.24 spines/10 μm ± 0.14; Sema3F-Fc-treated, 2.45 spines/10 μm ± 0.11, *p < 0.001). Sema3F-Fc-induced spine retraction in *Ank2*-null neurons was rescued by re-expressing AnkB-220 cDNA (Fc-treated, 3.08 spines/10 μm ± 0.14; Sema3F-Fc-treated, 2.30 spines/10 μm ±0.10, *p=0.001). AnkB-440 cDNA was not able to rescue Sema3F-induced spine retraction (Fc, treated, 3.59 spines/10 μm ± 0.20; Sema3F-Fc -treated, 3.43 spines/10 μm ±0.18, p = 0.99). Immunostaining with pan-AnkB antibodies verified equivalent levels of isoform expression. Points represent mean spine density of individual neurons.

(C) WT cortical neuronal cultures were transfected with pCAG-IRES-EGFP, immunostained, and imaged confocally (DIV14). Representative confocal images of neurons with apical dendritic spines (a-f) immunostained with pan-AnkB antibodies and Alexafluor-555 (red; a, d) and GFP with Alexafluor-488 (green; b, e). Merged images (c, f). Control staining with secondary antibodies alone is shown in the upper right dotted

inset in (a). Top panels are maximum intensity projections, with a single optical section of AnkB immunofluorescence staining of neurons shown in the lower right dotted inset in (a). AnkB immunofluorescence labeling was observed in spines with filopodial (fp), mushroom (mr), and stubby (st) morphology (d, arrows). At higher magnification AnkB immunolabeling with Alexafluor-594 was evident within a spine head domain (arrows, pseudocolored red) by (g) confocal or (h) STED superresolution microscopy. Scale bars = 100 μm (a-c), 5 μm (d-f), 0.2 μm (g,h).

(D) *NrCAM*-null cortical neuronal cultures were transfected with pCAG-IRES-EGFP vector alone, or plasmids containing WT or NrCAM $Y^{1231}H$ cDNAs. Cultures were treated with 5 nM Fc or Sema3F-Fc for 30 min at DIV14, immunostained for EGFP and apical dendrites imaged confocally (scale bar = 5 μm). Representative images of EGFP-labeled apical dendrites show that Sema3F-Fc promotes spine retraction only on *NrCAM*-null neurons re-expressing WT NrCAM.

(E) Sema3F-Fc induces spine retraction on apical dendrites of *NrCAM*-null cortical neurons re-expressing WT NrCAM but not NrCAM $Y^{1231}H$ or empty vector. Two-factor ANOVA with Tukey post hoc testing showed a significant difference (*p=0.003) in spine density in neurons re-expressing WT NrCAM following treatment with control Fc (4.28 spines/10 μm ± 0.16) vs. Sema3F-Fc (3.28 spines/10 μm ± 0.15). In contrast there was not a significant difference (p = 0.47) in spine density in neurons re-expressing NrCAM $Y^{1231}H$ following treatment with control Fc (4.25 spines/10 μm ± 0.22) vs. Sema3F-Fc (4.67 spines/10 μm ± 0.25). There was also not a significant difference (p > 0.99) in spine density in NrCAM-null neurons transfected with vector alone.

Figure 4: AnkB-220 Associates with NrCAM at the FIGQY Motif.

(A) Representative immunoblotting (IB) of AnkB-220 (220kDa) and NrCAM (130 kDa) in cortical lysates from postnatal mouse brain (P22-P105) (20 µg protein). Relative amounts of AnkB-220 or NrCAM were determined by densitometric scanning of respective protein bands. AnkB-220 and NrCAM expression were equivalent from P22-P105. Similar results were obtained in duplicate experiments.

(B) Co-immunoprecipitation of AnkB-220 (220kDa) and NrCAM (130 kDa) from equal amounts of protein (1mg) in cortex lysates of postnatal mouse forebrain (P22-P105). NrCAM was immunoprecipitated (IP) and AnkB-220 detected by immunoblotting (IB). AnkB immunoblots (above) were reprobed for NrCAM (below). Densitometric scanning was performed and the ratio of immunoprecipitated AnkB-220 to NrCAM indicated below.

(C) Co-immunoprecipitation of AnkB-220 with NrCAM from P28 mouse synaptoneurosomes, shown by immunoprecipitation (IP) with NrCAM antibodies or nonimmune IgG (NIg) and immunoblotting (IB) with AnkB antibodies. Blots were reprobed by immunoblotting with NrCAM antibodies (lower panels). Inputs represent synaptoneurosome lysates that were not subjected to IP. Example is representative of replicate blots.

(D) Filtered homogenate (FH), first supernatant (S1), and synaptoneurosome (Syn) samples (equal protein, 25µg) were blotted for Ankyrin B (AnkB). Membranes were stripped and reprobed for post-synaptic density protein 95 (PSD95, known synaptic protein), tubulin (a non-synaptic protein), and actin (loading control). AnkB and PSD95 showed enrichment in Syn samples, whereas tubulin showed decreased expression in Syn samples.

(E) Co-immunoprecipitation of AnkB-220 with WT NrCAM or mutant NrCAM $Y^{1231}H$ from transfected HEK293T cells (equal amounts protein). NrCAM was immunoprecipitated from HEK293T cell lysates and immunoblotted for AnkB. Blots were reprobed for NrCAM (lower panels). Densitometric scanning of bands yielded ratios of AnkB-220 to NrCAM in the IP samples (below). Input lysates (equal protein) are shown at right.

(F) Schematic of spine pruning process initiated by Sema3F. Sema3F binds the holoreceptor complex formed by NrCAM, Npn-2, and PlexA3. This binding event activates Rap-GAP activity and subsequent downstream signaling leads to spine elimination via Rac1 and RhoA GTPase-governed pathways. AnkB recruitment to NrCAM cytoplasmic domain may stabilize the Sema3F complex, enhancing the signaling leading to spine pruning.

Funding: This work was supported by the National Institutes of Health (R01 MH113280 to PFM and by a UNC School of Medicine Biomedical Research Core Project Award to PFM, the Carolina Institute for Developmental Disabilities Center Grant NIH P50HD103573 to Dr. Joseph Piven, and National Institutes of Health NRSA T32 HD040127 to KEM. Dr. Pablo Ariel, Director of the Microscopy Services Laboratory, UNC Department of Pathology and Laboratory Medicine, provided expert advice and assistance with microscopy, as supported by National Institutes of Health Cancer Center Core Grant P30 CA016086.

Acknowledgements:

We gratefully acknowledge Dr. Peter Mohler for providing Ank2 floxed mice, and Dr. Gordon Fishell for RCE mice. We acknowledge Dr. Paul Jenkins and Dr. Kathryn

Walder for antibodies and assistance with Ankyrin analysis. We thank Dr. Young Truong

(Department of Biostatistics at UNC) for multinomial logistic regression analysis of spine

morphology. Dr. Damaris Lorenzo provided assistance with mouse breeding.

References

Agarwal A, Dibaj P, Kassmann CM, Goebbels S, Nave KA, Schwab MH. 2012. In vivo imaging and noninvasive ablation of pyramidal neurons in adult NEX-CreERT2 mice. Cereb Cortex. 22:1473-1486.

Alvarez VA, Sabatini BL. 2007. Anatomical and physiological plasticity of dendritic spines. Annu Rev Neurosci. 30:79-97.

Anastasiades PG, Carter AG. 2021. Circuit organization of the rodent medial prefrontal cortex. Trends Neurosci. 44:550-563.

Bennett V, Healy J. 2009. Membrane domains based on ankyrin and spectrin associated with cell-cell interactions. Cold Spring Harb Perspect Biol. 1:1-19.

Berry KP, Nedivi E. 2017. Spine Dynamics: Are They All the Same? Neuron. 96:43-55.

Bhatt DH, Zhang S, Gan WB. 2009. Dendritic spine dynamics. Annu Rev Physiol. 71:261-282.

Bourne J, Harris KM. 2007. Do thin spines learn to be mushroom spines that remember? Curr Opin Neurobiol. 17:381-386.

Brzdak P, Wojcicka O, Zareba-Koziol M, Minge D, Henneberger C, Wlodarczyk J, Mozrzymas JW, Wojtowicz T. 2019. Synaptic Potentiation at Basal and Apical Dendrites of Hippocampal Pyramidal Neurons Involves Activation of a Distinct Set of Extracellular and Intracellular Molecular Cues. Cereb Cortex. 29:283-304.

Buhusi M, Schlatter MC, Demyanenko GP, Thresher R, Maness PF. 2008. L1 interaction with ankyrin regulates mediolateral topography in the retinocollicular projection. J Neurosci. 28:177-188.

Chen K, Li J, Wang C, Wei Z, Zhang M. 2017. Autoinhibition of ankyrin-B/G membrane target bindings by intrinsically disordered segments from the tail regions. Elife. 6.

Chen K, Yang R, Li Y, Zhou JC, Zhang M. 2020. Giant ankyrin-B suppresses stochastic collateral axon branching through direct interaction with microtubules. J Cell Biol. 219.

Culotta L, Penzes P. 2020. Exploring the mechanisms underlying excitation/inhibition imbalance in human iPSC-derived models of ASD. Mol Autism. 11:32.

Dai J, Buhusi M, Demyanenko GP, Brennaman LH, Hruska M, Dalva MB, Maness PF. 2013. Neuron glia-related cell adhesion molecule (NrCAM) promotes topographic retinocollicular mapping. PLoS One. 8:e73000.

Dai J, Dalal JS, Thakar S, Henkemeyer M, Lemmon VP, Harunaga JS, Schlatter MC, Buhusi M, Maness PF. 2012. EphB regulates L1 phosphorylation during retinocollicular mapping. Mol Cell Neurosci. 50:201-210.

De Rubeis S, He X, Goldberg AP, Poultney CS, Samocha K, Cicek AE, Kou Y, Liu L, Fromer M, Walker S, Singh T, Klei L, Kosmicki J, Shih-Chen F, Aleksic B, Biscaldi M, Bolton PF, Brownfeld JM, Cai J, Campbell NG, Carracedo A, Chahrour MH, Chiocchetti AG, Coon H, Crawford EL, Curran SR, Dawson G, Duketis E, Fernandez BA, Gallagher L, Geller E, Guter SJ, Hill RS, Ionita-Laza J, Jimenz Gonzalez P, Kilpinen H, Klauck SM, Kolevzon A, Lee I, Lei I, Lei J, Lehtimaki T, Lin CF, Ma'ayan A, Marshall CR,

McInnes AL, Neale B, Owen MJ, Ozaki N, Parellada M, Parr JR, Purcell S, Puura K, Rajagopalan D, Rehnstrom K, Reichenberg A, Sabo A, Sachse M, Sanders SJ, Schafer C, Schulte-Ruther M, Skuse D, Stevens C, Szatmari P, Tammimies K, Valladares O, Voran A, Li-San W, Weiss LA, Willsey AJ, Yu TW, Yuen RK, Study DDD, Homozygosity Mapping Collaborative for A, Consortium UK, Cook EH, Freitag CM, Gill M, Hultman CM, Lehner T, Palotie A, Schellenberg GD, Sklar P, State MW, Sutcliffe JS, Walsh CA, Scherer SW, Zwick ME, Barett JC, Cutler DJ, Roeder K, Devlin B, Daly MJ, Buxbaum JD. 2014. Synaptic, transcriptional and chromatin genes disrupted in autism. Nature. 515:209-215.

Demyanenko GP, Mohan V, Zhang X, Brennaman LH, Dharbal KE, Tran TS, Manis PB, Maness PF. 2014. Neural Cell Adhesion Molecule NrCAM Regulates Semaphorin 3F-Induced Dendritic Spine Remodeling. J Neurosci. 34:11274-11287.

Demyanenko GP, Tsai AY, Maness PF. 1999. Abnormalities in neuronal process extension, hippocampal development, and the ventricular system of L1 knockout mice. J Neurosci. 19:4907-4920.

Duncan BW, Mohan V, Wade SD, Truong Y, Kampov-Polevoi A, Temple BR, Maness PF. 2021a. Semaphorin3F Drives Dendritic Spine Pruning Through Rho-GTPase Signaling. Mol Neurobiol. 58:3817-3834.

Duncan BW, Murphy KE, Maness PF. 2021b. Molecular Mechanisms of L1 and NCAM Adhesion Molecules in Synaptic Pruning, Plasticity, and Stabilization. Front Cell Dev Biol. 9:625340.

Fame RM, MacDonald JL, Dunwoodie SL, Takahashi E, Macklis JD. 2016. Cited2 Regulates Neocortical Layer II/III Generation and Somatosensory Callosal Projection Neuron Development and Connectivity. J Neurosci. 36:6403-6419.

Forrest MP, Parnell E, Penzes P. 2018. Dendritic structural plasticity and neuropsychiatric disease. Nat Rev Neurosci. 19:215-234.

Garver TD, Ren Q, Tuvia S, Bennett V. 1997. Tyrosine phosphorylation at a site highly conserved in the L1 family of cell adhesion molecules abolishes ankyrin binding and increases lateral mobility of neurofascin. J Cell Biol. 137:703-714.

Golonzhka O, Nord A, Tang PLF, Lindtner S, Ypsilanti AR, Ferretti E, Visel A, Selleri L, Rubenstein JLR. 2015. Pbx Regulates Patterning of the Cerebral Cortex in Progenitors and Postmitotic Neurons. Neuron. 88:1192-1207.

Hinton VJ, Brown WT, Wisniewski K, Rudelli RD. 1991. Analysis of neocortex in three males with the fragile X syndrome. Am J Med Genet. 41:289-294.

Holtmaat A, Svoboda K. 2009. Experience-dependent structural synaptic plasticity in the mammalian brain. Nat Rev Neurosci. 10:647-658.

Holtmaat AJ, Trachtenberg JT, Wilbrecht L, Shepherd GM, Zhang X, Knott GW, Svoboda K. 2005. Transient and persistent dendritic spines in the neocortex in vivo. Neuron. 45:279-291.

Hortsch M, Nagaraj K, Mualla R. 2014. The L1 family of cell adhesion molecules: a sickening number of mutations and protein functions. Adv Neurobiol. 8:195-229.

Hutsler JJ, Zhang H. 2010. Increased dendritic spine densities on cortical projection neurons in autism spectrum disorders. Brain Res. 1309:83-94.

Huttenlocher PR. 1979. Synaptic density in human frontal cortex - developmental changes and effects of aging. Brain Res. 163:195-205.

Iossifov I, Levy D, Allen J, Ye K, Ronemus M, Lee YH, Yamrom B, Wigler M. 2015. Low load for disruptive mutations in autism genes and their biased transmission. Proc Natl Acad Sci U S A. 112:E5600-5607.

Iossifov I, Ronemus M, Levy D, Wang Z, Hakker I, Rosenbaum J, Yamrom B, Lee YH, Narzisi G, Leotta A, Kendall J, Grabowska E, Ma B, Marks S, Rodgers L, Stepansky A, Troge J, Andrews P, Bekritsky M, Pradhan K, Ghiban E, Kramer M, Parla J, Demeter R, Fulton LL, Fulton RS, Magrini VJ, Ye K, Darnell JC, Darnell RB, Mardis ER, Wilson RK, Schatz MC, McCombie WR, Wigler M. 2012. De novo gene disruptions in children on the autistic spectrum. Neuron. 74:285-299.

Irwin SA, Patel B, Idupulapati M, Harris JB, Crisostomo RA, Larsen BP, Kooy F, Willems PJ, Cras P, Kozlowski PB, Swain RA, Weiler IJ, Greenough WT. 2001. Abnormal dendritic spine characteristics in the temporal and visual cortices of patients with fragile-X syndrome: a quantitative examination. Am J Med Genet. 98:161-167.

Jenkins PM, Kim N, Jones SL, Tseng WC, Svitkina TM, Yin HH, Bennett V. 2015. Giant ankyrin-G: a critical innovation in vertebrate evolution of fast and integrated neuronal signaling. Proc Natl Acad Sci U S A. 112:957-964.

Kroon T, van Hugte E, van Linge L, Mansvelder HD, Meredith RM. 2019. Early postnatal development of pyramidal neurons across layers of the mouse medial prefrontal cortex. Sci Rep. 9:5037.

Kunimoto M. 1995. A neuron-specific isoform of brain ankyrin, 440-kD ankyrinB, is targeted to the axons of rat cerebellar neurons. J Cell Biol. 131:1821-1829.

Kunimoto M, Otto E, Bennett V. 1991. A new 440-kd isoform is the major ankyrin in neonatal brain. J Cell Biol. 115:1319-1331.

Laviola G, Macri S, Morley-Fletcher S, Adriani W. 2003. Risk-taking behavior in adolescent mice: psychobiological determinants and early epigenetic influence. Neurosci Biobehav Rev. 27:19-31.

Lim ET, Raychaudhuri S, Sanders SJ, Stevens C, Sabo A, MacArthur DG, Neale BM, Kirby A, Ruderfer DM, Fromer M, Lek M, Liu L, Flannick J, Ripke S, Nagaswamy U, Muzny D, Reid JG, Hawes A, Newsham I, Wu Y, Lewis L, Dinh H, Gross S, Wang LS, Lin CF, Valladares O, Gabriel SB, dePristo M, Altshuler DM, Purcell SM, State MW, Boerwinkle E, Buxbaum JD, Cook EH, Gibbs RA, Schellenberg GD, Sutcliffe JS, Devlin B, Roeder K, Daly MJ. 2013. Rare complete knockouts in humans: population distribution and significant role in autism spectrum disorders. Neuron. 77:235-242.

Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C. 2004. Interneurons of the neocortical inhibitory system. Nat Rev Neurosci. 5:793-807.

Miller DS, Wright KM. 2021. Neuronal Dystroglycan regulates postnatal development of CCK/cannabinoid receptor-1 interneurons. Neural Dev. 16:4.

Mohan V, Sullivan CS, Guo J, Wade SD, Majumder S, Agarwal A, Anton ES, Temple BS, Maness PF. 2019a. Temporal Regulation of Dendritic Spines Through NrCAM-Semaphorin3F Receptor Signaling in Developing Cortical Pyramidal Neurons. Cereb Cortex. 29:963-977.

Mohan V, Wade SD, Sullivan CS, Kasten MR, Sweetman C, Stewart R, Truong Y, Schachner M, Manis PB, Maness PF. 2019b. Close Homolog of L1 Regulates Dendritic Spine Density in the Mouse Cerebral Cortex Through Semaphorin 3B. J Neurosci. 39:6233-6250.

Murphy KE, Wade SD, Sperringer JE, Mohan V, Duncan BW, Zhang EY, Pak Y, Lutz D, Schachner M, Maness PF. 2023a. The L1 cell adhesion molecule constrains dendritic spine density in pyramidal neurons of the mouse cerebral cortex. Front Neuroanat. 17:1111525.

Murphy KE, Zhang EY, Wyatt EV, Sperringer JE, Duncan BW, Maness PF. 2023b. Doublecortin-Like Kinase 1 Facilitates Dendritic Spine Growth of Pyramidal Neurons in Mouse Prefrontal Cortex. Neuroscience. 508:98-109.

Petanjek Z, Judas M, Simic G, Rasin MR, Uylings HB, Rakic P, Kostovic I. 2011. Extraordinary neoteny of synaptic spines in the human prefrontal cortex. Proc Natl Acad Sci U S A. 108:13281-13286.

Peters A, Kaiserman-Abramof IR. 1970. The small pyramidal neuron of the rat cerebral cortex. The perikaryon, dendrites and spines. Am J Anat. 127:321-355.

Phillips M, Pozzo-Miller L. 2015. Dendritic spine dysgenesis in autism related disorders. Neurosci Lett. 601:30-40.

Qu F, Lorenzo DN, King SJ, Brooks R, Bear JE, Bennett V. 2016. Ankyrin-B is a PI3P effector that promotes polarized alpha5beta1-integrin recycling via recruiting RabGAP1L to early endosomes. Elife. 5:1-25.

Roberts JD, Murphy NP, Hamilton RM, Lubbers ER, James CA, Kline CF, Gollob MH, Krahn AD, Sturm AC, Musa H, El-Refaey M, Koenig S, Aneq MA, Hoorntje ET, Graw SL, Davies RW, Rafiq MA, Koopmann TT, Aafaqi S, Fatah M, Chiasson DA, Taylor MR, Simmons SL, Han M, van Opbergen CJ, Wold LE, Sinagra G, Mittal K, Tichnell C, Murray B, Codima A, Nazer B, Nguyen DT, Marcus FI, Sobriera N, Lodder EM, van den Berg MP, Spears DA, Robinson JF, Ursell PC, Green AK, Skanes AC, Tang AS, Gardner MJ, Hegele RA, van Veen TA, Wilde AA, Healey JS, Janssen PM, Mestroni L, van Tintelen JP, Calkins H, Judge DP, Hund TJ, Scheinman MM, Mohler PJ. 2019. Ankyrin-B dysfunction predisposes to arrhythmogenic cardiomyopathy and is amenable to therapy. J Clin Invest. 129:3171-3184.

Sakurai T, Lustig M, Babiarz J, Furley AJ, Tait S, Brophy PJ, Brown SA, Brown LY, Mason CA, Grumet M. 2001. Overlapping functions of the cell adhesion molecules Nr-CAM and L1 in cerebellar granule cell development. J Cell Biol. 154:1259-1273.

Sakurai T, Ramoz N, Reichert JG, Corwin TE, Kryzak L, Smith CJ, Silverman JM, Hollander E, Buxbaum JD. 2006. Association analysis of the NrCAM gene in autism and in subsets of families with severe obsessive-compulsive or self-stimulatory behaviors. Psychiatr Genet. 16:251-257.

Satterstrom FK, Walters RK, Singh T, Wigdor EM, Lescai F, Demontis D, Kosmicki JA, Grove J, Stevens C, Bybjerg-Grauholm J, Baekvad-Hansen M, Palmer DS, Maller JB, i P-BC, Nordentoft M, Mors O, Robinson EB, Hougaard DM, Werge TM, Bo Mortensen P, Neale BM, Borglum AD, Daly MJ. 2019. Autism spectrum disorder and attention deficit hyperactivity disorder have a similar burden of rare protein-truncating variants. Nat Neurosci. 22:1961-1965.

Scotland P, Zhou D, Benveniste H, Bennett V. 1998. Nervous system defects of AnkyrinB (-/-) mice suggest functional overlap between the cell adhesion molecule L1 and 440-kD AnkyrinB in premyelinated axons. J Cell Biol. 143:1305-1315.

Sellier C, Buijsen RAM, He F, Natla S, Jung L, Tropel P, Gaucherot A, Jacobs H, Meziane H, Vincent A, Champy MF, Sorg T, Pavlovic G, Wattenhofer-Donze M, Birling MC, Oulad-Abdelghani M, Eberling P, Ruffenach F, Joint M, Anheim M, Martinez-Cerdeno V, Tassone F, Willemsen R, Hukema RK, Viville S, Martinat C, Todd PK, Charlet-Berguerand N. 2017. Translation of Expanded CGG Repeats into FMRpolyG Is Pathogenic and May Contribute to Fragile X Tremor Ataxia Syndrome. Neuron. 93:331-347.

Smith KR, Kopeikina KJ, Fawcett-Patel JM, Leaderbrand K, Gao RQ, Schurmann B, Myczek K, Radulovic J, Swanson GT, Penzes P. 2014. Psychiatric Risk Factor ANK3/Ankyrin-G Nanodomains Regulate the Structure and Function of Glutamatergic Synapses. Neuron. 84:399-415.

Smith SA, Sturm AC, Curran J, Kline CF, Little SC, Bonilla IM, Long VP, Makara M, Polina I, Hughes LD, Webb TR, Wei Z, Wright P, Voigt N, Bhakta D, Spoonamore KG,

Zhang C, Weiss R, Binkley PF, Janssen PM, Kilic A, Higgins RS, Sun M, Ma J, Dobrev D, Zhang M, Carnes CA, Vatta M, Rasband MN, Hund TJ, Mohler PJ. 2015. Dysfunction in the betaII spectrin-dependent cytoskeleton underlies human arrhythmia. Circulation. 131:695-708.

Sousa VH, Miyoshi G, Hjerling-Leffler J, Karayannis T, Fishell G. 2009. Characterization of Nkx6-2-derived neocortical interneuron lineages. Cereb Cortex. 19 Suppl 1:i1-10.

Spruston N. 2008. Pyramidal neurons: dendritic structure and synaptic integration. Nat Rev Neurosci. 9:206-221.

Stevens SR, Rasband MN. 2021. Ankyrins and neurological disease. Curr Opin Neurobiol. 69:51-57.

Tai Y, Gallo NB, Wang M, Yu JR, Van Aelst L. 2019. Axo-axonic Innervation of Neocortical Pyramidal Neurons by GABAergic Chandelier Cells Requires AnkyrinG-Associated L1CAM. Neuron. 102:358-372.

Tang G, Gudsnuk K, Kuo SH, Cotrina ML, Rosoklija G, Sosunov A, Sonders MS, Kanter E, Castagna C, Yamamoto A, Yue Z, Arancio O, Peterson BS, Champagne F, Dwork AJ, Goldman J, Sulzer D. 2014. Loss of mTOR-dependent macroautophagy causes autistic-like synaptic pruning deficits. Neuron. 83:1131-1143.

Tasic B, Menon V, Nguyen TN, Kim TK, Jarsky T, Yao Z, Levi B, Gray LT, Sorensen SA, Dolbeare T, Bertagnolli D, Goldy J, Shapovalova N, Parry S, Lee C, Smith K, Bernard A, Madisen L, Sunkin SM, Hawrylycz M, Koch C, Zeng H. 2016. Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. Nat Neurosci. 19:335-346.

Trachtenberg JT, Chen BE, Knott GW, Feng G, Sanes JR, Welker E, Svoboda K. 2002. Long-term in vivo imaging of experience-dependent synaptic plasticity in adult cortex. Nature. 420:788-794.

Tran TS, Rubio ME, Clem RL, Johnson D, Case L, Tessier-Lavigne M, Huganir RL, Ginty DD, Kolodkin AL. 2009. Secreted semaphorins control spine distribution and morphogenesis in the postnatal CNS. Nature. 462:1065-1069.

Villasana LE, Klann E, Tejada-Simon MV. 2006. Rapid isolation of synaptoneurosomes and postsynaptic densities from adult mouse hippocampus. J Neurosci Methods. 158:30-36.

Wang D, Liu S, Warrell J, Won H, Shi X, Navarro FCP, Clarke D, Gu M, Emani P, Yang YT, Xu M, Gandal MJ, Lou S, Zhang J, Park JJ, Yan C, Rhie SK, Manakongtreecheep K, Zhou H, Nathan A, Peters M, Mattei E, Fitzgerald D, Brunetti T, Moore J, Jiang Y, Girdhar K, Hoffman GE, Kalayci S, Gumus ZH, Crawford GE, Psych EC, Roussos P, Akbarian S, Jaffe AE, White KP, Weng Z, Sestan N, Geschwind DH, Knowles JA, Gerstein MB. 2018. Comprehensive functional genomic resource and integrative model for the human brain. Science. 362.

Wang Q, Chiu SL, Koropouli E, Hong I, Mitchell S, Easwaran TP, Hamilton NR, Gustina AS, Zhu Q, Ginty DD, Huganir RL, Kolodkin AL. 2017. Neuropilin-2/PlexinA3 Receptors Associate with GluA1 and Mediate Sema3F-Dependent Homeostatic Scaling in Cortical Neurons. Neuron. 96:1084-1098.

Wang T, Guo H, Xiong B, Stessman HA, Wu H, Coe BP, Turner TN, Liu Y, Zhao W, Hoekzema K, Vives L, Xia L, Tang M, Ou J, Chen B, Shen Y, Xun G, Long M, Lin J, Kronenberg ZN, Peng Y, Bai T, Li H, Ke X, Hu Z, Zhao J, Zou X, Xia K, Eichler EE. 2016. De novo genic mutations among a Chinese autism spectrum disorder cohort. Nat Commun. 7:1-10.

Wang T, Hoekzema K, Vecchio D, Wu H, Sulovari A, Coe BP, Gillentine MA, Wilfert AB, Perez-Jurado LA, Kvarnung M, Sleyp Y, Earl RK, Rosenfeld JA, Geisheker MR, Han L, Du B, Barnett C, Thompson E, Shaw M, Carroll R, Friend K, Catford R, Palmer EE, Zou X, Ou J, Li H, Guo H, Gerdts J, Avola E, Calabrese G, Elia M, Greco D, Lindstrand A, Nordgren A, Anderlid BM, Vandeweyer G, Van Dijck A, Van der Aa N, McKenna B, Hancarova M, Bendova S, Havlovicova M, Malerba G, Bernardina BD, Muglia P, van Haeringen A, Hoffer MJV, Franke B, Cappuccio G, Delatycki M, Lockhart PJ, Manning MA, Liu P, Scheffer IE, Brunetti-Pierri N, Rommelse N, Amaral DG, Santen GWE, Trabetti E, Sedlacek Z, Michaelson JJ, Pierce K, Courchesne E, Kooy RF, Consortium S, Nordenskjold M, Romano C, Peeters H, Bernier RA, Gecz J, Xia K, Eichler EE. 2020. Large-scale targeted sequencing identifies risk genes for neurodevelopmental disorders. Nat Commun. 11:1-13.

Wang Y, Pascoe HG, Brautigam CA, He H, Zhang X. 2013. Structural basis for activation and non-canonical catalysis of the Rap GTPase activating protein domain of plexin. Elife. 2:e01279.

Willsey AJ, Sanders SJ, Li M, Dong S, Tebbenkamp AT, Muhle RA, Reilly SK, Lin L, Fertuzinhos S, Miller JA, Murtha MT, Bichsel C, Niu W, Cotney J, Ercan-Sencicek AG,

Gockley J, Gupta AR, Han W, He X, Hoffman EJ, Klei L, Lei J, Liu W, Liu L, Lu C, Xu X, Zhu Y, Mane SM, Lein ES, Wei L, Noonan JP, Roeder K, Devlin B, Sestan N, State MW. 2013. Coexpression networks implicate human midfetal deep cortical projection neurons in the pathogenesis of autism. Cell. 155:997-1007.

Yang R, Walder-Christensen KK, Kim N, Wu D, Lorenzo DN, Badea A, Jiang YH, Yin HH, Wetsel WC, Bennett V. 2019. ANK2 autism mutation targeting giant ankyrin-B promotes axon branching and ectopic connectivity. Proc Natl Acad Sci U S A. 116:15262-15271.

Yizhar O. 2012. Optogenetic insights into social behavior function. Biol Psychiatry. 71:1075-1080.

Yoon S, Parnell E, Kasherman M, Forrest MP, Myczek K, Premarathne S, Sanchez Vega MC, Piper M, Burne THJ, Jolly LA, Wood SA, Penzes P. 2020. Usp9X Controls Ankyrin-Repeat Domain Protein Homeostasis during Dendritic Spine Development. Neuron. 105:506-521.

Zhou X, Feliciano P, Shu C, Wang T, Astrovskaya I, Hall JB, Obiajulu JU, Wright JR, Murali SC, Xu SX, Brueggeman L, Thomas TR, Marchenko O, Fleisch C, Barns SD, Snyder LG, Han B, Chang TS, Turner TN, Harvey WT, Nishida A, O'Roak BJ, Geschwind DH, Consortium S, Michaelson JJ, Volfovsky N, Eichler EE, Shen Y, Chung WK. 2022. Integrating de novo and inherited variants in 42,607 autism cases identifies mutations in new moderate-risk genes. Nat Genet. 54:1305-1319.