TITLE: PlantConnectome: knowledge graph encompassing >70,000 plant articles

AUTHORS: Shan Chun Lim[1], Kevin Fo[1], Rohan Shawn Sunil[1], Manoj Itharajula[1], Yu
Song Chuah[1], Herman Foo[1], Emilia Emmanuelle Davey[1], Melissa Fullwood[1], Guillaume
Thibault[1], Marek Mutwil[1*]

[1]School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive,
Singapore, 637551

*Corresponding author:
Marek Mutwil
School of Biological Sciences,
Nanyang Technological University,60 Nanyang Drive,
Singapore 637551
Email: mutwil@ntu.edu.sg

Abstract

One of the main quests of plant biology is understanding how genes and metabolites
work together to form complex networks that drive plant growth, development, and
responses to environmental stimuli. However, the ever-growing volume and diversity of
scientific literature make it increasingly challenging to stay current with the latest
advances in gene function studies. Here, we tackle the challenge by deploying the text-
mining capacities of large language models to process over 71,000 plant biology
abstracts. Our approach unveiled nearly 5 million functional relationships between a
wide array of biological entities—genes, metabolites, tissues, and others—with a high
accuracy of over 85%. We encapsulated these findings in PlantConnectome, a user-
friendly database, and demonstrated its diverse utility by providing insights into gene
regulatory networks, protein-protein interactions, and stress responses. We believe this
innovative use of AI in the life sciences will allow plant scientists to keep up to date with
the rapidly growing corpus of scientific literature. PlantConnectome is available at
https://plant.connectome.tools/.

Introduction

Despite decades of research, only ~15% of *Arabidopsis thaliana*'s genes have been
comprehensively characterized, and the rate of new articles reporting gene functions
has dropped to <30% in 2023 since the peak in 2008 (Sunil et al. 2024). Due to the
time-consuming experiments and increased requirements to publish in premier journals,
the time needed to characterize a gene can take several years. Thus, choosing which
gene to start characterizing requires a strong hypothesis, which is typically based on
previous work reported in the literature. However, staying up to date with the

continuously growing scientific literature, and integrating the numerous pieces of the
gene function puzzle can be time-consuming and limit our ability to form strong
hypotheses.

Alternatively, computational gene function prediction can suggest which genes
have a specific function and are invaluable in generating new gene function hypotheses
(Brown et al. 2005; Persson et al. 2005). Predicting gene function requires two
components: i) omics data that captures gene properties (e.g., coding sequence,
expression patterns, and protein structure) and ii), gold standard data (i.e., genes with
experimentally verified functions) (Radivojac et al. 2013; Rhee and Mutwil 2014). The
omics data is firstly used to connect uncharacterized and characterized genes based on
sequence or expression similarity. Then, uncharacterized genes are labeled according
to the functions of the characterized genes (i.e., the gold standard data) to which they
were connected (Rhee and Mutwil 2014).

Nonetheless, gene function prediction remains highly challenging due to the
complexity and vastness of biological data, plateauing our understanding of gene
functions (Radivojac et al. 2013). Specifically, establishing the gold standard
necessitates manual, work-intensive extraction of gene functional information from
scientific articles (Oughtred et al. 2021), preventing public repositories that harbor the
gold standard data, such as BioGRID (protein-protein interactions, or PPIs) and AGRIS
(gene regulatory networks, or GRNs)(Yilmaz et al. 2011; Oughtred et al. 2021), from
keeping up to date with state-of-the-art knowledge. Furthermore, such repositories are
typically restricted to specific data types (e.g., PPI or GRNs), precluding the integration
of various data kinds that is critical to deepening our understanding of plant biology.

Several methods that extract gene functional information from literature have
been developed to address these challenges. PL-PPF (Predicate Logic for Predicting
Protein Functions) uses statistical methods to infer if a protein and a molecular term that
describes protein function are semantically related (Taha et al. 2019). However, the
method requires constructing complex statistical and linguistic models to link protein to
function and only considers protein-function relationships. The EVEX database
processes abstracts and full texts to identify regulatory relationships, posttranslational
modifications, gene expression patterns and other features of genes (Landeghem et al.
2013). However, the method also requires a manually constructed complex set of rules
to extract and categorise the relationships, and the database has not been updated for
a while. Another approach uses non-negative matrix factorization (NMF) for feature
reduction and then classifies the function of genes using K-nearest neighbor
(KNN)(Fodeh and Tiwari 2018). While the approach can reveal gene functions (e.g.,
gene A is a transcription factor), it does not reveal gene-gene relationships (e.g., gene A
regulates gene B). STRING is a popular database that integrates protein-protein data,
genomic features, co-expression and text mining to build gene co-function networks
(Szklarczyk et al. 2023). However, the text mining approach only identifies genes that

are frequently mentioned together and cannot reveal the type of the relationship (e.g., interaction, regulation, activation) or identify relationships between genes and other entities (e.g., treatments, hormones). KnetMiner uses a rule-based approach to build knowledge graphs capturing relationships between various entities (Hassani-Pak et al. 2016). However, the rule-based system requires the integration of multiple heterogenous datasets (e.g., 12 types of data)(Hassani-Pak et al. 2016), making it difficult to include new data, species and evidence types into the network. Furthermore, the database is gene-centric and does not allow searching the knowledge graph to understand how the different types of entities are related (e.g., traits and hormones).

In this paper, we aim to address the two fundamental challenges in gene function prediction: integrating the burgeoning information from scientific literature and using it to generate gold standard data for gene function prediction approaches. To achieve this, we seized the recent developments in Large Language Models (LLMs) to process over 71,000 research papers from leading journals in plant biology. Our approach excavated 4.8 million functional relationships between more than 2.7 million entities comprising genes, metabolites, tissues, organs, and other biological components. The manual inspection of these relationships revealed not only their high accuracy but also their ability to identify functional relationships between biological entities, even doubling the amount of functional information relative to the current coverage of gene regulatory networks. To provide access to this data, we constructed PlantConnectome, a user-friendly database containing knowledge graphs that can illuminate gene function, organ development, gene regulatory networks, protein-protein interactions, and other biological entities. PlantConnectome is available at https://plant.connectome.tools/.

Materials and Methods
Retrieval of articles
Using BioPython version 1.81, we downloaded papers containing *Arabidopsis thaliana* species name and gene identifiers (e.g., *At4g32410*). For each gene identifier, we also searched with gene aliases (e.g., *CESA1, RSW1*) retrieved from www.arabidopsis.org (Table S1). The NCBI query was:
query = f'(Arabidopsis thaliana[Title/Abstract] AND {query_term}[tw]) OR (Arabidopsis[Title/Abstract] AND {query_term}[tw]) OR (Thale cress[Title/Abstract] AND {query_term}[tw] OR (Mouse ear cress[Title/Abstract] AND {query_term}[tw] OR (Mouse-ear cress[Title/Abstract] AND {query_term}[tw])'. Query_term are the genes in Table S1, and other alternative names of Arabidopsis were included in the search. The code to perform this analysis is available in the Colab Notebook in Supplementary Data 1.

Large language model analysis of articles

We used GPT4 models to extract entities and relationships (GPT-4o), entity definitions
(GPT-4o-mini) and to identify the species that the article uses as a model (GPT-4o).
Furthermore, for each entity relationship, we asked GPT-4o to identify the evidence
(e.g., yeast two-hybrid, bioinformatical prediction) underpinning the relationship. We
iterated over several prompts to arrive at prompts that yielded consistently accurate
results on selected papers, such as Brinngmann et al., 2012, and others. In total,
71,136 articles were processed using OpenAI's batch API (Application Programming
Interface)(https://openai.com/api/pricing/). The code to perform this analysis is available
as Colab Notebook in Supplementary Data 1.
129
Entity and relationship disambiguation
To disambiguate relationships (e.g., 'caused', 'cause', 'causes') and entities (e.g.,
'Arabidopsis plants', 'Arabidopsis', 'Arabidopsis thaliana'), we identified the top 100
most common relationships and entities and devised a rule-based method to map the
various synonyms or variations to a canonical form. Passive edges (e.g., 'is regulated
by') were converted to active form ('regulates'). Entities that differed by casing (e.g.,
'Genes', 'genes') were represented by one canonical form. The code used for this
section is available as Supplementary Data 1.
138
Construction of PlantConnectome database
The PlantConnectome is hosted on a Google Cloud server. The backend was
implemented using the Python framework Flask and the Python packages networkx
version 3.1, pickle version 3.11.4, json version 3.11.4, and regex *version 3.11.4*. We
used JavaScript dependencies jQuery v3.6, Cytoscape.js v3.23, ChartJS v4.3, and
FileSaver v2.0.5 to visualize the knowledge graphs. The GitHub repository containing
the   source   code   of   the   database   is   available   at
https://github.com/mutwil/plant_connectome_latest.git
147
API for PlantConnectome
PlantConnectome also has an application programming interface that allows users to
conduct search queries remotely. The API accepts GET requests and is implemented
using the same set of packages described earlier. For each successful call to
PlantConnectome's API, a JSON (JavaScript Object Notation) object is returned,
containing the functional abbreviations, GO terms, other nodes, and text summaries
associated with the search query. To perform searches using the API, users can add
"/api/<search type>/<search query>" to the web address, where "<search type>" and
"<search query>" are placeholders representing the type of search and user's query,
respectively.

Figure 1. Meta-analysis of the 71,136 article abstracts. Meta-analysis of plant

literature. A) Top 20 journals of the 71,136 articles analyzed in this study. The red and

blue bars indicate the abstract-only or full-text articles, respectively. B) Clustermap of top 30 journals (rows) and topics (columns). The colormap corresponds to the fraction of a maximum value found in a row (journal) across the 71,136 articles. C) The number of abstracts (x-axis) with a given keyword (y-axis) of the analyzed papers. Each abstract can contain multiple keywords. D) Co-occurrence network of keywords in abstracts. Nodes represent keywords, while edges connect keywords found in at least 600 abstracts. Edge width is proportional to the number of abstract where any two keywords are co-mentioned. E) t-SNE visualization of the abstracts with a focus on plant organs. Each point represents an article, and the colors indicate the different organs. The plots for F) 'membranes' and G) 'RNA biosynthesis' are shown to the right.

Results
Meta-analysis of 71,136 Paper Abstracts
To retrieve articles that focus on *Arabidopsis thaliana* genes and how these genes are related to other biological entities, we searched for articles that mention *Arabidopsis thaliana* and gene IDs in the abstracts (code available in Supplementary Data 1, gene IDs Table S1). In total, 71,136 articles, of which 19,809 and 51,327 were accessible as full-text articles or abstracts only, respectively (Figure 1A, Table S2). The top 20 journals comprise Plant Physiology, the Plant Journal and Plant Cell, for which most articles were not available for high-throughput download as full text (Figure 1A, red bars). Conversely, the open access policies and the option to programmatically download the articles of the Frontiers in Plant Science, PLOS One, New Phytologist, BMC Plant Biology and Scientific Reports allowed us to download full-text articles from these journals.

To investigate whether the top 20 journals tend to publish specific topics, we determined the surveyed journals' discussion of cellular compartments, organs, and biological functions to assess their considered research topics. To this end, we defined a list of keywords pertaining to organs (e.g., roots = [root, hair, nodule, mycorrhizae]), biological processes (photosynthesis = [photosynthesis, photorespiration, photosystem]), and cellular compartments (e.g., nucleus = [nucleus, nucleolus, chromosome, nuclear pore])(Table S3). We counted the number of these keywords in each abstract (Supplementary Data 2). Most journals did not show particular specificity for any topic, except Development (focus on reproduction, red cell), Journal of Biological Chemistry (membranes) and Molecular Plant-Microbe Interactions (external stimuli responses)(Figure 1B-C). The most commonly studied organs were: roots, leaves, flowers and seeds, and the most studied pathways were: phytohormone action (how hormones work), external stimuli response (how plants respond to the environment), RNA biosynthesis (how gene expression is regulated) and plant reproduction and most studied subcellular compartments were: membranes, and Golgi apparatus (Figure 1B-C). Next, we investigated which keywords tend to co-occur in abstracts (Table S4,

Supplementary Data 3), which revealed, e.g., that pathogen research focuses on salicylic acid and transcriptional responses and uses leaves and roots as model organs (Figure 1D).

To visualize the relationships among the abstracts, we generated a two-dimensional t-distributed Stochastic Neighbor Embedding (t-SNE) plot (Macosko et al. 2015), of the keyword counts (Supplementary Data 2). This technique allows us to represent high-dimensional data in a way that preserves local similarities between abstracts. We used a perplexity value of 40, which balances the attention between local and global aspects of the data, and ran 1,000 iterations to ensure convergence to a stable configuration (Figure S1 shows the influence of t-SNE parameters). The resulting plot provided an interpretable layout that highlighted clusters of abstracts with similar content or themes. The plots demonstrate clear groupings by biological processes (Figure S2), subcellular compartments (Figure S3) and organs (Figure S4), providing a bird's eye view of plant literature (Figure 1E).

Text Mining Research Papers with Large Language Model Reveals 4,819,469 Relationships between 2,771,008 Entities

To extract information pertaining genes, metabolites, organs, environmental conditions, and other entities, we tasked OpenAI's GPT models with identifying functional relationships between pairs of entities (e.g., 'gene A' - interacts with - 'gene B')(Figure 2A, prompt 1 with GPT-4o), and also identifying the types of each entity (genes, metabolites, organs, treatments, others)(Supplementary Data 1). The output of this analysis was a Knowledge Graph (KG), where nodes represent entities and edges represent relationships (e.g., 'interacts with', 'regulates', 'causes'). To better understand which types of evidence underpin each relationship (e.g., 'pull-down assay', 'co-expression analysis'), we also asked GPT-4o model to reveal the relationship basis and species the experiments were performed in (Figure 2A, prompt 2 with GPT-4o). Finally, we tasked GPT-4o-mini model to annotate the extracted entities (e.g., 'CESA' - is - 'Cellulose Synthase A')(Figure 2A, prompt 3 with GPT-4o-mini). The process yielded a large KG comprising 4,819,469 relationships between 2,771,008 entities (Supplementary Data 4, Table S5 contains the three prompts).

Large language models are known to hallucinate and misunderstand the text, and to evaluate the accuracy of the identified entity types, we randomly selected 300 edges from the KG (code for random selection in Supplementary Data 1). We manually evaluated whether the identified entity types (e.g., 'flavonol' type is 'metabolite') are correct, by comparing the entity types with their known biological function. Overall, we observed >90% accuracy in entity type classification, with the exception of 'subcellular compartment' and 'treatment' (Figure 2B, Table S6). The incorrectly classified 'subcellular compartment' entity types comprised genomic features such as 'distal enhancer elements', and protein domains ('13 TM helices')(Table S6), while incorrect

'treatment' types comprised methods and resources (e.g., 'microsomal preparations',
'Genbank').

To evaluate the accuracy of the edges, we manually compared them to the text
from which they were extracted. Furthermore, since GPT-4 models have an October
2023 knowledge cutoff and could have been trained on the analysed articles, we chose
edges from articles published before 2024 and from 2024. Overall, we observed a high
accuracy of 85% (before 2024, Figure 2B red bar) and 93% (2024, orange bar),
showing that the models can extract information from any text and not just regurgitate
training data. The incorrect relationships typically misunderstood a hypothesis of the
authors (source sentence: "(results) made us wonder whether this tissue-specific
polarization of PINs is conserved in other bryophytes"), incorrectly producing a fact
edge ('tissue-specific polarization of PINs conserved in other bryophytes')(Table S6).
Since the analyzed articles were shortlisted by term '*Arabidopsis thaliana*', the majority
of the relationships were identified in the model plant, but we also identified other
models and crops, such as rice, wheat, soybean, tobacco and even yeast (Figure S5).

To investigate the relationship between the number of articles and the number of
identified entities and relationships, we randomly removed 10-90% of articles 100 times,
and recounted the number of retrieved items. Overall, we observed a linear relationship
between the number of articles and the retrieved data (Figure 2C), indicating that more
articles would expand the KG further. The amount of information extracted from the
introduction (median 50 and 63 relationships and entities extracted, Figure 2D-E),
results (59, 63) and discussion (47, 57) was higher than from abstracts (16, 21). Thus,
increasing the number of full text articles would further expand the KG.

Finally, we investigated which types of evidence are present in the top 20
journals. We categorized the evidence into 'literature' (article citing findings from other
articles), 'advanced wet' (article using advanced experimental approaches, such as pull-
down, transgenic lines), 'high throughput' (evidence based on, e.g., RNA-seq analysis,
differential gene expression) and 'bioinformatics' (evidence based on sequence
alignment, phylogenetic tree, Supplementary Data 1 contains the used code and
keywords). Overall, the evidence profiles of the different journals were similar, with Plant
Cell and Plant Physiology on average using more advanced wet lab methods (advanced
wet > 0.35). Plant Science, Plant Physiology and Biochemistry contain extensive
literature-based evidence. In contrast, Methods in Molecular Biology which focuses
typically on one method, contained the least diversity of the used evidence types.

Figure 2. Evaluation of the plant knowledge graph. A) The pipeline to extract: 1. the knowledge graph from the literature, 2. species and relationship basis and 3. entity definitions. B) The percentage of correct entities (blue bars) and relationships (red and orange bars). The x-axis indicates the percentage of correct items inferred from manual curation. C) The number of unique entities (red bars, left y-axis) and relationships (i.e., number of edges, blue bars, right y-axis) as a function of % articles removed (x-axis). The error bars represent the standard deviation. The data was generated by randomly removing a given percentage of articles 100 times. D) and E) the number of relationships and entities extracted from abstracts, introductions, results and discussions, respectively. F) Average score depicting the diversity of methods extracted from the top 20 journals (Figure 1A). The method categories include 'literature' citations, 'advanced wet' lab techniques, 'high throughput' experiments and 'bioinformatics'. A

289 score of 1 indicates that a given journal has, on average, used all types of methods
290 under one category, while a score of 0 indicates that no methods have been used. A
291 method category is comprised of keywords. For example, 'bioinformatics' is comprised
292 of: 'sequence', 'phylogenetic', 'genomic', 'alignment', and 'differential expression'. The
293 code and keywords are available in Supplementary Data 4.
294
Properties of the Connectome Network
We used the KG to construct the Connectome network, and visual inspection of the
whole graph revealed cluster-like structures of densely connected entities (Figure 3A).
Certain networks, such as protein-protein interactions, display scale-free behavior,
where most nodes have few connections, and few nodes have many connections
(Broido and Clauset 2019). To investigate whether the Connectome is scale-free, we
constructed a scatterplot of its log-transformed node frequency ($p(k)$) and node degree
(k)(Figure 3B). The points formed a line with a negative slope, indicating a typical power
law distribution (Mutwil et al. 2010), indicating that most entities have a few
relationships, while a small number of entities act as hubs with a large number of
connections.
We next investigated which entities and relationships are most important in the
KG. General entities, such as 'A. thaliana' and 'plants', and general relationships, such
as 'had', 'show' were most frequently observed (Figure 3C), but we also observed more
specific entities (e.g., auxin, ABA) and relationships (regulate, interacts with). The most
common entity types comprised 'gene' and 'phenotype', in line with our selecting articles
containing gene names and '*Arabidopsis thaliana*' (Figure 3D). A closer look at the
entity types revealed the most common entities for genes (e.g., *FT, FLC, PIF4*),
phenotypes (growth, flowering, germination, gene expression), organism (A. thaliana),
metabolite (ABA, auxin, ethylene, ROS), protein (general transcription factors,
phytochromes, enzymes), mutants (general 'transgenic plants'), treatment (drought,
ABA, salt stress), process (photosynthesis, DNA methylation, autophagy), organ (root,
leaf) and subcellular compartment (nucleus, plasma membrane)(Figure 3E). Finally, we
investigated how often the different entity types are connected in the network. We
observed most connections between 'gene'--'phenotype', 'mutant'--'phenotype' and
'gene'--'treatment' (Figure 3F), likely reflecting the typical function studies that
characterize genes in terms of mutant phenotypes and responses to various treatments.

Figure 3. Properties of the Connectome knowledge graph. A) Gephi visualization of the Connectome knowledge graph colored by node degrees. The ForceAtlas 2 algorithm was run until convergence with a stronger gravity law and a scaling factor of 0.5 to separate the graph's nodes. Light blue nodes represent nodes with the fewest degrees, and light green nodes are the nodes with the highest degrees. B) The degree distribution of the knowledge graph, where nodes represent entities and edges represent relationships between these entities. The x-axis (degree (k)) represents the number of connections (relationships) each entity has. In contrast, the y-axis (frequency P(k)) shows the frequency of an entity having exactly k connections. C) Top most frequently-appearing entities (top) and relationship (bottom) types. The size of the

lettering is proportional to the number of relationships. D) Top 10 entity types, with entities (x-axis) and their numbers (y-axis). E) Top 20 entities found in the top 10 entity types. F) The number of edges between the top 10 entity types.

Evaluation of the coverage and accuracy of the PlantConnectome
Our main motivation in this study was to expand the amount of the gold standard data capturing experimentally-verified gene functions. We, thus, investigated the overlap of relationships in our KG with data provided in the public repositories.

To compare the coverage and the accuracy of gene regulatory networks (GRNs), we obtained the *Arabidopsis thaliana* gene regulatory network from AGRIS (https://agris-knowledgebase.org/downloads.html, updated March 2019) (Yilmaz et al. 2011), comprising 4,409 confirmed transcription factor -> target edges. We also identified 3,695 edges from a study investigating responses to jasmonic acid (Zander et al. 2020). Next, we identified 15,009 transcription factor -> target edges in the Connectome (Supplementary Data 4). The edges shared between AGRIS and the Connectome comprised of 'regulate', 'binds', 'activates', 'targets', demonstrating the Connectome's ability to identify the various functions of transcription factors (Figure 4A). However, we observed a very minor overlap (e.g., 206 edges between AGRIS and the Connectome) between the three GRNs (Figure 4B), indicating the high dissimilarity between the GRNs. We identified 14,736 Connectome-specific edges between transcription factors and target genes, and while the most frequent association was 'interacts with', we also identified typical transcription factor-related terms, such as 'regulate', 'binds', 'activates', 'represses', and others (Figure 4C). This indicates that the Connectome transcription factor networks seamlessly integrate protein-protein interactions (PPIs) and GRN networks.

Furthermore, we compared the protein-protein (PPI) network from BioGRID to the Connectome's network. We found 1,369 edges shared between BioGRID and the Connectome that were not of class 'interacts with, forms a complex with' (or similar), and found that the edges comprised of types such as 'phosphorylates', 'associates with', but also 'does not interact with' (Figure 4D). This indicates that the Connectome can provide additional nuances to the PPIs or even contradictory information that should be investigated further. Overall, we also found a relatively poor overlap between BioGRID and Connectome, with only 2,783 'interacts with' edges shared (Figure 4E), and the large majority of edges being specific to each database.

To validate the accuracy of the Connectome-specific edges, we randomly sampled 100 GRN (out of 14,736) and 100 PPI (out of 8,234) edges (code in Supplementary Data 1), and inspected their accuracy by reading the corresponding text (Table S7). Overall, we observed 97% and 94% accuracy for the GRN and PPI networks, respectively (Figure 4F). In agreement with the misclassified entities and relationships (Figure 2B), GPT-4o confused future directions (e.g., 'In this regard, it will

be interesting to investigate whether FUS3 interacts with IDD8 through the fourth ZF domain.) as a fact ('FUS3 interacts with IDD8')(Table S7). These results indicate Connectome's valuable companionship and alternative role to not only AGRIS but also BioGRID.

Figure 4. Evaluation of the gene regulatory and protein-protein interaction networks identified by the Connectome. A) Word cloud of the 206 relationship types shared between AGRIS and the Connectome. B) Venn diagram showing the overlap between AGRIS, the Connectome and the Jasmonic Acid Gene Regulatory Network (GRN). The numbers indicate edges found between transcription factors and putative target genes. C) Word cloud of the 14736 Connectome-specific edges. *D) Word cloud of the 1369 edges shared between BIOGRID and the Connectome. The edges do not belong to the 'interacts with group'. E) The Venn diagram shows the overlap between*

386 *BIOGRID and the Connectome. All gene-gene edges in Connectome are indicated with*
387 *a green circle, while the blue circle shows genes connected by 'interacts with'*
388 *relationships.* F) Accuracy evaluation of the protein-protein interaction (PPI) and gene
389 regulatory network (GRN) edges that are specific to the Connectome. The blue and red
390 fields indicate correctly inferred or 'future direction' relationships, respectively.
391
Features of PlantConnectome
To provide access to the Connectome, we constructed dedicated database
(https://plant.connectome.tools/), which offers numerous methods of searching for
genes, metabolites, organs, and other entities by terms, author names, and PubMed
IDs, alongside a catalogue page (accessible under the
https://plant.connectome.tools/catalogue) listing all entities in the database. An entire
information page is also provided for each entity in the connectome, containing its
definitions (e.g., CESA: 'A large family of genes encoding cellulose synthases and
related enzymes') and source article.
To detail PlantConnectome's search result page, we performed a standard query
with the gene "Psad1" ('Mutant affecting photosystem I complex in plants',
https://plant.connectome.tools/alias/psad1), which is involved in the formation of
photosystem I (Ihnatowicz et al. 2004). The entity's landing page displays the number of
nodes in the knowledge graph and the number of papers used to construct the KG,
together with the extracted definitions of the entity (Figure 5A).
The knowledge graph is represented as an interactive network, depicting the
various relationships the search query shares with other entities in the database (Figure
5B). Upon clicking on a node, the user is provided with a 'Node properties' tooltip
displaying the node's definitions, and a set of options enabling the removal of the node
or isolation of the node's neighborhood. Clicking on an edge opens an 'Edge properties'
tooltip that displays the PubMed ID underpinning an edge and shows the experimental
basis (if available) of the edge. Users can select the node and relationship types by
clicking on the 'Node select' and 'Edge select' tools and thus focus on the entity and
relationship types of interest (Figure 5B). The current network view can be downloaded
as an image (SVG) or as a tab-delimited table, ready for further processing in, e.g.,
Cytoscape. The network is also available as a text summary (Figure 5C) and a table
(Figure 5D), where clicking on a given PubMed ID will prompt a popup containing the
corresponding abstract. The text summary adjusts its content to the selections specified
in the 'Node select' and 'Edge select' tools.

Figure 5. Outline of the Connectome entity page. A) The top of the page contains the
entity name and the graph size, comprising the number of nodes (entities) and papers

that are used to build the current graph. Note that the graph size changes dynamically when the user adjusts the node and edge selection. The entity definitions extracted from the various papers mentioning the entity are found in the paginated table. B) The graph is visualized as an interactive network implemented in cytoscape.js, where the nodes can be moved around, and the different elements can be clicked on to open tooltips that display additional information. The entities in the network can be searched by writing the entity's name and clicking submit (blue button). The users can select the entity types (e.g., genes/proteins, metabolites) and edge types (e.g., regulates, interacts with) of interest by clicking on the Node and Edge select tools. The network can be downloaded as a vector graphic image and tab-delimited table. C) The text summary provides an organized, textual representation of the network and PubMed IDs that underpin each edge. The text summary is dynamic and responds to the node and edge selection performed by the user. D) The network is also summarized as a table, where each edge is found as a row in the table.

Finally, PlantConnectome enables users to perform searches through an API, which returns a JSON object containing relevant network and functional information, extending its functionality to bioinformaticians who desire programmatic access to our database. As an example, an alias search on the PSAD1 gene may be performed by accessing the URL "https://plant.connectome.tools/api/alias/psad1".

Examples of how to use PlantConnectome

We provide three case studies, comprising protein complexes, gene regulatory networks, and stress responses, to demonstrate how the Connectome can be used to rapidly summarize available knowledge.

Example 1: Chloroplast Protein Translocation and Channel Member TOC75

To exemplify how the Connectome can be used to study protein-protein interactions, we used TOC75 as a query for the 'alias' search (*https://plant2.connectome.tools/alias/TOC75*), which searched for entities labelled as *TOC75* and it's aliases: *AT3G46740, MAR-01, TOC75-III*. This identified a knowledge graph containing 355 nodes based on 80 papers. We then narrowed down the graph to genes/proteins with the 'Node selection' tool, resulting in 199 nodes based on 61 papers.

Translocase complexes on the outer and inner envelope membranes (TOC and TIC, respectively) are used to import proteins into the chloroplast (Stengel et al. 2009). We compared the TOP75 graph to a review on the translocase complexes (Richardson and Schnell 2020), which revealed known TOC75 interactions such as *TOC 22, 34, 159*, and *TIC236* (Figure 5A). The associated nodes also provide additional genes relevant for TOC75 function, such as dek5 mutant, which is reducing the levels of TOC75 (experimental organism: maize, evidence: proteomics analysis and immunoblotting of chloroplast envelope proteins)(Zhang et al. 2019), and chaperone

466 HSP90C that interacts with many of the translocon proteins (experimental evidence:
467 Coprecipitation experiments with protein import components)(Inoue et al. 2013). Thus,
468 the Connectome allows a rapid elucidation of protein-protein interactions, and provides
469 source literature and evidence types supporting these interactions.
470
Example 2: Secondary Cell Wall Master Regulator
To demonstrate how our database can be used to study gene regulatory networks, we
selected the secondary cell wall biosynthesis regulator, *MYB46*
(https://plant.connectome.tools/alias/MYB46). The initial network contained 618 nodes
based on 129 papers, but we selected edges capturing typical gene regulatory
relationships (e.g., 'regulate', 'directly activate') from the 'Edge filter' menu, and arrived
at a network comprising 211 nodes based on 75 papers (Figure 6B). A literature search
on the gene regulatory network underlying secondary cell wall formation revealed a
large overlap between the output of the Connectome and the figure in the review article
(Xiao et al. 2021).
Genes regulated by *MYB46* included the secondary cell wall cellulose synthases
(CESA4,7,8), lignin biosynthesis genes, and a panel of downstream transcription factors
(*MYB4, 7, 32, 43, 52, 54, 58, 63*). Genes regulating *MYB46* comprised transcription
factor *SND1* and micro-RNA miR395c (Figure 6B). Interestingly, our Connectome, but
not the review, identified *VND1-7* and *NST2* (evidence: Direct target analysis using the
estrogen-inducible system)(Zhong et al. 2008) and *VNDs* (evidence: Quantitative PCR
analysis of *VND1* overexpressors showing induction of *MYB46* gene expression)(Zhou
et al. 2014) as regulators of *MYB46*. On the other hand, our Connectome missed
downstream *MYB20, 42, 75, 103* (Xiao et al. 2021).
Since the KnowledgeNetwork is summarized in a human- and machine-readable
format, we investigated whether ChatGPT can generate publication-ready passages. To
this end, the network summary (exemplified in Figure 5C) was fed into ChatGPT-4o with
the following prompt:
'Summarize the function of MYB46, using the text below. Indicate which genes it
regulates. Indicate which genes regulate it. Indicate which processes it regulates. For
each statement, make sure to indicate the PubMed ID, for example: 'Gene A regulates
gene B (PubMed ID)'. Use only the information provided by this text:
498
499 *<Network summary text pasted in here>*
500
501 *Write the text as if it was a paragraph for a review paper. List ALL of the genes*
502 *mentioned in text.'*
503
504 The outcome is, in our opinion, on par with some review articles:
505

Figure 6. Usage examples of the Connectome database. A) Gene/protein neighborhood of TOC75 Translocon at the outer envelope-membrane of chloroplasts 75 protein, found by alias search using 'TOC75'. Nodes represent gene/protein entities, while edges depict relationships. The node and edge legend indicate the different types of entities and relationship types, respectively. B) Gene/protein neighborhood of MYB46. Transcription factors regulating MYB46 are indicated with red stars. C) Neighborhood of 'heat stress' entities. The nodes were selected to only comprise of 'chemical' entity type.

1) Function of MYB46

MYB46 (At5g12870) is a transcription factor that functions as a master regulator in secondary cell wall (SCW) biosynthesis (PMID: 24984711, 24009617). It directly regulates the expression of genes involved in the biosynthesis of cellulose, hemicellulose, and lignin (PMID: 24984711, 34493336). MYB46 plays a key role in activating the secondary wall biosynthetic program (PMID: 25148240), promotes SCW thickening (PMID: 33086706), and is crucial for secondary wall formation in fibers and xylem vessels (PMID: 25535195, 34879821). Furthermore, MYB46 has been shown to drive differentiation of xylem vessel cells (PMID: 34971462), enhance resistance to Botrytis cinerea (PMID: 32214345), and regulate plant immune response (PMID: 34208611). The protein also acts in abiotic and biotic stress responses, including its upregulation under stress (PMID: 31077628, 33826618). MYB46 functions in a transcriptional regulatory cascade (PMID: 19808805) and activates transcription of lignin, cellulose, and hemicellulose biosynthetic genes (PMID: 33537043). Its overexpression can induce ectopic secondary wall biosynthesis (PMID: 34493336), leading to stunted growth (PMID: 25535195).

2) Genes Regulating MYB46

MYB46 is regulated by several transcription factors. It is a direct target of SND1 (secondary wall-associated NAC domain protein 1), NST1, NST2, VND6, and VND7, which are part of the NAC transcription factor family involved in secondary wall biosynthesis (PMID: 37372306, 29973639). The VND family members (VND1-7) also induce the expression of MYB46 (PMID: 25148240, 35154217). Additionally, third-tier SWNs (NAC TFs) regulate MYB46 (PMID: 24009617), while MPK6-mediated phosphorylation is involved in post-translational regulation of MYB46 (PMID: 33826618). MYB46 is upregulated under stress by various stimuli (PMID: 33826618), and SnRK2.2/3/6 kinases regulate its expression during stress responses (PMID: 33495344). GbTCP5 (PMID: 34880892) and ERF041-VP16 (PMID: 33995450) are also involved in the transcriptional regulation of MYB46.

3) Genes Regulated by MYB46

MYB46 regulates a wide array of downstream genes involved in secondary wall biosynthesis. These include cellulose synthase genes such as CESA4, CesA7, and CesA8 (PMID: 23726771, 26011122, 37372306), as well as xylan and lignin biosynthetic genes (PMID: 34493336, 24885077). It activates the expression of other transcription factors like MYB58, MYB63, MYB85, and MYB43, which further control SCW biosynthesis (PMID: 24009617, 31881731). MYB46 also regulates genes involved in hemicellulose synthesis, such as AtCSLA9 and GAUT12 (PMID: 24243147, 25120548). Additionally, it directly activates secondary wall biosynthetic genes (PMID: 22197883, 34862967), including mannan synthase and other SCW-associated enzymes

556   (PMID: 24879533, 32775951). MYB46 also influences genes linked to lignin
557   biosynthesis, such as CSE and CCoAOMT7 (PMID: 27390589, 35154217), and
558   promotes SCW-related gene expression (PMID: 36077531).
559
560   Importantly, since ChatGPT was asked to use the existing PubMed IDs, it does not
561   hallucinate non-existing references.
562
Example 3: Heat stress
To demonstrate how the Connectome can be used to study entities that are not
necessarily genes, we investigated chemicals, hormones and metabolites involved in
heat stress. We searched for 'heat' with 'normal' search, which took us to the page of
entities   containing   'heat'   in   their   description
(https://plant2.connectome.tools/normal/heat). The resulting large knowledge graph
comprised 2069 nodes from 1294 papers, which is expected as heat stress is one of the
most studied abiotic stresses in plants (Koh et al. 2024). We further focused the search
by selecting 'Chemical' in the 'Node filter' tool, which shrank the graph to 103 nodes
based on 57 papers (Figure 6C). This revealed two central 'heat stress' nodes,
categorized as 'phenotype' and 'treatment'. The graph revealed multiple chemicals that
are important for heat stress tolerance, such as isoprene (found in Discussion in
(Weraduwage et al. 2023), evidence: Transcriptomic studies on Arabidopsis thaliana
fumigated with isoprene), trehalose (introduction in (Jin et al. 2016), no evidence
available), AITC (Allyl isothiocyanate, found in Introduction of (Øverby et al. 2015), no
evidence available) and flavonols and flavones (found in Discussion in (Liu et al. 2021),
evidence: LC-MS measurements on QE HF-X coupled to Vanquish UHPLC). The graph
also reveals compounds that change their levels under heat stress, such as
triacylglycerols (Introduction of (Higashi et al. 2015), evidence: LC-MS-based lipidomic
analysis), phosphatidic acid (Discussion in (Kocourková et al. 2021), no evidence
available), reactive oxygen species (ROS, Discussion in (Cocetta et al. 2022), evidence:
histochemical analysis using 3,3-diaminobenzidine (DAB) staining and literature
references). The graph also mentions several hormones, such as salicylic acid (SA),
auxin and ethylene, cellular structures such as cutin and starch content, and other
entities that were classified as chemicals by GPT-4o. To conclude, the dynamic
selection option of edge types in the network enables scrutinizing different relationship
types between the entities found in PlantConnectome.
590

Discussion
We have illustrated GPT's text mining capacities in the context of scientific literature,
processing over 71,000 research abstracts at a moderate cost (~5000 USD) and
harvesting invaluable functional information therein. GPT could extract key entities and
relationships from research paper abstracts with high accuracy (Figure 2B, 4E) and few

prompts (Table S5). The amount of functional information excavated from the abstracts
increased the amount of machine-readable data, as demonstrated by our gene
regulatory networks that nearly tripled the available data (Figure 4B). Moreover,
PlantConnectome overcomes the limitations of typical databases that employ only one
data type, as it draws upon numerous data sources in establishing gene functions,
organ development, gene regulatory networks, protein-protein interactions, and other
phenomena, all in a user-friendly manner.

Our evaluation has shown that PlantConnectome is not only comprehensive and
accurate but also complementary to existing databases (Figure 4). Comparing
PlantConnectome's gene regulatory networks against AGRIS and its protein-protein
interaction networks against BioGRID demonstrates that PlantConnectome's retrieved
networks do not largely overlap with these reference databases. Rather, the GPT-
extracted networks complement them, showing the effectiveness of our text-mining
approach in utilizing the vast amount of literature that has not been captured by manual
curation.

However, GPT's outputs are not entirely accurate and still warrant manual
verification, as GPT-4o models have a tendency to misidentify entities and relationships
(Figure 2B, 4E), which is perhaps attributable to the varying language and content of the
>71,000 processed articles. *The correction of errors may be carried out by fine-tuning*
the models with manually curated examples containing the expected output (as, for
instance, that found in Table S6). Thus, the users of our database and knowledge graph
are encouraged to click on nodes and edges to further validate these entities' accuracy.

In conclusion, PlantConnectome is an innovative tool, combining the power of a
state-of-the-art language model with the comprehensive information embedded in a
massive collection of research articles. The tool offers an efficient and diversified way to
retrieve information for genes, metabolites, tissues, organs, and other biological
components. The potential applications of PlantConnectome are wide-ranging and
extend beyond those we have highlighted in this article. Furthermore, since we only
analyzed articles mentioning *Arabidopsis thaliana* and it's genes, the inclusion of all
plant scientific literature together with the inclusion of more full-text papers is bound to
increase the completeness of the knowledge graph, help us stay up to date with the
plant literature, and provide gold standard data for gene function prediction studies. We
anticipate that PlantConnectome will become a valuable resource for the plant science
community to facilitate various research activities, from a preliminary investigation of
gene functions to an in-depth study of a particular biological process.

631
632

633 **Supplemental Figures**

634
635 **Figure S1. tSNE analysis of the abstracts at the different perplexity and iteration**
636 **values.** The evolution of the plot at a perplexity of 5 (first row), 50 (second row) and 100
637 (third row) and different ranges of iterations: 10 (first column), 100 (second column) and
638 1000 (third column).

639
640
641 **Figure S2. tSNE analysis of the abstracts of the different biological processes, as**
642 **defined by MapMan.** A red point indicates an abstract that contains a keyword (e.g.,
643 pollen is a keyword for plant reproduction), while grey point indicates an absence of the
644 keyword match.
645

Figure S3. tSNE analysis of the abstracts of the different cellular compartments.

649
650 **Figure S4. tSNE analysis of the abstracts of the different major organs and cell**

651 **types.**

652
653 **Figure S5. Wordcloud depicts the species in which the relationships (edges) were**
654 **reported in.**
655
656
Supplemental Tables
Table S1. Gene IDs (column A) and aliases (column B) used to query the NCBI
database for articles.
Table S2. Journals used to construct the database. The columns indicate the journal
name, article title, publication year, PubMed ID, and whether the article was analyzed
as full text (yes/no).
Table S3. The three categories, their sub-categories, and keywords used to
perform topic analysis of paper abstracts.
Table S4. Keyword co-occurrence analysis. The two keywords are in columns A and
B, and the number of articles in which these keywords co-ocurred are shown in column
C.
Table S5. The prompts used to build the knowledge graph, extract entity
definitions and edge basis.
Table S6. Edge and entity type accuracy evaluation. Each row represents and edge
in the knowledge graph. Each edge contains the PubMed ID, source and target nodes,
source and target entity types and relationship description. The evaluations of the
source types (column G), target types (Column J), and relationship (Column M) are
indicated. The comments and the sentences that underpin the edge are found in the
other columns.
Table S7. Evaluation of the accuracies of the gene regulatory edges (top sub-
table) and protein-protein interactions (bottom sub-table).
678
679 **Supplementary Data**

Supplementary Data 1. The code to download and process the papers, and generate the figures in the manuscript is available at https://colab.research.google.com/drive/1gENxLK2172Bq1sV_dO6JhvORsFgaq0fS?usp=sharing

Supplementary Data 2. Keywords present in each abstract analyzed in this study. The keywords are defined in Table S3. https://figshare.com/ndownloader/files/49392538

Supplementary Data 3. Co-occurence network of keywords found in each abstract. The network can be viewed in Cytoscape. https://figshare.com/ndownloader/files/49392595

Supplementary Data 4. The knowledge graph used to build the Plant Connectome. The edges and nodes have been disambiguated by the code found in Supplementary Data 1. https://figshare.com/ndownloader/files/49198933

Data availability

The Plant Connectome database source code is available at: https://github.com/mutwil/plant_connectome_latest/

References

Broido AD and Clauset A. Scale-free networks are rare. Nat Commun. 2019:**10**(1):1017. https://doi.org/10.1038/s41467-019-08746-5

Brown DM, Zeef LAH, Ellis J, Goodacre R, and Turner SR. Identification of novel genes in Arabidopsis involved in secondary cell wall formation using expression profiling and reverse genetics. Plant Cell. 2005:**17**(8):2281–2295. https://doi.org/10.1105/tpc.105.031542

Cocetta G, Landoni M, Pilu R, Repiso C, Nolasco J, Alajarin M, Ugena L, Levy CCB, Scatolino G, Villa D, et al. Priming Treatments with Biostimulants to Cope the Short-Term Heat Stress Response: A Transcriptomic Profile Evaluation. Plants (Basel). 2022:**11**(9):1130. https://doi.org/10.3390/plants11091130

Fodeh SJ and Tiwari A. Exploiting MEDLINE for gene molecular function prediction via NMF based multi-label classification. Journal of Biomedical Informatics. 2018:**86**:160–166. https://doi.org/10.1016/j.jbi.2018.08.009

Hassani-Pak K, Castellote M, Esch M, Hindle M, Lysenko A, Taubert J, and Rawlings C. Developing integrated crop knowledge networks to advance candidate gene discovery. Applied & Translational Genomics. 2016:**11**:18–26. https://doi.org/10.1016/j.atg.2016.10.003

Hassani-Pak K, Singh A, Brandizi M, Hearnshaw J, Parsons JD, Amberkar S, Phillips AL, Doonan JH, and Rawlings C. KnetMiner: a comprehensive approach for supporting evidence-based gene discovery and complex trait analysis across species. Plant Biotechnol J. 2021:**19**(8):1670–1678. https://doi.org/10.1111/pbi.13583

Higashi Y, Okazaki Y, Myouga F, Shinozaki K, and Saito K. Landscape of the lipidome and transcriptome under heat stress in Arabidopsis thaliana. Sci Rep. 2015:**5**(1):10533. https://doi.org/10.1038/srep10533

Ihnatowicz A, Pesaresi P, Varotto C, Richly E, Schneider A, Jahns P, Salamini F,

and Leister D. Mutants for photosystem I subunit D of Arabidopsis thaliana:
Effects on photosynthesis, photosystem I stability and expression of nuclear
genes for chloroplast functions. Plant Journal. 2004:**37**(6):839–852.
https://doi.org/10.1111/j.1365-313X.2004.02011.x

Inoue H, Li M, and Schnell DJ. An essential role for chloroplast heat shock protein 90
(Hsp90C) in protein import into chloroplasts. Proc Natl Acad Sci U S A.
2013:**110**(8):3173–3178. https://doi.org/10.1073/pnas.1219229110

Jin Q, Hu X, Li X, Wang B, Wang Y, Jiang H, Mattson N, and Xu Y. Genome-Wide
Identification and Evolution Analysis of Trehalose-6-Phosphate Synthase Gene
Family in Nelumbo nucifera. Front Plant Sci. 2016:**7**:1445.
https://doi.org/10.3389/fpls.2016.01445

Kocourková D, Kroumanová K, Podmanická T, Daněk M, and Martinec J.
Phospholipase Dα1 Acts as a Negative Regulator of High Mg2+-Induced Leaf
Senescence in Arabidopsis. Front Plant Sci. 2021:**12**:770794.
https://doi.org/10.3389/fpls.2021.770794

Koh E, Sunil RS, Lam HYI, and Mutwil M. Confronting The Data Deluge: How Artificial
Intelligence Can Be Used in the Study of Plant Stress. Computational and
Structural Biotechnology Journal. 2024.
https://doi.org/10.1016/j.csbj.2024.09.010

Landeghem SV, Björne J, Wei C-H, Hakala K, Pyysalo S, Ananiadou S, Kao H-Y,
Lu Z, Salakoski T, Peer YV de, et al. Large-Scale Event Extraction from
Literature with Multi-Level Gene Normalization. PLOS ONE. 2013:**8**(4):e55814.
https://doi.org/10.1371/journal.pone.0055814

Liu B, Kong L, Zhang Y, and Liao Y. Gene and Metabolite Integration Analysis
through Transcriptome and Metabolome Brings New Insight into Heat Stress
Tolerance in Potato (Solanum tuberosum L.). Plants (Basel). 2021:**10**(1):103.
https://doi.org/10.3390/plants10010103

Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas
AR, Kamitaki N, Martersteck EM, et al. Highly Parallel Genome-wide
Expression Profiling of Individual Cells Using Nanoliter Droplets. Cell.
2015:**161**(5):1202–1214. https://doi.org/10.1016/j.cell.2015.05.002

Mutwil M, Usadel B, Schütte M, Loraine A, Ebenhöh O, and Persson S. Assembly of
an interactive correlation network for the Arabidopsis genome using a novel
Heuristic Clustering Algorithm. Plant Physiology. 2010:**152**(1):29–43.
https://doi.org/10.1104/pp.109.145318

Oughtred R, Rust J, Chang C, Breitkreutz B, Stark C, Willems A, Boucher L,
Leung G, Kolas N, Zhang F, et al. The BioGRID database: A comprehensive
biomedical resource of curated protein, genetic, and chemical interactions.
Protein Sci. 2021:**30**(1):187–200. https://doi.org/10.1002/pro.3978

Øverby A, Stokland RA, Åsberg SE, Sporsheim B, and Bones AM. Allyl
isothiocyanate depletes glutathione and upregulates expression of glutathione S-
transferases in Arabidopsis thaliana. Front Plant Sci. 2015:**6**:277.
https://doi.org/10.3389/fpls.2015.00277

Persson S, Wei H, Milne J, Page GP, and Somerville CR. Identification of genes
required for cellulose synthesis by regression analysis of public microarray data
sets. Proceedings of the National Academy of Sciences of the United States of

America. 2005:**102**(24):8633–8638. https://doi.org/10.1073/pnas.0503392102

Radivojac P, Clark WT, Oron TR, Schnoes AM, Wittkop T, Sokolov A, Graim K, Funk C, Verspoor K, Ben-Hur A, et al. A large-scale evaluation of computational protein function prediction. Nature Methods. 2013:**10**(3):221–227. https://doi.org/10.1038/nmeth.2340

Rhee SY and Mutwil M. Towards revealing the functions of all genes in plants. Trends in Plant Science. 2014:**19**(4):212–221. https://doi.org/10.1016/j.tplants.2013.10.006

Richardson LGL and Schnell DJ. Origins, function, and regulation of the TOC–TIC general protein import machinery of plastids. Journal of Experimental Botany. 2020:**71**(4):1226–1238. https://doi.org/10.1093/jxb/erz517

Stengel A, Benz JP, Buchanan BB, Soll J, and Bölter B. Preprotein Import into Chloroplasts via the Toc and Tic Complexes Is Regulated by Redox Signals in Pisum sativum. Molecular Plant. 2009:**2**(6):1181–1197. https://doi.org/10.1093/mp/ssp043

Sunil RS, Lim SC, Itharajula M, and Mutwil M. The gene function prediction challenge: large language models and knowledge graphs to the rescue. arXiv.org. 2024. https://arxiv.org/abs/2408.07222v1. Retrieved September 26, 2024

Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, Gable AL, Fang T, Doncheva NT, Pyysalo S, et al. The STRING database in 2023: protein–protein association networks and functional enrichment analyses for any sequenced genome of interest. Nucleic Acids Research. 2023:**51**(D1):D638–D646. https://doi.org/10.1093/nar/gkac1000

Taha K, Iraqi Y, and Al Aamri A. Predicting protein functions by applying predicate logic to biomedical literature. BMC Bioinformatics. 2019:**20**(1):71. https://doi.org/10.1186/s12859-019-2594-y

Weraduwage SM, Sahu A, Kulke M, Vermaas JV, and Sharkey TD. Characterization of promoter elements of isoprene-responsive genes and the ability of isoprene to bind START domain transcription factors. Plant Direct. 2023:**7**(2):e483. https://doi.org/10.1002/pld3.483

Xiao R, Zhang C, Guo X, Li H, and Lu H. MYB Transcription Factors and Its Regulation in Secondary Cell Wall Formation and Lignin Biosynthesis during Xylem Development. International Journal of Molecular Sciences. 2021:**22**(7):3560. https://doi.org/10.3390/ijms22073560

Yilmaz A, Mejia-Guerra MK, Kurz K, Liang X, Welch L, and Grotewold E. AGRIS: the Arabidopsis Gene Regulatory Information Server, an update. Nucleic Acids Res. 2011:**39**(Database issue):D1118-1122. https://doi.org/10.1093/nar/gkq1120

Zander M, Lewsey MG, Clark NM, Yin L, Bartlett A, Saldierna Guzmán JP, Hann E, Langford AE, Jow B, Wise A, et al. Integrated multi-omics framework of the plant response to jasmonic acid. Nat Plants. 2020:**6**(3):290–302. https://doi.org/10.1038/s41477-020-0605-7

Zhang J, Wu S, Boehlein SK, McCarty DR, Song G, Walley JW, Myers A, and Settles AM. Maize defective kernel5 is a bacterial TamB homologue required for chloroplast envelope biogenesis. J Cell Biol. 2019:**218**(8):2638–2658. https://doi.org/10.1083/jcb.201807166

Zhong R, Lee C, Zhou J, McCarthy RL, and Ye Z-H. A Battery of Transcription
Factors Involved in the Regulation of Secondary Cell Wall Biosynthesis in
Arabidopsis. Plant Cell. 2008:**20**(10):2763–2782.
https://doi.org/10.1105/tpc.108.061325
Zhou J, Zhong R, and Ye Z-H. Arabidopsis NAC domain proteins, VND1 to VND5, are
transcriptional regulators of secondary wall biosynthesis in vessels. PLoS One.
2014:**9**(8):e105726. https://doi.org/10.1371/journal.pone.0105726

A

Relationship types of the 206 shared edges

B

AGRIS

4201

14736

3616

JA GRN

Connectome - TF->Target

D

Relationship types of 1369 not 'interacts with' edges shared with BioGRID

E

BioGRID

Connectome - interacts with edges

69143

2783

8234

1369

35637

Connectome - all edges

C

Relationship types of 14736 Connectome-specific edges

F

Accurate   Future direction

PPI (100)

GRN (100)

0   20   40   60   80   100

% correct

A

Search results for: PSAD1 ← Entity name

Graph size: 98 nodes; Edges are based on: 17 paper(s) ← Graph size: (nodes, edges, source papers)

For small to medium networks (fewer than 500 nodes), the random layout is applied by default. Larger networks are loaded with random layout. Tip: You can apply filters to reduce the size of large networks.

Extracted Definitions (Page 1):

- A mutant allele of the PSAD1 gene, impacting photosystem I stability. (PMID: 14996217)
- Thylakoids from the psad1-1 mutant, adapted to light conditions. (PMID: 17968587)
- Mutant affecting photosystem I complex in plants (PMID: 22639613)
- A specific mutant of Arabidopsis thaliana with photosystem I defects. (PMID: 17968587)
- A gene encoding the D-subunit of photosystem I in Arabidopsis thaliana. (PMID: 14996217)

Entity definitions

B

Network search tool (not matching nodes become transparent)

Search a node by its name

| Submit | Go back | Download as SVG | Download complete network as TSV | ← Download network (as graphics (svg), table (tsv)) |

Node properties (definition extracted from text)

Edge properties (Experimental/computational basis extracted from text)

Node select tool (e.g., gene, metabolite, organ)

Edge select tool (e.g., interacts with, enhances)

C

Text summary of the network:

PSAD1 [mutant] has the following relationships: exhibit retarded growth [phenotype] (PMID: 22639613), light green phenotype [phenotype] (PMID: 22639613), reduction of chlorophylls and carotenes [phenotype] (PMID: 22639613), increased antheraxanthin [metabolite] (PMID: 22639613), increased tyrosine [metabolite] (PMID: 22639613), increased threonine [metabolite] (PMID: 22639613), increased asparagine [metabolite] (PMID: 22639613), increased histidine [metabolite] (PMID: 22639613), increased alpha-tocopherol [metabolite] (PMID: 22639613), increased gamma-tocopherol [metabolite] (PMID: 22639613); confirmed stunted phenotype [phenotype]

D

Table summary of the network:

Source	Interaction Type	Target	Section	Pubmed ID
Photosystem I reaction center subunit II-1 (psaD1) [protein]	involved in	photosynthesis [process]	37076046_results2	37076046
Transcription of gene(s) of psaD1/D2 [gene]	was up-regulated at	T1 vs T0 [treatment]	26865323_discuss1	26865323

A

Nodes Legend:
- Gene/Protein (blue circle)
- Phenotype (green square)
- Cell/Organ/Organism (pink circle)
- Chemical (yellow diamond)
- Treatment (light blue square)
- Process (purple square)
- Method (brown triangle)
- Genomic/Transcriptomic feature (teal square)
- Biological process (green diamond)
- Others (orange star)

Edges Legend:
- Activation/Induction/Causation/Result (green)
- Repression/Inhibition/Negative Regulation (red)
- Regulation/Control (blue)
- Expression/Detection/Identification (orange)
- Association/Interaction/Binding (purple)
- Localization/Containment/Composition (green)
- Requirement/Activity/Function/Participation (dark red)
- Encoding (purple)
- Synthesis/Formation (orange)
- Others (gray)

B

C

Graph size: 2069 nodes; Edges are based on: 1294 paper(s)

↓

Node select: 'Chemicals'

Graph size: 103 nodes; Edges are based on: 57 paper(s)