A flexible high-throughput cultivation protocol to assess the response of individuals' gut microbiota to diet-, drug-, and host-related factors

Janina N. Zünd[1], Serafina Plüss[1], Denisa Mujezinovic[1], Carmen Menzi[1,2], Philipp R. von Bieberstein[1,2], Tomas de Wouters[2], Christophe Lacroix[1], Gabriel E. Leventhal[2], Benoit Pugin[1,3,*]

[1] Laboratory of Food Biotechnology, Department of Health Sciences and Technology, ETH Zürich, Zürich, Switzerland

[2] PharmaBiome AG, Zürich, Switzerland

[3] Lead contact

* Correspondence: benoit.pugin@hest.ethz.ch

SUMMARY

The anaerobic cultivation of fecal microbiota is a promising approach for investigating how gut microbial communities respond to specific intestinal conditions and perturbations (e.g., altered gut physiology, diet, and drugs). Here, we describe a simple high-throughput protocol using 96-deepwell plates to cultivate stool-derived gut microbiota. Our protocol addresses key challenges in high-throughput culturing within an anaerobic chamber, including a thorough assessment of the impact of the gas phase on the medium's chemistry, a modular medium preparation process to enable the testing of several conditions in parallel, a medium formulation design to maximize the compositional similarity of fecal cultures with the donor microbiota, and the creation of a step-by-step protocol detailing all practical procedures from material preparation until sample handling for analyses. Finally, we validated our cultivation protocol by demonstrating that cultivated fecal microbiotas responded similarly to dietary fibers (resistant dextrin, soluble starch) and drugs (ciprofloxacin, 5-fluorouracil) as reported in vivo. This simple high-throughput cultivation protocol has the potential to facilitate culture-dependent studies, accelerate the discovery of gut microbiota-diet-drug-host interactions, and pave the way toward personalized and microbiota-centered interventions.

Keywords: high-throughput cultivation, gut microbiota, gut microbes, ex vivo, ecology, physiology, microbial community dynamics, anaerobic chamber, Hungate technique, 16S rRNA sequencing, short-chain fatty acids,

MOTIVATION

The human gut microbiota is a complex ecosystem unique to each individual. The extent of this diversity has limited our capacity to fully comprehend microbial dynamics that apply to the entire human population. To probe the response of donor-specific microbial communities to certain intestinal conditions or perturbations, *in vitro* cultivation of stool-derived gut microbiota can be employed. However, the cultivation of gut microbiota in strictly anaerobic conditions is commonly performed using individual gas-tight tubes, a highly time-consuming strategy limiting the number of conditions and donor microbiota that can be tested in parallel. The development of a flexible high-throughput protocol to cultivate and test donor-specific gut microbiota is therefore required. Hence, we developed a robust procedure for cultivating stool-derived microbiota (and pure gut microbial cultures) in 96-deepwell plates within an anaerobic chamber.

INTRODUCTION

The human gut is inhabited by a diverse and dynamic ecosystem of microbes, unique to each individual[1], and contributing to key health functions such as nutrient metabolism, pathogen colonization resistance, maintenance of gut barrier integrity, and education of the host immune system, among others[2]. Despite advances in understanding human gut microbiota's diversity and functionality, the mechanisms governing ecological and physiological dynamics in response to specific intestinal conditions and perturbations remain poorly understood. Numerous studies have documented changes in the gut microbial community in response to dietary interventions[3–5], antibiotic/non-antibiotic medications[6–8], or alterations of the host physiology (*e.g.,* intestinal pH, oxidative stress, transit time, antimicrobial peptide production, and osmotic pressure)[9–11]. However, high inter- and intra-individual variability of the gut microbiota combined with the complex interactions between diet-, drug-, and host-related factors *in vivo* have limited our capacity to generalize microbial responses to a broader population[12].

In vitro fecal cultivation holds promise for investigating the direct impact of relevant intestinal conditions on individuals' gut microbiota. However, the traditional cultivation of intestinal communities using anaerobic tube-based techniques (*e.g.,* Hungate tubes, Balch tubes or serum flasks)[13–15] is highly time-consuming and limits the number of donor microbiota and conditions that can be evaluated simultaneously. This is a major drawback as many responses and activities are specific to the donor[16–19], the strain[20–23], and the physicochemical environment of the gut[9–11]. Cultivation with a higher throughput

is thus essential to expand our understanding of gut microbiota-diet-drug-host interactions and facilitate the development of personalized microbiota-centered interventions[4,24]. Although high-throughput cultivation in multi-well plates is not new[25–28], several gaps remain to be addressed. The effect mediated by the anaerobic chamber gas phase on the chemistry of the growth medium and on microbial physiology has not been directly evaluated. The development of a highly modular medium preparation process for simultaneously testing several culture conditions has yet to be established. The formulation of a standardized growth medium maintaining the microbial composition of the donor feces in fecal cultures is still required. Lastly, a cultivation protocol detailing all technical procedures and practical solutions is still lacking, which is crucial to ensure ease of implementation and reproducibility in cultivation experiments.

In this study, we report on the development and validation of a simple, flexible, and reproducible high-throughput protocol to cultivate stool-derived gut microbiota and pure intestinal microbes in 96-deepwell plates within an anaerobic chamber. This protocol builds upon a previous tube-based bacterial enrichment study, where specific bacterial taxa were associated with the functional niches of the human gut[24]. Here, we transitioned the enrichment tube-based conditions to a high-throughput format, and we expanded the applicability of the protocol for testing microbiota responses to a range of diets- (*e.g.,* fibers, nitrogen sources, vitamins), drugs- (*e.g.,* antibiotics and non-antibiotic medication), and host-related factors (*e.g.,* pH, redox potential, oxidative stress). We validated our protocol by showing similar *in vitro* responses to known dietary fibers (resistant dextrin, soluble starch) and drugs (ciprofloxacin, 5-fluorouracil) as reported *in vivo.* A comprehensive protocol detailing the necessary steps and procedures for material preparation, high-throughput cultivation, and sample handling for analyses is provided in **Supplementary file 1.**

RESULTS

Optimizing the procedures for high-throughput cultivation within the anaerobic chamber

We aimed to create a standardized high-throughput cultivation protocol that is compatible with standard anaerobic chamber setups and utilizes commonly available laboratory equipment. The first critical steps to transition our previous tube-based cultivation conditions[24] to a high-throughput format included i) selecting a versatile and reusable multi-well vessel, ii) designing a robust yet modular medium preparation process, and iii) carefully assessing confounding factors associated with cultivation in the anaerobic chamber.

To replace the gas-tight rubber-stoppered tubes used with the gold-standard Hungate technique[14], we opted for 96-deepwell plates combined with gas-permeable sealing membranes. The use of 96-deepwell plates enables the simultaneous testing of multiple conditions, with 60 wells allocated for experiments surrounded by 36 water-filled wells to minimize evaporation and edge effects[29] (**Supplementary file 1, Step 4**). The working volume of 2 mL per well provides sufficient samples to conduct a range of analyses on the whole culture (*e.g.,* OD_{600}, pH), the cell-free supernatant (*e.g.,* metabolites), and the cell pellet (*e.g.,* DNA-based analyses) (**Supplementary file 1, Step 8**).

To prepare the medium and culture conditions, we designed a modular procedure in which an anaerobic basal medium solution (containing the nitrogen (N)-sources, fatty acids, salts, buffers, trace elements, and heat-stable vitamins) can be flexibly complemented within the chamber with a carbon (C)-sources solution, or any other solution with the components (heat stable/sensitive) to be tested (**Figure 1A; Supplementary file 1**). For the basal medium, we formulated bYCFA and mM2, which are modified, C-depleted versions of yeast extract casitone and fatty acids (YCFA) and M2GSC media, respectively, two widely used media for cultivating gut microbes and microbiota[30,31].

One first challenge when cultivating intestinal microbes within an anaerobic chamber is the maintenance of strict anaerobiosis. During plate-based cultivation, cultures are exposed to the chamber's atmosphere, which can be contaminated with O_2 that diffuses in or is introduced with consumables. O_2 is removed by reacting with H_2 and the palladium catalyst to produce H_2O. To show the impact of incomplete O_2 removal on bacterial growth, we cultivated two intestinal bacteria, one oxygen tolerant (*Bacteroides thetaiotaomicron* DSM 2079) and one strict anaerobic (*Faecalibacterium prausnitzii* DSM 17677), in a "strict" and "loose" O_2-handling regime. In the loose regime, the chamber was improperly prepared with non-rejuvenated catalysts and low H_2 (<1.5%), whereas the strict regime had rejuvenated catalysts and H_2 was >2.5%. The growth rate (μ_{max}) of *B. thetaiotaomicron* was significantly higher in the strict compared to the loose regime (0.75±0.02 vs. 0.65±0.03 h^{-1}; p<0.05). Similarly, the maximal OD_{600} for *F. prausnitzii* was significantly higher in the strict compared to the loose regime (0.27±0.02 vs. 0.06±0.04; p<0.01) (**Figure 1B; Table S1**). This illustrates that meticulously preparing the chamber (**Supplementary file 1, Step 2**) is crucial for reproducing anaerobic microbial growth.

A second challenge during cultivation in the anaerobic chamber is the control of pH, which is influenced by the equilibrium between gaseous and dissolved CO_2, with the latter forming carbonic acid (H_2CO_3) in an aqueous solution[32]. When preparing the basal medium, 100% CO_2 or N_2 is flushed to remove traces of O_2, resulting in dissolved CO_2 levels that are out of equilibrium with the chamber atmosphere (10% CO_2, 5% H_2, and 85% N_2).

Figure 1: Overview of the high-throughput cultivation protocol and effect of the anaerobic chamber atmosphere on the medium's chemistry and microbial physiology

A) Overview of the step-by-step cultivation protocol adapted for the anaerobic chamber setup. In a 96-deepwell plate, heat-stable/sensitive supplements (1) are mixed with HCl-titrated bYCFA (2) and inoculated with diluted feces (3). **B)** Growth kinetics (OD_{600}) of *B. thetaiotaomicron* DSM 2079 and *F. prausnitzii* DSM 17677 in bYCFA (30 mM glucose, pH 6.5, 37°C, 32 h) in an anaerobic chamber that was handled "strictly" (turquoise dots; >2.5% H_2) and "loosely" (coral dots; <1.5% H_2). Each condition was tested in biological triplicates. **C)** Characterization of the pH dynamics of CO_2 and N_2-flushed bYCFA (uninoculated) in a 96-deepwell plate in an anaerobic chamber. **D)** Impact of the gas phase of the anaerobic chamber on the pH of basal medium previously flushed with CO_2 or N_2. **E)** Relationship between pH, aqueous $NaHCO_3$ concentration, and gaseous CO_2 levels (~10% in the anaerobic chamber, ~100% in gas-tight tube;) in bYCFA after 48 h equilibration. **F)** Change of pH over time upon titration of bYCFA with HCl, with subsequent pH increase over 48 h. The resulting titration curve (at time 0 h) can be used to adjust the pH to different values. **G)** pH and growth (OD_{600}) during 48 h batch cultivation in bYCFA (initial pH 6.8 and 5.8; containing 4 g/L $NaHCO_3$ and 3 g/L resistant dextrin) titrated with HCl immediately before inoculation with a fecal microbiota (donor CDE; 1% inoculation of 10^{-4} fecal dilution). Each condition was tested in technical triplicates.

As a result, introducing CO_2-flushed medium in the chamber led to an abiotic increase of pH (+0.16±0.05 after 4 h, and +0.83±0.03 after 48 h in bYCFA). In contrast, a rapid drop of pH was observed when introducing N_2-flushed medium in the chamber (-0.36±0.02 after 4 h in bYCFA, **Figure 1C**; similar pH dynamics in mM2, **Figure S1**). These changes in pH indicate the release of CO_2 into the gas phase (CO_2-flushed) or the solubilization of CO_2 into the medium (N_2-flushed) to equilibrate with the levels in the chamber's atmosphere (**Figure 1D**). Regardless of the gas employed during media preparation, it is thus recommended to equilibrate in the chamber for at least 24 h prior to the experiment (**Supplementary file 1, Step 3**). The pH after equilibrium will depend on CO_2 level in the atmosphere as well as the concentration of $NaHCO_3$ in the medium, *i.e.,* the conjugate base of H_2CO_3 in the bicarbonate buffer system (Davenport diagram, **Figure 1E**). To establish a standard $NaHCO_3$-containing basal medium adapted to a range of CO_2-containing anaerobic chambers and that can be flexibly adjusted to a range of pH values, we chose a fixed concentration of 4 g/L $NaHCO_3$, and opted to adjust pH within the chamber via HCl titration (**Supplementary file 1, Step 6**). This approach allows for the rapid adjustment of the initial pH for cultivation, though with a gradual increase in pH over time (**Figure 1F**) as HCl titration disbalances the bicarbonate buffering system, leading to H_2CO_3 formation and CO_2 release. Therefore, pH adjustment should be performed right before inoculation. To validate the feasibility of this approach, we titrated bYCFA with HCl (to pH 6.8 and pH 5.8) immediately before inoculation with a fecal microbiota and incubated the cultures at 37°C for 48 h (**Figure 1G**). After an initial pH increase of +0.48±0.07 (pH 6.8) and +0.45±0.02 (pH 5.8) after 8 h, pH decreased and reached final values of 5.90±0.04 (pH 6.8) and 5.51±0.04 (pH 5.8) after 48 h (**Figure 1G**). The degree of alkalinization resulting from the disrupted bicarbonate buffer was small compared to the acidification caused by microbial activity (*i.e.,* organic acid production), and pH remained within the range of the bicarbonate buffering system (pH 5.1 to 7.1)[32] throughout the cultivation. Therefore, HCl-titration was integrated into the protocol for establishing the final culture conditions. This step requires the presence of a pH meter in the chamber and the utilization of a pre-determined titration curve (**Supplementary file 1, Step 6; Figure 1F**) to rapidly set the initial pH to the desired values.

Effect of basal medium and gas phase compositions on the metabolism and community structure of fecal cultures

Having selected a high-throughput vessel and established a modular procedure for preparing the medium and final culture conditions, we next aimed to evaluate whether the basal medium composition could influence fecal cultures' metabolic activity and taxonomic composition. We compared bYCFA and mM2, with bYCFA considered a potential substitute for mM2 to reduce batch-to-batch variability, as rumen fluid contained in mM2 (30% v/v) varies in composition and is difficult to procure. Rumen fluid was replaced with a mixture of hemin, vitamins and short-chain fatty acids (SCFA) in bYCFA. Using both the plate-based and tube-based cultivation techniques, we cultured two fecal microbiota for 48 h, 37°C, in bYCFA and mM2 supplemented with two different C-sources (resistant

dextrin, soluble starch, or H_2O as control). Our results showed that the concentration of produced metabolites after 48 h strongly correlated between bYCFA and mM2, both in terms of total metabolites (R=1, **Figure S2A**) and individual metabolites (*i.e.,* SCFA and intermediate metabolites; R=0.97, **Figure S2B**). Similarly, the richness of fecal cultures determined via observed ASVs (alpha diversity) strongly correlated between the two basal media (R=0.94, **Figure S2C**). The community composition, assessed via Bray-Curtis distances (beta diversity), showed closely overlapping confidence ellipses when visualized on a Principal Coordinate Analysis (PCoA; **Figure S2D**). Altogether, these data indicate that the effect mediated by the basal medium was minimal. Therefore, to ensure reproducibility and minimize batch-to-batch variability, we recommend the use of bYCFA as the standard basal medium for cultivating fecal microbiota (**Supplementary file 1**).

Next, we aimed to quantify to what degree the gas phase conditions of both cultivation techniques could impact fecal cultures, *i.e.,* an open system with gas-permeable 96-deepwell plates exposed to 10% CO_2, 5% H_2 and 85% N_2 and a close system with gas-tight tubes containing 100% CO_2. In bYCFA (**Figure 2A;** mM2 in **Figure S3**), the concentration of produced total metabolites (R=0.99, **Figure 2B**) and individual metabolites (R=0.97, **Figure 2C**) strongly correlated between the two cultivation techniques. The only noticeable difference was a lower accumulation of the intermediate metabolite formate in plates as compared to tubes when starch was added (**Figure 2C**). The richness of fecal cultures (observed ASVs) was similar or higher in plates than in tubes (**Figure 2D**). Permutational multivariate analysis (Adonis2) of Bay-Curtis distances revealed that the cultivation technique had the smallest effect size (F) on community composition (F=3.9, p≤0.01), as highlighted by the closely overlapping confidence ellipses for the two cultivation techniques (**Figure 2E**). Variations were mostly explained by the donor microbiota (F=70.1, p<0.001), followed by the supplemented fiber (F=24.1, p≤0.001). Finally, specific taxa enriched with each fiber also correlated between plates and tubes (R=0.85, **Figure 2F**).

Though the fecal cultures profiles were similar when grown using the two cultivation protocols, subtle differences in the produced metabolites were observed (*i.e.,* reduction of formate concentration both with bYCFA (**Figure 2C**) and mM2 (**Figure S3C**)). To identify whether these differences might be attributed to slight variations in the metabolic outputs of individual strains exposed to the chamber's atmosphere, we cultured a panel of pure gut bacteria in bYCFA using both cultivation techniques (**Table S2, Figure S4A**). The selected bacteria covered a range of nutritional preferences (*e.g.,* primary degraders of complex fibers, intermediate metabolites utilizers) and metabolic outputs (*e.g.,* acetate-, propionate- and butyrate-producers). All bacteria were able to grow in the anaerobic chamber using the plate-based technique, though with reduced growth (OD_{600}) after 48 h compared to the tubes (**Figure S4B**).

Figure 2: Impact of the cultivation technique (96-deepwell plates vs. gas-tight tubes) on the metabolism and composition of fecal cultures (donor ABX and BCY) cultivated in bYCFA for 48 h, 37°C, in the presence of resistant dextrin (3 g/L), soluble starch (3 g/L) or H2O as control (technical triplicates).
A) Experimental setup for comparing the plate- and tube-based techniques. All analyses were performed after 48 h cultivation. B) Correlation of the total metabolite production. Total metabolites were calculated by summing the medium-corrected concentrations of organic acids (SCFA and intermediates). C) Correlation of individual metabolites. Points represent means of blank corrected concentrations and bars represent standard deviations. D) Correlation of observed ASVs. Points represent means and bars represent standard deviations. E) PCoA plot of Bray-Curtis distances with 95% confidence ellipses (calculated separately for each donor) for the two cultivation techniques. F) Median clr-differences of taxa (>1% abundance) that are significantly different between each fiber (dextrin, starch) and H2O (p ≤0.05). Labels indicate highly enriched taxa (clr-differences >5).

The cultivation technique appeared to influence the carbohydrate metabolism of the tested strains, with all cultures exhibiting higher levels of end-metabolites (the SCFA acetate, propionate and butyrate) in plates as compared to tubes (**Figure S4C**). The two butyrate-producing species *Agathobacter rectalis* and *Eubacterium ramulus* yielded significantly lower levels of formate in plate compared to tubes (9.5±2 vs. 12.5±1 mM, p≤0.05, and 1.1±0.03 vs. 3.5±0.1 mM,

p≤0.001, respectively; **Figure S4C**). When grown in the chamber, the propionate-producing species *Phocaeicola dorei* showed higher conversion of succinate (1.8±0.03 vs. 3.0±0.1 mM, p≤0.001) to propionate (5.7±0.3 vs. 3.9±0.1 mM, p≤0.05; **Figure S4C**) as compared to tubes. Collectively, these data highlight that the difference of atmosphere between the two cultivation techniques can impact the metabolic output during microbial cultivation. These differences were more pronounced in pure cultures than in complex communities. Specific procedures for testing fecal microbiota (**Supplementary file 1, Step 5.1**) or pure cultures (**Supplementary file 1, Step 5.2**) have been included in the high-throughput protocol.

Formulating a growth medium to maintain the microbial composition of healthy adult feces in cultures

Next, we sought to formulate an optimal mix of C-sources for complementing bYCFA basal medium to maximize the compositional similarity between fecal cultures and feces from healthy human adults. By maintaining the main characteristics of the donor's community in culture, we aimed to enhance the relevance of *in vitro* testing for evaluating the effect mediated by diet-, drug- and host-related factors. To this end, we cultivated fecal samples from eight healthy adults and examined whether the bacterial composition of feces could be maintained using complex mixtures of carbohydrates (3 g/L total; 3C and 6C) and mucin (+Muc), as compared to glucose and H_2O (C-depleted control; **Figure 3A**). The composition of 3C was designed to mirror the C-sources present in the M2GSC medium (33% starch, 33% cellobiose and 33% glucose)[30], while 6C contained the carbohydrates found in the Macfarlane medium (15% starch, 15% pectin, 15% xylan, 8% arabinogalactan, 8% guar gum and 38% inulin)[33], a nutrient-rich medium that closely mimics the gut chyme of healthy adults, and is commonly used in continuous intestinal *in vitro* models.

In terms of bacterial richness, the number of observed ASVs in feces (median of 178 among donors) was best maintained in culture conditions containing both mucin and complex C-sources, with the 6C+Muc condition exhibiting the highest number of ASVs (148), followed by 3C+Muc (134), 3C (129), 6C (124), and glucose (113.5) (**Figure 3B**). Though 135.5 ASVs were detected in the C-depleted control, a significant portion likely originated from resting cells from the fecal inoculum, as limited growth (OD_{600}) was observed after 48 h (**Figure S5A**). When comparing the taxonomic composition of the fecal cultures to the original feces using Bray-Curtis distances (ranging from 0: all ASVs shared, to 1: no ASVs shared), the conditions containing complex C-sources demonstrated higher similarity to the feces (6C+Muc (0.55), 3C+Muc (0.55), 6C (0.56), and 3C (0.57)) than those supplemented with glucose (0.69) or H_2O (0.83; **Figure 3C**). In line, shared ASVs (relative abundance >0.1%) between feces and cultures were highest in the presence of 6C+Muc (80±5%), followed 3C+Muc (74±4%), 6C (73±5%), 3C (70±4%), and lowest with glucose (53±7%) or in the absence of C-source (55±7%; **Figure 3D**). At the family level, the main fecal taxa were retained in cultures containing the complex C-

sources (**Figure 3D**), except for donor JHB, with a loss of *Prevotellaceae* abundance observed after 48 h cultivation (0.6±0.3% among all complex C-source cultures vs. 32.2% in feces; **Figure 3D**). Notably, the addition of complex C-sources limited the abundance of *Enterobacteriaceae*, whose members have been associated with dysbiosis and were often reported to bloom during *in vitro* cultivation in the chamber[25,27]. The difference of *Enterobacteriaceae* levels between cultures and feces was smallest in the presence of 6C (clr-difference of 2.1, *i.e.*, 4.3-fold ($2^{2.1}$) increase) and 6C+Muc (4.5-fold increase), and the largest in the presence of glucose (13.3-fold increase) and H_2O (32.1-fold increase) (**Figure S5B**). Overall, among all cultures containing 6C+Muc, the maximum relative abundance of *Enterobacteriaceae* detected after 48 h was only 7.2% (Donor KPJ; **Figure 3D**).

When evaluating the metabolic outputs of fecal cultures, a significant increase in metabolite production was observed in the presence of C-sources (**Figure S5C**). Yet, substantial amounts of formate accumulated at the end of the cultivation (**Figure S5D**), indicating incomplete fermentation of the supplied C-source. We thus assessed whether the addition of N-source could shift fermentation profiles towards end-metabolites[34]. We cultivated fecal samples from seven healthy adults in bYCFA with 6C+Muc and supplemented an additional 7.2 g/L Amicase and 1 g/L yeast extract (**Figure 3E**). With this additional N-source, a reduction of formate levels after 48 h (from 14.4±3.8% to 1.3±2.6%) was observed, and a representative SCFA ratio[35] was produced, with 61.5±5.1% acetate, 18.9±1.9% butyrate, and 16.4±5.4% propionate among all donors (**Figure S6A**). However, higher N-source concentration was not superior in maintaining the composition of the original feces. Though a minor effect was observed in terms of richness (extra N-source: 123 ASVs; control: 119 ASVs; **Figure S6B**), N-source addition augmented Bray-Curtis distances between cultures and feces (extra N-source: 0.64; control: 0.56; **Figure 3E**). Similarly, *Enterobacteriaceae* relative abundance was promoted with the supplementation of additional N-source (**Figure 3E**). Therefore, we recommend complementing bYCFA with 6C+Muc without additional N-source to retain the compositional characteristics from healthy adult feces during cultivation (**Supplementary file 1**).

Lastly, we assessed whether communities cultured in bYCFA with 6C+Muc could be maintained through repeated passages to provide a setup for studying, *e.g.,* post-perturbation recovery patterns[25]. Two fecal cultures were re-inoculated (1% v/v) into fresh medium over three successive passages (P1, P2, P3) at two different intervals (24 or 48 h; **Figure 3F**). The successive passages increased the dissimilarity between the cultured communities and the original fecal samples (Bray-Curtis distances, **Figure 3F**). Despite a similar overall divergence of community structure between 24 and 48 h intervals, there was a notable difference in taxonomic composition and metabolic activity. Shorter incubation of 24 h compared to 48 h resulted in an increase in the clr-abundance of *Enterobacteriaceae* between initial cultures and subsequent passages (**Figure S7A**) and mitigated the decline of *Prevotellaceae* in both donors (**Figure S7A**).

Figure 3: Formulation of a growth medium capable of maintaining the microbial composition of healthy adult feces in cultures during high-throughput cultivation (in bYCFA, 48 h, 37°C).

A) Experimental setup for determining the optimal C-source mix to maintain the composition of the original feces in cultures. All analyses were performed after 48 h cultivation. B) Number of observed ASVs in feces and cultures with different C-sources. C) Bray-Curtis distances between the feces and the respective culture. D) Visual representation of the community composition (top) and the shared ASVs (with relative abundance >0.1%) between feces and cultures (bottom). E) Evaluation of the impact of additional N-source supplementation (Amicase and yeast extract) on Bray-Curtis distances and on the difference in Enterobacteriaceae clr-abundance as compared to the feces. F) Bray-Curtis distances and composition of cultures re-inoculated (1%) into fresh bYCFA with 6C+Muc (24/48 h, 37°C, 3 passages, donors n=2).

All analyses were performed on pooled samples from three independent replicates for each donor microbiota. A total of 11 distinct donors were used, *i.e.*, AXZ, BFW, GCB, JHB, KPJ, SUT, TLM, WPL, KHG, OIP and ZTR.

Conversely, higher levels of intermediate metabolites (*i.e.,* formate, succinate) were detected in 24 h compared to 48 h passages (**Figure S7B**). Although successive re-inoculation of fecal cultures induced shifts in community structure after each passage, this approach still shows promise for studying recovery dynamics in a high-throughput manner.

Treating fecal cultures with dietary fibers or drugs results in physiologically relevant microbial responses

Finally, we aimed to validate our *in vitro* high-throughput cultivation protocol by testing whether the supplementation of specific dietary fibers (resistant dextrin and soluble starch; **Figure 4A**) or drugs (omeprazole, ciprofloxacin and 5-fluorouracil (5-FU); **Figure 4D**) could induce microbial changes that are consistent with previous *in vivo* findings.

We chose resistant dextrin (3 g/L) and soluble starch (3 g/L) as dietary fibers due to their well-known ability to modulate the gut microbiota and mediate propiogenic[36] and butyrogenic[37–39] effects, respectively. Using the feces from eight healthy donors, we performed enrichment experiments in bYCFA containing each fiber as the sole C-source. Among all donors, we found significantly higher levels of propionate after 48 h in the presence of dextrin (10.9±2.9 mM) as compared to starch (6.0±2.4 mM; p≤0.05) or the control without C-sources (3.9±1.2 mM; p≤0.001). Conversely, starch supplementation led to significantly higher levels of butyrate (10.1±2.1 mM) compared to dextrin (4.5±1.1 mM; p≤0.001) or the control (2.4±1.5 mM; p≤0.01) (**Figure 4B**). At the taxonomic level, resistant dextrin specifically increased the abundance of *Fusicatenibacter* (p≤0.001) and *Lachnospiraceae* ND3007 group (p≤0.01) (**Figure 4C**) among all donors, consistent with previous *in vitro* and *in vivo* studies[40] (+ref NicheMap). Soluble starch supplementation enriched *Faecalibacterium* (p≤0.001) and *Agathobacter* (p≤0.001; formerly *Eubacterium*; **Figure 4C**) as shown in previous human trials[17,38,39,41]. *Blautia* was enriched by both fibers (p≤0.001; **Figure 4C**).

Many commonly used medications have been shown to disrupt gut microbial communities[6,42]. Reciprocally, microbial activities can alter drug efficacy and toxicity[43–45]. To test the direct interaction between the gut microbiota and drugs, we exposed the feces from seven healthy donors (in bYCFA with 6C+Muc) to a single dose of 11 µM omeprazole (proton-pump inhibitor), 50 µM ciprofloxacin (antibiotic) or 50 µM 5-FU (cytostatic), at their expected intestinal concentrations[25,46,47]. Compared to the control cultures, treatment with ciprofloxacin and 5-FU negatively impacted the microbial communities' diversity (Figure 4E) and structure (Figure 4F). Ciprofloxacin led to a relative increase in *Butyricicoccaceae* (p≤0.01) and *Lachnospiraceae* (p≤0.001) (**Figure 4G**). In line, the reduction in community diversity and the persistence of *Lachnospiraceae* was previously shown in mice[25] and ciprofloxacin-treated patients[48]. Treatment with 5-FU inhibited *Sutterellaceae* (p≤0.01) and *Erysipelatoclostridiaceae* (p≤0.01) and promoted the relative abundance of *Akkermansiaceae* (p≤0.05) (**Figure 4G**). The reduction in diversity and the drastic bloom of *Akkermansiaceae* in

response to 5-FU have been demonstrated in several mice studies[49–51]. On the other hand, we observed no major effect of omeprazole on fecal cultures at the tested concentration (**Figures 4E and 4F**). This common proton pump inhibitor was shown to decrease the richness, enrich the abundance of oral bacteria (e.g*., Streptococcaceae*) in the gut, and increase the risk of enteric infections *in vivo*[52]. The effect of omeprazole on the gut microbiota was proposed to be primarily mediated by the increase in gastric pH, which disrupts the barrier between the upper and lower gastrointestinal tract, rather than through a direct effect of the molecule itself[52]. Our data align with this hypothesis, with no direct impact observed on the microbiota. It is worth noting that omeprazole was dissolved in DMSO at a final concentration of 0.2% DMSO in fecal culture. At this concentration, DSMO did not affect the community diversity and structure (**Figure S8**), indicating that it can serve as an optimal solvent for testing compounds with low solubility in an aqueous phase.

DISCUSSION

Developing a well-controlled protocol for cultivating stool-derived microbial communities can significantly help advance our understanding of the causal interactions between gut microbes, diet, drugs, and host physiology[53]. However, the throughput of current anaerobic cultivation techniques is insufficient to capture the complex nature of the gut microbiota, with heterogenous compositions among individuals[16–19], variations in activities at the strain level[20–23], and the influence of environmental factors on ecological and physiological dynamics[6,10]. A few pioneering studies have reported the development of anaerobic high-throughput systems for cultivating stool-derived gut microbiota to generate personalized culture collections of gut microbes[54,55], map the ability of the human gut microbiota to metabolize small molecule drugs[27] or evaluate the response of fecal communities to specific stimuli[25,26]. These studies have highlighted the promises of high-throughput *in vitro* modeling by demonstrating a strong correlation between *in vitro* and *in vivo* gut microbiota responses[25,26]. Here, we addressed practical, technical, and biological gaps in fecal cultivation that we consider essential to create a widely applicable, reproducible and accessible protocol to advance the field of gut microbiota research.

First, we developed a protocol that relies solely on commonly available laboratory materials and equipment, making it easily implementable in various research settings. The core requirement is an anaerobic chamber equipped with a pH meter, an incubator, and multichannel pipettes. We have selected 96-deepwell plates as the cultivation vessel as they offer various advantages; they are reusable and provide a high throughput capacity (60 wells available for testing) with a sufficient volume of samples for analyses (2 mL). Further, the small surface area in each well minimizes the evaporation and gas transfer rate, thus reducing the effect of potential O_2 contamination[56]. The main limitation of 96-deepwell plates is their inability to support online monitoring of absorbance or fluorescence using plate readers.

Figure 4: Treatment with dietary fibers (donor n=8) or drugs (donor n=7) elicits physiologically relevant changes in the fecal cultures' metabolism and community composition (bYCFA, 48 h, 37°C).

A) Experimental setup to test the fiber-specific response of fecal cultures. All analyses were performed after 48 h cultivation. B) SCFA production by fecal cultures when bYCFA was supplemented with resistant dextrin (3 g/L), soluble starch (3 g/L) or no C-source (H2O as control). C) Median clr-differences of taxa significantly differed between each fiber (dextrin, starch) and H2O. Genera significantly different (p≤0.05 and clr-difference >3) are labeled and located above the red line. D) Experimental setup to test the effect of drugs (*i.e.*, omeprazole, ciprofloxacin and 5-FU) as compared to the control (H2O) in bYCFA supplemented with 6C+Muc. All analyses were performed after 48 h cultivation. E) The number of observed ASVs. F) Bray-Curtis distances between controls and the drug-treated cultures. G) Clr-differences of the 15 most abundant families. Significantly altered families, as compared to the control, are marked by stars.

All the analyses were performed on pooled samples from technical triplicates for each donor microbiota. Statistically significant results are marked by stars, with * indicating p≤0.05, ** p≤0.01 and *** p≤0.001.

Second, we comprehensively characterized confounding factors associated with anaerobic chamber cultivation. We showed that the gas phase of the chamber i) might contain residual O_2, ii) alters the pH of the medium, and iii) modulates the C-metabolism of intestinal microbes. To prevent oxygen contamination, we recommend properly preparing the chamber by monitoring H_2 levels and rejuvenating the palladium catalysts. Our protocol also includes a step of boiling and flushing (CO_2 or N_2) of the basal medium prior to sterilization (**Supplementary file 1**), similar to what is done with the Hungate technique[14]. While it is theoretically possible to skip these steps and generate anaerobiosis by pre-reducing a sterile aerobic media inside the chamber for a few days prior to the experiments[25,27], we observed a limited bloom of facultative aerobes, such as *Enterobacteriaceae*, with our protocol compared to other high-throughput cultivation studies[25,27]. We also highlighted the negative impact of a weak reducing atmosphere on the growth of intestinal microbes (**Figure 1B**). Therefore, we strongly recommend boiling and flushing the basal medium prior to sterilizing to ensure strict anaerobiosis. In regard to pH, we highlight that the equilibrium between gaseous and dissolved CO_2 and the concentration of $NaHCO_3$ in the solution influenced the pH in the chamber. Thus, the bicarbonate buffer is more challenging to control in an open system where CO_2 can escape. With a buffering range between pH 5.1 to 7.1[32], it can maintain the pH in the range of the human adult colon, ranging from 5.7 (proximal) to 6.7 (distal)[57], with values as low as 5.0 in some healthy individuals[58]. In addition, bYCFA basal medium contains a phosphate buffer maintaining pH between 6.4 and 7.4. Concerning the impact of the gas phase on microbial metabolism, we highlighted a few biological differences between the plate-based anaerobic chamber and the classic tube-based Hungate technique. The typically lower CO_2 concentration in the chamber (~10%) compared to tubes (~100%) mediated an increase of succinate to propionate conversion in pure cultures of *P. dorei* (**Figure S4C**)[59,60]. Moreover, the plate-based system allows for releasing microbially produced gases, such as H_2, a common growth-limiting factor in tube cultures when it builds up[61], promoting the accumulation of intermediate metabolites[62]. In agreement, tube cultures exhibited a higher accumulation of intermediate products (*i.e.,* formate) than plate cultivation (**Figures 2C, S3C and S4C**). Overall, the typical gas composition in the anaerobic chamber (*i.e.,* 5% H_2, 10% CO_2, and 85% N_2) is close to the physiological conditions reported *in vivo* for H_2 and CO_2, with concentrations of 2.9±0.7% and 9.9±1.6%, respectively[63].

Third, we designed a reproducible and modular medium. The basal medium bYCFA was formulated to be heat-stable (sterilization via autoclave) and contain as few undefined ingredients as possible to increase batch-to-batch reproducibility. This basal medium can be flexibly complemented with C-sources or any other molecules to test, and the pH can be easily adjusted, allowing for a wide range of culture conditions that can be covered with minimal medium preparation effort. Therefore, the effects of a combination of factors can be easily tested and incorporated during the experimental design (**Supplementary file 1**). Here, we showcased two examples of experimental design. We complemented bYCFA with defined fibers to identify taxa associated with functional niches (**Figures 4A, 4B and 4C**). This approach corroborated results from a previous tube-based enrichment study (+Ref Nichemap) while allowing the evaluation of more donors and conditions simultaneously. Then, we maintained fecal culture communities resembling the original feces by complementing bYCFA with an optimized mix of C-sources (6C+Muc; **Figure 3B, 3C and 3D**), and we tested the microbial responses to antibiotic and non-antibiotic drugs (**Figures 4D, 4E, 4F and 4G**). This approach can be extended to study the effects of other xenobiotics (*e.g.,* pollutants), dietary compounds (*e.g.*, N-sources, vitamins), or host-derived factors (*e.g.*, pH, redox potential, oxidative stress, bile salts, antimicrobial peptides). Note that molecules insoluble in water can be dissolved in DSMO, with no effect on the microbial community at a concentration of 0.2% (**Figure S8**) and potentially up to 1%[64].

Finally, we integrated all technical procedures and practical solutions into a simple yet comprehensive step-by-step protocol (**Supplementary file 1**). This protocol provides a high level of detail and encompasses key features like flexibility (*e.g.*, testing of various molecules and conditions alone or in combination; complex communities or pure cultures), reproducibility (*e.g.*, reduced media complexity; preparation of stable stock solutions), and simplicity (*e.g.*, minimized hands-on time; easy pH adjustment; reduce evaporation rate). In conclusion, our high-throughput cultivation protocol is a valuable tool for fundamental and applied gut microbiota research. It enables the rapid and reproducible characterization of diet-, host-, or drug-specific effects, microbe-microbe interactions[65], strain-specific[20,23] and donor-specific[19] responses and activities. The protocol can also be applied for the cost-efficient screening of drug metabolism by the gut microbiota[45,46], pre-selection of treatments or dosage prior to costly animal or human trials[66], development of personalized microbiota-targeted nutritional interventions[67], identification of responders and non-responders to drugs and prebiotics[19] and for the discovery and investigation of novel live biotherapeutics[68].

Acknowledgments

This work was supported by a grant from the Swiss Innovation Agency (51128.1 IP-LS) in collaboration with PharmaBiome AG. This work was also supported by an ETH Research Grant ETH-38 20-1. We thank Alfonso Die (ETH Zürich) and Leandra Knecht (Pharmabiome AG) for their assistance with HPLC-RI analyses. We thank Florentin Constancias for his support with statistical analysis. We thank Anna Greppi for her valuable feedback during the preparation of the manuscript. Data were generated in collaboration with the Genetic Diversity Centre (GDC), ETH Zurich.

Author contributions

BP, CL, TW, and GL conceptualized the project and provided financial support. JZ, BP and GL planned the experiments. JZ,

SP, DM, CM performed wet-lab experiments and analyses. JZ, PB, GL curated the data. JZ and PB performed data analysis. JZ, BP, GL, CL interpreted the data. JZ, SP and BP wrote the step-by-step protocol. JZ and BP wrote the manuscript. All authors critically reviewed the manuscript. All authors contributed to the article and approved the submitted version.

Declaration of interest

Authors CM, PB, TW and GL are or were employees of PharmaBiome AG. TW and CL are founders of PharmaBiome AG. The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Benoit Pugin (benoit.pugin@hest.ethz.ch).

METHODS

Experimental model and subject details

Human fecal samples

14 healthy donors (7 females, 7 males, ages ranging between 26 and 36) provided one or more fresh fecal samples. Donors were excluded if they had taken any antibiotics in the past six months. Sample collection was processed in an anonymized and non-interventional manner. Donor identity was ensured by using randomly generated three-letter codes. The ethics committee of ETH Zurich exempted this study from review.

Bacterial strains

A panel of ten human intestinal strains, representing a panel of fiber and intermediate metabolite utilizers, was acquired from the German Collection of Microorganisms and Cell Culture GmbH (DSMZ, Braunschweig, Germany) and the strain collection of PharmaBiome AG (Zürich, Switzerland; **Table S2**). Strains were stored at −80°C in anaerobic glycerol stocks (25%, v/v) and reactivated by inoculating (1.25%, v/v) Hungate tubes containing bYCFA (pH 6.8) supplemented with the respective carbon sources (final 3 g/L for fibers, 30 mM for glucose and intermediate metabolites; **Table S2**). Pre-cultures were incubated for 24 h at 37°C before experiments.

Method details

Growth media and anaerobic procedures

To cultivate fecal microbiota and pure cultures, we modified the yeast casitone fatty acid medium (YCFA)[31] and the M2GSC[30] medium. Modifications included a tenfold reduction of N-sources, as previously reported[24]. The resulting basal minimal media, referred to as basal YCFA (bYCFA) and modified M2GSC (mM2), consisted of (L^{-1}): 1 g Amicase (Sigma-Aldrich Chemie GmbH, Buchs, Switzerland), 1.25 g yeast extract (Procelys; Lesaffre, Marcq-en-Barœul, France), 0.5 g meat extract (Sigma-Aldrich), 150 mL mineral solution I

(0.52 g K_2HPO_4), 150 mL mineral solution II (0.52 g KH_2PO_4, 0.9 g NaCl, 0.9 g $(NH_4)_2SO_4$, 90 mg $MgSO_4$, 90 mg $CaCl_2$), 1 g L-cysteine HCl, and 4 g $NaHCO_3$. The medium mM2 further contained 300 mL rumen fluid (centrifuged and stored at -20°C after collection; Vetsuisse Faculty University of Zurich, Zürich, Switzerland). The rumen fluid was replaced in bYCFA with defined compounds, *i.e.,* 0.2 mL hemin stock solution (10 μg hemin from bovine, Sigma-Aldrich), 0.1 mL vitamin solution (10 μg biotin, 10 μg cobalamin, 30 μg 4-aminobenzoic acid, 50 μg folic acid and 150 μg pyridoxamine, Sigma-Aldrich) and 6.2 mL of volatile fatty acids solution (2.11 mL acetic acid (Sigma-Aldrich), 0.74 mL propionic acid (Sigma-Aldrich), 0.12 mL valeric acid (VWR International AG, Dietikon, Switzerland), 0.12 mL isovaleric acid (Sigma-Aldrich), and 0.12 mL isobutyric acid (Sigma-Aldrich) in 3.1 mL 5 M NaOH). Since the basal medium was complemented with (heat-stable/-sensitive) supplements after sterilization (**Figure 1A**), it was thus prepared in a more concentrated form (**Table S3**). All compounds except L-cysteine.HCl and $NaHCO_3$ were dissolved in 75% of the final volume (**Table S3**). The basal medium was then adjusted to pH 7, boiled (10 min), and the remaining compounds were added after cooling to approximately 60°C while constantly flushing with CO_2 (or N_2). After 10 min, the medium was transferred into DURAN® Pressure flasks (Huberlab AG, Aesch, Switzerland) or Hungate tubes (Milian, Vernier, Switzerland) and autoclaved.

To complement the basal medium, a range of heat-stable supplement solutions were prepared. For fecal microbiota cultivation, we supplemented the media with different heat-stable C-source solutions (concentration of 3 g/L in the final media), *i.e.,* glucose (Sigma-Aldrich), resistant dextrin (NUTRIOSE® FB06; Roquette, Lestrem, France), starch (soluble starch from potato; Sigma-Aldrich), a mix of six sugars (referred to as 6C; 15% starch, 15% pectin (pectin from citrus peel; Sigma-Aldrich), 15% xylan (xylan from oat spelt; Angene, London, United Kingdom), 8% arabinogalactan (from larch wood; Sigma-Aldrich), 8% guar (Sigma-Aldrich) and 38% inulin (Orafti GR; Beneo, Mannheim, Germany) or a mix of three sugars (referred to as 3C; 33% starch, 33% cellobiose (Sigma-Aldrich) and 33% glucose). The 6C and 3C mixes were also tested with mucin from porcine stomach type II (0.3 g/L final concentration; Sigma-Aldrich). Heat-stable carbohydrates were also supplemented to basal bYCFA for cultivating pure bacterial strains, as described in **Table S2**. Additional heat-stable supplements included N-source solutions, which resulted in final concentration of 8.2 g/L Amicase and 2.25 g/L yeast extract. Heat-stable compounds were dissolved in dH_2O, adjusted to pH 7, boiled for 10 min, cooled down under N_2 and autoclaved soon after.

In addition, heat-sensitive compounds were tested, including antibiotic and non-antibiotic drugs at final concentrations of 50 μM ciprofloxacin (Sigma-Aldrich), 11 μM omeprazole (Sigma-Aldrich) and 50 μM 5-fluorouracil (Sigma-Aldrich). Concentrated drugs were dissolved in anaerobic dH_2O (except for omeprazole which was dissolved in dimethyl sulfoxide (DMSO); 0.2% final concentration) in the anaerobic chamber, adjusted to pH 7 and filter sterilized.

The day before the experiment, the autoclaved basal medium was equilibrated with the gas phase of the anaerobic chamber (10% CO_2, 5% H_2, and 85% N_2, Coy Laboratory Products Inc., Grass Lake, MI, United States) under constant stirring, without the lid but covered with Breathe-Easier Sealing film (Diversified Biotech, Dedham, MA, United States). On the day of the experiment, pH was adjusted to 6.8 (if not indicated otherwise) by adding HCl 3 M based on a pre-determined titration curve (**Figure 1E**), and was mixed with the corresponding heat-stable/sensitive supplements (**Table S3**).

A complete step-by-step description of the preparation of the basal bYCFA medium and heat-stable/sensitive supplement solutions for the high-throughput cultivation in 96-deepwell plates (Milian, Vernier, Switzerland) is presented in **Supplementary file 1**. For gas-tight tube-based culture using the Hungate technique, anaerobiosis was maintained by applying a strict anaerobic technique as described before[69]. Syringes and needles were pre-flushed with sterile CO_2 before being used to inoculate carbon source solutions or bacteria.

Fecal sample inoculation, cultivation and sampling procedure

Healthy adult donors provided a fresh fecal sample in a plastic container containing an Oxoid™ AnaeroGen™ pouch (Thermo Fisher Diagnostics AG, Pratteln, Switzerland) to generate anaerobiosis during transport to the laboratory. Within 3 hours, the samples were transferred into the anaerobic chamber. Approximately 1 g of fecal matter was transferred using a plastic spoon into a 50 mL Falcon tube and homogenized with 9 mL of anaerobic phosphate buffer (pH 7.0). The resulting 10^{-1} dilution was then further diluted to 10^{-4} and was inoculated (1%, v/v) into a plate or tube containing previously pH-adjusted basal medium (pH 6.8) complemented with respective supplements (**Supplementary file 1, Step 5.2**). Unless otherwise stated, working volumes of 2 mL and 8 mL of medium were used in 96-deepwell plates and gas-tight Hungate tubes, respectively. After incubation for 48 h at 37°C, cultures were transferred out of the chamber and centrifuged (4'700 rpm, 20 min and 4°C). Cell pellets and supernatants were recovered and stored separately (-80°C) until further processing (**Supplementary file 1, Step 8**).

Anaerobic chamber preparation and its impact on bacterial growth

We assessed the effects of anaerobic chamber preparation on the growth of gut microbes in pure cultures. Growth kinetics of *Bacteroides thetaiotaomicron* DSM 2079 and *Faecalibacterium prausnitzii* DSM 17677 were monitored in an anaerobic chamber that was handled "strictly" or "loosely". A "strictly" handled chamber (**Supplementary file 1**, **Step 2**) consisted of rejuvenated palladium catalysts and a gas atmosphere replaced with fresh gas mix (10% CO_2, 5% H_2, and 85% N_2), resulting in 0-20 ppm O_2 and >2.5% H_2 (monitored with an Anaerobic Monitor CAM-12, Coy Laboratory Products Inc.).

To demonstrate the effect of a "loosely" handled chamber, cultures (adjusted to OD_{600} ~0.2) were inoculated (1% v/v) into 200 µL bYCFA (30 mM glucose, pH 6.5) in a 96-well plate (SPL Life Sciences Co. Ltd., Gyeonggi-do, South Korea). Then, non-rejuvenated palladium catalysts were introduced into the chamber, and gas mix was replaced with N_2 until the H_2 level was less than 1.5%. Growth kinetics of *F. prausnitzii* and *B. thetaiotaomicron* were then monitored for 36 h in a Tecan Infinite M200 PRO plate reader (Tecan Group Ltd., Männedorf, Switzerland) and compared to growth kinetics in a "strictly" handled anaerobic chamber. Each condition was tested in three biological replicates. The maximum specific growth rate (μ_{max}) was calculated according to first-order growth kinetics.

pH dynamics and bicarbonate buffering system in the anaerobic chamber

We characterized the relationship between pH, $NaHCO_3$ concentration, and CO_2 levels (0% or 100% in gas-tight tubes, compared to 10% in the anaerobic chamber) by measuring the pH of the basal medium containing different $NaHCO_3$ concentrations (0.4, 1, 2, 4 g/L). When required, pH dynamics were monitored over time. Measurements were performed using a pH microelectrode (diameter 6 mm; Metrohm AG, Herisau, Switzerland) and a 913 pH meter (Metrohm AG). pH was measured in a 96-deepwell plate filled with 1.5 mL basal medium. A titration with HCl 3 M was performed and served as a calibration to support the pH adjustment of bYCFA in the chamber (**Supplementary file 1**, **Step 6**).

Comparison of the cultivation techniques (96-deepwell plates vs. gas-tight tubes) using fecal cultures and pure bacterial cultures

Two stool-derived gut microbiota (donors ABX and BCY) and eight intestinal strains (**Table S2**) were assessed for their growth and metabolic capabilities using plate- and tube-based techniques. Plates and tubes were filled with bYCFA or mM2, complemented with the corresponding C-sources (**Table S3**), and inoculated (1% v/v) with fecal dilutions or pre-cultures of pure strains. After the incubation (48 h, 37°C), 200 µL of cultures was transferred into a 96-well plate for OD_{600} measurement. Supernatants and the cell pellets were recovered (4'700 rpm, 20 min and 4°C) for organic acids quantification and 16S rRNA sequencing analysis, respectively.

Specifically for this experiment, the following 16S rRNA sequencing pipeline was used: DNA extraction of fecal cultures was performed using the Maxwell® RSC instrument and the Maxwell® RSC PureFood GMO and Authentication Kit (Promega AG, Dübendorf, Switzerland). Library preparation, amplification, and paired-end (2x 300) sequencing of the hyper-variable regions V3/V4 of the genomic 16S rRNA gene with the primers (341F: 5"- CCTACGGGNBGCASCAG-3"; 806bR: 5"- GGACTACNVGGGTWTCTAAT-3") was performed by StarSeq GmbH (Mainz, Germany) using Illumina MiSeq. Cell pellets originating from later experiments were sequenced as described below.

Optimal medium composition to culture healthy adult feces and response of individuals' gut microbiota to fibers and drugs during high-throughput cultivation

We inoculated fecal dilutions from eight (C-source evaluation) or seven (N-source evaluation; fibers and drugs testing) adult donors in 96-deepwell plate containing bYCFA complemented with the corresponding supplements listed in **Table S3**. Given small functional and compositional variability across independent replicates in previous experiments, technical replicates were pooled by transferring 0.5 mL from each culture into a fresh plate (**Supplementary file 1**, **Step 8)**. Two pools were prepared, one of which was stored immediately at -80°C, while the other was processed as described above.

Organic acids quantification

Short-chain fatty acids (SCFA; acetate, propionate and butyrate) and intermediate metabolites (formate, lactate and succinate) were quantified by high-performance liquid chromatography equipped with a refractive index detector (HPLC-RI). Briefly, around 300 µL supernatant was passed through a 0.2 µm nylon membrane filter (AcroPrep™ Filter plate; Pall Corporation, New York, United States; or syringe filters; Merck & Cie KmG, Schaffhausen, Switzerland). Analysis with a sample volume of 40 µL was carried out using a Chromaster HPLC-System (Hitachi, Tokyo, Japan) equipped with a SecurityGuard Cartridge Carbo-H (4×3.0 mm; Phenomenex Inc., Torrance, CA, United States) and a Rezex ROA-Organic Acid H+ (300×7.8 mm; Phenomenex Inc.) column. Elution was performed at 80°C under isocratic conditions (10 mM H_2SO_4, flow rate 0.6 mL/min), and analytes were quantified using a refractive index detector Chromaster 5450 (Hitachi). Data were processed using the Chromaster software (Hitachi).

All listed compound concentrations were corrected with blank values of the basal medium to estimate the net production of SCFA and intermediates. To quantify the total metabolites, the concentration of the medium-corrected concentrations of measured metabolites were summed. Metabolite fractions were calculated by dividing the medium-corrected concentration of the respective metabolite by total metabolites.

DNA extraction and 16S rRNA sequencing analysis

Cell pellets were pre-processed using the Maxwell® HT 96 gDNA Blood Isolation Kit (Promega AG, Dübendorf, Switzerland) combined with a FastPrep homogenizer (FastPrep-24TM; MP Biomedicals, Illkirch-Graffenstaden, France), and followed by DNA purification using KingFisher Flex instrument (Thermo Fisher Scientific, Waltham, MA, United States), according to the manufacturers' instructions. Amplification and barcoding of the V4 hypervariable region of the 16S rRNA gene was performed using a PCR with the barcoded primer 515F (5′-GTGCCAGCMGCCGCGGTAA-3′) and 806R (5′-GGACTACHVGGGTWTCTAAT-3′)[70,71]. The pooled library was then sequenced using the amplifying primers (515F and 806R) and the Illumina MiSeq platform[72] at the Genetic Diversity Centre (ETH Zurich, Switzerland).

Sequencing reads were processed using the metabaRpipe R package (version 0.9)[73]. In brief, reads were inferred with the DADA2 R package (version 1.14.1)[74] with read filtering set to c(260, 250) and maximal error rates set to c(3,4). Then, bimeras were removed and taxonomy was assigned to ASVs using the SILVA database (v138.1)[75,76]. The communities were then analyzed using the phyloseq package (v1.40.0) of the R software (v4.2.0)[77]. Alpha and beta-diversity analyses were done on rarefied samples (with a minimal rarefaction depth of 1556 reads). Data meeting the criteria of homoscedasticity were evaluated with a permutational multivariate analysis using the adonis2 function of the vegan package (2.6-4). Differential abundance analysis was performed on non-rarified data using ALDEx2[78]. Using the aldex.clr function from the mia package (1.5.17), the data was centered log(2)-ratio (clr) transformed using 1000 Dirichlet Monte-Carlo (DMC) instances. Significantly altered taxa were identified based on the p-value of Welch's t-test.

Data analysis and visualization

Data were analyzed using R software (v4.2.0) and visualized using the ggplot package (3.3.6). Correlation lines were plotted using ggpmisc (0.5.1). Data were evaluated for normality using the Shapiro-Wilk Normality Test and when p>0.05, the comparison of means was performed using a t-test. Otherwise, a Wilcoxon test using the rstatix package (0.7.0) was used.

REFERENCES

1. Arumugam, M., Raes, J., Pelletier, E., Paslier, D. Le, Yamada, T., Mende, D.R., Fernandes, G.R., Tap, J., Bruls, T., Batto, J.M., et al. (2011). Enterotypes of the human gut microbiome. Nature *473*, 174–180. 10.1038/NATURE09944.

2. Valdes, A.M., Walter, J., Segal, E., and Spector, T.D. (2018). Role of the gut microbiota in nutrition and health. BMJ *361*, 36–44. 10.1136/BMJ.K2179.

3. Poeker, S.A., Geirnaert, A., Berchtold, L., Greppi, A., Krych, L., Steinert, R.E., De Wouters, T., and Lacroix, C. (2018). Understanding the prebiotic potential of different dietary fibers using an in vitro continuous adult fermentation model (PolyFermS). Sci. Rep. *8*, 1–12. 10.1038/s41598-018-22438-y.

4. Wastyk, H.C., Fragiadakis, G.K., Perelman, D., Dahan, D., Merrill, B.D., Yu, F.B., Topf, M., Gonzalez, C.G., Van Treuren, W., Han, S., et al. (2021). Gut Microbiota-Targeted Diets Modulate Human Immune Status. Cell *184*, 4137. 10.1016/J.CELL.2021.06.019.

5. Djekic, D., Shi, L., Brolin, H., Carlsson, F., Särnqvist, C., Savolainen, O., Cao, Y., Bäckhed, F., Tremaroli, V., Landberg, R., et al. (2020). Effects of a Vegetarian Diet on Cardiometabolic Risk Factors, Gut Microbiota, and Plasma Metabolome in Subjects With Ischemic Heart Disease: A Randomized, Crossover Study. J. Am. Hear. Assoc. Cardiovasc. Cerebrovasc. Dis. *9*, e016518. 10.1161/JAHA.120.016518.

6. Jackson, M.A., Verdi, S., Maxan, M.E., Shin, C.M., Zierer, J., Bowyer, R.C.E., Martin, T., Williams, F.M.K., Menni, C., Bell, J.T., et al. (2018). Gut microbiota associations with common diseases and prescription medications in a population-based cohort. Nat. Commun. *9*, 2655. 10.1038/s41467-018-05184-7.

7. Palleja, A., Mikkelsen, K.H., Forslund, S.K., Kashani, A., Allin, K.H., Nielsen, T., Hansen, T.H., Liang, S., Feng, Q., Zhang, C., et al. (2018). Recovery of gut microbiota of healthy adults following antibiotic exposure. Nat. Microbiol. *3*, 1255–1265. 10.1038/s41564-018-0257-9.

8. Klünemann, M., Andrejev, S., Blasche, S., Mateus, A., Phapale, P., Devendran, S., Vappiani, J., Simon, B., Scott, T.A., Kafkia, E., et al. (2021). Bioaccumulation of therapeutic drugs by human gut bacteria. Nature *597*, 533–538. 10.1038/s41586-021-03891-8.

9. Hashimoto, T., Perlot, T., Rehman, A., Trichereau, J., Ishiguro, H., Paolino, M., Sigl, V., Hanada, T., Hanada, R., Lipinski, S., et al. (2012). ACE2 links amino acid malnutrition to microbial ecology and intestinal inflammation. Nature *487*, 477–481. 10.1038/nature11228.

10. Tropini, C., Moss, E.L., Merrill, B.D., Ng, K.M., Higginbottom, S.K., Casavant, E.P., Gonzalez, C.G., Fremin, B., Bouley, D.M., Elias, J.E., et al. (2018). Transient Osmotic Perturbation Causes Long-Term Alteration to the Gut Microbiota. Cell *173*, 1742–1754. 10.1016/J.CELL.2018.05.008.

11. Roager, H.M., Hansen, L.B.S., Bahl, M.I., Frandsen, H.L., Carvalho, V., Gøbel, R.J., Dalgaard, M.D., Plichta, D.R., Sparholt, M.H., Vestergaard, H., et al. (2016). Colonic transit time is related to bacterial metabolism and mucosal turnover in the gut. Nat. Microbiol. *1*, 1–9. 10.1038/nmicrobiol.2016.93.

12. Murga-Garrido, S.M., Hong, Q., Cross, T.W.L., Hutchison, E.R., Han, J., Thomas, S.P., Vivas, E.I., Denu, J., Ceschin, D.G., Tang, Z.Z., et al. (2021). Gut microbiome variation modulates the effects of dietary fiber on host metabolism. Microbiome *9*, 117. 10.1186/S40168-021-01061-6.

13. Hungate, R.E. (1969). Chapter IV A Roll Tube Method for Cultivation of Strict Anaerobes. Methods Microbiol. *3*, 117–132. 10.1016/S0580-9517(08)70503-8.

14. Hungate, R.E. (1944). Studies on Cellulose Fermentation: I. The Culture and Physiology of an Anaerobic Cellulose-digesting Bacterium. J. Bacteriol. *48*, 499–513. 10.1128/JB.48.5.499-513.1944.

15. Kundra, P., Geirnaert, A., Pugin, B., Morales Martinez, P., Lacroix, C., and Greppi, A. (2022). Healthy adult gut microbiota sustains its own vitamin B12 requirement in an in vitro batch fermentation model. Front. Nutr. *9*, 2940. 10.3389/FNUT.2022.1070155.

16. Sandberg, J., Kovatcheva-Datchary, P., Björck, I., Bäckhed, F., and Nilsson, A. (2019). Abundance of gut Prevotella at baseline and metabolic response to barley prebiotics. Eur. J. Nutr. *58*, 2365–2376. 10.1007/S00394-018-1788-9.

17. Walker, A.W., Ince, J., Duncan, S.H., Webster, L.M., Holtrop, G., Ze, X., Brown, D., Stares, M.D., Scott, P., Bergerat, A., et al. (2011). Dominant and diet-responsive groups of bacteria within the human colonic microbiota. ISME J. *5*, 220–230. 10.1038/ismej.2010.118.

18. Cantu-Jungles, T.M., Bulut, N., Chambry, E., Ruthes, A., Iacomini, M., Keshavarzian, A., Johnson, T.A., and Hamaker, B.R. (2021). Dietary fiber hierarchical specificity: The missing link for predictable and strong shifts in gut bacterial communities. MBio *12*, e01028-21. 10.1128/MBIO.01028-21.

19. Hjorth, M.F., Christensen, L., Larsen, T.M., Roager, H.M., Krych, L., Kot, W., Nielsen, D.S., Ritz, C., and Astrup, A. (2020). Pretreatment Prevotella-to-Bacteroides ratio and salivary amylase gene copy number as prognostic markers for dietary weight loss. Am. J. Clin. Nutr. *111*, 1079–1086. doi.org/10.1093/ajcn/nqaa007.

20. Hall, A.B., Yassour, M., Sauk, J., Garner, A., Jiang, X., Arthur, T., Lagoudas, G.K., Vatanen, T., Fornelos, N., Wilson, R., et al. (2017). A novel Ruminococcus gnavus clade enriched in inflammatory bowel disease patients. Genome Med. *9*, 1–12. 10.1186/S13073-017-0490-5.

21. Karcher, N., Pasolli, E., Asnicar, F., Huang, K.D., Tett, A., Manara, S., Armanini, F., Bain, D., Duncan, S.H., Louis, P., et al. (2020). Analysis of 1321 Eubacterium rectale genomes from metagenomes uncovers complex phylogeographic population structure and subspecies functional adaptations. Genome Biol. *21*, 138. 10.1186/S13059-020-02042-Y.

22. Sorbara, M.T., Littmann, E.R., Fontana, E., Moody, T.U., Kohout, C.E., Gjonbalaj, M., Eaton, V., Seok, R., Leiner, I.M., and Pamer, E.G. (2020). Functional and Genomic Variation between Human-Derived Isolates of Lachnospiraceae Reveals Inter- and Intra-Species Diversity. Cell Host Microbe *28*, 134–146. 10.1016/J.CHOM.2020.05.005.

23. Otaru, N., Ye, K., Mujezinovic, D., Berchtold, L., Constancias, F., Cornejo, F.A., Krzystek, A., de Wouters, T., Braegger, C., Lacroix, C., et al. (2021). GABA Production by Human Intestinal Bacteroides spp.: Prevalence, Regulation, and Role in Acid Stress Tolerance. Front. Microbiol. *12*, 656895. 10.3389/FMICB.2021.656895.

24. Anthamatten, L., Bieberstein, P.R. von, Thabuis, C., Menzi, C., Reichlin, M., Meola, M., Rodriguez, B., Cordero, O.X., Lacroix, C., Wouters, T. de, et al. (2023). Mapping gut bacteria into functional niches reveals the ecological structure of human gut microbiomes. bioRxiv, 2023.07.04.547750. 10.1101/2023.07.04.547750.

25. Aranda-Díaz, A., Ng, K.M., Thomsen, T., Real-Ramírez, I., Dahan, D., Dittmar, S., Gonzalez, C.G., Chavez, T., Vasquez, K.S., Nguyen, T.H., et al. (2022). Establishment and characterization of stable, diverse, fecal-derived in vitro microbial communities that model the intestinal microbiota. Cell Host Microbe *30*, 260–272. 10.1016/J.CHOM.2021.12.008.

26. Li, L., Abou-Samra, E., Ning, Z., Zhang, X., Mayne, J., Wang, J., Cheng, K., Walker, K., Stintzi, A., and Figeys, D. (2019). An in vitro model maintaining taxon-specific functional activities of the gut microbiome. Nat. Commun. *10*, 4146. 10.1038/s41467-019-12087-8.

27. Javdan, B., Lopez, J.G., Chankhamjon, P., Lee, Y.C.J., Hull, R., Wu, Q., Wang, X., Chatterjee, S., and Donia, M.S. (2020). Personalized mapping of drug metabolism by the human gut microbiome. Cell *181*, 1661. 10.1016/J.CELL.2020.05.001.

28. Li, L., Ning, Z., Zhang, X., Mayne, J., Cheng, K., Stintzi, A., and Figeys, D. (2020). RapidAIM: A culture-And metaproteomics-based Rapid Assay of Individual Microbiome responses to drugs. Microbiome *8*, 1–16. 10.1186/S40168-020-00806-Z.

29. Douglas S. Auld, P.D., Peter A. Coassin, B.S., Nathan P. Coussens, P.D., Hensley, P., Klumpp-Thomas, C., Michael, S., G. Sitta Sittampalam, P.D., O. Joseph Trask, B.S., Bridget K. Wagner, P.D., Jeffrey R. Weidner, P.D., et al. (2020). Microplate Selection and Recommended Practices in High-throughput Screening and Quantitative Biology (Eli Lilly & Company and the National Center for Advancing Translational Sciences).

30. Miyazaki, K., Martin, J.C., Marinsek-Logar, R., and Flint, H.J. (1997). Degradation and Utilization of Xylans by the Rumen AnaerobePrevotella bryantii(formerlyP. ruminicolasubsp.brevis) B14. Anaerobe *3*, 373–381. 10.1006/ANAE.1997.0125.

31. Duncan, S.H., Hold, G.L., Harmsen, H.J.M., Stewart, C.S., and Flint, H.J. (2002). Growth requirements and fermentation products of Fusobacterium prausnitzii, and a proposal to reclassify it as Faecalibacterium prausnitzii gen. nov., comb. nov. Int. J. Syst. Evol. Microbiol. *52*, 2141–2146. 10.1099/00207713-52-6-2141.

32. Kohn, R.A., and Dunlap, T.F. (1998). Calculation of the buffering capacity of bicarbonate in the rumen and in vitro. J. Anim. Sci. *76*, 1702–1709. 10.2527/1998.7661702X.

33. Macfarlane, G.T., Macfarlane, S., and Gibson, G.R. (1998). Validation of a three-stage compound continuous culture system for investigating the effect of retention time on the ecology and metabolism of bacteria in the human colon. Microb. Ecol. *35*, 180–187. 10.1007/S002489900072.

34. Walker, A.W., Duncan, S.H., Carol McWilliam Leitch, E., Child, M.W., and Flint, H.J. (2005). pH and peptide supply can radically alter bacterial populations and short-chain fatty acid ratios within microbial communities from the human colon. Appl. Environ. Microbiol. *71*, 3692–3700. 10.1128/AEM.71.7.3692-3700.2005.

35. Cummings, J.H., Pomare, E.W., Branch, H.W.J., Naylor, E., and Macfarlane, G.T. (1987). Short chain fatty acids in human large intestine, portal, hepatic and venous blood. Gut *28*, 1221–1227. 10.1136/gut.28.10.1221.

36. Guerin-Deremaux, L., Ringard, F., Desailly, F., and Wils, D. (2010). Effects of a soluble dietary fibre NUTRIOSE® on colonic fermentation and excretion rates in rats. Nutr. Res. Pract. *4*, 470–476. 10.4162/NRP.2010.4.6.470.

37. Louis, P., Scott, K.P., Duncan, S.H., and Flint, H.J. (2007). Understanding the effects of diet on bacterial metabolism in the large intestine. J. Appl. Microbiol. *102*, 1197–1208. 10.1111/J.1365-2672.2007.03322.X.

38. Venkataraman, A., Sieber, J.R., Schmidt, A.W., Waldron, C., Theis, K.R., and Schmidt, T.M. (2016). Variable responses of human microbiomes to dietary supplementation with resistant starch. Microbiome *4*, 1–9. 10.1186/S40168-016-0178-X.

39. Baxter, N.T., Schmidt, A.W., Venkataraman, A., Kim, K.S., Waldron, C., and Schmidt, T.M. (2019). Dynamics of human gut microbiota and short-chain fatty acids in response to dietary interventions with three fermentable fibers. MBio *10*, e02566-18. 10.1128/MBIO.02566-18.

40. Barber, C., Sabater, C., Ávila-Gálvez, M.Á., Vallejo, F., Bendezu, R.A., Guérin-Deremaux, L., Guarner, F., Espín, J.C., Margolles, A., and Azpiroz, F. (2022). Effect of Resistant Dextrin on Intestinal Gas Homeostasis and Microbiota. Nutrients *14*, 4611. 10.3390/NU14214611.

41. Vital, M., Howe, A., Bergeron, N., Krauss, R.M., Jansson, J.K., and Tiedje, J.M. (2018). Metagenomic Insights into the Degradation of Resistant Starch by Human Gut Microbiota. Appl. Environ. Microbiol. *84*, e.01562-18. 10.1128/AEM.01562-18.

42. Vich Vila, A., Collij, V., Sanna, S., Sinha, T., Imhann, F., Bourgonje, A.R., Mujagic, Z., Jonkers, D.M.A.E., Masclee, A.A.M., Fu, J., et al. (2020). Impact of commonly used drugs on the composition and metabolic function of the gut microbiota. Nat. Commun. *11*, 362. 10.1038/s41467-019-14177-z.

43. van Kessel, S.P., Frye, A.K., El-Gendy, A.O., Castejon, M., Keshavarzian, A., van Dijk, G., and El Aidy, S. (2019). Gut bacterial tyrosine decarboxylases restrict levels of levodopa in the treatment of Parkinson's disease. Nat. Commun. *10*, 310. 10.1038/s41467-019-08294-y.

44. Yuan, L., Zhang, S., Li, H., Yang, F., Mushtaq, N., Ullah, S., Shi, Y., An, C., and Xu, J. (2018). The influence of gut microbiota dysbiosis to the efficacy of 5-Fluorouracil treatment on colorectal cancer. Biomed. Pharmacother. *108*, 184–193. 10.1016/J.BIOPHA.2018.08.165.

45. Zimmermann, M., Zimmermann-Kogadeeva, M., Wegmann, R., and Goodman, A.L. (2019). Mapping human microbiome drug metabolism by gut bacteria and their genes. Nature *570*, 462–467. 10.1038/s41586-019-1291-3.

46. Maier, L., Pruteanu, M., Kuhn, M., Zeller, G., Telzerow, A., Anderson, E.E., Brochado, A.R., Fernandez, K.C., Dose, H., Mori, H., et al. (2018). Extensive impact of non-antibiotic drugs on human gut bacteria. Nature *555*, 623–628. 10.1038/nature25979.

47. LaCourse, K.D., Zepeda-Rivera, M., Kempchinsky, A.G., Baryiames, A., Minot, S.S., Johnston, C.D., and Bullman, S. (2022). The cancer chemotherapeutic 5-fluorouracil is a potent Fusobacterium nucleatum inhibitor and its activity is modified by intratumoral microbiota. Cell Rep. *41*. 10.1016/J.CELREP.2022.111625.

48. Stewardson, A.J., Gaïa, N., François, P., Malhotra-Kumar, S., Delémont, C., Martinez de Tejada, B., Schrenzel, J., Harbarth, S., Lazarevic, V., Vervoort, J., et al. (2015). Collateral damage from oral ciprofloxacin versus nitrofurantoin in outpatients with urinary tract infections: a culture-free analysis of gut microbiota. Clin. Microbiol. Infect. *21*, 344.e1. 10.1016/J.CMI.2014.11.016.

49. Li, H.L., Lu, L., Wang, X.S., Qin, L.Y., Wang, P., Qiu, S.P., Wu, H., Huang, F., Zhang, B.B., Shi, H.L., et al. (2017). Alteration of gut microbiota and inflammatory cytokine/chemokine profiles in 5-fluorouracil induced intestinal mucositis. Front. Cell. Infect. Microbiol. *7*, 455. 10.3389/FCIMB.2017.00455.

50. Hamouda, N., Sano, T., Oikawa, Y., Ozaki, T., Shimakawa, M., Matsumoto, K., Amagase, K., Higuchi, K., and Kato, S. (2017). Apoptosis, Dysbiosis and Expression of Inflammatory Cytokines are Sequential Events in the Development of 5-Fluorouracil-Induced Intestinal Mucositis in Mice. Basic Clin. Pharmacol. Toxicol. *121*, 159–168. 10.1111/BCPT.12793.

51. Carvalho, R., Vaz, A., Pereira, F.L., Dorella, F., Aguiar, E., Chatel, J.M., Bermudez, L., Langella, P., Fernandes, G., Figueiredo, H., et al. (2018). Gut microbiome modulation during treatment of mucositis with the dairy bacterium Lactococcus lactis and recombinant strain secreting human antimicrobial PAP. Sci. Rep. *8*, 15072. 10.1038/s41598-018-33469-w.

52. Jackson, M.A., Goodrich, J.K., Maxan, M.E., Freedberg, D.E., Abrams, J.A., Poole, A.C., Sutter, J.L., Welter, D., Ley, R.E., Bell, J.T., et al. (2016). Proton pump inhibitors alter the composition of the gut microbiota. Gut *65*, 749–756. 10.1136/GUTJNL-2015-310861.

53. Clavel, T., Horz, H.P., Segata, N., and Vehreschild, M. (2022). Next steps after 15 stimulating years of human gut microbiome research. Microb. Biotechnol. *15*, 164–175. 10.1111/1751-7915.13970.

54. Hitch, T.C.A., Afrizal, A., Riedel, T., Kioukis, A., Haller, D., Lagkouvardos, I., Overmann, J., and Clavel, T. (2021). Recent advances in culture-based gut microbiome research. Int. J. Med. Microbiol. *311*, 151485. 10.1016/J.IJMM.2021.151485.

55. Goodman, A.L., Kallstrom, G., Faith, J.J., Reyes, A., Moore, A., Dantas, G., and Gordon, J.I. (2011). Extensive personal human gut microbiota culture collections characterized and manipulated in gnotobiotic mice. Proc. Natl. Acad. Sci. U. S. A. *108*, 6252–6257. 10.1073/PNAS.1102938108.

56. Berg, M., Undisz, K., Thiericke, R., Zimmermann, P., Moore, T., and Posten, C. (2001). Evaluation of Liquid Handling Conditions in Microplates. SLAS Discov. *6*, 47–56. 10.1177/108705710100600107.

57. Fallingborg, J. (1999). Intraluminal pH of the human gastrointestinal tract. Dan. Med. Bull. *46*, 183–196.

58. Koziolek, M., Grimm, M., Becker, D., Iordanov, V., Zou, H., Shimizu, J., Wanke, C., Garbacz, G., and Weitschies, W. (2015). Investigation of pH and Temperature Profiles in the GI Tract of Fasted Human Subjects Using the Intellicap(®) System. J. Pharm. Sci. *104*, 2855–2863. 10.1002/JPS.24274.

59. Fischbach, M.A., and Sonnenburg, J.L. (2011). Eating For Two: How Metabolism Establishes Interspecies Interactions in the Gut. Cell Host Microbe *10*, 336. 10.1016/J.CHOM.2011.10.002.

60. Macy, J.M., Ljungdahl, L.G., and Gottschalk, G. (1978). Pathway of Succinate and Propionate Formation in Bacteroides fragilis. J. Bacteriol. *134*,

84–91. 10.1128/JB.134.1.84-91.1978.

61. Smith, N.W., Shorten, P.R., Altermann, E.H., Roy, N.C., and McNabb, W.C. (2019). Hydrogen cross-feeders of the human gastrointestinal tract. Gut Microbes *10*, 270–288. 10.1080/19490976.2018.1546522.

62. Van Lingen, H.J., Plugge, C.M., Fadel, J.G., Kebreab, E., Bannink, A., and Dijkstra, J. (2016). Thermodynamic Driving Force of Hydrogen on Rumen Microbial Metabolism: A Theoretical Investigation. PLoS One *11*. 10.1371/JOURNAL.PONE.0161362.

63. Modesto, A., Cameron, N.R., Varghese, C., Peters, N., Stokes, B., Phillips, A., Bissett, I., and O'Grady, G. (2022). Meta-Analysis of the Composition of Human Intestinal Gases. Dig. Dis. Sci. *67*, 3842–3859. 10.1007/S10620-021-07254-1.

64. Chauveau, A., Treyer, A., Geirnaert, A., Bircher, L., Babst, A., Abegg, V.F., Simões-Wüst, A.P., Lacroix, C., Potterat, O., and Hamburger, M. (2023). Intestinal permeability and gut microbiota interactions of pharmacologically active compounds in valerian and St. John's wort. Biomed. Pharmacother. *162*, 114652. 10.1016/J.BIOPHA.2023.114652.

65. Shetty, S.A., Kuipers, B., Atashgahi, S., Aalvink, S., Smidt, H., and de Vos, W.M. (2022). Inter-species Metabolic Interactions in an In-vitro Minimal Human Gut Microbiome of Core Bacteria. npj Biofilms Microbiomes *8*, 21. 10.1038/s41522-022-00275-2.

66. Liu, Y., Gibson, G.R., and Walton, G.E. (2016). An In Vitro Approach to Study Effects of Prebiotics and Probiotics on the Faecal Microbiota and Selected Immune Parameters Relevant to the Elderly. PLoS One *11*, e0162604. 10.1371/JOURNAL.PONE.0162604.

67. Gibbons, S.M., Gurry, T., Lampe, J.W., Chakrabarti, A., Dam, V., Everard, A., Goas, A., Gross, G., Kleerebezem, M., Lane, J., et al. (2022). Perspective: Leveraging the Gut Microbiota to Predict Personalized Responses to Dietary, Prebiotic, and Probiotic Interventions. Adv. Nutr. *13*, 1450–1461. 10.1093/ADVANCES/NMAC075.

68. Relizani, K., Le Corf, K., Kropp, C., Martin-Rosique, R., Kissi, D., Déjean, G., Bruno, L., Martinez, C., Rawadi, G., Elustondo, F., et al. (2022). Selection of a novel strain of Christensenella minuta as a future biotherapy for Crohn's disease. Sci. Rep. *12*. 10.1038/S41598-022-10015-3.

69. Bircher, L., Geirnaert, A., Hammes, F., Lacroix, C., and Schwab, C. (2018). Effect of cryopreservation and lyophilization on viability and growth of strict anaerobic human gut microbes. Microb. Biotechnol. *11*, 721–

733. 10.1111/1751-7915.13265.

70. Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S.M., Betley, J., Fraser, L., Bauer, M., et al. (2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J. *6*, 1621–1624. 10.1038/ismej.2012.8.

71. Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A., Turnbaugh, P.J., Fierer, N., and Knight, R. (2011). Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. Proc. Natl. Acad. Sci. U. S. A. *108*, 4516–4522. 10.1073/PNAS.1000080107.

72. Walters, W., Hyde, E.R., Berg-Lyons, D., Ackermann, G., Humphrey, G., Parada, A., Gilbert, J.A., Jansson, J.K., Caporaso, J.G., Fuhrman, J.A., et al. (2015). Improved Bacterial 16S rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed Spacer Marker Gene Primers for Microbial Community Surveys. mSystems *1*. 10.1128/MSYSTEMS.00009-15.

73. Constancias, F., and Mahé, F. (2022). fconstancias/metabaRpipe: v0.9 (v0.9). Zenodo. 10.5281/zenodo.6423397.

74. Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., and Holmes, S.P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. Nat. Methods *13*, 581–583. 10.1038/nmeth.3869.

75. Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F.O. (2013). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res. *41*, D590–D596. 10.1093/NAR/GKS1219.

76. Yilmaz, P., Parfrey, L.W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J., Ludwig, W., and Glöckner, F.O. (2014). The SILVA and "All-species Living Tree Project (LTP)" taxonomic frameworks. Nucleic Acids Res. *42*, D643–D648. 10.1093/NAR/GKT1209.

77. McMurdie, P.J., and Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. PLoS One *8*, e61217. 10.1371/JOURNAL.PONE.0061217.

78. Fernandes, A.D., Reid, J.N.S., Macklaim, J.M., McMurrough, T.A., Edgell, D.R., and Gloor, G.B. (2014). Unifying the analysis of high-throughput sequencing datasets: Characterizing RNA-seq, 16S rRNA gene sequencing and selective growth experiments by compositional data analysis. Microbiome *2*, 1–13. 10.1186/2049-2618-2-15.

SUPPLEMENTARY FIGURES

Figure S1: pH dynamics of CO_2- and N_2-flushed mM2 in 96-deepwell plate during equilibration in the anaerobic chamber (uninoculated).

A

Total metabolite production

$R = 1.00$

B

Individual metabolite production

$R = 0.97$

C

Alpha diversity

$R = 0.94$

D

Bray-Curtis dissimilarity

Figure S2: Impact of basal media (bYCFA vs. mM2) on the metabolism and community structure of fecal-derived gut microbiota (donor ABX and BCY) cultivated 96-deepwell plates and gas-tight tubes for 48 h, 37°C, in the presence of resistant dextrin (3 g/L), soluble starch (3 g/L) or H_2O as control (technical triplicates).
All analyses were performed after 48 h cultivation. **A)** Correlation of the total metabolite production. Total metabolites were calculated by summing the medium-corrected concentrations of organic acids (SCFA and intermediates). **B)** Correlation of individual metabolites. Points represent means of blank corrected concentrations and bars represent standard deviations. **C)** Correlation of observed ASVs. Points represent means and bars represent standard deviations. **D)** PCoA plot of Bray-Curtis distances with 95% confidence ellipses (calculated separately for each donor) for the two basal medium.

Figure S3: Impact of the cultivation technique (96-deepwell plates vs. gas-tight tubes) on the metabolism and composition of fecal cultures (donor ABX and BCY) cultivated in mM2 for 48 h, 37°C, in the presence of resistant dextrin (3 g/L), soluble starch (3 g/L) or H₂O as control (technical triplicates).

A) Experimental setup for comparing the plate- and tube-based techniques. All analyses were performed after 48 h cultivation. **B)** Correlation of the total metabolite production. Total metabolites were calculated by summing the medium-corrected concentrations of organic acids (SCFA and intermediates). **C)** Correlation of individual metabolites. Points represent means of blank corrected concentrations and bars represent standard deviations. **D)** Correlation of observed ASVs. Points represent means and bars represent standard deviations. **E)** PCoA plot of Bray-Curtis distances with 95% confidence ellipses (calculated separately for each donor) for the two cultivation techniques. **F)** Median clr-differences of taxa (>1% abundance) that are significantly different between each fiber (dextrin, starch) and H₂O (p ≤0.05). Labels indicate highly enriched taxa (clr-differences >5).

Figure S4: Impact of the cultivation technique (96-deepwell plates vs. gas-tight tubes) on the physiology of pure bacterial strains cultivated in bYCFA for 48 h, 37°C, in the presence of different carbon sources (8 strains; biological triplicates).
A) Experimental setup for comparing the high-throughput protocol with the Hungate technique using pure cultures. All analyses were performed after 48 h cultivation. **B)** Growth of eight intestinal strains. Bars represent mean endpoint OD_{600} values. **C)** Individual metabolite production. Bars represent mean endpoint concentrations.
For this experiment, bYCFA was supplemented with strain-specific substrates (**Table S2**): *E. ramulus* with arabinogalactan, *P. dorei* with pea, *P. succinatutens* with succinate, *Acutalibacter* sp. with resistant dextrin, *B. adolescentis* with xylan, *A. rectalis* with soluble starch, *Frisingicoccus sp.* with DL-lactate, and *L. eligens* with pectin. Each condition was tested in biological triplicates. Significance was assessed using a t-test with * indicating p≤0.05, ** p≤0.01 and *** p≤ 0.001.

Figure S5: Characteristics of fecal cultures (in bYCFA, 48 h, 37°C, donors n=8) grown in the presence of different C-sources (6C+Muc, 3C+Muc, 6C, 3C and glucose) or in control conditions (C-depleted).

All analyses were performed after 48 h cultivation. **A)** Growth (OD_{600}) of fecal cultures (200 µL sample; path length ~5 mm). **B)** Median difference of clr-abundance of *Enterobacteriaceae* in cultures compared to feces. **C)** Total metabolites were calculated by summing the medium-corrected concentrations of organic acids (SCFA and intermediates) in the supernatant of cultures. **C**) Relative metabolite production was calculated by dividing the concentration of each metabolite by the total amount of metabolites produced.

A

Individual metabolite production

Figure S6: Characteristics of fecal cultures grown in bYCFA with 6C+Muc (48 h, 37°C, donors n=7) supplemented with additional N-source (Amicase and yeast extract) or control (H₂O) conditions
A) Relative metabolite production was calculated by dividing the concentration of each metabolite by the total amount of metabolites produced. **B)** Number of observed ASVs in feces and cultures.
All analyses were performed after 48 h on pooled samples from three technical replicates for each donor microbiota.

B

Alpha diversity

A

Family-level differences

Figure S7: Characteristics of cultures derived from donor microbiota BFW and GCB repeatedly passaged by re-inoculating 1% (3 passages, 24/48 h, 37°C, donors n=2) into bYCFA and 6C+Muc.
A) Difference of clr-abundance of the families *Enterobacteriaceae* and *Prevotellaceae* between the initial culture and the successive passages (P1, P2, P3). **B)** Relative metabolite production was calculated by dividing the concentration of each metabolite by the total amount of metabolites produced.

B

Individual metabolite production

Figure S8: Characteristics of fecal cultures grown in bYCFA with 6C+Muc (48 h, 37°C, donors n=7) and exposed to 0.2 % DMSO compared to control (H$_2$O). A) Number of observed ASVs in cultures treated with 0.2% DMSO or H$_2$O (control). **B)** Bray-Curtis distances between cultures treated with DMSO (0.2%) and H$_2$O visualized in a PCoA plot.
All analyses were performed after 48 h on pooled samples from three technical replicates for each donor microbiota.

SUPPLEMENTARY TABLES

Table S1: Effect of a "loosely" and "strictly" handled chamber on the maximum specific growth rate (μ_{max}) and cell density of *F. prausnitzii* DSM 17677 and *B. thetaiotaomicron* DSM 2079. Normality assumption was verified using the Shapiro-Wilk normality test, and significance was determined using unpaired T-test. Significance level was set to $p < 0.05$.

Strain	Chamber handling	μ_{max} (h^{-1})			max. cell density (OD_{600})		
		Mean	SD	*p*	Mean	SD	*p*
F. prausnitzii DSM 17677	$H_2 > 1.5\%$	0.33	0.01	0.3893	0.27	0.02	0.0023
	$H_2 < 1.5\%$	0.28	0.07		0.06	0.04	
B. thetaiotaomicron DSM 2079	$H_2 > 1.5\%$	0.75	0.02	0.0145	1.03	0.01	0.0755
	$H_2 < 1.5\%$	0.65	0.03		1.00	0.01	

Table S2: Bacterial strains used in this study and corresponding carbon source supplemented for growth.
Respective fibers were supplemented to reach final 3 g/L, and glucose and organic acids were applied at a concentration of 30 mM.

Genus	Species	Strain	Culture collection	Supplemented C-source
Faecalibacterium	*prausnitzii*	DSM 17677	DSMZ	Glucose
Bacteroides	*thetaiotaomicron*	DSM 2079	DSMZ	Glucose
Eubacterium	*ramulus*	PB-SKBSD	PB	Arabinogalactan
Phocaeicola	*dorei*	PB-SXBFK	PB	Pea
Phascolarctobacterium	*succinatutens*	PB-SDVAP	PB	Succinate
Acutalibacter	sp.	PB-SMJER	PB	Dextrin
Bifidobacterium	*adolescentis*	PB-SKRVN	PB	Xylan
Agathobacter	*rectalis*	PB-SWUPP	PB	Soluble starch
Frisingicoccus	sp.	PB-SKUSI	PB	DL-Lactate
Lachnospira	*eligens*	PB-SJATG	PB	Pectin

DSMZ: German Collection of Microorganisms and Cell Culture GmbH; PB: PharmaBiome AG.

Table S3: The modular components of the final medium and their composition.
The basal medium, heat-stable and heat-sensitive supplements were prepared based on the respective concentration factor and mixed according to % of the final volume.

Component type	Composition	Concentration factor	% of the final volume
Comparison of 96-deep well plates and Hungate tubes using fecal microbiota (Figures 2, S3)			
Basal medium	bYCFA, mM2	1.33X	75%
Heat-stable supplement	dextrin, starch, H_2O	4X	25%
Comparison of 96-deep well plates and Hungate tubes using pure cultures (Figure S4)			
Basal medium	bYCFA	1.33X	75%
Heat-stable supplement	arabinogalactan, pea fiber, dextrin, xylan, starch, pectin	4X	25%
Optimization of C-source composition to maintain *in vivo*-like communities (Figures 3, S5, S7)			
Basal medium	bYCFA	1.33X	75%
Heat-stable supplement	glucose, 6C, 6C+Mucin, 3C, 3C+Mucin, H_2O	4X	25%
Optimization of metabolic profile by additional N-source (Figures 3, S6)			
Basal medium	bYCFA	1.33X	75%
Heat-stable supplement	6C+Mucin	8X	12.5%
Heat-stable supplement	N-source, H_2O	8X	12.5%
Treatment of fecal cultures with dietary fibers (Figure 4)			
Basal medium	bYCFA	1.33X	75%
Heat-stable supplement	Dextrin, starch, H_2O	4X	25%
Treatment of fecal cultures with drugs (Figure 4, S8)			
Basal medium	bYCFA	1.33X	75%
Heat-stable supplement	6C+Mucin	8X	12.5%
Heat-sensitive supplement	omeprazole, ciprofloxacin, 5-FU, H_2O, DMSO	8X	12.5%