# Title: Bacterial sepsis triggers stronger transcriptomic immune responses in larger primates

**Authors:** Ryan McMinds[1,2]*, Rays H.Y. Jiang[1,2], Swamy R. Adapa[1,2], Emily Cornelius Ruhs[1,3,4], Rachel A. Munds[1], Jennifer W. Leiding[5,6], Cynthia J. Downs[7]†, Lynn B. Martin[1]†*

**Affiliations:**
[1]Center for Global Health and Infectious Diseases Research (GHIDR), University of South Florida; Tampa, Florida, USA
[2]USF Genomics program, University of South Florida College of Public Health, Tampa, Florida USA
[3]Department of Ecology and Evolution, University of Chicago; Chicago, Illinois, USA
[4]Grainger Bioinformatics Center, Field Museum of Natural History; Chicago, IL.
[5]Division of Allergy and Immunology, Department of Pediatrics, Johns Hopkins University, Baltimore, FL
[6]Institute for Clinical and Translational Research, Johns Hopkins All Children's Hospital, St. Petersburg, FL
[7]Department of Environmental Biology, SUNY College of Environmental Science and Forestry; Syracuse, New York, USA
†Shared senior authorship
*Corresponding authors. Emails: r.mcminds@gmail.com, lbmartin@usf.edu

**Abstract:** Empirical data relating body mass to immune defense against infections remain limited. Although the Metabolic Theory of Ecology predicts that larger organisms would have weaker immune responses, recent studies have suggested that the opposite may be true. These discoveries have led to the Safety Factor hypothesis, which proposes that larger organisms have evolved stronger immune defenses because they carry greater risks of exposure to pathogens and parasites. In this study, we simulated sepsis by exposing blood from nine primate species to a bacterial lipopolysaccharide (LPS), measured the relative expression of immune and other genes using RNAseq, and fit phylogenetic models to determine how gene expression was related to body mass. In contrast to non-immune-annotated genes, we discovered hypermetric scaling in the LPS-induced expression of innate immune genes, such that large primates had a disproportionately greater increase in gene expression of immune genes compared to small primates. Hypermetric immune gene expression appears to support the Safety Factor hypothesis, though this pattern may represent a balanced evolutionary mechanism to compensate for lower per-transcript immunological effectiveness. This study contributes to the growing body of immune allometry research, highlighting its importance in understanding the complex interplay between body size and immunity over evolutionary timescales.

**Author Summary:** Understanding the relationship between an animal's size and its ability to defend against disease can inform predictions about evolutionary tradeoffs and susceptibility to infection. Two major theories – the Metabolic Theory of Ecology (MTE) and the Safety Factor hypothesis – offer opposing views on how body size influences it immune defenses. In this study, we compared the immune gene expression of nine primate species to a simulated bacterial infection. We found that larger species mounted a stronger transcriptional immune response, consistent with either the Safety Factor hypothesis or an evolutionary pressure to compensate for and balance the effects expected in the context of the MTE.

**Introduction:**

The body size of organisms profoundly affects most of their life processes [1]. Partly, these effects are due to physical forces affecting how materials are moved into and out of cells, but evolutionary factors, too, can influence how size affects life [2]. For instance, reaching a large size inherently requires time, so selection should act strongly on developmental rates and age-specific breeding traits, which could lead to innovative molecular or physiological mechanisms that circumvent some physical constraints. Relationships between size and biological traits are often modeled as trait $\sim \alpha * \text{mass}^\beta$, which can be linearized via log-transformation of both body mass and the response variable (i.e., $\log(\text{trait}) \sim \log(\alpha) + \beta * \log(\text{mass})$). Relationships between body mass and response variables are then represented by the slope coefficients, $\beta$, which can be estimated with linear regression. Traits such as blood volume and length of the digestive tract scale proportionally to body size (i.e., isometrically), with $\beta$ approximately equal to 1 [3,4]. Other traits, such as hematocrit, also scale isometrically [5,6], but because they are measures of concentration that are inherently scaled by body mass, they have coefficients approximately equal to 0. Allometric relationships, where $\beta$ deviates from the proportional expectation, are also quite common. A trait with $\beta$ greater than isometry is called hypermetric whereas one with $\beta$ less than isometry is hypometric. A prime example of hypoallometry is organismal metabolic rate, which scales at ¾ the rate of body mass change. In other words, the total metabolic output of an organism increases as body mass increases, but at less than a one-to-one ratio. Equivalently, mass-normalized metabolic rate has $\beta = -¼$ (i.e., the metabolic rate *per unit mass* **decreases** as body mass itself increases) [7–9].

Biological scaling has been investigated extensively [6,10,11], and metabolic allometries in particular have been so powerful as to lead to the Metabolic Theory of Ecology (MTE) [7,12]. Although contentious [13–16], the MTE has been useful for everything from improving the conservation of wildlife to understanding the structure of human cities [1]. Nevertheless, the MTE so far appears to hold little utility for understanding how defenses, specifically immune system processes, relate to body size. Whereas the MTE predicts hypometric immune cell concentrations ($\beta \approx -¼$) [17], granulocyte concentrations in birds and mammals in fact scale hypermetrically (heterophils and neutrophils scale with $\beta = 0.19$ and $0.11$, respectively) [18,19]. These unexpected results suggest that other processes must come into play to explain how

80  defenses scale with body size, and the Safety Factor hypothesis was recently offered as an eco-
81  evolutionary explanation [20]. This hypothesis proposes that large organisms are exposed to
82  more numerous and more diverse pathogens within and between generations, such that the risks
83  and consequences of infection are much greater compared to smaller species. As a result, large
84  species may be selected to evolve exceptionally strong defenses compared to small species [19].
85
Like most physiological processes involving energy metabolism, gene expression (both the rate
of gene transcription and the abundance of gene transcripts) is predicted by the MTE to scale
hypometrically. A recent study including nematodes, insects, crustaceans, and fish provides
evidence that this is broadly true across the transcriptome [21], but whether immune genes are an
exception to this trend is unknown. Immune gene expression is of particular interest because it
could give more insight into the granulocyte hyperallometry previously described. If immune
gene expression scaling is hypometric, then hypermetric granulocyte concentrations could
simply act as compensation for relatively poor per-cell function. Alternatively, hypermetric
immune gene expression could provide large species with disproportionately stronger defenses.
Other recent studies produced data consistent with immune gene hyperallometry, but due to their
limited phylogenetic sampling, they could not disambiguate allometric effects from effects of
shared ancestry or other confounding variables [22,23].
98
In principle, the prediction for exceptionally strong defenses in large species could be fulfilled by
hypermetric baseline expression (i.e., the immune system maintains *consistently* greater
constitutive expression of defensive genes), by hypermetric response to infection (i.e., immune
gene expression is increased disproportionately *only after detection* of a pathogen), or by both
(Fig. 1A-C). In plants, large, long-lived species tend to have stronger constitutive barriers than
small, short-lived species [24]. Similar patterns have also been suggested across vertebrates
[25,26], and such a strategy could be necessary in the context of the slower metabolism observed
in larger animals. However, greater baseline gene expression would be more costly, and the
balance of these tradeoffs is unclear.
108
To assess the allometry of immune gene expression in these distinct contexts, we compared the
transcriptomic profiles of replicate aliquots of live blood of adult male individuals from nine
primate species after i) a standard dose of lipopolysaccharide (LPS) from *Escherichia coli* to
simulate sepsis (LPS(+)), and ii) a null (LPS(-)) treatment of LPS-free growth media to assess
baseline expression. Unlike a recent similar study [22], which included only four relatively large
catarrhine species, our study included platyrrhine and strepsirrhine species as well, and spanned
almost the full range of body masses of species in the Primates order (Fig. 2). We used Bayesian
phylogenetic generalized linear mixed models (pGLMMs) to link raw transcript abundance data
with fixed effects of the form $\alpha + \beta_1 *$ treatment $+ \beta_2 *$ log(mass) $+ \beta_3 *$ treatment $*$ log(mass),
which links the allometry of baseline (LPS(-)) transcript abundance to $\beta_2$, the allometry of LPS-
induced gene expression ($\Delta$-LPS ) to $\beta_3$, and the allometry of transcript abundance after LPS

(LPS(+)) to the sum of $\beta_2$ and $\beta_3$ (Fig. 1). Models also included an offset to compensate for
transcript compositionality, additional fixed effects to control for within-species differences in
body size, and random effects to control for shared phylogenetic history, repeated measures, and
statistical overdispersion. By standardizing our simulated sepsis methods and comparing
transcriptomic responses in a phylogenetically informed context, we revealed that innate immune
gene expression in response to simulated sepsis scales hypermetrically among Primates.

126
**Fig. 1. Immune genes are characterized by distinct allometries that often result in hypermetric relative**
**transcript abundance during simulated sepsis.** (**A-C**) Example expression patterns demonstrate how phylogenetic
generalized linear mixed model (pGLMM) coefficients correspond to different forms of allometry. (**A**) Hypometric

baseline abundance ($\beta_2 < 0$) is compensated for with hypermetric LPS-induced expression ($\beta_3 > 0$), resulting in
isometric abundance after LPS ($\beta_2 + \beta_3 \sim 0$). (**B**) Isometric baseline abundance ($\beta_2 \sim 0$) plus hypermetric LPS-
induced expression ($\beta_3 > 0$) results in hypermetric abundance after LPS ($\beta_2 + \beta_3 > 0$). (**C**) Hypermetric baseline
abundance ($\beta_2 > 0$) plus isometric LPS-induced expression ($\beta_3 \sim 0$) results in hypermetric abundance after LPS ($\beta_2 +$
$\beta_3 > 0$). (**D-H**) Log abundance (relative to the per-sample size factor and after adding a pseudocount of 0.5) of five
example genes (HGNC gene names *TLR4*, *CTLA4*, *IFNG*, *IL6*, and *IL1B*, respectively) as a function of species'
average body mass in kilograms (plotted on log scale). Symbols correspond to species (▢= *Microcebus murinus*, ○ =
*Callithrix jacchus*, × = *Lemur catta*, ▢+ = *Sapajus apella*, △ = *Daubentonia madagascariensis*, ▢ = *Macaca*
*mulatta*, ▢ = *Papio anubis*, ∗ = *Pongo abelii*, + = *Homo sapiens*). Black points correspond to samples with control
treatment, and red points correspond to samples with LPS treatment. Fitted lines are plotted using coefficients
extracted from the pGLMMs, with dashed lines for 95% credible intervals. pGLMM coefficients are listed, with an
asterisk (*) if the 95% credible interval did not include zero.

**Fig. 2. Body mass and phylogenetic relationships among samples.** (**A**) Primates phylogeny derived from [27]. Tips are annotated with the numbers of individuals sampled for the present study (the pool for *M. murinus* consisted of 8 individuals). (**B**) Body mass (kg) in the context of their phylogenetic relationships (° = individuals in our study, + = species mean from [28].

**Fig. 3. LPS-induced expression of immune genes scales hypermetrically among nine primate species.** (**A**) Log abundance (relative to the per-sample size factor and after adding a pseudocount of 0.5) of all reads belonging to immune-annotated 1:1 orthologous genes as a function of species' average body mass in kilograms (plotted on log scale). Symbols correspond to species (▽ = *Microcebus murinus*, ○ = *Callithrix jacchus*, × = *Lemur catta*, ⊕ = *Sapajus apella*, △ = *Daubentonia madagascariensis*, ◇ = *Macaca mulatta*, ⊠ = *Papio anubis*, ∗ = *Pongo abelii*, + = *Homo sapiens*). Black points correspond to samples with control treatment, and red points to samples with LPS treatment. Fitted lines are plotted using coefficients extracted from the pGLMMs, with dashed lines for 95% credible intervals. pGLMM coefficients are listed in the lower left legend, with the estimate followed by the 95% credible interval. (**B**) Log relative abundance of non-immune-annotated 1:1 orthologous genes.

## Results and Discussion:

***The total immune gene transcriptional response to LPS scales hypermetrically among Primate species.*** To assess the overall allometric effects on immune gene expression, we first pooled all transcript counts that mapped to immune-annotated one-to-one orthologous genes. Baseline

relative abundance (LPS(-)) of this sum of immune-annotated transcripts did not scale
significantly differently from isometry ($\beta_2 \sim -0.03$), but the LPS-induced expression ($\Delta$-LPS) did
scale hypermetrically ($\beta_3 \sim 0.13$) with 95% credibility. Although the relative abundance of
immune gene transcripts after LPS exposure (LPS(+)) did not scale with 95%-credible
hyperallometry ($\beta_2 + \beta_3 \sim 0.10$), the bulk of the posterior was positive (Table 1; Figs. 3, S1, S4-5).

One persistent complication inherent to comparative transcriptomics is how to deal with the
evolution of gene copy number. Because gene duplication is sometimes, but not always,
associated with punctuated (non-Brownian) evolution and neofunctionalization, fewer
assumptions are required if we limit an analysis to only one-to-one orthologous genes, as in our
primary analysis. However, varying copy number is one mechanism by which a gene may vary
its expression levels, so filtering out genes with duplications and losses has the potential to bias
comparative analyses of expression. We thus repeated our analysis of the overall expression of
immune genes using the sum of all reads that mapped to immune orthogroups, including
duplicated and paralogous genes. This analysis returned hypermetric estimates for all of baseline
relative abundance ($\beta_2 \sim 0.05$), LPS-induced expression ($\beta_3 \sim 0.08$), and relative abundance after
LPS ($\beta_2 + \beta_3 \sim 0.13$), although only relative abundance after LPS was 95% credible (Table 1;
Figs. S2-3, S6-7). Together, these analyses suggest that larger primates have disproportionally
stronger responses to LPS, leading to higher relative abundances of immune gene transcripts
during sepsis, than do smaller primates.

| Data subset | Gene expression context | Parameter | Posterior 2.5% CI | Posterior median | Posterior 97.5% CI |
|---|---|---|---|---|---|
| **9-sp, 1:1 orthologs** | LPS(-) | $\beta_2$ | -0.17 | -0.03 | 0.10 |
| **9-sp, 1:1 orthologs** | $\Delta$-LPS | $\beta_3$ | **0.01** | 0.13 | 0.25 |
| **9-sp, 1:1 orthologs** | LPS(+) | $\beta_2 + \beta_3$ | -0.04 | 0.10 | 0.23 |
| **9-sp, orthogroups** | LPS(-) | $\beta_2$ | -0.07 | 0.05 | 0.17 |
| **9-sp, orthogroups** | $\Delta$-LPS | $\beta_3$ | -0.0020 | 0.08 | 0.16 |
| **9-sp, orthogroups** | LPS(+) | $\beta_2 + \beta_3$ | **0.01** | 0.13 | 0.25 |
| **6-sp, 1:1 orthologs** | LPS(-) | $\beta_2$ | -0.05 | 0.02 | 0.09 |
| **6-sp, 1:1 orthologs** | $\Delta$-LPS | $\beta_3$ | **0.04** | 0.13 | 0.23 |
| **6-sp, 1:1 orthologs** | LPS(+) | $\beta_2 + \beta_3$ | **0.08** | 0.15 | 0.22 |
| **6-sp, orthogroups** | LPS(-) | $\beta_2$ | -0.25 | -0.13 | **-0.01** |
| **6-sp, orthogroups** | $\Delta$-LPS | $\beta_3$ | **0.01** | 0.17 | 0.34 |
| **6-sp, orthogroups** | LPS(+) | $\beta_2 + \beta_3$ | -0.07 | 0.04 | 0.16 |

**Table 1. Pooled analyses of immune-annotated genes support hypermetric LPS-induced expression.** Medians
and 95% credible intervals for betas extracted from Bayesian pGLMMs fit to different pools of immune-annotated
genes. 9-sp refers to analyses with all 9 sampled species, and 6-sp refers to analyses of only the 6 species with high-quality reference genomes. Orthogroups include 1:1 orthologous genes, but also genes that appear to have had

196 duplicates or deletions in the sampled species. Baseline relative abundance = LPS(-), LPS-induced expression =
197 Δ-LPS, and relative abundance after LPS = LPS(+).
198
***Individual immune-annotated genes are more likely than non-immune genes to scale***
***hypermetrically.*** Although the total expression of immune-annotated genes is an important
measure of the total activity of cells, it is possible that such a pattern is driven by the expression
of a few highly expressed genes. Thus, to assess broader trends across the transcriptome, we
additionally analyzed the expression of each one-to-one orthologous gene in our dataset and
classified them as hypometric, hypermetric, or of uncertain relation to body mass. Using Fisher's
exact tests on the resulting contingency tables, we found that individual immune-annotated genes
were more likely than non-immune-annotated genes to scale hypermetrically in terms of their
LPS-induced expression and relative abundance after LPS, but not their baseline relative
abundance (Table 2; Data S2).
209

| Parameter | Gene expression context | Allometry | Immune genes | Non-immune genes | P-value (Fisher) |
|---|---|---|---|---|---|
| $\beta_2$ | LPS(-) | hypermetric | 56 (12%) | 532 (13%) | 0.4703 |
| $\beta_2$ | LPS(-) | uncertain | 371 (78%) | 3162 (78%) | 0.9534 |
| $\beta_2$ | LPS(-) | hypometric | 48 (10%) | 366 (9%) | 0.4483 |
| $\beta_3$ | Δ-LPS | hypermetric | 69 (15%) | 362 (9%) | **0.0002** |
| $\beta_3$ | Δ-LPS | uncertain | 370 (78%) | 3265 (80%) | 0.2016 |
| $\beta_3$ | Δ-LPS | hypometric | 36 (8%) | 433 (11%) | **0.0381** |
| $\beta_2 + \beta_3$ | LPS(+) | hypermetric | 69 (15%) | 245 (6%) | **< 0.0001** |
| $\beta_2 + \beta_3$ | LPS(+) | uncertain | 376 (79%) | 3527 (87%) | **< 0.0001** |
| $\beta_2 + \beta_3$ | LPS(+) | hypometric | 30 (6%) | 288 (7%) | 0.5701 |

**Table 2. Number of immune- or non-immune-annotated genes with hypometric, hypermetric, or uncertain**
**relation to body mass.** Genes were considered hypometric in each context (baseline relative abundance = LPS(-),
LPS-induced expression = Δ-LPS, and relative abundance after LPS = LPS(+)) if the pGLMM 95% credible interval
for the corresponding beta was entirely negative, hypermetric if it was entirely positive, and uncertain if it included
zero. P-values correspond to Fisher's exact test comparing the proportions of immune or non-immune genes with
each allometric pattern vs. the sum of the other two patterns.
216
***Alternate analyses demonstrate similar results for LPS-induced expression.*** Of the nine
primate species we sampled, six had high-quality reference genomes with standardized
ENSEMBL annotations, but three did not. Because it is possible that the differences in assembly
and annotation quality are associated with unknown biases, we filtered out the three species with
lower-quality references and conducted the same analyses as before. Results of the analyses of
pooled of immune reads were consistent with the full 9-species analysis regarding LPS-induced
expression: both the pool of one-to-one orthologs and the pool of full orthogroups produced
95%-credible hypermetric estimates ($\beta_2 \sim 0.13$ & 0.17). The 6-species pool of one-to-one
orthologs also produced a 95%-credible hypermetric estimate for relative abundance after LPS

($\beta_2 + \beta_3 \sim 0.08$). However, analysis of whole immune orthogroups in this reduced set of species
produced a 95%-credible *hypo*metric estimate of baseline transcript abundance ($\beta_2 \sim -0.13$)
(Table 1, Figs. S8-11). Analysis of individual gene patterns had a similar result: immune genes
were significantly more likely to have hypermetric LPS-induced expression and relative
abundance after LPS, but were significantly *less* likely to have hypermetric baseline relative
abundances (Table S1, Data S3). These slightly different outcomes could result from greater
statistical power in the smaller dataset due to the larger number of genes identified as one-to-one
orthologs, or from a bias in the intercept for lower-quality reference genomes, which would add
noise to baseline abundance across species but leave estimates of LPS-induced expression
unbiased.
236
***Exaggerated LPS response in larger animals supports the Safety Factor hypothesis.*** Our
results show that immune gene expression is indeed associated with body mass in a
disproportionate (i.e., allometric) manner. Large primates increase their immune gene expression
more during simulated sepsis than small primates, which may result in disproportionately greater
relative abundance of immune-related gene transcripts during infection in large species (Fig. 3).
This result suggests that immune defense genes have greater priority in larger animals relative to
other genes, consistent with the Safety Factor hypothesis.
244
Genes with hypermetric responses and abundances were represented by diverse immune
functions, such as pathogen and damage sensors, transcription factors, and effectors (Data S2).
The allometry of immune gene transcription thus appears to be quite general. However, the
mechanisms of regulation still warrant further investigation. Immune genes did vary in their
individual patterns. For example, the LPS sensor toll-like receptor 4 (*TLR4*) and the pro-
inflammatory cytokines interferon gamma (*IFNG*), interleukin 6 (*IL6*), and interleukin 1 beta
(*IL1B*) were all estimated to have hypermetric relative abundance after LPS, consistent with the
Safety Factor hypothesis and the analyses of the sum of all immune genes. The hypermetric LPS-
induced expression and isometric or slightly hypometric baseline relative abundances of *TLR4*,
*IFNG*, and *IL1B* were also consistent with those results (Fig. 1 D,F,H). However, *IL6* contrasted
the general trend, with a strong, 95%-credible signal of hypermetric baseline relative abundance
and slightly hypometric LPS-induced expression (Fig. 1G). This could be related to the complex
roles that *IL6* plays in both immunity and non-immune metabolic functions, which contrasts with
more strictly-immune functions of *TLR4*, *IFNG*, and *IL1B*. We also note that not all immune-
annotated genes would be predicted to result in greater defense capabilities when expressed
more; for example, the immune regulatory protein cytotoxic T-lymphocyte associated protein 4
(*CTLA4*) is a *negative* regulator of inflammation and immune response [29]. This gene was a
clear exception to the general pattern of immune genes: it had 95%-credible hypermetric baseline
relative abundance and a hypometric estimate for LPS-induced expression (Fig. 1E). The
allometry of *CTLA4* was also notable because the direction of LPS response appears to reverse
across the range of primate body sizes: in the smallest species, LPS exposure resulted in

increased relative transcript abundance, while in the largest species, LPS exposure resulted in decreased relative transcript abundance. These patterns could have important implications for the interpretation of studies that use model organisms to draw conclusions about this gene in humans [30]. The variable patterns of these highlighted genes are non-exhaustive examples; further genes and details can be found in Data S2.

***Future directions: sex and intraspecific effects on allometry***. There is extensive evidence that across most species, males tend to have weaker immune responses than females and are more vulnerable to infection [31–34]. How these conditions interact with our results, especially given the patterns of sexual dimorphism displayed by our primate study species, is unknown. Because our samples were taken only from males, it is possible that the allometric coefficients that we describe are different for females. Depending on the physical and regulatory mechanisms that drive our results, we could imagine independence between sex- and allometric- effects, or stronger allometric effects in either sex. Similarly, other intraspecific variation, such as differences in adult size or age within a species, are difficult to assess with the current data. Although we studied adult animals and included intraspecific body size effects in our linear models, these were intended primarily as controls to ensure our primary analysis was focused on the evolutionary-scale differences among species, and our design is not ideal for interpreting the corresponding coefficients [35]. Exploring these questions should be a priority in the future.

***Eco-evolutionary implications of our results.*** Allometry in the magnitude of immune response may affect pathogen and parasite evolution [20]. Studies have shown that host tolerance can reduce selective pressures on pathogens, allowing them to evolve increased growth rates and increased pathology upon spillover to less-tolerant hosts [36]. Evidence in favor of the Safety Factor hypothesis suggests that smaller species may indeed tend to be more tolerant of pathogens; thus, spillover from small hosts to large may tend to result in infections with more severe pathologies than spillovers in the reverse direction. However, a straight line cannot necessarily be drawn from greater immune gene expression (or greater immune cell concentrations) to better clearance of pathogens. In another recent study, we found isometry in the constitutive ability of blood to kill three very different microbes [37]. In larger animals, metabolic constraints may cause each transcript and each granulocyte to be less effective. Paired with the present results, perhaps gene expression must be engaged hypermetrically to produce functional isometry. Thus, evolutionary mechanisms may simply compensate for the physical constraints imposed by body size variation.

**Materials and Methods:**
***Sample collection***. We collaborated with San Antonio Primate Center, the Duke Lemur Center, the Sedgwick County Zoo and the USF Health Clinic to obtain fresh whole blood samples from nine primate species, including humans, spanning more than three orders of magnitude of body mass, ranging from a 64 g *Microcebus murinus* to a 124,100 g *Pongo abelii*. Our final dataset

included two aye-ayes (*Daubentonia madagascariensis*), four ring-tailed lemurs (*Lemur catta*),
four tufted capuchins (*Sapajus apella*), four Rhesus macaques (*Macaca mulatta*), two olive
baboons (*Papio anubis*), two hybrid olive/yellow baboons (*Papio anubis cynocephalus*, treated
as *P. anubis* for all analyses), four common marmosets (*Callithrix jacchus*), eight mouse lemurs
(in a single pooled sample; *M. murinus*), one human (*Homo sapiens*), and one Sumatran
orangutan (*P. abelii*). We focused sampling on males to minimize sex differences in gene
expression. (Fig. 2, Data S1).
313
A fresh, venous whole blood sample (2 mL) was collected from each individual animal by
collaborators and the date and time were noted. One milliliter of fresh blood was then transferred
to each of two TruCulture tubes (Myriad RBM) containing 2 mL of a proprietary media mixture
(1-part blood to 2-parts media). One tube (LPS treatment) contained 1 μg/mL of 055:B5 strain
LPS (782-001087) and the other (null treatment) contained only the growth media (782-001086).
Each tube was then inverted three times to ensure adequate mixing and incubated at 37 °C for 8
hours. We chose 8 hours to balance the initial burst of pro-inflammatory cytokines, which peak
in humans 4 hours after an LPS challenge, and the later increase of other cytokines such as IL8,
which peaks at 24 hours [38]. After incubation, tubes were centrifuged at 400-500 g for 10 min.
The seraplas filter was then inserted into the tube and the supernatant (plasma/media layer) was
removed. The cells left at the bottom of the tube were then transferred to a new 2 mL cryovial,
RNAlater (Qiagen) was added and frozen at -80°C until RNA extraction could be completed. In
the case where a 2 mL blood draw was not possible (only for 1 marmoset and 1 capuchin), the
ratios inside the TruCulture tube were adjusted so that the 1:2 blood:media was consistent across
samples. Because of the small body mass of the mouse lemurs, we pooled blood from eight
individuals and distributed that 2 mL blood pool equally across the two samples (null and LPS).
De-identified blood was collected from a single human. This study was performed in accordance
with Institutional Review Board Approved Protocols as well as guidelines in the 1964
Declaration of Helsinki and its later amendments, with written informed consent obtained from
the patient.
334
***Whole blood RNA extraction.*** We used a Trizol-based protocol to extract RNA from primate
samples using the Direct-zol RNA Microprep kit (R2060, Zymo Research). All steps were
centrifuged at 13,000 g for 30 s unless otherwise noted. Briefly, 9 μl Proteinase K was added to a
300 μl sample stored in RNAlater and incubated for 30 min. Then, 2.0 mL Trizol was added to
the sample and incubated while vortexing for 15 min. We then added 2.3 mL ethanol to the
mixture and vortexed it. The sample was added to a Zymo-spin IC Column fitted with a
collection tube in 700 μl increments (the maximum loading volume of the column). After each
increment, the sample was centrifuged, and the flow-through discarded. We then performed the
DNase treatment steps. We added 400 μl RNA wash buffer and centrifuged the sample, and then
added 5 μl DNAse 1 and 35 μl DNA digestion buffer and incubated it for 15 min. We added 400
μl Direct-zol RNA prewash buffer to the column, centrifuged and discarded flow-through (x2).

Then 700 µl of RNA wash buffer was added to the column and centrifuged for 1.5 min to ensure
removal of buffers. Finally, 10 µl of RNase-free water was then added directly to the column
matrix, incubated for 1 min, and centrifuged. Concentrations and quality of each sample were
checked via RNA broad range Qubit fluorometric quantification (Q10210, Thermo Fisher
Scientific) and Agilent tape station 2200 (Agilent), respectively.

***Library preparation and RNAseq.*** Extracted RNA samples were submitted to the University of
Illinois at Urbana-Champaign Roy J. Carver Biotechnology Center for library preparation.
Libraries were prepared with the Ovation SOLO RNAseq kit (M01406, Tecan) with probes for
human rRNA and globin RNA. They were sequenced with the NovaSeq 6000 with an S4 flow-
cell and 150 bp paired-end reads. Reads were quality-trimmed before delivery and further
analysis, resulting in an average of 90 bp.

***Reference transcriptomes, orthogroups, and gene expression quantification.*** ENSEMBL
(release 109) [39] reference transcriptomes (*.cdna.all.fa.gz) were used for *M. murinus*, *C.
jacchus*, *P. anubis*, *M. mulatta*, *P. abelii*, and *H. sapiens*. Because *L. catta*, *D. madagascariensis*,
and *S. apella* are not yet integrated into ENSEMBL, reference transcriptomes were created with
the following workflow. First, NCBI GenBank genomes were acquired (GCF_020740605.2,
GCA_023783475.1, and GCF_009761245.1, respectively), and HISAT2 v2.1.0 [40] indexes
were built. Then, sequencing reads for each sample were assembled into 'superreads' using the
superreads.pl script provided for use with StringTie v1 [41]. Superreads and unassembled paired-
end reads were then mapped to their respective genome using HISAT2 with options --no-
discordant --no-mixed --dta-cufflinks. StringTie v2.2.1[42] was then used to generate
transcriptomes with default settings. TransDecoder v5.5.0 [43] was used to predict peptide
sequences. The longest predicted peptide was isolated for each gene, and these peptides were
input to Orthofinder v2.5.4 [44] in combination with ENSEMBL reference peptides (filtered
from Compara.109.protein_default.aa.fasta.gz) to identify orthogroups across all nine species.
The phylogeny from Hedges *et al.* 2015 was subsetted and used as a fixed species tree in the
Orthofinder analysis. Orthogroups were identified using Orthofinder's root-level
'Phylogenetic_Hierarchical_Orthogroups' output file (N0.tsv), and and 1:1 orthologs were
determined by filtering the rows of this table. Gene expression estimates were generated using
Salmon v1.6.0 [45] with default settings and raw sequencing reads from each sample.

***Differential abundance analysis of genes.*** To assess differential abundance of each gene, count
data were analyzed with phylogenetic generalized linear mixed models using the function
pglmm() from the R package *phyr* v1.1.0 [46]. To approximately account for the discrete and
compositional nature of RNASeq count data, a Poisson likelihood with a log link was used with
an offset term unique to each observation. This offset was determined by importing Salmon
quantification outputs into an R object with the package *tximport* v1.26.1 [47], filtering the genes
to include only 1:1 orthologs, and using the DESeq2 v1.38.2 [48] function *estimateSizeFactors*.

The resulting 'normalization factors' are designed to compensate for varying sequencing depth,
compositionality, and average transcript length across samples. Binary LPS treatment, the
standardized log of each species' average body mass [28], the standardized log of species-
normalized individual body size, and the interaction of LPS treatment with each body size effect,
were modeled as fixed effects ($\beta_1$-$\beta_5$), with random effects for individual, independent and
identically distributed (i.i.d.) species, species phylogeny, and the interactions between the
treatment and i.i.d. species and species phylogeny. *phyr* also automatically includes an
observation-level random effect to account for overdispersion relative to the expected variance of
the Poisson likelihood. The aforementioned phylogenetic tree from Hedges *et al.* 2015 [27] was
used for phylogenetic random effects. In this model, the main effect of LPS treatment ($\beta_1$) can be
interpreted as the change in expression after LPS treatment for each gene at the ancestral body
size (somewhat arbitrary for the purposes of this study, and determined by optimization with the
*ace* function from the R package *ape* 5.7-1 [49]), the main effect of species' average body mass
($\beta_2$) can be interpreted as the allometric effect on the baseline relative abundance of each gene
(e.g., LPS(-)), the interaction between LPS treatment and species' average body mass ($\beta_3$) can be
interpreted as the allometric effect on change in expression after LPS treatment (e.g., $\Delta$-LPS), the
main effect of species-normalized individual body size ($\beta_4$) may be caused by either aging-
related changes or within-species allometric differences in the baseline relative abundance of
each gene, and the interaction between LPS treatment and species-normalized individual body
size ($\beta_5$) may be caused by either aging-related changes or within-species allometric differences
in the change in expression after LPS treatment. The random effect for individuals enables the
model to focus on paired differences between and LPS and null treatment, the main species
effects control for incidentally correlated evolution of body mass and baseline gene expression,
and the species interaction effects control for incidentally correlated evolution of body mass and
the change in expression during LPS treatment. The model was fit using approximate Bayesian
inference with default priors with *INLA* v21.02.23 [50]. The sums of $\beta_2$ and $\beta_3$ and of $\beta_4$ and $\beta_5$
were calculated during model fitting using *INLA*'s 'lincomb' option, to represent the actual
allometric slopes among LPS-treated samples (e.g., LPS(+)). Models that did not converge were
retried up to 15 times, and genes were filtered from further analysis if they did not converge.
These genes (55 out of 4590 1:1 orthologs) tended to have some samples with low counts, which
can exacerbate overfitting and lead to singular model matrices. Fixed effect coefficients were
extracted, and individual genes were determined to have significant differential relative
abundance if their 95% credible interval did not contain zero. Two-by-two contingency tables
were constructed summarizing the number of genes that were allometric (or not) and immune-
annotated (or not), and Fisher's exact tests were used to determine if each allometric pattern was
more likely in immune vs. non-immune genes.
422
***Differential abundance analysis of total immune genes*.** Orthologous gene sets were annotated
with their corresponding immune functions as defined by Deschamps *et al*. 2016 [51]. External
gene names were determined for each ortholog by querying the Ensembl BioMart [39] with the

R package *biomaRt* v2.54.1 [52], using human members of each orthogroup, and these gene
names were used to extract immune function annotations from Table S1 from Deschamps *et al*.
Two workflows were used to assess the total expression of immune genes. In the first, counts for
all immune-annotated 1:1 orthologs were summed, and average gene lengths for the pool were
calculated for use in DESeq2's normalization factor calculations. This method is conservative in
focusing only on identifiable 1:1 orthologs and can utilize observation-specific normalization
factors so that results may be interpreted as differences in the number of transcripts across
samples. In the second, counts of all genes that belonged to immune-annotated orthogroups were
pooled, and 'size factors' that were consistent for a given sample were calculated by taking the
geometric mean of the normalization factors of all 1:1 orthologs. This method is more
comprehensive, covering all genes with likely immune functions, but because it does not
compensate for differences in average transcript length across samples, it is *less confidently*
interpretable as differences in numbers of transcripts. The choice to use 'size factors' instead of
'normalization factors' here was made because it is likely that gene predictions are fragmented
for the three species with lower-quality references, and apparent gene duplications or losses in
these species could artificially influence the predicted average transcript length. This version is
better interpreted as the amount of RNA with a given annotation, rather than the abundance of
transcripts with a given annotation. For both analyses, total immune gene abundance was
modeled with *phyr* and *INLA* using the same protocol and formula as analyses of individual
genes. Model diagnostics were performed using *DHARMa* [53]

To analyze only the six species with higher-quality ENSEMBL reference genomes, all
differential abundance methods were repeated after filtering the previous Salmon and
Orthofinder outputs.

All analyses that used R throughout the manuscript used version 4.2.2 [54]. The packages *ggtree*
v3.6.2 [55] and *ggplot2* v3.4.1 [56] were used for plotting.

**Acknowledgements:** The authors thank Debbie Christian, Deborah Chavez, and Chris Chen at
the Southwest National Primate Research Center at the Texas Biomedical Research Institute;
Erin Ehmke, Kay Welser, and Megan Davison at the Duke Lemur Center; and Jennifer Kropf
and Sandy Wilson at the Sedgwick County Zoo for their collaboration and support in obtaining
and processing these primate samples. We thank Manfred Schmolz at Myriad RBM for help with
TruCulture techniques and Alvaro Hernandez in troubleshooting RNA extractions and
sequencing. Finally, thanks to Samantha Oakey, Matthew Mercurio, Min Zhang, and the
University of South Florida Genomics Program Sequencing Core and Computational
Core/Omics Hub for laboratory and computational support.
**Funding:**
National Science Foundation Division of Integrative Organismal Systems grant NSF-IOS
1656551 (CJD)

National Science Foundation Division of Integrative Organismal Systems grant NSF-IOS
1656618 (LBM)

**Author contributions:**
468
Conceptualization: CJD, LBM, RHYJ, RM
Data curation: RM, ECR, SRA, RAM
Formal analysis: RM
Funding acquisition: CJD, RHYJ, LBM
Investigation: ECR, SRA
Methodology: RM, ECR, SRA, CJD, LBM, RHYJ
Project administration: CJD, LBM
Resources: ECR, CJD, LBM, JWL
Supervision: CJD, LBM, RHYJ
Visualization: RM
Writing – original draft: RM, CJD, LBM
Writing – review & editing: RM, CJD, LBM, RHYJ, RAM, ECR, SRA, JWL

**Competing interests:** J.W.L. is an employee and shareholder of Bluebird Bio, a speaker,
consultant, and research funding recipient of Horizon Therapeutics, a speaker and consultant
for Sobi, and on the advisory board of ADMA.

**Data and materials availability:** Quality-trimmed RNASeq data is available in NCBI
BioProject PRJNA834816. All code used for analysis is available at
https://github.com/rmcminds/McMinds_et_al_2022 and at Zenodo (doi:
10.5281/zenodo.10779069).

**Supplementary Materials:**
Figs. S1 to S11
Table S1
Data S1 to S3
493

**Total immune gene expression difference between LPS and Null**

495  **Fig. S1. Summed counts of all immune-annotated 1:1 orthologous genes show hypermetric response**

496  **to sepsis.** This figure draws on the same data as Figure 3 in the main text, but shows the difference

497  between LPS- and Null-treated samples from each individual rather than the raw abundance values. Body

498  mass is plotted on the log scale.

**Total immune gene expression**

499
**Fig S2. Summed counts of all immune-annotated reads show hypermetric concentrations after LPS.**
This figure is analogous to Figure 3 in the main text, but is the sum of all reads annotated with immune
functions, even if they belong to orthogroups that had duplications or gene deletions. This analysis is
more comprehensive as it was able to include more genes (>4000 compared to >1200), but it is probably
more susceptible to differences in gene lengths, or apparent gene lengths due to lower quality
transcriptome assemblies. As an alternate method with different assumptions that shows something
similar, it's a nice backup. Theoretically, this figure might be best interpreted as an increase in total
immune-related RNA, whereas Fig. 3 depicts an increase in the total number of immune-annotated
transcripts (but in the absence of widescale biases in gene or transcript lengths, the interpretation should
be similar).
510

**Total immune gene expression difference between LPS and Null**

511
512 **Fig. S3. Sum of all counts of reads that belong to immune-annotated orthogroups.** This figure is the
513 same as Figure S1, but this figure includes all immune reads rather than just 1:1 orthologs.
514

515
**Fig. S4. Density plots for 9-species 1:1 ortholog analysis. (A)** Posterior distributions of the residual
variance for each random effect. Effects are ordered by median estimates. Names are in *phyr* notation:
1|obs is the observation-level extra-poisson variance; 1|species__ is the (Brownian) phylogenetic effect on
baseline abundance; 1|species__@treatment is the phylogenetic effect on LPS-induced expression;
1|individual is the effect of individual; 1|species is the non-phylogenetic effect of species on baseline
abundance; and 1|species@treatment is the non-phylogenetic effect of species on LPS-induced
expression. **(B)** Posterior distributions of the fixed effect coefficients. The grand intercept (alpha) has
been removed, and coefficients are ordered by median estimates. The median and 95% credible intervals
are shown on each axis. Density plots are blue if the 95% credible interval does not include zero.

525
**Fig. S5. DHARMa plots for 9-species 1:1 ortholog analysis.** The significant results from the Kolgorov-
Smirnov and dispersion tests are consistent with an overfit model applied to count data. This outcome
results from the ability of the continuous parameters, including observation-level random effects, to
predict each observation almost perfectly, leaving little residual variance to be explained by the Poisson
count model. This result is common for Poisson models with observation-level random effects and is
usually associated with lower statistical power. This suggests that our analysis is conservative and should
not strongly influence the interpretation of effects that are found to be significant.
533

Fig. S6. Density plots for 9-species orthogroup analysis. (A) Posterior distributions of the residual variance for each random effect. Effects are ordered by median estimates. Names are in *phyr* notation: 1|obs is the observation-level extra-poisson variance; 1|species__ is the (Brownian) phylogenetic effect on baseline abundance; 1|species__@treatment is the phylogenetic effect on LPS-induced expression; 1|individual is the effect of individual; 1|species is the non-phylogenetic effect of species on baseline abundance; and 1|species@treatment is the non-phylogenetic effect of species on LPS-induced expression. (B) Posterior distributions of the fixed effect coefficients. The grand intercept (alpha) has been removed, and coefficients are ordered by median estimates. The median and 95% credible intervals are shown on each axis. Density plots are blue if the 95% credible interval does not include zero.

545
546 **Fig. S7. DHARMa plots for 9-species orthogroup analysis.** This is analogous to Fig S5, with similar
547 interpretations of overfitting and the conservative nature of the tests.
548
549

| Parameter | Gene expression context | Allometry | Immune genes | Non-immune genes | P-value (Fisher) |
|---|---|---|---|---|---|
| $\beta_2$ | LPS(-) | hypermetric | 186 (19%) | 2494 (24%) | **0.0012** |
| $\beta_2$ | LPS(-) | uncertain | 674 (69%) | 7010 (66%) | 0.0824 |
| $\beta_2$ | LPS(-) | hypometric | 116 (12%) | 1071 (10%) | 0.0876 |
| $\beta_3$ | Δ-LPS | hypermetric | 116 (12%) | 765 (7%) | **< 0.0001** |

| | | | | | |
|---|---|---|---|---|---|
| $\beta_3$ | $\Delta$-LPS | uncertain | 751 (77%) | 8136 (77%) | 1.0000 |
| $\beta_3$ | $\Delta$-LPS | hypometric | 109 (11%) | 1674 (16%) | **0.0001** |
| $\beta_2 + \beta_3$ | LPS(+) | hypermetric | 129 (15%) | 676 (6%) | **< 0.0001** |
| $\beta_2 + \beta_3$ | LPS(+) | uncertain | 770 (79%) | 8948 (85%) | **< 0.0001** |
| $\beta_2 + \beta_3$ | LPS(+) | hypometric | 77 (6%) | 951 (9%) | 0.2644 |

**Table S1. Number of immune- or non-immune-annotated genes with hypometric, hypermetric, or uncertain relation to body size (6-species analysis).** Genes were considered hypometric in each context (baseline abundance = LPS(-), LPS-induced expression = $\Delta$-LPS, and abundance after LPS = LPS(+)) if the pGLMM 95% credible interval for the corresponding beta was entirely negative, hypermetric if it was entirely positive, and uncertain if it included zero. Innate immune annotations were derived from Deschamps *et al*. 2016. P-values correspond to Fisher's exact test comparing the proportions of immune or non-immune genes with each allometric pattern vs. the sum of the other two patterns.

**Fig. S8. Density plots for 6-species 1:1 ortholog analysis. (A)** Posterior distributions of the residual
variance for each random effect. Effects are ordered by median estimates. Names are in *phyr* notation:
1|obs is the observation-level extra-poisson variance; 1|species__ is the (Brownian) phylogenetic effect on
baseline abundance; 1|species__@treatment is the phylogenetic effect on LPS-induced expression;
1|individual is the effect of individual; 1|species is the non-phylogenetic effect of species on baseline
abundance; and 1|species@treatment is the non-phylogenetic effect of species on LPS-induced
expression. **(B)** Posterior distributions of the fixed effect coefficients. The grand intercept (alpha) has
been removed, and coefficients are ordered by median estimates. The median and 95% credible intervals
are shown on each axis. Density plots are blue if the 95% credible interval does not include zero.

569
570 **Fig. S9. DHARMa plots for 6-species 1:1 ortholog analysis.** This is analogous to Fig S5, with similar
571 interpretations of overfitting and the conservative nature of the tests.
572

573
**Fig. S10. Density plots for 6-species orthogroup analysis. (A)** Posterior distributions of the residual
variance for each random effect. Effects are ordered by median estimates. Names are in *phyr* notation:
1|obs is the observation-level extra-poisson variance; 1|species__ is the (Brownian) phylogenetic effect on
baseline abundance; 1|species__@treatment is the phylogenetic effect on LPS-induced expression;
1|individual is the effect of individual; 1|species is the non-phylogenetic effect of species on baseline
abundance; and 1|species@treatment is the non-phylogenetic effect of species on LPS-induced
expression. **(B)** Posterior distributions of the fixed effect coefficients. The grand intercept (alpha) has
been removed, and coefficients are ordered by median estimates. The median and 95% credible intervals
are shown on each axis. Density plots are blue if the 95% credible interval does not include zero.
583

584
585 **Fig. S11. DHARMa plots for 6-species orthogroup analysis.** This is analogous to Fig S5, with similar
586 interpretations of overfitting and the conservative nature of the tests.

587
**References**

1. West GB. 2017 *Scale: the universal laws of growth, innovation, sustainability, and the pace of*
*life in organisms, cities, economies, and companies.*

2. Savage VM, Gillooly JF, Woodruff WH, West GB, Allen AP, Enquist BJ, Brown JH. 2004
The predominance of quarter☐power scaling in biology. *Funct Ecol* **18**, 257–282.
(doi:10.1111/j.0269-8463.2004.00856.x)

3. Calder WA. 1981 Scaling of Physiological Processes in Homeothermic Animals. *Annu. Rev.*
*Physiol.* **43**, 301–322. (doi:10.1146/annurev.ph.43.030181.001505)

4. Calder WA. 1987 Scaling Energetics of Homeothermic Vertebrates: an Operational
Allometry. *Annu. Rev. Physiol.* **49**, 107–120. (doi:10.1146/annurev.ph.49.030187.000543)

5. Prothero JW. 2015 *The design of mammals*. (doi:10.1017/cbo9781316275108)

6. Calder WA. 1996 *Size, function, and life history*. Harvard University Press.

7. Brown JH, Gillooly JF, Allen AP, Savage VM, West GB. 2004 Toward a metabolic theory of
ecology. *Ecology* **85**, 1771–1789. (doi:10.1890/03-9000)

8. Feldman HA, McMahon TA. 1983 The 3/4 mass exponent for energy metabolism is not a
statistical artifact. *Resp Physiol* **52**, 149–163. (doi:10.1016/0034-5687(83)90002-6)

9. Kleiber M. 1932 Body size and metabolism. *Hilgardia* **6**, 315–353.
(doi:10.3733/hilg.v06n11p315)

10. Schmidt-Nielsen K. 1984 *Scaling: Why is Animal Size so Important?* Cambridge University
Press.

11. Peters RH. 1983 *The Ecological Implications of Body Size*. Cambridge University Press.
(doi:10.1017/cbo9780511608551)

12. West GB, Brown JH, Enquist BJ. 1997 A general model for the origin of allometric scaling
laws in biology. *Science* **276**, 122–126. (doi:10.1126/science.276.5309.122)

13. Horn HS. 2004 Commentary on Brown et al.'s "Toward a metabolic theory of ecology."
*Ecology* **85**, 1816–1818. (doi:10.1890/03-0757)

14. O'Connor MP, Kemp SJ, Agosta SJ, Hansen F, Sieg AE, Wallace BP, McNair JN, Dunham
AE. 2007 Reconsidering the mechanistic basis of the metabolic theory of ecology. *Oikos* **116**,
1058–1072. (doi:10.1111/j.0030-1299.2007.15534.x)

15. Downs CJ, Hayes JP, Tracy CR. 2008 Scaling metabolic rate with body mass and inverse
body temperature: a test of the Arrhenius fractal supply model. *Funct Ecol* **22**, 239–244.
(doi:10.1111/j.1365-2435.2007.01371.x)

16. Glazier DS. 2015 Is metabolic rate a universal 'pacemaker' for biological processes? *Biol*
*Rev* **90**, 377–407. (doi:10.1111/brv.12115)

17. Dingli D, Pacheco JM. 2006 Allometric scaling of the active hematopoietic stem cell pool
across mammals. *Plos One* **1**, e2. (doi:10.1371/journal.pone.0000002)

18. Cornelius Ruhs E, Martin LB, Downs CJ. 2020 The impacts of body mass on immune cell
concentrations in birds. *Proc Royal Soc B Biological Sci* **287**, 20200655.
(doi:10.1098/rspb.2020.0655)

19. Downs CJ, Dochtermann NA, Ball R, Klasing KC, Martin LB. 2020 The effects of body
mass on immune cell concentrations of mammals. *Am Nat* **195**, 107–114. (doi:10.1086/706235)

20. Downs CJ, Schoenle LA, Han BA, Harrison JF, Martin LB. 2019 Scaling of host
competence. *Trends Parasitol* **35**, 182–192. (doi:10.1016/j.pt.2018.12.002)

21. Lopez MLD, Lin Y, Schneider SQ, Hsieh C, Shiah F, Machida RJ. 2023 Allometric scaling
of interspecific RNA transcript abundance to extend the use of metatranscriptomics in
characterizing complex communities. *Mol Ecol Resour* **23**, 52–63. (doi:10.1111/1755-
0998.13711)

22. Hawash MBF, Sanz-Remón J, Grenier J-C, Kohn J, Yotova V, Johnson Z, Lanford RE,
Brinkworth JF, Barreiro LB. 2021 Primate innate immune responses to bacterial and viral
pathogens reveals an evolutionary trade-off between strength and specificity. *Proc National
Acad Sci* **118**, e2015855118. (doi:10.1073/pnas.2015855118)

23. Vaure C, Liu Y. 2014 A comparative review of Toll-like receptor 4 expression and
functionality in different animal species. *Front Immunol* **5**, 316.
(doi:10.3389/fimmu.2014.00316)

24. Coley PD, Bryant JP, III FSC. 1985 Resource availability and plant antiherbivore defense.
*Science* **230**, 895–899. (doi:10.1126/science.230.4728.895)

25. Martin LB, Weil ZM, Nelson RJ. 2007 Immune defense and reproductive pace of life in
Peromyscus mice. *Ecology* **88**, 2516–2528. (doi:10.1890/07-0060.1)

26. Martin LB, Weil ZM, Nelson RJ. 2008 Fever and sickness behaviour vary among congeneric
rodents. *Funct Ecol* **22**, 68–77. (doi:10.1111/j.1365-2435.2007.01347.x)

27. Hedges SB, Marin J, Suleski M, Paymer M, Kumar S. 2015 Tree of Life Reveals Clock-Like
Speciation and Diversification. *Mol Biol Evol* **32**, 835–845. (doi:10.1093/molbev/msv037)

28. Burger JR, George MA, Leadbetter C, Shaikh F. 2019 The allometry of brain size in
mammals. *J Mammal* **100**, 276–283. (doi:10.1093/jmammal/gyz043)

29. Nakajima T, Suarez CJ, Lin K-W, Jen KY, Schnitzer JE, Makani SS, Parker N, Perkins DL,
Finn PW. 2010 T Cell Pathways Involving CTLA4 Contribute To a Model of Acute Lung Injury.
*J Immunol* **184**, 5835–5841. (doi:10.4049/jimmunol.0903238)

30. Martin LB, Hanson HE, Hauber ME, Ghalambor CK. 2021 Genes, Environments, and
Phenotypic Plasticity in Immunology. *Trends Immunol.* **42**, 198–208.
(doi:10.1016/j.it.2021.01.002)

31. Zuk M. 2009 The Sicker Sex. *PLoS Pathog.* **5**, e1000267.
(doi:10.1371/journal.ppat.1000267)

32. Oliva M *et al.* 2020 The impact of sex on gene expression across human tissues. *Science* **369**,
eaba3066. (doi:10.1126/science.aba3066)

33. Klein SL, Flanagan KL. 2016 Sex differences in immune responses. *Nat. Rev. Immunol.* **16**,
626–638. (doi:10.1038/nri.2016.90)

34. Rolff J. 2002 Bateman's principle and immunity. *Proc. Biol. Sci.* **269**, 867–72.
(doi:10.1098/rspb.2002.1959)

35. Westreich D, Greenland S. 2013 The Table 2 Fallacy: Presenting and Interpreting
Confounder and Modifier Coefficients. *Am J Epidemiol* **177**, 292–298. (doi:10.1093/aje/kws412)

36. Brook CE, Rozins C, Guth S, Boots M. 2023 Reservoir host immunology and life history
shape virulence evolution in zoonotic viruses. *PLOS Biol.* **21**, e3002268.
(doi:10.1371/journal.pbio.3002268)

37. Downs CJ, Schoenle LA, Goolsby EW, Oakey SJ, Ball R, Jiang RHY, Martin LB. 2023
Large Mammals Have More Powerful Antibacterial Defenses Than Expected from Their
Metabolic Rates. *Am. Nat.* **201**, 287–301. (doi:10.1086/722504)

38. DeForge LE, Remick DG. 1991 Kinetics of TNF, IL-6, and IL-8 gene expression in LPS-
stimulated human whole blood. *Biochem Bioph Res Co* **174**, 18–24. (doi:10.1016/0006-
291x(91)90478-p)

39. Cunningham F *et al.* 2021 Ensembl 2022. *Nucleic Acids Res* **50**, D988–D995.
(doi:10.1093/nar/gkab1049)

40. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019 Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907–915.
(doi:10.1038/s41587-019-0201-4)

41. Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. 2015 StringTie
enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol* **33**,
290–295. (doi:10.1038/nbt.3122)

42. Kovaka S, Zimin AV, Pertea GM, Razaghi R, Salzberg SL, Pertea M. 2019 Transcriptome
assembly from long-read RNA-seq alignments with StringTie2. *Genome Biol* **20**, 278.
(doi:10.1186/s13059-019-1910-1)

43. Haas B. 2022 *TransDecoder.*

44. Emms DM, Kelly S. 2019 OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol* **20**, 238. (doi:10.1186/s13059-019-1832-y)

45. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017 Salmon provides fast and bias-
aware quantification of transcript expression. *Nat Methods* **14**, 417–419.
(doi:10.1038/nmeth.4197)

46. Li D, Dinnage R, Nell LA, Helmus MR, Ives AR. 2020 phyr: An r package for phylogenetic
species☐distribution modelling in ecological communities. *Methods Ecol Evol* **11**, 1455–1463.
(doi:10.1111/2041-210x.13471)

47. Soneson C, Love MI, Robinson MD. 2016 Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences. *F1000research* **4**, 1521.
(doi:10.12688/f1000research.7563.2)

48. Love MI, Huber W, Anders S. 2014 Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* **15**. (doi:10.1186/s13059-014-0550-8)

49. Paradis E, Claude J, Strimmer K. 2004 APE: Analyses of Phylogenetics and Evolution in R
language. *Bioinformatics* **20**, 289–290. (doi:10.1093/bioinformatics/btg412)

50. Rue H, Riebler A, Sørbye SH, Illian JB, Simpson DP, Lindgren FK. 2016 Bayesian
computing with INLA: a review. *Arxiv*

51. Deschamps M, Laval G, Fagny M, Itan Y, Abel L, Casanova J-L, Patin E, Quintana-Murci L.
2016 Genomic signatures of selective pressures and introgression from archaic hominins at
human innate immunity genes. *Am J Hum Genetics* **98**, 5–21. (doi:10.1016/j.ajhg.2015.11.014)

52. Durinck S, Moreau Y, Kasprzyk A, Davis S, Moor BD, Brazma A, Huber W. 2005 BioMart
and Bioconductor: a powerful link between biological databases and microarray data analysis.
*Bioinformatics* **21**, 3439–3440. (doi:10.1093/bioinformatics/bti525)

53. Hartig F. 2022 *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)*
*Regression Models*. See https://CRAN.R-project.org/package=DHARMa.

54. R Core Team. 2021 *R: a language and environment for statistical computing*. Vienna,
Austria: R Foundation for Statistical Computing. See https://www.R-project.org/.

55. Yu G, Smith DK, Zhu H, Guan Y, Lam TT. 2017 ggtree: an R package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol*
*Evol* **8**, 28–36. (doi:10.1111/2041-210x.12628)

56. Wickham H. 2016 *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
See https://ggplot2.tidyverse.org.