***Emiliania huxleyi* virus arrests host calcification and nutrient consumption, and**
triggers shifts in organic stoichiometry

Tamar Dikstein[a,b], Gilad Antler[c,b,], Andre Pellerin[d], Shlomit Sharoni[e], Miguel J. Frada[a,b,1]

[a]Department of Ecology, Evolution and Behavior, Silberman Institute of Life Sciences, The
Hebrew University of Jerusalem, Jerusalem, Israel

[b]The Interuniversity Institute for Marine Sciences in Eilat, Eilat, Israel

[c]Department of Earth & Environmental Sciences, Ben-Gurion University of the Negev,
Beersheba, Israel

[d]Institut des sciences de la mer, Université du Québec à Rimouski and GEOTOP, Rimouski,
Québec, Canada

[e]Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of
Technology, Cambridge, United States of America

Corresponding author: miguel.frada@mail.huji.ac.il

Keywords: Coccolithophore, Virus, Nutrient uptake, Calcification, Elemental
stoichiometry.

Abstract

Blooms of the dominant coccolithophore *Emiliania huxleyi* are routinely infected by a
specific lytic virus (EhV), which rapidly kills host cells triggering bloom termination and
organic and inorganic carbon export. However, the impact of EhV on the dynamic of
resource acquisition and cellular stoichiometry remains unknown, limiting the current
understanding of the ecological and biogeochemical significance of *E. huxleyi* blooms.
To tackle this knowledge gap, we used algal and EhV cultures to determine over the
course of infections the dynamics of alkalinity, modulated by calcification, nitrate and
phosphate consumption and organic matter stoichiometry. We found that within 24hr
alkalinity concentration stabilized and nutrient uptake declined to background levels. In
parallel, the stoichiometric ratio of carbon to nitrogen increased and the nitrogen to
phosphorus ratio declined. These variations likely resulted from lipid accumulation

required for viral replication and the differential retention of phosphorus-rich

macromolecular pools in decaying cells, respectively. Finally, during post-infection and

decline of the host population, a progressive enrichment in phosphorus relative to

nitrogen and carbon was measured, detected in the remaining cell lysates. We estimate

that this stoichiometric shift is driven by the accumulation of heterotrophic bacteria

involved in the degradation of organic material. Viral-mediated cell remodeling and

consequent shifts in biomass stoichiometry likely impacts the patterns of nutrient cycling

and biological carbon pump efficiency during large-scale blooms in the oceans.

Introduction

Phytoplankton are a taxonomically diverse group of photosynthetic microorganisms that

inhabit the sunlit layer of the oceans where they are responsible for about 45% of global

primary production and for driving the biological carbon pump (1). This latter process

results in a net drawdown of atmospheric CO_2 into the ocean's interior, henceforth

influencing global climate (2). Biogeochemical cycles of the major nutrients (nitrogen; N

and phosphorus; P) and carbon (C) are interlinked (3, 4). The ratio of N to P in seawater

determines which nutrient limits biological productivity in the ocean, with cascading

effects on the marine food web (5–7). In addition, the ratio of carbon to the limiting

nutrient in organic matter, determines the maximum amount of carbon that can be fixed

and exported (8). By consequence, phytoplankton elemental stoichiometry is a main

component regulating the efficiency of the biological carbon pump.

Viruses are ubiquitous regulators of microbial growth and evolution in the oceans (9, 10).

Mortality caused by lytic viruses is estimated to influence marine primary production,

food-web dynamics and nutrient cycles in the oceans (11, 12). However, to our

knowledge little is known on the impact of viral infections on phytoplankton and

particulate organic carbon stoichiometry. Quantifying the impact of viral infection on

nutrient cycling in phytoplankton cells is critical to improve our understanding of marine

ecosystem functioning, biogeochemical cycles and the potential feedback on the global

climate. Here, we address the impact of viral infections in *Emiliania huxleyi*, a

cosmopolitan unicellular calcifying photoautotrophic organism belonging to the

coccolithophore clade and capable of forming large-scale blooms in the oceans.

Coccolithophores are an important group of marine phytoplankton estimated to

contribute 1% to 10% of primary production and phytoplankton biomass, increasing up to

40% during bloom conditions (13, 14). As a characteristic, coccolithophores produce an

outer covering of calcite plates termed coccoliths (15, 16). This attribute confers

coccolithophores a unique functional role in pelagic ecosystems. The biosynthesis of

coccoliths is an energy costly process involving the active transport of the calcification

substrates, Ca^{2+} and HCO^-_3, dissolved in seawater into the cell where coccoliths are

precipitated (16). In doing so, coccolithophores reduce both seawater total dissolved

inorganic carbon (DIC) and alkalinity, which changes the equilibrium between the

different DIC forms and reduces the potential for seawater CO_2 uptake and to buffer for

pH changes (17). Moreover, the biological precipitation and export of calcite results in a

net release of CO_2 to the atmosphere, a mechanism referred to as the alkalinity pump

(or carbonate counter pump), which counter acts the biological carbon pump.

Nonetheless, the impact of any CO_2 production during calcification may be offset through

the ballasting of organic carbon by heavy coccoliths entangled in marine snow, which

can increase the efficiency of the biological carbon pump (17, 18). The relative strength

of the two pumps, represented by the so-called rain ratio, determines to a large extent

the air/sea exchange fluxes of CO_2 and the magnitude of oceanic carbon sequestration,

with important effects on the biogeochemical cycling of carbon and climate (17, 19).

In this context, blooms of *E. huxleyi* bear a significant impact. These blooms are typically

extensive and develop over several weeks in temperate and subpolar waters during late

spring/early summer periods (20–22). The underlying conditions for bloom formation are

numerous and may vary between sites, thus a matter of debate (21, 23). However,

bloom decline is linked to an important extent to lytic infections by a specific *E. huxleyi*

virus (EhV) (24–27), a large double-stranded DNA Coccolithovirus (28). At the cellular

level, EhV infection leads to profound remodelling of the *E. huxleyi* metabolism to

support viral proliferation. These include the activation and recruitment of the host's

oxidative stress (29–31), upregulation of autophagy, nuclear degradation (32), and both

enhance glycolytic fluxes and lipid biosynthesis (33). The latter include triacylglycerols

(33, 34) and specific virus-derived glycosphingolipids that are important structural

components of EhV particles and chemical signals triggering host's programmed cell

death (30, 33). Additionally, EhV infection prompts coccolith shedding from the cell

surface, changing the optical signature of infected populations (35–37), and stimulates

the production of transparent exopolymer particles (TEP) that can form large aggregates

entrapping cells and coccoliths and enhance the vertical export flux of both particulate

organic and inorganic carbon (38, 39).

Although focus has been directed to viral-mediate changes in metabolic and structural

properties of host *E. huxleyi* cells, the impact of EhV on nutrient uptake, calcification

dynamics and ultimately on the basal elemental stoichiometric compositions (C:N:P) of

the viral infected cells (i.e., virocells, (40)) remains to be tested. This limits our

understanding of basic feature of the host-virus interplay between *E. huxleyi* and EhV

and bloom turnover, and the impact of viral infection on carbon export. To tackle this

knowledge-gap, we examined the dynamic of macronutrient (N and P) consumption and

calcification during viral infections, as well as the C:N:P stoichiometric composition of

infected populations in controlled culture conditions. Our study demonstrates the impact

of virus on the regulation of nutrient incorporation and cellular composition of *E. huxleyi*

virocells and provide key insights on ensuing ecological and biochemical consequences

of large-scale viral infections during oceanic blooms.

Results

Algal growth and variations in optical properties

Detailed survey of viral infections was monitored during 72 hr (Fig. 1). The cell growth

developed comparably in infections and controls during the first 36 hr reaching a density

of around 1.25×10^6 cells mL^{-1}. The growth rate over first day averaged ~0.5 ± 0.06 day$^-$

^{1} in both treatments. During the second day, concurrent exponential accumulation of

virions was detected in the infected cultures. Peak viral densities were detected 48 hr

post-infection (~ 2.7×10^8 virions mL^{-1}). The estimated viral burst size was estimated to

be ~230 viral particles cell^{-1}. In parallel, cell density declined down to 0.5×10^6 cells mL^{-1}

by 72 hr, while the control cultures reached the stationary growth phase (~ 2.0×10^6 cells

mL^{-1}) (Fig. 1A).

Average chlorophyll autofluorescence and side scatter measured by flow cytometry were

used to as indicators of cell health and degree of cell calcification. During exponential

growth, both parameters commonly oscillate through day-night cycles due to cell

enlargement and coccolith buildup during exponential growth (41). This was detected

during the first day in both treatments. However, during the second day these diel

oscillation in optical signals stopped and overall declined (Fig. 1B–C). A slight increase

in the average side scatter signal was detected at 72hr, can be explained by the

formation of aggregates of decaying cellular materials with high scattering properties, not
from the growth of living cells.
129

130
Figure 1. Algal and viral growth and variations in chlorophyll autofluorescence and side scatter.
E. huxleyi cultures were infected with EhV 201 and compared with noninfected control cells. (A)
Cell growth and extracellular viral production. (B) Mean population values of Red
autofluorescence. (C) Mean population values of side scatter. The x-axes correspond to hours
post infection. Grey sections represent the night period and dots indicate mean values of each
treatment (Control/Infection), Error bars represent standard error of replicates, n=3.
137

Alkalinity, nitrate and phosphate consumption

During the first day, the consumption of alkalinity (Fig. 2A), nitrate (Fig. 2B) and
phosphate (Fig. 2C) was broadly similar in infected and control cultures. During the first
light phase alkalinity, nitrate and phosphate consumption averaged 13.40 ± 0.90 µmol L^{-1}
h^{-1}, 3.1 ± 0.4 µmol L^{-1} h^{-1} and 0.12 ± 0.06 µmol L^{-1} h^{-1}, respectively. During the first
dark period the consumption of alkalinity was much lower (avg. 1.6 ± 0.8 µmol L^{-1} h^{-1}),
nitrate was about half (avg. 1.20 ± 0.20 µmol L^{-1} h^{-1}), and phosphate identical to light
period (avg. 0.11 ± 0.01 µmol L^{-1}). Overall, this was a period where cell growth was
identical in infected and control cultures. However, between 24hr and 36hr post-infection
coincident with the arrest of cellular optical signals, alkalinity was halted and nutrient

consumption declined to very low background levels (Fig. 2A). After 36hr nitrate and
phosphate uptake were both arrested.

Figure 2. Dynamic of Alkalinity, nitrate, and phosphate in control and infected *E. huxleyi* cultures.
The measured concentration of control and EhV infected cultures; (A) Alkalinity. (B) Dissolved
nitrate. (C) Dissolved phosphate. The x-axes correspond to hours post-infection. Grey sections
represent the night period and dots indicate the mean values of each treatment
(Control/Infection), Error bars represent the standard error of replicates, n=3.

Particulate organic matter and stoichiometry dynamics

To determine the effect of viral infection on the elemental composition of coccolithophore
cells, POC, PON, and POP were quantified over the length of the infection (72hr). An
additional experiment was set to determine the longer-term variations in the elemental
composition of cell-derived lysates over 13 days post-infection (333 hr). Both
experiments were set up at the exact conditions and the output was nearly identical, thus
presented together (unless otherwise indicated). POC, PON, and POP per cell as
determined at the start of the experiment were 5.45 ± 0.15 pg cell^{-1}, 1.22 ± 0.06 pg cell^{-1}
and 0.14 ± 0.003 pg cell^{-1}, respectively. However, EhV infections result in very
heterogeneous cultures, including cells in different physiological states, dead cells (33),
TEP (38), and aggregates (42), all of which bear a particulate organic matter (POM)
signature. Thus, the particulate organic C, N, and P (POC, PON, and POP, respectively)
are presented per unit volume to illustrate the temporal variations in the ensemble of
particulate components produced during viral infected and control cultures and not per
cell basis (Fig.3). The stoichiometric ratios are presented in Fig. 4, focusing separately
on the infection period and post-infection for better visual details of variations during
infections.

Over the infection period, the variations in cell and viral densities (Fig. 3A) were broadly

identical to what was shown before in Fig. 1A. Briefly, the infected cultures increased in

density equally to controls over the first 24hr and then decline to background densities

$<10^3$ cell mL^{-1}. Free EhV rapidly peaked by the end of the second day (~3.50 x 10^8

virions mL^{-1}). Viral densities remained relatively stable until day 13. The control

populations multiplied exponentially over the first 120hr and then transitioned to

stationary growth phase at a density of about 4 x 10^6 cell mL^{-1}. In parallel, cohabiting

bacteria progressively accumulated in the cultures (Fig. 3B). The increase in the

bacterial population was higher in infected cultures doubling by t = 96hr (~0.40 x 10^8

bacteria mL^{-1}) the density from the controls. The bacterial density kept increasing,

reaching in infected cultures twice as higher densities (~0.9 x 10^8 bacteria mL^{-1}) than in

the controls by day 13.

In control cultures, POC, PON, and POP concentrations progressively increased during

the exponential phase as biomass accumulated (Fig. 3C–-E). Then PON and POP

stabilized as cultures entered stationary growth. In parallel, the molar stoichiometric

ratios remained relatively stable during exponential growth and then C:N and C:P both

increase during exponential growth (Fig. 4). By contrast, in the presence of EhV the

POC, PON, and POP concentrations increased during the first day of infection, but then

rapidly declined concomitant with cell demise. During this first infection period, variations

in the elemental stoichiometry relative to the controls were identified. Namely, the C:N

ratio was significantly higher (C:N > 8.0) in infected cultures at 48hr and 72 hr post-

infection than in the controls (C:N ~ 6.7) (Fig. 4A). By contrast, over the same period the

average N:P was lower in infected cultures (N:P 14.0–15.5) than in the control ones (N:P

15.7–16.2) (Fig. 4E). In the post-infection period, a slight increase of POC, PON, and

POP density was detected until the end (Fig. 4B, D and F). However, in parallel,

negative trends were detected for the C:N ratio (R^2 = 0.91) and C:P ratio (R^2 = 0.37),

indicating a progressive enrichment of N and P relative to carbon in organic biomass

post-infection.

Bacterial stoichiometry often deviates from the typical phytoplankton Redfield Ratio.

Average bacterial C:P is 77 and C:N is 4.9, while N:P is 17, similar to phytoplankton (43,

44). Thus, to test whether the observed shift in POM stoichiometry post-infection could

result from the recycling of algal to bacterial biomass and accumulation of high bacterial

titers, we calculated the contribution of bacteria over the course of our experiment. The

 results indicate that by the end of the infection on day 3 (72 hr), bacteria biomass
contributed already around 30%, at day 7 (168 hr) nearly 60%, and by day 13 nearly
80% of the total measured POM (Fig. 5).

Figure 3. Variations in particulate organic matter (POC, PON, POP) in control and EhV-infected
E. huxleyi cultures. Between 0 hr and 72 hr concerns the infection period. After 96 hr concerns
the post-infection period (see the main text for details). (A) Cell and viral densities. (B) Free
bacterial densities (C–E) POC, PON, and POP concentrations, respectively. For higher detail,
inset figures overlapping C, D and E highlight the variations in POC, PON, and POP during the
infection period. Error bars represent the standard error of replicates, n=3.

Figure 4. Variations in the elemental stoichiometric ratios (C:N, C:P, and N:P) in control and EhV-infected *E. huxleyi* cultures. The infection period (0-72 hr; A, C, and E) and post-infection period (from 96 hr to 333 hr; B, D, and F) are presented separately for better visualization. Respectively, the equations and R^2 values of the linear regressions during the post-infection period for C:N, C:P and N:P are: y = -0.16x + 7.98, R^2 = 0.91; y = −4.28x + 138.39, R^2 = 0.37; y = −0.26x + 17.65, R^2 = 0.12. Error bars represent the standard error of replicates, n=3. Statistical significance is indicated by the superscript letter a (p < 0.05) and b (p < 0.1).

228 **Figure 5.** Mathematical estimates of the contribution of bacteria to particulate organic matter over
229 the course of the infection and post-infection periods as defined in Fig. 3 and 4. The quadratic
230 fitting describes the rate of bacterial accumulation, which is a saturation function ($f_{bacteria}$ cannot
231 exceed 1; y = 0.054x + 0.054, R^2=0.85).

Discussion

Discussion

EhV halts calcification and nutrient uptake in infected host cells

A typical dynamic of *E. huxleyi* (RCC 1216) by the specific EhV 201 set at an MOI of 0.2
was detected (45, 46). Broadly, during the second day of infection, the daily oscillations
of Chl fluorescence and side scattering signals were both halted and gradually declined
in intensity (Fig. 1). These responses are consistent with the decline in the cellular
chlorophyll content, impaired photophysiology (47–50), and coccolith shedding triggered
during EhV infections (37). Then, after 36hr, the cell growth abruptly declined as a
consequence of EhV-mediated cell death (29, 30). A primary goal of this study, was to
test the impact of EhV infection on 1) calcification as detected by determining the draw-
down of alkalinity levels in the medium and 2) on the intake of the major macronutrients
(nitrate and phosphate). During the first day, calcification was comparable between
infected and control cultures. Alkalinity draw-down occurred during the light period, but
not through the night period (Fig. 2A). However, on the second day of infection,
calcification was completely arrested. This coincided with the onset of viral burst and the
overall shift in Chl fluorescence and side scatter signals in infected populations (Fig. 1).
Coccolith biosynthesis is a complex energy-demanding mechanism requiring abundant
lipid membranes for the coccolith vesicles and the secretion system that ultimately
results in large inorganic structures seemingly unnecessary to viral replication (16).
Thus, it is plausible that the viral-mediated breakdown of photosynthesis and the rapid
progression of EhV, requiring large fluxes of nucleotides, lipidic resources, and energy,

naturally results in the cessation of calcification and the incapability of coccolithophore

virocells to keep and restore the disintegrating coccosphere.

As for calcification, nitrate and phosphate were both consumed during the first day in

both control and infected cultures (Fig. 2B–C). However, in contrast to calcification,

nutrient consumption continued through the dark period. Dark uptake of nitrate is known

for *E. huxleyi* (41). Reports on the dark uptake of phosphate are less clear (41, 51).

However, a higher affinity for phosphate has been described during the night, likely due

to higher requirements for cell division (52). Nutrients acquired up to this stage through

the infection are naturally incorporated into EhV biosynthesis as has been recently

demonstrated in isotopic labeling experiments with carbon and N (53, 54). The reliance

of EhV on external phosphate has not been thus far assessed, but likely follows

analogous trends facilitated by the activity of a viral phosphate permease gene encoded

by some EhV strains (55, 56). During the second day nutrient consumption was arrested.

This occurred as side scatter and chlorophyll signals stopped oscillating, but prior to viral

burst and cell decline. This suggests that at this stage the virocell reached the final

stages of viral maturation, and that any further metabolic and energetic requirements

mostly relied on intracellular recycling of available macromolecules.

EhV-mediated shifts in biomass stoichiometry during infection and post-infection

In control cultures, as expected, POC, PON, and POP per unit volume increased over

time as biomass accumulated until the cultures reached the stationary phase and PON

and POP stabilized. During the exponential growth phase, the C:N, C:P and N:P

stoichiometry were fairly constant and close to the Redfield ratio through the exponential

growth phase, but once the cultures entered the stationary phase the C:N and C:P

rapidly increased (Fig. 4). This is likely caused by the accumulation of C-rich lipids in

nitrate and phosphate limited cells (57). By contrast, the concentration of POC, PON,

and POP declined during the infection period, which can be essentially attributed to the

viral-mediated cell lysis and loss of cellular contents to the medium as dissolved organic

matter. Over this first infection period, a consistent increase in C:N and a decline in N:P

ratios were detected over multiple time points, indicating both an enrichment of C and P

during infection (Fig. 4). The first stoichiometric shift likely results from the well-

documented intracellular accumulation of C-rich lipids in infected *E. huxleyi* cells (33, 34,

58). The second stoichiometric shift is less clear. A comparable shift in elemental

composition during viral lysis has been detected in the "brown tide" chrysophyte

Aureococcus anaphagefferens (59). In that system, carbon showed the largest

proportional release to the dissolved phase following viral lysis. It was argued that this

likely derived from the relative predominance of carbon in algal cytoplasm that could

easily leak out to the medium following cell decay and bacteria consumption. An

important loss of dissolved carbon in *E. huxleyi* during infection has also been also

recently estimated (60). By contrast, in *A. anaphagefferens*, N and P were both more

retained in the particulate fraction. Nitrogen was possibly rapidly transferred to the

bacterial community, which we cannot argue for *E. huxleyi* during infection here the

relative importance of N to the elemental stoichiometry declined, while P was suggested

to be retained in large structural components of the algae (59), as for example lipid

membranes. This may be the case in *E. huxleyi* as well, which have been shown to bear

a more important fraction of phospholipids than control cells (34). The contribution from

mRNA that likely builds up during viral propagation may also be important to P

enrichment during infection. In contrast, viral biomass likely did not influence the overall

organic matter proportions and ratios. For instance, using the similar *Paramecium*

bursaria Chlorella virus-1 (PBCV-1) with estimated C:N:P stoichiometry of 17:5:1 and

mass of 1 x 10^9 Da (61, 62), we estimate that during the second day, the contribution of

Ehv accumulated inside the infected cells prior to burst (~230 virions per cell) would

represent a nearly undetectable fraction (< 1 x 10^{-8} %) to the total particulate organic C,

N or P harvested. In any case, a future detailed examination of the elemental

redistribution between cell, viral, and bacterial compartments and the composition and

dynamic of macromolecules should enable us to decipher the underlying process of

stoichiometric shifts detected here. However, the fact that overall stoichiometric changes

occurred during day 2 and 3 (48 hr-72 hr), which was after nutrient consumption

stopped, further suggest the prevalence of the internal recycling of cellular and

macromolecular pools to fuel viral proliferation relative to *de novo* intake of resources.

Post-infection, when remaining coccolithophore cell abundance was very low (10^3–10^4

cell mL^{-1}) and the cultures were visibly full of large aggregates (see (42)), POM

concentrations progressively bounced back to nearly initial values. In parallel, the C:N

and C:P significantly declined, indicating an enrichment in N and P relative to C in the

remaining organic matter. We use a stoichiometric approach to estimate bacterial

accumulation, as we counted free bacteria and not POM-associated bacteria. Our

estimates (Fig. 5) indicate that these trends likely resulted from the rampant

accumulation of heterotrophic bacteria that have considerably higher demands in N and

P relative to C (C:N= 4.9; C:P=77.0) than phytoplankton (43, 44). Thus, heterotrophic

bacteria, triggered by viral-mediated lysis, accumulate at high rates in remaining cell-

aggregated. Such an accumulation, leads to a significant replacement of algal biomass

by bacterial biomass in the decaying particulate organic matter.

Potential ecological and biogeochemical implications of EhV infections

During *E. huxleyi* infections, large fractions of organic matter are lost to the dissolved

phase as detected by marked POC, PON, and POP concentration decline (Fig. 3). The

rapid release of cell contents during infections, as part of the so-called 'viral shunt' (9,

10), triggers the rapid rise of bacterial communities that can actively benefit from the

break-down and reuse of dissolved elements and lysates as it has been shown in other

host-viral systems (e.g., (59)). In parallel, stoichiometric variations were detected relative

to non-infected cells (Fig. 4). In the short term, during the course of infection and

population decline, an enrichment in C and P relative to N was detected, which can be

directly attributed to the EhV replication cycle and underlying metabolic remodeling of

virocells. In the longer term, decaying organic matter of *E. huxleyi* cells appears to be

progressively replaced to a large extent by bacterial biomass likely bound to aggregates

(42) produced during the earlier infection. The current view is that the formation of

coccolith-dense aggregates during *E. huxleyi* blooms infected by EhV can enhance the

carbon export efficiency, following the 'viral shuttle' model (39, 63, 64). However,

modifications in the elemental composition can impact the efficiency of carbon export.

Rapid export of recently infected populations with high carbon ratios may indeed benefit

the efficiency of carbon export. By contrast, slow export or resuspension of lysates in

more turbulent settings could enable time for the accumulation of bacteria and

concomitant enrichment of P relative to carbon in exported material, which then could in

fact attenuate the magnitude of the biological carbon pump by lowering the relative

contribution of carbon to exported biomass (6, 8, 65).

Taken together, it seems that the counteracting effect of viral shunt and viral shuttle

likely operate simultaneously during the interplay between *E. huxleyi* and EhV, as it is

most likely the case for most viral-host systems in the oceans. Yet, their relative

importance and efficiencies as well as comprehensive exploration in natural blooms

stoichiometry and export dynamics are required to validate our culture-based results and

determine the significance of *E. huxleyi* blooms in export production in the oceans. To

this, one must join the impact of grazing on this complex dynamic. In fact, *E. huxleyi*

blooms are targeted by microbial and zooplankton consumers (39, 57, 66–68) as well as

eukaryotic osmotrophs (60), all of which can contribute to the consumption of infected

cells and lysates and can further influence carbon export efficiencies. On this topic,

differences in regard to susceptibility to grazing between infected and non-infected cells

in *E. huxleyi* have been observed (69–71). In a recent study, lower ingestion rates by the

heterotrophic dinoflagellate *Oxyrrhis marina* in EhV-infected cells were determined.

However, despite lower ingestion, *O. marina* displayed higher growth rates, which

suggests higher nutritional values and/or great assimilation of infected cells (71). The

stoichiometric shift observed in our study indicates however that such nutritional value

can rapidly change depending on the state of the infection, which in turn could affect the

efficiency of the marine food web feeding during blooms.

Finally, as detailed above, calcification rapidly stalled during infection. Consequently, the

consumption of dissolved carbonate ions also should stop as it stops the draw-down of

seawater alkalinity as we observed experimentally. This highlights the antagonistic role

of EhV infection on the marine carbon counter pump, reducing the net increase of

dissolved CO_2 by demising growing populations with high rates of calcification and

consequently upholding the buffer capacity of the oceans (17).

Concluding remarks

In this study, we demonstrate that EhV infection arrests *E. huxleyi* calcification, nutrient

acquisition and caused ensuing shifts in the stoichiometry of organic matter. These

included both a short-term shift during active viral infection and a long-term shift driven

by cohabiting bacterial communities actively consuming cell-derived lysates and

aggregates. During natural blooms, the dynamic of EhV infections can be rapid, but also

heterogeneous between different blooming areas (39). Thus, one can expect the

stoichiometric seascape to be very variable. Nonetheless, such viral-mediated changes

in the nature of exported materials, likely influence the cycling of elemental in the photic

layer and the export efficiency of biomass to the deep ocean during large-scale

coccolithophore blooms. Further studies are required to validate our observation in

natural coccolithophore populations. However, our study demonstrates that assessing

the impact of viral infections on the elemental biogeochemistry of phytoplankton biomass

is crucial to determine their role and impact on the marine ecosystem.

Materials and Methods

Cell culturing and viral-infections

The xenic calcifying *E. huxleyi* strain RCC1216 (diploid) was used in all experiments.

Cell cultures were set in seawater-based K/2 (minus ammonium) medium (72), at 18°C,

13h:11h light: dark cycle, respectively, 80 µmol quanta m^{-2} s^{-1} irradiance using a warm

white LED system and gently air-bubbled through 0.2 µm filter. A 6 L stock culture was

grown for 3 days and then subdivided into six replicate glass bottles of 1 L each. Three

of the replicates were infected with EhV 201 at a Multiplicity of infection (MOI) of 0.2, and

three other replicates were used as non-infected controls. Sub-samples for cell, bacterial

and viral counts, nitrate, phosphate, and alkalinity, as well as POC, PON, and POP,

were collected at different time points with high frequency in the first 72hr post-infection.

An additional experiment run over 13 days was used to determine long term changes in

POC, PON, and POP. See the details for sampling and analyses below.

Cell, virus, and bacterial counts

E. huxleyi, bacteria, and EhV particles were enumerated by flow cytometry (Attune NxT,

Life Technology). Algal cells were identified based on chlorophyll fluorescence

(excitation 488 nm) and side scatter (46, 73). The mean side scatter and chlorophyll

autofluorescence values were used as indicators for the degree of cell calcification

(coccolith covering) and cell health, respectively. For viral and bacterial counts, samples

of 0.5 mL were fixed with 0.5% glutaraldehyde and counterstained with 1:10000 SYBR

Green (Invitrogen). Viruses and bacteria were identified by flow cytometry based on the

green fluorescence (excitation: 488 nm and emission: 530–737 nm) and side scatter

(73). The cell growth rate was determined as the natural logarithm of the cell density at 0

and 24hr divided by time. Viral burst size, was determined as the number of viruses

released per lysed cell (estimated as the ratio of the maximal number of viruses

produced to the maximum cell concentration reached by the specific host before cell

decrease) (24, 35).

Inorganic Nutrients and alkalinity

Duplicate samples of 10 mL were filtered through 0.2 µm filters (Sterivex, Millipore) and kept at 4°C until further analysis. Alkalinity was measured by acid-base titration following (74), via automatic titrator (Titroline 7000 Si analytics), and calculated using a gran plot with a precision lower than 20 mEq. Phosphate (as dissolved reactive phosphate) was measured by colorimetric spectroscopy using the Mo-Blue method (Cary 8454 UV-vis spectrometer, Agilent) with a precision of 0.1 mM and detection limit of 0.1 µM (75). Nitrate was measured by second-derivative spectroscopy (Cary 8454 UV-vis spectrometer, Agilent) with a precision of 0.1 mM and detection limit of 0.1 µM (76).

Particulate organic carbon, nitrogen, and phosphorus

For POC and PON, 20 mL duplicate samples were collected and filtered under low-pressure vacuum filtration (100 mm Hg) onto pre-combusted (450°C, 5hr) Whatman GF/F filters (25 mm diameter, 0.7 µm effective pore size). Then the filters were rinsed with 230 µL of 1M hydrochloric acid (HCl) to remove particulate inorganic carbon (PIC), oven-dried overnight at 65°C, and folded into tin capsules. Carbon and nitrogen content analyses were conducted in Continuous-flow Isotope Ratio Mass Spectrometry (CF-IRMS) using a Deltaplus XP mass spectrometer (ThermoScientific) coupled to an elemental analysis (EA) COSTECH 4010 (Costech Analytical) equipped with a zero blank autosampler. For every 36 samples or less, 10 secondary standards are used to correct the data using a linear correction. Caffeine, nannochlopsis, and Mueller Hinton broth were used as secondary standards calibrated from NIST primary standard. Analytical error (n=50) on measurement was 0.06‰ and 0.08‰ for 6N and 6C, respectively. System suitability prior to analysis was evaluated using the standard deviation of zero reference gas (N and carbon dioxide) over 10 measurements and the maximum acceptable variation was set to 0.06%. Certified reference material from Elemental Microanalysis treated as a sample is used as quality control. Quantification is based on external secondary standards with a calibration range of 0.025 mg to 0.250 mg and 0.040 mg to 0.400 mg for nitrogen and carbon respectively according to (Agnihotri et al., 2014). For POP, 50 mL duplicate samples were filtered under low-pressure vacuum filtration (100 mm Hg) onto pre-combusted (450°C, 5hr) Whatman GF/F filters (25 mm diameter, 0.7 µm effective pore size). They were then combusted at 450°C, for 5hr in a muffle furnace, cooled down, and immersed in 10 ml of 0.5N of HCl for 60 min at 90°C in

a heating block (77). Finally, 5 ml of supernatant was volumetrically subsampled into 15

ml centrifuge tubes for colorimetric spectroscopy using the Mo-Blue method (Cary 8454

UV-vis spectrometer, Agilent) as mentioned earlier for the analysis of inorganic P.

Mathematical estimation of bacterial contribution to particulate organic matter

Given the rampant accumulation of free bacteria during the infection experiments (see

Fig. 3), it is most likely that there was a concomitant accumulation of bacteria attached to

particulate organic matter (60). POM-bound bacteria were not measured. Thus, we

estimated the contribution of bacteria to total particulate organic matter over the course

of the experiment using a stoichiometric approach. These calculations were initialized

two days post-infection, which coincided with the time of viral burst and increment in

bacterial densities relative to control cultures. To calculate the fraction of bacteria out of

the total particulate organic matter we used the following equation:

$f_{\mathrm{phyto}} \cdot (\mathrm{C{:}N})_{\mathrm{phyto}} + f_{\mathrm{bacteria}} \cdot (\mathrm{C{:}N})_{\mathrm{bacteria}} = (\mathrm{C{:}N})_{\mathrm{POM}}$,

where $(\mathrm{C{:}N})_{\mathrm{phyto}}$, $(\mathrm{C{:}N})_{\mathrm{bacteria}}$, and $(\mathrm{C{:}N})_{\mathrm{POM}}$ are the C:N ratio in phytoplankton,

bacteria, and POM, respectively. f_{phyto}, and f_{bacteria} are the fraction of phytoplankton

and bacteria, respectively. Assuming that $f_{\mathrm{phyto}} + f_{\mathrm{bacteria}} = 1$, and rearranging, we get

the following equation:

$$f_{\mathrm{bacteria}} = \frac{(\mathrm{C{:}N})_{\mathrm{POM}} - (\mathrm{C{:}N})_{\mathrm{phyto}}}{(\mathrm{C{:}N})_{\mathrm{bacteria}} - (\mathrm{C{:}N})_{\mathrm{phyto}}}$$

Here, $(\mathrm{C{:}N})_{\mathrm{phyto}}$= 8.32 is the ratio measured 48 hr post-infection (see Fig. 3 in the

Results section). $(\mathrm{C{:}N})_{\mathrm{bacteria}}$= 4.94 is the average of elemental ratio of heterotrophic

bacteria (Table S1; (43, 44) The $(\mathrm{C{:}N})_{\mathrm{POM}}$ is based on the results of our experimental

findings (Fig. 5).

Acknowledgments

This study was supported by a scholarship from the Bester Award for Oceanography

from the Hebrew University of Jerusalem attributed to TD, the Fulbright and Yad Hanadiv

Fellowships attributed to SS, the Israel Science Foundation grant number 2921/20

attributed to MJF, the Israel Science Foundation grant number 2361/19 attributed to GA,

and by the Natural Sciences and Engineering Research Council of Canada Discovery

grant (RGPIN-2022-04305) and Fonds de Recherche du Québec Relève Professorale

grant (314248) attributed to AP. We thank Tania Rivlin for undertaking inorganic nutrient

measurements and Gil Koplovitz for help during experiments. We finally thank Maoz

Fine and Nir Keren for constructive feedback during the preparatory stages of the work.

Author Contributions: T.D., G.A. and M.J.F. conceptualized and designed the

research; T.D., G.A., A.P., S.S. and M.J.F. performed research; G.A., A.P. and M.J.F.

contributed analytic tools and funding; T.D., G.A., A.P., S.S. and M.J.F. analyzed data

and revised paper drafts; T.D. and M.J.F. wrote the paper.

References

1.   C. B. Field, M. J. Behrenfeld, J. T. Randerson, P. Falkowski, Primary Production

494   of the Biosphere: Integrating Terrestrial and Oceanic Components. *Science (80).*

495   **281**, 237–240 (1998).

496   2.   J. L. Sarmiento, M. Bender, Carbon biogeochemistry and climate change.

497   *Photosynth. Res.* **39**, 209–234 (1994).

498   3.   D. A. Hutchins, M. R. Mulholland, F. Fu, Nutrient cycles and marine microbes in a

499   CO2-enriched ocean. *Oceanography* **22**, 128–145 (2009).

500   4.   P. G. Falkowski, T. Fenchel, E. F. Delong, The microbial engines that drive

501   earth's biogeochemical cycles. *Science (80-.).* **320**, 1034–1039 (2008).

502   5.   Z. V. Finkel, *et al.*, Phytoplankton in a changing world: Cell size and elemental

503   stoichiometry. *J. Plankton Res.* **32**, 119–137 (2010).

504   6.   C. Deutsch, T. Weber, Nutrient Ratios as a Tracer and Driver of Ocean

505   Biogeochemistry. *Ann. Rev. Mar. Sci.* **4**, 113–141 (2012).

506   7.   S. Sharoni, I. Halevy, Nutrient ratios in marine particulate organic matter are

507   predicted by the population structure of well-adapted phytoplankton. *Sci. Adv.* **6**,

508   1–10 (2020).

509   8.   E. D. Galbraith, A. C. Martiny, A simple nutrient-dependence mechanism for

510   predicting the stoichiometry of marine ecosystems. *Proc. Natl. Acad. Sci. U. S. A.*

511   **112**, 8199–8204 (2015).

512   9.   J. A. Fuhrman, Marine viruses and their biogeochemical and ecological effects.

513   *Nature* **399**, 541–548 (1999).

10. S. W. Wilhelm, C. A. Suttle, Viruses and Nutrient Cycles in the Sea. *Bioscience*
49, 781–788 (1999).

11. K. D. A. Mojica, J. Huisman, S. W. Wilhelm, C. P. D. Brussaard, Latitudinal
variation in virus-induced mortality of phytoplankton across the North Atlantic
Ocean. *ISME J.* **10**, 500–513 (2016).

12. A. E. Zimmerman, *et al.*, Metabolic and biogeochemical consequences of viral
infection in aquatic ecosystems. *Nat. Rev. Microbiol.* **18**, 21–34 (2020).

13. A. J. Poulton, T. R. Adey, W. M. Balch, P. M. Holligan, Relating coccolithophore
calcification rates to phytoplankton community dynamics: Regional differences
and implications for carbon export. *Deep. Res. Part II Top. Stud. Oceanogr.* **54**,
538–557 (2007).

14. A. J. Poulton, *et al.*, The 2008 *Emiliania huxleyi* bloom along the Patagonian
Shelf: Ecology, biogeochemistry, and cellular calcification. *Global Biogeochem.*
Cycles **27**, 1023–1033 (2013).

15. J. R. Young, S. A. Davis, P. R. Bown, S. Mann, Coccolith ultrastructure and
biomineralisation. *J. Struct. Biol.* **126**, 195–215 (1999).

16. F. M. Monteiro, *et al.*, Why marine phytoplankton calcify. *Sci. Adv.* **2** (2016).

17. B. Rost, U. Riebesell, Coccolithophores and the biological pump: responses to
environmental changes. *Coccolithophores*, 99–125 (2004).

18. C. Klaas, D. E. Archer, Association of sinking organic matter with various types of
mineral ballast in the deep sea: Implications for the rain ratio. *Global*
Biogeochem. Cycles **16**, 63-1-63–14 (2002).

19. J. L. Sarmiento, N. Gruber, Sinks for Anthropogenic Carbon. *Phys. Today* **55**,
30–36 (2002).

20. P. M. Holligan, *et al.*, A biogeochemical study of the coccolithophore, *Emiliania*
huxleyi, in the North Atlantic. *Global Biogeochem. Cycles* **7**, 879–900 (1993).

21. T. Tyrrell, A. Merico, "*Emiliania huxleyi*: bloom observations and the conditions
that induce them" in *Coccolithophores*, (Springer Berlin Heidelberg, 2004), pp.
75–97.

22.   A. Winter, J. Henderiks, L. Beaufort, R. E. M. Rickaby, C. W. Brown, Poleward
expansion of the coccolithophore *Emiliania huxleyi*. *J. Plankton Res.* **36**, 316–325
(2014).

23.   E. J. Lessard, A. Merico, T. Tyrrell, Nitrate:phosphate ratios and *Emiliania huxleyi*
blooms. *Limnol. Oceanogr.* **50**, 1020–1024 (2005).

24.   G. Bratbak, J. K. Egge, M. Heldal, Viral mortality of the marine alga *Emiliania*
huxleyi (Haptophyceae) and termination of algal blooms. *Mar. Ecol. Prog. Ser.*
93, 39–48 (1993).

25.   C. P. D. Brussaard, R. S. Kempers, A. J. Kop, R. Riegman, M. Heldal, Virus-like
particles in a summer bloom of *Emiliania huxleyi* in the North Sea. *Aquat. Microb.*
Ecol. **10**, 105–113 (1996).

26.   W. H. Wilson, *et al.*, Isolation of viruses responsible for the demise of an
Emiliania huxleyi bloom in the English Channel. *J. Mar. Biol. Assoc. United*
Kingdom **82**, 369–377 (2002).

27.   Y. Lehahn, *et al.*, Decoupling physical from biological processes to assess the
impact of viruses on a mesoscale algal bloom. *Curr. Biol.* **24**, 2041–2046 (2014).

28.   D. C. Schroeder, J. Oke, G. Malin, W. H. Wilson, Coccolithovirus
(Phycodnaviridae): Characterisation of a new large dsDNA algal virus that infects
Emiliania huxleyi. *Arch. Virol.* **147**, 1685–1698 (2002).

29.   K. D. Bidle, L. Haramaty, J. Barcelos, P. Falkowski, Viral activation and
recruitment of metacaspases in the unicellular coccolithophore, *Emiliania huxleyi*.
104 (2007).

30.   A. Vardi, *et al.*, Viral Glycosphingolipids Induce Lytic Infection and Cell Death in
Marine Phytoplankton. *Science (80-.).* **326**, 861–865 (2009).

31.   U. Sheyn, S. Rosenwasser, S. Ben-Dor, Z. Porat, A. Vardi, Modulation of host
ROS metabolism is essential for viral infection of a bloom-forming
coccolithophore in the ocean. *ISME J.* **10**, 1742–1754 (2016).

32.   D. Schatz, *et al.*, Hijacking of an autophagy-like process is critical for the life
cycle of a DNA virus infecting oceanic algal blooms. *New Phytol.* **204**, 854–863
(2014).

33.   S. Rosenwasser, *et al.*, Rewiring host lipid metabolism by large viruses
determines the fate of Emiliania huxleyi, a bloom-forming alga in the ocean. *Plant*
Cell **26**, 2689–2707 (2014).

34. S. Malitsky, *et al.*, Viral infection of the marine alga *Emiliania huxleyi* triggers lipidome remodeling and induces the production of highly saturated triacylglycerol. *New Phytol.* **210**, 88–96 (2016).

35. S. Jacquet, *et al.*, Flow cytometric analysis of an *Emiliana huxleyi* bloom terminated by viral infection. *Aquat. Microb. Ecol.* **27**, 111–124 (2002).

36. M. Trainic, *et al.*, Infection Dynamics of a Bloom-Forming Alga and Its Virus Determine Airborne Coccolith Emission from Seawater. *iScience* **6**, 327–335 (2018).

37. C. T. Johns, *et al.*, The mutual interplay between calcification and coccolithovirus infection. *Environ. Microbiol.* **21**, 1896–1915 (2019).

38. A. Vardi, *et al.*, Host-virus dynamics and subcellular controls of cell fate in a natural coccolithophore population. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 19327–19332 (2012).

39. C. P. Laber, *et al.*, Coccolithovirus facilitation of carbon export in the North Atlantic. *Nat. Microbiol.* **3**, 537–547 (2018).

40. S. Rosenwasser, C. Ziv, S. G. van Creveld, A. Vardi, Virocell Metabolism: Metabolic Innovations During Host–Virus Interactions in the Ocean. *Trends Microbiol.* **24**, 821–832 (2016).

41. M. N. Müller, A. N. Antia, J. LaRoche, Influence of cell cycle phase on calcification in the coccolithophore *Emiliania huxleyi*. *Limnol. Oceanogr.* **53**, 506–512 (2008).

42. F. Vincent, *et al.*, Viral infection switches the balance between bacterial and eukaryotic recyclers of organic matter during algal blooms. *bioRxiv*, 2021.10.25.465659 (2021).

43. A. E. Zimmerman, S. D. Allison, A. C. Martiny, Phylogenetic constraints on elemental stoichiometry and resource allocation in heterotrophic marine bacteria. *Environ. Microbiol.* **16**, 1398–1410 (2014).

44. A. E. White, S. J. Giovannoni, Y. Zhao, K. Vergin, C. A. Carlson, Elemental content and stoichiometry of SAR11 chemoheterotrophic marine bacteria. *Limnol. Oceanogr. Lett.* **4**, 44–51 (2019).

45. M. Frada, I. Probert, M. J. Allen, W. H. Wilson, C. De Vargas, The "Cheshire Cat" escape strategy of the coccolithophore *Emiliania huxleyi* in response to viral infection. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 15944–15949 (2008).

609   46.   M. J. Frada, *et al.*, Morphological switch to a resistant subpopulation in response
610   to viral infection in the bloom-forming coccolithophore *Emiliania huxleyi*. *PLOS*
611   *Pathog.* **13**, e1006775 (2017).

612   47.   C. Evans, G. Malin, G. P. Mills, W. H. Wilson, Viral infection of *Emiliania huxleyi*
613   (Prymnesiophyceae) leads to elevated production of reactive oxygen species. *J.*
614   *Phycol.* **42**, 1040–1047 (2006).

615   48.   J. Kegel, *et al.*, Pilot study of an EST approach of the coccolithophorid *Emiliania*
616   *huxleyi* during a virus infection. *Gene* **406**, 209–216 (2007).

617   49.   S. A. Kimmance, M. J. Allen, A. Pagarete, J. M. Martínez, W. H. Wilson,
618   Reduction in photosystem II efficiency during a virus-controlled *Emiliania huxleyi*
619   bloom. **495**, 65–76 (2014).

620   50.   C. A. Llewellyn, *et al.*, The response of carotenoids and chlorophylls during virus
621   infection of *Emiliania huxleyi* (Prymnesiophyceae). *J. Exp. Mar. Bio. Ecol.* **344**,
622   101–112 (2007).

623   51.   R. Riegman, W. Stolte, A. A. M. Noordeloos, D. Slezak, Nutrient uptake and
624   alkaline   phosphatase   (ec   3:1:3:1)   activity   of   emiliania   huxleyi
625   (Prymnesiophyceae)
626   during growth under n and p limitation in continuous cultures. *J. Phycol.* **36**, 87–
627   96 (2000).

628   52.   R. Geider, J. La Roche, Redfield revisited: variability of C:N:P in marine
629   microalgae and its biochemical basis. *Eur. J. Phycol.* **37**, 1–17 (2002).

630   53.   A. L. Pasulka, *et al.*, Interrogating marine virus-host interactions and elemental
631   transfer with BONCAT and nanoSIMS-based methods. *Environ. Microbiol.* **20**,
632   671–692 (2018).

633   54.   E. Yakubovskaya, T. Zaliznyak, J. M. Martínez, G. T. Taylor, Raman
634   Microspectroscopy Goes Viral: Infection Dynamics in the Cosmopolitan
635   Microalga, *Emiliania huxleyi*. *Front. Microbiol.* **12** (2021).

636   55.   M. J. Allen, D. C. Schroeder, W. H. Wilson, Preliminary characterisation of repeat
637   families in the genome of EhV-86, a giant algal virus that infects the marine
638   microalga *Emiliania huxleyi*. *Arch. Virol.* **151**, 525–535 (2006).

639   56.   W. H. Wilson, *et al.*, Complete genome sequence and lytic phase transcription
640   profile of a Coccolithovirus. *Science (80-.).* **309**, 1090–1092 (2005).

57. M. J. Frada, *et al.*, Zooplankton may serve as transmission vectors for viruses infecting algal blooms in the ocean. *Curr. Biol.* **24**, 2592–2597 (2014).

58. A. Pagarete, *et al.*, Dip in the gene pool: Metagenomic survey of natural coccolithovirus communities. *Virology* **466**–**467**, 129–137 (2014).

59. C. J. Gobler, D. A. Hutchins, N. S. Fisher, E. M. Cosper, S. A. Sañudo-Wilhelmy, Release and bioavailability of C, N, P, Se, and Fe following viral lysis of a marine chrysophyte. *Limnol. Oceanogr.* **42**, 1492–1504 (1997).

60. F. Vincent, *et al.*, Viral infection switches the balance between bacterial and eukaryotic recyclers of organic matter during coccolithophore blooms. *Nat. Commun.* **14** (2023).

61. J. L. Clasen, J. J. Elser, The effect of host *Chlorella* NC64A carbon:phosphorus ratio on the production of *Paramecium bursaria Chlorella* Virus-1. *Freshw. Biol.* **52**, 112–122 (2007).

62. L. F. Jover, T. C. Effler, A. Buchan, S. W. Wilhelm, J. S. Weitz, The elemental composition of virus particles: Implications for marine biogeochemical cycles. *Nat. Rev. Microbiol.* **12**, 519–528 (2014).

63. M. G. Weinbauer, Ecology of prokaryotic viruses. *FEMS Microbiol. Rev.* **28**, 127–181 (2004).

64. M. B. Sullivan, J. S. Weitz, S. Wilhelm, Viral ecology comes of age. *Environ. Microbiol. Rep.* **9**, 33–35 (2017).

65. Z. V. Finkel, *et al.*, Phytoplankton in a changing world: cell size and elemental stoichiometry. *J. Plankton Res.* **32**, 119–137 (2010).

66. J. C. Nejstgaard, H. J. Witte, P. van der Wal, A. Jacobsen, Copepod grazing during a mesocosm study of an *Emiliania huxleyi* (Prymnesiophyceae) bloom. *Sarsia* **79**, 369–377 (1994).

67. M. B. Olson, S. L. Strom, Phytoplankton growth, microzooplankton herbivory and community structure in the southeast Bering Sea: insight into the formation and temporal
persistence of an *Emiliania huxleyi* bloom. *Deep Sea Res. Part II Top. Stud. Oceanogr.* **49**, 5969–5990 (2002).

68. K. M. J. Mayers, *et al.*, Growth and mortality of coccolithophores during spring in a temperate Shelf Sea (Celtic Sea, April 2015). *Prog. Oceanogr.* **177**, 101928 (2019).

69.   C. Evans, W. H. Wilson, Preferential grazing of *Oxyrrhis marina* on virus-infected
Emiliania huxleyi. Limnol. Oceanogr. **53**, 2035–2040 (2008).

70.   A. I. Vermont, *et al.*, Virus infection of *Emiliania huxleyi* deters grazing by the
copepod *Acartia tonsa. J. Plankton Res.* **38**, 1194–1205 (2016).

71.   A. G. Goode, D. M. Fields, S. D. Archer, J. M. Martínez, Physiological responses
of oxyrrhis marina to a diet of virally infected *Emiliania huxleyi. PeerJ* **2019**, 1–25
(2019).

72.   M. D. Keller, R. C. Selvin, W. Claus, R. R. L. Guillard, Media for the culture of
oceanic ultraphytoplankton 1,2. *J. Phycol.* **23**, 633–638 (1987).

73.   D. Marie, C. P. D. Brussaard, R. Thyrhaug, G. Bratbak, D. Vaulot, Enumeration
of marine viruses in culture and natural samples by flow cytometry. *Appl. Environ.*
Microbiol. **65**, 45–52 (1999).

74.   M. Neder, R. Saar, A. Malik, G. Antler, T. Mass, New Insights on the Diurnal
Mechanism of Calcification in the Stony Coral, Stylophora pistillata. *Front. Mar.*
Sci. **8**, 1–15 (2022).

75.   W. I. M. Holman, A new technique for the determination of phosphorus by the
molybdenum blue method. *Biochem. J.* **37**, 256–259 (1943).

76.   W. G. Crumpton, T. M. Isenhart, P. D. Mitchell, Nitrate and organic N analyses
with second-derivative spectroscopy. *Limnol. Oceanogr.* **37**, 907–913 (1992).

77.   D. M. Karl, B. D. Tilbrook, G. Tien, Seasonal coupling of organic matter
production and particle flux in the western Bransfield Strait, Antartica. *Deep Sea*
Res. Part A. Oceanogr. Res. Pap. **38**, 1097–1126 (1991).