Resolving ancient gene transfers clarifies the early co-evolution of eukaryotes and giant

viruses

Sangita Karki [1], Frank O. Aylward [1,2*]

[1] Department of Biological Sciences, Virginia Tech, Blacksburg, United States of America

[2] Center for Emerging, Zoonotic, and Arthropod-borne Pathogens, Virginia Tech, Blacksburg,

United States of America

*Email for correspondence: faylward@vt.edu

Abstract

Members of the phylum *Nucleocytoviricota*, also called "giant viruses" due to their large physical dimensions and genome lengths, are a diverse group of dsDNA viruses that infect a wide range of eukaryotic hosts. Nucleocytoviruses likely evolved from smaller viruses, but the timing of their emergence and its relationship to the early evolution of eukaryotes remains unclear. Recent work has shown that the genomes of nucleocytoviruses often encode Eukaryotic Signature Proteins (ESPs) - including histones, vesicular trafficking factors, cytoskeletal components, and elements of RNA and DNA processing - that occur only rarely outside of eukaryotes. To investigate patterns of gene exchange between viruses and eukaryotes and possibly shed light on the early evolution of both, we examined the occurrence of viral-encoded ESPs (vESPs) and performed a comprehensive phylogenetic reconstruction on a subset that are widespread in nucleocytoviruses. Our results demonstrate that vESPs involved in cytoskeletal structure, ubiquitin system, and vesicular trafficking were acquired multiple times independently by nucleocytoviruses at different timepoints after the emergence of the eukaryotic supergroups. In contrast, vESPs involved in DNA and RNA processing are placed deep in their respective phylogenies, indicative of ancient gene exchange between nucleocytoviruses and eukaryotes. Examination of vESPs that could be rooted in archaea revealed that nucleocytoviruses likely acquired some of these genes prior to the emergence of the last eukaryotic common ancestor (LECA). Importantly, our findings also suggest that the eukaryotic delta DNA polymerase was acquired from nucleocytoviruses sometime during eukaryogenesis, underscoring the importance of viruses for early eukaryotic evolution. Collectively, these results suggest that gene exchange between nucleocytoviruses and eukaryotes played important roles in the evolution of both prior to the emergence of LECA.

45   **Keywords:** giant viruses, *Nucleocytovoricota*, eukaryotic signature proteins, viral origins, viral

46   diversity

Introduction

The discovery of giant viruses, also known as nucleocytoplasmic large DNA viruses (NCLDVs), has dramatically expanded our understanding of the limits of viral size and genome complexity. These viruses belong to the phylum *Nucleocytoviricota* and comprise 6 orders, 11 families, and a rapidly-expanding number of family-level lineages [1]. Although they are found in diverse environments, they are particularly abundant in the oceans [2–5] and infect a wide range of hosts, ranging from the smallest known free-living eukaryote [6] to metazoans [7]. Members of families such as *Poxviridae*, *Asfarviridae*, and *Iridoviridae* generally include well-known vertebrate pathogens [8–10] while members of the *Mimiviridae, Phycodnaviridae,* and *Marseilleviridae* primarily infect protists and algae [11–13]. Phylogenomic studies across these groups have provided evidence that they have a shared evolutionary origin and potentially evolved from a tectivirus-like ancestor [14–16]. Gene acquisition from their hosts coupled with gene duplication likely contributed to the subsequent expansion of genome size and complexity in this lineage [14,17,18].

A resonant theme in recent years has been the discovery of numerous "cell-like" genes in nucleocytoviruses that are common in cellular life but rare or absent from other viral lineages. These include genes involved in translation [19,20], transcription [21], DNA replication and repair pathways [22–24], diverse metabolic pathways including the TCA cycle and glycolysis [25], cytoskeletal structure [26–28], membrane trafficking [29], and ubiquitin signaling [30]. In some cases viruses have acquired these genes from their hosts recently, and viral homologs form well-defined clades within broader eukaryotic lineages. This appears to be the case for ammonia transporters, some rhodopsins, and sphingolipid metabolism genes [31–33]. In other cases, such

as histones and DNA polymerase subunits, viral genes are the result of ancient gene transfer events

and their evolutionary relationship to cellular homologs is less clear [34,35]. Recent studies have

shown that nucleocytoviruses often endogenize their genomes into those of their host [25,36,37]

demonstrating that some of these genes may have originated in viruses and subsequently

transferred to eukaryotes. Phylogenetic trees of multi-subunit RNA polymerase [21], DNA

topoisomerase IIA [38], and actin [28] are examples of ancient virus-to-eukaryote gene transfers

that have been proposed, although the direction of these transfer events is difficult to definitively

ascertain. Regardless of the direction of gene transfer, these studies have begun to elucidate ancient

patterns of gene transfer between viruses and eukaryotes that underscore the long co-evolutionary

history of eukaryotes and nucleocytoviruses. This dynamic interplay between viruses and their

hosts has shaped the early evolution of both and likely began prior to the emergence of the Last

Eukaryotic Common Ancestor (LECA) [21].

The evolution of eukaryotes represents a major evolutionary transition in the evolution of life on

Earth [39] and yet the details of this process remain a riddle [40]. Many complex features of

eukaryotes emerged in the stem eukaryotic lineage prior to the emergence of LECA, but the order

in which these traits emerged is largely unknown. Given the pre-LECA origin of

nucleocytoviruses, it has been proposed that the co-evolution of these viruses and their hosts

played a role in eukaryogenesis [41]. Interestingly, many nucleocytoviruses encode Eukaryotic

Signature Proteins (ESPs) that are involved in central processes in eukaryotic cells, such as DNA

and RNA processing, vesicular trafficking, cytoskeletal structure, and ubiquitin signaling. ESPs

are common in eukaryotes but typically rare or absent in bacteria and archaea, and examination of

their evolution therefore can provide key clues into the emergence of eukaryotes [42,43]. To

provide deeper insight into the early co-evolution of nucleocytoviruses and eukaryotes we

performed an in-depth survey of viral-encoded ESPs (vESPs) and performed a comprehensive

phylogenetic reconstruction of their evolutionary histories. Our results provide key insights into

the origin of nucleocytoviruses and the extent of virus-eukaryote gene transfer that occurred prior

to the emergence of LECA.

Results

Occurrence of vESPs

We identified a set of 763 Pfam domains that are present in greater than 95 percent of the

eukaryotic genomes surveyed, and 500 of these could be identified in at least one nucleocytovirus

genome. These Pfams could be clustered into six groups depending on their occurrence across

bacteria and archaea (S1 Data). Clusters 1 and 2 are broadly distributed across all cellular lineages

(pan-cellular group), clusters 3 and 4 are found primarily in eukaryotes (true vESPs), and clusters

5 and 6 are found primarily in eukaryotes and archaea (archaea-vESPs) (Figure 1A). Many vESPs

were only found in a small number of viral genomes and are not suitable for phylogenetic

reconstruction; we therefore examined the prevalence of vESPs from different clusters across the

Nucleocytoviricota to identify those that are broadly represented in this viral phylum (Figure 1B,

1C). To identify ESP that would be most likely to shed light on the early evolution of both

nucleocytoviruses and eukaryotes, we searched both Pfams that were represented in greater than

ten nucleocytovirus genomes and were true ESPs (i.e., they belonged to clusters 3 or 4). A

preponderance of these proteins are involved in vesicular trafficking, ubiquitin signaling,

cytoskeletal structure, mRNA biogenesis, and DNA processing, and we therefore focused our

analysis on protein families belonging to these functional categories. We focused our analysis on

true vESPs, but in some cases we also included archaeal vESPs because of the opportunity to root

these trees in the archaea.

Given the divergent nature of many vESPs that we examined here, we used several quality-control

filters to examine the reliability of the trees that we generated. In order to evaluate the phylogenetic

strength of the vESP trees we constructed, we calculated the Tree Certainty (TC) matric for each

tree. This metric varies between zero and one, with higher values having fewer contrasting

bipartitions and a more reliable topology [44]. Longer genes with a stronger phylogenetic signal

typically exhibit larger TC values [45], and here we report only trees with TC > 0.5. All TC scores

are provided for each tree and are also provided in S2 Data. In addition, for all vESPs we

constructed trees using both the best fit site-homogeneous substitution model and the best fit

complex model (i.e. site heterogeneous). The topologies of both trees were examined, and results

are only reported when they are largely consistent (see Methods for details and S2 and S3 Text).

Fig 1. Clustering and functional annotation of ESPs. A) Unsupervised clustering of 763

eukaryotic proteins revealed six different clusters based on their prevalence in different groups. B)

Functional categories of protein domains present in greater than 5 nucleocytovirus genomes

irrespective of clusters. C) Functional annotation of protein domains belonging to different

clusters.

Evolution of vESPs involved in ubiquitin signaling, cytoskeletal dynamics, and vesicular trafficking

Phylogenetic analysis of vESPs involved in ubiquitin signaling, vesicular trafficking, and cytoskeletal structure revealed topologies in which vESPs were nested within broader clades of eukaryotic proteins (Fig 2), clearly indicative of eukaryote-to-virus gene transfer events. It is not possible to root these trees given the lack of homologs in other cellular domains, but the placement of almost all vESPs within multiple distinct lineages nested within eukaryotic clades is strong evidence that viruses acquired these vESPs after the establishment of the main eukaryotic supergroups. Moreover, the occurrence of multiple distinct vESP clades demonstrates that viruses acquired most of these proteins multiple times independently throughout their evolution. It is also notable that the breadth of the vESP clades vary between trees. For example, whereas the SNARE and synaptobrevin vESPs are scattered broadly among eukaryotic homologs (Fig 2A), the Ub-activating enzyme and Ub-protease form some large clades (Fig 2B). Collectively this suggests that the vESPs were acquired by viruses over a range of different timescales, with shallow vESP clades representing recent acquisitions and deeper clades representing older gene transfer events.

A. Signal Transduction, Membrane Trafficking, and Transport

TC:0.83 — PI-3-kinase [PF00613]
TC:0.68 — SNARE [PF05739]
TC:0.67 — Synaptobrevin [PF00957]
TC:0.81 — Sec1 [PF00995]

Groups
Bacteria
Eukaryote
Giant Virus

B. Ubiquitin (Ub)/ Protein Degradation System

TC:0.73 — Ulp1 protease [PF02902]
TC:0.84 — Ub activating enzyme [PF10585]
TC:0.68 — Ub hydrolase [PF13423]
TC:0.7 — U-box domain [PF04564]

C. Cytoskeleton proteins

TC:0.8 — Dynamin central region [PF01031]
TC:0.81 — Myosin head [PF00063]
TC:0.7 — Microtubule binding [PF16796]
TC:0.73 — Tubulin tyrosine ligase [PF03133]

Fig 2. Phylogenetic trees for vESPs involved in (A) membrane trafficking, (B) ubiquitin system, and (C) cytoskeletal structure.

Our findings for a relatively recent viral acquisition membrane trafficking proteins is consistent with a phylogenetic study where a highly divergent Mimivirus Rab GTPase, a crucial regulator of membrane trafficking in eukaryotes, was reported to be acquired from unicellular eukaryotes [46]. Moreover, a recent study that confirmed the activity of some viral vesicular trafficking

components also suggested that these proteins had been acquired multiple times by viruses [29].

Several phylogenetic studies of viral cytoskeleton homologs also suggested that the viral myosin

and kinesin homologs form distinct clades nested within broader eukaryotic lineages [26,27].

Interestingly, a recent study of viral actin (i.e. viractin) homologs found evidence that viruses

encoded this protein prior to its origin in modern eukaryotes [28], which stands in contrast to the

general patterns we observed for cytoskeletal proteins. Viractin is relatively rare in

nucleocytoviruses genomes and was therefore excluded from our analysis, but this previous study

stands as a reminder that the relatively recent viral acquisition of eukaryotic cytoskeletal homologs

may not hold for all protein families. Indeed, exceptions for these late branching clades were

notable for some ubiquitin related proteins and a protein involved in membrane trafficking and

biogenesis (ATG8) (Fig 1 in S1 Text).

Phylogeny on DNA and RNA processing proteins belonging to true vESPs

The vESPs involved in mRNA and DNA processing formed deep-branching lineages in their

respective trees, contrasting sharply with vESPs involved in other processes (Fig 3). This suggests

that these proteins have a much more ancient evolutionary origin in the *Nucleocytoviricota*. Once

again, it is not possible to root these trees due to the lack of homologs in other cellular domains,

and it is therefore not possible to confidently determine the precise timing and direction of transfer

that led to these topologies. Nonetheless, the unrooted trees still exhibit a stark distinction to those

of vESPs involved in vesicular trafficking, cytoskeletal structure, and ubiquitin signaling,

indicating that vESPs involved in mRNA and DNA processing have an ancient origin in the

Nucleocytoviricota that can potentially be used to date the origin of this phylum.

DNA and RNA processing proteins

XRN exonuclease [PF03159] RNA pol Rpb1, domain 6 [PF04992] RNA pol Rpb5, N- terminal [PF03871] Translation initiation 4E [PF01652]

mRNA guanylyltransferase [PF01331] DNA primase domain [PF16898] Guanine-N(7)-methyltransferase [PF03291]

Groups
Bacteria
Archaea
Asgard archaea
Eukaryote
Giant Virus

Fig 3. Phylogenetic trees for true vESPs involved in DNA and RNA processing.

Specifically, DNA primase, XRN endonuclease, different RNA polymerase subunits, enzymes of mRNA capping complex (mRNA cap guanylyltransferase, mRNA cap methyltransferase), and eukaryotic initiation factor 4E, all showed deep branching topology suggestive an ancient gene transfer between nucleocytoviruses and eukaryotes. Indeed, proteins belonging to the mRNA capping complex showed deep branching placement suggestive of ancient gene transfers between nucleocytoviruses and its hosts (Fig 3). Capping and methylation mechanisms are crucial steps in mRNA processing and initiation of protein synthesis for eukaryotes as well as giant viruses. A previous phylogenetic study on mRNA capping enzymes of nucleocytoviruses suggested that these two mRNA capping domains were present in the ancestor of nucleocytoviruses [47].

Translation initiation factor 4E (eIF4e) is involved in directing ribosomes to bind the cap structure

of mRNA and therefore acts as a bridge between RNA processing and translation. Compared to

other proteins involved in DNA or RNA processing, the tree of eIF4e provided strong evidence

for multiple transfers of this protein from eukaryotes to viruses (Fig 3). Similar findings were

reported in a previous study that broadly examined translation-associated genes in giant viruses

and found that viral proteins were typically acquired multiple times independently from different

eukaryotic lineages [18].

Overall, phylogenetic analysis of DNA and RNA processing protein domains belonging to cluster

3 and 4 (true vESPs) shows that these vESPs are mostly ancient. Although multiple viral clades

were sometimes present in these trees, they were often placed deep in the tree in ancient nodes that

are difficult to resolve. We therefore consider it more likely that this is a product of difficulties

resolving deep nodes in a phylogeny rather than multiple gene transfer events. It is likely that the

acquisition of core machinery for DNA and mRNA synthesis were critical for the early

establishment of the *Nucleocytoviricota*, and it is therefore not surprising that these are the most

ancient protein families in this phylum.

Rooting gene transfer events with vESPs found in Archaea (Archaea-vESPs)

In order to infer direction of transfer, we expanded the analysis to include genes from the Archaea-

vESP protein families because these can be rooted with the Archaea as an outgroup (Clusters 5

and 6 in Fig 1A). This is critical because it allows for a more precise evaluation of when giant

viruses acquired these genes relative to the Last Eukaryotic Common Ancestor (LECA).

Moreover, in some cases it has been proposed that early eukaryotes acquired certain genes from

viruses prior to their diversification [21,28,38], and basal-branching placement of vESPs would

provide evidence that this is a possibility. Although Archaea-vESPs are not true ESPs due to their

presence in Archaea, the phylogenetic proximity of Asgard Archaea to early eukaryotes leaves

open the possibility of gene transfer between some archaeal lineages and giant viruses during the

early emergence of the *Nucleocytoviricota* [48]. Moreover, the Archaea-vESPs were enriched in

genes involved in DNA and RNA processing, which provides the opportunity to examine how

giant viruses acquired genes involved in these processes.

Similar to the phylogenies with the other vESPs, Archaea-vESPs involved in DNA and RNA

processing exhibited deep branching viral clades indicative of ancient gene transfers with

eukaryotes. Among the nine Archaea-vESPs we analyzed in detail, our rooted trees revealed basal-

branching eukaryotic lineages in seven and basal-branching nucleocytovirus clades in two (Fig 4).

The phylogeny of the RNA polymerase subunit 5- C terminal, which is common in Archaea and

eukaryotes but absent in bacteria, revealed an early-branching viral clade. This suggests that this

protein family already existed in the *Nucleocytoviricota* prior to the emergence of LECA,

consistent with a previous study that focused on the two major RNA polymerase subunits and

found that both of these protein families were found in the *Nucleocytoviricota* prior to the

diversification of modern eukaryotes [21]. Similarly, a phylogeny of the DNA sliding clamp also

revealed a deep branching viral clade indicative of an ancient presence in the *Nucleocytoviricota,*

although several deep nodes in this tree had low bootstrap support (<80, Fig 9 in S2 and S3 Text)

and must be interpreted with caution. For the other seven Archaea-vESPs, we recovered clear

basal-branching eukaryotic clades with strong bootstrap support (S2 and S3 Text).

Fig 4. Phylogenetic trees for Archaea-vESPs involved in DNA and RNA processing.

Some protein families, such as multimeric RNA polymerase subunits and family B polymerases, diversified in eukaryotes through a complex process of duplication and divergence that took place prior to the emergence of LECA. It therefore remains a possibility that giant viruses acquired these

257
258
259
260

proteins and then transferred them back to eukaryotes (i.e. reciprocal gene transfer), thereby

participating in early diversification of these protein families. This has already been suggested for

the two largest subunits of multimeric RNA polymerase, which are found in three copies in all

modern eukaryotes but only a single copy in bacteria, archaea, and giant viruses. A previous study

found that eukaryotic RNAP subunits form three distinct clades (I, II, and III), consistent with their

early diversification in eukaryotes, and that RNAP II is nested within a clade of viral homologs.

Our topology for RNAP subunit 2, domain 4, interestingly consists of II, III, IV&V clades and

nonetheless suggests that giant viruses acquired RNAP before all three subunits had been

established in eukaryotes (Fig 4). Similarly, we identified a case of reciprocal gene transfer for

Transcription elongation factor IIS (C-terminal) which is responsible for regulating RNAP II,

indicating that modern eukaryotes may have derived this key element of the transcriptional

apparatus from nucleocytoviruses. The TC support values surrounding the nested eukaryotic clade

are low (<0.6) and therefore should be interpreted with some caution, however.

Importantly, our phylogenetic analysis of family B DNA polymerases also suggests that reciprocal

eukaryote-virus gene transfers have played a role in their evolution. Eukaryotes typically encode

four main family B polymerases that were likely present in LECA [49]. These four PolBs form

distinct clades in our tree, consistent with their ancient origin. Interestingly, the eukaryotic delta

family PolB (PolDelta) clade - which includes representatives from all major eukaryotic

supergroups- is clearly nested within the major PolB clade of the *Nucleocytoviricota*. This strongly

suggests that the modern eukaryotic PolDelta subunit is derived from an ancient acquisition from

giant viruses that took place during the early diversification of the PolB family in eukaryotes.

Several other smaller eukaryotic clades are nested within the *Nucleocytoviricota* in this tree,

demonstrating that more recent transfers from viruses to eukaryotes have further complicated the

evolutionary history of this protein family. This is consistent with the hypothesis presented in an

early comparative genomic study that suggested eukaryotic PolDelta arose through interactions

with viruses [35]. Similarly, the phylogeny of DNA topoisomerase IV showed acquisition of this

gene by the eukaryotic host from nucleocytoviruses, consistent with a previous study which

showed viral origin of eukaryotic DNA topoisomerase IIA [38]. Together with the possible pre-

LECA origin of the DNA sliding clamp, which associates with PolDelta, these results suggest that

the DNA replication machinery of the *Nucleocytoviricota* originated prior to the diversification of

the major eukaryotic supergroups.

One last example of possible eukaryote-virus reciprocal gene transfer involves the core histones.

All modern eukaryotes have four major types of histones (H2A, H2B, H3 and H4), all of which

have also been found in giant viruses [34]. Phylogenetic analysis of core histone proteins revealed

the presence of viruses near the base of the major core histone diversification, consistent with a

previous study that suggested that giant viruses had acquired these protein families prior to the

emergence of LECA [50]. The overall topology of this tree suggests that multiple eukaryotes to

virus gene transfers occurred, and it is possible that some of the eukaryotic clades arose through

acquisition of viral homologs. The broad distribution of viral histone homologs, together with the

overall low TC value for this tree (0.6) somewhat obscures these conclusions, however (Fig 4).

Comparison with the *Mirusviricota*

A recent study discovered a novel phylum of large DNA viruses referred to as the *Mirusviricota*

that has evolutionary links to the Nucleocytoviricota. Mirusviruses encode a herpesvirus-like

morphogenetic module but otherwise have many components of DNA and RNA processing

machinery that resemble nucleocytoviruses, and it remains unclear which lineage emerged first.

We therefore conducted separate phylogenetic analysis of several vESPs that are also found in

mirusviruses, including a DNA primase, RNA polymerase, TFIIS, ERCC4, PolB, DNA sliding

clamp, and core histones. Our phylogenetic interpretation of these genes did not change when

mirusviruses were included. Moreover, in most cases mirusvirus sequences clustered clearly

within broader clades of nucleocytoviruses, with the exception of DNA topoisomerase IV and

DNA primase (Fig 2 in S1 Text).

Quantitative comparison of tree depth across vESP categories

In order to quantitatively compare the timing at which different genes emerged in the

Nucleocytoviricota, we examined ultrametric trees of all protein families analyzed in this study

and identified the depth at which the first appeared in this viral phylum. The true vESPs trees were

midpointed while Archea-vESPs were archeal rooted, and scenarios in which the most basal-

branching lineage was viral were given a depth of zero (Fig 5). As expected, the DNA and RNA

processing vESPs were the most ancient. Our tree depth analysis showed that a giant viral clade

of mRNA cap guanylyltransferase, DNA primase, DNA sliding clamp as well as RNAP subunits

are at the root, indicative of an ancient viral origin. By comparison, vESPs involved in cytoskeletal

structure, vesicular trafficking, and ubiquitin signaling appeared in the *Nucleocytoviricota* more

recently (Fig 5).

Fig 5. Phylogenetic tree depth for Membrane trafficking, Ubiquitin system, Cytoskeleton, and DNA and RNA associated proteins.

1. X-axis value 0 indicates root while 1 indicates tip in the phylogenetic tree and 2. Black border around point on DNA and RNA processing proteins represent Archaea-vESPs.

Discussion

In this study we present a comprehensive phylogenetic analysis of ESPs encoded in nucleocytoviruses to evaluate virus-host gene transfers that may have occurred early in the evolution of eukaryotes. Phylogenies on cytoskeletal proteins, membrane trafficking associated proteins, and ubiquitin signaling proteins generally showed late branching topologies indicative of

multiple recent viral acquisitions. In contrast, phylogenies of DNA and RNA processing vESPs

revealed deeper branching topologies indicative of more ancient gene exchange between the

eukaryotes and giant viruses.

A defining feature of the *Nucleocytoviricota* is the virus factory, also called the viroplasm, which

is a membrane-bound intracellular structure that forms during infection and is the location of DNA

synthesis, viral transcription, and virion morphogenesis. Although not all members of the

Nucleocytoviricota form virus factories during infection, it is a prevalent feature of giant viruses

spanning all six orders and two classes of this phylum and is most likely a trait that was present in

their last common ancestor. Central importance has been placed on virus factories in the biology

of nucleocytoviruses [51], and any study that seeks to clarify the origin of the *Nucleocytoviricota*

must therefore consider the origin of this viral structure. Proper functioning of the virus factory

would necessarily require a viral mechanism for DNA polymerization, transcription, and mRNA

processing, and it is therefore fascinating to note that vESPs involved in these processes exhibit

the most ancient evolutionary origins of any proteins in the *Nucleocytoviricota*. We therefore

propose that the most parsimonious scenario that explains the deep-branching topologies the

family B DNA polymerase, DNA sliding clamp, topoisomerase, multimeric RNA polymerase, and

mRNA capping machinery is pre-LECA origin of the virus factory. Recent evidence suggests that

stem Eukaryotes diversified for a long period of time prior to the emergence of LECA [52], and it

is therefore not only plausible but likely that these stem Eukaryotes housed a complex array of

viruses that gave rise to the modern *Nucleocytoviricota* [53].

It is possible that the virus factory was a key innovation that potentially allowed early

nucleocytoviruses to evade host defenses by creating a physical barrier between the host cytoplasm

and the site of viral replication and transcription. This is analogous to the infection strategy used

by many large bacteriophages [54], and is therefore a mechanism employed by several distinct

lineages in the virosphere. It is unclear whether early eukaryotes possessed a fully developed

nucleus at the time that the virus factory originated, and it is possible that the emergence of the

virus factory pre-dates the nucleus. Due to the mechanistic similarity of nuclei and virus factories,

it has been proposed the former may have evolved from the latter [55,56]. While our analysis

cannot confirm or refute this hypothesis, evidence for a viral origin of eukaryotic machinery

involved in DNA replication and transcription suggests that nucleocytoviruses played a role in the

emergence of modern eukaryotes.

A recent study discovered a novel phylum of large DNA viruses, the *Mirusviricota*, that are related

to the *Nucleocytoviricota* but encode a herpesvirus like capsid [57]. The Mirusviricota encompass

a broad phylogenetic diversity and it has been suggested that they may represent a progenitor to

the *Nucleocytoviricota*. According to this scenario, nucleocytoviruses potentially emerged through

loss of the herpesvirus-like capsid and acquisition of a double-jellyroll major capsid protein and

other associated structural components. To examine this possibility, we included Mirusviruses in

the trees of DNA and RNA processing vESPs, which are the most ancient protein families in the

Nucleocytoviricota (Fig 2 in S1 Text). In most cases the *Mirusviricota* formed well-supported

clades within the *Nucleocytoviricota*, providing evidence that the latter have a more ancient

evolutionary origin and that the *Mirusviricota* likely emerged from within the *Nucleocytoviricota*.

We still know little of the enigmatic origins of the *Mirusvirocita* due to their recent discovery and

a paucity of studies that have examined this lineage, however the evolutionary origins of the

Mirusviricota should therefore be revisited as more genomes become available.

Although many vESPs involved in DNA and RNA processing have likely pre-LECA origins in

the *Nucleocytoviricota*, this is not the case for all protein families in these categories. For example,

the topologies of XPG and ERCC4 endonucleases genes clearly have basal-branching eukaryotic

lineages that suggests nucleocytoviruses acquired these genes after the emergence of the major

eukaryotic supergroups. We therefore propose that the viral acquisition of cellular genes was a

stepwise process that took place in multiple stages (Fig 6). vESPs critical for virus factory

functioning were acquired first at the pre-LECA origin of the *Nucleocytoviricota* (stage I), and

further acquisition of additional genes involved in DNA and RNA processing and transcriptional

regulation continued shortly after the emergence of modern eukaryotic diversity (stage II). In the

third stage, multiple distinct lineages of nucleocytoviruses independently acquired genes involved

in cytoskeletal dynamics, ubiquitin signaling, vesicular transport, and a wide range of other

processes. Collectively these stages explain both the evolutionary origins of the *Nucleocytoviricota*

as well as the subsequent emergence of remarkably large and complex genomes within multiple

lineages of this group. Moreover, these findings highlight examples of reciprocal gene exchange

between viruses and early eukaryotes and provide strong evidence for a viral origin of the

eukaryotic family B DNA polymerase PolDelta. Our results underscore the importance of

understanding cellular evolution in the context of co-evolution between cells and viruses.

Fig 6. Schematic diagram showing gene exchange between the eukaryotes and giant viruses.

Our results indicate three scenarios (1) gene exchange between ancient giant viruses and proto-eukaryotes (2) gene exchange between giant viruses and eukaryotes post-LECA (3) recent gene exchange.

Materials and Methods

Compilation of genomic datasets used in this study

Prior to phylogenetic analysis we compiled a set of eukaryotic, archaeal, bacterial, and nucleocytovirus genomes. We downloaded high-quality eukaryotic genomes from the eggNOG v5.0 database. Because eggNOG has relatively few protist genomes, we also included complete and chromosome-level genomes for select protists available on the National Center for Biotechnology Information (NCBI) databases as of October 8, 2021. For bacterial and archaeal genomes, we retrieved genomes from the Genome Taxonomy Database (GTDB, v95). To enrich our database in asgard archaeal genomes, we also included Asgard archaea genomes as reported in [58] that were not already present in the GTDB. For giant viruses, we used a set of curated genomes available on the Giant Virus database (https://faylward.github.io/GVDB/) that have been described previously [1]. For mirusviruses, we used genomes that were recently reported [57]. For eukaryotic genomes we used protein predictions available on EggNOG, and for all other taxa we predicted proteins using Prodigal v. 2.6.3 with default parameters [59]. A full list of all genomes used is available on https://doi.org/10.5281/zenodo.8128475 .

Eukaryotic Signature Proteins (ESPs) detection, Clustering, and Functional annotation

For *de novo* identification of ESPs, we searched bacterial, archaeal, viral, and eukaryotic genomes against the Pfam database and searched for domains that were enriched in eukaryotes. For this we used the hmmsearch command in the HMMER3 v.3.3 package with the "--cut_nc" parameter [60], with the Pfam database version 32.0 [61] as a query. To determine which Pfams were ESPs, we first removed all Pfams that were not present in >95% of all eukaryotic genomes, leaving us with a set of 763 protein families (See S1 Data). We then sought to cluster these Pfams into groups

based on their distribution across the taxonomic groups in our analysis. For this we employed an

unsupervised learning approach using Self-Organising Maps (SOMs) in R version 3.6.3,

implemented with the kohonen package (script for this is available in

https://github.com/sangitakarki/Self-Organising-Maps). To choose the optimal number of clusters

we used the heuristic "elbow method", and ultimately arrived at six groups (Fig 1A). To determine

the role of these eukaryotic proteins we performed functional annotation of proteins using

hmmsearch with 1e-5 threshold against the KEGG database (KEGG release 99.0, ver. 2021-08-

01) [62]. Since the same genes had hits to multiple KEGG categories, we manually curated the

results from the KEGG annotation to refine categories.

Phylogenetic Analysis and Evaluation of Complex Models

For phylogenetic reconstruction, we retrieved protein sequences with hits to the Pfams under

investigation, aligned them together using Clustal Omega v1.2.4 with default parameters [63], and

used trimAl v1.4. rev15 for alignment trimming (parameter -gt 0.1) [64]. We then reconstructed

maximum likelihood phylogenetic trees using IQtree v1.6.12 [65] with the option -bb 1000 to

generate 1,000 ultrafast bootstraps [66], -m MFP to determine the best-fit model [67], -nt AUTO

and --runs 5 to select the highest likelihood tree. Several best fit substitution models for different

trees were selected based on Bayesian Information Criterion (BIC) [67] (full list of best fit models

is reported in S3 Data). Most DNA and RNA processing protein trees were inferred under the LG

[68] and VT [69] amino-acid exchange rate matrices, amino-acid frequencies (F), and variation in

evolutionary rates across sites (R) [70], with different numbers of categories. Because amino acid

substitution rates likely vary across alignments, we also inferred trees using complex models (C-

models) that have different substitution matrices for every position in the alignment [71]. Since

running complex models is time and memory consuming we used a new posterior mean site

frequency (PMSF) model [72] to compare our models from -MFP option to C-models (C10 to

C60). Although the trees inferred with the -MFP option generally had lower BICs, we still

examined the trees inferred with complex models to assess any differences in topology that could

be detected using the different methods. In almost all cases, we observed similar topologies

between the two methods. One notable example is RNA pol subunit 5C, for which different groups

of nucleocytoviruses were placed as basal branching when comparing the trees inferred with

different methods (S3 Text).

All trees were visualized using the Interactive Tree of Life (iTOL) [73]. If we observed long

branches that potentially represented rogue taxa, we removed these sequences from the analysis

and re-performed alignment, trimming, and phylogenetic reconstruction. In some cases, the

number of viral proteins outnumbered those of all other taxa by a factor of two or more. In these

cases, we downsampled the viral proteins by choosing one representative nucleocytoviral genome

from each family, with the genome with the highest N50 contig size chosen as the representative.

This was done on the grounds that a higher N50 contig size is indicative of a higher quality genome

assembly. In some cases, down sampling was also done for bacteria and archaea to class level and

eukaryotes and Asgard archaea to respective representatives.

Assessing Tree Quality and Tree Depth

476   In order to evaluate the phylogenetic congruency of the trees we used the Tree Certainty (TC)

477   metric, which provides a measure of overall tree quality [44]. This was done in RaxML version

478   8.2.12 [74] (options -t and -z used to input the tree file and bootstrap file, respectively). All TC

479   scores along with alignment length are available in S2 Data. We calculated the depth of viral clades

480   in trees using a script that utilizes ape package [75] in R, bio-python and python.

Data Availability

All genomes, proteins, and alignment used in this study can be found here:

https://doi.org/10.5281/zenodo.8128475

All phylogenetic trees are available on iTOL using the username sangita1:

https://itol.embl.de/shared_projects.cgi

All codes used in this study can be found here:

https://github.com/sangitakarki/NCLDV_coevolution

Acknowledgments

491   We acknowledge the use of the Virginia Tech Advanced Research Computing Center for

492   bioinformatic analyses performed in this study. We are also thankful to the members of Aylward

493   lab for their helpful suggestions.

References

1. Aylward FO, Moniruzzaman M, Ha AD, Koonin EV. A phylogenomic framework for charting the diversity and evolution of giant viruses. PLoS Biol. 2021;19: e3001430.

2. Schulz F, Alteio L, Goudeau D, Ryan EM, Yu FB, Malmstrom RR, et al. Hidden diversity of soil giant viruses. Nat Commun. 2018;9: 4881.

3. Mihara T, Koyano H, Hingamp P, Grimsley N, Goto S, Ogata H. Taxon Richness of "Megaviridae" Exceeds those of Bacteria and Archaea in the Ocean. Microbes Environ. 2018;33: 162–171.

4. Farzad R, Ha AD, Aylward FO. Diversity and genomics of giant viruses in the North Pacific Subtropical Gyre. Front Microbiol. 2022;13: 1021923.

5. Karki S, Moniruzzaman M, Aylward FO. Comparative Genomics and Environmental Distribution of Large dsDNA Viruses in the Family Asfarviridae. Frontiers in Microbiology. 2021. doi:10.3389/fmicb.2021.657471

6. Weynberg KD, Allen MJ, Ashelford K, Scanlan DJ, Wilson WH. From small hosts come big viruses: the complete genome of a second *Ostreococcus tauri* virus, OtV-1. Environmental Microbiology. 2009. pp. 2821–2839. doi:10.1111/j.1462-2920.2009.01991.x

7. Yang Z, Gray M, Winter L. Why do poxviruses still matter? Cell & Bioscience. 2021. doi:10.1186/s13578-021-00610-8

8. Sánchez-Vizcaíno JM, Laddomada A, Arias ML. African Swine Fever Virus. Diseases of Swine. 2019. pp. 443–452. doi:10.1002/9781119350927.ch25

9. Damon IK. Poxviruses. Manual of Clinical Microbiology. 2011. pp. 1647–1658. doi:10.1128/9781555816728.ch105

10. Whittington RJ, Becker JA, Dennis MM. Iridovirus infections in finfish - critical review with emphasis on ranaviruses. Journal of Fish Diseases. 2010. pp. 95–122. doi:10.1111/j.1365-2761.2009.01110.x

11. Oliveira G, La Scola B, Abrahão J. Giant virus vs amoeba: fight for supremacy. Virol J. 2019;16: 126.

12. Van Etten JL, Graves MV, Müller DG, Boland W, Delaroque N. Phycodnaviridae– large DNA algal viruses. Archives of Virology. 2002. pp. 1479–1516. doi:10.1007/s00705-002-0822-6

13. Boyer M, Yutin N, Pagnier I, Barrassi L, Fournous G, Espinosa L, et al. Giant Marseillevirus highlights the role of amoebae as a melting pot in emergence of chimeric microorganisms. Proc Natl Acad Sci U S A. 2009;106: 21848–21853.

14. Filée J, Chandler M. Gene Exchange and the Origin of Giant Viruses. Intervirology. 2010.
pp. 354–361. doi:10.1159/000312920

15. Yutin N, Wolf YI, Koonin EV. Origin of giant viruses from smaller DNA viruses not from
a fourth domain of cellular life. Virology. 2014. pp. 38–52. doi:10.1016/j.virol.2014.06.032

16. Krupovic M, Koonin EV. Polintons: a hotbed of eukaryotic virus, transposon and plasmid
evolution. Nat Rev Microbiol. 2015;13: 105–115.

17. Koonin EV, Yutin N. Multiple evolutionary origins of giant viruses. F1000Res. 2018;7.
doi:10.12688/f1000research.16248.1

18. Koonin EV, Yutin N. Evolution of the Large Nucleocytoplasmic DNA Viruses of
Eukaryotes and Convergent Origins of Viral Gigantism. Adv Virus Res. 2019;103: 167–
202.

19. Abergel C, Rudinger-Thirion J, Giegé R, Claverie J-M. Virus-encoded aminoacyl-tRNA
synthetases: structural and functional characterization of mimivirus TyrRS and MetRS. J
Virol. 2007;81: 12406–12417.

20. Raoult D, Audic S, Robert C, Abergel C, Renesto P, Ogata H, et al. The 1.2-Megabase
Genome Sequence of Mimivirus. Science. 2004. pp. 1344–1350.
doi:10.1126/science.1101485

21. Guglielmini J, Woo AC, Krupovic M, Forterre P, Gaia M. Diversification of giant and large
eukaryotic dsDNA viruses predated the origin of modern eukaryotes. Proc Natl Acad Sci U
S A. 2019;116: 19585–19592.

22. Blanc-Mathieu R, Ogata H. DNA repair genes in the Megavirales pangenome. Curr Opin
Microbiol. 2016;31: 94–100.

23. Srinivasan V, Schnitzlein WM, Tripathy DN. Fowlpox virus encodes a novel DNA repair
enzyme, CPD-photolyase, that restores infectivity of UV light-damaged virus. J Virol.
2001;75: 1681–1688.

24. Fischer MG, Kelly I, Foster LJ, Suttle CA. The virion of Cafeteria roenbergensis virus
(CroV) contains a complex suite of proteins for transcription and DNA repair. Virology.
2014;466-467: 82–94.

25. Moniruzzaman M, Martinez-Gutierrez CA, Weinheimer AR, Aylward FO. Dynamic
genome evolution and complex virocell metabolism of globally-distributed giant viruses.
Nat Commun. 2020;11: 1710.

26. Kijima S, Delmont TO, Miyazaki U, Gaia M, Endo H, Ogata H. Discovery of Viral Myosin
Genes With Complex Evolutionary History Within Plankton. Front Microbiol. 2021;12:
683294.

27. Ha AD, Moniruzzaman M, Aylward FO. High Transcriptional Activity and Diverse

Functional Repertoires of Hundreds of Giant Viruses in a Coastal Marine System.
doi:10.1101/2021.03.08.434518

28. Cunha VD, Da Cunha V, Gaia M, Ogata H, Jaillon O, Delmont TO, et al. Giant Viruses
Encode Actin-Related Proteins. Molecular Biology and Evolution. 2022.
doi:10.1093/molbev/msac022

29. Khalifeh D, Neveu E, Fasshauer D. Megaviruses contain various genes encoding for
eukaryotic vesicle trafficking factors. Traffic. 2022;23: 414–425.

30. Barrado-Gil L, Galindo I, Martínez-Alonso D, Viedma S, Alonso C. The ubiquitin-
proteasome system is required for African swine fever replication. PLoS One. 2017;12:
e0189741.

31. Monier A, Chambouvet A, Milner DS, Attah V, Terrado R, Lovejoy C, et al. Host-derived
viral transporter protein for nitrogen uptake in infected marine phytoplankton. Proc Natl
Acad Sci U S A. 2017;114: E7489–E7498.

32. Rozenberg A, Oppermann J, Wietek J, Fernandez Lahore RG, Sandaa R-A, Bratbak G, et
al. Lateral Gene Transfer of Anion-Conducting Channelrhodopsins between Green Algae
and Giant Viruses. Curr Biol. 2020;30: 4910–4920.e5.

33. Monier A, Pagarete A, de Vargas C, Allen MJ, Read B, Claverie J-M, et al. Horizontal gene
transfer of an entire metabolic pathway between a eukaryotic alga and its DNA virus.
Genome Res. 2009;19: 1441–1449.

34. Yoshikawa G, Blanc-Mathieu R, Song C, Kayama Y, Mochizuki T, Murata K, et al.
Medusavirus, a Novel Large DNA Virus Discovered from Hot Spring Water. J Virol.
2019;93. doi:10.1128/JVI.02130-18

35. Takemura M, Yokobori S-I, Ogata H. Evolution of Eukaryotic DNA Polymerases via
Interaction Between Cells and Large DNA Viruses. J Mol Evol. 2015;81: 24–33.

36. Filée J. Multiple occurrences of giant virus core genes acquired by eukaryotic genomes: the
visible part of the iceberg? Virology. 2014;466-467: 53–59.

37. Moniruzzaman M, Weinheimer AR, Martinez-Gutierrez CA, Aylward FO. Widespread
endogenization of giant viruses shapes genomes of green algae. Nature. 2020. pp. 141–145.
doi:10.1038/s41586-020-2924-2

38. Guglielmini J, Gaia M, Da Cunha V, Criscuolo A, Krupovic M, Forterre P. Viral origin of
eukaryotic type IIA DNA topoisomerases. Virus Evol. 2022;8: veac097.

39. Gabaldón T. Origin and Early Evolution of the Eukaryotic Cell. Annu Rev Microbiol.
2021;75: 631–647.

40. Koonin EV, Dolja VV, Krupovic M. Origins and evolution of viruses of eukaryotes: The
ultimate modularity. Virology. 2015;479-480: 2–25.

41. Forterre P, Gaïa M. Giant viruses and the origin of modern eukaryotes. Curr Opin Microbiol. 2016;31: 44–49.

42. Hartman H, Fedorov A. The origin of the eukaryotic cell: a genomic investigation. Proc Natl Acad Sci U S A. 2002;99: 1420–1425.

43. Kurland CG, Collins LJ, Penny D. Genomics and the irreducible nature of eukaryote cells. Science. 2006;312: 1011–1014.

44. Salichos L, Stamatakis A, Rokas A. Novel information theory-based measures for quantifying incongruence among phylogenetic trees. Mol Biol Evol. 2014;31: 1261–1271.

45. Martinez-Gutierrez CA, Aylward FO. Phylogenetic Signal, Congruence, and Uncertainty across Bacteria and Archaea. Mol Biol Evol. 2021;38: 5514–5527.

46. Zade A, Sengupta M, Kondabagil K. Extensive in silico analysis of Mimivirus coded Rab GTPase homolog suggests a possible role in virion membrane biogenesis. Front Microbiol. 2015;6: 929.

47. Yutin N, Koonin EV. Hidden evolutionary complexity of Nucleo-Cytoplasmic Large DNA viruses of eukaryotes. Virol J. 2012;9: 161.

48. Zaremba-Niedzwiedzka K, Caceres EF, Saw JH, Bäckström D, Juzokaite L, Vancaester E, et al. Asgard archaea illuminate the origin of eukaryotic cellular complexity. Nature. 2017;541: 353–358.

49. Kazlauskas D, Krupovic M, Guglielmini J, Forterre P, Venclovas Č. Diversity and evolution of B-family DNA polymerases. Nucleic Acids Res. 2020;48: 10142–10156.

50. Grau-Bové X, Navarrete C, Chiva C, Pribasnig T, Antó M, Torruella G, et al. A phylogenetic and proteomic reconstruction of eukaryotic chromatin evolution. Nat Ecol Evol. 2022;6: 1007–1023.

51. Claverie J-M. Viruses take center stage in cellular evolution. Genome Biol. 2006;7: 110.

52. Brocks JJ, Nettersheim BJ, Adam P, Schaeffer P, Jarrett AJM, Güneli N, et al. Lost world of complex life and the late rise of the eukaryotic crown. Nature. 2023. doi:10.1038/s41586-023-06170-w

53. Krupovic M, Dolja VV, Koonin EV. The virome of the last eukaryotic common ancestor and eukaryogenesis. Nat Microbiol. 2023;8: 1008–1017.

54. Chaikeeratisak V, Nguyen K, Khanna K, Brilot AF, Erb ML, Coker JKC, et al. Assembly of a nucleus-like structure during viral replication in bacteria. Science. 2017;355: 194–197.

55. Takemura M. Medusavirus Ancestor in a Proto-Eukaryotic Cell: Updating the Hypothesis for the Viral Origin of the Nucleus. Front Microbiol. 2020;11: 571831.

56. Bell PJL. Evidence supporting a viral origin of the eukaryotic nucleus. Virus Res. 2020;289:
198168.

57. Gaïa M, Meng L, Pelletier E, Forterre P, Vanni C, Fernandez-Guerra A, et al. Mirusviruses
link herpesviruses to giant viruses. Nature. 2023;616: 783–789.

58. Liu Y, Makarova KS, Huang W-C, Wolf YI, Nikolskaya AN, Zhang X, et al. Expanded
diversity of Asgard archaea and their relationships with eukaryotes. Nature. 2021;593: 553–
557.

59. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic
gene recognition and translation initiation site identification. BMC Bioinformatics. 2010;11:
119.

60. Eddy SR. Accelerated Profile HMM Searches. PLoS Comput Biol. 2011;7: e1002195.

61. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, et al.
Pfam: The protein families database in 2021. Nucleic Acids Res. 2021;49: D412–D419.

62. Kanehisa M, Furumichi M, Sato Y, Kawashima M, Ishiguro-Watanabe M. KEGG for
taxonomy-based analysis of pathways and genomes. Nucleic Acids Res. 2023;51: D587–
D592.

63. Sievers F, Higgins DG. Clustal omega. Curr Protoc Bioinformatics. 2014;48: 3.13.1–
3.13.16.

64. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25: 1972–
1973.

65. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol.
2015;32: 268–274.

66. Minh BQ, Nguyen MAT, von Haeseler A. Ultrafast approximation for phylogenetic
bootstrap. Mol Biol Evol. 2013;30: 1188–1195.

67. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: fast
model selection for accurate phylogenetic estimates. Nat Methods. 2017;14: 587–589.

68. Le SQ, Gascuel O. An Improved General Amino Acid Replacement Matrix. Molecular
Biology and Evolution. 2008. pp. 1307–1320. doi:10.1093/molbev/msn067

69. Müller T, Vingron M. Modeling Amino Acid Replacement. Journal of Computational
Biology. 2000. pp. 761–776. doi:10.1089/10665270050514918

70. Soubrier J, Steel M, Lee MSY, Der Sarkissian C, Guindon S, Ho SYW, et al. The influence
of rate heterogeneity among sites on the time dependence of molecular rates. Mol Biol Evol.

2012;29: 3345–3358.

71. Quang LS, Gascuel O, Lartillot N. Empirical profile mixture models for phylogenetic
reconstruction. Bioinformatics. 2008;24: 2317–2323.

72. Wang H-C, Minh BQ, Susko E, Roger AJ. Modeling Site Heterogeneity with Posterior
Mean Site Frequency Profiles Accelerates Accurate Phylogenomic Estimation. Syst Biol.
2018;67: 216–235.

73. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. Nucleic Acids Research. 2021. pp. W293–W296.
doi:10.1093/nar/gkab301

74. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics. 2014;30: 1312–1313.

75. Popescu A-A, Huber KT, Paradis E. ape 3.0: New tools for distance-based phylogenetics
and evolutionary analysis in R. Bioinformatics. 2012. pp. 1536–1537.
doi:10.1093/bioinformatics/bts184

Supporting information

S1 Text. Supporting figures.

Fig 1. Phylogeny for proteins belonging to DNA processing, ubiquitin signaling,

transport/trafficking and cytoskeleton dynamics respectively. Figure 2. Phylogeny with

Mirusvirus for DNA and RNA processing proteins using a complex model (C60+F+G).

S2 Text. Supporting figures.

Fig 1: ERCC4 domain phylogeny reconstructed using VT+F+R8 model (generated by -MFP

parameter) with bootstrap support value represented by black dots (bootstrap > 80% is shown).

Fig 2: Histone phylogeny reconstructed using VT+R8 model (generated by -MFP parameter)

with bootstrap support value represented by black dots (bootstrap > 80% is shown).

Fig 3: Polymerase family B phylogeny reconstructed using VT+F+R10 model (generated by -

MFP parameter) with bootstrap support value represented by black dots (bootstrap > 80% is

shown).

Fig 4: XPG endonuclease phylogeny reconstructed using VT+F+R10 model (generated by -MFP

parameter) with bootstrap support value represented by black dots (bootstrap > 80% is shown).

Fig 5: Transcription factor TFIIB phylogeny reconstructed using LG+R8 model (generated by -

MFP parameter) with bootstrap support value represented by black dots (bootstrap > 80% is

shown).

Fig 6: Transcription factor TFIIS phylogeny reconstructed using VT+F+R8 model (generated by

-MFP parameter) with bootstrap support value represented by black dots (bootstrap > 80% is

shown).

Fig 7: Phylogeny for RNA pol Rpb2, domain 4 reconstructed using LG+F+R10 model

(generated by -MFP parameter) with bootstrap support value represented by black dots

(bootstrap > 80% is shown).

Fig 8: Phylogeny for RNA pol Rpb5, C terminal reconstructed using VT+F+R7 model

(generated by -MFP parameter) with bootstrap support value represented by black dots

(bootstrap > 80% is shown).

Fig 9: Phylogeny for DNA sliding clamp reconstructed using LG+F+R8 model (generated by -

MFP parameter) with bootstrap support value represented by black dots (bootstrap > 80% is

shown).

S3 Text. Supporting figures.

Fig 1: ERCC4 domain phylogeny reconstructed using a complex model (C60 + F +G) with

bootstrap support value represented by black dots (bootstrap > 80% is shown).

Fig 2: Core Histone phylogeny reconstructed using a complex model (C60 + F +G) with

bootstrap support value represented by black dots (bootstrap > 80% is shown).

Fig 3: Polymerase family B phylogeny reconstructed using a complex model (C60 + F +G) with

bootstrap support value represented by black dots (bootstrap > 80% is shown).

Fig 4: XPG endonuclease phylogeny reconstructed using a complex model (C60 + F +G) with

bootstrap support value represented by black dots (bootstrap > 80% is shown).

Fig 5: Transcription factor TFIIB phylogeny reconstructed using a complex model (C60 + F +G)

with bootstrap support value represented by black dots (bootstrap > 80% is shown).

Fig 6: Transcription factor TFIIS phylogeny reconstructed using a complex model (C60 + F +G)

with bootstrap support value represented by black dots (bootstrap > 80% is shown).

Fig 7: Phylogeny for RNA pol Rpb2, domain 4 reconstructed using a complex model (C60 + F

+G) with bootstrap support value represented by black dots (bootstrap > 80% is shown).

Fig 8: Phylogeny for RNA pol Rpb5, C-terminal reconstructed using a complex model (C60 + F

+G) with bootstrap support value represented by black dots (bootstrap > 80% is shown).

Fig 9: Phylogeny for PCNA reconstructed using a complex model (C60 + F +G) with bootstrap

support value represented by black dots (bootstrap > 80% is shown).

S1 Data. Clusters, annotations, genome hits and other statistics for 763 proteins reported in

this study.

S2 Data. TC scores and alignment length for phylogeny presented in this study.

S3 Data. Lists of site homogenous best fit models according to Bayesian criteria for

phylogeny presented in this study.

744