Beyond Correlation: Optimal Transport Metrics For Characterizing Representational Stability and Remapping in Neurons Encoding Spatial Memory

Andrew Aoun [1,2,3]**, Oliver Shetler** [1,2,3] **and S. Abid Hussaini** [1,2,*]

[1] *Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Columbia University Medical Center, New York, NY 10032, USA*
[2] *Department of Pathology and Cell Biology, Columbia University Medical Center, New York, NY 10032, USA*
[3] *Co-first author*

Correspondence*:
Corresponding Author
sah2149@cumc.columbia.edu

2 ABSTRACT

Spatial representations in the entorhinal cortex (EC) and hippocampus (HPC) are fundamental to
cognitive functions like navigation and memory. These representations, embodied in spatial field
maps, dynamically remap in response to environmental changes. However, current methods, such
as Pearson's correlation coefficient, struggle to capture the complexity of these remapping events,
especially when fields do not overlap, or transformations are non-linear. This limitation hinders
our understanding and quantification of remapping, a key aspect of spatial memory function. To
address this, we propose a family of metrics based on the Earth Mover's Distance (EMD) as
a robust metric for characterizing remapping. Applied to both normalized and unnormalized
distributions, the EMD provides a granular, noise-resistant, and rate-robust description of
remapping. This approach enables the identification of specific cell types and the characterization
of remapping in various scenarios, including disease models. Furthermore, the EMD's properties
can be manipulated to identify spatially tuned cell types and to explore remapping as it relates
to alternate information forms such as spatiotemporal coding. By employing approximations of
the EMD, we present a feasible, lightweight approach that complements traditional methods. Our
findings underscore the potential of the EMD as a powerful tool for enhancing our understanding
of remapping in the brain and its implications for spatial navigation, memory studies and beyond.

Keywords: remapping, stability, place cell, grid cell, activity maps, optimal transport, spatial coding, spatiotemporal

1 INTRODUCTION

The entorhinal cortex (EC) and hippocampus (HPC) have been shown to play a crucial role in spatial
navigation and memory (Frank et al. (2000), Chrobak et al. (2000), Fyhn et al. (2004), Buzsáki and Moser
(2013), Lever et al. (2002)). Specifically, through a variety of cell types across these regions, the EC-HPC
network encodes a neural representation of the spatial environment (Fyhn et al. (2004), Hafting et al.
(2005), Moser et al. (2017)). Most notable are place cells in the HPC, which integrate spatial and temporal
information received from entorhinal areas including, but not limited to, grid cells, head-direction (HD)
cells and speed cells in medial entorhinal cortex (MEC), border cells in MEC and lateral entorhinal cortex

(LEC), object cells and more recently object-trace cells and 'time' cells in LEC (Iwase et al. (2020), Lisman (2007), Steffenach et al. (2005), Moser et al. (2008), Solstad et al. (2008), Diehl et al. (2017), Hardcastle et al. (2017), Tsao et al. (2013)). However, the spatial map generated from the coordinated activity of these cell types is not necessarily stable. In fact, multiple studies demonstrated that place cells are able to remap their activity dynamically and in response to small changes in their environment (Anderson and Jeffery (2003), Colgin et al. (2008), Wilson and McNaughton (1993)). Evidence suggests this remapping of firing activity patterns enables flexible encoding of different spatial representations, a mechanism that underlies episodic memory (Leutgeb et al. (2005b), Leutgeb et al. (2004), Ferbinteanu and Shapiro (2003)). More recently, there are growing bodies of evidence indicating the EC may be the origin of place cell remapping with realignments of grid cells in MEC and global remapping of cells in wider EC also contributing (Latuske et al. (2018), Fyhn et al. (2007)). In addition, different spatial cell types such as border cells and other cells throughout EC and HPC have also been observed to "remap" by changing their firing rate or their firing field locations in response to experimental changes (Schlesiger et al. (2018), Latuske et al. (2018)). It is therefore of particular interest to be able to quantify the degree of remapping occurring in different spatial cell types and to be able to characterize the spatiotemporal changes in a given firing rate map. Such firing rate maps can also be used outside the EC-HPC circuit and are not necessarily restricted to spatial maps. Spatiotemporal maps as seen in memory circuits are used in other areas as well. Notably, such spatial maps have been used to demonstrate remapping in visual cortex where firing rate maps can be created using the position of visual objects in a subject's field of view (Wang and Sainburg (2005), Neupane et al. (2016), Merriam et al. (2007)). Yet the spatial position, or location, of a subject/stimulus is not the only modality that gives rise to firing fields. In fact, remapping is also observed in cases where spatial maps using (x,y) positions may not be as appropriate such as with motor remapping, tactile remapping, temporal remapping, angular remapping (e.g. HD cells) and other observed occurrences of neural driven remapping. However, these spatial or non-spatial maps, and the firing fields contained within them, still need to be characterized to understand the functional role remapping plays throughout the brain.

Currently, remapping has been segregated under two categories thought to represent distinct environmental changes; these are rate remapping and global remapping (Leutgeb et al. (2005b), Colgin et al. (2008)). Rate remapping is observed when testing animals in the same location but changing contextual cues within the environment (e.g. object color) and is denoted by a change in firing rate unaccompanied by a shift in place field location (Leutgeb et al. (2005b)). Global remapping, however, involves both firing rate changes and shifts in firing fields and can appear both when testing animals in different locations and with certain salient cues (Wood et al. (2000), Kentros et al. (1998), Bostock et al. (1991)). As such the boundaries between global and rate remapping are not strictly delineated with respect to the degree of change necessary to trigger them. The underlying processes do however differ in that rate remapping supports continuous information streams and has been observed to occur gradually in response to environmental changes such as morphing of different arena shapes (Leutgeb et al. (2005a), Wills et al. (2005)), while global remapping is an abrupt process where all fields for a given cell remap entirely without intermediate steps (Leutgeb et al. (2005b), Leutgeb et al. (2005a))

Although global remapping is all-or-none within a cell, this is not necessarily the case for the broader population. The presence of partial remapping, involving separate reference frames maintained by different subsets of cells, suggests that global remapping is used to transition between stable and unstable states, encoding multiple maps with different reference frames, thus supporting continuous-like information streams through population responses (Wills et al. (2005), Tsodyks (1999)). These differing subsets of reference frames within global remapping need to be characterized to understand how these transitions between states occur and the environmental influences driving them. Additionally, global remapping seems

to be a product of both a rate component and a spatial component where, in the former, firing rate is altered
and, in the latter, firing place fields can be translated, rotated, scaled or otherwise reshaped. To better
understand the specifics of these spatial transformations we need to be able to thoroughly characterize the
transitions between states. This requires disentangling the different components involved in remapping
such as spatial remapping without rate driven influences. In fact, there are three main components that
contribute to the spatial maps involved in remapping studies. These are a rate component, a temporal
component and a spatial component. The first is well defined in remapping studies however, given that rate
coding is not the only information coding schema present in EC and HPC spatial cells, it is important to be
able to describe remapping as it relates to alternate information forms such as spatiotemporal coding.

The main methods to identify rate and global remapping are based on Pearson's r correlation applied to
spatial bins on a firing rate map (Leutgeb et al. (2005b), Wills et al. (2005), Hussaini et al. (2011)). This
approach is sufficient to identify linear relationships in transitions between firing rate maps on different
experimental sessions or conditions. However, Pearson's r is vulnerable to outliers and cannot effectively
capture non-linear transitions nor can it allow for a segregation of remapping types beyond that of pure
rate remapping or joint rate-spatial remapping (i.e. global remapping with a rate change and a spatial
transformation of varying complexity). This correlation approach compares spatial bins at the same position
across different ratemaps for a given cell and is most informative when only linear rate changes are
occurring. Pearson's r can quantify simple rate changes but is unable to quantify global remapping beyond
its presence or lack thereof. Therefore the profile for global remapping incorporates both a change in
firing rate and any type of non-linear shift in the spatial map density that cannot easily be described
by a correlation metric. Pearson's r also requires distributions to be the same in size which restricts the
information that can be carried in a sample of the rate map.

We can see that existing methods are only able to distinguish simple changes in rate from broader whole
field changes in rate and space without an explicit characterization of the varied spatial transformations
observed in global remapping. One can imagine, in a simple example, that a place field retaining its rate
while changing the shape or dispersion of the field will be poorly distinguished from a place field retaining
its rate without distortion or other in-place modifications of the cell field. In fact, seemingly random
distortion in place fields and grid fields is seen in multiple disease models emphasizing the need for a
stronger metric to capture spatial dissimilarity (Jun et al. (2020), Fu et al. (2017)). In a further example,
one can imagine two distinct place fields that remap in different ways such that both change their firing rate
but one is translated while another is rotated. Given current approaches, both these sets of transformations
would be deemed global remapping despite the clear difference in how the remapping occurred and the
exact origins behind the distinct outcomes. This range of spatiotemporal changes within global remapping
is complex and holds necessary information for understanding the mechanism of spatial cell remapping, a
key component of episodic memory (Knierim et al. (2006), Alexander et al. (2020)). We therefore believe a
more rigorous approach focused on spatiotemporal distances and capturing non-linear rate transformations
is needed to further probe the complexities of remapping clearly visible in spatial navigation and memory
studies. One such method that can complement Pearson's r correlations for rate remapping and extend
our ability to identify and describe both spatial and temporal transformations in global remapping is the
Earth Mover's Distance (EMD). The EMD is a distance computed on a pair of distributions (Vasserstein
(1969), Panaretos and Zemel (2019)), often applied in computer science for image analysis tasks (Rubner
et al. (2000)). This distance when computed using unnormalized distributions can also be referred to as the
discrete Wasserstein distance; for normalized distributions it reduces to the Wasserstein distance, but will
be described as the 'normalized EMD' in this paper to avoid confusing the EMD and Wasserstein distance
as wholly separate metrics. The EMD has been shown to be a highly robust spike train distance metric to

quantify temporal similarity patterns in spike trains with varying rate profiles, enabling us to probe the
pure temporal component of remapping (Sihn and Kim (2019), Sotomayor-Gómez et al. (2021)). However,
the rate component remains the primary source of evidence in current remapping studies and the spatial
component, as well as the joint spatio-temporal component, remain poorly quantified.

This study aims to address the challenges in quantifying and characterizing the remapping process of
spatial cell types observed in navigation studies, memory studies, and beyond, with particular focus on
non-linear transformations and transitions between states. Current methods, such as Pearson's r correlation,
have limitations in distinguishing simple rate changes from broader whole field changes in rate and space,
and in characterizing the varied spatial transformations observed in global remapping. Therefore, we
propose a more rigorous, comprehensive approach that focuses on spatiotemporal distances and captures
non-linear rate transformations. We employ the Earth Mover's Distance (EMD), a robust metric traditionally
used in computer science for image analysis tasks, to delve deeper into the complexities of remapping that
are evident in spatial navigation and memory studies. The EMD quantifies the minimum optimal transport
distance between two 2D firing rate distributions capturing non-linear non-overlapping transformations and
changes in shape and dispersion of fields. We aim to explore how the EMD can enhance our understanding
of mechanisms underlying spatial cell remapping and provide a valuable metric for quantifying remapping
across different experimental trials/sessions. We consider the use of such a metric for neurodegenerative or
otherwise impaired remapping studies by exploring spatial sensitivity and noise robustness. We investigate
how generalized approximations of the EMD could be used to manipulate spatial sensitivity properties
and identify various cell types associated with points, areas, or other mapped stimuli, such as object and
trace cells in LEC or point-driven attention mechanisms in visual areas. We also consider the feasibility
of applying the EMD given the computational cost by comparing the sliced EMD approximation, which
allows for efficient applications of the normalized and unnormalized 2D EMD by computing the average of
many 1D EMD values along random image slices (Bonneel et al. (2015)). This approximation is used in
all computations in this paper apart from the single-point Wasserstein generalizations (see methods). We
further assess the feasibility of applying the EMD given the nature of stability and remapping data where
rate effects are varied, and rate changes commonplace. The EMD's performance is also assessed against
the non-linear spearman rank correlation coefficient. Through this comprehensive analysis, we underscore
the superiority of the EMD over traditional metrics, highlighting its unique ability to elucidate complex
spatiotemporal transformations, effectively distinguish various global remapping patterns, and maintain
robustness in the face of noise, rate fluctuations, and other potential distortions.

2 METHODS

2.1 Synthetic Fields

Place cells and grid cells were modeled as gaussian blobs with fixed standard deviation. Place cell
centroids were restricted to bins inside the square map. Place cells were modeled as (17,17) ratemaps with
standard deviation varying between 1 and 3 for different figures where standard deviation was consistent.
For elliptical fields, two standard deviation parameters are used for the y and x direction. These were set to
1 and 3 respectively. To model remapping, we shift the location of the centroid on a wider map (N*3, N*3)
and slice the relevant portion to center the field as needed.

Grid cells consisted of multiple place fields with kernel size and standard deviation parameters. They
were organized in a hexagonal pattern across a wider grid (N*8 + kernel * 2) with gapN bins separating
place field edges both horizontally and vertically. Slices were taken across this wider grid to obtain (17,17)

rate maps of grid 'cells'. To model remapping, we shift the initial sliced grid map by N*N pairs (from 0 to
N) on a wider map (N*8 + kernel*2) and slice the relevant portion to shift the grid phase as needed.

In generating heatmaps, while place field centroids are shifted across the map (left/right/up/down) and
tested against a fixed field at the center, grid fields are shifted right by 0 to N and down by 0 to N creating
stepwise slices across the wider grid. These slices are tested against the initial sliced grid map. Therefore
grid cell examples are not shifted around the absolute center of the grid map but rather moving away from
the top left. Given that the wider grid pattern is consistent and symmetrical, the information provided by
the results should be no different than if a single point at the center of the grid map was chosen to shift
around.

Both wider and tighter spaced grid modules were considered by varying standard deviation and gapN
parameters. Grid field centroids were not necessarily inside the (17,17) ratemap at all slices taken and a
part of the field excluding the centroid could be included in the slice..

All fields were interpolated to (257, 257) for plotting only. Additional examples of synthetic fields without
interpolation are provided in the supplementary documents (S1-S3).

2.2 EMD

173 The Earth Mover's Distance (EMD) is a measure of dissimilarity between two arbitrary un-normalized
174 distributions defined over a metric space with a distance metric $d(x, y)$. Intuitively, EMD can be thought of
175 as the minimum cost required to transform one distribution into another, where the cost is proportional to
176 the amount of "earth" moved and the distance it is moved. The general formula for the EMD is:

$$EMD(a, b) = \min_{f \in F(a,b)} \sum_{x,y \in D} f(x, y) d(x, y) \tag{1}$$

In this equation, $f(x, y)$ represents the flow from element x to element y, and $d(x, y)$ denotes the ground
distance between $\times$ and y. $F(a, b)$ is the set of all feasible flows satisfying the following constraints:

$$\sum_{y \in D} f(x, y) \leq a(x), \quad \forall x \in D$$

$$\sum_{x \in D} f(x, y) \leq b(y), \quad \forall y \in D$$

$$\sum_{x,y \in D} f(x, y) = \min \left(\sum_{x \in D} a(x), \sum_{y \in D} b(y) \right)$$

These constraints ensure that the total flow from each element in a does not exceed its value, the total flow
to each element in b does not exceed its value, and the total flow between the two distributions is equal to
the smaller sum of either distribution.

To compute the EMD, one must solve a transportation problem, which is an instance of a minimum-cost
flow problem. In the one- dimensional case, the EMD has a closed-form solution that is much simpler to
compute compared to the multi-dimensional case.

Let $P(x)$ and $Q(x)$ be two one-dimensional distributions defined over the same domain, with cumulative
distribution functions $F_P(x)$ and $F_Q(x)$. The EMD between them can be computed as:

$$EMD(P,Q) = \int \left| F_P(x) - F_Q(x) \right| dx \tag{2}$$

185 In the discrete case, it can be calculated as:

$$EMD(P,Q) = \sum_{i=1}^{n} \left| F_P(i) - F_Q(i) \right| \tag{3}$$

where $F_P(i)$ and $F_Q(i)$ are the cumulative sums of the respective distributions up to index 1. The one-
dimensional EMD has a closed-form solution because the optimal transport plan is unique and easy to
find. In contrast, the multidimensional EMD, which extends the one- dimensional EMD algorithm to
multiple-dimensional signals such as images or videos, does not have a closed-form solution. The optimal
transport plan is more complicated, and the problem becomes a linear optimization problem. Common
approaches to computing the multi-dimensional EMD include the Hungarian algorithm or other linear
programming techniques, which can solve the problem in polynomial time.

The algorithmic complexity of the multi-dimensional EMD depends on the chosen method for solving
the linear optimization problem. For the Hungarian algorithm, the complexity is $O\left(n^3\right)$, where n is the
number of elements in each distribution. For two dimensional distributions, the complexity increases to
$O\left((mn)^3\right)$, for a $m \times n$ distribution. For example, rate-maps of size 16×16 ($n = 16^2 = 256$) would
require $16^6 = 16,777,216$ operations and a 32×32 ratemap would require $1,073,741,824$ operations.
The computational cost can be reduced in practice by using approximations. In this paper, we use the Sliced
Earth Mover's Distance (Sliced EMD) (Bonneel et al. (2015)).

The Sliced EMD is an efficient algorithm to estimate the EMD between multi-dimensional distributions,
such as 2D distributions, by leveraging the closed-form solution for the 1D case. The main idea behind the
sliced EMD is to project the multi- dimensional distributions onto multiple one-dimensional lines (slices)
and then compute the 1D EMD on each of these slices. The average of the 1D EMDs across all slices
provides an approximation of the multi-dimensional EMD.

The sliced EMD algorithm does the following steps:

• Choose a set of random directions (slices) in the 2D space.

• Project the 2D distributions onto each of these slices.

• For each slice, sort the projections of both distributions.

• Calculate the 1D EMD between the sorted projections using the closed-form solution.

• Average the 1D EMDs across all slices to obtain the sliced EMD.

The algorithmic complexity of the sliced EMD is determined by the number of slices (L), the number of
points in each distribution (N), and the sorting complexity. Since sorting has an average complexity of
$O(N \log N)$, the total complexity of the sliced EMD algorithm is $O(LN \log N)$. For a 16×16 ratemap
and $L = 10^3$ projections would be $2,048,000$. For a 32×32 ratemap, the complexity scales to $10,240,000$.
The percent complexity of the Sliced EMD compared to the EMD computed with the Hungarian Algorithm
for 16×16 and 32×32 ratemaps respectively is 1.22% and 0.95% respectively.

Compared to the Hungarian algorithm and other linear programming techniques, the sliced EMD provides
a much more computationally efficient approximation for multi-dimensional EMD, especially when the
number of points in the distributions is large. While the accuracy of the sliced EMD may not be as high as
the exact EMD computed using the Hungarian algorithm, it often provides a very good balance between
computational efficiency and accuracy, making it suitable for various applications where an exact EMD
calculation would be too computationally expensive. For two dimensional distributions, the sliced EMD
generally converges with between one hundred and ten thousand projections (Figure S11). Even at the upper
bound of ten thousand projections, the sliced EMD is substantially more efficient than Optimal Transport
techniques. We used the sliced EMD approximation with 10**4 slices to reproduce near-theoretical EMD
scores on all figures in our analysis except for field localization figures (10**2).

2.3 Single point Wasserstein

In this paper, we primarily use the Sliced Earth Mover's Distance (EMD) to compare two distributions.
However, there is a special use case where the EMD complexity can be further reduced. This use case
arises when comparing a normalized ratemap with a distribution that has all its mass concentrated at a
single point.

In this case, the EMD formula simplifies to the sum of the distances between the point of interest and each
point in the distribution, weighted by the normalized proportion of mass at that point in the distribution.
Specifically, let P be a normalized ratemap and Q be a distribution that has all its mass concentrated at a
single point q. The EMD between P and Q is given by:

$$EMD(P, Q) = \sum_{x \in D} \|x - q\|_2 \cdot P(x) \tag{4}$$

Here, $\|x - q\|_2$ denotes the Euclidean distance between x and q, and P(x) represents the proportion of
mass at point $\times$ in the normalized ratemap P. This formula reduces the complexity of computing the EMD
from $O\left(n^3 \backslash \log n\right)$ to $O(n)$, where n is the number of elements in the distribution.

This simplified EMD formula is useful for object remapping, where one is interested in comparing a
ratemap to a distribution that represents a single object. For example, it can be used to compare the ratemap
of a rodent's environment with a distribution that represents the location of a reward or danger zone.

2.4 Polymorphisms

243 Here, we outline specific polymorphisms of the EMD that we compute for several use cases.

2.4.1 Whole map EMD

245 The whole map EMD is computed with two ratemaps. For this use case, the Sliced EMD is used with
246 between 10^2 and 10^4 projections.

2.4.2 Field EMD

248 The field EMD is a special use case where only firing fields are considered. In this case, rates are imputed
249 to zero outside of the firing field while rates within the firing field are retained. The Sliced EMD is then
250 used to estimate the EMD on these masked ratemaps.

2.4.3 Binary EMD

The binary EMD is computed using ratemaps where one is imputed for the bins inside the firing field and zero is imputed for those outside the firing field.

2.4.4 Centroid distance

The centroid distance refers to the euclidean distance between two points (e.g. field centroid-object or field centroid- field centroid)

2.5 Firing blob extraction

Firing fields were estimated as in (Fyhn et al. (2007)). The peak firing rate was chosen as any bin in the rate map with the largest value (rate). Firing fields were determined as a contiguous region where the firing rate was above 20% of the peak rate. Therefore the top 80% of bins were considered and a blob extraction procedure was then used to extract contiguous regions.

2.6 Correlation measures

Pearson's r correlation coefficients were computed using the python package Scipy. Correlations were computed between the raw and normalized 2D gaussian distributions by comparing the value at each bin in the NxN ratemap. Spearman-rank correlation coefficients were computed in the same way using the spearman function instead.

2.7 Code

All analyses were done in python using custom code on Jupyter notebooks. All plots were made using matplotlib. Sliced Wasserstein distances on whole maps and field restricted maps as well as binary EMD distances were computed using the python optimal transport package (ot). Map to point distances were computed using custom single-point Wasserstein functions. All functions are integrated as part of our custom analysis toolkit Neuroscikit. A prototype to analyze remapping under different EMD metrics for whole map to whole map cases, object/quadrant cases, field to field cases and specific session grouping cases is provided and being developed as part of this toolkit.

3 RESULTS

In order to understand why EMD outperforms Pearson's r, we need to explore remapping concepts with fine grained control of spatial fields. We therefore opted to synthetically generate examples of field maps where manipulations would allow us to vary 1) field size (stdev) 2) field noise (stdev) 3) field shape (ellipse or circle) 4) field count (1,2 or grid) and, most importantly, 5) field location (x,y) (see methods). We did this by modeling fields as gaussian blobs with centroid coordinates (x,y) on a square map of size N and a standard deviation parameter to vary the field intensity. When necessary and wherever specified, we added normally or uniformly distributed noise across the ratemap. These synthetic fields allow us to approximate certain cell types and different firing field situations that come about in experimentally recorded data. We use these synthetic fields to demonstrate how EMD outperforms Pearson's r in single, dual and multi-field cases.

3.1 EMD captures linear and non-linear transformations resulting in non-overlapping fields

While single field maps can be thought of as a modeling of place cells, the insights derived from the EMD in such situations are applicable to any type of single field map. Therefore, to first compare the Earth Mover's Distance (EMD) metric to the Pearson's r correlation coefficient, we considered the simplest remapping case which involves two maps with one identical field in each, but at different locations. This would approximate the simplest case of spatial remapping where every aspect of a field is unchanged except the position on a map. More importantly however, this manipulation allows us to consider remapping transformations that result in non-overlapping fields, a case that Pearson's r would quantify as no correlation ($p = 0$). Such cases are critical to quantify since remapping resulting in non-overlap still has a biological significance and/or a driver beyond that of having remapped or not. These non-overlapping transformations are often seen in cases of global remapping.

Our synthetic place fields were modeled as 2D-gaussian distributions ($\sigma = 1$) with a single centroid (center bin) in the middle of a square map. To allow for a single point that is at the true center of the square map, we used N = 17 bins for height and width as opposed to the more traditional (16,16) ratemaps. While one ratemap had its field, and centroid, fixed to the middle of the map, the other was translated across the square such that the centroid had visited every possible bin in the (NxN) ratemap. At every point in a bin, the remapping scores were computed between the fixed and translated rate map. Weights from the ratemap were normalized resulting in a normalized EMD score (Wasserstein distance). As such, we tested remapping across all possible transformations resulting in a field, at least partially, still in the map but not necessarily overlapping with the fixed field. In doing so, we demonstrate that the EMD metric shows greater sensitivity to spatial transformations than the Pearson's r correlation coefficient (Figure 1). Specifically, the EMD metric is able to capture all possible linear and non-linear transformations resulting in either overlapping or non-overlapping receptive fields. We show that, for a given pair of identical receptive fields, circular or elliptical, the EMD score is non-zero for all possible place field centroids whereas, as one place field from the pair is gradually translated outside the field of the other, the Pearson's r coefficient accelerates to 0 (Figure 1). In fact, Pearson's r is unable to quantify any remapping that results in wholly non-overlapping receptive fields (r = 0) while the EMD is non-zero at all possible centroids. Therefore the coverage of information provided by Pearson's r is restricted to remapping that results in overlap between fields and varies depending on the size and placement of the field in a map.

Experimentally however, single field cells are not the only observation and various cell types exhibit multiple fields. One such prominent example is the observed pattern in grid cells where multiple fields are laid out in a hexagonal pattern. Combinations of these grid cells coordinate as grid modules with various orientation, spacing and other properties that enable advanced spatial processing using the overlap of multiple grid cells (Hafting et al. (2005), Fyhn et al. (2007)). As such, we then considered the case of overlapping grid modules, both tight and wide in their spacing. In the wide case, we modeled N=3 grid fields with $\sigma = 2$ and 10 bins of separation. In the tight case, we consider $N = 3$ grid fields with standard deviation = 1 and only 6 bins of separation. Given that the map size was N=17, and the requirement to not overlap fields in a grid cell, the decrease in bin separation from 10 to 6 is non-trivial and leads to significantly less spacing between grid fields. In both cases we follow a similar procedure as above and shift the grid module horizontally and vertically for N*N pairs of positions creating a map of EMD and Pearson's r scores (Figure 2). We demonstrate that EMD also surpasses Pearson's r in its sensitivity to phase transitions. Specifically, we see that both EMD and Pearson's r suitably capture the transition from in phase to out of phase but EMD provides a more specific quantification of this boundary. For ease of

comparison, we inverted the color scheme of the EMD heatmaps so that hotter colors represent similarity and colder colors represent dissimilarity in a way aligned with how Pearson's r shows up on comparable heatmaps. In doing so we see that the inverted EMD heatmap demonstrates a broader range of EMD values which allow for a narrower determination of the phase crossing boundary (thinner shaded region at boundary crossing). Additionally, Pearson's r also results in 0 correlation scores when grid modules are completely non-overlapping, EMD however can still quantify these regions of no overlap. Therefore EMD also offers more spatial sensitivity in the case of grid fields.

The EMD is especially useful in cases of no overlap where global remapping transformations struggle to be defined with Pearson's r. The application of the EMD on place cell transformations resulting in non-overlap demonstrates this (Figure 1, Figure S4). We see reflective properties in that all four corners of the square rate map show the same EMD score gradient. Similarly, the top, bottom, left and right (N,S,W,E) locations show highly similar EMD values. This suggests that the EMD can facilitate the characterization of complex non-linear transformations involving rotations of fields, both overlapping and non-overlapping. This can also be extended to object fields, object trace fields, border fields and any other point or area driven field remapping. Additionally, given that the gradient of EMD scores extends outside the area covered by the gaussian field, this metric may be able to describe remapping and distortions of the underlying field, either in the form of simple scaling or complex degeneration of fields. The increased spatial sensitivity demonstrated by the EMD is also reflected in the elliptical and circular nature of place fields being captured in the underlying EMD distribution (Figure 1, Figure Figure S4). EMD is equally sensitive and informative in the case of non identical place fields, where field shape is varied, as seen with the circle-ellipse pair and the rotated ellipse-ellipse pair further suggesting that the EMD may be used to capture complex changes in the field shape, size and distribution on a rate map (Figure S4).

These complex changes and field distortions are commonplace in neurodegenerative studies such as the observed progressive degeneration of spatial maps in AD mouse models during spatial memory and navigation tasks (Jun et al. (2020), Fu et al. (2017), Rodriguez et al. (2020)). Despite the impairments to the underlying spatial map, the remapping of such a map still needs to be quantified to assess differences in stability, learning and memory in these disease states. Additionally, inactivation studies, such as those involving muscimol to inactivate medial septum (MS) or hippocampal input to EC resulting in impaired grid cells and grid cell disappearance respectively (Brandon et al. (2011), Bonnevie et al. (2013)), are commonplace and assessing the stability of spatial maps post inactivation is important. To confirm the benefit of using the EMD in cases of impaired cell maps, we considered two more remapping scenarios involving no translations to the field centroid but rather in-place transformations on the field itself resulting in noisy or degenerate spatial maps.

3.2 EMD is more stable to noise and field degeneration

In the first case, we considered a scaling remapping of the place field where a gradual increase in the standard deviation of the gaussian field on a fixed (33x33) map results in an increasingly large field (σ range $= 0 - 6$). We found that while Pearson's r was poor at quantifying spatial scaling and insensitive to minor changes, the EMD maintained its spatial sensitivity, and symmetry, allowing for a quantification of both scaling up and scaling down of place fields relative to a single fixed field ($\sigma = 3$) (Figure 3). The EMD is therefore sensitive across a broader range of standard deviations than Pearson's r which can only really distinguish large shifts in scaling magnitudes. Pearson's r is also skewed such that larger standard deviations are less distinguishable than smaller ones while the EMD is robust in both directions and symmetrical.

In the next case, we considered a more complex 'remapping' of place fields (17 by 17 pixels) involving
progressive degeneration of spatial maps as commonly seen in spatial memory studies of disease models
(Jun et al. (2020),Fu et al. (2017)). To model this progressive degeneration of a spatial map and the
increased 'noise' associated with this degeneration, we added 17 by 17 pixel noise maps in a stepwise
manner. The noise fields were sampled from a normal distribution ($\mu = 0$) with increasing standard
deviation. We did this for two pairs of place fields ($\sigma = 1$), non-overlapping and partially overlapping place
fields (Figure 3). For each we computed the whole map EMD, the Pearson's r correlation coefficient and a
field-restricted EMD using blob extraction based on 20% of the peak firing rate as in (Fyhn et al. (2007)).
In the former non-overlapping case, we see significant deviation of the Pearson's r correlation coefficient
from the baseline remapping value of -0.0454 computed at no noise. The value is close to 0 since there is
no overlap between the fields and bins outside the field are close to 0. We see however that as we increase
the added noises, the Pearson's r score accelerates away from the true score and begins to plateau around
$r = 0.8$ to 1, solely due to extraneous noise. This same logarithmic trend in Pearson's r scores is evident in
the overlapping case but with a larger base correlation given the initial overlap (base: 0.339, μ: 0.814, σ:
0.190).

The whole map EMD on the other hand remains more stable relative to the baseline no-noise score and
deviates significantly both fewer times and at larger standard deviations. This same trend is seen in the
overlapping (base: 0.208, μ: 0.369, σ: 0.171) and non-overlapping (base: 1.17, μ: 1.29, σ: 0.247) cases.
We also find that the whole map EMD is a more stable choice at progressively larger standard deviations
than the field EMD, which restricts itself to the area occupied by a field. In the partially overlapping case
the field EMD appears more stable than the overlapping case in that it fluctuates on a similar scale and to
a similar degree relative to the whole map EMD (base: 0.183, μ: 0.369, σ: 0.222). We also see that they
have the same mean (0.369) and similar baseline and standard deviations (0.208 vs 0.183 and 0.171 vs
0.222) but with larger deviations for the field EMD. This contrasts with the non-overlapping case where
the field EMD, at larger standard deviations, seems to deviate further from baseline than in the overlapping
case (base: 1.04, μ: 1.03, σ: 0.202). We find that the field EMD in the non-overlapping case has a lower
standard deviation than the whole map EMD (0.247 vs 0.202) but also varies on a smaller scale. These
differences in the field EMD and whole map EMD can be understood through the interference of added
noise on the detection of firing fields based on peak rate (discussion section). Even so, we find that the
EMD is more robust to noise and stays closer to the baseline EMD score. As such the EMD scores provide
a more stable metric to capture remapping than the Pearson's r score, which is highly susceptible to noise.

Given that Pearson's r is so sensitive to outliers and these ratemaps were un-smoothed and un-normalized
post added noise, we repeated the procedure by smoothing ratemaps with a gaussian kernel (size= 5,
$\sigma = 1$) and normalizing post smoothing (Figure S5, S6). In doing so, we demonstrate that the observed
trend holds and the normalized EMD (Wasserstein distance) again proves to be a more robust metric. We
first find that the logarithmic distribution of Pearson's r scores across noise standard deviations seen in
the unnormalized case remains but shifts to more linear in the smoothed-normalized case (base: 0.563, μ:
0.763, σ: 0.120). We observe this for both the non-overlapping and partially overlapping cases. EMD on
the other hand is again a more stable choice and deviates less from the baseline score for the overlapping
(base: 0.0831, μ: 0.189, σ: 0.318) and non overlapping cases (base: 0.475, μ: 0.621, σ: 0.464). The whole
map EMD scores also have noticeably fewer outliers in that there are fewer large peaks away from baseline.
We also see that the field EMD evolves similarly to the unnormalized example for the partially overlapping
and non overlapping case. In the former however, we see that the fluctuation is at a different range of values
to the whole map EMD (base: 0.0830, μ: 0.0789, σ: 0.0264), reflective of normalization inside the detected
field area, and in the latter, we see that the field EMD deviates more significantly from the baseline score

than the normalized example (base: 0.472, μ: 0.380, σ: 0.120). We see similar trends in a third test with
normalized but unsmoothed ratemaps where Pearson's r is unchanged from the unnormalized case owing to
the inherent normalization procedure of computing Pearson's r. Whole map EMD is similarly stable across
different noise standard deviations, with a greater range due to an extreme outlier while field EMD is also
stable relative to baseline with a sharper deviation from baseline as is seen in the non-overlapping case.
Therefore EMD is less sensitive to degeneration for both unnormalized and normalized ratemaps, with or
without smoothing, emphasizing the need for it in identifying regions of low remapping for particularly
noisy or disease-driven degenerate spatial maps

We therefore demonstrate that the EMD computed between two ratemaps is robust to noise and
degeneration while Pearson's r is not. The emphasis however is on a robust distance between two non-
identical ratemaps (two different cells/sessions) in the presence of increasing noise. While remapping
stability is often assessed between different cells or a single cell across different sessions, this is not the
only case that can benefit from informed stability analyses. In fact, the EMD should also maintain this noise
robustness in a within-cell or within-session comparison as opposed to an across cell/session comparison.
That is, the EMD should be able to assess cell stability within a session for a single cell map. This is an
important property in order to determine regions of low or high spatial remapping. Effectively, such a
property is important to confidently localize a field on a map and so that we can describe how remapping
regions are related to spatial locations of different objects, points or even other centroids/fields on that
same map.

3.3   EMD quantiles extend field localization and capture remapping relative to a point

To test such within-session stimulus/location-relative stability, we use an adjusted EMD metric computed
between the true whole map and a 'fake' map where all the density has been placed in a single bin (single
point Wasserstein distance). This process is repeated across all possible bins in the 17x17 spatial map such
that a map of EMD scores should be lowest (least dissimilar/most similar) at the point of lowest remapping.
In the case of place fields or other single field cells, the point of lowest remapping is presumed to be at or
near the centroid of the field.We do this for a rate map with different standard deviations of added noise. To
account for border and numerical issues that reflect the scores, we pad the ratemap with N=2 bins (Figure
S8). To reduce the influence of outliers on Pearson's r and firing rates, we smooth the ratemaps after adding
noise (Figure S8). We note that smoothing primarily recovers Pearson's r's accuracy but not the EMD
which remains robust without smoothing (Figure S8). We also find that, relative to Pearson's r, the EMD is
more robust to noise with the spatial bins holding the top 20% of EMD scores being less spread out and less
sparse than the spatial bins holding the top 20% of Pearson's r scores as the amount of noise is increased
(Figure 4, FigureS7, Figure S8). This is particularly evident on noise heavy maps where Pearson's r loses
its specificity towards the region of lowest remapping while EMD retains it. Pearson's r and the EMD do
however demonstrate similar specificity towards the 'true' field centroid. In fact, computing the centroid of
the field based on the 80th percentile of EMD and Pearson's r scores results in a similar approximation to
the true centroid for EMD scores (Figure 4, Figure S7, Figure S8).

When we compare the whole-map to point-map EMD scores with a blob detection approach based on
firing bins that cross 20% of the peak firing rate (Fyhn et al. 2007), we see a similar result for the case of a
single place field. That is, the 80th quantile of scores from the map to point approach perform similarly to
the blob detection approach at low noise levels, especially the top 20% of pearson scores and the top 90%
of EMD scores (Figure S8). However as noise increases, the specificity of the top 20% Pearson's r scores
and the 20% peak threshold firing scores is degraded. That is, the scores become less specific and therefore

less sensitive as a tool to determine the true centroid of a field and locate that field on a map. At higher standard deviations of noise, the quality is sufficiently degraded to result in multiple fields being detected despite there being a single place field on the original map. These often smaller fields that are outside the region encompassed by the top 80% of EMD scores (and top 90%) would be removed experimentally if they were below a certain area threshold. However it is not uncommon for noise blobs (noise fields) to cross this threshold and the number/shape can often vary with ratemap smoothing parameters that are part of the field detection process (Grijseels et al. 2021) (Figure S9, Figure S10). As such blob detection algorithms based on a percentage threshold of the peak firing rate may be more robust than other existing methods but can result in too many fields being detected (Figure S9, Figure S10). In the case of single fields, the EMD is a more robust metric to measure remapping relative to a point than the Pearson's r score and may be even used as a filter/validation method for field detection in cases of noisy rate maps where multiple dispersed noise fields are extracted.

In the case of dual fields however, the top 80% of EMD values span a contiguous region that includes the area spanned by both fields. While EMD is still robust to degeneration and noise in the dual field case, we can see that it does not capture the separate nature of each field at larger standard deviations (Figure 5). In fact, while the top 90% of EMD scores is robust to field shape at low standard deviations, this is not the case for larger standard deviations where the map is sufficiently distorted for the EMD to miss the dual field relationship. Despite this we still observe that the relative positioning of the highest EMD quantile region is more consistent and less degenerate at higher standard deviations (Figure 5). We see that the EMD peak regions are dependent on the placement of fields on the map. With multiple fields, the peak EMD will be located at a weighted average of all fields, which is the lowest point of remapping, and requires the least amount of work in order to shift all the density to that point as opposed to any other point on the rate map. As such, the map-to-point EMD approach is still informative in the case of multiple fields and the quantiles generated by comparing remapping at every possible point allows for a localization of low and high remapping regions. This could be used to identify the region or quantile of remapping relative to a known stimulus region (e.g. object location) and could even help filter out detected fields that fall far outside of the contiguous regions of lowest remapping, Additionally, the quantiles generated by the whole map to point EMD enable a suitable reference distribution to compare experimentally recorded distances. These quantiles provide us with a percentile that describes how far a remapping distance falls relative to all, or a sample of all, possible points that a field or cell could remap to. This can be used to describe the region of remapping a field falls in relative to objects, regions of interest or general areas on a spatial map. These quantiles also create a common scale to compare in situations where EMD distances are highly influenced by occupancy and coverage distances or even different arena sizes where there are inherently different total distances to move as part of the EMD computation. We also see that these quantiles are sustained through increasing noise providing a robust metric to assess spatially correlated and/or driven remapping.

3.4 EMD captures object or trace cells and other position/rotation driven remapping

The robustness of the EMD to noise seen using the map-to-point approach, coupled with the symmetrical/reflective properties shown and the quantile reference distribution, lends itself to the investigation of specific entorhinal and hippocampal cell types. Specifically, the EMD map-to-point approach involving a quantile reference distribution can be used to track remapping relative to a stimulus point (object location), both current and past (object and object-trace), relative to a stimulus region, both general (N,S,E,W) and specific (single point or multi-location), and relative to current position, both discrete (place cell with single preferred x,y location) and continuous (stepwise sliding window). With the findings of object, location and stimulus encoding playing a central role in EC-HPC circuitry, it is

critical that we are able to quantify such remapping to understand its role in episodic memory and changes
associated with impairments in this role (Wilson et al. (2013a), Chao et al. (2016), Wilson et al. (2013b),
Tsao et al. (2013), Wang et al. (2018)). Additionally, point driven remapping is especially important given
theories of EC functioning and mounting evidence for relative temporal tracking of stimulus locations and
identities in LEC (Wilson et al. (2013a), Tsao et al. (2013), Wang et al. (2018)). Such point remapping is
also found in visual areas where remapping of pointers is suggested to underlie an attention mechanism
(Cavanagh et al. (2010)). This evidence further supports the notion of attention-modulated salient-stimulus
tracking in the LEC where relative changes in stimulus location, identity or simply intensity are likely
encoded and cells need thorough characterization to understand their functional role.

In the case of object location tracing, we can use a combination of the whole map-to-point map EMD
and a centroid distance (euclidean distance between centroids) from a known object location to determine
whether, on the current session, remapping is driven towards that location. Specifically, for a given NxN
ratemap, we compute the EMD between the true rate map and a fake rate map where a single bin (object
location) holds all the density. We can do this relative to the current session/trial or a previous session/trial.
We consider a model experiment setup where an object is shifted across four possible locations and a
synthetic object 'cell' traces the current location of the object with the placement of its field ($N = 17$,
$\sigma = 1$) (Figure 6). At each object 'session', we compute the centroid distance between the field and all
4 possible locations as well as map-to-point EMD values. We see that these values are at their lowest
when the field is overlapping with the object location, at their largest when in the furthest of the 4 possible
locations and the same value when in one of the remaining two corners. The two corners with the same
EMD value (identical distance away from current location) can be distinguished from one another using
the centroid distances. The single point EMD (Wasserstein) quantiles can be used to determine how close a
field/cell is to the point of lowest remapping and where that point is relative to the recorded object locations.
This demonstrates that we can quantify remapping towards the current object location on a trial, previous
trial location or even future trial location (if we suspect the subject's cell of predicting the change). Thus
the EMD enables us to identify object, object-trace and/or other types of cell tracking mechanisms.

Importantly however, we again notice the reflective property of the EMD where two possible object
positions share the same EMD value despite having different centers. The location of the objects on the
map (quadrant/region) and the vector transformations from the field centroids towards these different
locations (magnitude + angle) will allow us to differentiate between them despite the similarity in EMD
scores. Therefore, a substantial shift in centroid location (large magnitude + particular angle) associated
with a low remapping value can be indicative of rotational remapping for a map/field. We can see that this
property is not present with Pearson's r scores which are unable to quantify any rotational remapping or
object tracing resulting in non-overlap with r=0 at all points outside of current object location ($N = 33$,
$\sigma = 3$ for rotating field) (Figure 6). Identifying rotational remapping and describing the angle and direction
of a rotation is important to understand how these rotations come about and the specific influences that may
be driving them or that they may be driven towards. These rotations, either whole map or for a specific field
on a map, have been observed experimentally in different contexts including, but not limited to, grid cell
rotational realignment in entorhinal cortex (Fyhn et al. (2007)) and place cell cue-driven rotations (Fenton
et al. (2000), Muller and Kubie (1987)).

EMD is even more suitable to quantify these rotations and other stimulus driven remappings as it can be
computationally simplified for the point map case. Specifically, we show that the properties seen in the
whole map to single point map EMD computation are held in cases where the object location is poorly
defined or when we are restricted to a field. In the former case, we computed the EMD between only the

field on a map and a general quadrant encompassing ¼ of the square arena and including the current object
location. In the latter, we computed a simplified, normalized EMD (single point Wasserstein) between
only the field on a map and a singular point rather than the entire 'fake' point map. In both cases the
reflective property that supports complex remapping involving rotations and/or tracing is maintained.
Therefore the concept of a whole-map to point-map approach can be generalized to any combination of
whole-map/field-map to point-map/quadrant-map/single-point remapping.

The generalization of the whole map - point map EMD into a whole map - single point Wasserstein
significantly reduces computation time and provides a simpler and more flexible way to apply the EMD
computation. This is highly useful for less specific remapping cases involving no object locations but
broader environment differences spanning multiple spatial bins. For example, EMD can also be used to
characterize rotational remapping of border cells or remapping towards/driven by entire quadrants/regions
of an arena/environment. EMD is also unrestricted by non-overlapping rotational remapping. If we consider
remapping in the form of field rotation for a pair of overlapping place fields, we see that Pearson's r cannot
effectively describe rotations. We show that the angle of rotation or presence of a different angle of rotation
cannot effectively be distinguished. We find that Pearson's r is close to 0 for around 100 to 250 degrees of
rotation and is blown up below and above those limits. EMD on the other hand is much smoother across all
tested angles demonstrating its effectiveness both for rotational remapping that results in either overlapping
or non-overlapping fields (Figure 6).

3.5 EMD on spatial maps is robust to rate changes

The feasibility of the EMD is not only supported by the various simplifications of its computation (map
to point, sliced map, field to field), but also by the rate robustness seen with the EMD in a previous study
(Sihn and Kim (2019)) and in our results. Given that firing rate can vary within a session and even more so
across sessions separated by days, the EMD benefits from having rate robustness properties which allow
for a characterization of spatiotemporal changes alongside the traditional rate remapping quantifications.
In the previous study, EMD was shown to be rate robust for temporal coding by varying the total number
of spikes. The authors demonstrated that the variability of the EMD across different firing ratios of spike
trains was negligible whereas the distance variability for varying degrees of temporal similarity was not,
thus allowing for a rate-robust pure-temporal spike train distance (Sihn and Kim (2019)). We also see this
rate-robustness in our data where EMD values do not vary significantly with different firing rate ratios in
normalized EMD distances. However, it is common practice in field comparison studies to discard low
firing rates to reduce spurious correlations caused by noise. While this can help reduce noise effects, there
is an inherent loss of information associated with the change in computed firing distribution. Therefore
having a rate robust metric that can include all spikes with minimal impact from noise is especially
important. As such, we propose the use of a binary EMD to describe field dispersion regardless of the
rate distribution. We demonstrate this rate robustness using the binary EMD generalization alongside
normalized and unnormalized whole-map and field-map EMD distances (Figure 7). To do this, we used the
scaling of place fields as a type of rate transformation. We can view this transformation as an increase in
field intensity where larger intensities indicate higher firing rates while smaller intensities indicate lower
firing rate. For a fixed field, scaling its activity across sessions, we computed three measures of EMD
remapping involving whole map to map EMD, field to field EMD and a binary EMD that imputes 0 outside
a field and 1 inside a field thereby silencing rate effects and varying solely with density dispersion. In
practice, one could compute two separate 'binary' metrics where one is field restricted and imputes 0 or 1
as described while another spike density EMD would use all raw spike positions (x,y for a given spike)
directly without distributing them into a weighted (NxN) firing rate map (whole map dispersion). Since we

synthesized our data for these simulated cases, we can only apply the binary EMD and not the whole map
spike density EMD. We repeat the selected EMD measures on our rate maps for normalized (Wasserstein)
and unnormalized (EMD) intensity changes (place field scaling) (Figure 7). Given that the binary EMD
uses raw spikes rather than firing rate maps, we find that the scores are the same in both the normalized and
unnormalized case. As such, the binary EMD provides a rate robust metric to quantify the dispersion of rate
maps and/or the dispersion of the underlying field in the ratemap, regardless of the rate distribution across
the map/field. This binary EMD approach can thus be used to describe remapping without the influence of
the rate distribution.

In the case of EMD measures including rate influences, we test both the whole map EMD and the field
restricted EMD. In the normalized case, we see that the field EMD (Wasserstein) and the whole map EMD
(Wasserstein) show similar gradients/rates of change across intensity scaling factors (standard deviations).
We also see that the whole map EMD and field EMD vary on a similar scale to the binary EMD since
rate effects have been normalized. In the unnormalized case, the scales change as the whole map EMD
and field EMD are weighted by the firing rates in their respective bins (whole map = all bins, field = field
bins only). This scale of EMD values better reflects the use of raw firing rate scores. Additionally, as we
increase the intensity, the whole map EMD and field EMD increase their separation. When compared to
the normalized case, this is again reflective of the use of raw firing rate scores which, at larger standard
deviations, result in larger differences between whole map EMD and field EMD. Therefore the use of the
normalized and unnormalized EMD results in similarly evolving trends for the distance metric enabling
a rate robust interpretation of the results. However, we find that, when comparing two identical spatial
maps with varying firing rate ratios, the EMD is highly stable if using the binary EMD or the normalized
EMD but varies when using unnormalized rates (Figure 7). These differences therefore allow the EMD
to be manipulated in such a way that rate effects can be separated from spatiotemporal changes and the
underlying shifts in shape and dispersion described regardless of intensity changes. As such, we can
investigate remapping both with and without rate effects. We can characterize remapping for reduced rate
effects (normalized EMD == Wasserstein distance) and raw rate effects (unnormalized EMD) and compare
these to binary or spike density EMD scores to separately quantify the contribution to remapping provided
by rate and/or spatiotemporal changes.

3.6 EMD outperforms linear and non-linear metrics

While Pearson's r was the primary metric tested alongside EMD, owing to its popular use in remapping
studies, it is not the only plausible metric to apply. As such, we also thought it relevant to compare the
EMD and Wasserstein metric to another common metric. Since Pearson's r is a strong linear metric, we
opted for a non-linear metric found in the Spearman rank correlation coefficient. We repeated all figures
with the spearman rank coefficient instead of the Pearson's r coefficient and provide a sample of key
figures in the supplementary documents (S12 - S18). We find that the EMD remains the superior choice as
an overall more sensitive and robust metric. Specifically, we see that Spearman-ρ actually outperforms
Pearson's r in its spatial sensitivity but still cannot effectively capture all transformations resulting in
non-overlap. While in the case of two overlapping circular fields of identical shape Spearman-ρ is fully
symmetric like the EMD, it does not maintain this property across different field shapes with the overlapping
ellipse pair (same angle of rotation) only allowing for a distinction of left and right as opposed to all four
corners. In fact, despite being a non-linear metric, Spearman-ρ cannot effectively quantify all complex,
non-linear transformations. This is further seen in its inability to quantify overlapping identical fields
across different scales. Spearman-ρ is however capable of describing transformations such as rotations or
single point/quadrant remapping if the two fields are identical circles. In such cases, Spearman-ρ appears

to support rotations and object remapping. This however is highly restricted by noise, as is the case with
Pearson's r. We find that Spearman-ρ is not robust to noise and quickly degenerates away from the baseline,
no-noise Spearman score. Spearman-ρ is therefore similarly susceptible to Pearson's r despite the added
nonlinear sensitivity. Overall however, the EMD's non-linear selectivity is more encompassing, more robust
and more effective at quantifying and characterizing the spatial transformations underlying remapping.

4   DISCUSSION

Through these simulated, general cases of remapping, we demonstrate that the Earth Mover's Distance
(EMD) is more spatially sensitive in characterizing remapping than the Pearson's r correlation coefficient
and other plausible metrics like the non-linear Spearman rank correlation coefficient. We find that, both
Pearson's r and EMD are suitable for cases of remapping where fields are still overlapping from session to
session. However, we demonstrate that Pearson's r is unsuitable in describing remapping that results in
non-overlapping receptive fields whereas EMD can numerically quantify remapping at any point in the
ratemap. We show that EMD holds its sensitivity for circular/elliptical place fields and wide/narrow spaced
grid fields. Additionally, we find that the symmetry offered by EMD allows us to describe more complex
non-linear translations such as rotations and scaling of fields. In fact, we find that Pearson's r is less suitable
than the EMD in describing either scaling or rotations and does not offer the same sensitivity that the
EMD provides. More importantly however, we show that this spatial sensitivity of the EMD to linear and
non-linear transformations is robust to noise and field degeneration. By manipulating fixed fields across a
range of standard deviations of added noise, we find that, for both normalized and unnormalized ratemaps,
the EMD is more stable relative to the 'true' EMD score. That is, as we increase the standard deviation
of noise, EMD varies less frequently and at later standard deviations than the Pearson's r score. Notably,
we see that the EMD performs similarly in the normalized and unnormalized cases also demonstrating a
robustness to rate. While the raw EMD values change for the normalized and unnormalized ratemaps, the
distribution across standard deviations evolves in the same ways, with fewer outliers in the normalized and
smoothed-then-normalized cases than the unnormalized case, further demonstrating a type of robustness to
rate. The application of a binary EMD where all spikes are considered with even weight by imputing 0
and 1 for outside and inside the field respectively or by using raw spike positions directly (spike density)
further disentangles rate effects by exclusively describing the underlying dispersion of the field or map,
regardless of firing rate. This highlights the use of such a metric in describing field or whole map distortions
and degenerations where the spatial distribution of spikes can change separately to rate and even the field
centroid.

This application of a spike density EMD where all spikes are considered with even weight by using raw
spike positions enables us to effectively disentangle rate effects by exclusively describing the underlying
dispersion of the map, regardless of firing rate. This can also be extended to the case of individual fields
and highlights the use of such a metric in describing field distortions and field degenerations where the
actual distribution of the field can change separately to rate and even the centroid. Such dispersion metrics
can be thought of as purely spatial and can be extended to include the entire distribution of spike positions
as part of a spike density EMD. This 'pure spatial' metric may be most useful in the case of linear tracks
with identical navigation structure across animals but can also be extended to arena navigation studies
where coverage and occupancy vary from animal to animal. In the latter, caution should be taken to account
for coverage where arena sizes may be consistent but map exploration varies from animal to animal.
Normalization of distances by coverage limits would reduce such effects. Additionally, care should be taken
in interpreting dispersion metrics (binary and spike density) in the case of extremely biased occupancy

where most of the spiking distribution will fall in the same area purely due to biased behavior. This argument also extends to the use of EMD to assess temporal similarity profiles as part of a 'pure temporal' metric (Sihn and Kim 2019). These pure metrics may be best interpreted in such a way where especially stable distances (binary/density or temporal EMD) along with fair occupancy, and while accounting for coverage limits, would be a non-trivial result, as would differing trends among the rate, temporal and spatial components of remapping. In fact, given that rate remapping is well defined and temporal remapping of firing profiles can be quantified (Sihn and Kim 2019), we further suggest the EMD and single point Wasserstein metric to describe the spatial component in firing rate maps enabling a much more detailed and flexible framework for remapping. Through various manipulations, this would allow for an understanding of how different components of neural coding interact by separately and concurrently considering the rate, spatial and temporal components.

We also see in these examples that the whole map EMD is more suitable than the field-restricted EMD in capturing remapping despite degeneration. The reason behind this is likely two fold. The first being that the field-restricted EMD does not capture noise across the entire spatial map, therefore, because in our example, randomly sampled normally distributed values are added at every position in the ratemap (NxN noise), the specific subset added to the indices of a given field may be particularly large/small relative to the rest of the distribution. As such, from step to step (each increase in std dev), restricting the EMD metric to the single field can cause larger deviations than would be seen with the whole map case. The second reason lies in the methods for detection of gaussian fields and the sensitivity of these approaches to noise and spatial degeneration. The best approach is often to select a contiguous region from a given spatial map, with a minimum size, where the firing rate is above a certain peak threshold (Fyhn et al. (2007), Grijseels et al. (2021)). Although this is suitable in the cases of low noise, degenerate and noisy spatial maps are often experimentally recorded and need more granular characterization of place field centroids and area. Traditional field characterization methods are sensitive to noise and often involve smoothing of the ratemap as part of this procedure (Fyhn et al. (2007), Grijseels et al. (2021)). With minimal smoothing, localization of place fields in a noisy map results in multiple 'noise' fields detected with centroids deviating from the true field positions (Figure S9, Figure S10). With greater smoothing, localization of place fields on a noisy map can result in too wide an area being characterized as a field and multiple fields being incorrectly merged (Figure S9, Figure S10). In the presence of similar noise throughout the rate map, smoothing can even result in a contiguous region crossing detection thresholds despite being outside the fields of interest (Figure S9, Figure S10). Therefore the field EMD can be particularly susceptible to the accuracy of the detected fields and by extension to the underlying field detection process. However both the whole map EMD and field EMD still offer more stable and robust alternatives to Pearson's r, especially in cases of high noise and for disease states.

While we find that the whole map to point map EMD does not replace the map blobs (fields) detection approach, we show that it can help inform classification of valid blobs by highlighting the region of lowest remapping. In the case of known single field cells (e.g. place cells), the map to point EMD will identify the region of lowest remapping which falls near the true field centroid and can therefore be used to filter out extracted noise blobs that may have crossed the area/size threshold but fall far from the region of lowest remapping. In the case of dual fields, the EMD can only separately identify fields at low noise levels and using the top 90% threshold as opposed to the top 20% of Pearson's r or 20% peak firing rates. In fact, in the case of dual fields, the EMD gravitates to the center point between the two fields which offers the lowest point of remapping. While it cannot be used for field detection in this case, it may yet be used to filter out blobs that fall far from this region. Therefore the whole map to point map EMD approach is suitable for localizing in the case of a known single field (e.g place cell) or to identify the region of lowest

remapping in the case of multiple fields. In fact, the EMD is not as susceptible to noise as Pearson's r
and peak firing rates in that it performs more consistently and with less degradation in either the case of
unnormalized ratemaps or smoothed then normalized ratemaps.

Moreover, the remapping quantiles that are generated across the entire map for single and multi-field
cases can be used as a standard reference distribution. That is, single point EMD (Wasserstein) scores can
be converted into a quantile below which all values are smaller (easier point locations to remap to) and
above which all values are larger (harder point locations to remap to). We see that these quantiles are a result
of computing remapping scores between the whole map and every possible point on the rate map (Figure
4). In doing so, single point EMD values, and by extension individual fields, can be localized on a rate map
through the 'region' of lowest remapping that they fall in. These quantiles also enable comparisons across
cells where field locations, arena sizes and firing rates are sufficiently different that the raw EMD values
cannot be effectively compared. For example, in practice, while arena size may be consistent, coverage and
occupancy can vary, as can firing rates as a result of exploration. Despite the rate robustness of the EMD
distance, the single point Wasserstein generalization involves a direct multiplication using weights from
the rate map bins (normalized firing rates) where excessive differences due to purely behavioral aspects
(such as low firing rates due to low coverage/occupancy in an area of the map) can result in EMD values
varying on different scales but not for a physiological reason. In such a scenario, one might observe a
comparable trend in remapping within a cell's own sessions but the raw difference in EMD scores is not
comparable across different cells if the behavioral profiles are substantially different for the single point
Wasserstein. In fact, while two separate cells might increase or decrease their remapping score across a
group of sessions, the scale within which they vary can be different due to inherent differences in coverage
where the range of positions covered by two ratemaps is not identical. Even with consistent coverage of
arenas, several experimental contexts make use of changing arena shapes thus changing the total distance
available in the EMD computation. Providing a quantile alongside a remapping value allows for a standard
scale to compare different cells and describes how far/close the computed remapping score is relative to
all the possible points of remapping in a rate map. If a point were the lowest point of remapping (e.g.
exactly at the object location for an object cell or at the centroid of an individual field), then the quantile
would be sufficiently small such that most or all of the other possible locations result in a larger remapping
score. These quantiles are unique to the EMD and appear across noise standard deviations allowing for
a quantification of remapping regions despite degeneration, instability and substantial behavioral/rate
influences.

The advantages of the whole map to point map approach are pertinent to a wide array of remapping
investigations and spatial navigation research. Numerous studies have reported complex remapping in
relation to objects and locations, highlighting the need for a more nuanced description and characterization
of these changes beyond the two prevailing categories of rate and global remapping. By generalizing
the whole map to point map Earth Mover's Distance (EMD) computation into a whole map to single
point or whole map to quadrant methodology, the EMD enables a more detailed analysis and serves as an
instrumental tool for identifying object cells, object-trace cells, and other stimulus or location-driven cells
exhibiting intricate rotational remapping or transformations resulting in non-overlap. Consequently, the
EMD emerges as a pivotal metric in delineating these non-linear, non-overlapping remapping outcomes.
We advocate the employment of the EMD to further scrutinize cases of global remapping and discern
the impact of stimuli on cell remapping. The EMD's viability as a remapping metric is bolstered by the
computational expediency of its simplified cases. The modified map to single point approach allows for
swift and adaptable application of the EMD, while the sliced EMD approximation can adequately estimate
the true EMD value. Utilizing the sliced EMD across varying numbers of projections, with $10^{**}2$ offering

a favorable balance of speed and accuracy, renders the EMD an optimal choice for cell remapping analyses.
Nonetheless, the sliced EMD may still be computationally demanding in instances involving exceptionally
large rate maps (e.g., 256 x 256) and a substantial number of cells.

While the EMD applied to pairs of whole spatial maps requires the sliced EMD approximation due
to having large 2 dimensional distributions, EMD can be applied faster and more efficiently on smaller
distributions. EMD on 2 dimensional maps is better equipped to describe the spatiotemporal patterns
seen than Pearon's r correlation coefficient however the temporal aspect of these maps could be further
characterized by applying the EMD in a stepwise manner. In a basic example, one could take the spiking
activity of a given cell and iterate over time windows of activity (with window size and step size as
parameters) to produce a continuous string of EMD values describing remapping as it evolves over an
experimental session or trial. While this approach is highly unsuitable with Pearson's r because of the need
for same size distributions and the over-susceptibility to spurious correlations at lower sample sizes, EMD
has been shown in this paper to be more robust and in previous work to support continuous application
(Zhao et al. (2010)). Given that the EMD can be applied on raw spike positions (binary EMD) and does not
require a rate map to describe the dispersion of density across the map, we further propose it as a tool to
investigate remapping continuously. And given that we do not require overlapping fields nor same size
distributions to apply the EMD, we argue that remapping can be described in a continuous manner for
window sizes with sufficiently large spike sample rates. As such, EMD offers additional flexibility for
disentangling spatiotemporal components and can be manipulated in a continuous-like setup to describe
remapping as it relates to specific temporal events and/or markers. With sufficiently long experimental
sessions and high sample rates, EMD will support a characterization of remapping on smaller timescales
than separate sessions and can enlighten intersession and intertrial remapping dynamics that are triggered
or otherwise shaped by time. For example, one can consider an experimental setup where object location is
rotated continuously or otherwise transformed during the session, as was seen in a previous study (Shapiro
et al. (1997), Knierim et al. (1995)). In such a case, applying EMD across a sliding window of spike activity
can be highly informative in identifying how remapping evolves over time for a given cell and for the
broader population. This can be especially useful given the varying results seen in morph experiments
where partial and complete remapping are seen to occur in different ratios across different studies (Wills
et al. (2005), Leutgeb et al. (2005a), Colgin et al. (2010)). Using EMD we can better describe the quantity
of remapping over time, the periods or triggers before/after which the gradient of remapping increases or
decreases, and the amount of remapping across the population of cells (or ensembles of cells) at specific
points in time.

In summary, the Earth Mover's Distance (EMD) offers a more comprehensive and spatially sensitive
approach to characterizing remapping in comparison to the Pearson's r correlation coefficient. EMD's ability
to handle non-overlapping receptive fields and intricate non-linear transformations, such as rotations and
scaling, renders it a powerful tool for understanding the complexities of spatial navigation and remapping.
Although Pearson's r might remain useful in specific cases with linear relationships and overlapping fields,
EMD's versatility makes it applicable to a broader range of scenarios. EMD estimators such as sliced EMD
are computationally expensive with respect to correlation metrics. However, most modern computers can
easily handle the additional computational load and this should not be a hindrance to the adoption of EMD
in most use-cases. The application of EMD in spatial remapping research has far-reaching implications in
the study of memory and neurodegenerative disorders, such as Alzheimer's Disease. By providing a more
detailed analysis of spatial remapping, EMD can shed light on the intricate relationships between spatial
representations, memory formation, and the influence of various factors on these processes. The enhanced
understanding of remapping dynamics facilitated by EMD may contribute to the identification of potential

therapeutic targets for memory-related disorders, thereby opening new avenues for Alzheimer's Disease
research and treatment.

CONFLICT OF INTEREST STATEMENT

814 The authors declare that the research was conducted in the absence of any commercial or financial
815 relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

816 A.A, O.S and S.A.H conceptualized the project. O.S defined the statistical methods. A.A designed the
817 experiments and performed the data analysis. A.A wrote the manuscript with inputs and edits from O.S
818 and S.A.H.

FUNDING

819 NIH/NIA grants: R01AG064066 and supplement: R01AG064066-S1

ACKNOWLEDGMENTS

820 We would like to thank Dr. Gustavo Rodriguez and Dr. Radha Raghuraman for their advice and assistance
821 in developing this framework and for their helpful comments throughout the writing of this manuscript.

DATA AVAILABILITY STATEMENT

822 The jupyter notebooks used to synthesize data and generate the figures can be found at the Hussaini Lab
823 github page in the Neuroscikit project under _prototypes/cell_remapping /remapping_paper.

REFERENCES

Alexander, A. S., Robinson, J. C., Dannenberg, H., Kinsky, N. R., Levy, S. J., Mau, W., et al. (2020).
Neurophysiological coding of space and time in the hippocampus, entorhinal cortex, and retrosplenial
cortex 4, 2398212820972871. doi:10.1177/2398212820972871
Anderson, M. I. and Jeffery, K. J. (2003). Heterogeneous modulation of place cell firing by changes
in context 23, 8827–8835. doi:10.1523/JNEUROSCI.23-26-08827.2003. Publisher: Society for
Neuroscience Section: Behavioral/Systems/Cognitive
Bonneel, N., Rabin, J., Peyré, G., and Pfister, H. (2015). Sliced and radon wasserstein barycenters of
measures 51, 22–45. doi:10.1007/s10851-014-0506-3
Bonnevie, T., Dunn, B., Fyhn, M., Hafting, T., Derdikman, D., Kubie, J. L., et al. (2013). Grid cells require
excitatory drive from the hippocampus 16, 309–317. doi:10.1038/nn.3311. Number: 3 Publisher: Nature
Publishing Group
Bostock, E., Muller, R. U., and Kubie, J. L. (1991). Experience-dependent modifications
of hippocampal place cell firing 1, 193–205. doi:10.1002/hipo.450010207. _eprint:
https://onlinelibrary.wiley.com/doi/pdf/10.1002/hipo.450010207
Brandon, M. P., Bogaard, A. R., Libby, C. P., Connerney, M. A., Gupta, K., and Hasselmo, M. E. (2011).
Reduction of theta rhythm dissociates grid cell spatial periodicity from directional tuning 332, 595–599.
doi:10.1126/science.1201652. Publisher: American Association for the Advancement of Science

Buzsáki, G. and Moser, E. I. (2013). Memory, navigation and theta rhythm in the hippocampal-entorhinal
system 16, 130–138. doi:10.1038/nn.3304. Number: 2 Publisher: Nature Publishing Group

Cavanagh, P., Hunt, A. R., Afraz, A., and Rolfs, M. (2010). Visual stability based on remapping of attention
pointers 14, 147–153. doi:10.1016/j.tics.2010.01.007. Publisher: Elsevier

Chao, O. Y., Huston, J. P., Li, J.-S., Wang, A.-L., and de Souza Silva, M. A. (2016).
The medial prefrontal cortex—lateral entorhinal cortex circuit is essential for episodic-like
memory and associative object-recognition 26, 633–645. doi:10.1002/hipo.22547. _eprint:
https://onlinelibrary.wiley.com/doi/pdf/10.1002/hipo.22547

Chrobak, J. J., Lörincz, A., and Buzsáki, G. (2000). Physiological patterns in the
hippocampo-entorhinal cortex system 10, 457–465. doi:10.1002/1098-1063(2000)10:4⟨457::
AID-HIPO12⟩3.0.CO;2-Z. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/1098-
1063%282000%2910%3A4%3C457%3A%3AAID-HIPO12%3E3.0.CO%3B2-Z

Colgin, L. L., Leutgeb, S., Jezek, K., Leutgeb, J. K., Moser, E. I., McNaughton, B. L., et al. (2010).
Attractor-map versus autoassociation based attractor dynamics in the hippocampal network 104, 35–50.
doi:10.1152/jn.00202.2010. Publisher: American Physiological Society

Colgin, L. L., Moser, E. I., and Moser, M.-B. (2008). Understanding memory through hippocampal
remapping 31, 469–477. doi:10.1016/j.tins.2008.06.008

Diehl, G. W., Hon, O. J., Leutgeb, S., and Leutgeb, J. K. (2017). Grid and nongrid cells in medial entorhinal
cortex represent spatial location and environmental features with complementary coding schemes 94,
83–92.e6. doi:10.1016/j.neuron.2017.03.004

Fenton, A. A., Csizmadia, G., and Muller, R. U. (2000). Conjoint control of hippocampal place cell
firing by two visual stimuli: I. the effects of moving the stimuli on firing field positions 116, 191–210.
doi:10.1085/jgp.116.2.191

Ferbinteanu, J. and Shapiro, M. L. (2003). Prospective and retrospective memory coding in the hippocampus
40, 1227–1239. doi:10.1016/S0896-6273(03)00752-9

Frank, L. M., Brown, E. N., and Wilson, M. (2000). Trajectory encoding in the hippocampus and entorhinal
cortex 27, 169–178. doi:10.1016/S0896-6273(00)00018-0

Fu, H., Rodriguez, G. A., Herman, M., Emrani, S., Nahmani, E., Barrett, G., et al. (2017). Tau pathology
induces excitatory neuron loss, grid cell dysfunction, and spatial memory deficits reminiscent of early
alzheimer's disease 93, 533–541

Fyhn, M., Hafting, T., Treves, A., Moser, M.-B., and Moser, E. I. (2007). Hippocampal remapping and
grid realignment in entorhinal cortex 446, 190–194. doi:10.1038/nature05601. Number: 7132 Publisher:
Nature Publishing Group

Fyhn, M., Molden, S., Witter, M. P., Moser, E. I., and Moser, M.-B. (2004). Spatial representation in the
entorhinal cortex 305, 1258–1264. doi:10.1126/science.1099901. Publisher: American Association for
the Advancement of Science

Grijseels, D. M., Shaw, K., Barry, C., and Hall, C. N. (2021). Choice of method of place cell classification
determines the population of cells identified 17, e1008835. doi:10.1371/journal.pcbi.1008835. Publisher:
Public Library of Science

Hafting, T., Fyhn, M., Molden, S., Moser, M.-B., and Moser, E. I. (2005). Microstructure of a spatial
map in the entorhinal cortex 436, 801–806. doi:10.1038/nature03721. Number: 7052 Publisher: Nature
Publishing Group

Hardcastle, K., Ganguli, S., and Giocomo, L. M. (2017). Cell types for our sense of location: where we are
and where we are going 20, 1474–1482. doi:10.1038/nn.4654. Number: 11 Publisher: Nature Publishing
Group

Hussaini, S. A., Kempadoo, K. A., Thuault, S. J., Siegelbaum, S. A., and Kandel, E. R. (2011). Increased size and stability of CA1 and CA3 place fields in HCN1 knockout mice 72, 643–653. doi:10.1016/j.neuron.2011.09.007

Iwase, M., Kitanishi, T., and Mizuseki, K. (2020). Cell type, sub-region, and layer-specific speed representation in the hippocampal–entorhinal circuit 10, 1407. doi:10.1038/s41598-020-58194-1. Number: 1 Publisher: Nature Publishing Group

Jun, H., Bramian, A., Soma, S., Saito, T., Saido, T. C., and Igarashi, K. M. (2020). Disrupted place cell remapping and impaired grid cells in a knockin model of alzheimer's disease 107, 1095–1112.e6. doi:10.1016/j.neuron.2020.06.023

Kentros, C., Hargreaves, E., Hawkins, R. D., Kandel, E. R., Shapiro, M., and Muller, R. V. (1998). Abolition of long-term stability of new hippocampal place cell maps by NMDA receptor blockade 280, 2121–2126. doi:10.1126/science.280.5372.2121. Publisher: American Association for the Advancement of Science

Knierim, J. J., Kudrimoti, H. S., and McNaughton, B. L. (1995). Place cells, head direction cells, and the learning of landmark stability 15, 1648–1659. doi:10.1523/JNEUROSCI.15-03-01648.1995. Publisher: Society for Neuroscience Section: Articles

Knierim, J. J., Lee, I., and Hargreaves, E. L. (2006). Hippocampal place cells: Parallel input streams, subregional processing, and implications for episodic memory 16, 755–764. doi:10.1002/hipo.20203. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/hipo.20203

Latuske, P., Kornienko, O., Kohler, L., and Allen, K. (2018). Hippocampal remapping and its entorhinal origin 11

Leutgeb, J. K., Leutgeb, S., Treves, A., Meyer, R., Barnes, C. A., McNaughton, B. L., et al. (2005a). Progressive transformation of hippocampal neuronal representations in "morphed" environments 48, 345–358. doi:10.1016/j.neuron.2005.09.007

Leutgeb, S., Leutgeb, J. K., Barnes, C. A., Moser, E. I., McNaughton, B. L., and Moser, M.-B. (2005b). Independent codes for spatial and episodic memory in hippocampal neuronal ensembles 309, 619–623. doi:10.1126/science.1114037. Publisher: American Association for the Advancement of Science

Leutgeb, S., Leutgeb, J. K., Treves, A., Moser, M.-B., and Moser, E. I. (2004). Distinct ensemble codes in hippocampal areas CA3 and CA1 305, 1295–1298. doi:10.1126/science.1100265. Publisher: American Association for the Advancement of Science

Lever, C., Wills, T., Cacucci, F., Burgess, N., and O'Keefe, J. (2002). Long-term plasticity in hippocampal place-cell representation of environmental geometry 416, 90–94. doi:10.1038/416090a. Number: 6876 Publisher: Nature Publishing Group

Lisman, J. E. (2007). Role of the dual entorhinal inputs to hippocampus: a hypothesis based on cue/action (non-self/self) couplets. In *Progress in Brain Research*, ed. H. E. Scharfman (Elsevier), vol. 163 of *The Dentate Gyrus: A Comprehensive Guide to Structure, Function, and Clinical Implications*. 615–818. doi:10.1016/S0079-6123(07)63033-7

Merriam, E. P., Genovese, C. R., and Colby, C. L. (2007). Remapping in human visual cortex 97, 1738–1755. doi:10.1152/jn.00189.2006. Publisher: American Physiological Society

Moser, E. I., Kropff, E., and Moser, M.-B. (2008). Place cells, grid cells, and the brain's spatial representation system 31, 69–89. doi:10.1146/annurev.neuro.31.061307.090723. _eprint: https://doi.org/10.1146/annurev.neuro.31.061307.090723

Moser, E. I., Moser, M.-B., and McNaughton, B. L. (2017). Spatial representation in the hippocampal formation: a history 20, 1448–1464. doi:10.1038/nn.4653. Number: 11 Publisher: Nature Publishing Group

Muller, R. U. and Kubie, J. L. (1987). The effects of changes in the environment on the spatial firing
of hippocampal complex-spike cells 7, 1951–1968. doi:10.1523/JNEUROSCI.07-07-01951.1987.
Publisher: Society for Neuroscience Section: Articles

Neupane, S., Guitton, D., and Pack, C. C. (2016). Two distinct types of remapping in primate cortical area
v4 7, 10402. doi:10.1038/ncomms10402. Number: 1 Publisher: Nature Publishing Group

Panaretos, V. M. and Zemel, Y. (2019). Statistical aspects of wasserstein distances 6, 405–431. doi:10.1146/
annurev-statistics-030718-104938. _eprint: https://doi.org/10.1146/annurev-statistics-030718-104938

Rodriguez, G. A., Barrett, G. M., Duff, K. E., and Hussaini, S. A. (2020). Chemogenetic attenuation of
neuronal activity in the entorhinal cortex reduces a and tau pathology in the hippocampus 18, e3000851.
doi:10.1371/journal.pbio.3000851. Publisher: Public Library of Science

Rubner, Y., Tomasi, C., and Guibas, L. J. (2000). The earth mover's distance as a metric for image retrieval

Schlesiger, M. I., Boublil, B. L., Hales, J. B., Leutgeb, J. K., and Leutgeb, S. (2018). Hippocampal global
remapping can occur without input from the medial entorhinal cortex 22, 3152–3159. doi:10.1016/j.
celrep.2018.02.082

Shapiro, M. L., Tanila, H., and Eichenbaum, H. (1997). Cues that hippocampal
place cells encode: Dynamic and hierarchical representation of local and distal
stimuli 7, 624–642. doi:10.1002/(SICI)1098-1063(1997)7:6⟨624::AID-HIPO5⟩3.0.
CO;2-E. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/%28SICI%291098-
1063%281997%297%3A6%3C624%3A%3AAID-HIPO5%3E3.0.CO%3B2-E

Sihn, D. and Kim, S.-P. (2019). A spike train distance robust to firing rate changes based on the earth
mover's distance 13

Solstad, T., Boccara, C. N., Kropff, E., Moser, M.-B., and Moser, E. I. (2008). Representation of geometric
borders in the entorhinal cortex 322, 1865–1868. doi:10.1126/science.1166466. Publisher: American
Association for the Advancement of Science

[Dataset] Sotomayor-Gómez, B., Battaglia, F. P., and Vinck, M. (2021). SpikeShip: A method for fast,
unsupervised discovery of high-dimensional neural spiking patterns. doi:10.1101/2020.06.03.131573.
Pages: 2020.06.03.131573 Section: New Results

Steffenach, H.-A., Witter, M., Moser, M.-B., and Moser, E. I. (2005). Spatial memory in the rat requires
the dorsolateral band of the entorhinal cortex 45, 301–313. doi:10.1016/j.neuron.2004.12.044

Tsao, A., Moser, M.-B., and Moser, E. I. (2013). Traces of experience in the lateral entorhinal cortex 23,
399–405. doi:10.1016/j.cub.2013.01.036

Tsodyks, M. (1999). Attractor neural network models of spatial maps in hippocampus
9, 481–489. doi:10.1002/(SICI)1098-1063(1999)9:4⟨481::AID-HIPO14⟩3.0.CO;
2-S. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/%28SICI%291098-
1063%281999%299%3A4%3C481%3A%3AAID-HIPO14%3E3.0.CO%3B2-S

Vasserstein, L. N. (1969). Markov processes over denumerable products of spaces, describing large systems
of automata

Wang, C., Chen, X., Lee, H., Deshmukh, S. S., Yoganarasimha, D., Savelli, F., et al. (2018). Egocentric
coding of external items in the lateral entorhinal cortex 362, 945–949. doi:10.1126/science.aau4940.
Publisher: American Association for the Advancement of Science

Wang, J. and Sainburg, R. L. (2005). Adaptation to visuomotor rotations remaps movement vectors,
not final positions 25, 4024–4030. doi:10.1523/JNEUROSCI.5000-04.2005. Publisher: Society for
Neuroscience Section: Behavioral/Systems/Cognitive

Wills, T. J., Lever, C., Cacucci, F., Burgess, N., and O'Keefe, J. (2005). Attractor dynamics in the
hippocampal representation of the local environment 308, 873–876. doi:10.1126/science.1108905.
Publisher: American Association for the Advancement of Science

Wilson, D. I., Langston, R. F., Schlesiger, M. I., Wagner, M., Watanabe, S., and Ainge, J. A. (2013a). Lateral
entorhinal cortex is critical for novel object-context recognition 23, 352–366. doi:10.1002/hipo.22095.
_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/hipo.22095

Wilson, D. I., Watanabe, S., Milner, H., and Ainge, J. A. (2013b). Lateral entorhinal cortex is necessary for
associative but not nonassociative recognition memory 23, 1280–1290. doi:10.1002/hipo.22165. _eprint:
https://onlinelibrary.wiley.com/doi/pdf/10.1002/hipo.22165

Wilson, M. A. and McNaughton, B. L. (1993). Dynamics of the hippocampal ensemble code for space 261,
1055–1058. doi:10.1126/science.8351520

Wood, E. R., Dudchenko, P. A., Robitsek, R. J., and Eichenbaum, H. (2000). Hippocampal neurons encode
information about different types of memory episodes occurring in the same location 27, 623–633.
doi:10.1016/S0896-6273(00)00071-4

Zhao, Q., Yang, Z., and Tao, H. (2010). Differential earth mover's distance with its applications to visual
tracking 32, 274–287. doi:10.1109/TPAMI.2008.299

FIGURE CAPTIONS

Figure 1. Identical place field translation. Stepwise horizontal linear translation of identical, overlapping place fields ($N = 17$, $\sigma = 3$) moving from the center to the right (**A, C**). EMD score is shown on the left while Pearson's r is shown on the right (top panel - EMD vs Pearson). 12 steps are shown and scores are rounded for display. Scores from remapping tested at all possible centroids along a single row on the rate map (bottom panel). EMD and Pearson's r scores tested at all possible centroids in the rate map (N*N) (**B, D**). Scores for horizontal and diagonal translations along the rate map are shown for all rows ($N = 17$) (top panel). Heatmap showing the gradient of scores for both raw and inverted EMD (left and center) and for Pearson's r (right) (bottom panel).

Figure 2. Identical grid field translation. Stepwise horizontal linear translation of identical, overlapping grid fields ($N = 3$, $\sigma = 1$) moving from the top left corner to the right and/or downwards on a rate map ($N = 17$) (**A, C**). EMD score is shown on the left while Pearson's r is shown on the right. 12 steps are shown and scores are rounded for display (top panel - EMD vs Pearson). Grid maps were sliced from a larger map with sufficient fields and bins to support N*N steps. Initial grid maps were chosen by taking a slice from the wider map. Scores from remapping tested across N*N different shifts from the initial grid map (0 to N combinations) (bottom panel). EMD and Pearson's r scores tested at N*N different centroid positions on the wider grid (**B, D**). Scores for horizontal and diagonal translations along the rate map are shown for all rows ($N = 17$) (top panel). Heatmap showing the gradient of scores for both raw and inverted EMD (left and center) and for Pearson's r (right) (bottom panel).

Figure 3. Incremental field degeneration. Stepwise nonlinear translation of overlapping fields ($N = 33$) relative to a fixed field at $\sigma = 3$. 12 steps are shown with 6 scaling down and 6 scaling up relative to the fixed field (**A**). EMD score is shown on the left while Pearson's r is shown on the right (left panel). Scores from remapping tested across a range of standard deviations for the scaling field (right panel). Incremental field degeneration for a pair of fields, non-overlapping and overlapping (**B,C**). Left panels show the stepwise degradation in the rate map due to randomly sampled normally distributed noise with varying standard deviations. Noise standard deviations are shown above the rate map plots. The distribution plots show the computed remapping score between the pair of fields for the overlapping and non-overlapping cases. Both cases have values for the EMD (red), field EMD (green) and Pearson's r scores (blue) displayed (right panels).

Figure 4. Single field localization. Field localization plots across two different noise levels (rows: low noise 0.1 and high noise 0.5). For each row in a plot, the first column shows the ratemap post added noise with padding, smoothing and normalizing. The second column shows the EMD distribution on the padded rate map with scores being relative to a fake point map with all the density placed in the bin at which the EMD score is found. The third column shows the same map to point computation for Pearson's r scores. The fourth column shows the 80th percentile scores for the EMD (red) and Pearson's r distributions (blue). The fifth column shows the top 20% firing rates in the cell. The last column (sixth) holds the extracted blobs (fields) from the padded ratemap with the centroid of each blob shown in red. The circle represents the true field centroid. The star represents the centroid computed on the peak EMD scores. The triangle is the centroid computed from the peak Pearson's r scores. The diamond is the centroid from the peak firing rates. The red dots are the centroids of a given field

Figure 5. Dual field localization. Field localization plots across two different noise levels (rows: low noise 0.1 and high noise 0.5). For each row in a plot, the first column shows the ratemap post added noise with padding, smoothing and normalizing. The second column shows the EMD distribution on the padded rate map with scores being relative to a fake point map with all the density placed in the bin at which the EMD score is found. The third column shows the same map to point computation for Pearson's r scores. The fourth column shows the 80th percentile scores for the EMD (red) and Pearson's r distributions (blue). The fifth column shows the top 20% firing rates in the cell. The last column (sixth) holds the extracted blobs (fields) from the padded ratemap with the centroid of each blob shown in red. The circle represents the true field centroid. The star represents the centroid computed on the peak EMD scores. The triangle is the centroid computed from the peak Pearson's r scores. The diamond is the centroid from the peak firing rates. The red dots are the centroids of a given field

Figure 6. Complex non-linear field remapping. Stepwise nonlinear translation of overlapping fields ($N = 33$) relative to a fixed field at $\sigma = 3$. 12 steps are shown with 6 scaled down and 6 scaled up relative to the fixed field (A). EMD score is shown on the left while Pearson's r is shown on the right (top panel). Scores from remapping tested across a range of rotation angles (bottom panel). Four corner point driven remapping with top left, top right, bottom right and bottom left tested. Fields were positioned so as to be fully encompassed by the rate map area. The first column shows the field location, four possible object/point/stimulus locations (stars), and distances from the field centroid to each of the four positions. The second column shows the whole map to whole map EMD scores with the full rate map and a fake pointmap (1 at object location, 0 everywhere else). The third shows a field restricted EMD between a field and a quadrant of multiple bins. The fourth column shows an approximation to the whole map sliced EMD using only the field and the single point object point (single point Wasserstein). The last column holds the Pearson's r scores. Heatmaps demonstrate the scores in the four possible corners.

Figure 7. EMD robustness to intensity changes. Incremental field scaling (increasing intensity) across a (17,17) ratemap with a field at the center (**A**). Field is scaled from $= 1$ to $\sigma = 6$ (left). Intensity changes are considered using whole map EMD (red), field restricted EMD (green) and a binary EMD (blue) using raw spike positions. Normalized (top right panel) and unnormalized (bottom right panel) weights are shown for both. EMD scores tested against different firing rate ratios for two identical fields (**B**). Ratios greater than 1 and less than 1 were tested.