The bigger picture: global analysis of solubilization performance of classical detergents versus new synthetic polymers utilizing shotgun proteomics

Stefan Mueller[a]*, Jan Kubicek[b], Felipe Merino[b], Philipp Hanisch[b], Barbara Maertens[b] and Jan-Wilm Lackmann[c]

[a]CECAD/ CMMC Proteomics Facility, Center for Molecular Medicine Cologne (CMMC) Universitaet Koeln, Robert-Koch-Str. 21, 50931 Köln, Germany

[b]Cube Biotech GmbH, Creative Campus Allee 12, 40789 Monheim, Germany

[c]CECAD/ CMMC Proteomics Facility, CECAD - Cluster of Excellence, Universitaet Koeln, Robert-Koch-Str. 21, 50931 Köln, Germany

*Correspondence to: stefan.mueller@uni-koeln.de

Abstract

Integral membrane proteins are critical for many cellular functions. Roughly 25% of all human genes code for membrane proteins, and about 70% of all approved drugs target them. Despite their importance, laborious and harsh purification conditions often hinder their characterization. Traditionally, they are removed from the membrane using detergents, thereby taking the proteins out of their native environment, affecting their function. Recently, a variety of synthetic polymers have been introduced, which can extract membrane proteins together with their native lipids into a so-called native nanodisc. However, they usually show lesser solubilization capacity than detergents, and their general applicability for membrane protein biochemistry is poorly understood. Here, we used Hek293 cell membrane extracts and LC-MS-based proteomics to compare the ability of nanodisc-forming polymers against state-of-the-art detergents to solubilize the membrane proteome. Our data demonstrates the general ability of synthetic co-polymers to extract membrane proteins, rivalling the efficacy of commonly used detergents. Interestingly, each class of solubilization agent presents specific solubilization profiles. Importantly, we found no correlation between efficiency and number of transmembrane domains, isoelectric point, or GRAVY score for any compound. Combined, our data show that these polymers are a versatile alternative to detergents for the biochemical and structural study of membrane proteins, functional proteomics or as components of a native lysis/solubilization buffers. Our work here represents the first attempt at a proteome-scale comparison of the efficacy of nanodisc-forming polymers. These data should serve as starting reference to researchers looking to purify membrane proteins in near native conditions.

Introduction

Integral membrane proteins perform key cellular functions, such as cell signaling, selective transport and generally they mediate the interaction between cells and their environment. Accordingly, although they represent about a quarter of all proteins encoded in the human genome (Fagerberg et al., 2010) they are the target of the majority of the currently approved drugs (Santos et al., 2017). Despite their clear relevance, they remain very poorly characterized. For instance, up until 2021, only 3% of all structures in the PDB were of membrane proteins (Le Bon et al., 2021). Many factors contribute to this, including low production yields, but perhaps the biggest challenge is that they need to be solubilized from the membrane. Traditionally, this is done using detergents, which remove the majority, if not all, lipids around the protein. Being stripped of their native environment, membrane proteins become usually unstable as lipids are known to be important co-factors for protein function and stability (Barrera et al., 2013; Bogdanov et al., 2008; Contreras et al., 2011; Laganowsky et al., 2014; Pollock et al., 2018; Saliba et al., 2015; Schmidpeter et al., 2022), and detergents can replace the lipid bilayers only to a very limited extent (Seddon et al., 2004). The choice of an appropriate detergent depends on the protein and the application. For example, quantitative and deep profiling of entire proteomes often requires either strong detergents such as sodium dodecylsulfate (SDS), sodium deoxycholate (SDC) or RapiGest™, high concentrations of chaotropic agents such as 8 M Urea or 6 M Guanidinium Hydrochloride, or mixture of them(Hughes et al., 2014; Vit et al., 2016; Wiśniewski et al., 2009). Furthermore, enrichment of membranes by ultracentrifugation followed by washing with alkaline sodium carbonate or 8 M Urea has been described to improve the identification of low abundant membrane proteins (Fujiki et al., 1982; Kongpracha et al., 2022). In the case of quantitative proteomics, highly denaturing extraction conditions are extremely helpful, since they instantly reduce intrinsic protease and phosphatase activity in extracts, disrupt supramolecular protein networks and unfold proteins making them amenable for efficient tryptic digestion. In clear contrast, structural and functional characterization requires proteins in near native state, and enzymatic activity as well molecular interactions should be preserved as far as possible. While many soluble proteins can be easily extracted in simple aqueous buffers, the addition of mild detergents such as decyl maltoside (DM), dodecyl maltoside (DDM) and lauryl maltose neopentyl glycol (LMNG) (Choy et al., 2021) is required for membrane proteins. To stabilize extracted membrane proteins, detergents must be kept above their critical micelle concentration (CMC) over the entire workflow, significantly increasing reagent costs, experiment complexity, and negatively affecting many downstream applications.

More than two decades ago, the group of Jean-Luc Popot (Tribet et al., 1996) developed a new kind of amphipathic polymers or "amphipols", capable of stabilizing integral membrane proteins. Amphipols can provide the hydrophobic environment the protein needs and bind it so tightly that no additional

amphipol or detergent is needed in downstream applications. For this reason, classical amphipols like

A8-35, have been particularly popular for single particle electron cryo-microscopy (Cryo-EM) structures

(Popot, 2018; Zoonens and Popot, 2014). Unfortunately, these polymers lack the ability to directly

solubilize proteins from the membrane, and an initial detergent extraction is required. In 2009, the

Overduin lab introduced a new class of co-polymer to tackle the challenge of solubilizing while

preserving the native lipid environment (Knowles et al., 2009). Made by repeating units of styrene and

maleic acid (SMA), SMAs were known to act as a molecular "cookie cutter" (Parmar et al., 2016) taking

integral membrane proteins together with a belt of their own native lipids out of the membrane,

forming a so-called SMALP (Styrene and Maleic acid lipid particle). To date, SMALPs have been used for

many applications, including radioligand and spectroscopic binding assays, NMR based assays, and

measurement of channel activity (see (Unger et al., 2021) for a summary). Unfortunately, SMALPs are

unstable a pH values below 6.5, precipitate in the presence of divalent ions, and interfere with protein

quantification using aromatic side chain absorption. However, in recent years, the promise of

preserving the native lipid environment of the protein has led to the development of several alternative

co-copolymers that tackled some of the shortcomings of the original SMAs (for an overview of this

rapidly evolving field see (Brown et al., 2021; Krishnarjuna and Ramamoorthy, 2022; Orekhov et al.,

2022)). Variations in composition and manufacturing processed generate polymers with distinct

capabilities. For instance, replacing styrene by diisobutylene, Diisobutylene-maleic acid (DIBMA)

produces a gentler polymer without absorption at 280 nm and higher tolerance to divalent ions

(Oluwole et al., 2017). Although less efficient than SMAs, DIBMA could successfully solubilized ß2

adrenergic receptor producing a functional protein with higher stability compared to DDM (Harwood

et al., 2021). Recently, two new polymer classes have been reported. Smith and colleagues (Smith et

al., 2020) introduced a new co-polymers made of "acrylic acid and styrene" (AASTY) and synthesized

by reversible addition-fragmentation chain-transfer. AASTYs have a much precisely controlled

molecular weight and the use of acrylic acid confers them higher tolerance to divalent ions than SMAs

(Timcenko et al., 2022). Interestingly, the group of Manuela Zoonens recently demonstrated that

replacing the aliphatic group of traditional amphipols by cyclo-alkanes (CyclAPols, known commercially

as Ultrasolute™ Amphipols) resulted in polymers with excellent solubilization characteristics, capable

of forming native nanodiscs (Marconnet et al., 2020). Moreover, CyclAPols solubilized bateriorhodopsin

shows increased thermal satiability when compared to SMAs.

Lipidomics analysis of SMAs and DIBMAs has demonstrated that both polymer classes can successfully

extract native lipid from eukaryotic and bacterial cells, although the extracted lipidic composition is

somehow dependent on the exact chemical nature of the polymer. Interestingly, the lipid composition

of SMALPs depend also on the identity of the protein (Teo et al., 2019) suggesting that these polymers

can capture local compositional variations of the membrane. For most of these polymers, high-

resolution Cryo-EM structure are available for membrane proteins solubilized from bacterial (Catalano et al., 2021; Doyle et al., 2022; Flegler et al., 2020; Higgins et al., 2021; Qiu et al., 2018; Sun et al., 2018; Swainsbury et al., 2023; Tascón et al., 2020) or eukaryotic cells (Yoder and Gouaux, 2020; Yu et al., 2021), many of which showing densities consistent with the presence of native lipids (Catalano et al., 2021; Doyle et al., 2022; Flegler et al., 2020; Qiu et al., 2018; Yoder and Gouaux, 2020). Unfortunately, while there is clear evidence that all these polymers can form native nanodiscs, most of the information about their solubilization capabilities come from selected example proteins and systematic characterization of their solubilization capacity against other methods has only recently been performed for SMA (Carlson et al., 2019; Morrison et al., 2021).

Here, we used LC-MS analysis to compare the ability of nanodisc-forming polymers and detergents to extract membrane proteins from Hek293 cells. In contrast to in-depth characterization of selected protein examples, shotgun proteomics facilitates the quantification of thousands of individual proteins in parallel and allows for more generalized observations. Testing a total of 22 extraction conditions including different types of SMAs, DIBMAs, AASTYs, and Ultrasolute™ Amphipol, and the commonly used non-ionic detergents DM, DDM and LMNG, our data represents the most comprehensive proteome-scale study on membrane protein solubilization by polymers.

Methods

Cell culture

Human embryonic kidney 293 cells (Expi293F™, Thermo Fisher) were cultured in Expi293™ Expression

medium at a cell density between 5×10^5 cells/ml and 4×10^6 cells/ml in suspension. 100 ml of a 2 x

10^6 cells/ ml culture growing in exponential phase were harvested by centrifugation and the cell pellet

was stored at – 80 °C.

Preparation of membrane pellets

The cell pellet was resuspended in 150 mM NaCl and 20 mM HEPES, pH 7.4, cells were disrupted by

sonication. After a centrifugation step at 9,000 x g the protein concentration of the supernatant was

adjusted to 0.25 mg/ml using 150 mM NaCl and 20 mM HEPES pH 7.4. The adjusted supernatant was

pipetted into 100µl aliquots, and the aliquots centrifuged at 60,000 x g for 1 hour. The supernatant was

discarded and the cell pellets consisting of approximately 25 µg of total proteins were frozen at - 80 °C.

Protein extraction and digestion

For an initial protein extraction screen SDS was tested as 5% solution in 50 mM HEPES pH 7.4. RIPA

buffer was composed of 150 mM NaCl, 1% Nonidet P-40, 0.5% deoxycholate, 0.1% SDS in 50 mM HEPES

pH 7.4. All other tested synthetic polymers and detergents were used as 2.5% solutions in 50 mM HEPES

pH 7.4 (Table 1). In a subsequent dilution experiment the extraction properties of a selected panel of

detergents and polymers were tested using five different concentrations ranging from 0.125% to 2.5%

in 50 mM HEPES pH 7.4 (Table 1). All reagent solutions and samples were used at 5° C and kept in pre-

chilled cooling racks. Either 25 µl detergent or polymer solution were added to the membrane pellets,

each containing approximately 25 µg total protein. The tubes were briefly vortexed and placed in a

Thermomixer (Eppendorf) for 16 h at 800 rpm and 5°C. After extraction, the tubes were transferred to

a pre-chilled rotor and centrifuged for 35 min at 21,000 x g and 5°C. 20 µl supernatant were carefully

transferred to new tubes without touching their bottom. 80 µl of cold acetone (-20°C) were added, the

tubes were briefly vortexed and incubated for 120 min at -20°C to precipitate proteins. Supernatants

were removed and the remaining protein pellets were washed once with 250 µl cold acetone (-20°). 25

µl 2% deoxycholate in 50 mM HEPES TrisHCl pH 8.5, 5 mM TCEP and 40 mM chloroacetamide were

added to the airdried protein pellets. Proteins were re-solubilized, reduced and S-alkylated in a single

step for 2 h at 46°C. 0.5 µg trypsin were added in 5 µl 2% deoxycholate in 50 mM TrisHCl pH 8.5 and

proteins were digested at 37°C. After 16 h, 30 µl of 5% trifluoroacetic acid were added to stop the

digest. By following this precipitation-resolubilization protocol, we removed solubilization agents and

standardize the tryptic digestion step, which if fully compatible with SDC (Lin et al., 2008). Samples

were centrifuged for 5 min at 21,000 x g and 20°C to pellet precipitated deoxycholate and 50 µl clear

supernatant were removed and transferred to fresh tubes. 20 µl were subsequently loaded on DVB-

RPS mixed mode stage tips activated with 20 µl methanol and equilibrated to 0.1% formic acid (FA).

The loaded peptides were sequentially washed with 40 µl 0.1% FA in water and 40 µl 0.1% FA in 60%

acetonitrile followed by alkaline elution with 40 µl 1% ammonium hydroxide in 60% acetonitrile.

Eluates were immediately acidified with 10 µl 10% FA and dried in a vacuum centrifuge. Sample were

re-solubilized in 10 µl 5% FA, 2% acetonitrile and stored at -20°C until LC-MS analysis.

LC-MS Data acquisition and processing

Protein digests were analyzed on a Q-Exactive Plus (Thermo Scientific) mass spectrometer that was

coupled to an EASY nLC 1200 UPLC (Thermo Scientific). Re-solubilized protein digests were loaded onto

an in-house-made pulled tip column (30 cm length, 75 µm inner diameter, filled with 2.7 µm Poroshell

EC120 C18, Agilent) which has been equilibrated in 0.1% FA in water. Peptides were separated using

increasing acetonitrile concentrations in 0.1% FA followed by a 10 min column wash with 95%

acetonitrile in 0.1 % FA. 140 min gradients were used for data dependent (DDA) acquisitions and 90

min gradients with data independent (DIA) acquisitions. The protein extraction screen experiment was

acquired with the DDA method. Briefly, MS1 survey scans were acquired from 300 to 1,750 m/z at a

resolution of 70,000 and stored in profile mode. The top 10 most abundant peptides were isolated

within a 1.8 Th window and subjected to HCD fragmentation at a normalized collision energy setting of

27%. The AGC target was set to 5e5 charges, allowing a maximum injection time of 60 ms. Product ions

were detected in the Orbitrap at a resolution of 17,500. Precursors were dynamically excluded for 25 s.

A DIA method was used for the analysis of the dilution experiment. MS1 scans were acquired from 380

m/z to 1,020 m/z at 35,000 resolution. Maximum injection time was set to 60 ms and the AGC target

to 1E6. MS2 scans starting from 250 m/z were acquired with 17,500 resolution, maximum injection

time of 60 ms and the AGC target set to 1E6. 30 x 20 m/z staggered DIA windows covered the precursor

mass range from 400 m/z – 1,000 m/z and resulted in 60 nominal 10 m/z windows after deconvolution.

For spectral library generation, aliquots of the 2.5% condition of all extraction experiments were pooled

and analyzed with six consecutive gas phase fractionation (GPF) runs covering the mass range from 400

m/z to 1,000 m/z in sequential 100 m/z segments. MS2 scans were acquired with 25 x 4 m/z staggered

windows in a 100 m/z range. All other settings were identical to the standard DIA method used for

sample acquisition. DDA and DIA data were stored as centroid.

Data analysis

DDA runs were processed with MaxQuant version 1.5.3.8 (Tyanova et al., 2016a) using default

parameter. Briefly, MS2 spectra were searched against the human canonical swissprot reference

proteome database (UP5640, downloaded 08.26.2020), including a list of common contaminants. False

discovery rates on protein and PSM level were estimated by the target-decoy approach to 1% Protein

FDR and 1% PSM FDR respectively. The minimal peptide length was set to 7 amino acids and

carbamidomethylation at cysteine residues was considered as a fixed modification. Oxidation (M) and

Acetyl (Protein N-term) were included as variable modifications. The match-between runs option was

enabled within replicate groups. Protein and gene identifiers together with iBAQ intensities were

imported into Perseus version 1.6.5.0 (Tyanova et al., 2016b) for subsequent analysis. Decoys and

potential contaminants were removed. IBAQ values were log2 transformed and identified proteins

were filtered for at least 3 out of 3 values in at least one replicate group. Principal component analysis

was calculated on the imputed group means using Perseus defaults.

All DIA raw files were deconvoluted and transformed to mzML files using the msconvert module in

Proteowizard. DIA-NN 1.8.0 (Demichev et al., 2020) was used in library free mode to generate a project

specific spectral library from a human canonical Swissprot reference proteome fasta file and the 6 GPF

runs. The settings differing from defaults were: min precursor m/z set to 400, max precursor m/z set

to 1,000, min fragment m/z set to 250, max fragment m/z set to 1,800 and precursor charge states

ranging from 2 to 4. The resulting spectral library (11,214 proteins, 11,186 genes and 139,723

precursors) was used for the subsequent analysis of the sample runs with default parameters and the

relaxed protein inference option. Using the "diann" R-package (https://github.com/vdemichev/diann-

rpackage) MaxLFQ intensities were calculated from non-normalized intensities provided in the DIA-NN

output file and were imported into Perseus. For heatmap generation, data were filtered for at least 3

valid values in total and log2 transformed MaxLFQ values were normalized to an interval from 0-1.

Gravy Scores and Isoelectric points were calculated from the human canonical swissprot database

proteome (UP5640,) using the "Peptides" R-package (https://github.com/dosorio/Peptides/). Hek293

gene expression data were downloaded from https://www.proteinatlas.org (Uhlén et al., 2015). Entries

with noise level gene expression (TPM < 1) and non-membrane proteins (TMD count = 0) were

removed. The remaining entries were considered as expected membrane protein expression

background. Mean iBAQ intensities as well as protein level TMD counts were merged to the expression

background and the percentage of entries that were also identified by the proteomics experiment was

calculated to estimate the analytical depth of the proteomic analysis.

Results and Discussion

We used LC-MS to compare a panel of nanodisc-forming polymers and commonly used detergents for their ability to extract integral transmembrane proteins from Hek293 cell membranes. In total 22 extraction conditions including different types of SMAs, DIBMAs, AASTYs, and Ultrasolute^{TM}, and the commonly used non-ionic detergents DM, DDM and LMNG were tested. Our non-detergent control was HEPES buffer. As positive control we chose SDS, SDC, and Fos-12 as a panel of very aggressive detergents, as well as the ubiquitously used RIPA buffer. While these are not suited for native extraction, their high protein solubilization capacity should provide an upper limit of detectable membrane proteins in the current experimental setup. For an overview of the compounds tested (Table 1) and their physicochemical characteristics see Table S1.

Analytical depth of our study

We could identify a total of 7,197 proteins across all runs, including 5,665 non-membrane proteins (Table S2). We attribute this to loosely membrane-interacting proteins or soluble proteins trapped in vesicles. Normally, this is dealt by washing the samples with alkaline carbonate buffers(Fujiki et al., 1982; Kongpracha et al., 2022). However, the elevated pH needed for this (> 11) is incompatible with native protein extraction. Therefore, we filtered the data for proteins with at least one TMD, leaving 1,532 membrane protein identifications. For comparison, RNA-seq experiments of Hek293 cells report 2,371 transmembrane proteins to be expressed with transcript levels greater 1 transcript per million (Protein Atlas; https://www.proteinatlas.org/). Considering this number as expression background, we estimated the analytical depth of our proteome analysis to be better than 64 % for proteins with at least one TMD. Importantly, while mRNA levels are not always good predictors with protein amounts, they should be strongly correlated for the cell growth conditions we tested (Liu et al., 2016). Indeed, comparing transcript levels with our detected proteins (Figure S1), suggests that low gene expression is the major reason for proteins missing in our data set, strongly arguing for the high quality of the dataset.

Different solubilization compounds results in unique solubilization patterns

For our initial screening round, we assessed the solubilization capacity of all compounds at a final concentration of 2.5%, or 5% in the case of SDS. To get an initial overview of their solubilization differences, we used the mean abundance of the detected membrane proteins to performed a principal component analysis (PCA). Strikingly, the first two components (~70% of the variance) reveal

that individual reagents form tight cluster with other members of their respective classes (Figure 1A).
Higher components can separate ionic from non-ionic detergents, while to a lesser extent still
preserving the separate clustering of the different polymer classes (Figure 1B)

The clear separation between detergents and polymers, as well as the separation between different
polymer classes suggest that each solubilization agent class produces its own distinctive fingerprint.
However, a relation between the individual components of the PCA and specific physicochemical
characteristics of the compounds is not immediately obvious. To our surprise, although denaturation
potential and extraction properties of ionic and non-ionic detergents are clearly different, DM, DDM
and LMNG cluster closely with the much more aggressive SDS, SDC and Fos-12. As expected, the RIPA
buffer — a mixture of SDS, SDC and Nonidet P-40 (NP-40) — is located within this cluster as well.
Digitonin does not follow this trend, and is located far away from the main detergent cluster, near SMA
502-E.

Polymers can efficiently solubilize membrane proteins

Next, we sought to understand whether the differences in solubilization profiles we observe are a
consequence of differences in solubilization efficiency or versatility. In our negative control, we
detected an average of 253 membrane proteins, 181 (71.41%) of which are single span membrane
proteins (Figure 2). In clear contrast, the membrane protein counts for the ionic detergents SDC, SDS
and Fos-12 were 1250, 1,241, and 1,280 respectively. Unexpectedly, the ionic detergents performed
similarly to Ultrasolute^{TM} Amphipol-17 and Amphipol-18 (1,270 and 1,272 membrane proteins
respectively) and were outperformed by the non-ionic detergents DM, DDM, and LMNG (membrane
protein counts of 1,297, 1,308, and 1,315 respectively); the latter being the highest in the data set
(Figure 2). We obtained intermediate efficiency for SMAs, with counts of 1,175, 1,198 and 1,066 for
SMA-200, SMA-300 and SMA-502-E respectively. AASTYs and the DIBMAs had the lowest performance
with counts ranging from 991 to 1,010 and from 897 to 969 respectively (Figure 2). Nevertheless,
DIBMAs and AASTYs are efficient solubilization agents, showing significantly higher solubilization than
the background HEPES alone. Importantly, solubilization efficiency of membrane proteins is only one
piece of the puzzle. Purification strategy, stabilization of the membrane protein and downstream
applications are equally important aspects to consider, which makes screening of various reagents
indispensable. Indeed, polymer identity or even concentration can affect all of these parameters
(Szundi et al., 2021). Absolute solubilization efficiency may also not be critical for many applications
where a near-native state is the strongest constraint and sufficient initial material is present; the
obvious example would be after overexpression. For instance, the comparably milder DIBMA has been

successfully used to determine the Cryo-EM structure of the mechanosensitive-like channel YnaI

(Flegler et al., 2020) and has been successfully used to produce highly active and stable GPCR samples

(Harwood et al., 2021).

Polymers can efficiently extract diverse classes of membrane proteins

Our PCA analysis revealed that different compound classes we tested produce distinct solubilization

patterns. Although they clearly differ in their solubilization efficiency, this cannot be the only driver as

the clustering of detergent and polymer classes remains present for higher components in the PCA

analysis (Figure 1B). We first analyzed the potential impact of the membrane protein topology on

solubilization. For this, we grouped proteins according to their number of TMDs and calculated

identification rates as the fraction of membrane proteins we identified versus what we expected from

RNA data. Identification rate differs substantially among TMD groups (Figure 3). Interestingly, the

profiles are similar for all tested extraction conditions, with scale differences clearly proportional to

efficiency. Since median protein abundancies have a major impact on identification rates in proteomic

experiments, we tested if this was behind the identification rate pattern. Indeed, mRNA level shows a

similar pattern to our detection rates (Figure 3), with a significant correlation between identification

rates and median mRNA abundance (Figure S1). In turn this suggests that protein abundance and not

preferential solubilization is behind the identification rate pattern.

A clear candidate for the differential polymer preferences would be the physicochemical properties of

the proteins and their respective transmembrane regions. For instance, hydrophobicity and charge

substantially differ across membrane proteins and could easily affect the polymer's solubilization

efficiency. To test this, we calculated GRAVY scores and isoelectric points for either the full protein

sequences or their membrane spanning regions. Interestingly, the distributions of either characteristic

for the detected membrane proteins reveal no major difference among the solubilization agents,

strongly suggesting that isoelectric points and hydrophobicity cannot explain the differential extraction

patterns and that we observe (Figure 4).

Contrary to detergents, the polymers we tested solubilize the protein together with its lipidic

microenvironment. Far from being just a scaffold for these proteins, the exact lipidic around them is

matched to their specific requirements (Lee, 2004), with each of them possessing their own lipidic

fingerprint (Corradi et al., 2018). Interestingly, SMAs have already been shown to solubilize, from the

same membrane, proteins with different local lipidic composition (Teo et al., 2019). However, early

characterization of nanodisc-forming polymers has shown that their solubilization efficiency is

somehow dependent on membrane composition (Oluwole et al., 2017; Smith et al., 2020). Combined,

this suggests that, for a specific target, differences in solubilization ability between polymers would be
partially determined by the identity of the annular lipids surrounding the membrane protein.

Optimal solubilization conditions are specific to each compound and target combination

So far, we compared the capacity of polymers and detergents to extract membrane proteins using a
standard concentration of 2.5%. Although this concentration is effective for all conditions tested, it is
relatively high. In many cases, high concentrations like these can interfere with purification, or affect
the activity of the protein (Szundi et al., 2021). In those kinds of cases where gentle solubilization of
proteins is required, it is common practice to use additives at concentrations as low as possible in order
to minimize detrimental effects. DDM and LMNG are typically used from 0.1% to 1%, whereas SMAs
have been successfully used in concentrations as low as 0.25% (Kopf et al., 2020; Pitch et al., 2021). In
order to determine useful minimal concentrations, we performed a second extraction screen with
concentrations of 0.125%, 0.25%, 0.5%, 1% and 2.5% for a selected panel of polymers and non-ionic
detergents. The raw data from this experiment can be found in Table S2. In line with the first screening,
DDM and LMNG were the most efficient compounds and reached maximum membrane protein counts
at concentrations as low as 0.5% (Figure 5A) and maximum summed intensities at 1% (Figure 5B). In
agreement with our earlier experiments, Ultrasolute™ Amphipol-17 and 18 showed very high
solubilization efficiency reaching their maximum counts at 1% and plateauing at a maximum value of
summed intensities comparable to detergents. AASTY 11-45 and 6-50 also reached a plateau for
summed intensities at concentrations above 1%, although only AASTY 11-45 does not seem to find a
plateau for the number of individual membrane proteins (Figure 5A,B). DIBMAs in general seem to find
a plateau for both quantities, although with much lower values (Figure 5A,B), arguing for a generally
lower solubilization efficiency. For all other compounds the curves imply that the concentration for
maximum solubilization has not been reached and may clearly exceed 2.5%. Combined, this shows that
polymers can be efficiently used at concentrations significantly lower than 2.5%. We would recommend
0.25 – 1% for Ultrasolute™ Amphipols and 0.5 – 1% for AASTYs, while a higher concentration would be
required for DIBMAs and SMAs. To what extent these results from endogenously expressed targets in
this study can be transferred to overexpression conditions has to be tested individually for each target.

Finally, to further dissect the versatility of these polymers we created a heatmap of the normalized
protein intensities obtained for each individual detected membrane proteins, at each different
solubilization agent concentration (Figure 6). Unsurprisingly, detergents are able to solubilize most of
the proteins in the map with high efficiency. In addition, Ultrasolute™ Amphipols and to a lesser extent
AASTY 6-50 produce a very similar patter, both in terms of concentration dependency and number and

identity of the solubilized proteins. Combined this demonstrates that nanodisc-forming polymers

present a strong alternative to detergents, especially when the lipid environment is key for protein

function.

Concluding remarks

We used LC-MS to perform label-free quantification proteomics aimed at screening native-nanodisc-

forming polymers and compared them with state-of-the-art detergents for their ability to solubilize the

Hek293 membrane proteome. We omitted other membrane-mimetic strategies — such as MSP-based

nanodisc (Bayburt et al., 2002), peptidisc (Carlson et al., 2018) and saponins (Frauenfeld et al., 2016)

— as they all require the intermediate addition of detergents, thus altering the native environment of

the proteins under investigation. The study presented here is the first major attempt to compare the

ability of a comprehensive panel of polymers at the proteome scale. As such, we believe that this study

will become a reference for researches working with different aspects of the biochemistry of

membrane proteins.

We have demonstrated that these synthetic polymers exhibit solubilization performance that can rival

those observed in commonly used detergents, but have the advantage of preserving the protein in a

near-native environment; Ultrasolute^{TM} Amphiphol 17 and 18 even outperformed the denaturing

detergent group. Considering the separate clustering observed in Figure 1, plus the patterns observed

in Figure 6, we believe that, for any given target, it is likely that at least one polymer class should provide

efficient solubilizationWe are certain being able to preserve the native membrane environment will

become and invaluable tool for membrane biochemists. As such, we anticipate that these synthetic

polymers will have a broad range of application in the future, in particular as components of native

lysis and solubilization buffers. Moreover, with the ever-increasing access the electron cryo-microscopy

we expect that these polymers will make a big impact in the structural biology of integral membrane

proteins.

Acknowledgements

All mass spectrometric analysis was performed in the CMMC/CECAD Proteomics Facility, CECAD
University of Cologne.

Data Availability

379 The mass spectrometry data included in this manuscript is available in Table S2. For ease of navigation,

380 we have also created a solubilization database with the data from the manuscript, available at

381 https://cube-biotech.com/solubilization-database

Conflict of interest disclosure

JK, BM, PH, and FM are shareholders of Cube Biotech GmbH, which produces and sells synthetic co-
polymers for membrane protein characterization. SM and JWL declare no conflict of interest.

References

Barrera, N.P., Zhou, M., Robinson, C.V., 2013. The role of lipids in defining membrane protein
interactions: insights from mass spectrometry. Trends Cell Biol. 23, 1–8.
https://doi.org/10.1016/j.tcb.2012.08.007

Bogdanov, M., Mileykovskaya, E., Dowhan, W., 2008. Lipids in the Assembly of Membrane Proteins and
Organization of Protein Supercomplexes. Subcell. Biochem. 49, 197–239. https://doi.org/10.1007/978-
1-4020-8831-5_8

Brown, C.J., Trieber, C., Overduin, M., 2021. Structural biology of endogenous membrane protein
assemblies in native nanodiscs. Curr. Opin. Struct. Biol. 69, 70–77.
https://doi.org/10.1016/j.sbi.2021.03.008

Carlson, M.L., Stacey, R.G., Young, J.W., Wason, I.S., Zhao, Z., Rattray, D.G., Scott, N., Kerr, C.H., Babu,
M., Foster, L.J., Duong Van Hoa, F., 2019. Profiling the Escherichia coli membrane protein interactome
captured in Peptidisc libraries. eLife 8, e46615. https://doi.org/10.7554/eLife.46615

Catalano, C., Ben-Hail, D., Qiu, W., Blount, P., des Georges, A., Guo, Y., 2021. Cryo-EM Structure of
Mechanosensitive Channel YnaI Using SMA2000: Challenges and Opportunities. Membranes 11, 849.
https://doi.org/10.3390/membranes11110849

Choy, B.C., Cater, R.J., Mancia, F., Pryor, E.E., 2021. A 10-year meta-analysis of membrane protein
structural biology: Detergents, membrane mimetics, and structure determination techniques. Biochim.
Biophys. Acta BBA - Biomembr. 1863, 183533. https://doi.org/10.1016/j.bbamem.2020.183533

Contreras, F.-X., Ernst, A.M., Wieland, F., Brügger, B., 2011. Specificity of Intramembrane Protein–Lipid
Interactions. Cold Spring Harb. Perspect. Biol. 3, a004705.
https://doi.org/10.1101/cshperspect.a004705

Corradi, V., Mendez-Villuendas, E., Ingólfsson, H.I., Gu, R.-X., Siuda, I., Melo, M.N., Moussatova, A.,
DeGagné, L.J., Sejdiu, B.I., Singh, G., Wassenaar, T.A., Delgado Magnero, K., Marrink, S.J., Tieleman, D.P.,
2018. Lipid–Protein Interactions Are Unique Fingerprints for Membrane Proteins. ACS Cent. Sci. 4, 709–
717. https://doi.org/10.1021/acscentsci.8b00143

Demichev, V., Messner, C.B., Vernardis, S.I., Lilley, K.S., Ralser, M., 2020. DIA-NN: neural networks and
interference correction enable deep proteome coverage in high throughput. Nat. Methods 17, 41–44.
https://doi.org/10.1038/s41592-019-0638-x

Doyle, M.T., Jimah, J.R., Dowdy, T., Ohlemacher, S.I., Larion, M., Hinshaw, J.E., Bernstein, H.D., 2022.
Cryo-EM structures reveal multiple stages of bacterial outer membrane protein folding. Cell 185, 1143-
1156.e13. https://doi.org/10.1016/j.cell.2022.02.016

Flegler, V.J., Rasmussen, A., Rao, S., Wu, N., Zenobi, R., Sansom, M.S.P., Hedrich, R., Rasmussen, T.,
Böttcher, B., 2020. The MscS-like channel YnaI has a gating mechanism based on flexible pore helices.
Proc. Natl. Acad. Sci. 117, 28754–28762. https://doi.org/10.1073/pnas.2005641117

Fujiki, Y., Hubbard, A.L., Fowler, S., Lazarow, P.B., 1982. Isolation of intracellular membranes by means
of sodium carbonate treatment: application to endoplasmic reticulum. J. Cell Biol. 93, 97–102.
https://doi.org/10.1083/jcb.93.1.97

Harwood, C.R., Sykes, D.A., Hoare, B.L., Heydenreich, F.M., Uddin, R., Poyner, D.R., Briddon, S.J.,
Veprintsev, D.B., 2021. Functional solubilization of the β2-adrenoceptor using diisobutylene maleic
acid. iScience 24. https://doi.org/10.1016/j.isci.2021.103362

Higgins, A.J., Flynn, A.J., Marconnet, A., Musgrove, L.J., Postis, V.L.G., Lippiat, J.D., Chung, C., Ceska, T.,
Zoonens, M., Sobott, F., Muench, S.P., 2021. Cycloalkane-modified amphiphilic polymers provide direct
extraction of membrane proteins for CryoEM analysis. Commun. Biol. 4, 1–9.
https://doi.org/10.1038/s42003-021-02834-3

Hughes, C.S., Foehr, S., Garfield, D.A., Furlong, E.E., Steinmetz, L.M., Krijgsveld, J., 2014. Ultrasensitive
proteome analysis using paramagnetic bead technology. Mol. Syst. Biol. 10, 757.
https://doi.org/10.15252/msb.20145625

Knowles, T.J., Finka, R., Smith, C., Lin, Y.-P., Dafforn, T., Overduin, M., 2009. Membrane Proteins
Solubilized Intact in Lipid Containing Nanoparticles Bounded by Styrene Maleic Acid Copolymer. J. Am.
Chem. Soc. 131, 7484–7485. https://doi.org/10.1021/ja810046q

Kongpracha, P., Wiriyasermkul, P., Isozumi, N., Moriyama, S., Kanai, Y., Nagamori, S., 2022. Simple But
Efficacious Enrichment of Integral Membrane Proteins and Their Interactions for In-Depth Membrane
Proteomics. Mol. Cell. Proteomics 21. https://doi.org/10.1016/j.mcpro.2022.100206

Kopf, A.H., Dörr, J.M., Koorengevel, M.C., Antoniciello, F., Jahn, H., Killian, J.A., 2020. Factors influencing
the solubilization of membrane proteins from Escherichia coli membranes by styrene–maleic acid
copolymers. Biochim. Biophys. Acta BBA - Biomembr. 1862, 183125.
https://doi.org/10.1016/j.bbamem.2019.183125

Krishnarjuna, B., Ramamoorthy, A., 2022. Detergent-Free Isolation of Membrane Proteins and
Strategies to Study Them in a Near-Native Membrane Environment. Biomolecules 12, 1076.
https://doi.org/10.3390/biom12081076

Laganowsky, A., Reading, E., Allison, T.M., Ulmschneider, M.B., Degiacomi, M.T., Baldwin, A.J., Robinson,
C.V., 2014. Membrane proteins bind lipids selectively to modulate their structure and function. Nature
510, 172–175. https://doi.org/10.1038/nature13419

Le Bon, C., Michon, B., Popot, J.-L., Zoonens, M., 2021. Amphipathic environments for determining the
structure of membrane proteins by single-particle electron cryo-microscopy. Q. Rev. Biophys. 54, e6.
https://doi.org/10.1017/S0033583521000044

Lee, A.G., 2004. How lipids affect the activities of integral membrane proteins. Biochim. Biophys. Acta
BBA - Biomembr. 1666, 62–87. https://doi.org/10.1016/j.bbamem.2004.05.012

Lin, Y., Zhou, J., Bi, D., Chen, P., Wang, X., Liang, S., 2008. Sodium-deoxycholate-assisted tryptic digestion
and identification of proteolytically resistant proteins. Anal. Biochem. 377, 259–266.
https://doi.org/10.1016/j.ab.2008.03.009

Liu, Y., Beyer, A., Aebersold, R., 2016. On the Dependency of Cellular Protein Levels on mRNA
Abundance. Cell 165, 535–550. https://doi.org/10.1016/j.cell.2016.03.014

Marconnet, A., Michon, B., Le Bon, C., Giusti, F., Tribet, C., Zoonens, M., 2020. Solubilization and
Stabilization of Membrane Proteins by Cycloalkane-Modified Amphiphilic Polymers.
Biomacromolecules 21, 3459–3467. https://doi.org/10.1021/acs.biomac.0c00929

Morrison, K.A., Heesom, K.J., Edler, K.J., Doutch, J., Price, G.J., Koumanov, F., Whitley, P., 2021.
Development of Methodology to Investigate the Surface SMALPome of Mammalian Cells. Front. Mol.
Biosci. 8.

Oluwole, A.O., Danielczak, B., Meister, A., Babalola, J.O., Vargas, C., Keller, S., 2017. Solubilization of
Membrane Proteins into Functional Lipid-Bilayer Nanodiscs Using a Diisobutylene/Maleic Acid
Copolymer. Angew. Chem. Int. Ed. 56, 1919–1924. https://doi.org/10.1002/anie.201610778

Orekhov, P.S., Bozdaganyan, M.E., Voskoboynikova, N., Mulkidjanian, A.Y., Karlova, M.G., Yudenko, A.,
Remeeva, A., Ryzhykau, Y.L., Gushchin, I., Gordeliy, V.I., Sokolova, O.S., Steinhoff, H.-J., Kirpichnikov,
M.P., Shaitan, K.V., 2022. Mechanisms of Formation, Structure, and Dynamics of Lipoprotein Discs
Stabilized by Amphiphilic Copolymers: A Comprehensive Review. Nanomaterials 12, 361.
https://doi.org/10.3390/nano12030361

Parmar, M.J., Lousa, C.D.M., Muench, S.P., Goldman, A., Postis, V.L.G., 2016. Artificial membranes for
membrane protein purification, functionality and structure studies. Biochem. Soc. Trans. 44, 877–882.
https://doi.org/10.1042/BST20160054

Pitch, S.G., Yao, W., Szundi, I., Fay, J., Chen, E., Shumate, A., Kliger, D.S., Farrens, D.L., 2021. Functional
integrity of membrane protein rhodopsin solubilized by styrene-maleic acid copolymer. Biophys. J. 120,
3508–3515. https://doi.org/10.1016/j.bpj.2021.05.008

Pollock, N.L., Lee, S.C., Patel, J.H., Gulamhussein, A.A., Rothnie, A.J., 2018. Structure and function of
membrane proteins encapsulated in a polymer-bound lipid bilayer. Biochim. Biophys. Acta BBA -
Biomembr., Beyond the Structure-Function Horizon of Membrane Proteins 1860, 809–817.
https://doi.org/10.1016/j.bbamem.2017.08.012

Popot, J.-L., 2018. Membrane Proteins in Aqueous Solutions, Biological and Medical Physics, Biomedical
Engineering. Springer International Publishing, Cham. https://doi.org/10.1007/978-3-319-73148-3

Qiu, W., Fu, Z., Xu, G.G., Grassucci, R.A., Zhang, Y., Frank, J., Hendrickson, W.A., Guo, Y., 2018. Structure
and activity of lipid bilayer within a membrane-protein transporter. Proc. Natl. Acad. Sci. 115, 12985–
12990. https://doi.org/10.1073/pnas.1812526115

Saliba, A.-E., Vonkova, I., Gavin, A.-C., 2015. The systematic analysis of protein–lipid interactions comes
of age. Nat. Rev. Mol. Cell Biol. 16, 753–761. https://doi.org/10.1038/nrm4080

Santos, R., Ursu, O., Gaulton, A., Bento, A.P., Donadi, R.S., Bologa, C.G., Karlsson, A., Al-Lazikani, B.,
Hersey, A., Oprea, T.I., Overington, J.P., 2017. A comprehensive map of molecular drug targets. Nat. Rev.
Drug Discov. 16, 19–34. https://doi.org/10.1038/nrd.2016.230

Schmidpeter, P.A.M., Wu, D., Rheinberger, J., Riegelhaupt, P.M., Tang, H., Robinson, C.V., Nimigean,
C.M., 2022. Anionic lipids unlock the gates of select ion channels in the pacemaker family. Nat. Struct.
Mol. Biol. 1–9. https://doi.org/10.1038/s41594-022-00851-2

Seddon, A.M., Curnow, P., Booth, P.J., 2004. Membrane proteins, lipids and detergents: not just a soap
opera. Biochim. Biophys. Acta BBA - Biomembr., Lipid-Protein Interactions 1666, 105–117.
https://doi.org/10.1016/j.bbamem.2004.04.011

Smith, A.A.A., Autzen, H.E., Faust, B., Mann, J.L., Muir, B.W., Howard, S., Postma, A., Spakowitz, A.J.,
Cheng, Y., Appel, E.A., 2020. Lipid Nanodiscs via Ordered Copolymers. Chem 6, 2782–2795.
https://doi.org/10.1016/j.chempr.2020.08.004

Sun, C., Benlekbir, S., Venkatakrishnan, P., Wang, Y., Hong, S., Hosler, J., Tajkhorshid, E., Rubinstein, J.L.,
Gennis, R.B., 2018. Structure of the alternative complex III in a supercomplex with cytochrome oxidase.
Nature 557, 123–126. https://doi.org/10.1038/s41586-018-0061-y

Swainsbury, D.J.K., Hawkings, F.R., Martin, E.C., Musiał, S., Salisbury, J.H., Jackson, P.J., Farmer, D.A.,
Johnson, M.P., Siebert, C.A., Hitchcock, A., Hunter, C.N., 2023. Cryo-EM structure of the four-subunit
Rhodobacter sphaeroides cytochrome bc1 complex in styrene maleic acid nanodiscs. Proc. Natl. Acad.
Sci. 120, e2217922120. https://doi.org/10.1073/pnas.2217922120

Szundi, I., Pitch, S.G., Chen, E., Farrens, D.L., Kliger, D.S., 2021. Styrene-maleic acid copolymer effects
on the function of the GPCR rhodopsin in lipid nanoparticles. Biophys. J. 120, 4337–4348.
https://doi.org/10.1016/j.bpj.2021.09.012

Tascón, I., Sousa, J.S., Corey, R.A., Mills, D.J., Griwatz, D., Aumüller, N., Mikusevic, V., Stansfeld, P.J.,
Vonck, J., Hänelt, I., 2020. Structural basis of proton-coupled potassium transport in the KUP family.
Nat. Commun. 11, 626. https://doi.org/10.1038/s41467-020-14441-7

Teo, A.C.K., Lee, S.C., Pollock, N.L., Stroud, Z., Hall, S., Thakker, A., Pitt, A.R., Dafforn, T.R., Spickett, C.M.,
Roper, D.I., 2019. Analysis of SMALP co-extracted phospholipids shows distinct membrane
environments for three classes of bacterial membrane protein. Sci. Rep. 9, 1813.
https://doi.org/10.1038/s41598-018-37962-0

Timcenko, M., Autzen, A.A.A., Autzen, H.E., 2022. Characterization of Divalent Cation Interactions with
AASTY Nanodiscs. ACS Appl. Polym. Mater. 4, 1071–1083. https://doi.org/10.1021/acsapm.1c01507

Tribet, C., Audebert, R., Popot, J.-L., 1996. Amphipols: Polymers that keep membrane proteins soluble
in aqueous solutions. Proc. Natl. Acad. Sci. 93, 15047–15050.
https://doi.org/10.1073/pnas.93.26.15047

Tyanova, S., Temu, T., Cox, J., 2016a. The MaxQuant computational platform for mass spectrometry-
based shotgun proteomics. Nat. Protoc. 11, 2301–2319. https://doi.org/10.1038/nprot.2016.136

Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M.Y., Geiger, T., Mann, M., Cox, J., 2016b. The
Perseus computational platform for comprehensive analysis of (prote)omics data. Nat. Methods 13,
731–740. https://doi.org/10.1038/nmeth.3901

Uhlén, M., Fagerberg, L., Hallström, B.M., Lindskog, C., Oksvold, P., Mardinoglu, A., Sivertsson, Å.,
Kampf, C., Sjöstedt, E., Asplund, A., Olsson, I., Edlund, K., Lundberg, E., Navani, S., Szigyarto, C.A.-K.,
Odeberg, J., Djureinovic, D., Takanen, J.O., Hober, S., Alm, T., Edqvist, P.-H., Berling, H., Tegel, H., Mulder,
J., Rockberg, J., Nilsson, P., Schwenk, J.M., Hamsten, M., von Feilitzen, K., Forsberg, M., Persson, L.,
Johansson, F., Zwahlen, M., von Heijne, G., Nielsen, J., Pontén, F., 2015. Tissue-based map of the human
proteome. Science 347, 1260419. https://doi.org/10.1126/science.1260419

Unger, L., Ronco-Campaña, A., Kitchen, P., Bill, R.M., Rothnie, A.J., 2021. Biological insights from SMA-
extracted proteins. Biochem. Soc. Trans. 49, 1349–1359. https://doi.org/10.1042/BST20201067

Vit, O., Man, P., Kadek, A., Hausner, J., Sklenar, J., Harant, K., Novak, P., Scigelova, M., Woffendin, G.,
Petrak, J., 2016. Large-scale identification of membrane proteins based on analysis of trypsin-protected
transmembrane segments. J. Proteomics 149, 15–22. https://doi.org/10.1016/j.jprot.2016.03.016

Wiśniewski, J.R., Zougman, A., Nagaraj, N., Mann, M., 2009. Universal sample preparation method for
proteome analysis. Nat. Methods 6, 359–362. https://doi.org/10.1038/nmeth.1322

Yoder, N., Gouaux, E., 2020. The His-Gly motif of acid-sensing ion channels resides in a reentrant 'loop'
implicated in gating and ion selectivity. eLife 9, e56527. https://doi.org/10.7554/eLife.56527

Yu, J., Zhu, H., Lape, R., Greiner, T., Du, J., Lü, W., Sivilotti, L., Gouaux, E., 2021. Mechanism of gating
and partial agonist action in the glycine receptor. Cell 184, 957-968.e21.
https://doi.org/10.1016/j.cell.2021.01.026

Zoonens, M., Popot, J.-L., 2014. Amphipols for Each Season. J. Membr. Biol. 247, 759–796.
https://doi.org/10.1007/s00232-014-9666-8

Figures

Figure 1. Clustering of membrane protein extraction profiles of different solubilization agents.
Principal component analysis of the membrane protein iBAQ values. The means of three replicates per
extraction condition were used. Each class of solubilization agent is shown in a different color to
highlight their clustering.

Figure 2. Cumulative number of detected membrane proteins. For each replicate and condition, the summed number of membrane protein identifications is shown.

Figure 3. Membrane protein identification rate as a function of the number transmembrane domains.
For each solubilization agent class, we calculate the identification rate as fraction of identified membrane proteins versus the total number of expressed membrane proteins in Hek293 cells. We further subdivided the proteins according to their number of transmembrane domains. For reference, we also plotted the RNA level as median number of transcripts per million in grey bars.

Figure 4. Distribution of isoelectric points and GRAVY scores for the proteins detected in our solubilization experiments. For all detected membrane proteins we calculated the isoelectric points (**A,B**) and GRAVY scores (**C,D**) using either the transmembrane regions (**A,C**) or the full protein sequence (**B,D**).

Figure 5. Membrane protein extraction efficiencies of tested solubilization agents. A. Number of unique membrane proteins detected as a function of solubilization agent concentration concentrations (0.125 %, 0.25 %, 0.5 %, 1 % and 2.5 %). **B.** Summed membrane protein intensities as a function of solubilization agent concentration.

Figure 6. Membrane protein intensities as a function of solubilization agent concentrations. For each solubilization agent, we tested five increasing concentrations (0.125 %, 0.25 %, 0.5 %, 1 % and 2.5 %) which are plotted in columns from left to right. Hepes without any additives was used as control in triplicate. Row in the heat map represent an individual proteins, which were further grouped according to their number transmembrane domains.

625 **Table 1.** Summary of the solubilization agents used in our screening experiments.

Class	Copolymer/detergent	Primary screen	Dilution screen
		2.5%	0.125 – 2.5%
AASTY	AASTY 11-55	yes	No
AASTY	AASTY 6-55	yes	Yes
AASTY	AASTY 11-50	yes	No
AASTY	AASTY 6-50	yes	Yes
AASTY	AASTY 11-45	yes	Yes
Amphipol	Ultrasolute™ Amphipol 18 (CyclAPolC8C0-50)	yes	Yes
Amphipol	Ultrasolute™ Amphipol 17 (CyclAPolC6C2-50)	yes	Yes
DIBMA	DIBMA Glucosamine	yes	No
DIBMA	DIBMA Glycerol	yes	Yes
DIBMA	DIBMA 12	yes	Yes
DIBMA	DIBMA 10	yes	Yes
SMALP	SMALP 502-E	yes	Yes
SMALP	SMALP 300	yes	Yes
SMALP	SMALP 200	yes	Yes
Neutral detergents	Digitonin-A	yes	Yes
Neutral detergents	LMNG	yes	Yes
Neutral detergents	DM	yes	No
Neutral detergents	DDM	yes	Yes
Ionic detergents	RIPA buffer (1 % Nonidet P-40, 0.5 % deoxycholate, 0.1 % SDS)	yes	No
Ionic detergents	SDS	yes, 5%	No
Ionic detergents	SDC	yes	No
Ionic detergents	Fos-12	yes	No
Negative Control	HEPES	yes	Yes

Supplementary figures

Supplementary Figure 1 Correspondence between membrane protein mRNA and protein detection
in Hek293 cells. RNA transcript counts of membrane protein coding genes were plotted against binned
log2 transformed TPM. Transcripts with TPM < 1 were removed. In blue, we show the number of
proteins we could detect as a function of their corresponding TPM value obtained from Protein Atlas
(See methods).

Supplementary Table 1. Physicochemical characteristics of all solubilization agents used.

Copolymer/detergent	Apolar subunit	Polar subunit	Subunit ratio	CMC	Mw	Charge
				% (w/v)	g/mol	
AASTY 11-55	Styrene	Acrylic acid	55:45	n.a.	11,000	Negative
AASTY 6-55	Styrene	Acrylic acid	55:45	n.a.	5,500	Negative
AASTY 11-50	Styrene	Acrylic acid	1:1	n.a.	11,000	Negative
AASTY 6-50	Styrene	Acrylic acid	1:1	n.a.	5,500	Negative
AASTY 11-45	Styrene	Acrylic acid	45:55	n.a.	11,000	Negative
CyclAPolC8C0-50 Ultrasolute Amphipol 18	Cyclooctyl acrylamide	Acrylic acid	1:1	n.a.	7,500	Negative
CyclAPolC6C2-50 Ultrasolute Amphipol 17	2-Cyclohexyl-ethyl acrylamide	Acrylic acid	1:1	n.a.	7,500	Negative
DIBMA Glucosamine	Diisobutylidene	Maleamic acid-glucosamine	1:1	n.a.	14,900	Negative
DIBMA Glycerol	Diisobutylidene	Maleamic acid-3-amino-1,2-propanediol	1:1	n.a.	12,500	Negative
DIBMA 12	Diisobutylidene	Maleic acid	1:1	n.a.	12,000	Negative
DIBMA 10	Diisobutylidene	Maleic acid	1:1	n.a.	10,000	Negative
SMA 502-E	Styrene	Maleic acid monoethyl ester	1.5:1	n.a.	7,000	Negative
SMA 300	Styrene	Maleic acid	3:1	n.a.	10,000	Negative
SMA 200	Styrene	Maleic acid	2:1	n.a.	6,500	Negative
Digitonin-A			n.a	0.2 - 0.3	1,229.31	Neutral
LMNG			n.a	0.001	1,005.19	Neutral
DM			n.a	0.087	482.56	Neutral
DDM			n.a	0.0087	510.62	Neutral
RIPA buffer			n.a	n.a.	n.a.	Some negative (SDS, cholate), some positive (Tris)
SDS			n.a	0.2	288.38	Negative (sulfate)
Fos-12			n.a	0.047	351.5	Neutral (zwitterionic)
HEPES			n.a	n.a.	238.30	Positive (two positive and one negative charges)