Multi-tissue proteomics identifies a link between satellite DNA organization and

heritable transposon repression in *Drosophila*

Ankita Chavan[1,2,3¶], Lena Skrutl[1,2¶], Federico Uliana[1,3,#], Melanie Pfister[1], Franziska Brändle[1,2],

Laszlo Tirian[4], Delora Baptista[5], Dominik Handler[4], David Burke[6], Anna Sintsova[7], Pedro

Beltrao[8], Julius Brennecke[4], Madhav Jagannathan[1,3*]

[1]Institute of Biochemistry, ETH Zürich, Zürich, Switzerland

[2]Life Sciences Zürich Graduate School, Zürich, Switzerland

[3]Bringing Materials to Life Consortium, Zürich, Switzerland

[4]Institute of Molecular Biotechnology of the Austrian Academy of Sciences, Vienna

BioCenter, Vienna, Austria

[5]Instituto Gulbenkian de Ciência, Oeiras, Portugal

[6]European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI),

Cambridge, UK

[7]Institute of Microbiology, ETH Zürich, Zürich, Switzerland

[8]Institute of Molecular Systems Biology, ETH Zürich, Zürich, Switzerland

[#]Present address – Johannes Gutenberg University Mainz, Mainz, Germany

*Corresponding author

E-mail: madhav.jagannathan@bc.biol.ethz.ch

[¶]These authors contributed equally to this work.

ABSTRACT

Non-coding satellite DNA repeats are abundant at the pericentromeric heterochromatin of

eukaryotic chromosomes. During interphase, sequence-specific DNA-binding proteins

cluster these repeats from multiple chromosomes into nuclear foci known as

chromocenters. Despite the pivotal role of chromocenters in cellular processes like genome

encapsulation and gene repression, the associated proteins remain incompletely

characterized. Here, we use two satellite DNA-binding proteins, D1 and Prod, as baits to

characterize the chromocenter-associated proteome in *Drosophila* embryos, ovaries, and

testes through quantitative mass spectrometry. We identify D1- and Prod-associated

proteins, including known heterochromatin proteins as well as proteins previously unlinked

to satellite DNA or chromocenters, thereby laying the foundation for a comprehensive

understanding of cellular functions enabled by satellite DNA repeats and their associated

proteins. Interestingly, we find that multiple components of the transposon-silencing piRNA

pathway are associated with D1 and Prod in embryos. Using genetics, transcriptomics, and

small RNA profiling, we show that flies lacking D1 during embryogenesis exhibit transposon

expression and gonadal atrophy as adults. We further demonstrate that this gonadal

atrophy can be rescued by mutating the checkpoint kinase, *Chk2*, which mediates germ

cell arrest in response to transposon mobilization. Thus, we reveal that a satellite DNA-

binding protein functions during embryogenesis to silence transposons, in a manner that is

heritable across later stages of development.

INTRODUCTION

'Heterochromatin' is classically defined as regions on eukaryotic chromosomes that remain condensed throughout the cell cycle[1]. The sequences that constitute heterochromatin mainly consist of simple non-coding tandem repeats known as satellite DNA [2] and interspersed repeats known as transposable elements (TEs) [3]. Both types of repeats can account for significant fractions of eukaryotic genomes and previous studies have suggested that they may propagate selfishly at the expense of overall organismal fitness. In addition, unrestricted expression of satellite DNA repeats and TEs can lead to adverse outcomes such as genome instability, chromosome mis-segregation and cell death[4–7]. However, accumulating evidence suggests important functions for regions enriched in these repetitive sequences. For example, satellite DNA repeats at the centromere have been proposed to function in the stable propagation of centromere identity[8,9] while pericentromeric satellite DNA repeats are thought to be important for chromosome pairing[10,11] and chromosome segregation[12,13]. Similarly, regulated activity of TEs can benefit cellular function[14] through roles in telomere maintenance[15], repeat copy number stability[16,17], genome restructuring[18] and the birth of new genes.

Within the nucleus, constitutive heterochromatin is organized into distinct domains that are separate from the more transcriptionally active euchromatin. For example, pericentromeric satellite DNA repeats from multiple chromosomes are clustered by satellite DNA-binding proteins into DNA-dense nuclear foci known as chromocenters, which typically remain transcriptionally silent[19,20]. This transcriptional silencing is orchestrated, in part, by the deposition of repressive post-translational modifications (PTMs) at target chromatin loci[19]. Once established, this repressive chromatin state can be heritably propagated, both across cell divisions within an organism as well as across generations [21–24]. Intriguingly, abrogation of heterochromatin-associated PTMs does not always perturb the highly condensed state and transcriptional inactivity of the underlying repetitive sequences[24,25].

In these instances, sequence-specific satellite DNA-binding proteins have been proposed

as a complementary mechanism that maintains DNA compaction and transcriptional

silencing at constitutive heterochromatin[24,25]. However, experimental evidence as to

whether satellite DNA-binding proteins regulate transcriptional silencing at constitutive

heterochromatin is lacking.

In *Drosophila*, we have shown that two sequence-specific DNA-binding proteins, D1 and

Prod, contribute to chromocenter formation by clustering their cognate satellite DNA

repeats [20,26,27]. Strikingly, chromocenter disruption through mutation of *D1* or *prod*

leads to defective genome encapsulation in the form of micronuclei, which is associated

with DNA damage and cell death[26,27]. However, these phenotypes occur in distinct

tissues, with male germ cells acutely sensitive to *D1* mutation while larval imaginal discs fail

to develop in the absence of Prod [26,27]. These findings point to potential tissue-specific

differences in chromocenter composition and regulation. However, the chromocenter-

associated proteome in distinct tissues remains largely unidentified since most proteomic

studies have typically used constitutive heterochromatin proteins[28–30] or centromeric

proteins[29,31,32] as baits in cultured cells.

In this study, we set out to identify chromocenter-associated proteins in multiple *Drosophila*

tissues using quantitative mass spectrometry. We used the satellite DNA-binding proteins,

D1 and Prod, as baits and identified chromocenter-associated proteins from embryos, adult

ovaries and germline stem cell (GSC)-enriched adult testes. We have identified 495 proteins

significantly enriched after affinity purification of D1 and Prod across all tissues.

Interestingly, pathway analysis indicated that the D1- and Prod-associated proteome in

embryos was enriched in DNA repair and transposon repression proteins. Following up on

these data, we have identified an unanticipated molecular relationship between satellite

DNA organization and TE repression in *Drosophila*. Taken together, our study lays the

groundwork toward a comprehensive understanding of the multi-faceted contributions of

satellite DNA repeats and their binding proteins to cellular function.

RESULTS

Establishment and validation of strains carrying GFP-tagged satellite DNA-binding proteins

We performed affinity purification coupled to mass spectrometry (AP-MS) to identify chromocenter-associated proteins in *Drosophila* embryos, ovaries and GSC-enriched testes using D1 and Prod as baits (Fig. 1A). In nuclei from these tissues, D1 and Prod foci were often found in proximity to each other (Fig. 1B), and we expected to identify common interactors between the two baits. To minimize variability between baits, we used an affinity purification strategy involving pulldown of GFP-tagged baits using GFP nanobody-coupled beads. For D1, we made use of an available strain where GFP has been inserted upstream of the endogenous locus. To validate this strain, we assessed the number of chromocenters (D1 and Prod foci per cell) in male and female GSCs as well as fertility in both sexes in comparison to a wild type reference strain (*D. melanogaster yw*). We observed that the number of D1 and Prod foci per cell in male and female GSCs was identical to the wild type (Fig. 1C, D), indicating that chromocenter formation was unperturbed in the GFP-D1 strain in these cell types. In addition, we observed that homozygous GFP-D1 males and females were fertile, although the fertility of homozygous GFP-D1 females is reduced compared to the wild type (Fig. 1E, F). For Prod, we first generated a transgenic strain where a GFP-tagged codon-optimized *Prod* was placed downstream of a 400 bp promoter sequence and inserted into the attP40 locus (Fig. S1A). The *GFP-prod* transgene fully rescued the larval lethality associated with loss-of-function *prod* mutations, *prod^{k01180}* and *prod^U* [33](Fig. S1B). Hereafter, GFP-Prod refers to the *GFP-prod* transgene in a *prod* mutant background (*p400-GFP-prod, prod^{k01180}/prod^U*) (Fig. S1B). We observed that the number of GFP-Prod foci was at wild type levels in male and female GSCs (Fig. 1C, D), suggesting that the transgene does not interfere with chromocenter formation in these cells. Like the GFP-D1 strain, GFP-Prod males were fully fertile, while GFP-Prod females were sub-fertile in comparison to the

wild type (Fig. 1E, F). However, ovaries from both the GFP-D1 and GFP-Prod strains were

morphologically similar to the control (Fig. S1C), suggesting that early stages of oogenesis

are likely unaffected in these strains. As a negative control for GFP-D1 and GFP-Prod, we

used a strain ubiquitously expressing nucleoplasmic GFP (NLS-GFP). The NLS-GFP did not

enrich at chromocenters, and we observed that the number of chromocenters was

comparable to the wild type, GFP-D1 and GFP-Prod strains (Fig. 1C, D). Therefore, the

GFP-tagged D1 and Prod strains and the NLS-GFP control strain were used to characterize

the chromocenter-associated proteome in intact *Drosophila* tissues.

Multi-tissue characterization of the chromocenter-associated proteome

First, we optimized lysis and affinity purification conditions such that our bait proteins were

effectively solubilized (Fig. S1D) and immunoprecipitated (Fig. S1E). Following this, we

affinity purified GFP-D1, GFP-Prod (and NLS-GFP as a control) from the embryo, ovary and

testis lysates and identified the associated proteins by quantitative mass spectrometry.

While we were unable to analyze Prod pulldowns specifically in GSC-enriched testes due to

poor sample quality, we identified 495 proteins that were significantly enriched in the D1

and Prod pulldowns in comparison to the control pulldown of NLS-GFP across all tissues

($log_2FC>1$ and $p<0.05$, Fig. 2A-C, Fig. S2A, Table S1-3), with these hits likely representing

both direct and indirect interactions. In embryos, we identified that D1 co-purified known

chromocenter-associated or pericentromeric heterochromatin-associated proteins, such as

Su(var)3-9[34], ADD1[28,29], HP5[28,29], mei-S332[35], Mes-4[28], Hmr[29,36,37],

Lhr[29,36], and members of the chromosome passenger complex, such as borr and

Incenp[38] (Fig. 2A, Table S1). Prod pulldown in embryos co-purified chromocenter-

associated proteins such as Hcs[39] and Saf-B[40] (Table S1). Our proteomic data also

identified previously reported D1/Prod associations such as D1-ball (embryo), D1-RfC38

(ovary), D1-Smn (ovary), D1-Parp (embryo), Prod-CG15107 (ovary) and Prod-bocks

(embryo)[41,42]. Moreover, analysis of our proteomic data with STRING-DB[43] revealed

relationships between D1 and Prod co-purified proteins. For example, Prod co-purified

components of the replication protein A (RPA complex) (Fig. S2B), while D1 co-purified

multiple members of the origin recognition complex (ORC) (Fig. S2C) and the condensin I

complex (Fig. S2D).

We also observed a substantial overlap between D1- and Prod-associated proteins (yellow

points in Fig. 2A, B, Table S1-3), with 61 hits pulled down by both baits (blue arrowheads,

Fig. 2C) in embryos and ovaries. This observation is consistent with the fact that both D1

and Prod occupy sub-domains within the larger constitutive heterochromatin domain in

nuclei. Surprisingly, only 12 proteins were co-purified by the same bait (D1 or Prod) across

different tissues (magenta arrowheads, Fig. 2C, Table S1-3). In addition, only a few proteins

such as an uncharacterized DnaJ-like chaperone, CG5504, were associated with both D1

and Prod in embryos and ovaries (Fig. 2D). One interpretation of these results is that the

protein composition of chromocenters may be tailored to cell- and tissue-specific functions

in *Drosophila*. However, we also note that the large number of unidentified peptides in AP-

MS experiments means that more targeted experiments are required to validate whether

certain proteins are indeed tissue-specific interactors of D1 and Prod.

Finally, we used AlphaFold Multimer (AFM) to predict interaction interfaces for a subset D1

and Prod interactors ($\log_2$FC>2, p<0.01) from all tissues and potentially distinguish direct

versus indirect interactions[44]. AFM-based structural modeling has significantly advanced

identification of protein interaction interfaces and has been used across a wide variety of

organisms and protein types[45]. Moreover, the AFM-derived interface predicted template

modelling (ipTM) score has been used as a standard metric to evaluate the accuracy of the

structural models. Based on previous studies[44,46,47], ipTM scores above 0.5 represent a

medium confidence model of the interaction interface while ipTM scores above 0.8

represent high confidence models. Despite both D1 and Prod containing large intrinsically

disordered regions (IDRs), which hinder the accuracy of structural predictions, we found

that 6/78 D1 interactors and 16/79 Prod interactors yielded an ipTM score above 0.5 (Table

S4). Taken together, we have identified several proteins associated with the satellite DNA-

binding proteins, D1 and Prod across *Drosophila* tissues and AFM-based structural

modelling suggests that some of these hits could be direct interactors of D1 and Prod.

DNA repair and TE repression proteins are co-purified by D1 and Prod in embryos

Manual inspection of our data revealed that D1 and Prod pulled down multiple proteins

involved in DNA repair, such as Fancd2 (D1 and Prod), Top2 (D1 only), chrac-16 (D1 and

Prod) and the RPA complex (Prod only) in embryos (Fig. 2A, Table S5). Tandem repeats like

satellite DNA are inherently vulnerable to copy number changes[48] and are known to

employ specialized mechanisms during DNA repair[49–53]. Therefore, these interactions

may contribute to repeat homeostasis, especially during the rapid early embryonic

divisions. In addition, we identified that TE repression proteins (Table S6), such as the

canonical piRNA pathway members, Piwi (P-element induced wimpy testes) (Prod only) and

Maelstrom (Mael) (D1 and Prod) were also co-purified in embryos (Fig. 2A, Table S1).

Consistently, a hypergeometric test for pathway enrichment revealed that DNA repair and

TE repression proteins were enriched in the pulldowns from embryos and to a lesser extent

in other tissues (Fig. 2E).

TE repression proteins are associated with chromocenters

Transposable elements (TEs) are typically viewed as agents of genomic instability and their

unchecked mobilization is usually associated with DNA damage, mutation, cell death and

sterility[54]. In germline tissues from flies, as well as other metazoans, genome defense

against transposons requires a dedicated small RNA pathway known as the Piwi-interacting

RNA (piRNA) pathway[55,56]. In *Drosophila*, transcription of piRNA precursors is mediated

by a protein complex comprised of Rhino, Cutoff and Deadlock (RDC complex) [57–59],

followed by Nxf3-dependent nuclear export of precursor transcripts and processing into

mature piRNAs[55,56]. Mature piRNAs are loaded onto a family of RNA-binding Argonaute

proteins including Piwi (P-element induced wimpy testis), a canonical member of the piRNA

pathway [55,56]. piRNA-bound Piwi enters the nucleus, locates genomic TE loci through

sequence complementarity and mediates TE repression through local heterochromatin

formation and chromatin compaction. Heterochromatin-associated proteins such as Hybrid

male rescue (Hmr)[36], Lethal hybrid rescue (Lhr)[36] and small ovary (Sov)[60–62] are

implicated in heterochromatin formation and transcriptional repression at transposon

loci[60]. All three proteins were co-purified by D1 in embryos in this study. In addition,

recent studies in cultured cells of embryonic origin have also identified that both Piwi and

Hmr interact with Prod [63]. These data support the notion that satellite DNA-binding

proteins may contribute to piRNA pathway function, potentially through regulating

heterochromatin stability at TE loci.

In order to validate our proteomics data, we used strains where either Piwi, which is a

canonical piRNA pathway Argonaute protein, or Sov, which is implicated in piRNA-

dependent heterochromatin formation at TEs[60–62], were tagged with GFP. Although Piwi

mediates transcriptional repression at TE loci, it is localized throughout the entire nucleus

and the GFP-Piwi signal expectedly overlapped with the heterochromatic D1 and Prod foci

in embryonic nuclei (Fig. 3A). In contrast, Sov, which is recruited to transposon loci by the

piRNA pathway protein Panoramix[60], exhibited a punctate GFP signal in embryonic

nuclei, which was frequently present at the boundaries of D1/Prod-containing

chromocenters (Fig. 3A). We observed a similar pattern for GFP-Piwi and Sov-GFP

localization in adult germ cells from males and females (Fig. S3A-D). Moreover, the D1-

interacting heterochromatin protein, Hmr, which contributes to TE repression[36], was also

observed at the chromocenter boundary (Fig. S3E, F), in agreement with recent reports

suggesting that Hmr may bridge centromeric and pericentromeric heterochromatin[29,32].

To further test the potential associations between D1/Prod and Piwi, we performed a

reciprocal AP-MS experiment using embryonic lysates from the GFP-Piwi strain (and the

NLS-GFP strain as a control) (Fig. 3B, Table S7). Piwi co-purified proteins identified as

D1/Prod interactors in embryonic lysates (red points, Fig. 3B). Strikingly, four proteins

(CG14715, Fit, CG10208 and Ugt35D1) were high confidence hits from the Piwi, D1 and

Prod pulldowns (Fig. 3C). All these proteins are relatively unstudied. CG14715 is an

uncharacterized FK506-binding protein, while Ugt35D1 is a putative UDP-

glycosyltransferase. Female independent of transformer (Fit) is a sexually dimorphic small

peptide that has been implicated in feeding control[64], while CG10208 is an

uncharacterized protein with no known domains or function. To determine how these

proteins may potentially interact with each other and with D1, Prod and Piwi, we used AFM

to model interaction interfaces between all possible pairs (Table S8). Interestingly, 8/28 of

the protein pairs had ipTM scores above 0.5 (blue and red points, Fig. 3D), with these

medium and high confidence scores highlighting interfaces that may give rise to direct

interactions. Among these, both D1 and Prod had medium confidence interfaces with Piwi,

which were mediated by contacts between disordered regions on D1 and Prod that

threaded through a central cleft in the Piwi protein (Fig. S4A). At the same time, we were

unable to detect a Piwi-D1 interaction by co-immunoprecipitation from embryonic lysates

(Fig. S4B), suggesting that this interaction may only occur in a small subset of cells (e.g.

germ cells) in embryos or that the interaction may be mediated by weak/transient

associations between disordered regions. Overall, the data from the proteomic, microscopy

and structural modeling experiments strongly hint at a functional relationship between

satellite DNA organization and the piRNA pathway.

Satellite DNA clustering into chromocenters does not rely on piRNA pathway proteins

Repressive histone post-translational modifications, such as H3K9me2/3, are enriched at

pericentromeric heterochromatin[13,25] and recent studies identifying piRNAs

complementary to complex satellite DNA repeats [65] suggest that the piRNA pathway may

promote repressive PTMs at satellite DNA repeats and facilitate chromocenter formation.

This idea was supported by prior studies showing that an interaction between Piwi and the

heterochromatin protein, HP1, contributes to heterochromatin organization in both somatic

and germ cells [66–68]. Since Piwi mutation results in a severe agametic phenotype in

ovaries and testes, we instead used mutations of aubergine, which mediates piRNA

amplification through the ping-pong mechanism[56]. In the absence of aubergine, total

piRNA levels are decreased and piRNA-free Piwi fails to enter the nucleus[69]. We observed

that aubergine mutation (aub^{HN2}/aub^{QC42}) as well as germline knockdown of aubergine

resulted in a mild de-clustering of Prod foci in female GSCs but had no effect on the

number of D1 foci in the same cells (Fig. S5A-C). These findings suggest that the piRNA

pathway does not have a prominent role in chromocenter stability in female GSCs.

Satellite DNA-binding proteins make a subtle contribution to TE repression in adult

ovaries

For an initial assessment of whether D1 and Prod contribute to TE repression, we used

established reporters of transposon activity in the adult germline (*GFP-Burdock-lacZ*) and

adult soma (*gypsy-lacZ*) of *Drosophila* ovaries [70,71]. In the gonadal soma, we used the *tj-*

gal4 driver to express RNAi against D1 in a genetic background containing the *gypsy-lacZ*

reporter. Despite effective depletion of D1 (Fig. S6A), we did not observe increased

expression of the TE reporter (*gypsy-lacZ*) based on staining for beta-galactosidase (β-Gal)

in comparison to the negative control (mCherry RNAi) (Fig. S7A). Similar to D1 depletion,

Prod depletion did not result in de-repression of *gypsy-lacZ* in the somatic follicle cells (Fig.

S6B, Fig. S7A). In contrast, RNAi-mediated depletion of Sov, which is required for TE

repression in gonadal somatic cells[61,62], resulted in β-Gal signal in the follicle cells across

multiple stages of oogenesis, indicating expression of the somatic TE reporter (Fig. S7A).

Next, we tested whether loss of D1 in the germline resulted in increased expression of the

germline TE reporter, *GFP-Burdock-lacZ*. To do so, we used a previously validated mutant

allele (*D1^{LL03310}*) in trans to a chromosomal deficiency spanning the D1 locus (*Df(3R)^{Bsc666}*) in

a background containing *GFP-Burdock-lacZ*[26]. Here, we did not observe GFP signal in

the germline cells of both *D1* mutant ovaries and the heterozygous controls (Fig. S7B).

Similarly, RNAi-mediated Prod depletion (Fig. S6C) in a background containing *GFP-*

Burdock-lacZ also did not lead to observable GFP signal in the ovaries (Fig. S7B). In

contrast, depletion of aubergine, which is required for TE repression in germ cells, resulted

in a strong germline GFP signal (Fig. S7B). These data suggest that D1 and Prod do not

contribute to the repression of *gypsy* and *Burdock* during oogenesis. In addition, previous

studies have determined that Prod depletion in adult ovaries does not alter the expression

of other TEs such as *HeT-A*, *TAHRE* and *blood* [70,71]. However, we cannot exclude from

these data that D1 and Prod contribute to the repression of other TEs.

Therefore, we performed RNA sequencing to determine transcriptome-wide effects on TE

expression. Since Chk2 arrests germ cell development in response to TE de-repression and

DNA damage[72,73], we used heterozygous and *D1* mutant ovaries in a *chk2* mutant

(*chk2^{6006}*) background to prevent developmental arrest of potential TE-expressing germ

cells. We observed that both genotypes exhibited similar gonad morphology (Fig. 4A).

Consistent with our data from the TE reporters and previous studies, expression of most TE

families was not elevated in *D1* mutant ovaries in comparison to the heterozygous control

(Fig. 4B, Table S9). One exception was the LTR retrotransposon *tirant,* which was

expressed 5-6-fold higher in *D1* mutant ovaries. However, the magnitude of TE expression

in the *D1* mutant ovaries is not in the range of what has been observed for ovaries that lack

canonical piRNA pathway proteins[59,74]. The mild increase in *tirant* expression is therefore

unlikely due to compromised piRNA pathway function.

Levels of TE-targeting piRNAs are largely unperturbed in the absence of D1

Since transposon silencing by the piRNA pathway relies on the production of piRNAs and

their loading onto the Piwi, Aub and Ago3 Argonaute proteins[56], we performed small RNA

sequencing to determine whether the Argonaute-bound piRNA pool was altered in *D1*

mutant ovaries in comparison to heterozygous control ovaries. Our data show that total

levels of mature piRNAs, normalized to miRNA reads, are not substantially affected by the

loss of D1 (Fig. 4C, D). We also did not observe a marked reduction in antisense piRNAs

complementary to TEs in *D1* mutant ovaries, including those mapping to *tirant* (Fig. 4E). Our

data indicate that the biogenesis of TE-mapping piRNAs is unaffected in the absence of D1.

Furthermore, our RNA-seq data also demonstrated that none of the 347 genes that are

known to contribute to TE silencing (Table S6) were differentially expressed in *D1* mutant

ovaries (blue points, Fig. 4F, Table S9). Altogether, our data suggest that the subtle

changes in the expression of a few TEs in *D1* mutant ovaries are unlikely due to

compromised piRNA biogenesis or altered expression of piRNA pathway components.

Rather, we propose that impaired heterochromatin stability in the *D1* mutant is the

underlying cause.

The piRNA biogenesis machinery is mislocalized to satellite DNA in the absence of D1

In contrast to TE transcripts, we observed that transcripts corresponding to the abundant

1.688g/cm^3 satellite DNA repeat (denoted as *SAR DM* and *SAR2 DM*, Fig. 4B) were

upregulated in *D1* mutant ovaries in comparison to the heterozygous control. Conversely,

transcripts mapping to the Responder (Rsp) satellite DNA repeat were downregulated in the

absence of D1 (Fig. 4B). These changes in satellite DNA transcription were correlated with

changes in piRNAs originating from these repetitive loci (Fig. 4E). We further mapped the

small RNA reads to the different monomers that make up the 1.688 g/cm^3 satellite DNA and

found a specific upregulation of the 260bp and 359bp variants (Fig. 4E).

Recent studies have demonstrated that the Rhino-Deadlock-Cutoff complex stimulates

transcription of complex satellite DNA repeats, which are then processed into mature

piRNAs [65,75]. Interestingly, the absence of a ZAD-zinc finger DNA-binding protein known

as Kipferl causes the RDC complex to mislocalize to and transcribe the 359 and Rsp

satellite DNA repeats at higher levels in ovaries[75]. Based on our observation that *D1*

mutant ovaries exhibit upregulated transcripts and piRNAs corresponding to the 359bp

satellite DNA (Fig. 4B, E), we tested whether D1 may also regulate the localization of the

RDC complex during oogenesis. To do so, we used a strain expressing GFP-tagged Cutoff

(EGFP-Cuff) to assess RDC complex localization in ovarian nurse cells, where

pericentromeric satellite DNA repeats from all chromosomes are clustered into a single

prominent DAPI-dense and D1-enriched chromocenter (arrowhead, Fig. 5A). In the EGFP-

Cuff strain, we observed that Cuff was localized to the periphery of the nurse cell

chromocenter and was also present at many sites across the nucleus, consistent with the

presence of piRNA clusters at pericentromeric heterochromatin and chromosome arms

(Fig. 5A, Fig. S8A-B). We next examined Cuff localization in *D1* mutant nurse cells in

comparison to the heterozygous control. In both the heterozygous control and *D1* mutant,

DAPI-dense chromocenters were still detected in nurse cells (yellow arrowheads, Fig. 5B).

However, we observed an approximately 3-fold increase in chromocenter size in nurse cells

lacking D1 (Fig. 5B-C), likely due to impaired satellite DNA clustering. Strikingly, this

increase in chromocenter size in *D1* mutants was accompanied by a substantial mis-

localization of Cuff to satellite DNA (Fig. 5B, D), where we observed ~6-fold more Cuff at *D1*

mutant chromocenters in comparison to the heterozygous control (Fig. 5B, D). In addition,

we observed that the expanded chromocenters in the *D1* mutant nurse cells were also

positive for Nxf3 (Fig. 5E), which normally colocalizes with Cuff at piRNA clusters and

mediates nuclear export of piRNA precursors[76,77]. Consistently, RNA *in situ* hybridization

revealed increased transcription of the 359 satellite DNA repeat at the expanded Cuff-

positive chromocenters in *D1* mutant nurse cells (Fig. 5F, G). Finally, the 359bp probe signal

was abolished following RNase A treatment (Fig. 5H), which is further indication that the

359bp satellite DNA repeat is aberrantly transcribed following loss of D1. Our data indicate

that D1 inhibits RDC localization to certain complex satellite DNA repeats in *Drosophila*

ovaries. However, unlike in *Kipferl* mutant ovaries, the RDC mislocalization in *D1* mutant

ovaries did not have a substantial effect on the levels of piRNAs targeting TEs and TE

silencing. Altogether, the TE reporter experiments, gene expression analysis and small RNA

sequencing suggest that TE repression in the germ cells and somatic cells of adult gonads

is largely independent of D1 or Prod.

Maternal and zygotic D1 contribute to TE repression in adult gonads

Although D1 and Prod appear to be dispensable for TE repression in adult gonads, our

proteomic data identified that TE repression proteins were mainly co-purified by D1 and

Prod in embryos. Thus, testing the functional relevance of these interactions required

depleting satellite DNA-binding proteins during embryogenesis. This experiment is

challenging since F_1 mutants (the progeny of heterozygous parents) contain maternal

deposits during embryogenesis (Fig. S9A-C) and protein/RNA depletion experiments can be

ineffective in early embryos[78]. However, the progeny of F_1 mutants (F_2 mutants) lack both

maternal and zygotic protein (Fig. S9A-C) and allow us to test the role of satellite DNA-

binding proteins in TE repression during embryogenesis. Since *D1* mutants are viable and

fertile (unlike *prod* mutants, which are larval lethal[33]), we chose to assess TE repression in

F_1 *D1* mutants, F_2 *D1* mutants and their associated heterozygous controls (Fig. S9A-C).

Briefly, the F_1 *D1* mutant lacks zygotically expressed D1 but retains a maternal contribution

during embryogenesis and potentially early larval development (Fig. S9A, C). In comparison,

the F_1 *D1* heterozygote contains both maternally contributed and zygotically expressed D1

(Fig. S9A, C). In the F_2 *D1* mutants, which are the progeny of F_1 *D1* mutant females, both

maternal and zygotic D1 are absent during embryogenesis and adulthood (Fig. S9B, C),

while the corresponding F_2 *D1* heterozygote lacks maternal D1 in the early embryo but

expresses D1 upon zygotic genome activation (Fig. S9B, C).

We initially assessed female gonad morphology in the above genotypes since unrestricted

transposon expression can result in arrested germ cell development and agametic

ovaries[4,79]. Strikingly, we observed that F_2 *D1* mutant females predominantly exhibited

agametic ovaries (Fig. 6A), with the vast majority of ovarioles either lacking any discernible

Vasa-positive germ cells or containing only undifferentiated Vasa-positive germ cells, which

did not develop into cysts (Fig. 6B, C). Moreover, the failure of germline development and

agametic ovaries in the F2 *D1* mutant strongly correlated with loss of female fertility (Fig.

6D). Interestingly, the presence of either maternal or zygotic D1 was sufficient to rescue this

phenotype, as gonad morphology and early germ cell development of both the F_1 *D1*

mutant females (maternal D1 only) and the F_2 *D1* heterozygote females (zygotic D1 only)

were mostly unaffected (Fig. 6A-C), even though both genotypes exhibited more variable

fertility (Fig. 6D). We next tested whether certain transposons were expressed in F_2 *D1*

mutant ovaries using single molecule RNA *in situ* hybridization (smFISH). Here, we

observed that the *copia* retrotransposon, but not other TEs such as *burdock* or *gypsy*, was

highly expressed in ~70% of early germ cells from F_2 *D1* mutant ovaries (Fig. 6E-G). In

comparison, *copia* expression was not elevated in early germ cells from the *yw* wild type

strain or in the F_1 *D1* mutant (Fig. 6A). While ~90% of the *aubergine* mutant germaria

(positive control) contained *copia* expressing germ cells (Fig. 6E, F), this expression was

mostly restricted to differentiated germ cell cysts; undifferentiated *aubergine* mutant germ

cells (GSCs and cystoblasts, arrowheads, Fig. 6E) exhibited a markedly lower level of *copia*

expression (Fig. 6G). Thus, the high level of transposon expression, occurring in the early

germ cells of F_2 *D1* mutants, may contribute to the loss of GSCs or block their

differentiation, ultimately leading to agametic gonads. Consistently, we found that *chk2*

mutation rescues the agametic gonads in the F_2 *D1* mutant (Fig. 6H, I). Together, our data

reveal embryogenesis (or potentially early larval development) as a critical period when the

presence of maternal and/or zygotic D1 is sufficient for establishing TE silencing, which can

then be heritably maintained throughout development (Fig. 7).

Discussion

In this study, we have characterized the chromocenter-associated proteome across *Drosophila* embryos, ovaries and GSC-enriched testes using two known satellite DNA-binding proteins, D1 and Prod, as baits. We have identified 495 potential interactions across three different tissues encompassing known heterochromatin-associated proteins, DNA repair enzymes, proteins involved in transposon repression and a host of uncharacterized proteins. Interestingly, we found a tissue-specific signature for the chromocenter proteome, with D1 and Prod associating with many common proteins per tissue and relatively few common proteins per bait across the tissues. This finding suggests that the tissue-context or cellular function may be a primary determinant of the chromocenter proteome. In addition, our structural predictions using AFM highlight potential interfaces that may mediate direct interactions between D1, Prod and their associated proteins. We suggest that these high-scoring hits represent promising avenues of research that will expand our knowledge of satellite DNA biology and chromocenter function.

Since islands of TE are often embedded within vast tracts of satellite DNA[3,17,80], we were intrigued by the hits from the piRNA pathway, which were predominantly identified in embryos. Here, we observed that the absence of D1 during embryogenesis (the F_2 *D1* mutant) led to a striking elevation of TE expression and Chk2-dependent tissue atrophy in adult ovaries. Similarly deleterious phenotypes were not observed in adult ovaries, when either only maternally deposited D1 (the F_1 *D1* mutant) or only zygotically expressed D1 (the F_2 *D1* heterozygote) were present during embryogenesis. These data suggest that the presence of D1 throughout development is not *per se* required for TE repression. Rather, our findings point to embryogenesis as the likely window of time during which D1 contributes to heritable TE silencing in germ cells. They also explain why approaches such as RNAi-mediated knockdown or mutation in the F_1 generation, where D1 likely persists

during embryogenesis, do not result in TE expression. Our results also echo previous

observations that embryonic Piwi is important for heterochromatin-dependent gene

silencing in adults[68], although it should also be noted that the role of embryonic Piwi in

transposon silencing is more limited[81,82].

How could a satellite DNA-binding protein like D1 contribute to heritable silencing at TE

loci? While the D1-dependent clustering and compaction of satellite DNA repeats may

shield satellite DNA-embedded TE loci from transcription factors, this alone does not

explain how TEs remain silenced throughout development in the absence of D1 (e.g. the F_1

D1 mutant). Rather, we propose that biophysical properties associated with the

condensation of satellite DNA repeats into chromocenters could play an important role. For

example, the intrinsically disordered regions (IDRs) present in D1 and Prod[83], could

facilitate the selective partitioning of client proteins into chromocenters. Potential client

proteins could include piRNA-dependent heterochromatin proteins such as Hmr, Lhr and

Sov, canonical piRNA pathway proteins such as Piwi and Mael or other D1/Prod-interacting

heterochromatin proteins identified in this study. In essence, satellite DNA-binding proteins

may foster an environment at satellite DNA-embedded TE loci during embryogenesis that

facilitates the deposition of repressive PTMs and ensures heritable transcriptional

repression.

In summary, our study has characterized the chromocenter-associated proteome across

three *Drosophila* tissues. We have revealed an unanticipated relationship between satellite

DNA organization and heritable transposon repression. The as-of-yet uncharacterized hits

identified in this study establish a path toward revealing the full scope of satellite DNA and

chromocenter function in *Drosophila* as well as other eukaryotes.

Materials and Methods

Fly husbandry and strains. All fly stocks were raised on standard Bloomington medium at 25°C unless otherwise indicated. *D. melanogaster y w* was used as a wild type stock. *prod^U* (BDSC42686), *GFP-D1* (BDSC50850), *Ubi-NLS-GFP* (BDSC5826, BDSC5826), *Df(3R)^Bsc666* (BDSC26518), *aub^QC42* (BDSC4968), *aub^HN2* (BDSC8517), *piwi^1* (BDSC43637), *piwi^2* (BDSC43319), *UAS-Dcr-2* (BDSC24650), *mCherry ^RNAi* (BDSC35785) and *aub ^RNAi* (BDSC39026) were obtained from the Bloomington *Drosophila* stock center. *D1^LL03310* (DGRC140754) and *FRT42D prod^k08810* (DGRC111248) were obtained from the Kyoto stock center. *GFP-Burdock-lacZ* (v313212), *tj-gal4, gypsy-lacZ* (v313222), *GFP-Piwi* (v313321), *EGFP-Cuff* (v313269), *D1 ^RNAi* (v101946) and *Prod ^RNAi* (v106593) were obtained from the Vienna Drosophila Resource Center. *Hmr-HA*[36] was a gift from Daniel Barbash. Sov-GFP (*sov^def1*; *Dp(1;2)^FF025056*)[61] was a gift from Ferenc Jankovics. *nos-gal4* (second chromosome insertion) and *chk2^6006* were gifts from Yukiko Yamashita. *nos-gal4* (third chromosome insertion)[84] and UAS-Upd[85] have been previously described.

Construction of the GFP-Prod strain. First, a 400bp promoter upstream of the Prod start site from *D. melanogaster* was PCR amplified using the following primer pair, GATCAAGCTTCTGTTGTTATGCATATCGTTC and GATCGAATTCCCGGGTATCCTTGCTC and subcloned into the HindIII and EcoRI sites on *pUASt-GFP-attB*, replacing the UAS sequence. Subsequently, a codon-optimized Prod sequence was inserted downstream of the GFP, and transgenic flies were generated using PhiC31 integrase-mediated transgenesis into the *attP40* site (BestGene). The GFP-Prod allele was generated by recombining the *p400-GFP-Prod* transgene with *prod^k08810*. Two virgin *prod^U* females were crossed to single tester male carrying either the *p400-GFP-Prod*, *prod^k08810* allele or the *prod^k08810* allele as a control in vials at 25°C. The percent of trans-heterozygous *prod* mutant flies was quantified in each replicate.

Fertility assay. For male fertility assays, a single tester male was maintained with two virgin

y w females in vials for 7d and the number of resulting progenies was counted until 20 days

post setup. For female fertility assays, a single tester female was maintained with two *y w*

males in vials for 7d and the number of resulting progenies was counted until 20 days post

setup. Vials that contained any deceased parent flies were omitted from the analysis.

Immunofluorescence staining. Immunofluorescence staining of *Drosophila* tissues was

performed as follows. For fixation, adult tissues were dissected in PBS, transferred to 4%

formaldehyde in PBS and fixed for 30 minutes. Fixation of *Drosophila* embryos was

performed using a standard protocol. Briefly, collected embryos were moved to glass

scintillation vials and washed three times with 1x PBS, dechorionated using 3 ml of 50%

bleach solution for 90 secs by vigorous shaking and washed again in 1x PBS five times to

remove the bleach. Dechorionated embryos were fixed using a heptane: 4%

paraformaldehyde (PFA) (1:1) solution by vigorously shaking for 20 minutes. Following

fixation, the bottom aqueous phase was removed and ice cold methanol was added to

devitellinize the fixed embryos. After this step, embryos were washed three times in ice-cold

methanol and stored in methanol at -20°C for further use. Prior to immunofluorescence

staining, embryos were rehydrated using serial washes of methanol: PBS-T at 4:1, 1:1 and

1:4 volumes and followed by one wash in 1x PBS-T for 5 min each. For immunostaining,

fixed tissues were washed in 1x PBS-T (PBS containing 0.1% Triton-X) for at least 60

minutes, followed by incubation with primary antibody in 3% bovine serum albumin (BSA) in

PBS-T at 4°C overnight. Samples were washed three times in PBS-T, incubated with

secondary antibody in 3% BSA in PBS-T at 4°C overnight, washed as above, and mounted

in VECTASHIELD with DAPI (Vector Labs). Fluorescent images were taken using a Leica

TCS SP8 confocal microscope with 63x oil-immersion objectives (NA=1.4). Images were

processed using Adobe Photoshop or ImageJ.

RNA in situ hybridization. RNA *in situ* hybridization on ovaries was performed as previously described[86]. Briefly, 3-4 ovaries were dissected in RNase-free 1x PBS and fixed in 3.7% formaldehyde in PBS for 30 minutes on a nutator with gentle shaking. Following fixation, samples were washed with RNase-free 1x PBS three times and subsequently incubated in ice-cold ethanol at 4°C overnight on a nutator. The next day, samples were washed with RNase-free wash buffer (2XSSC, 10% formamide) for 3 minutes at room temperature and subsequently incubated with the hybridization mix (50-125 nM probes, 2x SSC, 10% dextran sulfate, 1g/L *E.coli* tRNA, 2mM vanadyl ribonucleoside complex, 0.5% Rnase-free BSA and 10% deionized formamide) at 37°C overnight. Samples were then washed twice (wash buffer) at 37°C for 30 minutes each and mounted in VECTASHIELD with DAPI (Vector Labs). The following probe was used to hybridize to the 359bp satellite plus strand (AGGATTTAGGGAAATTAATTTTTGGATCAATTTTCGCATTTTTTGTAAG). Stellaris RNA FISH probes against the full-length Burdock consensus sequence were synthesized (LGC Biosearch technologies). Stellaris RNA FISH probes targeting Het-A and Copia were gifts from Yukiko Yamashita. An oligo-dT probe was used to hybridize to polyadenylated RNA. Fluorescent images were taken using a Leica TCS SP8 confocal microscope with 63x oil-immersion objectives (NA=1.4). Images were processed using Adobe Photoshop.

Image Analysis. All image analysis was performed using ImageJ. For the quantitation of relative chromocenter size, we demarcated a region of interest (ROI^N) encompassing the entire nucleus as well as second ROI encompassing only the chromocenter (ROI^C) for each nurse cell (stages 4-6) analyzed. ROI^C/ROI^N equals the relative chromocenter size. For enrichment of EGFP-Cuff at chromocenters, we estimated the total EGFP-Cuff signal in ROI^C and divided it by the total EGFP-Cuff signal in ROI^N to give the percentage of EGFP-Cuff at chromocenters. For the normalized estimation of Copia expression, we estimated the average intensity of nuclear Copia (ROI^{Copia}) and normalized this value to the three

cytoplasmic ROIs that estimated the average intensity of total mRNA based on the oligo-dT

probe. For the normalized estimation of 359 transcripts at chromocenters, we estimated the

average intensity of 359 probe signal at chromocenters (ROI^C) for each nurse cell and

normalized this value to a cytoplasmic ROI that have background fluorescence.

Isolation of nuclei from embryos. Flies reared in population cages were allowed to lay

eggs on fresh yeast paste for ~18 hours. Embryos were collected using a set of sieves,

washed with 1X PBS and dechorionated using 50% bleach. Next, embryos were moved to a

dounce homogenizer containing light nuclear buffer (15 mM Tris-HCl pH 7.4, 60 mM KCl,

5 mM MgCl$_2$, 15 mM NaCl2 and 1 M sucrose) supplemented with 1x protease inhibitor

cocktail (PIC, Roche,11697498001)and 1mM PMSF. The homogenate was then incubated

on ice for 5 minutes and filtered through 2 layers of Miracloth tissue (Calbiochem). The

filtered homogenate was then centrifuged for 20 min at 480 rpm at 4°C to pellet any debris.

The supernatant containing nuclei was transferred into a fresh tube and the nuclei were

pelleted at 4000 g for 30 min at 4°C. The pellet was then resuspended in 1 ml light nuclear

buffer containing 0.1 mM EDTA and the nuclei were again centrifuged at 16000 g for 5

minutes before lysis.

Tissue lysis. Nuclei isolated from embryos were lysed for 90 min at 4°C with extraction

buffer (15 mM Tris-HCl pH 7.4, 60 mM KCl, 5 mM MgCl$_2$, 450 mM NaCl2, 1% Triton-X

[vol/vol], 500 units of benzonase) 1X PIC and 5 μl/ml PMSF). The resulting lysate was

separated from the insoluble chromatin by centrifugation for 20 min at 14000 g at 4°C. For

each ovary replicate, ~ 200 pairs were dissected from <1d old flies, flash frozen with liquid

nitrogen and stored at -80°C prior to lysis. 500µl of ice-cold lysis buffer (450 mM NaCl,

50mM Tris-HCl (pH-7.5), 5% glycerol, 1mM MgCl2, 1mM EDTA, 0.2% NP40, 0.5mM DTT,

1x PIC, 5 μl/ml PMSF) was added to frozen ovaries, which were then homogenized with a

motorized pestle until complete breakdown of the tissue. After homogenization, samples

were incubated with benzonase (Sigma Aldrich, 2 µl/500 µl of lysate) for 90 minutes at 4°C

on a rotating wheel. Following this, samples were first centrifuged for 3 min at 10000 g at

4°C to precipitate the debris from the tissues, homogenized again for 30 secs and

centrifuged for 5 min at ~18700 g at 4°C. The supernatant (without the surface lipid layer)

was transferred to a new 1.5 ml tube and centrifuged again for 5 min at ~18700g prior to

use. For each testis replicate, ~100 pairs of GSC-enriched (nos-gal4 > UAS-Upd) testis

tumors were dissected, flash frozen with liquid nitrogen and stored at -80°C prior to lysis.

300 µl RIPA lysis buffer [450 mM NaCl, 0.5 % NP-40 [vol/vol], 0.5 % Sodium deoxycholate

[mass/vol], 0.1 % SDS, 50 mM Tris pH 7.4] containing 1x protease inhibitor cocktail (PIC)

(Roche,11697498001) and 5 µl/ml PMSF (Sigma-Aldrich, P7626) was added to frozen

testes for 10 min. Tissue disruption was performed by applying 20 strokes with the tight

pestle in a 1 ml dounce homogenizer, the samples were incubated for another 20 minutes on

ice following which benzonase was added for 1 hour at 10°C. Homogenized lysates were

centrifuged at 14000 g for 20 min at 4°C prior to use. GFP-Piwi and NLS-GFP embryos were

collected as indicated in the previous section, dechorionated, flash frozen with liquid nitrogen

and stored at -80°C prior to lysis. Embryo lysis for these samples was performed in an

identical manner to the ovary lysis above.

Affinity purification and mass spectrometry. Affinity purification was performed using

anti-GFP nanobody coupled magnetic beads (~ 30µl slurry) (Chromotek AB_2631358).

Briefly, beads were collected using a magnetic rack, washed three times using the lysis

buffer and incubated with the lysate for 90 minutes on a rotating wheel at 4°C. After the

incubation, beads were washed three times for 5 minutes with a wash buffer (450 mM NaCl,

50mM Tris-HCl (pH-7.5), 1mM MgCl2, 1 mM EDTA, 0.5mM DTT) at room temperature.

Washed beads containing the protein complexes were subjected to on-beads proteolysis.

Samples were transferred into a 10kDa molecular weight cutoff spin column (Vivacon 500,

Sartorious), following the FASP protocol[87]. Beads in solution were dried, denaturated (8M

Urea), reduced (5mM TCEP, 30min 37C) and alkylated (10mM Iodoacetamide, 30min 37C).

Beads were then washed three times with 50mM ammonium bicarbonate (250µl). During

the buffer exchange, samples were centrifuged at 10000g. Subsequently, samples were

proteolyzed with 0.5µg of Trypsin (Promega, sequencing grade) for 16h at 37°C. The

proteolysis was quenched with 5% formic acid and peptides were subjected to C18

cleanup (BioPureSPN PROTO 300 C18, Nest group), following the manufacturer's

procedure. The eluted peptides were then dried using a speedvac and resuspended in 20µl

of 2% acetonitrile and 0.1% formic acid.

Mass spectrometry data acquisition. The MS datasets were acquired with two different

instrumental setups. The first setup (A) was utilized for the D1, Prod and NLS-GFP samples

from embryos, ovaries and testes. The second setup (B) was employed for acquiring the

Piwi and NLS-GFP samples from embryos. In setup A, LC-MS/MS acquisition was

performed on an Orbitrap QExactive+ mass spectrometer (ThermoFisher) coupled to an

EASY-nLC-1000 liquid chromatography system (ThermoFisher). Peptides were separated

using a reverse phase column (75µm ID x 400mm New Objective, in-house packed with

ReproSil Gold 120 C18, 1.9 µm, Dr. Maisch GmbH) across a gradient from 3 to 25% in 170

minutes and from 25 to 40% in 10 minutes (buffer A: 0.1% (v/v) formic acid; buffer B: 0.1%

(v/v) formic acid, 95% (v/v) acetonitrile). The DDA data acquisition mode was set to perform

one MS1 scan followed by a maximum of 20 scans for the top 20 most intense peptides

with MS1 scans (R=70'000 at 400m/z, AGC=1e6 and maximum IT=100 ms), HCD

fragmentation (NCE=25%), isolation windows (1.8 m/z) and MS2 scans (R=17'500 at

400m/z, AGC=1e5 and maximum IT=55ms). A dynamic exclusion of 30s was applied and

charge states lower than two and higher than seven were rejected for the isolation. In setup

B, LC-MS/MS acquisition was performed on an Orbitrap Exploris480 mass spectrometer

(ThermoFisher) coupled to an Vanquish Neo liquid chromatography system (ThermoFisher).

Peptides were separated using a reverse phase column (75µm ID x 400mm New Objective,

in-house packed with ReproSil Gold 120 C18, 1.9 μm, Dr. Maisch GmbH) across a linear

gradient from 7 to 35% in 120 minutes (buffer A: 0.1% (v/v) formic acid; buffer B: 0.1% (v/v)

formic acid, 95% (v/v) acetonitrile). The DDA data acquisition mode was set to perform one

MS1 scan followed by a maximum of 20 scans for the top 20 most intense peptides with

MS1 scans (R=60'000 at 400m/z, AGC=1e6 and auto maximum IT), HCD fragmentation

(NCE=28%), isolation windows (1.4 m/z) and MS2 scans (R=15'000 at 400m/z, AGC=2e5

and auto maximum IT). A dynamic exclusion of 25s was applied and charge states lower

than two and higher than six were rejected for the isolation.

Mass spectrometry data analysis. Acquired spectra were analyzed using the MaxQuant

software package version 1.5.2.8 embedded with the Andromeda search engine against

Drosophila proteome reference dataset (http://www.uniprot.org/, downloaded on

18.01.2021, 22'044 proteins including not reviewed proteins) extended with reverse decoy

sequences. The search parameters were set to include only full tryptic peptides, maximum

two missed cleavage, carbamidomethyl as static peptide modification, oxidation (M) and

deamidation (N-ter) as variable modification and "match between runs" option. The MS and

MS/MS mass tolerance was set to 10ppm. False discovery rate of <1% was used for

protein identification. Protein abundance was determined from the intensity of the top two

unique peptides. Intensity values were median normalized and imputed using random

sampling from a normal distribution-generated 1% less intense values. Statistical analysis

was performed using unpaired two-sided t-test. Hits identified from the differential analysis

between the bait pulldown versus the NLS-GFP control, with $\log_2 FC$ >1 and p-value < 0.05,

were considered as interacting proteins. Surprisingly, Prod spectra were not always reliably

identified in certain samples, even when used as bait. For pathway enrichment analysis, we

first generated a list of *Drosophila* DNA repair proteins (GO:0006281, Table S4) and a

custom list of transposon repression proteins based on the literature (Table S5) and used a

hypergeometric test to calculate the enrichment of proteins belonging to these lists in the

D1/Prod pulldowns.

Structural predictions using Alphafold2. To predict the structures of binary protein-protein

interactions, we used version 2.3.2 of Alphafold2, which contains updated parameters for

multimeric protein-protein interactions[44]. Default values for all parameters and for

generating multiple sequence alignments were used. To rank the models according to a

confidence metric, we used the ipTM score, which has been show to distinguish interacting

from non-interacting proteins[88]. Structural models are available upon request.

Co-immunoprecipitation and western blotting. For the co-immunoprecipitation, 125-

250mg of frozen embryos were thawed on ice and lysed in an identical manner to the ovary

lysis protocol detailed above. ~50μg of lysate was collected as input while ~970 μg of lysate

at a concentration of 2mg/ml was incubated with anti-GFP nanobody coupled beads (~ 50μl

slurry) (Chromotek AB_2631358) for 90 minutes at 4°C. After the incubation, beads were

washed three times for 5 minutes with a wash buffer (450 mM NaCl, 50mM Tris-HCl pH 7.5,

1mM MgCl2, 1 mM EDTA, 0.5mM DTT) at room temperature. Protein complexes bound to

the beads were eluted in 80μl of SDS sample buffer by heating at 95°C for 5 minutes. Input

samples were also brought to a final volume of 80μl by adding 30μl of SDS sample buffer.

Equal volumes of input and IP samples were loaded on a 4-12% SDS gradient gel (Sigma

Aldrich). Immunoprecipitated proteins were detected using mouse anti-GFP (1:1000, Roche

1181 4460001), guinea pig anti-D1 (1:1000, [26]) and rabbit anti-histone H3 (1:1000, Abcam

ab1791). Secondary antibodies used were Alexa Fluor Plus 800 donkey anti-mouse

(1:10,000, Invitrogen) and donkey anti-guinea pig IR dye 680 (1:10,000, Li-Cor). Western

blots were imaged using the Odyssey DLx fluorescence imager (Li-Cor).

RNA sequencing and data analysis. For each sample, approximately 5 pairs of ovaries from 4 day old females were dissected in RNase-free 1x PBS and flash frozen in liquid nitrogen until RNA extraction. RNA extraction for each replicate was performed using the RNeasy RNA extraction kit (Qiagen). Samples were treated with DNase post RNA extraction and purified using an RNA purification kit (Promega). RNA concentrations were assessed using a Nanodrop as well as a Qubit RNA analyzer for sample quality and RIN scores. rRNA-depleted libraries were prepared and sequenced by Novogene on an NovaSeq 6000 (Illumina) using paired-end 150 bp sequencing. The resulting raw reads were cleaned by removing adaptor sequences, low-quality-end trimming, and removal of low-quality reads using BBTools v 38.18 (Bushnell, B. *BBMap*. Available from: https://sourceforge.net/projects/bbmap/). The exact commands used for quality control can be found on the Methods in Microbiomics webpage (Sunagawa, S. *Data Preprocessing — Methods in Microbiomics 0.0.1 documentation*. Available from: https://methods-in-microbiomics.readthedocs.io/en/latest/preprocessing/preprocessing.html). The quality-controlled reads were aligned against BDGP6.32 using STAR aligner[89]. Transcript and transposon abundances were quantified using Tetranscripts[90]. Differential gene expression analysis was performed using Bioconductor R package DESeq2 v 1.37.4[91].

Small RNA sequencing and data analysis. Small RNA cloning was performed as previously described[75]. In brief, ~5 pairs of ovaries from three biological replicates per genotype were lysed and Argonaute-sRNA complexes were isolated using TraPR ion exchange spin columns. sRNAs were subsequently purified using acidic phenol. 3′ adaptors containing six random nucleotides plus a 5 nt barcode on their 5′ end and 5′ adaptors containing four random nucleotides at their 3′ end were subsequently ligated to the small RNAs before reverse transcription, PCR amplification, and sequencing on an Illumina NovaSeqSP instrument. Raw reads were trimmed for linker sequences, barcodes and the

4/6 random nucleotides before mapping to the *D. melanogaster* genome (dm6), using Bowtie

(version.1.3.0, settings: -f -v 3 -a --best --strata --sam) with 0 mismatches allowed. Genome-

mapping reads were crossreferenced with Flybase genome annotations (r6.40) using

BEDTools to allow the removal of reads mapping to rRNA, tRNA, snRNA, snoRNA loci, and

the mitochondrial genome. For TE mappings, all genome mappers were used allowing no

mismatches. Reads mapping to multiple elements were assigned to the best match. Reads

mapping equally well to multiple positions were randomly distributed. Libraries were

normalized to 1 million sequenced microRNA reads. For calculation of piRNAs mapping to

TEs, only antisense piRNAs were considered, and counts were normalized to TE length.

Data Availability. The mass spectrometry data have been deposited to the

ProteomeXchange Consortium via the PRIDE partner repository with the following dataset

identifiers: PXD044367 (IP-MS of D1 from Testis), PXD043237 (IP-MS of Piwi from Embryo),

PXD043236 (IP-MS of Prod and D1 in Ovary) and PXD043234 (IP-MS of Prod and D1 in

Embryo). We have also generated a web app to facilitate browsing of the proteomics data

(https://d1-prod-interactors-bcfffefc8a0d.herokuapp.com/).

References

References

1. Heitz E. Das Heterochromatin der Moose. I. Jb wiss Bot. 1928;69: 762–818.

2. Pardue ML, Gall JG. Chromosomal localization of mouse satellite DNA. Science.
1970;168: 1356–8.

3. Pimpinelli S, Berloco M, Fanti L, Dimitri P, Bonaccorsi S, Marchetti E, et al.
Transposable elements are stable structural components of Drosophila melanogaster
heterochromatin. Proc Natl Acad Sci U S A. 1995;92: 3804–8.

4. Kidwell MG, Kidwell JF, Sved JA. Hybrid Dysgenesis in DROSOPHILA
MELANOGASTER: A Syndrome of Aberrant Traits Including Mutation, Sterility and
Male Recombination. Genetics. 1977;86: 813–33.

5. Orsi GA, Joyce EF, Couble P, McKim KS, Loppin B. Drosophila I-R hybrid dysgenesis
is associated with catastrophic meiosis and abnormal zygote formation. Journal of Cell
Science. 2010;123: 3515–3524. doi:10.1242/jcs.073890

6. Zhu Q, Hoong N, Aslanian A, Hara T, Benner C, Heinz S, et al. Heterochromatin-
Encoded Satellite RNAs Induce Breast Cancer. Mol Cell. 2018;70: 842-853 e7.
doi:10.1016/j.molcel.2018.04.023

7. Ting DT, Lipson D, Paul S, Brannigan BW, Akhavanfard S, Coffman EJ, et al. Aberrant
Overexpression of Satellite Repeats in Pancreatic and Other Epithelial Cancers.
Science. 2011;331: 593–596. doi:10.1126/science.1200801

8. Mellone BG, Fachinetti D. Diverse mechanisms of centromere specification. Current
Biology. 2021;31: R1491–R1504. doi:10.1016/j.cub.2021.09.083

9. Kasinathan S, Henikoff S. Non-B-Form DNA Is Enriched at Centromeres. Mol Biol
Evol. 2018;35: 949–962. doi:10.1093/molbev/msy010

10. Dernburg AF, Sedat JW, Hawley RS. Direct Evidence of a Role for Heterochromatin in
Meiotic Chromosome Segregation. Cell. 1996;86: 135–146. doi:10.1016/S0092-
8674(00)80084-7

11. Dernburg AF, Broman KW, Fung JC, Marshall WF, Philips J, Agard DA, et al.
Perturbation of Nuclear Architecture by Long-Distance Chromosome Interactions. Cell.
1996;85: 745–759. doi:10.1016/S0092-8674(00)81240-4

12. Bouck DC, Bloom K. Pericentric Chromatin Is an Elastic Component of the Mitotic
Spindle. Current Biology. 2007;17: 741–748. doi:10.1016/j.cub.2007.03.033

13. Peters AH, O'Carroll D, Scherthan H, Mechtler K, Sauer S, Schofer C, et al. Loss of the
Suv39h histone methyltransferases impairs mammalian heterochromatin and genome
stability. Cell. 2001;107: 323–37.

14. Cosby RL, Chang N-C, Feschotte C. Host–transposon interactions: conflict,
cooperation, and cooption. Genes Dev. 2019;33: 1098–1116.
doi:10.1101/gad.327312.119

15. Cacchione S, Cenci G, Raffa GD. Silence at the End: How Drosophila Regulates
Expression and Transposition of Telomeric Retroelements. Journal of Molecular
Biology. 2020;432: 4305–4321. doi:10.1016/j.jmb.2020.06.004

16. Nelson JO, Slicko A, Yamashita YM. The retrotransposon R2 maintains Drosophila
ribosomal DNA repeats. Proceedings of the National Academy of Sciences. 2023;120:
e2221613120. doi:10.1073/pnas.2221613120

17. Chang C-H, Chavan A, Palladino J, Wei X, Martins NMC, Santinello B, et al. Islands of
retroelements are major components of Drosophila centromeres. Becker PB, editor.
PLoS Biol. 2019;17: e3000241. doi:10.1371/journal.pbio.3000241

18. Nowacki M, Higgins BP, Maquilan GM, Swart EC, Doak TG, Landweber LF. A
Functional Role for Transposases in a Large Eukaryotic Genome. Science. 2009;324:
935–938. doi:10.1126/science.1170023

19. Allshire RC, Madhani HD. Ten principles of heterochromatin formation and function.
Nat Rev Mol Cell Biol. 2018;19: 229–244. doi:10.1038/nrm.2017.119

20. Brändle F, Frühbauer B, Jagannathan M. Principles and functions of pericentromeric
satellite DNA clustering into chromocenters. Seminars in Cell & Developmental
Biology. 2022;128: 26–39. doi:10.1016/j.semcdb.2022.02.005

21. Seller CA, Cho C-Y, O'Farrell PH. Rapid embryonic cell cycles defer the establishment
of heterochromatin by Eggless/SetDB1 in Drosophila. Genes Dev. 2019;33: 403–417.
doi:10.1101/gad.321646.118

22. Yuan K, O'Farrell PH. TALE-light imaging reveals maternally guided, H3K9me2/3-
independent emergence of functional heterochromatin in Drosophila embryos. Genes
Dev. 2016;30: 579–593. doi:10.1101/gad.272237.115

23. Tang R, Zhou M, Chen Y, Jiang Z, Fan X, Zhang J, et al. H3K14ac facilitates the
reinstallation of constitutive heterochromatin in Drosophila early embryos by engaging
Eggless/SetDB1. Proceedings of the National Academy of Sciences. 2024;121:
e2321859121. doi:10.1073/pnas.2321859121

24. Atinbayeva N, Valent I, Zenk F, Loeser E, Rauer M, Herur S, et al. Inheritance of H3K9
methylation regulates genome architecture in Drosophila early embryos. The EMBO
Journal. 2024;43: 2685–2714. doi:10.1038/s44318-024-00127-z

25. Montavon T, Shukeir N, Erikson G, Engist B, Onishi-Seebacher M, Ryan D, et al.
Complete loss of H3K9 methylation dissolves mouse heterochromatin organization.
Nat Commun. 2021;12: 4359. doi:10.1038/s41467-021-24532-8

26. Jagannathan M, Cummings R, Yamashita YM. A conserved function for
pericentromeric satellite DNA. VijayRaghavan K, editor. eLife. 2018;7: e34122.
doi:10.7554/eLife.34122

27. Jagannathan M, Cummings R, Yamashita YM. The modular mechanism of
chromocenter formation in Drosophila. VijayRaghavan K, Gerbi SA, editors. eLife.
2019;8: e43938. doi:10.7554/eLife.43938

28. Alekseyenko AA, Gorchakov AA, Zee BM, Fuchs SM, Kharchenko PV, Kuroda MI.
Heterochromatin-associated interactions of *Drosophila* HP1a with dADD1, HIPP1, and
repetitive RNAs. Genes Dev. 2014;28: 1445–1460. doi:10.1101/gad.241950.114

29. Kochanova NY, Schauer T, Mathias GP, Lukacs A, Schmidt A, Flatley A, et al. A multi-
layered structure of the interphase chromocenter revealed by proximity-based
biotinylation. Nucleic Acids Research. 2020;48: 4161–4178. doi:10.1093/nar/gkaa145

30. Swenson JM, Colmenares SU, Strom AR, Costes SV, Karpen GH. The composition
and organization of Drosophila heterochromatin are heterogeneous and dynamic.
Shilatifard A, editor. eLife. 2016;5: e16096. doi:10.7554/eLife.16096

31. Thomae AW, Schade GOM, Padeken J, Borath M, Vetter I, Kremmer E, et al. A Pair of
Centromeric Proteins Mediates Reproductive Isolation in Drosophila Species.
Developmental Cell. 2013;27: 412–424. doi:10.1016/j.devcel.2013.10.001

32. Lukacs A, Thomae AW, Krueger P, Schauer T, Venkatasubramani AV, Kochanova NY,
et al. The Integrity of the Speciation Core Complex is necessary for centromeric
binding and reproductive isolation in Drosophila. bioRxiv. 2021; 2021.02.05.429932.
doi:10.1101/2021.02.05.429932

33. Torok T, Harvie PD, Buratovich M, Bryant PJ. The product of proliferation disrupter is
concentrated at centromeres and required for mitotic chromosome condensation and
cell proliferation in Drosophila. Genes & Development. 1997;11: 213–225.
doi:10.1101/gad.11.2.213

34. Schotta G, Ebert A, Krauss V, Fischer A, Hoffmann J, Rea S, et al. Central role of
Drosophila SU(VAR)3-9 in histone H3-K9 methylation and heterochromatic gene
silencing. EMBO J. 2002;21: 1121–1131. doi:10.1093/emboj/21.5.1121

35. Kerrebrock AW, Moore DP, Wu JS, Orr-Weaver TL. Mei-S332, a drosophila protein
required for sister-chromatid cohesion, can localize to meiotic centromere regions.
Cell. 1995;83: 247–256. doi:10.1016/0092-8674(95)90166-3

36. Satyaki PR, Cuykendall TN, Wei KH, Brideau NJ, Kwak H, Aruna S, et al. The Hmr and
Lhr hybrid incompatibility genes suppress a broad range of heterochromatic repeats.
PLoS Genet. 2014;10: e1004240. doi:10.1371/journal.pgen.1004240

37. Blum JA, Bonaccorsi S, Marzullo M, Palumbo V, Yamashita YM, Barbash DA, et al.
The Hybrid Incompatibility Genes Lhr and Hmr Are Required for Sister Chromatid
Detachment During Anaphase but Not for Centromere Function. Genetics. 2017;207:
1457–1472. doi:10.1534/genetics.117.300390

38. Ainsztein AM, Kandels-Lewis SE, Mackay AM, Earnshaw WC. INCENP Centromere
and Spindle Targeting: Identification of Essential Conserved Motifs and Involvement of
Heterochromatin Protein HP1. Journal of Cell Biology. 1998;143: 1763–1774.
doi:10.1083/jcb.143.7.1763

39. Reyes-Carmona S, Valadéz-Graham V, Aguilar-Fuentes J, Zurita M, León-Del-Río A.
Trafficking and chromatin dynamics of holocarboxylase synthetase during
development of Drosophila melanogaster. Molecular Genetics and Metabolism.
2011;103: 240–248. doi:10.1016/j.ymgme.2011.03.005

40.   Huo X, Ji L, Zhang Y, Lv P, Cao X, Wang Q, et al. The Nuclear Matrix Protein SAFB
Cooperates with Major Satellite RNAs to Stabilize Heterochromatin Architecture
Partially through Phase Separation. Molecular Cell. 2020;77: 368-383.e7.
doi:10.1016/j.molcel.2019.10.001

41.   Guruharsha KG, Rual JF, Zhai B, Mintseris J, Vaidya P, Vaidya N, et al. A protein
complex network of Drosophila melanogaster. Cell. 2011;147: 690–703.
doi:10.1016/j.cell.2011.08.047

42.   Tang H-W, Spirohn K, Hu Y, Hao T, Kovács IA, Gao Y, et al. Next-generation large-
scale binary protein interaction network for Drosophila melanogaster. Nat Commun.
2023;14: 2162. doi:10.1038/s41467-023-37876-0

43.   Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The
STRING database in 2023: protein-protein association networks and functional
enrichment analyses for any sequenced genome of interest. Nucleic Acids Res.
2023;51: D638–D646. doi:10.1093/nar/gkac1000

44.   Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, et al. Protein complex
prediction with AlphaFold-Multimer. bioRxiv; 2022. p. 2021.10.04.463034.
doi:10.1101/2021.10.04.463034

45.   Durham J, Zhang J, Humphreys IR, Pei J, Cong Q. Recent advances in predicting and
modeling protein–protein interactions. Trends in Biochemical Sciences. 2023;48: 527–
538. doi:10.1016/j.tibs.2023.03.003

46.   Burke DF, Bryant P, Barrio-Hernandez I, Memon D, Pozzati G, Shenoy A, et al.
Towards a structurally resolved human protein interaction network. Nat Struct Mol
Biol. 2023;30: 216–225. doi:10.1038/s41594-022-00910-8

47.   Kim A-R, Hu Y, Comjean A, Rodiger J, Mohr SE, Perrimon N. Enhanced Protein-
Protein Interaction Discovery via AlphaFold-Multimer. bioRxiv; 2024. p.
2024.02.19.580970. doi:10.1101/2024.02.19.580970

48.   Charlesworth B, Sniegowski P, Stephan W. The evolutionary dynamics of repetitive
DNA in eukaryotes. Nature. 1994;371: 215–220. doi:10.1038/371215a0

49.   Chiolo I, Minoda A, Colmenares SU, Polyzos A, Costes SV, Karpen GH. Double-strand
breaks in heterochromatin move outside of a dynamic HP1a domain to complete
recombinational repair. Cell. 2011;144: 732–744. doi:10.1016/j.cell.2011.02.012

50.   Ryu T, Spatola B, Delabaere L, Bowlin K, Hopp H, Kunitake R, et al. Heterochromatic
breaks move to the nuclear periphery to continue recombinational repair. Nat Cell Biol.
2015;17: 1401–1411. doi:10.1038/ncb3258

51.   Mitrentsi I, Lou J, Kerjouan A, Verigos J, Reina-San-Martin B, Hinde E, et al.
Heterochromatic repeat clustering imposes a physical barrier on homologous
recombination to prevent chromosomal translocations. Molecular Cell. 2022;82: 2132-
2147.e6. doi:10.1016/j.molcel.2022.03.033

52.   Hughes SE, Hawley RS. Topoisomerase II Is Required for the Proper Separation of
Heterochromatic Regions during Drosophila melanogaster Female Meiosis. PLOS
Genetics. 2014;10: e1004650. doi:10.1371/journal.pgen.1004650

53.   Caridi CP, D'Agostino C, Ryu T, Zapotoczny G, Delabaere L, Li X, et al. Nuclear F-
actin and myosins drive relocalization of heterochromatic breaks. Nature. 2018;559:
54–60. doi:10.1038/s41586-018-0242-8

54.   Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, et al. Ten
things you should know about transposable elements. Genome Biology. 2018;19: 199.
doi:10.1186/s13059-018-1577-z

55.   Ozata DM, Gainetdinov I, Zoch A, O'Carroll D, Zamore PD. PIWI-interacting RNAs:
small RNAs with big functions. Nat Rev Genet. 2019;20: 89–108. doi:10.1038/s41576-
018-0073-3

56.   Czech B, Munafò M, Ciabrelli F, Eastwood EL, Fabry MH, Kneuss E, et al. piRNA-
Guided Genome Defense: From Biogenesis to Silencing. Annual Review of Genetics.
2018;52: 131–157. doi:10.1146/annurev-genet-120417-031441

57.   Klattenhoff C, Xi H, Li C, Lee S, Xu J, Khurana JS, et al. The Drosophila HP1 Homolog
Rhino Is Required for Transposon Silencing and piRNA Production by Dual-Strand
Clusters. Cell. 2009;138: 1137–1149. doi:10.1016/j.cell.2009.07.014

58.   Zhang Z, Wang J, Schultz N, Zhang F, Parhad SS, Tu S, et al. The HP1 Homolog
Rhino Anchors a Nuclear Complex that Suppresses piRNA Precursor Splicing. Cell.
2014;157: 1353–1363. doi:10.1016/j.cell.2014.04.030

59.   Mohn F, Sienski G, Handler D, Brennecke J. The Rhino-Deadlock-Cutoff Complex
Licenses Noncanonical Transcription of Dual-Strand piRNA Clusters in Drosophila.
Cell. 2014;157: 1364–1379. doi:10.1016/j.cell.2014.04.031

60.   Andreev VI, Yu C, Wang J, Schnabl J, Tirian L, Gehre M, et al. Panoramix
SUMOylation on chromatin connects the piRNA pathway to the cellular
heterochromatin machinery. Nat Struct Mol Biol. 2022;29: 130–142.
doi:10.1038/s41594-022-00721-x

61.   Jankovics F, Bence M, Sinka R, Faragó A, Bodai L, Pettkó-Szandtner A, et al.
Drosophila small ovary gene is required for transposon silencing and heterochromatin
organization, and ensures germline stem cell maintenance and differentiation.
Development. 2018;145: dev170639. doi:10.1242/dev.170639

62.   Benner L, Castro EA, Whitworth C, Venken KJT, Yang H, Fang J, et al. Drosophila
Heterochromatin Stabilization Requires the Zinc-Finger Protein Small Ovary. Genetics.
2019;213: 877–895. doi:10.1534/genetics.119.302590

63.   Onishi R, Sato K, Murano K, Negishi L, Siomi H, Siomi MC. Piwi suppresses
transcription of Brahma-dependent transposons via Maelstrom in ovarian somatic
cells. Science Advances. 2020;6: eaaz7420. doi:10.1126/sciadv.aaz7420

64.   Sun J, Liu C, Bai X, Li X, Li J, Zhang Z, et al. Drosophila FIT is a protein-specific satiety
hormone essential for feeding control. Nat Commun. 2017;8: 14161.
doi:10.1038/ncomms14161

65.   Wei X, Eickbush DG, Speece I, Larracuente AM. Heterochromatin-dependent
transcription of satellite DNAs in the Drosophila melanogaster female germline. Weigel
D, editor. eLife. 2021;10: e62375. doi:10.7554/eLife.62375

66.   Giauque CC, Bickel SE. Heterochromatin-Associated Proteins HP1a and Piwi
930   Collaborate to Maintain the Association of Achiasmate Homologs in Drosophila
931   Oocytes. Genetics. 2016;203: 173–189. doi:10.1534/genetics.115.186460

932   67.   Brower-Toland B, Findley SD, Jiang L, Liu L, Yin H, Dus M, et al. Drosophila PIWI
933   associates with chromatin and interacts directly with HP1a. Genes Dev. 2007;21:
934   2300–2311. doi:10.1101/gad.1564307

935   68.   Gu T, Elgin SCR. Maternal Depletion of Piwi, a Component of the RNAi System,
936   Impacts Heterochromatin Formation in Drosophila. PLOS Genetics. 2013;9: e1003780.
937   doi:10.1371/journal.pgen.1003780

938   69.   Ma X, Wang S, Do T, Song X, Inaba M, Nishimoto Y, et al. Piwi Is Required in Multiple
939   Cell Types to Control Germline Stem Cell Lineage Development in the Drosophila
940   Ovary. PLOS ONE. 2014;9: e90267. doi:10.1371/journal.pone.0090267

941   70.   Handler D, Meixner K, Pizka M, Lauss K, Schmied C, Gruber FS, et al. The Genetic
942   Makeup of the Drosophila piRNA Pathway. Molecular Cell. 2013;50: 762–777.
943   doi:10.1016/j.molcel.2013.04.031

944   71.   Czech B, Preall JB, McGinn J, Hannon GJ. A Transcriptome-wide RNAi Screen in the
945   Drosophila Ovary Reveals Factors of the Germline piRNA Pathway. Molecular Cell.
946   2013;50: 749–761. doi:10.1016/j.molcel.2013.04.007

947   72.   Ghabrial A, Schüpbach T. Activation of a meiotic checkpoint regulates translation of
948   Gurken during Drosophila oogenesis. Nat Cell Biol. 1999;1: 354–357.
949   doi:10.1038/14046

950   73.   Moon S, Cassani M, Lin YA, Wang L, Dou K, Zhang ZZ. A Robust Transposon-
951   Endogenizing Response from Germline Stem Cells. Dev Cell. 2018;47: 660-671 e3.
952   doi:10.1016/j.devcel.2018.10.011

953   74.   Sienski G, Dönertas D, Brennecke J. Transcriptional Silencing of Transposons by Piwi
954   and Maelstrom and Its Impact on Chromatin State and Gene Expression. Cell.
955   2012;151: 964–980. doi:10.1016/j.cell.2012.10.040

956   75.   Baumgartner L, Handler D, Platzer SW, Yu C, Duchek P, Brennecke J. The Drosophila
957   ZAD zinc finger protein Kipferl guides Rhino to piRNA clusters. Bourc'his D, Struhl K,
958   Zhang Z, editors. eLife. 2022;11: e80067. doi:10.7554/eLife.80067

959   76.   Kneuss E, Munafò M, Eastwood EL, Deumer U-S, Preall JB, Hannon GJ, et al.
960   Specialization of the *Drosophila* nuclear export family protein Nxf3 for piRNA precursor
961   export. Genes Dev. 2019;33: 1208–1220. doi:10.1101/gad.328690.119

962   77.   ElMaghraby MF, Andersen PR, Pühringer F, Hohmann U, Meixner K, Lendl T, et al. A
963   Heterochromatin-Specific RNA Export Pathway Facilitates piRNA Production. Cell.
964   2019;178: 964-979.e20. doi:10.1016/j.cell.2019.07.007

965   78.   Staller MV, Yan D, Randklev S, Bragdon MD, Wunderlich ZB, Tao R, et al. Depleting
966   Gene Activities in Early Drosophila Embryos with the "Maternal-Gal4–shRNA" System.
967   Genetics. 2013;193: 51–61. doi:10.1534/genetics.112.144915

79. Lin H, Spradling AC. A novel group of pumilio mutations affects the asymmetric
division of germline stem cells in the Drosophila ovary. Development. 1997;124: 2463–
2476. doi:10.1242/dev.124.12.2463

80. Sun X, Wahlstrom J, Karpen G. Molecular structure of a functional Drosophila
centromere. Cell. 1997;91: 1007–19.

81. Fabry MH, Falconio FA, Joud F, Lythgoe EK, Czech B, Hannon GJ. Maternally
inherited piRNAs direct transient heterochromatin formation at active transposons
during early Drosophila embryogenesis. Struhl K, editor. eLife. 2021;10: e68573.
doi:10.7554/eLife.68573

82. Gonzalez LE, Tang X, Lin H. Maternal Piwi regulates primordial germ cell development
to ensure the fertility of female progeny in Drosophila. Genetics. 2021;219: iyab091.
doi:10.1093/genetics/iyab091

83. Erdel F. Phase transitions in heterochromatin organization. Current Opinion in
Structural Biology. 2023;80: 102597. doi:10.1016/j.sbi.2023.102597

84. Van Doren M, Williamson AL, Lehmann R. Regulation of zygotic gene expression in
Drosophilaprimordialgerm cells. Current Biology. 1998;8: 243–246.
doi:10.1016/S0960-9822(98)70091-0

85. Zeidler MP, Perrimon N, Strutt DI. Polarity determination in the Drosophila eye: a novel
role for Unpaired and JAK/STAT signaling. Genes Dev. 1999;13: 1342–1353.

86. Lu KL, Nelson JO, Watase GJ, Warsinger-Pepe N, Yamashita YM. Transgenerational
dynamics of rDNA copy number in Drosophila male germline stem cells. Spradling AC,
editor. eLife. 2018;7: e32421. doi:10.7554/eLife.32421

87. Wiśniewski JR, Zougman A, Nagaraj N, Mann M. Universal sample preparation method
for proteome analysis. Nat Methods. 2009;6: 359–362. doi:10.1038/nmeth.1322

88. Bryant P, Pozzati G, Elofsson A. Improved prediction of protein-protein interactions
using AlphaFold2. Nat Commun. 2022;13: 1265. doi:10.1038/s41467-022-28865-w

89. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013;29: 15–21.
doi:10.1093/bioinformatics/bts635

90. Jin Y, Tam OH, Paniagua E, Hammell M. TEtranscripts: a package for including
transposable elements in differential expression analysis of RNA-seq datasets.
Bioinformatics. 2015;31: 3593–3599. doi:10.1093/bioinformatics/btv422

91. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology. 2014;15: 550. doi:10.1186/s13059-014-
0550-8

1003

1004

Acknowledgements

We thank members of the Jagannathan lab, Ulrike Kutay, Ruth Kroschewski and Hugo Stocker for discussion and comments on the manuscript. We thank Benjamin Frühbauer for help with representing the protein models. We thank Daniel Barbash, Ferenc Jankovics, Yukiko Yamashita, the Bloomington Drosophila Stock Center, the Vienna Drosophila Resource Center, the Kyoto Drosophila Stock Center and the Developmental Studies Hybridoma Bank for reagents and resources. We thank the Scientific Center for optical and Electron Microscopy (ScopeM) and the Mass Spectrometry facility at the Institute of Biochemistry at ETH Zurich for technical support. We thank Shinichi Sunagawa for providing bioinformatics resources. M.J is supported by a project grant (310030_189131) from the Swiss National Science Foundation. A.C acknowledges support from Genetics Society of America in the form of a DeLill Nasser Award for Professional Development in Genetics.

1019 **Figures and Legends**

1020

Figure 1. Experimental setup for the characterization of the chromocenter-associated

proteome. (A) Experimental scheme for the identification of D1- and Prod-associated

proteins across multiple *Drosophila* tissues using quantitative mass spectrometry. (B) D1

and Prod are in proximity within the nucleus in multiple cell types including gastrulating

embryos, female germ cells and male germ cells. (C) Bar plot of the total number of D1 foci

present in female GSCs and male GSCs in the wild type (*yw*), *NLS-GFP*, *GFP-D1* and *GFP-*

Prod strains. n indicates the number of cells analyzed. (D) Bar plot of the total number of

Prod foci present in female GSCs and male GSCs in the wild type (*yw*), *NLS-GFP*, *GFP-D1*

and *GFP-Prod* strains. n indicates the number of cells analyzed. (E, F) Bar plot of the

number of progeny sired by a single tester male (E) or female (F) when mated with two wild

type flies of the opposite sex over the course of one week. **** represents $p < 0.0001$ from an

ordinary one-way ANOVA. All error bars are SD and all scale bars are 5µm.

1033

Figure 2. The chromocenter-associated proteome across *Drosophila* tissues. (A-B)

Volcano plots of the D1-associated proteome in embryos (A) and ovaries (B). The dashed

lines mark $\log_2$FC>1 and p<0.05; blue points indicate proteins involved in transposon

repression, red points indicate proteins involved in DNA repair and yellow points indicate

proteins pulled down by D1 and Prod. Number of biological replicates are indicated in the

plot. (C) Intersection plot representing number of overlapping proteins in the D1 and Prod

pulldowns from different tissues. Blue arrowheads indicate number of proteins common to

D1 and Prod pulldown in the same tissue while magenta arrowheads indicate number of

common proteins pulled down by D1 or Prod across different tissues (D) Enrichment of the

selected proteins pulled down by D1 and Prod from more than one tissue. The size and

colour of the circles are proportional to the $\log_2$FC while the black outline indicates $p < 0.05$.

(E) Hypergeometric test of DNA repair or transposon repression pathway proteins from the

multi-tissue D1 and Prod interactome. Dashed line indicates $p = 0.05$.

1047

Figure 3. Molecular relationships between proteins involved in transposon repression

and chromocenter formation (A) GFP-Piwi and Sov-GFP embryos were stained with

D1/Prod (magenta) and DAPI (blue). Arrowheads indicate Sov-GFP foci in proximity to

chromocenter boundaries. (B) Volcano plot of the Piwi-associated proteome in embryos.

The dashed lines mark $\log_2$FC>1 and p<0.05 while the red points indicate proteins identified

in pulldowns of D1 or Prod from embryos. Number of biological replicates are indicated. (C)

Enrichment of the selected proteins pulled down by D1, Prod and Piwi from embryos. The

size and colour of the circles are proportional to the $\log_2$FC while the black outline indicates

p<0.05. (D) ipTM scores of the pairwise protein structural models based on Alphafold

Multimer. Gray points indicate low confidence structural models, blue points indicate

medium confidence structural models (ipTM>0.5) and red points indicate high confidence

structural models (ipTM>0.8). All scale bars are 5μm.

1060

Figure 4. The effect of D1 on piRNA biogenesis and TE expression in ovaries. (A)

Brightfield images of ovaries from *D1* heterozygous and *D1* mutant females in a *chk2^6006*

background. Scale bars: 100μm. (B) Volcano plot of differentially expressed transposon

families in ovaries from *D1* heterozygous and *D1* mutant females in a *chk2^6006* background

(n=3). Statistically significant changes ($\log_2$FC >1, p_{adj} <0.05) are indicated with the black

outlines. (C, D) Size profiles of Argonaute-bound small RNAs and miRNAs isolated from *D1*

heterozygous and *D1* mutant ovaries. Reads from three replicates per genotype were

normalized per million miRNA reads. (E) Average FPKM for piRNAs mapping antisense to

transposon consensus sequences from D1 heterozygous and D1 mutant ovaries (n=3).

Black outline indicates statistically significant changes in antisense piRNAs ($\log_2$FC >1,

p_{adj}<0.05) (F) Volcano plot of differentially expressed genes in ovaries from *D1* heterozygous

and *D1* mutant females in a *chk2^6006* background (n=3). Statistically significant changes

($\log_2$FC >1, p_{adj} <0.05) are indicated with the black outlines. Blue points indicate the 347

genes associated with transposon repression (Table S6).

Figure 5. D1 blocks localization of the piRNA biogenesis machinery to chromocenters.

(A) A single nurse cell from an EGFP-Cuff ovary stained with D1 (magenta), Vasa (blue) and

DAPI (gray). The arrowhead points to the DAPI-dense chromocenter. The insets highlight

Cuff localization at the periphery of D1-containing chromocenters. (B) Localization of EGFP-

Cuff in nurse cell nuclei from control ($D1^{LL03310}$/TM6C) and D1 mutant ($D1^{LL03310}$/$Df(3R)^{Bsc666}$)

ovaries. Yellow arrowheads point to the location of the chromocenter. (C) Box-and-whisker

plot of the size of nurse cell chromocenters relative to total nuclear size in control

($D1^{LL03310}$/TM6C) and D1 mutant ($D1^{LL03310}$/$Df(3R)^{Bsc666}$) ovaries. **** represents p<0.0001 from

a student's t-test. (D) Box-and-whisker plot of EGFP-Cuff enrichment at chromocenters

relative to the rest of the nurse cell nucleus in control ($D1^{LL03310}$/TM6C) and D1 mutant

($D1^{LL03310}$/$Df(3R)^{Bsc666}$) ovaries. **** represents p<0.0001 from a student's t-test. (E)

Localization of EGFP-Cuff (green) and Nxf3 (magenta) in nurse cell nuclei from control

(*D1^{LL03310}*/TM6C) and D1 mutant (*D1^{LL03310}*/*Df(3R)^{Bsc666}*) ovaries. Yellow arrowheads point to

the location of the chromocenter. (F) RNA *in situ* hybridization against the 359bp satellite

DNA transcripts (plus strand, magenta) in control (*D1^{LL03310}*/TM6C) and D1 mutant

(*D1^{LL03310}*/*Df(3R)^{Bsc666}*) ovaries in an EGFP-Cuff (blue) background that were also stained for

DAPI (grey). (G) Box-and-whisker plot of chromocenter-localized 359bp satellite DNA

transcripts in nurse cells from control (*D1^{LL03310}*/TM6C) and D1 mutant (*D1^{LL03310}*/*Df(3R)^{Bsc666}*)

ovaries in an EGFP-Cuff background. **** represents $p < 0.0001$ from a student's t-test. All

scale bars are 5µm. (H) RNA *in situ* hybridization against the 359bp satellite DNA transcripts

(plus strand, magenta) in control (*D1^{LL03310}*/TM6C) and D1 mutant (*D1^{LL03310}*/*Df(3R)^{Bsc666}*)

ovaries in an EGFP-Cuff (green) background that were also stained for DAPI (blue) following

RNase A treatment post fixation.

Figure 6. Loss of D1 during embryogenesis triggers transposon expression in adult ovaries. (A) Brightfield images of ovaries from the indicated genotypes. Scale bars: 100µm. (B) Ovaries from the indicated genotypes were stained with Vasa (green) and DAPI (magenta). Scale bars: 5µm. (C) Percentage of germaria containing a normal population of germ cells, only germline stem cells/cystoblasts (GSC/CB only) or completely lacking germ cells in *yw* (n=48), *aubergine* mutant (n=66), F_1 *D1* heterozygous (n=51), F_1 *D1* mutant (n=65), F_2 *D1* heterozygous (n=81) and F_2 *D1* mutant (n=109). (D) Bar plot of the number of progeny sired by a single female of the indicated genotypes when mated with two wild type males over the course of one week. CV represents coefficient of variation. **** represents $p < 0.0001$ from an ordinary one-way ANOVA. (E) RNA *in situ* hybridization against Copia (green) and total polyadenylated mRNA (blue) was performed in ovaries of the indicated genotypes that were also stained for DAPI (magenta). Yellow and black arrowheads indicate undifferentiated germ cells in the *aubergine* mutant germarium. Scale bars: 5µm. (F) Percentage of germaria with germ cells expressing Copia transcripts in *yw* (n=62), *aubergine* mutant (n=67), F_1 *D1* mutant (n=71) and F_2 *D1* mutant (n=50). (G) Box-and-whisker plot of copia expression normalized to total polyA mRNA in undifferentiated germ cells (GSCs/CBs) from *yw* (n=16), *aubergine* mutant (n=22), F_1 *D1* mutant (n=18) and F_2 *D1* mutant (n=29). **** represents $p < 0.0001$ from one-way ANOVA. (H) Brightfield images of ovaries from the indicated genotypes. Scale bars: 100µm. (I) Percentage of germaria containing a normal population of germ cells, only germline stem cells/cystoblasts (GSC/CB only) or completely lacking germ cells in F_2 *D1* heterozygous (n=62) and F_2 *D1* mutant (n=38) ovaries in a *Chk2* mutant background.

1122

1123 **Figure 7. The role of D1 in heritable TE repression.** An embryonic pool of D1 that

1124 originates from either maternal deposition (*Mat +*) and/or zygotic expression (*Zyg +*) is

1125 important for TE repression in adult gonads. In the absence of embryonic D1 (F_2 *D1* mt, *Mat*

1126 *– Zyg –*), TE expression contributes to Chk2-mediated gonadal atrophy.

1127

Supplementary Figure 1. Schematic of the GFP-Prod allele and optimization of sample

lysis and immunoprecipitation. (A) A 400bp promoter sequence upstream of the Prod

translation start site was placed upstream of a GFP-tagged and codon-optimized Prod

sequence, inserted into the attP40 locus and recombined with the *prod*[k08810] loss-of-

function allele. (B) Rescue of prod mutant (*prod*[k08810/U]) lethality was assessed using p400-

GFP-Prod, *prod*[k08810] (p400-GFP-Prod, 447 flies were counted from 5 crosses) and *prod*[k08810]

(Ctrl, 314 flies were counted from 5 crosses). Red line indicates median while dashed line

indicates percent progeny for complete rescue. (C) Brightfield images of ovaries from the

indicated genotypes. Scale bars: 100µm. (D) Western blot of equivalent fractions of the

soluble supernatant and insoluble pellet from GFP-Piwi embryos using mouse anti-GFP,

guinea pig anti-D1 and rabbit anti-H3 (loading control). (E) Immunoprecipitation of GFP from

embryos expressing GFP-Prod or GFP-D1. Immunoprecipitated proteins were blotted using

mouse anti-GFP and are indicated by arrowheads. Expected sizes: GFP-Prod: 75kDa, GFP-

D1: 80kDa.

1142

1143

Supplementary Figure 2. Characterization of D1- and Prod-associated proteins in

GSC-enriched testes and embryos. (A) Volcano plot of the D1-associated proteome in

GSC-enriched testes. The dashed lines mark $\log_2 FC>1$ and $p<0.05$; blue points indicate

proteins involved in transposon repression and red points indicate proteins involved in DNA

repair. Number of biological replicates are indicated in the plot. (B-D) k-means clustering of

embryo hits using STRING-DB show relationships between Prod-associated (B) and D1-

associated (C, D) proteins.

1151

Supplementary Figure 3. Localization of D1 and Prod with respect to Piwi, Sov and Hmr in nurse cells and spermatogonia. GFP-Piwi (A, B), sov-GFP (C, D) and Hmr-HA (E, F) ovaries and testes were stained with D1/Prod (magenta), Vasa/Lamin (blue) and DAPI (gray). Arrowheads indicate sov or Hmr foci in proximity to chromocenter boundaries in individual nurse cells or spermatogonia. Scale bars: 5µm.

1157

Supplementary Figure 4. Assessing the interactions between Piwi and satellite DNA-

binding proteins by AFM structural modelling and co-immunoprecipitation. (A)

Representation of the D1-Piwi and D1-Prod interaction interfaces predicted by Alphafold

Multimer structural modelling. The following regions on D1 (120-143, 157-175 and 322-355)

and on Prod (120-181 and 320-343) are depicted in magenta. (B) Immunoprecipitation of

GFP and Piwi using GFP-trap beads from NLS-GFP (control) and GFP-Piwi embryos.

Immunoprecipitated proteins were blotted using mouse anti-GFP and guinea pig anti-D1.

1165

Supplementary Figure 5. Loss of *aubergine* has a mild effect on Prod foci in female

GSCs. (A) Female GSCs expressing *mCherry^RNAi* (*mCherry^GLKD*) or *aub^RNAi* (*aub^GLKD*) under the

control of *nos-gal4* stained for Prod (magenta), Lamin (green) and Vasa (blue). Scale bars:

5µm. (B, C) Box-and-whisker plot of the total number of Prod foci (B) or D1 foci (C) present

per female GSC in *mCherry^GLKD* (n=58 for Prod, n=60 for D1), *aub^GLKD* (n=42 for Prod, n=36

for D1), *aubergine* heterozygous control (Ctrl., n=38 for Prod, n=59 for D1) and *aubergine*

mutant (*aub^QC42/HN2*, n=52 for Prod, n=56 for D1). n indicates the number of cells analyzed. **

indicates p<0.01 based on a one-way ANOVA.

DAPI Prod

Supplementary Figure 6. Validation of D1 and Prod knockdowns in gonadal somatic

cells and germline cells. (A, B) Individual egg chambers stained for DAPI (magenta) and

Prod/D1 following *tj-gal4* mediated expression of *D1* ^RNAi^ (KK110307) or *Prod* ^RNAi^

(KK104361). *mCherry* ^RNAi^ was used as a control. (C, D) Anterior region of individual

ovarioles were stained for DAPI (blue) and Prod (magenta) following *nos-gal4* mediated

expression of *mCherry* ^RNAi^ (C) or *Prod* ^RNAi^ (D). Scale bars: 5μm.

1181

Supplementary Figure 7. Loss of D1 or Prod does not trigger expression of transposon reporters in gonadal somatic cells or germ cells. (A) Individual egg chambers stained for beta-galactosidase (indicating expression of *gypsy-lacZ*) following *tj-gal4* mediated expression of *mCherry* ^{RNAi}, *sov* ^{RNAi} (HMC04875), *D1* ^{RNAi} (KK110307) or *Prod* ^{RNAi} (KK104361). Arrowhead indicates follicle cells expressing *gypsy-lacZ*. (B) Anterior region of individual ovarioles were stained for DAPI in the following strains carrying GFP-Burdock-lacZ: *D1*^{LL03310}/*TM6B* (Ctrl.) and *D1*^{LL03310}/*Df(3R)*^{Bsc666} (*D1* mutant), *nos > mCherry* ^{RNAi} (*mCherry* ^{GLKD}), *nos > Prod* ^{RNAi} (*Prod* ^{GLKD}), *nos > aub*^{RNAi} (HMS01945, *aub* ^{GLKD}). Arrowheads indicate germ cells expressing GFP-Burdock. Scale bars: 5µm.

1191

1192 **Supplementary Figure 8. Localization of EGFP-Cuff relative to chromocenters.** (A, B)

1193 EGFP-Cuff ovaries and testes were stained for D1/Prod (magenta), Vasa/Lamin (blue) and

1194 DAPI (gray). Individual nurse cells or spermatogonia are depicted. Scale bars: 5µm.

1195

1196 **Supplementary Figure 9. Putative expression levels of maternal and zygotic D1**

1197 **through development.** (A, B) Crossing schemes used to generate F$_1$ *D1* mutants and F$_2$ *D1*

1198 mutants. (C) Theoretical depiction of maternal deposition and zygotic expression of D1

1199 across embryonic, larval and adult stages in the indicated genotypes.

Supplementary Tables

Supplementary Table 1. List of proteins detected by LC-MS/MS in embryonic lysates from GFP-D1, GFP-Prod and NLS-GFP strains. Fold change of average MS1 intensity and p-value for the GFP-D1 and GFP-Prod strains relative to the NLS-GFP control strain are shown for detected proteins. Proteins pulled down by GFP-D1 and GFP-Prod ($log_2FC>1$, $p<0.05$) are highlighted in yellow.

Supplementary Table 2. List of proteins detected by LC-MS/MS in ovary lysates from GFP-D1, GFP-Prod and NLS-GFP strains. Fold change of average MS1 intensity and p-value for the GFP-D1 and GFP-Prod strains relative to the NLS-GFP control strain are shown for detected proteins. Proteins pulled down by GFP-D1 and GFP-Prod ($log_2FC>1$, $p<0.05$) are highlighted in yellow.

Supplementary Table 3. List of proteins detected by LC-MS/MS in GSC-enriched testes lysates from GFP-D1 and NLS-GFP strains. Fold change of average MS1 intensity and p-value for the GFP-D1 strain relative to the NLS-GFP control strain are shown for detected proteins.

Supplementary Table 4. ipTM scores of D1 and Prod interactors based on AFM structural models.

Supplementary Table 5. List of proteins associated with DNA repair in *Drosophila*.

Supplementary Table 6. List of proteins associated with transposon repression in *Drosophila*.

1226

1227 **Supplementary Table 7. List of proteins detected by LC-MS/MS in embryonic lysates**

1228 **from GFP-Piwi and NLS-GFP strains.** Fold change of average MS1 intensity and p-value

1229 for the GFP-Piwi strain relative to the NLS-GFP control strain is shown for detected

1230 proteins.

1231

1232 **Supplementary Table 8. ipTM scores of protein pairs based on AFM structural models.**

1233

1234 **Supplementary Table 9. Differentially expressed genes and TE families between *D1***

1235 **heterozygous and *D1* mutant ovaries in a *chk2^{6006}* background.** Identified genes were

1236 cross-referenced against genes implicated in TE silencing from supplementary table 6.