The contribution of gene flow, selection, and genetic drift to five thousand years of human allele frequency change

Alexis Simon[1,2] and Graham Coop[1,2]

[1]Center for Population Biology, University of California, Davis, CA 95616
[2]Department of Evolution and Ecology, University of California, Davis, CA 95616

Abstract

Genomic time-series from experimental evolution studies and ancient DNA datasets offer us a chance to more directly observe the interplay of various evolutionary forces. Here we show how the genome-wide variance in allele frequency change between two time points can be decomposed into the contributions of gene flow, genetic drift, and linked selection. In closed populations, the contribution of linked selection is identifiable because it creates covariances between time intervals, and genetic drift does not. However, repeated gene flow between populations can also produce directionality in allele frequency change, creating covariances. We show how to accurately separate the fraction of variance in allele frequency change due to admixture and linked selection in a population receiving gene flow. We use two human ancient DNA datasets, spanning around 5,000 years, as time transects to quantify the contributions to the genome-wide variance in allele frequency change. We find that a large fraction of genome-wide allele frequency change is due to gene flow. In both cases, after correcting for known major gene flow events in those populations, we do not observe a signal of genome-wide linked selection. Thus despite the known role of selection in shaping long-term polymorphism levels, and an increasing number of examples of strong selection on single loci and polygenic scores from ancient DNA, it appears to be gene flow and drift, and not selection, that are the main determinants of recent genome-wide allele frequency change. Our approach should be applicable to the growing number of contemporary and ancient temporal population genomics datasets.

1 Introduction

One fundamental goal of evolutionary biology is to quantify the contribution of basic evolutionary forces to allele frequency change and how this differs across species, types of traits, and loci. For example, there is a long-standing debate about the contribution to genome-wide allele frequency change of genetic drift versus the direct or indirect effects of selection, where the indirect effects includes "linked selection", the impact of correlated selection at nearby selected sites (Barton, 2000; Charlesworth et al., 1993; Coop, 2016; Kaplan et al., 1989; Maynard-Smith & Haigh, 1974; Sella et al., 2009). However, these questions are difficult to answer with a single snapshot of contemporary genomes as it

requires teasing apart multiple interacting processes (gene flow, demography, hard or soft sweeps, background selection, and selective interference, etc).

Genomic time-series, from museum collections and ancient DNA, offer a potent reservoir of temporal genetic data to track the changes in allele frequencies, identify selected loci, and understand the impact of other evolutionary forces (e.g. He et al., 2023; Le et al., 2022; Mathieson & Terhorst, 2022). Ancient human DNA has already revolutionized our view of human history, revealing that large scale population movement and gene flow is pervasive with complex patterns of allele frequency change driven by population turnover.

Unlike genetic drift, allele frequency change due to either selection or gene flow is expected to be sustained and directional. Methods to look at selection at single loci and on polygenic scores in human ancient DNA now often account for the confounding effects of gene flow. These approaches have revealed persuasive signals of selection (Field et al., 2016; Irving-Pease et al., 2022; Ju & Mathieson, 2021; Mathieson et al., 2015; Mathieson & Terhorst, 2022; Souilmi et al., 2022; Wilde et al., 2014). However, these methods only capture outlier loci, and so cannot give us a full picture of how gene flow, selection, and genetic drift have driven genome-wide change. Linked selection is thought to have a critical role in shaping patterns of genetic diversity and divergence in humans on long time scales, with an autosomal reduction in diversity of upward of 20 % from background selection alone (McVicker et al., 2009; Murphy et al., 2021). Some authors have also argued for a pervasive role of selective sweeps in shaping genome-wide patterns of diversity (Enard et al., 2014; Schrider & Kern, 2017). Thus, it is an open question how much of allele frequency change genome-wide is driven by selection in humans.

Here, we set out to decompose the total variance in allele frequency change into the contributions of linked selection, gene flow, and drift. Unlike genetic drift, which generates uncorrelated allele frequency change between non-overlapping time intervals, ongoing selection creates covariance in allele frequency between time intervals. The use of genome-wide allele frequency change covariances in time-series data has recently been proposed to identify the proportion of genome-wide change due to selection in closed populations (Buffalo & Coop, 2019). In a panmictic population, a genome-wide positive covariance between a pair of time periods indicates the compounding of allele frequency change across generations, while negative covariance can potentially result from selection pressures in opposite directions. Many different modes of selection are expected to generate these covariance patterns (Buffalo & Coop, 2020; Santiago & Caballero, 1998). This approach has been applied to experimental evolution experiments (Brennan et al., 2022; Buffalo & Coop, 2020) and to natural populations where temporal sampling is available (in *Mimulus* and oaks; Kelly, 2022; Saleh et al., 2022). These applications, along with related methods (Bertram, 2021), have revealed that a reasonable proportion of total allele frequency change, especially in artificial selection experiments, can be potentially attributed to wide-spread selection beyond just a few outliers. However, their application when gene flow is present is complicated, as sustained gene flow across time periods can drive covariance in allele frequency change among those time periods.

Here, we develop theory and simulations to show how the effect of gene flow can be accounted for using an admixture model from a known set of sources to recover the genome-wide contribution of gene flow and linked selection. We demonstrate this approach using two European human ancient DNA time-series from the UK and the Bohemian region of Central-Europe to quantify the contributions of linked selection and gene flow to the total variance in allele frequency change between the Neolithic and a modern or recent

time point. In both cases, we find a major contribution of gene flow to allele frequency change. After correcting for known gene flow in these time transects, we do not observe any signal of genome-wide linked selection.

2 Results

2.1 Model

We consider a model of data from a population sampled at multiple discrete time points ($t \in [0, T]$) from an arbitrary geographic region. This population receives gene flow, modeled as single pulses of admixture, from other known source groups through time. We follow the population allele frequency over time, p_t, and use Δp_t, defined as $p_{t+1} - p_t$, to denote the change in allele frequency between adjacent time points. The focal population has mean ancestry fractions $\bar{\alpha}_{r,t}$ ($r \in [1, R]$) from R source populations that will change over time due to gene flow. We assume that allele frequencies in the source populations are constant and that good proxy samples for these sources are available, and discuss the implications of those assumptions later.

Given sample frequencies at some large set of SNPs, we can calculate the empirical variance-covariance matrix of allele frequency change for our time-series, $\text{Cov}(\Delta p_i, \Delta p_j)$, for all combinations of time intervals i and j, averaged over SNPs. In calculating these covariances we include adjustments for biases in the variance and covariance estimates due to shared sampling of an intermediate time point (see section 4.1 and appendix A).

Under our model, the total variance in allele frequency change between the first sampled time point (0) and any following time point (T) in the time-series can be decomposed into sums over time intervals of the contributions due to drift, selection and admixture:

$$
\begin{aligned}
\text{Var}(p_T - p_0) = &\underbrace{\sum_{i=0}^{T-1} \text{Var}(\Delta_D p_i)}_{\text{Drift}} + \underbrace{\sum_{i=0}^{T-1} \text{Var}(\Delta_S p_i)}_{\text{Selection}} + \underbrace{\sum_{i=0}^{T-1} \text{Var}(\Delta_A p_i)}_{\text{Admixture}} \\
&+ \underbrace{\sum_{i \neq j}^{T-1} \text{Cov}(\Delta_S p_i, \Delta_S p_j)}_{\text{Selection}} + \underbrace{\sum_{i \neq j}^{T-1} \text{Cov}(\Delta_A p_i, \Delta_A p_j)}_{\text{Admixture}}
\end{aligned}
\tag{1}
$$

For simplicity, here we omit an interaction between drift in one time period that admixture in later time periods subsequently erases, that adds an additional term to covariances (appendix C) that we account for.

The expected variance and covariance of allele frequency change due to admixture follow from the expected allele frequency change in ancestry proportions through time. Specifically, if in the t^{th} time interval admixture changes the ancestry proportion from the r^{th} source from $\bar{\alpha}_{r,t}$ to $\bar{\alpha}_{r,t+1}$ ($\Delta \bar{\alpha}_{r,t} = \bar{\alpha}_{r,t+1} - \bar{\alpha}_{r,t}$), then the expected change in frequency due to admixture is $\Delta_A p_t = \sum_{r=1}^{R} \Delta \bar{\alpha}_{r,t} f_r$, where f_r is the allele frequency in the source population r. Thus, the admixture contribution to covariance in allele frequency change between time periods i and j can be expressed as:

$$
\text{Cov}(\Delta_A p_i, \Delta_A p_j) = \text{Cov}\left(\sum_{r=1}^{R} \Delta \bar{\alpha}_{r,i} f_r, \sum_{r=1}^{R} \Delta \bar{\alpha}_{r,j} f_r\right)
\tag{2}
$$

As we only have sample allele frequencies from proxies of the sources of admixture, this matrix is corrected for sampling noise biases in f_r (appendix B). With this admixture covariance in hand, we can now calculate the contribution of gene flow to allele frequency change, and adjust for the contribution of admixture when looking for covariances induced by selection.

We express the estimated proportion of total variance in allele frequency change attributable to gene flow (admixture) up to time t ($A(t)$) as:

$$A(t) \equiv \frac{\sum_{i=0}^{t-1} \text{Var}(\Delta_A p_i) + \sum_{i \neq j}^{t-1} \text{Cov}(\Delta_A p_i, \Delta_A p_j)}{\text{Var}(p_t - p_0)} \tag{3}$$

where the terms in the numerator are given by eq. (2). Note that $A(t)$ might be an under-estimate as it excludes the contribution of gene flow from unmodeled sources, as well as gene flow events that leave the admixture proportions relatively unchanged.

The proportion of total variance in allele frequency change between 0 and t due to linked selection, $G(t)$, is defined as the ratio of the total covariance due to linked selection over the total variance (Buffalo & Coop, 2019). Under our model, $G(t)$ can be estimated by correcting the empirical covariance by the estimated covariance term due to admixture:

$$\begin{aligned} G(t) &\equiv \frac{\sum_{i \neq j}^{t-1} \text{Cov}(\Delta_S p_i, \Delta_S p_j)}{\text{Var}(p_t - p_0)} \\ &= \frac{\sum_{i \neq j}^{t-1} \text{Cov}(\Delta p_i, \Delta p_j) - \text{Cov}(\Delta_A p_i, \Delta_A p_j)}{\text{Var}(p_t - p_0)} \end{aligned} \tag{4}$$

We also report an estimate of G not controlling for admixture (G_{nc}). Note that G is a lower bound on the proportion as it does not account for the contribution of linked selection to the variance in allele frequency change within time periods. We attribute the residual proportion $1 - A - G$ of the total allele frequency variance to drift-like allele frequency change. This proportion of the temporal variance is consistent with genetic drift as it excludes covariances between time periods, which can not come from drift, and the contribution of known gene flow.

2.2 Simulations

To illustrate our decomposition of genome-wide allele frequency change, we simulated a simple scenario where a population receives pulses of admixture (arrows in fig. 1A and fig. S1). This repeated admixture results in positive covariances between time intervals due to the admixture-driven allele frequency change in generations 160-100 (measured before present) being in the same direction as those in generations 60-0 (fig. 1B & C). We can remove the covariance due to admixture (eq. (2)), resulting in covariances that are not significantly different from zero (fig. 1B & D).

We can calculate the total contribution of gene flow to allele frequency change in our simulated time-series (eq. (3)) and see that gene flow accounts for much of the allele frequency change. Because the repeated gene flow creates positive covariance in allele frequency change, failing to account for this gene flow generates a spurious signal of linked selection (large non-zero G_{nc}'s, dashed black line fig. 1E). However, when we account for gene flow, the signal of linked selection is almost completely removed from our neutral simulations (G, black line, fig. 1E). The remaining slightly non-zero G value in our final

Figure 1: Simulation scenario of admixture between two populations (0 and 1) under neutrality (B-E) and with selection (F). **(A)** Ancestry proportions of the focal admixed population through time in generations before present. Arrows indicate the migration pulses from Pop1 (at 150, 130, 110, 50, 30 and 10 generations before present). **(B)** Covariance between time intervals. Below diagonal values are before admixture correction, above diagonal are after admixture correction; stars indicate a 95% CI not overlapping 0. **(C)** and **(D)** pre- and post-admixture correction covariances with 95% CIs. X-axis values are slightly shifted for visualization and the bottom lines group together the points to their corresponding time. **(E)** Proportion of the total variance between the initial measured time (120) and t due to linked selection (G_{nc} and G are pre- and post-admixture correction respectively) and to gene flow (A) with 95% CIs. Points are slightly x-shifted for visualization. **(F)** Simulations for the decomposition of variance for neutral polymorphisms with selection around a gradually moving optimum starting at generation 140 BP for three independent traits. All CIs are computed using 100 replicates of the simulations.

time period (fig. 1E) results from a slight over-correction for the covariance due to the interaction of drift and gene flow (see figs. S2 and S3F).

To illustrate the effects of selection on the covariance, we extended the above admixture simulations to include a set of loci underlying traits under stabilizing selection around a moving optimum (see section 4.2). In these simulations we can see an increasing proportion of neutral allele frequency change being due to selection as covariances build up over time (black line fig. 1D, see also fig. S3). The effect of selection on the covariances in neutral allele frequencies is also well illustrated in a model without any gene flow (figs. S4 and S5).

2.3   Ancient DNA time transects

We investigated two time transects of allele frequencies in ancient humans in restricted geographical regions, the first one in the UK (Patterson et al., 2022, the England and Wales samples), and the second in the Central European region of Bohemia (Papac et al., 2021, samples from the current Czech Republic) spanning periods back to ∼5,600 years ago. Both these time-series cover major migrations of people, where an initial mixture of early-farmer-like ancestry (EEF-like) and Western hunter gatherer-like ancestry (WHG-like) is partially replaced by Steppe pastoralist-like ancestry (Steppe-like). This large turnover due to Steppe-like migration into Central and Western Europe was followed by a rebound in EEF-like ancestry over a longer time period.

We combined the Patterson et al. (2022) UK data from 793 ancient individuals with the present day GBR 1000 genomes samples ("people of European ancestry from Great Britain (GBR)"), to form a time-series that runs from ∼5,500 years ago to the present day. Following Patterson et al. (2022) we broke the samples into 7 time periods corresponding to transitions in ancestry proportions (fig. 2C). We used the individual ancestry proportions of Patterson et al. (2022) inferred from a `qpAdm` three population model. These ancestry proportions are calculated to reflect genetic similarity to a set of samples that are pre-specified proxies for sources of ancestry. Note that an increase in a particular ancestry likely does not reflect gene flow directly from that source but rather from more nearby groups who themselves were already mixtures. In turn, each of the three major putative source ancestries were themselves a product of admixture in the past.

The covariance matrix of allele frequency changes between time periods is shown in fig. 2A. The UK time transect shows a number of large negative covariances between allele frequency change in the first time period (Δp_0, 5424-4005 years ago) and subsequent time periods (see also black points in fig. 3A). This negative covariance largely reflects that the initial large population turnover due to Steppe-like migration (during the first time period), was followed by an increase in EEF-like ancestry (fig. 2C, Patterson et al., 2022) generating allele frequency changes in the opposite direction. After correction by admixture, covariances are strongly reduced with only a small subset differing significantly from 0 (fig. 3C). In our UK time transect, most of the total variance in allele frequency change across time is due to admixture based changes, with 60% of allele frequency change being driven by the Steppe-like gene flow, only for the contribution of admixture to drop gradually as the EEF-like ancestry rebounds (fig. 3E, G and A). If we do not adjust for admixture, our estimate of the contribution of linked selection (G) is negative (fig. 3E, G_{nc}), reflecting the negative covariances induced by admixture. After adjusting for admixture there is no signal of a long-term contribution of linked selection, with G not departing from zero, as there is no consistent pattern of residual positive covariances.

Our empirical covariance results match those produced by a UK-like neutral simulation (Pearson and Durbin, 2023 model with UK matched admixture pulses, figs. S13 and S14).

We investigated another time transect from the Bohemia region (Papac et al., 2021) spanning from 5606 to 3037 years ago, that we also split in 7 time periods following the original paper (fig. 2D). The largest covariance is negative, and an order of magnitude larger than in the UK dataset (fig. 3B), and again seems to be due to the large influx from a Steppe-like source between 3880 and 3120 years ago (between time points 3 and 4, fig. 2D) and subsequent recovery of EEF-like ancestry (between Δp_3 and Δp_4, fig. 2D). Again this large covariance due to admixture can be corrected for (fig. 3D). While the older time points have a large amount of uncertainty, due to small sample sizes (section 4.3), nearly all of the variance in allele frequency change across the full time period is accounted by admixture ($A \sim 0.9$) and we see little evidence of allele frequency change attributable to linked selection in this Bohemian time-series (fig. 3F, G).

In sum, we see little evidence, in either transect, of linked selection in the covariances in allele frequency change between time intervals, suggesting that having accounted for admixture, much of the residual change across time intervals is due to drift-like sampling processes. One caveat is that if selection operates over short time scales, e.g. selection pressures fluctuate or deleterious alleles are quickly lost, selection could generate substantial allele frequency change (variance) within time intervals but little to no covariance between the time intervals we consider. To address this, we first reran our covariance analysis on the larger UK dataset splitting each time period in half. With this finer dissection of short-term covariances, we still see no evidence of covariance due to linked selection (fig. S9).

Linked selection is also expected to have larger effects in low recombination regions than high recombination regions, and in regions with a greater density of functional sites (patterns that are seen in long term human polymorphism datasets, Cai et al., 2009; Hernandez et al., 2011; McVicker et al., 2009). In both time transects, we do not observe any significant variation in the G and A statistics recalculated in bins of recombination rate (Bhérer et al., 2017) or proxies of the linked-selection effect of the density of functional sites (B-value, McVicker et al., 2009; Murphy et al., 2021) (figs. S7 and S11). However, in the UK transect we do see a significant increase in the total variance in allele frequency change in the lowest B-value bin (corresponding to the largest decrease in effective population size due to background selection, fig. 4A, first bin mean is above the genome-wide 95 % CI) and in the variances of change within some of the time intervals (fig. 4B). Noise in the smaller Bohemia time transect precludes seeing such effects (figs. S10 to S12). The allele frequency temporal variance increase in the UK for the lowest quintile B-value compared to the genome-wide mean is of 14.8 %, suggesting a fifth of this, $\sim$3 %, is an estimate of the contribution of linked selection to allele frequency change.

Figure 2: Human time-series covariance matrices and ancestries (UK left column, Bohemia right column). **(A-B)** Covariance matrices with covariances significantly different from zero marked with a star. The covariances have only been corrected for sampling bias and not admixture. **(C-D)** Mean ancestry proportions from the three reference populations in the time transects samples (the mean of sample ages in each period is used for the representation).

Figure 3: Human time-series covariance corrections and variance decomposition (UK left column, Bohemia right column). The top panels show the time intervals on the years BP axis, note that the two time series have different axes. All 95% CIs are computed through a block bootstrap procedure and represented with vertical bars. **(A-B)** Covariance values pre-admixture correction. Each line corresponds to the covariance between a first time interval (Δp_i, color code) and a later time interval (Δp_j, x-axis). **(C-D)** Covariance values post-admixture correction. **(E-F)** Proportion of the total variance, between the initial measured time (0) and time t, due to linked selection (G_{nc} for non-corrected and G) and to gene flow (A).

Figure 4: Temporal variances for the UK time-series binned by a proxy for the strength of linked selection (B-value). **(A)** Total temporal variance and sum of covariance for each quintile bin of B-values. The blue dash-dotted line and interval are the genome-wide mean and 95% confidence interval for total variance computed with 1/5 of windows sampled on each bootstrap to be comparable to the binned values. **(B)** Variances by bin for each time interval, normalized by heterozygosity. B-value quintile bins: [0.536-0.755), [0.755-0.849), [0.849-0.902), [0.902-0.944), [0.944-0.997).

3 Discussion

Here we have shown how ancient DNA data can be used to decompose the contribution of gene flow, linked selection, and drift to genome-wide allele frequency change. Using two ancient DNA time transects, our results demonstrate that gene flow is the dominant force changing allele frequencies in the recent history of European human populations, and that selection-driven change is not common across the genome. This does not necessarily contradict the number of signals of temporal selection found to date, as a small fraction of loci could be subject to strong selection (e.g. Irving-Pease et al., 2022; Ju & Mathieson, 2021; Le et al., 2022; Mathieson et al., 2015; Mathieson & Terhorst, 2022; Wilde et al., 2014). Indeed, some of these methods apply similar admixture adjustments as ours, but look for genome-wide outliers and so only detect strong selection on single loci (e.g. Mathieson and Terhorst (2022) expect to be able to detect selection coefficients > 0.02). Another set of approaches look for ancient selection on polygenic scores constructed from genome-wide association studies (Le et al., 2022; Mathieson & Terhorst, 2022). These approaches account for admixture and can detect subtle shifts at loci in ancient DNA, but rely on the fraction of genetic variation for specific traits captured in modern day samples (Yair & Coop, 2022). Thus, our genome-wide method is complementary to both time-series outlier approaches as well as phenotype-motivated approaches.

The large contribution of gene flow to evolutionary change in the past few thousand years is not surprising given the dynamic picture of population movement that has emerged from ancient DNA. The lack of a substantial contribution of linked selection is perhaps more surprising. Linked selection has been estimated to account for upward of 20 % reduction in long-term patterns of human diversity (under models of background selection, McVicker et al., 2009; Murphy et al., 2021), and so we should expect a similar portion of the variance in allele frequencies to come from linked selection. Much of this effect should manifest itself in the compounding of positive covariances between allele frequency change across the generations. While this effect has been seen empirically in selection experiments and in some natural populations, we currently do not see any evidence of this in humans. One possibility is that time periods we consider are not long enough for strong covariances to build up, as the long term patterns of linked selection reflect dynamics over coalescent time scales of hundreds of thousands of years. In contrast, the other possibility is that negative selection generating background selection is fast enough that it does not contribute to covariances among the time periods used here. However, under this latter interpretation, we should see higher allele frequency variances in regions predisposed to stronger linked selection, but we see this effect only weakly when partitioning loci by B-value. Larger collections of ancient DNA will allow better temporal resolution of allele frequency covariances, which could be combined with more individual-level approaches to avoid the need for sample lumping in time periods. It is also possible that some signals of linked selection may be washed out at the fine geographic scale of our time-series, as our time-series approach may partially be picking up ephemeral change which may average out over the much larger meta-population within which our time-series are embedded.

Our approach uses ancestry proportions called from ancient DNA for the three major inferred waves of gene flow into Europe. The sparsity of ancient DNA means that we rely on the ancestry proportions of relatively small samples of ancient individuals to be representative of people living in the past. However, the periods that we divide our samples into reflect reasonably well established periods in the peopling of the regions.

We also rely on allele frequencies in a set of samples as proxy for sources of gene flow. As we discuss below, the misspecification of the sources of gene flow may appear as evolutionary change within our focal time-series. One future extension might be to use principal components analysis to learn about major axes of population structure involving samples in a time-series and then to regress these PCAs out of our genotypes to account for variation in ancestry composition in a more model-free manner.

Our admixture correction seems to perform well on time intervals involving the large ancestry shift in Steppe-like and EEF-like ancestry (compare black points between fig. 3A and C). However, we see a number of negative covariances that remain after adjustment for admixture (fig. 3A and C). In principle, these could reflect fluctuating selection, but that seems unlikely given the general lack of other evidence of selection. Rather, these covariances could reflect that our proxies for gene flow sources only capture part of the allele frequency change driven by gene flow. Indeed, while we use a sample of European Early farmers to account for the increase in this ancestry in the UK, the most probable hypothesis for this increase is migration from mainland Europe from populations with a larger proportion of EEF-like ancestry than in the UK, but being itself already an admixed population with Steppe-like ancestry. More detailed modeling with admixture graphs and tree sequences could help better resolve the sources of gene flow in time-series (e.g. Allentoft et al., 2022; Irving-Pease et al., 2022; Pearson and Durbin, 2023, although such inferences may currently not be fully robust, Maier et al., 2023).

We attribute residual variance after accounting for gene flow and temporal covariances to drift-like processes. Genetic drift from the compounded sampling of parents to form each generation in our geographic area will obviously contribute to this. However, as our focal geographic areas are not homogeneous populations, small changes in ancestry composition over time not captured by larger scale admixture analyses might be captured as drift-like processes. Finally, as we take a fixed sample to reflect the allele frequencies in the sources of gene flow, change in the actual groups contributing to gene flow can also contribute to the signal of drift; e.g. if the allele frequencies in the source of EEF-like ancestry early differs from those contributing EEF-like ancestry later in the time-series, that will appear as drift-like change in our focal time-series. Our drift-like change in allele frequencies are small, corresponding to relative large estimates of temporal effective population sizes (see section 4.3). However, further work is needed to separate the long term effect of drift and combined contribution of other drift-like processes to our estimates of allele frequency change.

Extensions of our approach to larger geographical areas would allow the contributions of local genetic drift and migration among regions to be more fully explored. Such analyses would also pose an interesting set of modeling challenges to measure evolutionary change across spatially spread populations experiencing both local migration and more long-range gene flow events.

Finally, while a large body of ancient DNA work has focused on humans, ancient DNA datasets for a wide range of other organisms are also being generated (e.g. dogs, Bergström et al., 2020; horses, Orlando, 2020; sticklebacks, Kirch et al., 2021). The spread of ancient DNA and museum DNA research as well as more widespread usage of genome-wide sequencing to temporally monitor contemporary natural populations will generate a rich set of resources of time-series data. This offers the chance for comparative studies to decompose the contribution of different forces to genome-wide evolutionary change across systems, time-scales, and ecological and selection regimes.

4 Materials and Methods

4.1 Calculating the covariance matrix

We bin our samples into a set of discrete time points, and then calculate the allele frequency change at SNP l between adjacent time points, t and $t+1$, $\Delta p_{t,l}$. We then calculate the empirical variance-covariance matrix of these allele frequency changes for all time points averaged across SNPs. We denote the raw covariance matrix by $\mathbf{R}$. We wish to quantify the expected contribution of admixture to this matrix, but in doing so we also have to correct for sampling noise in both the time-series allele frequencies and sources of admixture. The corrected covariance matrix is given by

$$\mathbf{C} = (\mathbf{R} - \mathbf{S}_s) - (\mathbf{A} + \mathbf{S}_A) - \mathbf{D} \tag{5}$$

$\mathbf{S}_s$ is the expected matrix of biases from using sample frequencies in calculating the empirical covariance matrix (see eq. (6) below). Our admixture adjustment, $\mathbf{A}$, is the expected admixture variance-covariance matrix (see eq. (2)), where proxy samples are used as references for the admixture sources. The matrix $\mathbf{S}_A$ is the expected bias in the admixture matrix due to the sample noise from using sample frequencies in our admixture correlation (see eq. (7) below). Finally, $\mathbf{D}$ is the expected drift/admixture interaction matrix (see appendix C).

Here we calculate the sampling biases in the specific case of pseudohaploid data in line with the ancient DNA datasets considered in this paper (appendix A), using n_i for the haploid sample size at time point i and $\tilde{p}_{i,l}$ the sample frequency at SNP l. The sampling noise from taking a small sample of individuals inflates the variance of allele frequency change and shared sample between adjacent time points creates covariance

$$
\begin{aligned}
S_{s,i=j} &= \mathbb{E}\left(\frac{1}{n_{i,l}-1}\tilde{p}_{i,l}(1-\tilde{p}_{i,l}) + \frac{1}{n_{i+1,l}-1}\tilde{p}_{i+1,l}(1-\tilde{p}_{i+1,l}) \right) \\
S_{s,j-i=1} &= \mathbb{E}\left(-\frac{1}{n_{i+1,l}-1}\tilde{p}_{i+1,l}(1-\tilde{p}_{i+1,l}) \right)
\end{aligned}
\tag{6}
$$

with all other terms in the matrix set to zero (appendix A and Buffalo and Coop, 2019). These matrices are calculated as an average over all our SNPs. Second, sampling noise is also present in frequencies of the samples used as proxies for admixture, and so this biases the admixture expectation as the same reference samples are used for multiple time points (appendix B):

$$S_{A,i,j} = \mathbb{E}\left(\sum_r \Delta\alpha_{r,i}\Delta\alpha_{r,j}\frac{1}{n_{r,l}-1}\tilde{f}_{r,l}(1-\tilde{f}_{r,l}) \right) \tag{7}$$

following eq. (B.3) with $\alpha_{r,i}$ the admixture proportion from reference population r at time i, and $\tilde{f}_r$ the empirical allele frequency in the reference population r.

Finally, the simple admixture covariance expectation is missing a term due to shared drift variances between time intervals. This can be estimated as shown in appendix C and requires the assumption that only one parental population is contributing to gene flow during each time interval. $\mathbf{D}$ is given by eq. (C.9) and is a dependent on the estimated drift variance terms in parental populations and admixture proportions at each time steps common between two time points.

4.2 Simulations

We used the Demes format to write inter-operable demographic scenarios (Gower et al., 2022). This allowed us to run the same model with either `msprime` for neutral simulations (Baumdicker et al., 2022) or `SLiM` (v3.7) for simulations including selection (Haller & Messer, 2019). Results were recorded as treesequences and analyzed in `Python` using `tskit` (Kelleher et al., 2018). All results are based on 100 replicates of each scenario. The simulations pipeline was built with `snakemake` (Mölder et al., 2021) and can be found in the zenodo archive https://doi.org/10.5281/zenodo.8093105 and includes the version of all software used.

In the main text simple scenario, an ancestral population splits into the source populations 1,500 generations before present (BP). All populations are kept at a constant size of 10,000 diploid individuals. 200 generations BP our focal population that receives the admixture pulses is created from the first parental population (pop0). Pulses of admixture from pop1 happen at regular 20 generations interval starting at 150 and finishing at 10 generations BP. We sample 30 individuals 10 generations before and after pulses in our focal population. For our admixture sources we sample 30 individuals from each parental population at 200 generations BP for allele frequency computations. Samples are rendered pseudohaploid to mimic ancient DNA results (though no missing data was inserted). A census event of all populations is performed in the source populations when the admixed population is created to allow us to compute admixture proportions of all descendants.

We simulated a chromosome 100 Mbp in length with a mutation rate of 1×10^{-8} per bp and per generation and a uniform recombination rate of 2×10^{-8}.

To simulate linked-selection in `SLiM`, we considered three independent polygenic traits with alleles having a random effect size of ± 0.01 evolving under a model of stabilizing selection around an initial optimum of 0 for each trait. The fitness landscape is a gaussian function centered on the optimum with a variance, V_s, of 1. The optimum is gradually shifted from 0 to 3 standard deviations between 140 generations BP to the present similarly in all extent populations (by steps of shift/time). The ancestral population has a burn-in of 0.1N generations in SLiM and the complete ancestral history has been recapitated with `pyslim`. Mutation under selection are not used in the downstream analyses and neutral mutations are added a posteriori with `msprime`. We note that these simulations are not intended to mimic a particular selection scenario, as the density of loci underlying different traits per chromosome is unknown. Rather the parameters were chosen to generate results where both admixture and selection made comparable contributions for illustration purposes.

4.3 Ancient DNA analyses

We used two datasets from Patterson et al. (2022) and Papac et al. (2021) for ancient DNA time transects in the UK and the Bohemian region respectively. Data from those papers were downloaded from the indicated sources and merged with a set of parental population proxies and modern samples from the AADR v50.0 1240k dataset (Mallick et al., 2023 using data from Allentoft et al., 2015; Bergström et al., 2020; Brace et al., 2019; Consortium, 2015; Coutinho et al., 2020; Fu et al., 2016; Haak et al., 2015; Jones et al., 2015; Lamnidis et al., 2018; Lazaridis et al., 2017; Lazaridis et al., 2016; Lipson et al., 2017; Mallick et al., 2016; Margaryan et al., 2020; Martiniano et al., 2016; Mathieson et al., 2018; Mathieson et al., 2015; Mittnik et al., 2018; Narasimhan et al., 2019; Olalde

et al., 2018; Patterson et al., 2012; Raghavan et al., 2014; Rivollat et al., 2020; Scheib et al., 2019; Schroeder et al., 2019; Skoglund et al., 2015; Villalba-Mouco et al., 2019; Wang et al., 2019). Modern samples were used to provide a modern time point in the UK time transect. The data analysis `snakemake` pipeline can be found in the zenodo archive https://doi.org/10.5281/zenodo.8093107 and includes the version of all software used.

Individuals from each time period defined in the original analyses are pooled together to compute allele frequencies and the mean estimated age is taken as the time point date. For the UK based dataset (Patterson et al., 2022), we merged the published data with AADR v50.0 (providing modern samples and parental population proxies), and with data from Fowler et al. (2022) to access 10 individuals missing from the other datasets. Only loci with more than 10 genotypes in each time point grouping and more than 5 genotypes in reference populations were kept, resulting in 474,554 SNPs kept over the initial 1,135,618. We used combined filters 0 and 1 from Table S5 of Patterson et al. (2022) as our quality and relevance filtering. This resulted in sample sizes of [37, 69, 26, 23, 273, 38, 62] for periods labeled ['Neolithic', 'Chalcolithic/EBA', 'Middle Bronze Age', 'Late Bronze Age', 'Iron Age', 'Post Iron Age', 'Modern'] and mean non-missing genotypes across all SNPs of [25.5, 46.3, 19.6, 14.8, 208.2, 25.5, 59.9]. Mean sample dates for those periods are [5424, 4005, 3326, 2929, 2215, 1180, 0] years BP. Reference sample sizes are [18, 21, 18] for groups labeled ['WHGA', 'Balkan_N', 'OldSteppe'] in the dataset interpreted as WHG-like, EEF-like and Steppe-like. Those reference samples have mean non-missing genotypes across all SNPs of [7.4, 15.4, 12.4] respectively.

For the Papac et al. (2021) dataset, only loci with more than 2 genotypes in each time point grouping and more than 2 genotypes in reference samples were kept, resulting in 461,844 SNPs kept over the initial 1,150,639. Sample sizes in this dataset are [3, 5, 29, 14, 48, 59, 84] for periods labeled ['Neotithic', 'Proto-Eneolithic', 'Early Eneolithic', 'Middle Eneolithic', 'Corded Ware', 'Bell Beaker', 'Unetice'] and mean non-missing genotypes across all SNPs of [3., 3.7, 23.4, 10.9, 33.1, 41.9, 55.1]. Mean sample dates for those periods are [5607, 5253, 4229, 3880, 3726, 3120, 3037] years BP. Reference sample sizes are [4, 17, 15] for groups labeled ['WHG', 'Anatolia_Neolithic', 'Yamnaya'] in the dataset interpreted as WHG-like, EEF-like and Steppe-like. Those reference samples have mean non-missing genotypes across all SNPs of [3.5, 13.1, 9.0] respectively.

We used admixture measures from both published papers produced by the `qpAdm` method, extracted from the supplementary Table S5 from Patterson et al. (2022) and Table S9 from Papac et al. (2021). In concordance with the literature on European Human demographic history during the last 5000 years, we consider the simplest three way admixture between populations genetically most similar to European early farmers (EEF-like, early migrants from Anatolia), Western hunter gatherers (WHG-like) and individuals associated to the Steppe pastoralists Yamnaya culture (Steppe-like).

We computed confidence intervals around estimates by block bootstrap sampling of windows of 1000 SNPs along the genome. Statistics computed for each window where re-sampled 10^4 times with replacement and a 95% confidence interval was computed by the pivot method as in Buffalo and Coop (2020). Statistics were computed through a weighted average to account for variability in the number of SNPs in each window (windows at the end of chromosomes often do not contain the required number of SNPs). When dealing with ratio statistics (like G), we computed separately the numerators and denominators and used the ratio of the weighted averages for the final values.

Each dataset was transformed from the `eigenstrat` to the `sgkit` format through a `plink` (Chang et al., 2015) conversion step. Sex chromosomes were removed from the

datasets. To investigate correlation of our statistics with recombination or background selection, we incorporated in the dataset recombination rates (Bhérer et al., 2017, sex-averaged version) and B-values (Murphy et al., 2021) for each SNP – by using the value of the window the SNP was in. We split all SNPs into five quantile bins and computed G and A proportions for each one, as well as the variance and covariances.

We can compute a simple estimate of the diploid effective population size, $2N$, by equating the expected variance due to drift after t generations,

$$\mathrm{Var}(p_t) = \overline{p_0(1 - p_0)} \left[1 - \left(1 - \frac{1}{2N} \right)^t \right] \tag{8}$$

with the residual variances for each time period in the studied datasets (having adjusted for the variance due to admixture and sampling). Using a generation time of 30 years, and using the number of generations between the mid-points of each time interval for the UK dataset we obtain $2N = [$ 351472, 3204693, 2827982, 2041971, 361192, 203114 $]$ for each time interval. For the Bohemia dataset time intervals we get $2N = [$ 768209, 2576065, 1392381, 1181146, 1275489, 1167632 $]$. We note that these effective population size estimates are only approximate as they do not account for the more continuous distribution of sampling times present in the data.

Data availability

No new data was produced for this work. Analyses pipelines are available at doi: 10.5281/zenodo.8093105 for simulations and at doi: 10.5281/zenodo.8093107 for data. Those analyses rely on a custom helper python module available at doi: 10.5281/zenodo.8093101.

Acknowledgments

We thank members of the Coop lab and Vince Buffalo for helpful discussions. AS and GC were supported by the National Institute of General Medical Sciences of the National Institutes of Health (NIH R35 GM136290 to GC).

References

Allentoft, M. E., Sikora, M., Refoyo-Martínez, A., Irving-Pease, E. K., Fischer, A., Barrie, W., Ingason, A., Stenderup, J., Sjögren, K.-G., Pearson, A., Mota, B. S. d., Paulsson, B. S., Halgren, A., Macleod, R., Jørkov, M. L. S., Demeter, F., Novosolov, M., Sørensen, L., Nielsen, P. O., … Willerslev, E. (2022). Population genomics of stone age eurasia. *bioRxiv.* https://doi.org/10.1101/2022.05.04.490594

Allentoft, M. E., Sikora, M., Sjögren, K.-G., Rasmussen, S., Rasmussen, M., Stenderup, J., Damgaard, P. B., Schroeder, H., Ahlström, T., Vinner, L., Malaspinas, A.-S., Margaryan, A., Higham, T., Chivall, D., Lynnerup, N., Harvig, L., Baron, J., Casa, P. D., Dąbrowski, P., … Willerslev, E. (2015). Population genomics of bronze age eurasia. *Nature, 522*(7555), 167–172. https://doi.org/10.1038/nature14507

Barton, N. H. (2000). Genetic hitchhiking. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences, 355*(1403), 1553–1562. https://doi.org/10.1098/rstb.2000.0716

Baumdicker, F., Bisschop, G., Goldstein, D., Gower, G., Ragsdale, A. P., Tsambos, G., Zhu, S., Eldon, B., Ellerman, E. C., Galloway, J. G., Gladstein, A. L., Gorjanc, G., Guo, B., Jeffery, B., Kretzschumar, W. W., Lohse, K., Matschiner, M., Nelson, D., Pope, N. S., … Kelleher, J. (2022). Efficient ancestry and mutation simulation with msprime 1.0. *Genetics*, *220*(3), iyab229. https://doi.org/10.1093/genetics/iyab229

Bergström, A., Frantz, L., Schmidt, R., Ersmark, E., Lebrasseur, O., Girdland-Flink, L., Lin, A. T., Storå, J., Sjögren, K.-G., Anthony, D., Antipina, E., Amiri, S., Bar-Oz, G., Bazaliiskii, V. I., Bulatović, J., Brown, D., Carmagnini, A., Davy, T., Fedorov, S., … Skoglund, P. (2020). Origins and genetic legacy of prehistoric dogs. *Science*, *370*(6516), 557–564. https://doi.org/10.1126/science.aba9572

Bertram, J. (2021). Allele frequency divergence reveals ubiquitous influence of positive selection in drosophila. *PLOS Genetics*, *17*(9), e1009833. https://doi.org/10.1371/journal.pgen.1009833

Bhérer, C., Campbell, C. L., & Auton, A. (2017). Refined genetic maps reveal sexual dimorphism in human meiotic recombination at multiple scales. *Nature Communications*, *8*(1), 14994. https://doi.org/10.1038/ncomms14994

Brace, S., Diekmann, Y., Booth, T. J., van Dorp, L., Faltyskova, Z., Rohland, N., Mallick, S., Olalde, I., Ferry, M., Michel, M., Oppenheimer, J., Broomandkhoshbacht, N., Stewardson, K., Martiniano, R., Walsh, S., Kayser, M., Charlton, S., Hellenthal, G., Armit, I., … Barnes, I. (2019). Ancient genomes indicate population replacement in early neolithic britain. *Nature Ecology & Evolution*, *3*(5), 765–771. https://doi.org/10.1038/s41559-019-0871-9

Brennan, R. S., deMayo, J. A., Dam, H. G., Finiguerra, M., Baumann, H., Buffalo, V., & Pespeni, M. H. (2022). Experimental evolution reveals the synergistic genomic mechanisms of adaptation to ocean warming and acidification in a marine copepod. *Proceedings of the National Academy of Sciences*, *119*(38), e2201521119. https://doi.org/10.1073/pnas.2201521119

Buffalo, V., & Coop, G. (2019). The linked selection signature of rapid adaptation in temporal genomic data. *Genetics*, *213*, 1007–1045. https://doi.org/10.1534/genetics.119.302581

Buffalo, V., & Coop, G. (2020). Estimating the genome-wide contribution of selection to temporal allele frequency change. *Proceedings of the National Academy of Sciences*, *117*(34), 20672–20680. https://doi.org/10.1073/pnas.1919039117

Cai, J. J., Macpherson, J. M., Sella, G., & Petrov, D. A. (2009). Pervasive hitchhiking at coding and regulatory sites in humans. *PLoS Genetics*, *5*(1), e1000336. https://doi.org/10.1371/journal.pgen.1000336

Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-generation PLINK: Rising to the challenge of larger and richer datasets. *GigaScience*, *4*(1), s13742–015–0047–8. https://doi.org/10.1186/s13742-015-0047-8

Charlesworth, B., Morgan, M. T., & Charlesworth, D. (1993). The effect of deleterious mutations on neutral molecular variation. *134*, 1289–1303.

Consortium, 1. G. P. (2015). A global reference for human genetic variation. *Nature*, *526*(7571), 68–74. https://doi.org/10.1038/nature15393

Coop, G. (2016). Does linked selection explain the narrow range of genetic diversity across species? *bioRxiv*. https://doi.org/10.1101/042598

Coutinho, A., Günther, T., Munters, A. R., Svensson, E. M., Götherström, A., Storå, J., Malmström, H., & Jakobsson, M. (2020). The neolithic pitted ware culture

foragers were culturally but not genetically influenced by the battle axe culture herders. *American Journal of Physical Anthropology*, *172*(4), 638–649. https://doi.org/10.1002/ajpa.24079

Enard, D., Messer, P. W., & Petrov, D. A. (2014). Genome-wide signals of positive selection in human evolution. *Genome Research*, *24*(6), 885–895. https://doi.org/10.1101/gr.164822.113

Field, Y., Boyle, E. A., Telis, N., Gao, Z., Gaulton, K. J., Golan, D., Yengo, L., Rocheleau, G., Froguel, P., McCarthy, M. I., & Pritchard, J. K. (2016). Detection of human adaptation during the past 2000 years. *Science*, *354*(6313), 760–764. https://doi.org/10.1126/science.aag0776

Fowler, C., Olalde, I., Cummings, V., Armit, I., Büster, L., Cuthbert, S., Rohland, N., Cheronet, O., Pinhasi, R., & Reich, D. (2022). A high-resolution picture of kinship practices in an early neolithic tomb. *Nature*, *601*(7894), 584–587. https://doi.org/10.1038/s41586-021-04241-4

Fu, Q., Posth, C., Hajdinjak, M., Petr, M., Mallick, S., Fernandes, D., Furtwängler, A., Haak, W., Meyer, M., Mittnik, A., Nickel, B., Peltzer, A., Rohland, N., Slon, V., Talamo, S., Lazaridis, I., Lipson, M., Mathieson, I., Schiffels, S., … Reich, D. (2016). The genetic history of ice age europe. *Nature*, *534*(7606), 200–205. https://doi.org/10.1038/nature17993

Gower, G., Ragsdale, A. P., Bisschop, G., Gutenkunst, R. N., Hartfield, M., Noskova, E., Schiffels, S., Struck, T. J., Kelleher, J., & Thornton, K. R. (2022). Demes: A standard format for demographic models. *Genetics*, *222*(3), iyac131. https://doi.org/10.1093/genetics/iyac131

Haak, W., Lazaridis, I., Patterson, N., Rohland, N., Mallick, S., Llamas, B., Brandt, G., Nordenfelt, S., Harney, E., Stewardson, K., Fu, Q., Mittnik, A., Bánffy, E., Economou, C., Francken, M., Friederich, S., Pena, R. G., Hallgren, F., Khartanovich, V., … Reich, D. (2015). Massive migration from the steppe was a source for indo-european languages in europe. *Nature*, *522*(7555), 207–211. https://doi.org/10.1038/nature14317

Haller, B. C., & Messer, P. W. (2019). SLiM 3: Forward genetic simulations beyond the wright–fisher model (R. Hernandez, Ed.). *Molecular Biology and Evolution*, *36*(3), 632–637. https://doi.org/10.1093/molbev/msy228

He, Z., Dai, X., Lyu, W., Beaumont, M., & Yu, F. (2023). Estimating temporally variable selection intensity from ancient DNA data. *Molecular Biology and Evolution*, msad008. https://doi.org/10.1093/molbev/msad008

Hernandez, R. D., Kelley, J. L., Elyashiv, E., Melton, S. C., Auton, A., McVean, G., 1000 GENOMES PROJECT, Sella, G., & Przeworski, M. (2011). Classic selective sweeps were rare in recent human evolution. *Science*, *331*(6019), 920–924. https://doi.org/10.1126/science.1198878

Irving-Pease, E. K., Refoyo-Martínez, A., Ingason, A., Pearson, A., Fischer, A., Barrie, W., Sjögren, K.-G., Halgren, A. S., Macleod, R., Demeter, F., Henriksen, R. A., Vimala, T., McColl, H., Vaughn, A., Speidel, L., Stern, A. J., Scorrano, G., Ramsøe, A., Schork, A. J., … Willerslev, E. (2022). The selection landscape and genetic legacy of ancient eurasians. *bioRxiv*. https://doi.org/10.1101/2022.09.22.509027

Jones, E. R., Gonzalez-Fortes, G., Connell, S., Siska, V., Eriksson, A., Martiniano, R., McLaughlin, R. L., Gallego Llorente, M., Cassidy, L. M., Gamba, C., Meshveliani, T., Bar-Yosef, O., Müller, W., Belfer-Cohen, A., Matskevich, Z., Jakeli, N., Higham, T. F. G., Currat, M., Lordkipanidze, D., … Bradley, D. G. (2015). Upper palae-

olithic genomes reveal deep roots of modern eurasians. *Nature Communications*, *6*(1), 8912. https://doi.org/10.1038/ncomms9912

Ju, D., & Mathieson, I. (2021). The evolution of skin pigmentation-associated variation in west eurasia. *Proceedings of the National Academy of Sciences*, *118*(1), e2009227118. https://doi.org/10.1073/pnas.2009227118

Kaplan, N. L., Hudson, R. R., & Langley, C. H. (1989). The "hitchhiking effect" revisited. *Genetics*, *123*(4), 887–899. https://doi.org/10.1093/genetics/123.4.887

Kelleher, J., Thornton, K. R., Ashander, J., & Ralph, P. L. (2018). Efficient pedigree recording for fast population genetics simulation. *PLOS Computational Biology*, *14*(11), e1006581. https://doi.org/10.1371/journal.pcbi.1006581

Kelly, J. K. (2022). The genomic scale of fluctuating selection in a natural plant population. *Evolution Letters*, *6*(6), 506–521. https://doi.org/10.1002/evl3.308

Kirch, M., Romundset, A., Gilbert, M. T. P., Jones, F. C., & Foote, A. D. (2021). Ancient and modern stickleback genomes reveal the demographic constraints on adaptation. *Current Biology*, *31*(9), 2027–2036.e8. https://doi.org/10.1016/j.cub.2021.02.027

Lamnidis, T. C., Majander, K., Jeong, C., Salmela, E., Wessman, A., Moiseyev, V., Khartanovich, V., Balanovsky, O., Ongyerth, M., Weihmann, A., Sajantila, A., Kelso, J., Pääbo, S., Onkamo, P., Haak, W., Krause, J., & Schiffels, S. (2018). Ancient fennoscandian genomes reveal origin and spread of siberian ancestry in europe. *Nature Communications*, *9*(1), 5018. https://doi.org/10.1038/s41467-018-07483-5

Lazaridis, I., Mittnik, A., Patterson, N., Mallick, S., Rohland, N., Pfrengle, S., Furtwängler, A., Peltzer, A., Posth, C., Vasilakis, A., McGeorge, P. J. P., Konsolaki-Yannopoulou, E., Korres, G., Martlew, H., Michalodimitrakis, M., Özsait, M., Özsait, N., Papathanasiou, A., Richards, M., … Stamatoyannopoulos, G. (2017). Genetic origins of the minoans and mycenaeans. *Nature*, *548*(7666), 214–218. https://doi.org/10.1038/nature23310

Lazaridis, I., Nadel, D., Rollefson, G., Merrett, D. C., Rohland, N., Mallick, S., Fernandes, D., Novak, M., Gamarra, B., Sirak, K., Connell, S., Stewardson, K., Harney, E., Fu, Q., Gonzalez-Fortes, G., Jones, E. R., Roodenberg, S. A., Lengyel, G., Bocquentin, F., … Reich, D. (2016). Genomic insights into the origin of farming in the ancient near east. *Nature*, *536*(7617), 419–424. https://doi.org/10.1038/nature19310

Le, M. K., Smith, O. S., Akbari, A., Harpak, A., Reich, D., & Narasimhan, V. M. (2022). 1,000 ancient genomes uncover 10,000 years of natural selection in europe. https://doi.org/10.1101/2022.08.24.505188

Lipson, M., Szécsényi-Nagy, A., Mallick, S., Pósa, A., Stégmár, B., Keerl, V., Rohland, N., Stewardson, K., Ferry, M., Michel, M., Oppenheimer, J., Broomandkhoshbacht, N., Harney, E., Nordenfelt, S., Llamas, B., Gusztáv Mende, B., Köhler, K., Oross, K., Bondár, M., … Reich, D. (2017). Parallel palaeogenomic transects reveal complex genetic history of early european farmers. *Nature*, *551*(7680), 368–372. https://doi.org/10.1038/nature24476

Maier, R., Flegontov, P., Flegontova, O., Isildak, U., Changmai, P., & Reich, D. (2023). On the limits of fitting complex models of population history to f-statistics (M. Nordborg, Ed.). *eLife*, *12*, e85492. https://doi.org/10.7554/eLife.85492

Mallick, S., Li, H., Lipson, M., Mathieson, I., Gymrek, M., Racimo, F., Zhao, M., Chennagiri, N., Nordenfelt, S., Tandon, A., Skoglund, P., Lazaridis, I., Sankararaman, S., Fu, Q., Rohland, N., Renaud, G., Erlich, Y., Willems, T., Gallo, C., … Reich,

D. (2016). The simons genome diversity project: 300 genomes from 142 diverse populations. *Nature*, *538*(7624), 201–206. https://doi.org/10.1038/nature18964

Mallick, S., Micco, A., Mah, M., Ringbauer, H., Lazaridis, I., Olalde, I., Patterson, N. J., & Reich, D. E. (2023). The allen ancient DNA resource (AADR): A curated compendium of ancient human genomes. https://doi.org/10.1101/2023.04.06.535797

Margaryan, A., Lawson, D. J., Sikora, M., Racimo, F., Rasmussen, S., Moltke, I., Cassidy, L. M., Jørsboe, E., Ingason, A., Pedersen, M. W., Korneliussen, T., Wilhelmson, H., Buś, M. M., de Barros Damgaard, P., Martiniano, R., Renaud, G., Bhérer, C., Moreno-Mayar, J. V., Fotakis, A. K., … Willerslev, E. (2020). Population genomics of the viking world. *Nature*, *585*(7825), 390–396. https://doi.org/10.1038/s41586-020-2688-8

Martiniano, R., Caffell, A., Holst, M., Hunter-Mann, K., Montgomery, J., Müldner, G., McLaughlin, R. L., Teasdale, M. D., van Rheenen, W., Veldink, J. H., van den Berg, L. H., Hardiman, O., Carroll, M., Roskams, S., Oxley, J., Morgan, C., Thomas, M. G., Barnes, I., McDonnell, C., … Bradley, D. G. (2016). Genomic signals of migration and continuity in britain before the anglo-saxons. *Nature Communications*, *7*(1), 10326. https://doi.org/10.1038/ncomms10326

Mathieson, I., Alpaslan-Roodenberg, S., Posth, C., Szécsényi-Nagy, A., Rohland, N., Mallick, S., Olalde, I., Broomandkhoshbacht, N., Candilio, F., Cheronet, O., Fernandes, D., Ferry, M., Gamarra, B., Fortes, G. G., Haak, W., Harney, E., Jones, E., Keating, D., Krause-Kyora, B., … Reich, D. (2018). The genomic history of southeastern europe. *Nature*, *555*(7695), 197–203. https://doi.org/10.1038/nature25778

Mathieson, I., Lazaridis, I., Rohland, N., Mallick, S., Patterson, N., Roodenberg, S. A., Harney, E., Stewardson, K., Fernandes, D., Novak, M., Sirak, K., Gamba, C., Jones, E. R., Llamas, B., Dryomov, S., Pickrell, J., Arsuaga, J. L., de Castro, J. M. B., Carbonell, E., … Reich, D. (2015). Genome-wide patterns of selection in 230 ancient eurasians. *Nature*, *528*(7583), 499–503. https://doi.org/10.1038/nature16152

Mathieson, I., & Terhorst, J. (2022). Direct detection of natural selection in bronze age britain. *Genome Research*, *32*(11), 2057–2067. https://doi.org/10.1101/gr.276862.122

Maynard-Smith, J., & Haigh, J. (1974). The hitch-hiking effect of a favourable gene. *Genetical Research*, *23*(1), 23–35. https://doi.org/10.1017/S0016672300014634

McVicker, G., Gordon, D., Davis, C., & Green, P. (2009). Widespread genomic signatures of natural selection in hominid evolution. *PLOS Genetics*, *5*(5), e1000471. https://doi.org/10.1371/journal.pgen.1000471

Mittnik, A., Wang, C.-C., Pfrengle, S., Daubaras, M., Zariņa, G., Hallgren, F., Allmäe, R., Khartanovich, V., Moiseyev, V., Tõrv, M., Furtwängler, A., Andrades Valtueña, A., Feldman, M., Economou, C., Oinonen, M., Vasks, A., Balanovska, E., Reich, D., Jankauskas, R., … Krause, J. (2018). The genetic prehistory of the baltic sea region. *Nature Communications*, *9*(1), 442. https://doi.org/10.1038/s41467-018-02825-9

Mölder, F., Jablonski, K., Letcher, B., Hall, M., Tomkins-Tinch, C., Sochat, V., Forster, J., Lee, S., Twardziok, S., Kanitz, A., Wilm, A., Holtgrewe, M., Rahmann, S., Nahnsen, S., & Köster, J. (2021). Sustainable data analysis with snakemake. *F1000Research*, *10*(33). https://doi.org/10.12688/f1000research.29032.2

Murphy, D., Elyashiv, E., Amster, G., & Sella, G. (2021). Broad-scale variation in human genetic diversity levels is predicted by purifying selection on coding and non-coding elements. https://doi.org/10.1101/2021.07.02.450762

Narasimhan, V. M., Patterson, N., Moorjani, P., Rohland, N., Bernardos, R., Mallick, S., Lazaridis, I., Nakatsuka, N., Olalde, I., Lipson, M., Kim, A. M., Olivieri, L. M., Coppa, A., Vidale, M., Mallory, J., Moiseyev, V., Kitov, E., Monge, J., Adamski, N., … Reich, D. (2019). The formation of human populations in south and central asia. *Science*, *365*(6457), eaat7487. https://doi.org/10.1126/science.aat7487

Olalde, I., Brace, S., Allentoft, M. E., Armit, I., Kristiansen, K., Booth, T., Rohland, N., Mallick, S., Szécsényi-Nagy, A., Mittnik, A., Altena, E., Lipson, M., Lazaridis, I., Harper, T. K., Patterson, N., Broomandkhoshbacht, N., Diekmann, Y., Faltyskova, Z., Fernandes, D., … Reich, D. (2018). The beaker phenomenon and the genomic transformation of northwest europe. *Nature*, *555*(7695), 190–196. https://doi.org/10.1038/nature25738

Orlando, L. (2020). The evolutionary and historical foundation of the modern horse: Lessons from ancient genomics. *Annual Review of Genetics*, *54*(1), 563–581. https://doi.org/10.1146/annurev-genet-021920-011805

Papac, L., Ernée, M., Dobeš, M., Langová, M., Rohrlach, A. B., Aron, F., Neumann, G. U., Spyrou, M. A., Rohland, N., Velemínský, P., Kuna, M., Brzobohatá, H., Culleton, B., Daněček, D., Danielisová, A., Dobisíková, M., Hložek, J., Kennett, D. J., Klementová, J., … Haak, W. (2021). Dynamic changes in genomic and social structures in third millennium BCE central europe. *Science Advances*, *7*(35), eabi6941. https://doi.org/10.1126/sciadv.abi6941

Patterson, N., Isakov, M., Booth, T., Büster, L., Fischer, C.-E., Olalde, I., Ringbauer, H., Akbari, A., Cheronet, O., Bleasdale, M., Adamski, N., Altena, E., Bernardos, R., Brace, S., Broomandkhoshbacht, N., Callan, K., Candilio, F., Culleton, B., Curtis, E., … Reich, D. (2022). Large-scale migration into britain during the middle to late bronze age. *Nature*, *601*(7894), 588–594. https://doi.org/10.1038/s41586-021-04287-4

Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., Genschoreck, T., Webster, T., & Reich, D. (2012). Ancient admixture in human history. *Genetics*, *192*(3), 1065–1093. https://doi.org/10.1534/genetics.112.145037

Pearson, A., & Durbin, R. (2023). Local ancestry inference for complex population histories. *bioRxiv*. https://doi.org/10.1101/2023.03.06.529121

Raghavan, M., Skoglund, P., Graf, K. E., Metspalu, M., Albrechtsen, A., Moltke, I., Rasmussen, S., Stafford Jr, T. W., Orlando, L., Metspalu, E., Karmin, M., Tambets, K., Rootsi, S., Mägi, R., Campos, P. F., Balanovska, E., Balanovsky, O., Khusnutdinova, E., Litvinov, S., … Willerslev, E. (2014). Upper palaeolithic siberian genome reveals dual ancestry of native americans. *Nature*, *505*(7481), 87–91. https://doi.org/10.1038/nature12736

Rivollat, M., Jeong, C., Schiffels, S., Küçükkalıpçı, İ., Pemonge, M.-H., Rohrlach, A. B., Alt, K. W., Binder, D., Friederich, S., Ghesquière, E., Gronenborn, D., Laporte, L., Lefranc, P., Meller, H., Réveillas, H., Rosenstock, E., Rottier, S., Scarre, C., Soler, L., … Haak, W. (2020). Ancient genome-wide DNA from france highlights the complexity of interactions between mesolithic hunter-gatherers and neolithic farmers. *Science Advances*, *6*(22), eaaz5344. https://doi.org/10.1126/sciadv.aaz5344

Saleh, D., Chen, J., Leplé, J.-C., Leroy, T., Truffaut, L., Dencausse, B., Lalanne, C., Labadie, K., Lesur, I., Bert, D., Lagane, F., Morneau, F., Aury, J.-M., Plomion, C., Lascoux, M., & Kremer, A. (2022). Genome-wide evolutionary response of european oaks during the anthropocene. *Evolution Letters*, *6*(1), 4–20. https://doi.org/10.1002/evl3.269

Santiago, E., & Caballero, A. (1998). Effective size and polymorphism of linked neutral loci in populations under directional selection. *Genetics*, *149*(4), 2105–2117. https://doi.org/10.1093/genetics/149.4.2105

Scheib, C. L., Hui, R., D'Atanasio, E., Wohns, A. W., Inskip, S. A., Rose, A., Cessford, C., O'Connell, T. C., Robb, J. E., Evans, C., Patten, R., & Kivisild, T. (2019). East anglian early neolithic monument burial linked to contemporary megaliths. *Annals of Human Biology*, *46*(2), 145–149. https://doi.org/10.1080/03014460.2019.1623912

Schrider, D. R., & Kern, A. D. (2017). Soft sweeps are the dominant mode of adaptation in the human genome. *Molecular Biology and Evolution*, *34*(8), 1863–1877. https://doi.org/10.1093/molbev/msx154

Schroeder, H., Margaryan, A., Szmyt, M., Theulot, B., Włodarczak, P., Rasmussen, S., Gopalakrishnan, S., Szczepanek, A., Konopka, T., Jensen, T. Z. T., Witkowska, B., Wilk, S., Przybyła, M. M., Pospieszny, Ł., Sjögren, K.-G., Belka, Z., Olsen, J., Kristiansen, K., Willerslev, E., … Allentoft, M. E. (2019). Unraveling ancestry, kinship, and violence in a late neolithic mass grave. *Proceedings of the National Academy of Sciences*, *116*(22), 10705–10710. https://doi.org/10.1073/pnas.1820210116

Sella, G., Petrov, D. A., Przeworski, M., & Andolfatto, P. (2009). Pervasive natural selection in the drosophila genome? *PLOS Genetics*, *5*(6), e1000495. https://doi.org/10.1371/journal.pgen.1000495

Skoglund, P., Mallick, S., Bortolini, M. C., Chennagiri, N., Hünemeier, T., Petzl-Erler, M. L., Salzano, F. M., Patterson, N., & Reich, D. (2015). Genetic evidence for two founding populations of the americas. *Nature*, *525*(7567), 104–108. https://doi.org/10.1038/nature14895

Souilmi, Y., Tobler, R., Johar, A., Williams, M., Grey, S. T., Schmidt, J., Teixeira, J. C., Rohrlach, A., Tuke, J., Johnson, O., Gower, G., Turney, C., Cox, M., Cooper, A., & Huber, C. D. (2022). Admixture has obscured signals of historical hard sweeps in humans. *Nature Ecology & Evolution*, 1–13. https://doi.org/10.1038/s41559-022-01914-9

Villalba-Mouco, V., van de Loosdrecht, M. S., Posth, C., Mora, R., Martínez-Moreno, J., Rojo-Guerra, M., Salazar-García, D. C., Royo-Guillén, J. I., Kunst, M., Rougier, H., Crevecoeur, I., Arcusa-Magallón, H., Tejedor-Rodríguez, C., García-Martínez de Lagrán, I., Garrido-Pena, R., Alt, K. W., Jeong, C., Schiffels, S., Utrilla, P., … Haak, W. (2019). Survival of late pleistocene hunter-gatherer ancestry in the iberian peninsula. *Current Biology*, *29*(7), 1169–1177.e7. https://doi.org/10.1016/j.cub.2019.02.006

Wang, J., Gaughan, S., Lamer, J. T., Deng, C., Hu, W., Wachholtz, M., Qin, S., Nie, H., Liao, X., Ling, Q., Li, W., Zhu, L., Bernatchez, L., Wang, C., & Lu, G. (2019). Resolving the genetic paradox of invasions: Preadapted genomes and post-introduction hybridization of bigheaded carps in the mississippi river basin. *Evolutionary Applications*, eva.12863. https://doi.org/10.1111/eva.12863

Wilde, S., Timpson, A., Kirsanow, K., Kaiser, E., Kayser, M., Unterländer, M., Hollfelder, N., Potekhina, I. D., Schier, W., Thomas, M. G., & Burger, J. (2014). Direct evidence for positive selection of skin, hair, and eye pigmentation in europeans during the last 5,000 y. *Proceedings of the National Academy of Sciences*, *111*(13), 4832–4837. https://doi.org/10.1073/pnas.1316513111

Yair, S., & Coop, G. (2022). Population differentiation of polygenic score predictions under stabilizing selection. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *377*(1852), 20200416. https://doi.org/10.1098/rstb.2020.0416

Appendix

A Pseudohaploid sampling

Following Buffalo and Coop (2020), the observed variance in allele frequency at time t can be decomposed with the law of total variance:

$$\text{Var}(\tilde{p}_t) = \mathbb{E}(\text{Var}(\tilde{p}_t|p_t)) + \text{Var}(\mathbb{E}(\tilde{p}_t|p_t)) \tag{A.1}$$

$$= \mathbb{E}(\text{Var}(\tilde{p}_t|p_t)) + \text{Var}(p_t) \tag{A.2}$$

This gives us a way to correct the observed variance for sampling noise.

Pseudohaploid representation is common in ancient DNA data to avoid errors when calling heterozygotes. Most often, one read (and therefore allele) is selected randomly among the mapped reads for each individual at a given position. Pseudohaploid calling can be modeled as a binomial sampling. We consider sampling n_t individuals in a population with frequency of the alternate allele p_t at time t. Pseudohaploid calling is equivalent to each individual drawing one allele from the pool of alleles. We define $X_t \sim \text{Binom}(n_t, p_t)$ the number of alternate alleles sampled and $\tilde{p}_t = X_t/n_t$. Then the sampling noise is

$$\text{Var}(\tilde{p}_t|p_t) = \frac{1}{n_t^2}\text{Var}(X_t) \tag{A.3}$$

$$= \frac{1}{n_t}p_t(1-p_t) \tag{A.4}$$

Correction of the variance $\text{Var}(\Delta\tilde{p}_t)$ is carried out by subtracting both sampling variances $\text{Var}(\tilde{p}_t|p_t)$ and $\text{Var}(\tilde{p}_{t+1}|p_{t+1})$. As in Buffalo and Coop (2020), the covariances between two overlapping time intervals, $\text{Cov}(\Delta\tilde{p}_t, \Delta\tilde{p}_{t+1})$ are negatively biased by the shared sampling noise in p_{t+1}, and this needs to be corrected by adding the shared time point sampling variance $\text{Var}(\tilde{p}_{t+1}|p_{t+1})$ back in. For these corrections we need an unbiased estimator of the half heterozygosity. We define the sample heterozygosity as $\tilde{H}/2 = \tilde{p}(1-\tilde{p})$, then

$$\mathbb{E}(\tilde{H}/2) = p(1-p)\left(\frac{n-1}{n}\right) \tag{A.5}$$

Therefore

$$\text{Var}(\tilde{p}_t|p_t) = \frac{1}{n_t-1}\tilde{p}_t(1-\tilde{p}_t) \tag{A.6}$$

Similarly, if needed, we can compute the diploid sampling bias estimator:

$$\text{Var}(\tilde{p}_t|p_t) = \frac{1}{2n_t-1}\tilde{p}_t(1-\tilde{p}_t) \tag{A.7}$$

B Pseudohaploid sampling noise in reference populations

Let's consider the allele frequency of reference population r

$$\tilde{f}_r = f_r + \delta f_r \tag{B.1}$$

The observed allele frequency is equal to the true allele frequency (f_r) plus sampling noise.

Therefore

$$\Delta p_{A,i} = \sum_{r=1}^{R} \Delta \bar{\alpha}_{i,r}(\tilde{f}_r - \delta f_r) = \Delta \tilde{p}_{A,i} - \Delta \delta f_{r,i} \tag{B.2}$$

Decomposing $\text{Cov}(\Delta p_{A,i}, \Delta p_{A,j})$ as $\mathbb{E}(\Delta p_{A,i} \Delta p_{A,j}) - \mathbb{E}(\Delta p_{A,i})\,\mathbb{E}(\Delta p_{A,j})$ and remembering that $\mathbb{E}(\Delta \delta f_{r,i}) = 0$, we end up with $\text{Cov}(\Delta p_{A,i}, \Delta p_{A,j}) = \text{Cov}(\Delta \tilde{p}_{A,i}, \Delta \tilde{p}_{A,j}) + \mathbb{E}(\Delta \delta f_{r,i} \Delta \delta f_{r,j})$

Therefore

$$\text{Cov}(\Delta p_{A,i}, \Delta p_{A,j}) = \text{Cov}(\Delta \tilde{p}_{A,i}, \Delta \tilde{p}_{A,j}) + \sum_{r} \Delta \alpha_{r,i} \Delta \alpha_{r,j} \, \text{Var}(\delta f_r) \tag{B.3}$$

with $\text{Var}(\delta f_r)$, the variance of sampling noise in the pseudohaploid case, equal to $\frac{1}{n_r - 1}\tilde{f}_r(1 - \tilde{f}_r)$ (similar to appendix A).

C Accounting for drift in the admixture correction

We consider a simple model where only the focal admixed population experiences drift and parental populations from which gene flow occurs are not. This is in line with our use of a single proxy sample for the sources of admixture. Under this model drift happening in any of our contributing populations is absorbed into the drift observed in the focal population. Drift that occurs in one time interval can partially be erased by admixture in subsequent time intervals. This interaction between drift and gene flow generates additional covariance that needs to be accounted for.

Let f_r be the frequencies of R parental populations for a particular SNP. At time 0, an admixed population of frequency p_0 is established with ancestry proportions q_{k0} from any of the R populations. Subsequently, between time points t and $t + 1$, this admixed population can receive a migration pulse from any of the R populations, where a proportion $\alpha_{r,t}$ of individuals in the focal population are replaced by migrants. Between time intervals t and $t + 1$, drift happens changing the frequency by Δd_t. p_t is the allele frequency at time t of our admixed population. $q_{r,t}$ are the ancestry proportions at time t of this admixed population.

We define the proportion of individuals replaced by admixture as:

$$A_t = \sum_{r} \alpha_{r,t} \quad , \text{ with } 0 \leq A_t \leq 1 \tag{C.1}$$

The change in allele frequency can then be written as:

$$p_{t+1} = (1 - A_t)p_t + \sum_{r} \alpha_{r,t} f_r + \Delta d_t \tag{C.2}$$

$$\Delta p_t = p_{t+1} - p_t \tag{C.3}$$

$$= -A_t p_t + \sum_{r} \alpha_{r,t} f_r + \Delta d_t \tag{C.4}$$

We can expand this out in terms of the change in allele frequency due to admixture and drift in the preceding time periods:

$$\Delta p_t = \sum_r \left[\alpha_{r,t} - A_t q_{r,t} \right] f_r$$
$$+ \Delta d_t - A_t \left[\Delta d_{t-1} + \sum_{k=0}^{t-2} \prod_{l=k+1}^{t-1} (1 - A_t) \Delta d_k \right] \tag{C.5}$$

We can express the ancestry fraction from source r at time in terms of the change due to admixture in previous time periods:

$$q_{r,t} = \alpha_{r,t-1} + \sum_{k=0}^{t-2} \alpha_{r,k} \prod_{l=k+1}^{t-1} (1 - A_l) \tag{C.6}$$

$$q_{r,t+1} = \alpha_{r,t} + (1 - A_t) q_{r,t}$$
$$\Delta q_{r,t} = \alpha_{r,t} - A_t q_{r,t} \tag{C.7}$$

As a constraint, if at each time step there is only one $\alpha_{rt} > 0$, then it simplifies to:

$$\Delta q_{r,t} = \alpha_{r,t} - \alpha_{r,t} q_{r,t}$$
$$\alpha_{r,t} = \frac{\Delta q_{r,t}}{1 - q_{r,t}} \tag{C.8}$$

allowing us to compute the admixture fraction from the ancestry proportions.

When computing the covariance between two time intervals i and j ($i < j$), a composite drift term depending on the admixture pulses will be shared between all time intervals between times 0 to i between the two Δp_i and Δp_j. This expected admixture-drift term D can be computed as:

$$D(\Delta p_i, \Delta p_j)_{i<j} = \sum_{k=0}^{i} \left[(-A_i)^{\mathbf{1}_{\mathbb{Z}_{>0}}(i-k)} \left[\prod_{l=k+1}^{i-1} (1 - A_l) \right]^{\mathbf{1}_{\mathbb{Z}_{>0}}(i-k-1)} \right.$$
$$\left. \times (-A_j) \left[\prod_{l=k+1}^{j-1} (1 - A_l) \right]^{\mathbf{1}_{\mathbb{Z}_{>0}}(j-k-1)} \mathrm{Var}(\Delta d_k) \right] \tag{C.9}$$

with

$$\mathbf{1}_{\mathbb{Z}_{>0}}(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{else} \end{cases} \tag{C.10}$$

D can then be subtracted from the empirical covariance to remove this effect.

To compute D, individual time interval drift variances need to be estimated ($\mathrm{Var}(\Delta d_k)$). We can use the fact that the variances at each time interval can be decomposed as a linear combination of drift variances to solve for them. Solving is only possible when considering that only one parental population is contributing to gene flow at a given time to be able to estimate the values of individual α_r (eq. (C.8)). The system for $0 \le i \le t$ is of the form:

$$\mathrm{Var}(\Delta p_i) = \mathrm{Var} \left(\sum_r \left[\alpha_{r,t} - A_t q_{r,t} \right] f_r \right) + \mathrm{Var}(\Delta d_i)$$
$$+ A_i^2 \left[\mathrm{Var}(\Delta d_{i-1}) + \sum_{k=0}^{i-2} \prod_{l=k+1}^{i-1} (1 - A_l)^2 \, \mathrm{Var}(\Delta d_k) \right] \tag{C.11}$$