***GIGANTEA* accelerates wheat heading time through gene interactions**

converging on *FLOWERING LOCUS T1*

Chengxia Li[1, 2*], Huiqiong Lin[1, 2], Juan M. Debernardi[1, 2], Chaozhong Zhang[1, 2] and Jorge

Dubcovsky[1, 2*]

[1] Department of Plant Sciences, University of California, Davis, CA 95616

[2] Howard Hughes Medical Institute, Chevy Chase, MD 20815, USA.

Chengxia Li: ORCID 0000-0003-4634-8451. Email: chxli@ucdavis.edu

Huiqiong Lin: ORCID 0000-0003-1833-2559. Email: hqlin@ucdavis.edu

Juan M. Debernardi: ORCID 0000-0002-4591-7061. Email: jmdebernardi@ucdavis.edu

Chaozhong Zhang: ORCID 0000-0002-0247-3771. Email: cazhang@ucdavis.edu

Jorge Dubcovsky: ORCID 0000-0002-7571-4345. Email: jdubcovsky@ucdavis.edu

16
17 **Running head**: *GIGANTEA* interactions regulate wheat heading time
18
19

* Corresponding authors: Jorge Dubcovsky. Email: jdubcovsky@ucdavis.edu

Chengxia Li. Email: chxli@ucdavis.edu

Abstract

Precise regulation of flowering time is critical for cereal crops to synchronize reproductive

development with optimum environmental conditions, thereby maximizing grain yield. The plant

specific gene *GIGANTEA* (*GI*) plays an important role in the control of flowering time, with

additional functions on the circadian clock and plant stress responses. In this study, we show that

GI loss-of-function mutants in a photoperiod sensitive tetraploid wheat background exhibit

significant delays in heading time under both long-day (LD) and short-day (SD) photoperiods,

with stronger effects under LD. However, this interaction between GI and photoperiod is no

longer observed in isogenic lines carrying either a photoperiod insensitive allele in the

PHOTOPERIOD1 (*PPD1*) gene or a loss-of-function allele in *EARLY FLOWERING 3* (*ELF3*), a

known repressor of *PPD1*. These results suggest that the normal circadian regulation of *PPD1* is

required for the differential effect of *GI* on heading time in different photoperiods. Using crosses

between mutants or transgenic of *GI* and those of critical genes in the flowering regulation

pathway, we show that *GI* accelerates wheat heading time by promoting *FLOWERING LOCUS*

T1 (*FT1*) expression via interactions with *ELF3, VERNALIZATION 2* (*VRN2*), *CONSTANS* (*CO*),

and the age-dependent microRNA172-*APETALA2* (*AP2*) pathway, at both transcriptional and

protein levels. Our study reveals conserved *GI* mechanisms between wheat and Arabidopsis, but

also identifies specific interactions of GI with the distinctive photoperiod and vernalization

pathways of the temperate grasses. These results provide valuable knowledge for modulating

wheat heading time and engineering new varieties better adapted to a changing environment.

Introduction

GIGANTEA (*GI*) is a plant-specific gene that encodes a large protein (1,173 amino acids in Arabidopsis) with no domains of known biochemical function (Fowler et al. 1999). The GI protein is predominantly localized to the nuclei where it forms nuclear bodies (Huq et al. 2000). Both *GI* transcription and GI protein abundance are regulated by the circadian clock (Fowler et al. 1999; Park et al. 1999; David et al. 2006) and can be detected throughout plant development (Fowler et al. 1999). In Arabidopsis plants grown under long-day photoperiod (16 h light/8 h dark, henceforth LD), the *GI* transcripts peak at Zeitgeber time 10 (ZT10); while under short-day (8 h light/16 h dark, henceforth SD), the peak is at ZT8 (Fowler et al. 1999; Park et al. 1999).

Defects in several components of the circadian clock have been shown to affect the *GI* transcription profile (Fowler et al. 1999; Lu et al. 2012). Chromatin Immunoprecipitation (ChIP) analysis showed that the evening complex (EC), including EARLY FLOWERING 3 (ELF3), EARLY FLOWERING 4 (ELF4) and LUX ARRYTHMO (LUX), regulates *GI* transcription by direct binding to its promoter (Ezer et al. 2017). ELF3 can also regulate GI protein degradation through its interaction with CONSTITUTIVELY PHOTOMORPHOGENIC1 (COP1) in an EC-independent manner (Yu et al. 2008). The Arabidopsis *elf3 gi* double mutant is unable to synchronize the circadian oscillator to the light-dark cycles, indicating that these two genes are essential for light entrainment of the clock (Anwer et al. 2020).

The multiple *gi* mutants identified in Arabidopsis (Rédei, 1962; Koornneef et al. 1991) show different effects on the length of the circadian clock period, but all of them exhibit a late flowering phenotype (Fowler et al. 1999; Park et al. 1999). The effect of *GI* on Arabidopsis flowering time is stronger under LD than under SD, indicating an interaction between *GI* and the photoperiod pathway (Araki and Komeda, 1993; Mizoguchi et al. 2005). In addition to its role in the circadian clock regulation and photoperiodic flowering response, *GI* has been shown to play important roles in multiple other physiological processes such as light signaling, chlorophyll accumulation, starch accumulation and stress tolerance (reviewed in Mishra & Panigrahi, 2015; Brandoli et al. 2020). Because GI can physically interact with many proteins of the circadian clock and photoperiodic flowering pathways, GI has been suggested to function as a scaffold or hub protein and orchestrates other protein interactions (Imaizumi and Kay, 2006).

In Arabidopsis, GI regulates the expression of *FT* and flowering time through both CONSTANS

(CO)-dependent and CO-independent mechanisms. In the CO-dependent pathway, GI acts

between the circadian oscillator and the main photoperiod gene *CO* to accelerate flowering by

promoting both *CO* and *FT* mRNA abundance (Mizoguchi et al. 2005). GI interacts with the

FLAVIN-BINDING, KELCH REPEAR, F-BOX1 (FKF1) protein to form a complex that

regulates the protein abundance of CYCLING DOF FACTOR 1 (CDF1), a transcriptional

repressor of *CO*. The degradation of CDF1 protein elevates the transcription of *CO*, thereby

promoting *FT* expression and flowering (Sawa et al. 2007). In addition, GI can interact with CO

directly and indirectly through interactions with FKF1 and ZEITLUPE (ZTL) to regulate CO

protein stability (Song et al. 2014; Hwang et al. 2019). Finally, the interaction between GI and

SPINDLY (SPY), which is a negative regulator of gibberellin signaling, can inhibit SPY activity

and promote *FT* transcription in a CO-dependent manner (Tseng et al. 2004).

GI also regulates *FT* expression in Arabidopsis through two CO-independent pathways. The first

one involves microRNA172 (miR172), which inhibits the expression of flowering repressors

TARGET OF EAT 1 (*TOE1*) and *APETALA 2* (*AP2*) (Jung et al. 2007). In the second CO-

independent pathway, GI activates *FT* transcription through direct binding to the *FT* promoter

alone or in a protein complex with *FT* repressors such as SHORT VEGETATIVE PHASE (SVP),

TEMPRANILLO 1 (TEM1) and TEM2 (Sawa and Kay, 2011).

In Arabidopsis, *CO* plays a central role in the photoperiodic response by inducing flowering

under LD conditions (Valverde et al. 2004), and *CO* homologs from rice and sorghum have also

been shown to play major roles in the photoperiodic control of flowering (Yano et al. 2000;

Yang et al. 2014). However, in the temperate grasses, the duplicated *CO1* and *CO2* genes show

limited effects on the photoperiodic regulation of flowering time in the presence of

PHOTOPERIOD1 (*PPD1*). In the absence of functional *PPD1* alleles in wheat (Shaw et al.

2020), or in the presence of hypomorphic *PPD1* alleles in barley (Alqudah et al. 2014), the effect

of *CO1* on the acceleration of heading time under LD is stronger than in the presence of *PPD1*.

In the temperate grasses, *PPD1* is the dominant gene in the photoperiodic pathway, showing

strong differences in heading time between LD and SD even in the absence of functional copies

of both *CO1* and *CO2* (Shaw et al. 2020). Although the interactions between *GI* and *CO* have

been extensively studied in Arabidopsis (Mizoguchi et al. 2005; Sawa et al. 2007) and rice

(Hayama et al. 2003), the interactions between *GI* and *PPD1* remain largely unexplored.

Another understudied aspect of the role of *GI* in the temperate cereals is its interaction with

VRN2, a LD repressor of *FT1* with no known homologs in eudicots (Yan et al. 2004). *VRN2*

encodes a protein comprising a zinc finger motif at the N-terminus and a CCT domain at the C-

terminus (Yan et al. 2004), which is orthologous to the rice protein GRAIN NUMBER, PLANT

HEIGHT, AND HEADING DATE 7 (GHD7) (Xue et al. 2008; Woods et al. 2016). In rice, the

GI protein has been shown to interact *in vivo* with GHD7 and to regulate its stability (Zheng et al.

2019). Loss-of-function mutations in *GHD7* result in earlier flowering under LD in rice (Xue et

al. 2008), and mutations in *VRN2* result in accelerated heading under LD and a spring growth

habit in wheat (Yan et al. 2004; Distelfeld et al. 2009).

In this study, we show that *GI* loss-of-function mutants in wheat exhibit significant delays in

heading time under both LD and SD. This delay is larger under LD than under SD in photoperiod

sensitive wheats (PS), but the differences disappear in photoperiod insensitive (PI) wheats and

elf3 mutants. Moreover, we show that GI regulates *FT1* expression and heading time through its

interactions with *ELF3*, *VRN2*, *CO*, and the age-dependent miR172-AP2 pathway, revealing both

conserved and specific mechanisms for the temperate grasses.

Results

***GI* loss-of-function mutants delay heading time in wheat under LD photoperiod**

From a sequenced EMS mutant population in the tetraploid spring wheat variety Kronos

(Krasileva et al. 2017), we identified two truncation mutants for the *GI* A-genome homoeolog

(henceforth, *A-401* and *A-2019*) and two truncation mutants for the B-genome homeolog

(henceforth *B-2205* and *B-3825*, Fig. 1A). Line *A-401* has a splice site mutation that leads to the

retention of intron 4 and generates a premature stop codon that eliminates 1,034 amino acids

(89.5%) of the GI protein (total 1,155 amino acids). The other three mutant lines all have

premature stop codon mutations: *A-2019* (Q801*), *B-3825* (Q41*) and *B-2205* (W754*).

Kronos carries the photoperiod insensitive allele *Ppd-A1a*, which has a 1,027 bp deletion in the

promoter region and is associated with misexpression of *PPD1*, accelerated induction of *FT1*

expression, and early flowering under SD (Wilhelm et al. 2009). To study the effect of the *GI*

mutations in a photoperiod sensitive background, we crossed the four *gi* mutants twice with a

near-isogenic Kronos line carrying the photoperiod sensitive *Ppd-A1b* allele (henceforth Kronos-

PS, Pearce et al. 2017), and intercrossed them to generate three different BC_1F_2 lines

homozygous for *gi* mutants in both the A- and B- genome homoeologs. The *gi-1* line, generated

from the cross between *A-2019* and *B-2205*, headed 12 d later than Kronos-PS, whereas the *gi-2*

(*A-401* x *B-2205*) and *gi-3* (*A-401* x *B-3825*) mutants headed 16 d later (Fig. 1B). These results

indicate that *GI* functions as a promoter of heading time in wheat under LD.

To further explore the function of *GI*, we generated wheat transgenic lines constitutively

expressing a fusion of the full-length coding region of the *GI* gene and a C-terminal 4xMYC tag

under the control of the maize *UBIQUITIN* promoter in a Kronos-PI background (hereafter

referred to as *UBI-GI-MYC*). Among the six independent transgenic events, transgenic line #13

showed the highest transcript levels of *GI* relative to the non-transgenic sister line (Fig. S1A, $P <$

0.001) and was selected for further characterization under both LD and SD conditions.

Transgenic T_1 plants of the *UBI-GI-MYC* line #13 headed 3.3 d earlier than the non-transgenic

Kronos-PI controls when grown under LD (Fig. S1B), but showed no difference in heading time

relative to the controls under SD (Fig. S1C). These results confirmed that *GI* is a LD promoter of

wheat heading time.

We next crossed the *UBI-GI-MYC* #13 transgenic plant (PI) with the *gi-2* mutant (PS) to test the

ability of the transgene to rescue the delayed heading time of the *gi-2* mutant. In the F_2 progeny,

we selected sister lines homozygous for the wild-type control (WT), the non-transgenic *gi-2*

mutant (*gi-2*), and the combined *UBI-GI-MYC* / *gi-2* (all homozygous for the *PPD1-A1b* allele,

PS). The *UBI-GI-MYC* / *gi-2* plants headed 12.4 d earlier than the *gi-2* mutant, but still 2.4 d later

than the wild type under LD (Fig. 1C). These results indicate that the *UBI-GI-MYC* #13

transgene can largely, but not completely, rescue the delayed heading time of the *gi-2* mutant

(Fig. 1C).

In addition to the late heading under LD, the *gi-2* mutants also showed an increased number of

spikelets per spike (Fig.1D-E) and tillers (Fig. 1F-G). Compared with the wild-type control

plants, the *gi-2* mutant produced on average 5.8 more spikelets per spike (Fig.1E) and had an

average of 5.6 more spike-bearing tillers (Fig. 1G), likely associated with its extended vegetative

phase.

GI shows strong interaction with photoperiod in Kronos-PS, but not in Kronos-PI

When grown under SD (8 h light), Kronos-PI plants head between 80 and 100 days, whereas Kronos-PS plants fail to head (Shaw et al. 2020; Alvarez et al. 2023). Under these conditions Kronos-PI plants produce normal spikes and viable grains, whereas Kronos-PS plants undergo spike and stem elongation arrest, and spikes fail to emergence from the leaf sheaths (Pearce et al. 2017; Shaw et al. 2020). Similarly, the *gi-2* mutants failed to head under SD before the experiment was terminated at 150 d (Fig. 2A), but showed fully developed young spikes wrapped inside the leaf sheath (Fig. 2B).

Although it is not possible to determine heading time in Kronos-PS under SD, the final number of leaves can be used to estimate the timing of the transition from the vegetative shoot apical meristem (SAM) to the reproductive phase (flowering time). In two separate SD experiments, one performed under 8 h of light (SD) and the other under 10 h of light (SD-10h), Kronos-PS primary shoots produced 13.9 and 13.7 leaves in SD and SD-10h, respectively (including leaves still wrapped inside the sheaths). The *gi-2*-PS mutant produced ~1 more leaf than the Kronos-PS control under both conditions: SD (14.9 leaves) and SD-10h (14.5 leaves, Fig. 2C). However, when the same lines were grown under LD, the difference between wild type and *gi-2* increased to 3.4 leaves, indicating a highly significant interaction ($P < 0.001$) between the effects of *GI* and photoperiod on leaf number (Supplemental Information S2).

This interaction between *GI* and photoperiod was no longer significant when we tested the effect of the *gi-2* mutant on heading time in a Kronos-PI background (henceforth *gi-2*-PI). The *gi-2*-PI mutant showed similar delays in heading time relative to Kronos-PI under both LD (15.4 d) and SD conditions (16.5 d, Fig. 2D). A factorial ANOVA for heading time using these lines, showed highly significant effects for *GI* and photoperiod ($P < 0.001$), but no significant interaction between them ($P = 0.4347$, Supplemental Information S2).

To explore the interactions between *GI* and *PPD1* under LD, we crossed Kronos-PS *gi-2* with a loss-of-function *ppd1* mutant (Pearce et al. 2017) and compared heading times among the four possible homozygous classes. A factorial ANOVA showed highly significant effects of *PPD1* and *GI* on heading times ($P < 0.001$), but no significant interaction between the two genes (Fig. 2E, $P = 0.3807$, Supplemental Information S2). We did not study these lines under SD because *gi-2* and *ppd1* mutants, as well as Kronos-PS controls do not head under SD. Taken together,

these results suggest that the *PPD1* PS-allele is important for the *GI* differential flowering

responses in LD and SD.

Lastly, we compared *PPD1* expression profiles in Kronos-PS and the *gi-2*-PS mutant under LD

(Fig. 2F). In Kronos-PS, *PPD1* transcripts showed diurnal rhythms with a peak expression 12

hours after the lights were turned on (zeitgeber time 12 = ZT12). In the *gi-2*-PS mutant, *PPD1*

transcript levels were significantly downregulated relative to Kronos-PS from the ZT12

expression peak to ZT20 (Fig. 2F).

_GI_ and _ELF3_ show strong interactions on the regulation of _FT1_ expression and heading

time

Since loss-of-function mutations in *elf3* greatly reduce the photoperiodic response and

significantly alter *PPD1* expression (Alvarez et al. 2016; 2023), we evaluated the effect of *gi-2*

in the *elf3* mutant background. In the F_2 progeny of a cross between *gi-2* and *elf3* (both in

Kronos-PS), we selected homozygous *elf3* and *elf3 gi-2* mutants and evaluated them under LD

and SD. Both *GI* and photoperiod showed highly significant effects on heading time ($P < 0.001$),

however, no interaction was observed between them ($P = 0.7608$, Supplemental Information S3).

The delayed heading caused by the *gi-2* mutant in the *elf3* background was similar under LD

(14.7 d) and SD (14.9 d, Fig. 3A), indicating that in the absence of a functional *ELF3* the effect

of *GI* is no longer modulated by photoperiod.

We further explored the interaction between *GI* and *ELF3* in an additional LD experiment

including Kronos-PS, *gi-2*, *elf3*, and *elf3 gi-2* mutants in the same PS genetic background.

Consistent with previous findings (Alverez et al. 2016; 2023), the *elf3* allele was associated with

a highly significant acceleration in heading time and a reduced number of leaves relative to

Kronos-PS, but the differences were larger in the *gi-2* than in the wild-type background (Fig. 3B

and C). By contrast, the *gi-2* allele was associated with significant increases in leaf number and

delayed heading, with larger differences in the presence of the functional *Elf3* allele than in the

elf3 mutant background (Fig. 3B and C). These differential responses were reflected in highly

significant interactions between GI and ELF3 for both traits ($P < 0.001$, Supplemental

Information S3).

We next examined the diurnal expression pattern of *FT1* transcripts in the same four genotypes.

RNA samples were extracted from the 4th fully expanded leaves (L4, before the SAM transition

to the reproductive phase) collected at 4 h intervals for a 24 h period. As expected, the highest

transcript levels of *FT1* were observed in the *elf3* mutant (earliest heading) and the lowest in the

gi-2 mutant (latest heading, Fig. 3D). Factorial ANOVAs at each ZT point revealed highly

significant effects of *ELF3* and *GI* and highly significant interactions on the regulation of *FT1* (*P*

< 0.001, Supplemental Information S3). These results suggest that the differences in heading

time and leaf number in the *elf3* and *gi-2* mutants are mediated by their effects and interactions

on the regulation of *FT1* expression.

We next explored the effect of the different mutant genotypes on the expression of *GI* and core

circadian oscillator genes *CIRCADIAN CLOCK ASSOCIATED1* (*CCA1*, Fig. 4B) and *TIMING*

OF CAB EXPRESSION 1 (*TOC1*, Fig. 4C). *GI* transcript levels in the *elf3* mutant were

significantly lower than in Kronos-PS at the ZT12 peak, but significantly higher at ZT0, ZT4 and

ZT20 (Fig. 4A). The diurnal expression patterns of *CCA1* (Fig. 4B) and *TOC1* (Fig. 4C) were

altered in the *gi-2* and *elf3* individual mutants, with an even larger disruption observed in the *elf3*

gi-2 combined mutant. This was particularly evident in the *CCA1* expression profile, which

showed no diurnal rhythms in the combined mutant (Fig. 4B). In summary, these results indicate

that the combined *elf3 gi-2* mutant has a very disruptive effect on the circadian clock, a feature

that is also conserved in Arabidopsis (Anwer et al. 2020).

Finally, because ELF3 was previously shown to physically interact with GI and modulate the

cyclic accumulation of the GI protein in Arabidopsis (Yu et al. 2008), we explored the physical

interactions between the wheat GI and ELF3 proteins. Indeed, we observed a strong interaction

between the two wheat proteins in yeast two-hybrid (Y2H) assays (Fig. S2). The GI and ELF3

interactions at the protein level, together with the interactions at the transcriptional level, likely

contribute to the observed genetic interactions between *GI* and *ELF3* on heading time observed

in this study.

Interactions between *GI* and the LD flowering repressor *VRN2* contribute to the regulation

of wheat heading time

To determine the relationship between *GI* and *VRN2*, we crossed *gi-2* with a Kronos *vrn2* loss-

of-function mutant (Distelfeld et al. 2009) and studied the effect of the four different

homozygous genotypes on heading time. When grown under LD, the *vrn2* mutant headed 5.3 d

earlier than the wild type, and this effect increased to 12.4 d when the same comparison was

performed in the *gi-2* mutant background (Fig. 5A). On the contrary, the 21.6 d delay in heading

time associated with the *gi-2* mutant in plants homozygous for the functional *Vrn2* allele was

reduced to 14.5 d in plants homozygous for the *vrn2* mutant allele (Fig. 5A). These differential

effects were reflected in a highly significant interaction between *GI* and *VRN2* on heading time

($P < 0.001$, Supplemental Information S5).

When we analyzed the same plants for total leaf number, we also found highly significant

differences associated with *VRN2* and *GI* (Fig. 5B), but the interaction was not significant ($P =$

0.2665, Supplemental Information S5). These results suggest that the interaction between *GI* and

VRN2 has a limited effect on the initial transition of the SAM from the vegetative to the

reproductive phase, but a stronger effect on the subsequent phase of spike development and stem

elongation.

To characterize better the genetic interaction between *GI* and *VRN2* on heading time, we

determined the effect of the *gi* mutation on the *VRN2* transcription profile by qRT-PCR (Fig. 5C).

In both the Kronos-PS control and the *gi-2* mutant, *VRN2* transcripts increased during the day to

reach a peak at ZT12, with slightly lower levels in *gi-2* at ZT0 and ZT8 ($P< 0.05$). (Fig. 5C).

However, after the lights were turned off, *VRN2* transcript levels decreased rapidly in Kronos-PS

but remained high in the *gi-2* mutant (Fig. 5C), suggesting a role of *GI* in the downregulation of

VRN2 at night.

Since VRN1 is a known repressor of *VRN2* (Chen & Dubcovsky, 2012), we also analyzed the

transcript levels of *VRN1* using the same leaf samples. *VRN1* transcript levels were significantly

higher in the wild type than in the *gi-2* mutant in most time points, but at ZT20 the two

genotypes showed no significant differences (Fig. 5D). This result suggests that the role of *GI* in

the downregulation of *VRN2* during the night is not mediated by *VRN1*, but we cannot

completely rule out the possibility of a residual effect of the higher transcript levels of *VRN1* at

dusk (ZT16) in Kronos-PS relative to *gi-2*.

In addition to their interactions at the transcriptional level, GI and VRN2 proteins show strong

interaction in Y2H assays (Fig. S2). A similar interaction has been reported between the two

orthologous rice proteins by Y2H and further validated by BiFC assays in *Nicotiana*

benthamiana and by Co-IP experiments in rice protoplasts (Zheng et al. 2019). In summary, the

observed interactions between GI and VRN2 at the protein and transcriptional levels likely

contribute to their significant genetic interactions on heading time.

GI shows strong interactions with *CO1* and *CO2* on the regulation of heading time in wheat

The regulation of flowering time by *GI* in Arabidopsis is mainly mediated through the *CO-*

dependent pathway (Sawa et al. 2007), so we studied the interactions between *GI* and the two

CO paralogs present in wheat: *CO1* and *CO2* (Shaw et al. 2020). We first determined the effect

of the Kronos-PS *gi-2* mutant on the diurnal expression patterns of *CO1* and *CO2* in the 4th

leaves. The transcript levels of *CO1* were significantly up-regulated in the *gi-2* mutant relative to

the Kronos-PS control at ZT0 and ZT4, but were mostly indistinguishable between the two

genotypes at other time points (Fig. 6A). Transcript levels of *CO2* were significantly higher in

the *gi-2* mutant relative to Kronos-PS at three (ZT4, ZT8 and ZT12) of the six analyzed time

points (Fig. 6B).

Interestingly, GI also showed strong physical interactions with both CO1 and CO2 proteins in

Y2H assays (Fig. S2). These results suggest that wheat GI may also regulate CO protein stability

through protein interaction, as previously described in Arabidopsis GI (Song et al. 2014; Hwang

et al. 2019). Interactions at both the transcriptional and protein levels may contribute to the

genetic interaction between *GI* and *CO1/CO2* on heading time described in the next section.

From the cross between *gi-2* and *co1co2* double mutant in the Kronos-PS background we

generated the eight possible homozygous classes and studied their effect on heading time (all

genotypes were grown in the same LD growth chamber). Among the four genotypes carrying the

Gi wild-type allele, *co1* flowered 3.8 d earlier, *co2* 2.7 d later and the *co1co2* combined mutant

4.3 d earlier than Kronos-PS (Fig. 6C), similarly to what was reported in a previous study (Shaw

et al. 2020). Among the four genotypes carrying the *gi-2* mutation, the differences between the

individual mutants and the wild type (*co1* 4.5 d earlier and *co2* 5.7 d later) were in the same

direction as in the corresponding lines with the *Gi* allele (Fig. 6C). However, the *co1 co2*

combined mutant was 6.1 d later than the wild type in the *gi-2* background and 4.3 d earlier in

the *Gi* background (Fig. 6C). These opposite results were reflected in highly significant

interactions among these three genes in a 3-way ANOVA ($P < 0.001$, Supplemental Information

S6). The effect of *GI* on heading time was stronger in the presence of the *co1* allele than in the

presence of the *Co1* allele, whereas the effect of *CO1* was stronger in *Gi* than in *gi-2* (Fig. 6C).

The effects of both *co2* and *gi-2* on heading time were stronger in the presence of the mutant

allele of the other gene than in the presence of the wild-type allele (Fig. 6C).

Interactions between *GI* and the *miR172-AP2* age-dependent pathway contribute to the

regulation of *FT1* and wheat heading time

In Arabidopsis, one of the pathways that interacts with *GI* to regulate the expression of *FT* is the

age-dependent miR172-*AP2/TOE* pathway (Jung et al. 2007), which is conserved in wheat

(Debernardi et al. 2022). To test if a similar interaction exists in wheat, we first determined the

effect of *gi-2* (in PS background) on the expression levels of *miR172*, *AP2* and *FT1*. In both L7

and L9 leaves, the mature *miR172* levels were significantly lower in the *gi-2* mutant than in the

Kronos-PS control (Fig. 7A). As expected from the known role of *miR172* as a repressor of

AP2L1 (= *TOE1*, Debernardi et al. 2022), the transcript levels of *AP2L1* were higher in the *gi-2*

mutant than in the Kronos-PS control (Fig. 7B). Transcript levels of *FT1* were significantly

lower in the *gi-2* mutants relative to the wild type at both stages, which is consistent with the

known role of *AP2L1* as a repressor of *FT1* (Debernardi et al. 2022) and with the later heading

time of the *gi-2* mutant relative to the wild type (Fig. 7C).

To investigate the genetic interaction between miR172 and GI, we crossed the *gi-2* mutant with a

UBI-miRNA172 transgenic line (hereafter *UBI-miR172*) developed previously (Debernardi et al.

2022), and evaluated the four different homozygous genotypes for heading time in a growth

chamber under LD. Plants overexpressing *miR172* showed a small but significant acceleration in

heading time (Fig. 7D), consistent with a previous study (Debernardi et al. 2022). In the *gi*

mutant background, the acceleration of heading time by *UBI-miR172* (6.3 d) was greater than in

the wild type (2.3 d), and both were highly significant (Fig. 7D). On the contrary, the delayed

heading associated with *gi-2* was smaller in the *UBI-miR172* transgenic plants (15.6 d) than in

the non-transgenic sister lines (19.6 d, Fig. 7D). A factorial ANOVA for heading time revealed a

highly significant interaction ($P < 0.001$) between *GI* and *UBI-miR172* (Supplemental

Information S8). Taken together, these results confirmed that interactions between *GI* and the

miR172-*AP2L1* pathway contribute to the regulation of *FT1* expression and to the *GI* effects on

wheat heading time.

Integration of *GI* into a working model of the regulation of wheat heading time

To summarize the multiple pathways through which *GI* regulates wheat heading time, and its

interactions with other flowering genes, we present a working model in Fig. 8, and a summary of

the protein interactions in Supplemental Fig. S2B. Both figures indicate a highly interconnected

regulatory network.

This model shows that GI promotes *FT1* expression and heading time through multiple pathways,

including the grass-specific *VRN2* gene, as well as through complex interactions with *CO1/CO2*

and the miR172-AP2 age-dependent pathway. The model also shows that the interactions

between *GI* and photoperiod are dependent on the presence of a wild-type *PPD1*-PS allele. Some

of the interactions are shown by bidirectional grey arrows because they are too complex to be

summarized by a promotion arrow or repression T-bar. For example, *VRN2* expression is

upregulated in the *gi-2* mutant at ZT20 but downregulated at ZT0-ZT8 (Fig. 5); and *GI* is

downregulated in the *elf3* mutant at ZT12 but upregulated at ZT20-ZT4 (Fig. 4).

This model presents only the strongest effects, but additional weaker effects are known to exist.

For example, *PHYB* affects heading time and expression of multiple flowering genes even in the

absence of *ELF3*, whereas *ELF3* affects heading time even in the absence of *PPD1* (Alvarez et al.

2023).

Discussion

Functional conservation of *GI* homologs in the regulation of flowering time

The role of *GI* as a flowering promoter is conserved among LD plant species grown under

inductive LD photoperiods, while some functional diversification can be observed when those

plants are grown under non-inductive SD conditions. In Arabidopsis, *GI* activates *FT* expression

and promotes flowering time, and most *gi* mutants show delayed flowering phenotypes in both

LD and SD conditions, with smaller delays under SD (summarized in Mishra and Panigrahi,

2015). In pea, mutations in *LATE BLOOMER1* (*LATE1*, *GI* homolog in pea) prevent the

induction of *FT* and result in late flowering under LD (Hecht et al. 2007), confirming the

function of *LATE1* as a LD flowering promoter. In photoperiod sensitive wheat genotypes, the

promoting effect of *GI* on heading time is also stronger under LD than under SD (Fig. 2C).

GI homologs from SD grass species, such as rice and sorghum, promote flowering under SD

photoperiods, with *gi* mutants showing reduced florigen expression and late flowering under this

condition. However, under LD photoperiods rice *gi* mutant shows no changes in heading time

(Izawa et al. 2011), whereas sorghum *gi* mutant still shows delayed flowering (Abdul-Awal et al.

2020). Although the *GI* promotion of heading in SD grasses seems opposite to what we

described above for LD species, in both groups, *GI*'s stronger promotion of flowering occurs

under the inductive photoperiods (LD for LD-plants and SD for SD-plants). This result is not

surprising, since *GI* promotes flowering though multiple pathways that converge on the

upregulation of *FT*, which is expressed under inductive photoperiods. *GI*'s stronger

photoperiodic response under inductive photoperiods is conserved in LD species from wheat (Fig.

8) to Arabidopsis (Araki and Komeda, 1993; Fowler et al. 1999), and SD species from rice

(Izawa et al. 2011) to sorghum (Abdul-Awal et al. 2020), suggesting a conserved role of *GI* in

the photoperiodic response.

The interaction between *GI* and photoperiod is disrupted in Kronos PI and the *elf3* mutant

Grass species have a unique *PPD1*/*PRR37*-dependent photoperiodic flowering pathway in

addition to the *CO*-dependent pathway, but their relative importance varies across species (Yano

et al. 2000; Koo et al. 2013; Murphy et al. 2011; Yang et al. 2014; Pearce et al. 2017; Shaw et al.

2020). In wheat and barley, *PPD1* plays a more dominant role than *CO1* and *CO2* in the

photoperiodic regulation of heading time (Alqudah et al. 2014; Shaw et al. 2020). Most of the

natural variation in the photoperiodic response in wheat is associated with either deletions in the

promoters of *PPD-A1* or *PPD-D1* or with copy number variation of *PPD-B1* (Wilhelm et al.

2009; Beales et al. 2007; Díaz et al. 2012; Würschum et al. 2019). The deletion in the promoter

of *PPD1* results in misexpression of *PPD1* during the night and dawn, and the induction of *FT1*

and earlier heading under SD (Alvarez et al. 2023; Wilhelm et al. 2009).

The misexpression of *PPD1* in Kronos-PI also resulted in the elimination of the significant

interaction on leaf number observed in Kronos-PS between *GI* and photoperiod (Fig. 2C). These

results indicate that the photoperiod sensitive *PPD1* allele is critical for conveying the

photoperiodic information to *GI* and for generating its differential flowering response in LD and

SD (Fig. 2C).

This hypothesis is further supported by the absence of a significant interaction between *GI* and

photoperiod on heading time in the *elf3* mutant, where the expression profiles of both *GI* (Fig.

4A) and *PPD1* (Alvarez et al. 2023) are significantly disrupted. In the absence of *ELF3* function,

GI can still promote flowering time, although its effect is weaker in *elf3* relative to the *ELF3*

wild-type allele (Fig. 3B-C). Moreover, in the absence of *ELF3*, the effect of *GI* on flowering is

similar in LD and SD, as *elf3 gi* mutant show similar delays in heading time relative to the *elf3*

mutant under both photoperiods (Fig. 3A). These results indicate that the differential effect of *GI*

on heading time in different photoperiods is also dependent on a functional *ELF3*.

The altered *PPD1* expression in *elf3* may contribute to the lost interaction between *GI* and

photoperiod, but additional post-translational mechanisms cannot be ruled out. In this study, we

show that GI and ELF3 physically interact with each other by Y2H assay (Fig. S2), an

interaction that has been also observed in Arabidopsis (Yu et al. 2008). In Arabidopsis, GI

protein stability and photoperiodic flowering is also regulated by the interaction between ELF3

and COP1 (Yu et al. 2008). ELF3 is essential for the interaction between GI and COP1, and the

Arabidopsis *elf3* mutant exhibits a disruption in the cyclic accumulation of GI associated with

early flowering and photoperiod-insensitivity (Zagotta et al. 1996; Yu et al. 2008). Similar

protein interactions may also contribute to the disrupted photoperiodic response of *GI* in the *elf3*

wheat mutant background.

***GI* interacts with the grass-specific *VRN2* repressor to regulate wheat heading time**

The effects of *VRN2* on heading time reported in this study are smaller than those reported in

previous studies in winter wheat (Yan et al. 2004; Distelfeld et al. 2009). In winter cereals, *VRN2*

is a LD flowering repressor that prevents *FT1* expression during the fall, and is repressed in the

spring by *VRN1*, which is upregulated by vernalization (Chen & Dubcovsky, 2012). However, in

spring wheats, such as the Kronos variety used in this study, *VRN1* induction does not require

vernalization and the earlier down-regulation of *VRN2* results in smaller effects of *VRN2* on

heading time in spring than in winter wheat varieties.

In this study, we show that the effects of *VRN2* on heading time are modulated by its interactions

with *GI* (Fig. 5A-B) both at the transcriptional and protein levels. In the *gi-2* mutant, the

expression levels of *VRN2* vary during the day, with significant increases at ZT20 and significant

decreases between ZT0 and ZT8 (Fig. 5C). These diurnal changes likely reflect *GI*'s own

circadian oscillation (Fig. 4A) and/or the effects of the *gi-2* mutants on the central oscillators

(Fig. 4B-C). GI and VRN2 also interact at the protein level (Fig. S2), a result that has been

validated for the rice orthologs (Zheng et al. 2019). A recent study in rice has shown that co-

expression of GI and GHD7 causes reduced accumulation of the GHD7 protein, and that GI and

phytochromes function antagonistically in the regulation of GHD7 protein stability (Zheng et al.

2019).

The existence of a similar negative effect of GI on the stability of the VRN2 protein in wheat is

indirectly supported by the stronger effect of *VRN2* on heading time in the *gi-2* mutant than in

the presence of the functional *Gi* allele (Fig. 5A-B). This significant genetic interaction between

GI and *VRN2* on heading time also resulted in larger differences between the *gi-2* mutant and the

wild type in the presence of the functional *Vrn2* allele than in the *vrn2* mutant. This result

indicates that part of the effect of *GI* in the acceleration of heading time in wheat is mediated by

its interactions with *VRN2*.

In summary, these results show that in the grasses, GI interacts with *VRN2*, a gene that has no

homologs in the eudicots plants. It will be interesting to determine if these genetic interactions

are the result of direct interactions between these two genes or an indirect effect of the multiple

interactions between *GI* and other flowering genes.

The conserved GI-CO pathway contributes to the regulation of wheat heading time

The mechanistic characterization of *GI*'s role in photoperiodic flowering regulation is mostly

from Arabidopsis, where GI promotes *CO* transcription and acts as a major mediator between the

circadian clock and CO. Under LD, the peak of GI protein accumulation coincides with FKF1 in

the middle of the day, resulting in the repression of CDF1 and, consequently, in increased

transcript levels of *CO* and *FT* (Sawa et al. 2007). Under SD, GI accumulation peaks three hours

earlier than that of FKF1, precluding the repression of CDF1, and resulting in reduced levels of

CO transcripts (Sawa et al. 2007). Therefore, the formation of the GI-FKF1 complex is required

for day-length measurement in Arabidopsis. In addition, GI can regulate CO temporal stability

through direct physical interactions and through its interactions with FKF1 and ZTL (Song et al.

2014; Hwang et al. 2019).

In this study, we show a direct physical interaction between GI and both CO1 and CO2 wheat

proteins in Y2H assays (Fig. S2). A similar interaction between GI and CO has been reported in

Arabidopsis (Song et al. 2014), suggesting similar mechanisms between the two species. This

hypothesis is also supported by the physical interactions observed between GI and ZTL, COP1

and FKF1 in *Brachypodium* (Hong et al. 2010) and GI-FKF1 interaction in sorghum (Abdul-

Awal et al. 2020). The physical interactions between GI and both CO1 and CO2 in wheat are

reflected in a strong genetic interaction, where the differences in heading time between Kronos-

PS and the combined *co1 co2* mutant are reverted in the *gi-2* mutant relative to the wild type

(Fig.6C).

We detected differences between wheat and Arabidopsis in the transcriptional regulation of *CO*

by GI. While transcript levels of *CO* are significantly reduced in the Arabidopsis *gi* mutant

relative to the wild type (Sawa et al. 2007; Sawa and Kay, 2011), *CO1* and *CO2* are significantly

upregulated at different times of the day in the wheat *gi-2* mutant (Fig. 6A-B). Therefore,

although the presence of the GI-CO pathway seems to be conserved in Arabidopsis and wheat,

the mechanisms by which they control photoperiodic flowering likely differ between these

two species.

These differences are also reflected in the more limited role of *CO1* and *CO2* in the regulation of

the photoperiodic response in the temperate grasses, where *PPD1* is the main photoperiod gene

(Shaw et al. 2020). Although both the *PPD1*-dependent pathway and the *CO*-dependent pathway

can perceive the differences in photoperiod in the absence of the other gene, there are still

significant genetic interactions among them that are important for the fine-tuning of the

flowering response in wheat (Shaw et al. 2020).

GI can regulate wheat heading time by CO-independent regulatory pathways

In addition to the CO-mediated pathway, we show that in wheat GI can also regulate *FT1*

expression through the age-dependent miR172-AP2 module. The interaction between GI and

miR172-AP2 seems to be conserved between Arabidopsis and wheat, with GI promoting

miR172 and repressing AP2 expression in both species (Jung et al. 2007; Fig. 7).

In Arabidopsis, GI has been shown to promote *FT* transcription through direct binding to its

promoter regions (Sawa & Kay, 2011). However, as SVP, TEM1 and TEM2 also bind to the

same *FT* promoter regions, it is also possible that the binding of GI to the *FT* promoter is

mediated by a complex with other proteins (Sawa & Kay, 2011). In wheat, we also observed a

strong downregulation of *FT1* in the *gi-2* mutant (Fig. 7C), which was observed even in the

presence of the *elf3* mutation (Fig. 3D). Although these results may reflect a direct interaction

between GI and the *FT1* promoter, we cannot rule out indirect effects mediated by the multiple

pathways through which GI regulates *FT1*.

Materials and methods

Plant materials

The reference sequences of *GI* (TraesCS3A02G116300 and TraesCS3B02G135400) were used

to screen for mutations in the sequenced ethyl methanesulfonate mutagenized Kronos population

(Krasileva et al. 2017) using BLASTN. A splicing site mutation (*A-401*) and three stop codon

mutations (*A-2019*, *B-2205* and *B-3825*) were selected to generate *GI* loss-of-function mutants.

To test genetic interactions between *GI* and *ELF3*, *CO1*, *CO2*, *PPD1* and *VRN2*, we combined

gi-2 with the respective loss-of-function mutants of these genes and generated *elf3 gi*, *co1 gi*, *co2*

gi, *co1 co2 gi*, *ppd1 gi* and *vrn2 gi* in the same Kronos-PS background. Primers used to genotype

and verify the presence of individual gene mutations are listed in Table S1. We also crossed the

gi-2 mutant with a *Ubi-miR172* transgenic line (Debernardi et al. 2022) to investigate the

interaction between miR172 and *GI* in the Kronos-PS background. Primers used to differentiate

the *Ppd-A1b* allele (PS) from *Ppd-A1a* allele (PI) are also included in Table S1.

Growth conditions

All experiments were performed in controlled environment conditions using the Conviron

PGR15 growth chambers. During the lights-on period, the growth chambers were set at 22°C,

but the first and last hour of this lights-on period were set at 20°C to provide a more gradual

change between temperatures. Night temperatures were set at 17°C. All PGR15 chambers used

similar metal halide and high-pressure sodium light configurations, and lights were set to the

same intensity in all experiments (~330µM m^{-2} s^{-1}). Lights were on for 8 hours for SD-8h, 10

hours for SD-10h experiments and 16 hours for LD experiments.

Generation of the transgenic line overexpressing *GI*

Transgenic Kronos plants overexpressing *GI* were generated at the UC Davis Plant

Transformation Facility (http://ucdptf.ucdavis.edu/) using the Japan Tobacco vector pLC41

(hygromycin resistance) and transformation technology licensed to UC Davis. The coding region

of *GI-A* gene from Kronos was cloned downstream of the maize UBIQUITIN promoter with a C-

terminal 4xMYC tag. Agrobacterium strain EHA105 was used to infect Kronos immature

embryos and all transgenic plants were genotyped by PCR using primers described in

Supplemental Table S2.

qRT-PCR analysis

For the expression analyses across developmental stages, we collected the newest fully expanded

leaf samples (L7, L9 and L11) 4 hours after lights were turned on (at ZT4). For the time-course

experiments analyzing the diurnal gene expression patterns, the 4th fully expanded leaves were

collected at 4h intervals for 24 hours. RNA samples were extracted using the Spectrum Plant

Total RNA Kit (Sigma-Aldrich). Total RNA was treated with RQ1 RNase-free DNase (Promega)

prior to cDNA synthesis. And 1µg of total RNA was used to synthesize cDNA using the High-

Capacity cDNA Reverse Transcription Kit (Applied Biosystems) according to the

manufacturer's instructions. The cDNA was then diluted 20 times and 5 µl of the dilution was

mixed with 2×VeriQuest Fast SYBR Green qPCR Master Mix (Affymetrix, 75690) and primers

for the real-time qPCR analysis. Primer sequences are listed in Table S2. *ACTIN* was used as

endogenous control except for miR172, the small nucleolar RNA 101 (SnoR) was used as

internal reference. Primers used in qRT-PCR analyses are included in Table S2.

Acknowledgments and Funding

This project was supported by the Howard Hughes Medical Institute (https://www.hhmi.org/) and by competitive Grants 2016-67013-24617 and 2022-68013-36439 from the United States Department of Agriculture, National Institute of Food and Agriculture (https://nifa.usda.gov/).

Author contributions

CL and HL designed and performed most of the experiments and analyses. CL wrote the first version of the manuscript. JMD contributed the *UBI-miR172* transgenic material and determined the effect of GI on the transcript levels of *miR172* and *AP2*. CZ contributed to the diurnal gene expression analyses. All authors reviewed the manuscript, contributed ideas and provided valuable suggestions. CL and JD were responsible for the conceptualization of the project. JD was responsible for obtaining funding, project coordination and supervision, statistical analyses and for the final revision of the manuscript.

Conflict of interest

The authors declare no conflict of interest

Data and materials availability

All Kronos mutants and transgenic lines are available from the authors upon requests without any restrictions for use. Kronos single mutants are also available from the Germplasm Resources Unit (GRU) at the John Innes Centre.

Supplemental Materials

Fig. S1. Expression analysis of the transgenic *UBI-GI-MYC* lines (in Kronos-PI background) and evaluation of the effect of the transgene on heading time under long- (LD) and short-day (SD).

Fig. S2. GI protein interacts with ELF3, CO1, CO2 and VRN2, but not with PPD1 in Y2H assays.

Table S1. Primers used for mutant genotyping.

Table S2. Primers used in the qRT-PCR analysis.

Supplemental Information S1-S8: raw data and statistics supporting figures.

References

Abdul-Awal SM, Chen J, Xin Z, Harmon FG (2020) A sorghum *gigantea* mutant attenuates
florigen gene expression and delays flowering time. Plant Direct. **4**(11): e00281.

Alqudah AM, Sharma R, Pasam RK, Graner A, Kilian B, Schnurbusch T (2014) Genetic
dissection of photoperiod response based on GWAS of pre-anthesis phase duration in
spring barley. PLoS ONE. **9**(11): e113120.

Alvarez MA, Li C, Lin H, Joe A, Padilla M, Woods DP, Dubcovsky J (2023) *EARLY*
FLOWERING 3 interactions with *PHYTOCHROME B* and P*HOTOPERIOD1* are critical
for the photoperiodic regulation of wheat heading time. PLoS Genet. **19**: e1010655.

Alvarez MA, Tranquilli G, Lewis S, Kippes N, Dubcovsky J (2016) Genetic and physical
mapping of the earliness *per se* locus *Eps-A^m1* in *Triticum monococcum* identifies *EARLY*
FLOWERING 3 (ELF3) as a candidate gene. Funct Integr Genomics. **16**: 365-382.

Anwer MU, Davis A, Davis SJ, Quint M (2020) Photoperiod sensing of the circadian clock is
controlled by *EARLY FLOWERING 3* and *GIGANTEA*. Plant J. **101**: 1397-1410.

Araki T, Komeda Y (1993) Analysis of the role of the late-flowering locus, *Gl*, in the flowering
of *Arabidopsis thaliana*. Plant Journal **3**: 231-239.

Beales J, Turner A, Griffiths S, Snape JW, Laurie DA (2007) A *pseudo-response regulator* is
misexpressed in the photoperiod insensitive *Ppd-D1a* mutant of wheat (*Triticum aestivum*
L.). Theor Appl Genet. **115**: 721-733.

Brandoli C, Petri C, Egae-Cortines M and Weiss J (2020) Gigantea: uncovering new
functions in flower development. Genes. **11**: 1142.

Chen A, Dubcovsky J. (2012) Wheat TILLING mutants show that the vernalization gene VRN1
down-regulates the flowering repressor *VRN2* in leaves but is not essential for flowering.
PLoS Genet. **8**(12): e1003134.

David KM, Armbruster U, Tama N, Putterill J (2006) Arabidopsis GIGANTEA protein is
post-transcriptionally regulated by light and dark. FEBS Lett. **580**: 1193-1197.

Debernardi JM, Woods DP, Li K, Li C, Dubcovsky J (2022) MiR172-*APETALA2-like* genes
integrate vernalization and plant age to control flowering time in wheat. PLoS Genet
18(4): e1010157

Díaz A, Zikhali M, Turner AS, Isaac P, Laurie DA (2012) Copy number variation affecting
the *Photoperiod-B1* and *Vernalization-A1* genes is associated with altered flowering time
in wheat (*Triticum aestivum*). PLoS One. **7**(3): e33234.

Distelfeld A, Tranquilli G, Li C, Yan L and Dubcovsky J (2009) Genetic and molecular
characterization of the *VRN2* loci in tetraploid wheat. Plant Physiol. **149**: 245-257.

Ezer D, Jung JH, Lan H, Biswas S, Gregoire L, Box MS, Charoensawan V, Cortijo S, Lai X,
Stöckle D, Zubieta C, Jaeger KE and Wigge PA (2017) The evening complex
coordinates environmental and endogenous signals in Arabidopsis. Nat. Plants. **3**: 17087.

Fowler S, Lee K, Onouchi H, Samach A, Richardson K, Morris B, Coupland G, Putterill J
(1999) *GIGANTEA*: A circadian clock-controlled gene that regulates photoperiodic
flowering in Arabidopsis and encodes a protein with several possible membrane-spanning
domains. EMBO J. **18**: 4679–4688.

Hayama R, Yokoi S, Tamaki S, Yano M, Shimamoto K (2003) Adaptation of photoperiodic
control pathways produces short-day flowering in rice. Nature. **422**: 719-722.

Hecht V, Knowles CL, Vander Schoor JK, Liew LC, Jones SE, Lambert MJ, Weller JL
(2007) Pea *LATE BLOOMER1* is a *GIGANTEA* ortholog with roles in photoperiodic
flowering, deetiolation, and transcriptional regulation of circadian clock gene homologs.
Plant Physiol. **144**: 648-661.

Hong SY, Lee S, Seo PJ, Yang MS, Park CM (2010) Identification and molecular
characterization of a *Brachypodium distachyon GIGANTEA* gene: functional
conservation in monocot and dicot plants. Plant Mol Biol. **72**: 485-497.

Huq E, Tepperman JM, Quail PH (2000) GIGANTEA is a nuclear protein involved in
phytochrome signaling in Arabidopsis. Proc Natl Acad Sci USA. **97**: 9789-9794.

Hwang DY, Park S, Lee S, Lee SS, Imaizumi T, Song YH (2019) GIGANTEA regulates the
timing stabilization of CONSTANS by altering the interaction between FKF1 and
ZEITLUPE. Mol Cells. **42**: 693-701.

Imaizumi T and Kay SA (2006) Photoperiodic control of flowering: not only by coincidence.
Trends Plant Sci. **11**: 550-558.

Izawa T, Mihara M, Suzuki Y, Gupta M, Itoh H, Nagano AJ, Motoyama R, Sawada Y,
Yano M, Hirai MY, Makino A, Nagamura Y (2011) Os-*GIGANTEA* confers robust
diurnal rhythms on the global transcriptome of rice in the field. Plant Cell. **23**:1741-1755.

Jung JH, Seo YH, Seo PJ, Reyes JL, Yun J, Chua NH, Park CM (2007) The *GIGANTEA*-
regulated microRNA172 mediates photoperiodic flowering independent of *CONSTANS*
in Arabidopsis. Plant Cell. **19**: 2736–2748.

Koo BH, Yoo SC, Park JW, Kwon CT, Lee BD, An G, Zhang Z, Li J, Li Z, Paek NC (2013)
Natural variation in *OsPRR37* regulates heading date and contributes to rice cultivation at
a wide range of latitudes. Mol Plant. **6**: 1877-1888.

Koornneef M, Hanhart CJ, van der Veen JH (1991) A genetic and physiological analysis of
late flowering mutants in *Arabidopsis thaliana*. Mol Gen Genet. **229**: 57-66.

Krasileva KV, Vasquez-Gross HA, Howell T, Bailey P, Paraiso F, Clissold L, Simmonds J,
Ramirez-Gonzalez RH, Wang X, Borrill P, Fosker C, Ayling S, Phillips AL, Uauy C,
Dubcovsky J (2017) Uncovering hidden variation in polyploid wheat. Proc Natl Acad
Sci USA. **114**: E913-E921.

Lu SX, Webb CJ, Knowles SM, Kim SH, Wang Z, Tobin EM (2012) CCA1 and ELF3
interact in the control of   hypocotyl length and flowering time in Arabidopsis. Plant Physiol. **158**:
1079-1088.

Mishra P and Panigrahi KC (2015) GIGANTEA – an emerging story. Front. Plant Sci. **6**: 8.

Murphy RL, Klein RR, Morishige DT, Brady JA, Rooney WL, Miller FR, Dugas DV, Klein
PE, Mullet JE (2011) Coincident light and clock regulation of *pseudoresponse regulator*
protein 37 (*PRR37*) controls photoperiodic flowering in sorghum. Proc Natl Acad Sci
USA. **108**: 16469-16474.

Park DH, Somers DE, Kim YS, Choy YH, Lim HK, Soh MS, Kim HJ, Kay SA, Nam HG
(1999) Control of circadian rhythms and photoperiodic flowering by the *Arabidopsis*
GIGANTEA gene. Science. **285**: 1579-1582.

Pearce S, Shaw LM, Lin H, Cotter JD, Li C, Dubcovsky J (2017) Night-break experiments
shed light on the Photoperiod1-mediated flowering. Plant Physiol. **174**: 1139-1150.

Rédei GP (1962) Supervital mutants of Arabidopsis. Genetics. **47**: 443-460.

Sawa M, Kay SA (2011) GIGANTEA directly activates *Flowering Locus T* in *Arabidopsis*
thaliana. Proc. Natl. Acad. Sci. **108**: 11698–11703.

Sawa M, Nusinow DA, Kay SA and Imaizumi T (2007) FKF1 and GIGANTEA complex
formation is required for day-length measurement in *Arabidopsis*. Science. **318**: 261-265.

Shaw LM, Li C, Woods DP, Alvarez MA, Lin H, Lau MY, Chen A, Dubcovsky J (2020)
Epistatic interactions between *PHOTOPERIOD1*, *CONSTANS1* and *CONSTANS2*
modulate the photoperiodic response in wheat. PLoS Genet. **16**(7): e1008812.

Song YH, Estrada DA, Johnson RS, Kim SK, Lee SY, MacCoss MJ, Imaizumi T (2014)
Distinct roles of FKF1, Gigantea, and Zeitlupe proteins in the regulation of Constans
stability in *Arabidopsis* photoperiodic flowering. Proc Natl Acad Sci USA. **111**: 17672-
17677.

Tseng TS, Salomé PA, McClung CR, Olszewski NE (2004) SPINDLY and GIGANTEA
interact and act in *Arabidopsis thaliana* pathways involved in light responses, flowering,
and rhythms in cotyledon movements. Plant Cell. **16**: 1550-1563.

Valverde F, Mouradov A, Soppe W, Ravenscroft D, Samach A, Coupland G (2004)
Photoreceptor regulation of CONSTANS protein in photoperiodic flowering. Science.
303: 1003-1006.

Wilhelm EP, Turner AS, Laurie DA (2009) Photoperiod insensitive *Ppd-A1a* mutations in
tetraploid wheat (*Triticum durum* Desf.). Theor Appl Genet. **118**: 285-294.

Woods DP, McKeown MA, Dong Y, Preston JC, Amasino RM (2016) Evolution of
VRN2/Ghd7-like genes in vernalization-mediated repression of grass flowering. Plant
Physiol. **170**: 2124-2135.

Würschum T, Rapp M, Miedaner T, Longin CFH, Leiser WL (2019) Copy number variation
of *Ppd-B1* is the major determinant of heading time in durum wheat. BMC Genet. **20**: 64.

Xue W, Xing Y, Weng X, Zhao Y, Tang W, Wang L, Zhou H, Yu S, Xu C, Li X, Zhang Q
(2008) Natural variation in *Ghd7* is an important regulator of heading date and yield
potential in rice. Nat Genet. **40**: 761-767.

Yan L, Loukoianov A, Blechl A, Tranquilli G, Ramakrishna W, SanMiguel P, Bennetzen
JL, Echenique V, Dubcovsky J (2004) The wheat *VRN2* gene is a flowering repressor
down-regulated by vernalization. Science. **303**: 1640-1644.

Yang S, Weers BD, Morishige DT, Mullet JE (2014) *CONSTANS* is a photoperiod regulated
activator of flowering in sorghum. BMC Plant Biol. **14**: 148.

Yano M, Katayose Y, Ashikari M, Yamanouchi U, Monna L, Fuse T, Baba T, Yamamoto
K, Umehara Y, Nagamura Y, Sasaki T (2000) *Hd1*, a major photoperiod sensitivity
quantitative trait locus in rice, is closely related to the Arabidopsis flowering time gene
CONSTANS. Plant Cell. **12**: 2473-2484.

Yu JW, Rubio V, Lee NY, Bai S, Lee SY, Kim SS, Liu L, Zhang Y, Irigoyen ML, Sullivan
JA, Zhang Y, Lee I, Xie Q, Paek NC, Deng XW (2008) COP1 and ELF3 control
circadian function and photoperiodic flowering by regulating GI stability. Mol Cell. **32**:
617–630.

Zagotta MT, Hicks KA, Jacobs CI, Young JC, Hangarter RP, Meeks-Wagner DR (1999).
The *Arabidopsis ELF3* gene regulates vegetative photomorphogenesis and the
photoperiodic induction of flowering. Plant J. **10**: 691-702.

Zheng T, Sun J, Zhou S, Chen S, Lu J, Cui S, Tian Y, Zhang H, Cai M, Zhu S, Wu M,
Wang Y, Jiang L, Zhai H, Wang H, Wan J (2019) Post-transcriptional regulation of
Ghd7 protein stability by phytochrome and OsGI in photoperiodic control of flowering in
rice. New Phytol. **224**: 306-320.

 Figure legends

Fig. 1. Effect of *gi* loss-of-function mutants and transgenic lines expressing *GI* under the maize

ubiquitin promoter (*UBI-GI-MYC*) on heading time under long-day (LD). A, Schematic diagram

showing *GI* gene structure and the positions of selected mutations. Exons are represented by

rectangles and introns by a line. Mutant line *A-2019* was combined with *B-2205* to generate *gi-1*,

A-401 combined with *B-2205* to generate *gi-2*, and *A-401* combined with *B-3825* to generate *gi-3*

(letters indicate the genomes, and numbers indicate the Kronos mutant identification). B, *gi*

mutants in Kronos-PS show a significant delay in heading time relative to their control sister

lines under LD. C, The *UBI-GI-MYC* transgene largely rescues the delayed heading time of *gi-2*

mutant in Kronos-PS background under LD. Different letters above the bars indicate significant

differences in Tukey test = $P < 0.01$. D-E, Kronos-PS spikes of *gi-2* mutants produce more

spikelets per spike (SNS) than those of the wild type. The scale bar in D = 1 cm. F, Kronos-PS

12-week-old *gi-2* mutant compared to wild-type sister lines. G, The *gi-2* mutant plants produce

more spike-bearing tillers than the control sister plants. Numbers inside bars indicate the number

of plants included in each analysis. Error bars are standard errors of the means. *** = $P < 0.001$

for differences with the wild-type control using two-tailed *t*-test in B, E and G. Raw data and

statistical analyses are available in Supplemental Data S1.

Fig. 2. Interactions between *GI* and photoperiod in Kronos-PS, Kronos-PI and *ppd1* mutants. A,

Heading time evaluation of Kronos-PS and *gi-2* mutant grown under SD (8h light). Arrows on

top of bars indicate that both genotypes failed to head before the experiment was terminated at

150 d. B, Pictures of developing young spikes of Kronos-PS and *gi-2* mutant dissected 100 d

after germination from the same batch of plants grown under SD as in A. Scale bar = 1000 μm. C,

Effect of *gi-2* on leaf number under SD-8h, SD-10h and LD-16h in a Kronos-PS background.

Numbers inside the bars in A and C indicate the number of plants included in the analysis for

each genotype. D, Interaction graph showing the effect of *gi-2* on days to heading under SD-8h

and LD-16h in a Kronos-PI background (n = 10 - 11). E, Interaction graph showing the effect of

gi-2 and *ppd1* mutants on heading time under LD. Error bars are standard errors of the means

based on n = 7 - 8. F, Effect of *gi* mutations on *PPD1* transcription profiles under LD. The gray

shaded area indicates the 8 h of darkness. Primers used in the qRT-PCR amplify both *PPD1*

homeologs. The 4[th] fully expanded leaves were collected at 4 h intervals for 24h. Expression

levels are expressed relative to *ACTIN* as endogenous control using the ΔCt method. Error bars

are standard errors of the means based on n = 5. ns = not significant, ** = $P < 0.01$, *** = $P <$

0.001 based on two-tail *t*-tests for each time point. Raw data and statistical analyses are available

in Supplemental Data S2.

Fig. 3. Interactions between *GI* and *ELF3* on the regulation of flowering/heading time and *FT1*

expression. A, Interaction graph showing the effect of the *gi-2* mutation on days to heading

under different photoperiods in a Kronos-PS-*elf3* mutant background *(elf3* vs. *elf3 gi-2).* SD = 8

h light and LD = 16 h light (n = 10-12). B and C, Interaction graphs for the effects of *GI* and

ELF3 on heading time (B) and total number of leaves produced by the primary shoots (C) under

LD. Error bars are standard errors of the means based on n =8-10. *** = $P < 0.001$ for the four

simple effect ANOVAs. The 2 x2 factorial ANOVAs are in Supplemental Data S3. D, Time-

course expression analysis of *FT1* in the 4th fully expanded leaves of Kronos-PS (WT), *elf3*, *gi-2*

and *elf3 gi-2* mutants under LD. The gray shaded area indicates the 8 h of darkness. The inset

shows *FT1* expression from the same experiment but excluding the *elf3* mutant to visualize better

the smaller differences among the other genotypes. Expression levels were calculated relative to

the endogenous control *ACTIN* using the ΔCt method (n = 5). Raw data and statistical analyses

are available in Supplemental Data S3.

Fig. 4 Effect of the *elf3* and *gi-2* mutations on the transcription profile of *GI* and the core

circadian oscillator genes. Samples were collected at 4h intervals for a 24h period from the 4th

fully expanded leaves. Expression levels of the target genes relative to the *ACTIN* endogenous

control were evaluated by qRT-PCR using the ΔCt method. A, Effect of the *elf3* mutation on *GI*

transcript levels. *** = $P < 0.001$ based on two-tailed t-test. The shaded gray area indicates the 8

h of darkness. Error bars are s.e.m based on 5 biological replications. B and C, Diurnal

expression patterns of the core circadian clock genes *CCA1* (B) and *TOC1* (C) in Kronos-PS

(WT), *elf3*, *gi-2,* and combined mutant *elf3 gi-2*. Five biological replicates were used per

genotype. Raw data and factorial ANOVAs at each time point are provided in Supplemental

Data S4.

Fig. 5. Interactions between *GI* and *VRN2*. A-B, Interaction graphs showing the effect of *GI* on

heading time (A) and leaf number (B) in the presence and absence of functional *VRN2* alleles (n

= 10-12). C-D, Effect of *gi-2* mutation on the transcript levels of *VRN2* (C) and *VRN1* (D) under

LD. Each data point represents the average of 5 biological replicates. Samples were collected at

4h intervals for a 24h period from the 4[th] fully expanded leaves and transcript levels were

evaluated by qRT-PCR and calculated relative to the *ACTIN* endogenous control using the ΔCt

method. Error bars are standard errors of the means. ns = not significant, * = $P < 0.05$, ** = $P <$

0.01, and *** = $P < 0.001$ were obtained from simple effect ANOVAS in A and B (complete

ANOVA in Supplemental Data S5) and two-tailed t-tests in C and D at each time point. Raw

data is available in Supplemental Data S5.

Fig. 6. Interactions among *GI, CO1* and *CO2* on the regulation of wheat heading time under LD

photoperiod. A-B, Effect of *gi-2* mutation on the expression profiles of *CO1* (A) and *CO2* (B)

under LD (16h light). The gray shadowed area indicates the hours of darkness. The 4[th] fully

expanded leaves were collected at 4 h intervals for a 24 h period. Expression levels were

calculated relative to the endogenous control *ACTIN* using the ΔCt method. Values are averages

of 5 biological replicates and error bars are standard errors of the means. ns = not significant, * =

$P < 0.05$, ** = $P < 0.01$, *** = $P < 0.001$ for differences between genotypes at each time point

using t-tests. C, Heading time of wild-type Kronos-PS (WT), *co1, co2, co1 co2, gi, co1 gi, co2 gi*

and *co1 co2 gi* homozygous mutants grown in the same growth chamber under LD. The number

of plants included in the experiment are indicated inside the bars. Error bars are standard errors

of the means. Different letters with the same color indicate significant differences in Tukey tests

($P < 0.05$) in separate ANOVAs for *Gi* (blue) and *gi-2* (orange) genotypes. Raw data and

statistical analysis are available in Supplemental Data S6.

Fig. 7. Characterization of interactions between *GI* and *miR172*. A-C, Evaluation of the effect of

gi-2 mutation on the transcript levels of (A) mature *miR172,* (B) *AP2L1*, and (C) *FT1* by qRT-

PCR analysis. The mature *miR172* levels were calculated using the small nucleolar RNA 101

(SnoR) as internal reference, and the transcript levels of *AP2L1* and *FT1* were calculated relative

to the internal reference *ACTIN* using the ΔCt method. Samples were collected at ZT4 from

levels 7 and 9 of Kronos-PS (WT) and *gi-2* mutant plants grown under LD. Values are averages

of 5 biological replicates and error bars are standard errors of the means. $* = P < 0.05$, $** = P$

<0.01, $*** = P < 0.001$ indicate differences between the wild type and *gi-2* mutant using two-tail

t-test. D, Interaction graph showing the effect of the *UBI-miR172* transgene on heading time

under LD in the presence and absence of *GI* functional alleles (n = 13-15). $*** = P < 0.001$

indicate the significance of the four simple effect ANOVAs. Raw data and statistical analysis of

the complete 2 x2 factorial ANOVA is available in Supplemental Data S7.

Fig. 8. Regulation of heading time in wheat. Blue arrows indicate transcriptional promotion, red

lines transcriptional repression, orange lines posttranscriptional regulation, and grey lines genetic

interactions.

Fig. 1. Effect of *gi* loss-of-function mutants and transgenic lines expressing *GI* under the maize ubiquitin promoter (*UBI-GI-MYC*) on heading time under long-day (LD). A, Schematic diagram showing *GI* gene structure and the positions of selected mutations. Exons are represented by rectangles and introns by a line. Mutant line *A-2019* was combined with *B-2205* to generate *gi-1*, *A-401* combined with *B-2205* to generate *gi-2*, and *A-401* combined with *B-3825* to generate *gi-3* (letters indicate the genomes, and numbers indicate the Kronos mutant identification). B, *gi* mutants in Kronos-PS show a significant delay in heading time relative to their control sister lines under LD. C, The *UBI-GI-MYC* transgene largely rescues the delayed heading time of *gi-2* mutant in Kronos-PS background under LD. Different letters above the bars indicate significant differences in Tukey test = *P* < 0.01. D-E, Kronos-PS spikes of *gi-2* mutants produce more spikelets per spike (SNS) than those of the wild-type. The scale bar in D = 1 cm. F, Kronos-PS 12-week-old *gi-2* mutant compared to wild-type sister lines. G, The *gi-2* mutant plants produce more spike-bearing tillers than the control sister plants. Numbers inside bars indicate the number of plants included in each analysis. Error bars are standard errors of the means. *** = *P* < 0.001 for differences with the wild-type control using two-tailed *t*-test in B, E and G. Raw data and statistical analyses are available in Supplemental Data S1.

Fig. 2. Interactions between *GI* and photoperiod in Kronos-PS, Kronos-PI and *ppd1* mutants. A, Heading time evaluation of Kronos-PS and *gi-2* mutant grown under SD (8h light). Arrows on top of bars indicate that both genotypes failed to head before the experiment was terminated at 150 d. B, Pictures of developing young spikes of Kronos-PS and *gi-2* mutant dissected 100 d after germination from the same batch of plants grown under SD as in A. Scale bar = 1000 μm. C, Effect of *gi-2* on leaf number under SD-8h, SD-10h and LD-16h in a Kronos-PS background. Numbers inside the bars in A and C indicate the number of plants included in the analysis for each genotype. D, Interaction graph showing the effect of *gi-2* on days to heading under SD-8h and LD-16h in a Kronos-PI background (n = 10 - 11). E, Interaction graph showing the effect of *gi-2* and *ppd1* mutants on heading time under LD. Error bars are standard errors of the means based on n = 7 - 8. F, Effect of *gi* mutations on *PPD1* transcription profiles under LD. The gray shaded area indicates the 8 h of darkness. Primers used in the qRT-PCR amplify both *PPD1* homeologs. The 4th fully expanded leaves were collected at 4 h intervals for 24h. Expression levels are expressed relative to *ACTIN* as endogenous control using the ΔCt method. Error bars are standard errors of the means based on n = 5. ns = not significant, ** = $P < 0.01$, *** = $P < 0.001$ based on two-tail *t*-tests for each time point. Raw data and statistical analyses are available in Supplemental Data S2.

Fig. 3. Interactions between *GI* and *ELF3* on the regulation of flowering/heading time and *FT1* expression. A, Interaction graph showing the effect of the *gi-2* mutation on days to heading under different photoperiods in a Kronos-PS-*elf3* mutant background *(elf3* vs. *elf3 gi-2)*. SD = 8 h light and LD = 16 h light (n = 10-12). B and C, Interaction graphs for the effects of *GI* and *ELF3* on heading time (B) and total number of leaves produced by the primary shoots (C) under LD. Error bars are standard errors of the means based on n =8-10. *** = *P* < 0.001 for the four simple effect ANOVAs. The 2 x2 factorial ANOVAs are in Supplemental Data S3. D, Time- course expression analysis of *FT1* in the 4th fully expanded leaves of Kronos-PS (WT), *elf3*, *gi-2* and *elf3 gi-2* mutants under LD. The gray shaded area indicates the 8 h of darkness. The inset shows *FT1* expression from the same experiment but excluding the *elf3* mutant to visualize better the smaller differences among the other genotypes. Expression levels were calculated relative to the endogenous control *ACTIN* using the ΔCt method (n = 5). Raw data and statistical analyses are available in Supplemental Data S3.

Fig. 4 Effect of the *elf3* and *gi-2* mutations on the transcription profile of *GI* and the core circadian oscillator genes. Samples were collected at 4h intervals for a 24h period from the 4th fully expanded leaves. Expression levels of the target genes relative to the *ACTIN* endogenous control were evaluated by qRT-PCR using the ΔCt method. A, Effect of the *elf3* mutation on *GI* transcript levels. *** = $P < 0.001$ based on two-tailed t-test. The shaded gray area indicates the 8 h of darkness. Error bars are s.e.m based on 5 biological replications. B and C, Diurnal expression patterns of the core circadian clock genes *CCA1* (B) and *TOC1* (C) in Kronos-PS (WT), *elf3*, *gi-2,* and combined mutant *elf3 gi-2*. Five biological replicates were used per genotype. Raw data and factorial ANOVAs at each time point are provided in Supplemental Data S4.

Fig. 5. Interactions between *GI* and *VRN2*. A-B, Interaction graphs showing the effect of *GI* on heading time (A) and leaf number (B) in the presence and absence of functional *VRN2* alleles (n = 10-12). C-D, Effect of *gi-2* mutation on the transcript levels of *VRN2* (C) and *VRN1* (D) under LD. Each data point represents the average of 5 biological replicates. Samples were collected at 4h intervals for a 24h period from the 4th fully expanded leaves and transcript levels were evaluated by qRT-PCR and calculated relative to the *ACTIN* endogenous control using the ΔCt method. Error bars are standard errors of the means. ns = not significant, * = $P < 0.05$, ** = $P < 0.01$, and *** = $P < 0.001$ were obtained from simple effect ANOVAS in A and B (complete ANOVA in Supplemental Data S5) and two-tailed t-tests in C and D at each time point. Raw data is available in Supplemental Data S5.

Fig. 6. Interactions among *GI, CO1* and *CO2* on the regulation of wheat heading time under LD photoperiod. A-B, Effect of *gi-2* mutation on the expression profiles of *CO1* (A) and *CO2* (B) under LD (16h light). The gray shadowed area indicates the hours of darkness. The 4th fully expanded leaves were collected at 4 h intervals for a 24 h period. Expression levels were calculated relative to the endogenous control *ACTIN* using the ΔCt method. Values are averages of 5 biological replicates and error bars are standard errors of the means. ns = not significant, * = $P < 0.05$, ** = $P < 0.01$, *** = $P < 0.001$ for differences between genotypes at each time point using t-tests. C, Heading time of wild-type Kronos-PS (WT), *co1, co2, co1 co2, gi, co1 gi, co2 gi* and *co1 co2 gi* homozygous mutants grown in the same growth chamber under LD. The number of plants included in the experiment are indicated inside the bars. Error bars are standard errors of the means. Different letters with the same color indicate significant differences in Tukey tests ($P < 0.05$) in separate ANOVAs for *Gi* (blue) and *gi-2* (orange) genotypes. Raw data and statistical analysis are available in Supplemental Data S6.

Fig. 7. Characterization of interactions between *GI* and *miR172*. A-C, Evaluation of the effect of *gi-2* mutation on the transcript levels of (A) mature *miR172*, (B) *AP2L1*, and (C) *FT1* by qRT-PCR analysis. The mature *miR172* levels were calculated using the small nucleolar RNA 101 (SnoR) as internal reference, and the transcript levels of *AP2L1* and *FT1* were calculated relative to the internal reference *ACTIN* using the ΔCt method. Samples were collected at ZT4 from levels 7 and 9 of Kronos-PS (WT) and *gi-2* mutant plants grown under LD. Values are averages of 5 biological replicates and error bars are standard errors of the means. * = $P < 0.05$, ** = $P < 0.01$, *** = $P < 0.001$ indicate differences between the wild type and *gi-2* mutant using two-tail *t*-test. D, Interaction graph showing the effect of the *UBI-miR172* transgene on heading time under LD in the presence and absence of *GI* functional alleles (n = 13-15). *** = $P < 0.001$ indicate the significance of the four simple effect ANOVAs. Raw data and statistical analysis of the complete 2 x 2 factorial ANOVA is available in Supplemental Data S7.

Fig. 8. Regulation of heading time in wheat. Blue arrows indicate transcriptional promotion, red lines transcriptional repression, orange lines posttranscriptional regulation, and grey lines genetic interactions.

Parsed Citations

Abdul-Awal SM, Chen J, Xin Z, Harmon FG (2020) A sorghum gigantea mutant attenuates florigen gene expression and delays flowering time. Plant Direct. 4(11): e00281.
Google Scholar: Author Only Title Only Author and Title

Alqudah AM, Sharma R, Pasam RK, Graner A, Kilian B, Schnurbusch T (2014) Genetic dissection of photoperiod response based on GWAS of pre-anthesis phase duration in spring barley. PLoS ONE. 9(11): e113120.
Google Scholar: Author Only Title Only Author and Title

Alvarez MA, Li C, Lin H, Joe A, Padilla M, Woods DP, Dubcovsky J (2023) EARLY FLOWERING 3 interactions with PHYTOCHROME B and PHOTOPERIOD1 are critical for the photoperiodic regulation of wheat heading time. PLoS Genet. 19: e1010655.
Google Scholar: Author Only Title Only Author and Title

Alvarez MA, Tranquilli G, Lewis S, Kippes N, Dubcovsky J (2016) Genetic and physical mapping of the earliness per se locus Eps-Am1 in Triticum monococcum identifies EARLY FLOWERING 3 (ELF3) as a candidate gene. Funct Integr Genomics. 16: 365-382.
Google Scholar: Author Only Title Only Author and Title

Anwer MU, Davis A, Davis SJ, Quint M (2020) Photoperiod sensing of the circadian clock is controlled by EARLY FLOWERING 3 and GIGANTEA. Plant J. 101: 1397-1410.
Google Scholar: Author Only Title Only Author and Title

Araki T, Komeda Y (1993) Analysis of the role of the late-flowering locus, GI, in the flowering of Arabidopsis thaliana. Plant Journal 3: 231-239.
Google Scholar: Author Only Title Only Author and Title

Beales J, Turner A, Griffiths S, Snape JW, Laurie DA (2007) A pseudo-response regulator is misexpressed in the photoperiod insensitive Ppd-D1a mutant of wheat (Triticum aestivum L.). Theor Appl Genet. 115: 721-733.
Google Scholar: Author Only Title Only Author and Title

Brandoli C, Petri C, Egae-Cortines M and Weiss J (2020) Gigantea: uncovering new functions in flower development. Genes. 11: 1142.
Google Scholar: Author Only Title Only Author and Title

Chen A, Dubcovsky J. (2012) Wheat TILLING mutants show that the vernalization gene VRN1 down-regulates the flowering repressor VRN2 in leaves but is not essential for flowering. PLoS Genet. 8(12): e1003134.
Google Scholar: Author Only Title Only Author and Title

David KM, Armbruster U, Tama N, Putterill J (2006) Arabidopsis GIGANTEA protein is post-transcriptionally regulated by light and dark. FEBS Lett. 580: 1193-1197.
Google Scholar: Author Only Title Only Author and Title

Debernardi JM, Woods DP, Li K, Li C, Dubcovsky J (2022) MiR172-APETALA2-like genes integrate vernalization and plant age to control flowering time in wheat. PLoS Genet 18(4): e1010157
Google Scholar: Author Only Title Only Author and Title

Díaz A, Zikhali M, Turner AS, Isaac P, Laurie DA (2012) Copy number variation affecting the Photoperiod-B1 and Vernalization-A1 genes is associated with altered flowering time in wheat (Triticum aestivum). PLoS One. 7(3): e33234.
Google Scholar: Author Only Title Only Author and Title

Distelfeld A, Tranquilli G, Li C, Yan L and Dubcovsky J (2009) Genetic and molecular characterization of the VRN2 loci in tetraploid wheat. Plant Physiol. 149: 245-257.
Google Scholar: Author Only Title Only Author and Title

Ezer D, Jung JH, Lan H, Biswas S, Gregoire L, Box MS, Charoensawan V, Cortijo S, Lai X, Stöckle D, Zubieta C, Jaeger KE and Wigge PA (2017) The evening complex coordinates environmental and endogenous signals in Arabidopsis. Nat. Plants. 3: 17087.
Google Scholar: Author Only Title Only Author and Title

Fowler S, Lee K, Onouchi H, Samach A, Richardson K, Morris B, Coupland G, Putterill J (1999) GIGANTEA: A circadian clock-controlled gene that regulates photoperiodic flowering in Arabidopsis and encodes a protein with several possible membrane-spanning domains. EMBO J. 18: 4679–4688.
Google Scholar: Author Only Title Only Author and Title

Hayama R, Yokoi S, Tamaki S, Yano M, Shimamoto K (2003) Adaptation of photoperiodic control pathways produces short-day flowering in rice. Nature. 422: 719-722.
Google Scholar: Author Only Title Only Author and Title

Hecht V, Knowles CL, Vander Schoor JK, Liew LC, Jones SE, Lambert MJ, Weller JL (2007) Pea LATE BLOOMER1 is a GIGANTEA ortholog with roles in photoperiodic flowering, deetiolation, and transcriptional regulation of circadian clock gene homologs.

Plant Physiol. 144: 648-661.
  Google Scholar: Author Only Title Only Author and Title

Hong SY, Lee S, Seo PJ, Yang MS, Park CM (2010) Identification and molecular characterization of a Brachypodium distachyon GIGANTEA gene: functional conservation in monocot and dicot plants. Plant Mol Biol. 72: 485-497.
  Google Scholar: Author Only Title Only Author and Title

Huq E, Tepperman JM, Quail PH (2000) GIGANTEA is a nuclear protein involved in phytochrome signaling in Arabidopsis. Proc Natl Acad Sci USA. 97: 9789-9794.
  Google Scholar: Author Only Title Only Author and Title

Hwang DY, Park S, Lee S, Lee SS, Imaizumi T, Song YH (2019) GIGANTEA regulates the timing stabilization of CONSTANS by altering the interaction between FKF1 and ZEITLUPE. Mol Cells. 42: 693-701.
  Google Scholar: Author Only Title Only Author and Title

Imaizumi T and Kay SA (2006) Photoperiodic control of flowering: not only by coincidence. Trends Plant Sci. 11: 550-558.
  Google Scholar: Author Only Title Only Author and Title

Izawa T, Mihara M, Suzuki Y, Gupta M, Itoh H, Nagano AJ, Motoyama R, Sawada Y, Yano M, Hirai MY, Makino A, Nagamura Y (2011) Os-GIGANTEA confers robust diurnal rhythms on the global transcriptome of rice in the field. Plant Cell. 23:1741-1755.
  Google Scholar: Author Only Title Only Author and Title

Jung JH, Seo YH, Seo PJ, Reyes JL, Yun J, Chua NH, Park CM (2007) The GIGANTEA-regulated microRNA172 mediates photoperiodic flowering independent of CONSTANS in Arabidopsis. Plant Cell. 19: 2736–2748.
  Google Scholar: Author Only Title Only Author and Title

Koo BH, Yoo SC, Park JW, Kwon CT, Lee BD, An G, Zhang Z, Li J, Li Z, Paek NC (2013) Natural variation in OsPRR37 regulates heading date and contributes to rice cultivation at a wide range of latitudes. Mol Plant. 6: 1877-1888.
  Google Scholar: Author Only Title Only Author and Title

Koornneef M, Hanhart CJ, van der Veen JH (1991) A genetic and physiological analysis of late flowering mutants in Arabidopsis thaliana. Mol Gen Genet. 229: 57-66.
  Google Scholar: Author Only Title Only Author and Title

Krasileva KV, Vasquez-Gross HA, Howell T, Bailey P, Paraiso F, Clissold L, Simmonds J, Ramirez-Gonzalez RH, Wang X, Borrill P, Fosker C, Ayling S, Phillips AL, Uauy C, Dubcovsky J (2017) Uncovering hidden variation in polyploid wheat. Proc Natl Acad Sci USA. 114: E913-E921.
  Google Scholar: Author Only Title Only Author and Title

Lu SX, Webb CJ, Knowles SM, Kim SH, Wang Z, Tobin EM (2012) CCA1 and ELF3 interact in the control of hypocotyl length and flowering time in Arabidopsis. Plant Physiol. 158: 1079-1088.
  Google Scholar: Author Only Title Only Author and Title

Mishra P and Panigrahi KC (2015) GIGANTEA – an emerging story. Front. Plant Sci. 6: 8.
  Google Scholar: Author Only Title Only Author and Title

Murphy RL, Klein RR, Morishige DT, Brady JA, Rooney WL, Miller FR, Dugas DV, Klein PE, Mullet JE (2011) Coincident light and clock regulation of pseudoresponse regulator protein 37 (PRR37) controls photoperiodic flowering in sorghum. Proc Natl Acad Sci USA. 108: 16469-16474.
  Google Scholar: Author Only Title Only Author and Title

Park DH, Somers DE, Kim YS, Choy YH, Lim HK, Soh MS, Kim HJ, Kay SA, Nam HG (1999) Control of circadian rhythms and photoperiodic flowering by the Arabidopsis GIGANTEA gene. Science. 285: 1579-1582.
  Google Scholar: Author Only Title Only Author and Title

Pearce S, Shaw LM, Lin H, Cotter JD, Li C, Dubcovsky J (2017) Night-break experiments shed light on the Photoperiod1-mediated flowering. Plant Physiol. 174: 1139-1150.
  Google Scholar: Author Only Title Only Author and Title

Rédei GP (1962) Supervital mutants of Arabidopsis. Genetics. 47: 443-460.
  Google Scholar: Author Only Title Only Author and Title

Sawa M, Kay SA (2011) GIGANTEA directly activates Flowering Locus T in Arabidopsis thaliana. Proc. Natl. Acad. Sci. 108: 11698–11703.
  Google Scholar: Author Only Title Only Author and Title

Sawa M, Nusinow DA, Kay SA and Imaizumi T (2007) FKF1 and GIGANTEA complex formation is required for day-length measurement in Arabidopsis. Science. 318: 261-265.
  Google Scholar: Author Only Title Only Author and Title

Shaw LM, Li C, Woods DP, Alvarez MA, Lin H, Lau MY, Chen A, Dubcovsky J (2020) Epistatic interactions between

PHOTOPERIOD1, CONSTANS1 and CONSTANS2 modulate the photoperiodic response in wheat. PLoS Genet. 16(7): e1008812.
Google Scholar: Author Only Title Only Author and Title

Song YH, Estrada DA, Johnson RS, Kim SK, Lee SY, MacCoss MJ, Imaizumi T (2014) Distinct roles of FKF1, Gigantea, and Zeitlupe proteins in the regulation of Constans stability in Arabidopsis photoperiodic flowering. Proc Natl Acad Sci USA. 111: 17672-17677.
Google Scholar: Author Only Title Only Author and Title

Tseng TS, Salomé PA, McClung CR, Olszewski NE (2004) SPINDLY and GIGANTEA interact and act in Arabidopsis thaliana pathways involved in light responses, flowering, and rhythms in cotyledon movements. Plant Cell. 16: 1550-1563.
Google Scholar: Author Only Title Only Author and Title

Valverde F, Mouradov A, Soppe W, Ravenscroft D, Samach A, Coupland G (2004) Photoreceptor regulation of CONSTANS protein in photoperiodic flowering. Science. 303: 1003-1006.
Google Scholar: Author Only Title Only Author and Title

Wilhelm EP, Turner AS, Laurie DA (2009) Photoperiod insensitive Ppd-A1a mutations in tetraploid wheat (Triticum durum Desf.). Theor Appl Genet. 118: 285-294.
Google Scholar: Author Only Title Only Author and Title

Woods DP, McKeown MA, Dong Y, Preston JC, Amasino RM (2016) Evolution of VRN2/Ghd7-like genes in vernalization-mediated repression of grass flowering. Plant Physiol. 170: 2124-2135.
Google Scholar: Author Only Title Only Author and Title

Würschum T, Rapp M, Miedaner T, Longin CFH, Leiser WL (2019) Copy number variation of Ppd-B1 is the major determinant of heading time in durum wheat. BMC Genet. 20: 64.
Google Scholar: Author Only Title Only Author and Title

Xue W, Xing Y, Weng X, Zhao Y, Tang W, Wang L, Zhou H, Yu S, Xu C, Li X, Zhang Q (2008) Natural variation in Ghd7 is an important regulator of heading date and yield potential in rice. Nat Genet. 40: 761-767.
Google Scholar: Author Only Title Only Author and Title

Yan L, Loukoianov A, Blechl A, Tranquilli G, Ramakrishna W, SanMiguel P, Bennetzen JL, Echenique V, Dubcovsky J (2004) The wheat VRN2 gene is a flowering repressor down-regulated by vernalization. Science. 303: 1640-1644.
Google Scholar: Author Only Title Only Author and Title

Yang S, Weers BD, Morishige DT, Mullet JE (2014) CONSTANS is a photoperiod regulated activator of flowering in sorghum. BMC Plant Biol. 14: 148.
Google Scholar: Author Only Title Only Author and Title

Yano M, Katayose Y, Ashikari M, Yamanouchi U, Monna L, Fuse T, Baba T, Yamamoto K, Umehara Y, Nagamura Y, Sasaki T (2000) Hd1, a major photoperiod sensitivity quantitative trait locus in rice, is closely related to the Arabidopsis flowering time gene CONSTANS. Plant Cell. 12: 2473-2484.
Google Scholar: Author Only Title Only Author and Title

Yu JW, Rubio V, Lee NY, Bai S, Lee SY, Kim SS, Liu L, Zhang Y, Irigoyen ML, Sullivan JA, Zhang Y, Lee I, Xie Q, Paek NC, Deng XW (2008) COP1 and ELF3 control circadian function and photoperiodic flowering by regulating GI stability. Mol Cell. 32: 617–630.
Google Scholar: Author Only Title Only Author and Title

Zagotta MT, Hicks KA, Jacobs CI, Young JC, Hangarter RP, Meeks-Wagner DR (1999). The Arabidopsis ELF3 gene regulates vegetative photomorphogenesis and the photoperiodic induction of flowering. Plant J. 10: 691-702.
Google Scholar: Author Only Title Only Author and Title

Zheng T, Sun J, Zhou S, Chen S, Lu J, Cui S, Tian Y, Zhang H, Cai M, Zhu S, Wu M, Wang Y, Jiang L, Zhai H, Wang H, Wan J (2019) Post-transcriptional regulation of Ghd7 protein stability by phytochrome and OsGI in photoperiodic control of flowering in rice. New Phytol. 224: 306-320.
Google Scholar: Author Only Title Only Author and Title