Title: Collaborative Cross mice have diverse phenotypic responses to infection with

Methicillin-resistant *Staphylococcus aureus* USA300.

Short Title: MRSA Infection in CC mice

Authors: Aravindh Nagarajan[1,3], Kristin Scoggin[1,3], Jyotsana Gupta[3], Manuchehr

Aminian[6,7], L. Garry Adams[8], Michael Kirby[6], David Threadgill[1,2,4,5], Helene Andrews-

Polymenis [1, 3] *

Affiliations

[1]Interdisciplinary Program in Genetics and Genomics, Texas A&M University, College

Station, Texas, United States of America

[2]Department of Molecular and Cellular Medicine, Texas A&M University, College Station,

Texas, United States of America

[3]Department of Microbial Pathogenesis and Immunology, Texas A&M University,

College Station, Texas, United States of America

[4]Texas A&M Institute for Genome Sciences and Society, Texas A&M University, College

Station, Texas, United States of America

[5]Department of Biochemistry & Biophysics and Department of Nutrition, Texas A&M

University, College Station, Texas, United States of America

[6]Department of Mathematics, Colorado State University, Fort Collins, Colorado, United

States of America

[7]Department of Mathematics and Statistics, California State Polytechnic University,

Pomona, California, United States of America

[8]Department of Veterinary Pathobiology, Texas A&M University, College Station, Texas,

United States of America

*Corresponding author

Keywords: Host-pathogen interaction, MRSA USA300, Collaborative Cross, QTL

mapping, *Npc1*, *Ifi44l*, tolerance, sepsis, sexual dimorphism in infection

Abstract:

Staphylococcus aureus (*S. aureus*) is an opportunistic pathogen causing diseases

ranging from mild skin infections to life threatening conditions, including endocarditis,

pneumonia, and sepsis. To identify host genes modulating this host-pathogen

interaction, we infected 25 Collaborative Cross (CC) mouse strains with methicillin-

resistant *S. aureus* (MRSA) and monitored disease progression for seven days using a

surgically implanted telemetry system. CC strains varied widely in their response to

intravenous MRSA infection. We identified eight 'susceptible' CC strains with high

bacterial load, tissue damage, and reduced survival. Among the surviving strains, six

with minimal colonization were classified as 'resistant', while the remaining six tolerated

higher organ colonization ('tolerant'). The kidney was the most heavily colonized organ,

but liver, spleen and lung colonization were better correlated with reduced survival.

Resistant strains had higher pre-infection circulating neutrophils and lower post-infection

tissue damage compared to susceptible and tolerant strains. We identified four CC

strains with sexual dimorphism: all females survived the study period while all males

met our euthanasia criteria earlier. In these CC strains, males had more baseline

circulating monocytes and red blood cells. We identified several CC strains that may be

useful as new models for endocarditis, myocarditis, pneumonia, and resistance to

MRSA infection. Quantitative Trait Locus (QTL) analysis identified two significant loci, on

Chromosomes 18 and 3, involved in early susceptibility and late survival after infection.

We prioritized *Npc1* and *Ifi44l* genes as the strongest candidates influencing survival

using variant analysis and mRNA expression data from kidneys within these intervals.

<u>Author Summary:</u>

Methicillin-resistant *Staphylococcus aureus* is a human opportunistic pathogen that can

cause life-threatening diseases. To study the influence of host genetics on the outcome

of MRSA infection, we infected a collection of genetically diverse mice. We identified

different phenotypes for survival, organ colonization, and tissue damage, and classified

CC strains into MRSA susceptible, tolerant, and resistant categories. We identified

several parameters that correlated with these phenotypes. Four CC strains exhibited

strong sexual dimorphism in infection outcome: females lived longer, and males had

higher baseline circulating monocytes and red blood cells. Several of the CC strains we

characterize may represent better animal models for diseases caused by MRSA. QTL

analysis identified two genes, *Npc1* and *Ifi44l*, as strong candidates for involvement in

early susceptibility and late survival after MRSA infection. Our data suggests a strong

involvement of host genetics in MRSA infection outcome.

Introduction

Staphylococcus aureus, a gram-positive organism, is a commensal and

opportunistic pathogen that can cause severe morbidity and mortality [1, 2]. This

organism colonizes 20-30% of humans permanently and another 50-60% intermittently

[3, 4]. *S. aureus* colonization is a risk factor for subsequent infections ranging from mild

skin and soft tissue infections to serious invasive infections and death [5-7]. *S. aureus* is

a multi-host pathogen that can colonize and cause disease in cows, chickens, pigs,

sheep, and wild animals [8]. Methicillin-resistant *S. aureus* is resistant to β-lactam

antibiotics [9], and was primarily associated with hospital-acquired infections until the

1980s [10, 11]. In the 1990s, MRSA spread rapidly in the community and is now

considered a global threat [12]. MRSA causes more than 300,000 hospitalizations and

10,000 deaths in the US annually, at an estimated cost of 1.7 billion dollars (CDC,

2018).

USA300 is a highly successful community-acquired clone of MRSA. First

reported in 1999, by 2011 USA300 became the primary cause of severe skin and soft-

tissue infections, bacteremia, and community-onset pneumonia [13]. The increased

disease burden caused by MRSA USA300 is attributed to the presence of multiple

mobile genetic elements (MGEs) carrying virulence factors. In humans, USA300

infections are associated with increased mortality and increased incidence of severe

sepsis than hospital-acquired MRSA infections [14, 15]. The differences in colonization

rates, diversity in disease progression and mortality, exhibited by *S. aureus* suggests

the involvement of host genetics.

Studies investigating the pathogenesis of *S. aureus* have relied heavily on a

limited number of inbred mouse strains [16, 17]. *S. aureus* infections in mice induce a

diverse spectrum of diseases similar to what is observed in humans [17]. Intravenous

infection in mice with *S. aureus* is commonly used as a model for sepsis with this

pathogen [17-20]. Following intravenous inoculation with up to 10^7 organisms, *S. aureus*

Newman (a methicillin-sensitive strain) eventually disseminates into various tissues,

leading to the formation of abscesses in the vasculature, lung, heart, liver, spleen, and

kidney [21]. It can take up to a month for these accumulating lesions to cause death in

mice [21]. At higher doses (5×10^7-5×10^8 CFU), mice develop septic shock within 12-48

hours leading to death [22]. Infection with MRSA USA300 in mice follows a similar

pattern of organ colonization [23].

Very little is known about the host genes and gender differences that influence *S.*

aureus infections in mice. Targeted deletion of MyD88 (myeloid differentiation primary

response protein, C57Bl6 x 129S1 intercross) and NOD2 (nucleotide-binding

oligomerization domain-containing protein 2) render mice highly susceptible to *S.*

aureus infection [24, 25]. Outcomes after infection with *S. aureus* differ depending on

the strain of laboratory mice used. In a bacteremia model, C57BL/6J mice are the most

resistant in controlling bacterial growth and survive, while A/J, DBA/2, and BALB/c mice

are very susceptible [26]. This information has been used to map a limited number of

loci involved in this differential outcome. Using C57BL/6J and A/J Chromosomal

substitution strains (CSS), six genes on three different chromosomes (Chr 11 – *Dusp3*,

Psme3; Chr 8 – *Crif1, Cd97*; Chr 18 – *Tnfaip8, Seh1l*) have been implicated to

susceptibility after infection with *S. aureus* (sanger 476) strain [27-29]. To our

knowledge, no study has looked at the host factors involved in USA300 pathogenesis, a

hypervirulent isolate compared to other strains [30-32]. Furthermore, no screen for

MRSA infection disease outcomes has been performed across a broad range of host

genetics.

Collaborative Cross mice have been successfully used to create disease models

and identify susceptibility genes for various infections [33]. The CC is a large panel of

recombinant inbred mouse strains created by interbreeding five classical inbred strains

(A/J, C57BL/6J, 129S1/SvlmJ, NOD/ShiLtJ, NZO/HlLtJ) and three wild-derived inbred

strains (CAST/EiJ, PWK/PhJ, and WSB/EiJ) using a funnel-breeding scheme [34, 35].

CC strains have more than 30 million Single Nucleotide Polymorphisms (SNPs), and

their genomes are fully sequenced, making them an ideal population for QTL mapping

[36, 37]. The CC has been heavily used to identify susceptibility to viral pathogens,

including Ebola virus [38], West Nile virus [39], Influenza A virus [40], Coronaviruses

[41, 42], Theiler's murine encephalomyelitis virus [43], Cytomegalovirus [44] and Rift

Valley fever virus [45]. The CC has also been used to study infection with bacterial

pathogens, including *Mycobacterium tuberculosis* [46, 47], *Klebsiella pneumonia* [48],

Pseudomonas aeruginosa [49], *Borrelia recurrentis* [50], *Salmonella* Typhimurium [51,

52] and *Salmonella* Typhi [53].

We were interested in how host genetics influences disease outcome after MRSA

bacteremia. Therefore, we screened 25 CC strains for disease outcome phenotypes

after intravenous infection with MRSA USA300. We chose a 7-day screening period and

very sensitive telemetry monitoring of MRSA infected animals in an attempt to capture

and identify range of disease phenotypes from highly susceptible to resistant. Using this

model system, we identified a wide range of disease outcomes, potential new MRSA

disease models, and genomic regions linked to survival using QTL mapping. In order to

prioritize the genes within these intervals we used variant analysis and mRNA

expression data from kidneys.

Results

<u>CC strains have diverse disease outcomes after MRSA infection.</u>

We screened twenty-five Collaborative Cross strains (three males and three

females per strain) for their phenotypes after intravenous infection with MRSA USA300

(Table S1). Because blood borne infections can become rapidly fatal, we used a 7-day

screening period to allow us to delineate sensitive and resistant CC lines. Animals were

monitored for clinical symptoms using a continuously monitored telemetry system and a

manual health scoring method up to one-week post-infection to identify those

susceptible to MRSA infection. At the time of necropsy spleen, liver, heart, lung, and

kidney were collected to enumerate the bacterial load and tissue damage (Table S2).

C57BL/6J mice, known to survive MRSA infection, served as controls [26]. All

C57BL/6J mice survived until the end of the experiment (day 7) and were thus classified

as a surviving strain (Fig 1A). CC strains varied widely in their response to MRSA sepsis

(Fig 1 and Fig S1). Median survival ranged from 2.5 to 7 days (Fig 1A). Median weight

loss ranged from losing 20% of their body weight (CC002, CC037) to losing less than

1% (CC003, CC017, CC024) (Fig 1B). Out of the 156 infected mice, 10 mice belonging

to 6 different strains appeared to clear MRSA from all organs we collected (Fig 1D). We

included these mice in our analysis based on disturbances in circadian pattern following

infection and histology scores, suggesting that they did become infected (Fig S5 and

S6).

CC strains that had four or more mice of the six infected meeting our euthanasia

criteria before day seven were classified as 'susceptible'. Eight of the 25 CC strains we

infected were classified in this category (CC002, CC006, CC027, CC037, CC036,

CC053, CC057, CC061) (Fig 1A). Susceptible strains, on average, lost 16% of their

body weight (Fig 1B). Two susceptible strains, CC027 and CC057, had median kidney

colonization on the order of 10^9, ~100-fold higher than the inoculation dose (Fig 1C).

Five CC strains had three mice that met our euthanasia criteria before day 7. Of

these five strains, the survival of four of these strains was uniformly sexually dimorphic

during our experimental time period: all the males met our euthanasia criteria early in

the experimental period, while all females of these strains survived the full 7-day

infection (CC013, CC015, CC043, CC058) (Fig 1). These four CC strains were

classified as sexually dimorphic in their survival after MRSA infection. CC001, classified

as 'surviving', may also display sexual dimorphism; 2/3 males met our euthanasia

criteria prior to day 7, and the remaining male never recovered to a pre-infection

circadian pattern of body temperature or activity. Infection experiments of longer

duration will be required to determine if the sensitivity of males in this strain is uniform.

Finally, although CC042 is highly susceptible to other bacterial pathogens [46, 51, 54],

couldn't easily be classified with respect to MRSA infection outcome as two females and

one male survived infection.

We classified the remaining 12 surviving CC strains into tolerant and resistant

phenotypes. We defined tolerance as surviving the infection with a high bacterial burden

while limiting the health impact caused by the pathogen [55-57]. Six strains with higher

total organ colonization compared to the inoculum dose were classified as tolerant

(CC001, CC005, CC019, CC024, CC038, CC051) (Fig 1D). These strains limited the

impact to their health (based on circadian pattern and health scores) despite high

colonization in several organs (Fig S5 and Table S2). Although CC051 mice infected

with MRSA survived and were classified here as tolerant, they exhibited some weight

loss and a persistent disruption of their circadian pattern of body temperature and

activity. Infections of longer duration will be needed to determine whether CC051 mice

are truly tolerant or display a delayed susceptibility phenotype [52].

Surviving the infection by preventing colonization or clearing the pathogen was

defined as resistance [55-57]. The remaining six strains, with lower (compared to the

inoculum dose) or no colonization, were classified as resistant (CC003, CC012, CC017,

CC023, CC025, CC041) (Fig 1D). MRSA infection had minimal impact on the health of

these CC strains (Fig S5 and Table S2). The wide range of disease severity, survival,

weight loss, and organ colonization we observed during blood borne MRSA infection

suggest diverse host responses across genetically different CC strains.

Correlation of colonization, tissue damage and immune parameters with survival

Pre-infection weight did not influence survival (R = -0.05), while weight loss after

infection correlated with poor survival (R = -0.65) (Fig 2A). Colonization in all five organs

was highly correlated with poor survival (Fig 2A). The inability to limit bacterial growth in

the kidney has previously been associated with susceptibility to intravenous *S. aureus*

infections in mice [26]. In our experiments, the kidney was the most highly colonized

organ across all CC strains and high kidney colonization correlated with poor survival (R

= -0.58) (Fig 1C, 2A). Unexpectedly however, colonization in the liver (R = -0.78),

spleen (R = -0.73), and lung (R = -0.67) was more highly correlated with poor survival

than colonization of the kidneys (Fig 2A), despite the fact that these organs had lower

bacterial load than the kidneys (Fig 1C, 1D) and less tissue damage (Fig 2B). These

data suggest that the spleen, liver and lung are more intolerant to colonization by MRSA

and acute damage than the kidney during MRSA infection.

S. aureus causes damage to host tissue in mice as early as 48 hours after

intravenous challenge [21]. To assess tissue damage in the CC mouse population, a

board-certified pathologist scored each tissue blindly for damage (0 = normal to 4 =

severe damage) (Table S3 and Fig S2). Across all organs, resistant strains had

significantly less tissue damage than tolerant (P < 0. 001) and susceptible strains (P <

0.010) (Fig 2B). Interestingly, there was no significant difference in tissue damage

between tolerant and susceptible strains (P = 0.57) (Fig 2B). This finding suggests that

tolerant strains may survive MRSA infection in the face of both high colonization and

severe tissue damage.

High bacterial burden in kidneys translated to more significant tissue damage. Of

the 25 CC strains we studied, 16 had a median pathology score of 2.5 or above for

kidneys, indicating moderate to severe tissue damage (Fig 2B, Scale of 0-4, 0= no

damage, 4= severe damage). This damage did not translate to susceptibility however,

as histology scores across all five tissues that we sampled had only a weak correlation

with poor survival (Fig 2A). Spleen damage had the highest correlation R = -0.23 with

poor survival, followed by kidney (R = -0.18) and heart (R = -0.18) damage. This weak

correlation may be due to the diverse phenotypic responses exhibited by the CC strains

to MRSA infection. Some infected CC strains met our euthanasia criteria very early in

infection, perhaps before extensive tissue damage had time to occur. Other CC strains

lived to the end of the experiment, allowing time for severe tissue damage to occur and

become cumulative.

Pre-infection inflammation and immune parameters play a role in disease

prediction and progression [42, 58]. We collected blood from mice five weeks before

infection to perform Complete Blood Counts (CBC). Interestingly, CC strains resistant to

MRSA infection had a higher number of pre-infection circulating neutrophils than

susceptible and tolerant strains (Fig 2C). There were no significant differences in other

pre-infection CBC parameters (Table S2).

Survival is sex-dependent in four CC strains.

In the four sexually dimorphic strains, females lived until the end of the

experiment (day 7), but all the males met our euthanasia criteria much earlier (Fig 3A).

Males lost more weight than females (Fig 3B) and had higher liver, spleen, and lung

colonization (Fig 3C). Colonization did not differ significantly between males and

females in the kidneys, the highest colonized organ in these strains (Fig 3C). Females

of these CC strains appear to limit MRSA replication in some organs or spread from the

kidneys by unknown mechanisms.

To assess baseline differences between males and females of the strains that

were sexually dimorphic with respect to MRSA infection, we evaluated pre-infection

complete blood counts (CBC). Males had higher circulating monocytes and red blood

cells before infection than females of these strains (Fig 3D). The number of circulating

monocytes (P = 0.83) and red blood cells (P = 0.10) were not significantly different

between males and females in the rest of the CC population (21 strains) (Table S2).

Thus, this difference in the starting blood parameters is unique to CC strains that

displayed sexual dimorphism in susceptibility to MRSA infection.

<u>Baseline body temperature and baseline activity did not correlate with survival.</u>

To track the disease progression in infected mice, we used a combination of

automated telemetry and manual health scoring (Table S2). A surgically implanted

telemetry device tracked body temperature and activity levels every minute before and

throughout the infection (Fig S5 and Fig S6). Data from five days before infection was

used to create a baseline circadian pattern for each mouse. A machine-learning

algorithm was also used to predict when infected mice deviated from their normal

circadian pattern after infection. No significant differences were identified in the elapsed

time between infection and deviation from the normal circadian pattern between

surviving and susceptible strains (Table S2). 90% of the infected mice deviated from

their normal circadian pattern of body temperature and activity within 24 hours of

infection (Fig S3). The first health check performed by laboratory staff, 24 hours after

infection, could only detect 56% of these mice. In the next two days, twice daily health

checks detected all the remaining mice as ill (Fig S3). These data suggest a significant

lag time between circadian pattern disruption detected by telemetry and signs of illness

detected by laboratory staff and the effectiveness of the telemetry system in accurately

predicting survival after infection.

<u>Two independent genomic regions influence survival.</u>

We used RQTL2 software with CC genotypes imputed from qtl2qpi to identify

genomic regions associated with survival after MRSA infection [59, 60]. Within the CC

strains we infected, survival was a highly heritable trait with a broad sense heritability

score of 0.38 (Table S4). While using only 25 CC strains reduces the power of QTL

mapping [61], using 6 animals per strain and continuous monitoring using a telemetry

device added more power to our study and allowed us to sensitively define the

phenotypes we observed.

For early susceptibility on day 2, there was a highly significant peak on

chromosome 18 (Fig 4A). We named this QTL peak ESMI (Early Susceptibility in MRSA

Infection). This peak disappeared after day 3 post-infection. For survival on day 7 (late

survival), there was a highly significant peak on Chromosome 3 (Fig 4B). We named

this peak LSMI (Late Survival in MRSA Infection). To understand the impact of sex on

survival, we performed the same analysis after removing the four strains that exhibited

sexual dimorphism in survival. The early susceptibility peak on Chr 18 and late survival

peak on Chr 3 were retained (Fig S4), suggesting that sex did not influence our QTL

peaks. We compared the phenotype at the top SNPs in both peaks for potential

interaction (Table S5). No significant interactions were identified, suggesting that our

QTL regions act independently of each other.

<u>*Npc1* is a strong candidate for influencing early susceptibility during MRSA infection.</u>

294   The advantage of the CC panel over other outbred mouse panels is the high

295 number of SNPs and structural variants present across the CC genomes [36, 37]. We

296 used this feature to shortlist the genes within the associated genomic regions to identify

297 those of most interest. The early susceptibility QTL peak, ESMI, on chromosome 18 is a

298 5 Mb region between 12-17 Mb with a peak around 14.67 Mb. This region contains 21

299 protein-coding genes and 49 non-coding RNA genes. Because the CC strains are a

300 mosaic of eight inbred founder strains, we looked at the founder effect plot in this region

301 to attempt to identify each founder's contribution to survival (Table S5). We found that

302 the WSB allele in this region contributes to reduced survival, while A/J and NZO alleles

303 contribute to prolonged survival (Fig 5A). Using all the SNPs, insertions and deletions

304 that matched this allele effect pattern in the Mouse Phenome Database (MPD), we

305 identified 20,582 variants across all 21 genes. We used Variant Effect Predictor (VEP)

306 to further prioritize these variants based on their predicted effect on protein structure

307 and function, identifying 52 high and moderate-impact variants in 7 genes (Table S6).

308 The highest impact variant for each gene is listed (Fig 5B). Four of the seven genes we

309 shortlisted have previously been associated with immune function in the Mouse

310 Genome Informatics Portal (highlighted in red, Fig 5B) [62]. We sequenced mRNA from

311 kidneys collected from CC strains at necropsy (at least one male and one female per

312 strain) to further prioritize genes. Then, we looked for differentially expressed genes that

313 matched the founder effect pattern within this region. For the early susceptibility QTL

314 peak, two genes, Cadherin2 (*Cdh2*) and NPC intracellular cholesterol transporter 1

315 (*Npc1*), matched the founder effect pattern (Fig 5C). *Npc1* is our top candidate in this

316 region due to its previous association with immune functions [63, 64].

Ifi44l is a strong candidate for involvement in late survival after MRSA infection.

The late survival QTL peak, LSMI, on chromosome 3 is a 4 Mb region between

146-150 Mb with a peak near 146.66 Mb. This region contains 13 protein-coding genes

and 39 non-coding RNA genes. The A/J allele in this region reduced survival

significantly, while CAST, NOD, NZO, and WSB alleles promoted longer survival (Fig

6A). Using this founder effect pattern, we shortlisted five genes with variants predicted

to have a high impact on the function of the encoded proteins (Table S6). The top

variant for each gene is listed (Fig 6B). Two genes in the Interferon Stimulated Gene

family (ISG), Interferon induced proteins 44 and 44l genes (*Ifi44* and *Ifi44l*), are our

primary candidates at this locus. *Ifi44* has a missense mutation, and *Ifi44l* has a high-

impact mutation in the gene's promoter region. In mice and humans, *Ifi44l* is a paralog

of *Ifi44, but it remains* unclear if *Ifi44* and *Ifi44l* have redundant or distinct functions [65].

Using gene expression data, we determined that *Ifi44l*, is highly expressed in strains

with the A/J allele but not with the alleles from the other founders (Fig 6C). Thus, *Ifi44l*

became our top candidate gene influencing late survival during MRSA infection.

Discussion

We used a genetic screen of CC mouse strains to determine whether mice of

different genetic backgrounds have diverse disease outcomes after intravenous

infection with MRSA USA300, and ultimately to identify regions of the genome linked to

survival. We identified 8 CC strains that were highly susceptible to fatal infection, 4

strains with strong sexually dimorphic survival, 6 strains that appeared tolerant, and 6

strains that were resistant to MRSA infection (even totally clearing the pathogen in

several cases). Survival, bacterial colonization, and weight loss differed even within a

given phenotypic category. Across 8 CC strains that we determined to be susceptible to

MRSA sepsis based on reduced survival after infection, survival time ranged from 2.5 to

4 days, and kidney colonization varied as much as 10-fold. Two susceptible strains,

CC061 and CC036 had very similar kidney colonization, yet CC061 appeared to have a

more severe disease and lost twice as much weight as CC036 (19% vs. 9%). We also

observed similar diversity in weight loss and colonization within tolerant and resistant

strains. Our data support both the importance of host genetics in disease outcome after

MRSA infection, but also the hypothesis that there may be multiple mechanisms of

susceptibility, tolerance and resistance to MRSA infection across a broad set of host

genetics.

In both animals and humans, males are more susceptible than females to

bacterial infections [66]. Some of these differences can be attributed to the differences

in X-chromosome number and inactivation process between males and females. The X-

chromosome also encodes several critical immune genes that may play important roles

in sexually dimorphic infection outcomes [67]. Males have a higher overall risk for

MRSA carriage [68] and bloodstream infections [69, 70]. However, there is no

consensus on the role of gender on mortality [70-74]. Across all CC strains in our study

females survived longer than males. We determined that this difference was driven by 4

CC strains that exhibited strong sexual dimorphism in survival, despite similar levels of

kidney colonization in both sexes.

Two blood parameters, pre-infection baseline numbers of circulating monocytes

and red blood cells, were higher only in males of the CC strains that exhibited sexually

dimorphic survival phenotypes in response to MRSA infection. This finding suggests

that increased numbers of circulating monocytes and red blood cells may be linked to

susceptibility of males of these CC strains to severe MRSA infection. *S. aureus* persists

within monocyte-derived macrophages and can use these cells as a vehicle for

dissemination [75, 76]. Furthermore, heme iron is the preferred source of iron for *S.*

aureus during infection [77], and this organism can bind and lyse RBCs [78-80] and

directly bind hemoglobin [81] [82]. Thus, in males of these CC strains, increased pre-

infection circulating monocytes may facilitate more efficient MRSA dissemination while

increased heme availability may provide an additional source of iron for MRSA during

infection.

Neutrophils are the primary defense against *S. aureus* infections [7, 83], and

patients with impaired neutrophil numbers and/or function are predisposed to *S. aureus*

infections [84-86]. We noticed that pre-infection circulating neutrophil numbers were

significantly higher in our MRSA-resistant CC strains than in the tolerant and

susceptible strains. Our data is consistent with observations in humans, supporting a

critical role for neutrophils in resistance to MRSA infection in several CC strains.

Most of the CC strains we used in this study were also used to define disease

outcome after oral infection with non-typhoidal *Salmonella* [51, 52]. Six of the eight CC

strains that were susceptible to MRSA infection were also susceptible or delayed

susceptible to *Salmonella* infection (CC006, CC027, CC036, CC037, CC053, CC061).

The remaining two MRSA susceptible CC strains, CC002 and CC057, were resistant

and tolerant to *Salmonella* infection respectively [52]. One CC strain, CC038, was

tolerant to both MRSA and *Salmonella* infection. However, all of the remaining CC

strains we tested had different disease outcomes in MRSA infection than they did after

Salmonella infection. The phenotypes we discovered suggest that some of the

mechanisms of susceptibility to severe MRSA and *Salmonella* infection may be shared,

while the mechanisms of tolerance and resistance to these two pathogens are largely

distinct.

We compared our MRSA infection outcome data to a similar study of

Mycobacterium tuberculosis (Mtb) infection in CC strains [47]. While this study did not

classify survival phenotypes, we used the median lung colonization values as a rough

proxy for disease severity. For the 52 CC strains infected with Mtb, log lung CFU ranged

from 5.1 to 8.5, with a median of 6.7 CFU [47]. Among the six MRSA-susceptible CC

strains that were also infected in *Mtb* studies, two strains (CC027, CC037) were

sensitive to both *Salmonella* and MRSA infection and were also heavily colonized by

Mtb in the lung [47, 51, 52]. A common mechanism may underlie the susceptibility of

these CC strains to three very different bacterial pathogens, administered by different

routes.

We also identified CC strains that were not uniformly susceptible to all three

pathogens. Two CC strains susceptible to both MRSA and *Salmonella* (CC006, CC061)

were poorly colonized in the lung with *Mtb* (5.6 and 6.5 log CFU) [47]. Finally, all five

MRSA-resistant CC strains were highly colonized in the lung with *Mtb* (average – 8.4

log CFU), and thus appear to be potentially highly susceptible to *Mtb* infection. The

MRSA-tolerant strains had variable *Mtb* colonization in the lung. This differential

response to different bacterial pathogens suggests that most CC strains that we tested

do not have a significant general dysfunction of immunity, where we would expect broad

susceptibility to multiple pathogens. Future work will tease out specific mechanisms of

susceptibility and resistance to different pathogens.

Using the survival data that we generated in this study, we identified QTL linked

to both early susceptibility and late survival after intravenous MRSA infection. The

primary candidate gene in the early susceptibility QTL, *Npc1*, is in endosomal and

lysosomal membranes and mediates intracellular cholesterol trafficking [87]. In humans,

mutations in *NPC1* cause Niemann–Pick Type C disease, leading to massive

accumulation of cholesterol in lysosomes in all tissues and premature death [88].

Defects in *NPC1* can lead to increased lipid storage in macrophages and a chronic

inflammatory condition [89, 90]. *NPC1* is also the primary host factor responsible for

Ebola, and other filovirus, entry into host cells [91]. These viruses use *NPC1* on

endosomes and lysosomes to trigger a membrane fusion process that allows the

expulsion of the viral genome into the cytoplasm and subsequent viral replication [92].

Little is known about the role of *NPC1* in bacterial infections. In a hematopoietic

mouse model, *Npc1* mutant mice have a significant increase in the relative abundance

of *Staphylococcus* ssp. in their intestine [93], suggesting a role for *Npc1* in keeping

Staphylococcus ssp. at a low level in this niche. Mutations in *Npc1* impair steady-state

autophagosome maturation and interfere with the autophagic trafficking of bacteria to

lysosomes in macrophages [94]. *S. aureus* manipulates autophagy in both neutrophils

and macrophages to promote its survival and escape [95-97]. CC strains carrying the

WSB allele in this region have a missense mutation in *Npc1*. This altered form of *Npc1*

may interfere with autophagy in macrophages and neutrophils, such that *S. aureus* can

survive initial phagocytosis leading to the early susceptibility phenotype.

The A/J allele on Chromosome 18 contributed to poor survival of MRSA infected

mice on day 7. A/J mice are also highly susceptible to SH1000, a different strain of *S.*

aureus [26]. *Ifi44l* promotes macrophage differentiation during bacterial infection and

facilitates inflammatory cytokine secretion [98]. Overexpression of *Ifi44* or *Ifi44l* is

sufficient to control respiratory syncytial virus (RSV) infection [65]. *Ifi44l* represses the

interferon response, protecting the host from the negative effects of the innate immune

response during viral infection [99]. The A/J *Ifi44l* allele may be less effective in

controlling the interferon response, reducing the survival of *S. aureus* infected mice in

the later stages of infection.

A series of previous studies examined the genetic differences that influence *S.*

aureus Sanger-476 infection in A/J and C57BL/6J mice using a chromosome

substitution strain (CSS) model [27-29]. Although C57BL/6 and A/J mice are among the

CC founders, we did not identify any genes in common with this previous work. Like

USA300, Sanger-476 is a community-acquired strain but is sensitive to methicillin [100].

USA300 also has additional pathogenicity islands that are absent in the Sanger-476

strain [101] and also produces higher amounts of virulence factors compared to other

MRSA strains [30, 102, 103]. These genetic differences make USA300 distinct from,

and potentially more virulent than, other community-acquired clones of MRSA. On the

host side, the CC founders include six additional inbred strains, making the CC strain

collection considerably more genetically diverse than the CSS panel. The fact that the

genetic regions we identify as linked to survival in this study did not overlap with those

of the previous study is thus likely due to use of different *S. aureus* strains, more diverse

mouse strains and different mapping power [27]. However, given that earlier work using

the CSS panel identified increased inflammation and defective immune responses as

critical to MRSA resistance, we were encouraged to identify *Npc1* and *Ifi44l* as potential

influencers of survival, as these genes are also involved in modulating inflammation and

influencing immune responses.

The CC panel has previously been used to identify mouse strains for new model

generation [33]. Although it was not a goal of this study to identify new mouse models

for different MRSA caused syndromes, our analysis of pathology across different CC

strains suggested that several strains may be make suitable models to fill gaps in our

knowledge of cardiac disease, pneumonia, and recovery from *S. aureus* systemic

infections. *S. aureus* is the leading cause of infective endocarditis (IE), which is

associated with high mortality (40-50%) [104, 105]. Current models of IE require

anesthetizing C57BL6/J mice and physically damaging their heart valve using a catheter

[106, 107]. *S. aureus* is also the most common cause of bacterial myocarditis, with fatal

outcomes [108-110]. Currently, there are no murine models to study MRSA induced

myocarditis. In our study, three CC strains (CC013, CC036, C0042) had over 200-fold

greater colonization in the heart than C57BL/6J mice and had systemic embolic

bacterial purulent myocarditis. Similarly, pneumonia caused by *S. aureus* has a high

mortality rate [111]. In our study, CC013, CC058, and CC061, had significant lung

damage. Finally, we identified six MRSA-resistant CC strains (CC023, CC025, CC012,

CC003, CC041, and CC017) that robustly survive systemic infection, have lower

bacterial colonization, and in most cases less tissue damage than the traditional

resistant mouse model for MRSA: C57BL/6J mice [26, 29]. We hope that the lines we

list above will be studied in more detail and may provide a useful resource for various *S.*

aureus based syndromes.

In summary, using colonization, telemetry, survival, and histology data, we have

identified diverse disease outcomes after MRSA USA300 infection in and described two

novel genomic regions that influence survival at different stages after systemic infection.

In addition, our work suggests several interesting pre-infection correlates that appear to

influence survival after MRSA infection, and a connection between sex and

susceptibility to MRSA infection in certain genetic backgrounds. With several *S. aureus*

vaccines failing at human clinical trials [112, 113], our data support strong consideration

of host genetics as an important factor while designing therapeutics and vaccines

against pathogens. Finally, as the CC strains have now been infected with multiple

bacterial pathogens, future will work will use the phenotypic diversity of host responses

to infection of this mouse collection to define different mechanisms of susceptibility,

tolerance and resistance.

<u>Acknowledgements</u>

We thank Dr. Magnus Hook for providing the bacterial strain used in this study.

We also thank Dr. Karl Broman (University of Wisconsin) for helpful discussions and

troubleshooting problems in the rqtl2 google forum. We also would like to thank Dr.

Callie Kobayashi, Dr. Joana Rocha, Ms. Marissa Talamantes, Mr. Chris Bowden, Mr.

Connor Mathis, and Ms. Kaya Mariello for assistance with experiments. This work was

funded by the Defense Advanced Research Project Agency (DARPA), project DARPA

D17AP00004.

Materials and Methods

Ethics statement

All mouse studies followed the Guide for the Care and Use of Laboratory of Animals of the National Institutes of Health. The animal protocols (2015-0315 D and 2018-0488 D) were reviewed and approved by Texas A&M Institutional Animal Care and Use Committee (IACUC).

Bacterial strains and media

Methicillin-resistant *Staphylococcus aureus* isolate used in this study was the kind gift of Dr. Magnus Hook (Texas A&M Institute of Biosciences and Technology, Houston). USA300 is a fully virulent, community-acquired clone of MRSA. Strains were routinely cultured in Luria-Bertani (LB) broth and plates supplemented with antibiotics when needed at 50 mg/L Kanamycin Sulphate. For murine infections, strains were grown aerobically at 37^{0}C to a stationary phase in LB broth supplemented with Kanamycin.

Murine strains

Both conventional (C57BL/6J) and Collaborative Cross mice were used for these experiments. In total, three males and three females belonging to 25 different CC strains (chosen at random) and C57BL/6J were used (156 mice total). All mice were initially purchased from UNC's Systems Genetics Core Facility (SGCF). They were bred independently at the Division of Comparative Medicine at Texas A&M University (Table S1). Mice were fed Envigo Teklad Global 19% Protein Extrudent Rodent Diet (2919) or

Envigo Teklad Rodent Diet (8604) based on strain need. Mice were provided with

cardboard huts, food, and water ad libitum.

Placement of telemetry devices

Mice (5-7 weeks old) were anesthetized with an isoflurane vaporizer using the

Kent Scientific SomnoSuite Low-flow anesthesia system to implant telemetry devices. A

midline abdominal incision was made, and Starr Life Science G2 Emitter devices were

sutured to the ventral abdominal wall. The abdominal muscle layer was sutured with 5-0

vicryl, and the skin layer was closed using stainless steel wound clips. Buprenorphine

(0.0001 mg/g) was administered intraperitoneally before recovery from anesthesia. Mice

were monitored twice daily for surgical complications and humanely euthanized when

needed. Surgical clips were removed on the seventh day, and animals were allowed to

recover for one more week.

Infection with MRSA

After baseline scoring for health, 6 mice (now 8-12 weeks old) per CC strain were

infected with MRSA USA300. We chose a 7-day screening period and very sensitive

telemetry monitoring of MRSA infected animals in an attempt to capture and identify

range of disease phenotypes from highly susceptible to resistant. To minimize batch

effects, mice from a given CC strain were infected in different experiments across a two-

year period. This approach may have resulted in some within strain variability in

infection outcomes. Briefly, Mice were anesthetized using isoflurane and infected with 1

x 10^7 CFU in 50 µl of LB broth intravenously at the inferior fornix into the retro-orbital

sinus. For each experiment, mice were inoculated at the same time of day. Mice that

became moribund within 6 hours of infection were humanely euthanized and removed

from the experiment.

<u>Health monitoring</u>

After the recovery from surgery, mice were moved to a BSL-2 facility and

acclimated for 5-7 days. Individual cages containing implanted mice were placed onto

ER4000 receiver platforms and calibrated to receive signals from the implanted

telemetry devices. The outputs, body temperature (once per minute) and gross motor

activity (summation of movement per minute), were continuously fed to a computer

system for visualization (Fig S5 and Fig S6). This sensitive continuous monitoring

allowed us understand disease progression for each animal in real-time, as disruptions

of the normal circadian pattern of core body temperature after infection indicated

symptomatic MRSA infection. A machine-learning algorithm was used to identify the

time to deviation from the circadian pattern of body temperature and activity. Detailed

explanation of the methodology and calculations can be found here [51].

We manually scored four health parameters twice daily: physical appearance,

body conditioning, and provoked and unprovoked behavior. The scoring scale ranged

from 0-3, with zero = normal and three = abnormal (Table S2). Additionally, as a

measure of activity, four small nestlets were placed at each corner of the cage in the

evening. The following day, the number of nestlets moved from the cage corners into

the hut was noted.

<u>Euthanasia criteria</u>

After infection, mice were monitored continuously using telemetry readings and

twice daily using visual health scoring. For telemetry, euthanasia criteria were defined

by a sudden drop in temperature (3^0C or more). For manual heath scoring, the

euthanasia criteria were reached when the combined health score exceeded 8. Mice

that met one or both these criteria were humanely euthanized by CO2 asphyxiation.

Bacterial load determination

After euthanasia, whole blood, serum, and organs (spleen, liver, heart, lung, and

kidney) were collected. A consistent region of each organ was collected in 3mL ice-cold

PBS and homogenized. The serially diluted homogenate was plated on LB plates

supplemented with kanamycin for bacterial enumeration in each organ. Data are

expressed as CFU/g of tissue. Total organ colonization was calculated by adding CFU/g

values of all the five organs collected.

Histopathology

After euthanasia, portions of collected organs were fixed in 10% neutral buffered

formalin for 24 hours and stored in 70% ethanol. Fixed tissue was embedded in paraffin,

sectioned at 5 μm, and stained with hematoxylin and eosin (H&E). A board-certified

veterinary pathologist scored all the slides for tissue damage on a scale of 0 to 4 in a

blinded manner (Table S3). All the raw pathology data is available in the supplementary

file (Table S2). Whole slide images of H&E-stained tissue sections were captured as

digital files by scanning at 40X using a 3D Histech Pannoramic SCAN II FL Scanner

(Epredia, MI). Digital files were processed by Aiforia Hub (Cambridge, MA) software to

generate images with scaled bars.

Complete blood count

One week before the implantation of telemetry devices, whole blood was

collected from each mouse by submandibular bleeding. From the same mice, blood was

collected at necropsy after infection by cardiac puncture. Abaxis VetScan HM5,

optimized for rodents, was used to analyze blood collected in Ethylene Diamine Tetra

Acetic acid (EDTA) tubes. The ratio between the blood parameters after and before

infection (value >1 = increased after infection and value < 1 = decreased after infection)

was calculated and reported.

Heritability

Broad-sense heritability (H^2) was calculated using the formula $H^2 = V_G/V_P =$

$V_G/(V_E + V_G)$ as previous described [114, 115]. V_E and V_G are the variance explained by

the environmental and the genetic component respectively while V_P is the total

phenotypic variance for a given phenotype. Detailed explanation of the calculation can

be found here [116].

QTL analysis

QTL analysis was performed using R/qtl2 software [59]. This method accounts

for the complex population structure in CC strains. Briefly, genotype probabilities were

imputed from the QTL viewer [60]. Genome scans were performed on the transformed

phenotype using the scan1 function. The generated Logarithm of Odds (LOD) score is

the likelihood ratio comparing the hypothesis of a QTL at a given position versus that of

no QTL. We used the number of mice within a strain that survived at the end of each

day as the phenotype. The phenotype was randomly shuffled 999 times to establish

genome-wide significance, and LOD scores were calculated for each iteration using the

scan1perm function [117]. The 85[th] percentile of the scores was considered significant

for that phenotype. The genomic confidence interval was calculated by dropping the

LOD scores by 1.8 for each significant peak. Mouse Genome Informatics (MGI) was

used to find the genes and QTL features within each interval using mouse genome

version GRCm38 [118]. To further shortlist candidate genes, the founder strain

distribution pattern was queried against the CC variant database (V3 Version). The

Variant effect predictor [VEP] from the ensemble genome database was used to

calculate the impact score for the variants [119].

<u>RNA extraction and sequencing</u>

Total RNA was extracted from frozen tissues using Direct-zol RNA Miniprep plus

kit following the manufacturer's protocol (Zymo Research - R2073). The purity and

quantity of the extracted RNA were analyzed using RNA 6000 Nano LabChip Kit and

Bioanalyzer 2100 (Agilent CA, USA, 5067-1511). High-Quality RNA samples with RIN

number > 7.0 were used to construct the sequencing library. After total RNA was

extracted, mRNA was purified from total RNA (5μg) using Dynabeads Oligo (dT) with

two rounds of purification (Thermo Fisher, CA, USA). Following purification, the mRNA

was fragmented into short fragments using divalent cations under elevated temperature

(Magnesium RNA Fragmentation Module (NEB, cat. e6150, USA) under 94℃ 5-7min).

Then the cleaved RNA fragments were reverse-transcribed to create the cDNA by

SuperScript™ II Reverse Transcriptase (Invitrogen, cat. 1896649, USA), which were

next used to synthesize U-labeled second-stranded DNAs with E. coli DNA polymerase

I (NEB, cat.m0209, USA), RNase H (NEB, cat.m0297, USA) and dUTP Solution

(Thermo Fisher, cat. R0133, USA). An A-base was then added to the blunt ends of each

strand, preparing them for ligation to the indexed adapters. Each adapter contained a T-

base overhang for ligating the adapter to the A-tailed fragmented DNA. Dual-index

adapters were ligated to the fragments, and size selection was performed with

AMPureXP beads. After the heat-labile UDG enzyme (NEB, cat.m0280, USA) treatment

of the U-labeled second-stranded DNAs, the ligated products were amplified with PCR

by the following conditions: initial denaturation at 95℃ for 3 min; 8 cycles of

denaturation at 98℃ for 15 sec, annealing at 60℃ for 15 sec, and extension at 72℃ for

30 sec; and then final extension at 72℃ for 5 min. The average insert size for the final

cDNA libraries was 300±50 bp. At last, we performed the 2×150bp paired-end

sequencing (PE150) on an Illumina Novaseq™ 6000 following the vendor's

recommended protocol.

RNA sequencing data analysis

Reads were trimmed using trim galore (version 0.6.7) [120-123]. This removed

adapters, poly tails, more than 5% of unknown nucleotides, and low-quality reads

containing more than 20% of low-quality bases (Q-value <20). Both forward and reverse

hard trimmed at 100 base pairs. FastQC was used to verify data quality before and after

cleaning [123]. Cleaned reads were aligned and counted against the mouse reference

genome (GRCm39) using STAR (version 2.7.9a) aligner [124]. Downstream processing

of the data was performed using IDEP 1.0 [125, 126]. Gene counts were analyzed for

differentially expressed genes using DESeq2 [127]

Data availability

All of the data points collected on each mouse before and after the infection can

be found in the supplementary file (Table S2). Raw RNA sequencing files can be found

at SRA (PRJNA1002380).

Figure and Table descriptions

Main Figures

Fig 1: CC strains exhibit a diverse response to MRSA infection. After intravenous

infection with MRSA USA300 as described in the Materials and Methods section, we

show: A. survival time, B. percent weight change after infection, C. kidney colonization,

and D. Total organ colonization; horizontal dotted line represents inoculum dose. CC

strains are shown in the ascending order of survival. If survival was equal between two

or more strains, strains are arranged in descending order of total organ colonization (the

combined colonization of all organs collected). Dots represent individual mice; black

dots represent males; green dots represent females. The median and interquartile

range are shown for each strain.

Fig 2: Correlation of tissue colonization, histology scores and blood parameters

with survival. A. Heat map showing Spearman correlation 'R' values between survival,

weight change, organ colonization, and tissue damage scores. B. A stacked bar plot

showing the median tissue damage score for five organs (Kidney = green, spleen =

purple, liver = yellow, heart = blue, lung = orange). Pathology in each tissue was blindly

scored by a pathologist on a scale of 0 = no damage, to 4= severe damage. C. Mean

neutrophil levels in uninfected mice. Mann-Whitney tests were performed to determine

statistical significance (* = P < 0.05, ** = P < 0.01).

Fig 3: Sexual dimorphism in survival after MRSA infection. We present pooled data

for the four strains that exhibited sexual dimorphism in survival: A. survival time, B.

percent weight change after infection, C. colonization across organs, and D. pre-

infection monocyte and red blood cell counts (U – Uninfected). Each point represents an

individual mouse; filled circles represent males; triangles represent females. Each color

represents a CC strain as noted in the legend. Medians with 95% confidence interval

are shown for each strain. A, B, and D – Student's T-test and C

Fig 4. Genomic regions involved in survival after MRSA infection. LOD plots for

square root transformed survival phenotype (percentage of animals that survived at the

end of each day). A. Early susceptibility in MRSA infection (ESMI, day 2 post-infection)

peak on chromosome 18. B. Late survival in MRSA infection (LSMI, day 7 post-

infection) peak on chromosome 3. The dotted (Red – 95%, Blue – 90%, Green – 85%)

lines represent the significant LOD scores for 999 permutations.

Fig 5. Strongest candidate genes on chromosome 18 for early susceptibility. A.

Box plot showing the founder allele contribution at the highest marker on the peak. Each

dot represents a strain, lines represent minimum to maximum. B. Genes with high-

impact variants shortlisted using the founder effect pattern. C. Mean kidney mRNA

expression values after infection (log-transformed). The number of animals is given

within brackets. Lines represent standard error of the mean values. ANOVA with

multiple comparison correction using Tukey test was performed (**** = P < 0.0001).

Fig 6. Strongest candidate genes on chromosome 3 for late survival. A. Box plot showing the founder allele contribution at the highest marker on the peak. Each dot represents a strain, lines represent minimum to maximum. B. Genes with high-impact variants shortlisted using the founder effect pattern. C. Mean kidney mRNA expression values after infection (log-transformed). The number of animals sequenced for each founder allele is within the brackets. Lines represent standard error of the mean values. ANOVA with multiple comparison correction Tukey test was performed (**** = P <0.0001).

Supplementary Figures

Fig S1: Colonization in other organs. A. Liver colonization. B. spleen colonization, C. heart colonization. D. lung colonization. Strains are shown in ascending order of survival. Where survival was equal, strains are arranged in descending order of total organ colonization. Dots represent individual mice; black dots represent males; green dots represent females. The median and interquartile range are shown for each strain.

Fig S2. Representative images of histology scoring scheme. Tissues were sectioned and stained with H&E for scoring. Shows the 400X magnification of representative scoring for all the organs.

Fig S3: Time to deviation from the normal circadian pattern. Plot showing the number of animals that deviated from their normal pattern as identified by telemetry system (machine - dot) vs. Laboratory worker (human - squares)

Fig S4. Impact of sex on QTL regions. LOD plots for square root transformed survival phenotype excluding the four sexually dimorphic strains. A. Early susceptibility. B. Late

survival. The dotted (Red – 99%, Blue – 95%, Green – 90%) lines represent the

significant LOD scores for 999 permutations.

Fig S5. Circadian pattern for all mice involved in the study. Blue line represents

temperature, black line represents activity and red line represents the time of infection.

Fig S6. Temperature change after infection. Change in temperature for each mouse

after infection (for 48 hours). The data is grouped by CC strain and colored lines in each

plot represent the strain averages.

Supplementary Tables

Table S1: Mouse strains used in this study. Number of mice, origin, and location of

breeding for all the mice used in the study.

Table S2: Data associated with each mouse. All the pre-infection and post-infection

data associated with each mouse in the study.

Table S3: Tissue scoring matrix with descriptions. The final score per organ

corresponds to the highest score assigned to a single category.

Table S4: Heritability scores. Broad range heritability scores for various phenotypes.

Table S5: Founder contribution. Sheet Chr18 – Founder contribution at the highest

peak on chromosome 18 for ESMI. Sheet Chr3 – on chromosome 03 for LSMI.

Table S6: Shortlisted variants for survival peaks. Sheet Chr_18 – high impact

variants on chromosome 3 for ESMI. Sheet Chr_3 – high impact variants on

chromosome 3 for LSMI.

References

1. Lowy, F.D., *Staphylococcus aureus infections.* N Engl J Med, 1998. **339**(8): p. 520-32.
2. Krismer, B., et al., *The commensal lifestyle of Staphylococcus aureus and its interactions with the nasal microbiota.* Nat Rev Microbiol, 2017. **15**(11): p. 675-687.
3. Wertheim, H.F., et al., *The role of nasal carriage in Staphylococcus aureus infections.* The Lancet infectious diseases, 2005. **5**(12): p. 751-762.
4. Laux, C., A. Peschel, and B. Krismer, *Staphylococcus aureus Colonization of the Human Nose and Interaction with Other Microbiome Members.* Microbiol Spectr, 2019. **7**(2).
5. Tong, S.Y., et al., *Staphylococcus aureus infections: epidemiology, pathophysiology, clinical manifestations, and management.* Clin Microbiol Rev, 2015. **28**(3): p. 603-61.
6. Clarridge, J.E., 3rd, et al., *Impact of strain typing methods on assessment of relationship between paired nares and wound isolates of methicillin-resistant Staphylococcus aureus.* J Clin Microbiol, 2013. **51**(1): p. 224-31.
7. Howden, B.P., et al., *Staphylococcus aureus host interactions and adaptation.* Nature Reviews Microbiology, 2023.
8. Haag, A.F., J.R. Fitzgerald, and J.R. Penadés, *Staphylococcus aureus in Animals.* Microbiol Spectr, 2019. **7**(3).
9. Otto, M., *Community-associated MRSA: what makes them special?* Int J Med Microbiol, 2013. **303**(6-7): p. 324-30.
10. Lee, A.S., et al., *Methicillin-resistant Staphylococcus aureus.* Nat Rev Dis Primers, 2018. **4**: p. 18033.
11. Jevons, M.P., A.W. Coe, and M.T. Parker, *Methicillin resistance in staphylococci.* Lancet, 1963. **1**(7287): p. 904-7.
12. Turner, N.A., et al., *Methicillin-resistant Staphylococcus aureus: an overview of basic and clinical research.* Nat Rev Microbiol, 2019. **17**(4): p. 203-218.
13. Diekema, D.J., et al., *Continued emergence of USA300 methicillin-resistant Staphylococcus aureus in the United States: results from a nationwide surveillance study.* Infect Control Hosp Epidemiol, 2014. **35**(3): p. 285-92.
14. Kempker, R.R., et al., *Association of methicillin-resistant Staphylococcus aureus (MRSA) USA300 genotype with mortality in MRSA bacteremia.* J Infect, 2010. **61**(5): p. 372-81.
15. Kreisel, K.M., et al., *USA300 methicillin-resistant Staphylococcus aureus bacteremia and the risk of severe sepsis: is USA300 methicillin-resistant Staphylococcus aureus associated with more severe infections?* Diagn Microbiol Infect Dis, 2011. **70**(3): p. 285-90.
16. Parker, D., *Humanized Mouse Models of Staphylococcus aureus Infection.* Front Immunol, 2017. **8**: p. 512.
17. Kim, H.K., D. Missiakas, and O. Schneewind, *Mouse models for infectious diseases caused by Staphylococcus aureus.* J Immunol Methods, 2014. **410**: p. 88-99.

18. McAdow, M., et al., *Coagulases as determinants of protective immune responses against Staphylococcus aureus.* Infect Immun, 2012. **80**(10): p. 3389-98.

19. McAdow, M., et al., *Preventing Staphylococcus aureus sepsis through the inhibition of its agglutination in blood.* PLoS Pathog, 2011. **7**(10): p. e1002307.

20. Kuklin, N.A., et al., *A novel Staphylococcus aureus vaccine: iron surface determinant B induces rapid antibody responses in rhesus macaques and specific increased survival in a murine S. aureus sepsis model.* Infect Immun, 2006. **74**(4): p. 2215-23.

21. Cheng, A.G., et al., *Genetic requirements for Staphylococcus aureus abscess formation and persistence in host tissues.* FASEB J, 2009. **23**(10): p. 3393-404.

22. Cheng, A.G., et al., *Contribution of coagulases towards Staphylococcus aureus disease and protective immunity.* PLoS Pathog, 2010. **6**(8): p. e1001036.

23. Dyzenhaus, S., et al., *MRSA lineage USA300 isolated from bloodstream infections exhibit altered virulence regulation.* Cell Host Microbe, 2023. **31**(2): p. 228-242.e8.

24. Takeuchi, O., K. Hoshino, and S. Akira, *Cutting Edge: TLR2-Deficient and MyD88-Deficient Mice Are Highly Susceptible to Staphylococcus aureus Infection1.* The Journal of Immunology, 2000. **165**(10): p. 5392-5396.

25. Deshmukh, H.S., et al., *Critical role of NOD2 in regulating the immune response to Staphylococcus aureus.* Infect Immun, 2009. **77**(4): p. 1376-82.

26. von Kockritz-Blickwede, M., et al., *Immunological mechanisms underlying the genetic predisposition to severe Staphylococcus aureus infection in the mouse model.* Am J Pathol, 2008. **173**(6): p. 1657-68.

27. Yan, Q., et al., *Candidate genes on murine chromosome 8 are associated with susceptibility to Staphylococcus aureus infection in mice and are involved with Staphylococcus aureus septicemia in humans.* PLoS One, 2017. **12**(6): p. e0179033.

28. Yan, Q., et al., *Dusp3 and Psme3 are associated with murine susceptibility to Staphylococcus aureus infection and human sepsis.* PLoS Pathog, 2014. **10**(6): p. e1004149.

29. Ahn, S.H., et al., *Two genes on A/J chromosome 18 are associated with susceptibility to Staphylococcus aureus infection by combined microarray and QTL analyses.* PLoS Pathog, 2010. **6**(9): p. e1001088.

30. Montgomery, C.P., et al., *Comparison of virulence in community-associated methicillin-resistant Staphylococcus aureus pulsotypes USA300 and USA400 in a rat model of pneumonia.* J Infect Dis, 2008. **198**(4): p. 561-70.

31. Cheung, G.Y., et al., *Role of the accessory gene regulator agr in community-associated methicillin-resistant Staphylococcus aureus pathogenesis.* Infect Immun, 2011. **79**(5): p. 1927-35.

32. Li, M., et al., *Evolution of virulence in epidemic community-associated methicillin-resistant Staphylococcus aureus.* Proc Natl Acad Sci U S A, 2009. **106**(14): p. 5883-8.

33. Noll, K.E., M.T. Ferris, and M.T. Heise, *The Collaborative Cross: A Systems Genetics Resource for Studying Host-Pathogen Interactions.* Cell Host Microbe, 2019. **25**(4): p. 484-498.

34. Threadgill, D.W., et al., *The collaborative cross: a recombinant inbred mouse
population for the systems genetic era.* Ilar j, 2011. **52**(1): p. 24-31.
35. Churchill, G.A., et al., *The Collaborative Cross, a community resource for the
genetic analysis of complex traits.* Nature genetics, 2004. **36**(11): p. 1133.
36. Srivastava, A., et al., *Genomes of the Mouse Collaborative Cross.* Genetics,
2017. **206**(2): p. 537-556.
37. Shorter, J.R., et al., *Whole Genome Sequencing and Progress Toward Full
Inbreeding of the Mouse Collaborative Cross Population.* G3
Genes|Genomes|Genetics, 2019. **9**(5): p. 1303-1311.
38. Rasmussen, A.L., et al., *Host genetic diversity enables Ebola hemorrhagic fever
pathogenesis and resistance.* Science, 2014. **346**(6212): p. 987-991.
39. Graham, J.B., et al., *Genetic diversity in the collaborative cross model
recapitulates human West Nile virus disease outcomes.* MBio, 2015. **6**(3).
40. Ferris, M.T., et al., *Modeling host genetic regulation of influenza pathogenesis in
the collaborative cross.* PLoS Pathog, 2013. **9**(2): p. e1003196.
41. Xiong, H., et al., *Genomic profiling of collaborative cross founder mice infected
with respiratory viruses reveals novel transcripts and infection-related strain-
specific gene and isoform expression.* G3 (Bethesda), 2014. **4**(8): p. 1429-44.
42. Graham, J.B., et al., *Baseline T cell immune phenotypes predict virologic and
disease control upon SARS-CoV infection in Collaborative Cross mice.* PLoS
Pathog, 2021. **17**(1): p. e1009287.
43. Brinkmeyer-Langford, C.L., et al., *Host genetic background influences diverse
neurological responses to viral infection in mice.* Sci Rep, 2017. **7**(1): p. 12194.
44. Jensen, I.J., et al., *Novel Mouse Model of Murine Cytomegalovirus-Induced
Adaptive NK Cells.* Immunohorizons, 2022. **6**(1): p. 8-15.
45. Cartwright, H.N., et al., *Genetic diversity of collaborative cross mice enables
identification of novel rift valley fever virus encephalitis model.* PLoS Pathog,
2022. **18**(7): p. e1010649.
46. Smith, C.M., et al., *Tuberculosis Susceptibility and Vaccine Protection Are
Independently Controlled by Host Genotype.* mBio, 2016. **7**(5).
47. Smith, C.M., et al., *Host-pathogen genetic interactions underlie tuberculosis
susceptibility in genetically diverse mice.* Elife, 2022. **11**.
48. Vered, K., et al., *Susceptibility to Klebsiella pneumoaie infection in collaborative
cross mice is a complex trait controlled by at least three loci acting at different
time points.* BMC Genomics, 2014. **15**: p. 865.
49. Lorè, N.I., F.A. Iraqi, and A. Bragonzi, *Host genetic diversity influences the
severity of Pseudomonas aeruginosa pneumonia in the Collaborative Cross
mice.* BMC Genet, 2015. **16**: p. 106.
50. Rogovskyy, A.S., et al., *The First Immunocompetent Mouse Model of Strictly
Human Pathogen, Borrelia recurrentis.* Infect Immun, 2021. **89**(7): p. e0004821.
51. Scoggin, K., et al., *Genetic background influences survival of infections with
Salmonella enterica serovar Typhimurium in the Collaborative Cross.* PLoS
Genet, 2022. **18**(4): p. e1010075.
52. Scoggin, K., et al., *Elucidating Mechanisms of Tolerance to Salmonella
Typhimurium across Long-Term Infections Using the Collaborative Cross.* mBio,
2022. **13**(4): p. e0112022.

53. Alugupalli, K.R., et al., *Identification of collaborative cross mouse strains*
permissive to Salmonella enterica serovar Typhi infection. Sci Rep, 2023. **13**(1):
p. 393.
54. Zhang, J., et al., *Identification of new loci involved in the host susceptibility to*
Salmonella Typhimurium in collaborative cross mice. BMC Genomics, 2018.
19(1): p. 303.
55. Ayres, J.S. and D.S. Schneider, *Tolerance of infections.* Annu Rev Immunol,
2012. **30**: p. 271-94.
56. McCarville, J. and J. Ayres, *Disease tolerance: concept and mechanisms.*
Current opinion in immunology, 2018. **50**: p. 88-93.
57. Schneider, D.S. and J.S. Ayres, *Two ways to survive infection: what resistance*
and tolerance can teach us about treating infectious diseases. Nat Rev Immunol,
2008. **8**(11): p. 889-95.
58. Ngcobo, S., et al., *Pre-infection plasma cytokines and chemokines as predictors*
of HIV disease progression. Sci Rep, 2022. **12**(1): p. 2437.
59. Broman, K.W., et al., *R/qtl2: Software for Mapping Quantitative Trait Loci with*
High-Dimensional Data and Multiparent Populations. Genetics, 2019. **211**(2): p.
495-502.
60. Vincent, M., et al., *QTLViewer: an interactive webtool for genetic analysis in the*
Collaborative Cross and Diversity Outbred mouse populations. G3
Genes|Genomes|Genetics, 2022. **12**(8).
61. Keele, G.R., et al., *Determinants of QTL Mapping Power in the Realized*
Collaborative Cross. G3 Genes|Genomes|Genetics, 2019. **9**(5): p. 1707-1727.
62. Blake, J.A., et al., *Mouse Genome Database (MGD): Knowledgebase for mouse-*
human comparative biology. Nucleic Acids Res, 2021. **49**(D1): p. D981-d987.
63. Praggastis, M., et al., *A Murine Niemann-Pick C1 I1061T Knock-In Model*
Recapitulates the Pathological Features of the Most Prevalent Human Disease
Allele. 2015. **35**(21): p. 8091-8106.
64. Platt, N., et al., *Immune dysfunction in Niemann-Pick disease type C.* J
Neurochem, 2016. **136 Suppl 1**(Suppl Suppl 1): p. 74-80.
65. Busse, D.C., et al., *Interferon-Induced Protein 44 and Interferon-Induced Protein*
44-Like Restrict Replication of Respiratory Syncytial Virus. J Virol, 2020. **94**(18).
66. Dias, S.P., M.C. Brouwer, and D. van de Beek, *Sex and Gender Differences in*
Bacterial Infections. Infect Immun, 2022. **90**(10): p. e0028322.
67. Fish, E.N., *The X-files in immunity: sex-based differences predispose immune*
responses. Nat Rev Immunol, 2008. **8**(9): p. 737-44.
68. Harbarth, S., et al., *Evaluating the probability of previously unknown carriage of*
MRSA at hospital admission. The American journal of medicine, 2006. **119**(3): p.
275. e15-275. e23.
69. Laupland, K.B., et al., *The changing epidemiology of Staphylococcus aureus*
bloodstream infection: a multinational population-based surveillance study.
Clinical microbiology and infection, 2013. **19**(5): p. 465-471.
70. Kaasch, A.J., et al., *Staphylococcus aureus bloodstream infection: a pooled*
analysis of five prospective, observational studies. Journal of Infection, 2014.
68(3): p. 242-251.

71. Maclayton, D.O., et al., *Case-control study of the relationship between MRSA*
bacteremia with a vancomycin MIC of 2 µg/mL and risk factors, costs, and
outcomes in inpatients undergoing hemodialysis. Clinical therapeutics, 2006.
28(8): p. 1208-1216.
72. Mansur, N., et al., *Does sex affect 30-day mortality in Staphylococcus aureus*
bacteremia? Gender medicine, 2012. **9**(6): p. 463-470.
73. Smit, J., et al., *Gender differences in the outcome of community-acquired*
Staphylococcus aureus bacteraemia: a historical population-based cohort study.
Clinical Microbiology and Infection, 2017. **23**(1): p. 27-32.
74. Paulsen, J., et al., *Epidemiology and outcome of Staphylococcus aureus*
bloodstream infection and sepsis in a Norwegian county 1996–2011: an
observational study. BMC infectious diseases, 2015. **15**(1): p. 1-10.
75. Kubica, M., et al., *A potential new pathway for Staphylococcus aureus*
dissemination: the silent survival of S. aureus phagocytosed by human
monocyte-derived macrophages. PloS one, 2008. **3**(1): p. e1409.
76. Pidwill, G.R., et al., *The Role of Macrophages in Staphylococcus aureus*
Infection. Front Immunol, 2020. **11**: p. 620339.
77. Skaar, E.P., et al., *Iron-source preference of Staphylococcus aureus infections.*
Science, 2004. **305**(5690): p. 1626-8.
78. Croize, J., et al., *Improved identification of Staphylococcus aureus using a new*
agglutination test. Results of an international study. Apmis, 1993. **101**(6): p. 487-
91.
79. Wilkerson, M., et al., *Comparison of five agglutination tests for identification of*
Staphylococcus aureus. J Clin Microbiol, 1997. **35**(1): p. 148-51.
80. Shin, P.K., et al., *Characteristics of new Staphylococcus aureus-RBC adhesion*
mechanism independent of fibrinogen and IgG under hydrodynamic shear
conditions. Am J Physiol Cell Physiol, 2005. **289**(3): p. C727-34.
81. Gianquinto, E., et al., *Interaction of human hemoglobin and semi-hemoglobins*
with the Staphylococcus aureus hemophore IsdB: a kinetic and mechanistic
insight. Scientific Reports, 2019. **9**(1): p. 18629.
82. Spaan, A.N., et al., *Staphylococcus aureus Targets the Duffy Antigen Receptor*
for Chemokines (DARC) to Lyse Erythrocytes. Cell Host Microbe, 2015. **18**(3): p.
363-70.
83. Spaan, A.N., et al., *Neutrophils versus Staphylococcus aureus: a biological tug of*
war. Annu Rev Microbiol, 2013. **67**: p. 629-50.
84. Bouma, G., et al., *Recent advances in the understanding of genetic defects of*
neutrophil number and function. Br J Haematol, 2010. **151**(4): p. 312-26.
85. Andrews, T. and K.E. Sullivan, *Infections in patients with inherited defects in*
phagocytic function. Clin Microbiol Rev, 2003. **16**(4): p. 597-621.
86. Lakshman, R. and A. Finn, *Neutrophil disorders and their management.* J Clin
Pathol, 2001. **54**(1): p. 7-19.
87. Pfeffer, S.R., *NPC intracellular cholesterol transporter 1 (NPC1)-mediated*
cholesterol export from lysosomes. J Biol Chem, 2019. **294**(5): p. 1706-1709.
88. Pentchev, P.G., *Niemann–Pick C research from mouse to gene.* Biochimica et
Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids, 2004. **1685**(1): p.
3-7.

89. Hendrikx, T., et al., *Lysosomal cholesterol accumulation: driver on the road to inflammation during atherosclerosis and non-alcoholic steatohepatitis.* Obes Rev, 2014. **15**(5): p. 424-33.

90. Xu, X., et al., *Lysosomal cholesterol accumulation in macrophages leading to coronary atherosclerosis in CD38(-/-) mice.* J Cell Mol Med, 2016. **20**(6): p. 1001-13.

91. Carette, J.E., et al., *Ebola virus entry requires the cholesterol transporter Niemann–Pick C1.* Nature, 2011. **477**(7364): p. 340-343.

92. Morales-Tenorio, M., et al., *Potential pharmacological strategies targeting the Niemann-Pick C1 receptor and Ebola virus glycoprotein interaction.* European Journal of Medicinal Chemistry, 2021. **223**: p. 113654.

93. Houben, T., et al., *Hematopoietic Npc1 mutation shifts gut microbiota composition in Ldlr(-/-) mice on a high-fat, high-cholesterol diet.* Sci Rep, 2019. **9**(1): p. 14956.

94. Schwerd, T., et al., *Impaired antibacterial autophagy links granulomatous intestinal inflammation in Niemann-Pick disease type C1 and XIAP deficiency with NOD2 variants in Crohn's disease.* Gut, 2017. **66**(6): p. 1060-1073.

95. Prajsnar, T.K., et al., *The autophagic response to Staphylococcus aureus provides an intracellular niche in neutrophils.* Autophagy, 2021. **17**(4): p. 888-902.

96. Dai, Y., et al., *Heterogeneous Vancomycin-Intermediate Staphylococcus aureus Uses the VraSR Regulatory System to Modulate Autophagy for Increased Intracellular Survival in Macrophage-Like Cell Line RAW264.7.* Front Microbiol, 2019. **10**: p. 1222.

97. O'Keeffe, K.M., et al., *Manipulation of Autophagy in Phagocytes Facilitates Staphylococcus aureus Bloodstream Infection.* Infect Immun, 2015. **83**(9): p. 3445-57.

98. Jiang, H., et al., *IFI44L as a Forward Regulator Enhancing Host Antituberculosis Responses.* J Immunol Res, 2021. **2021**: p. 5599408.

99. DeDiego, M.L., L. Martinez-Sobrido, and D.J. Topham, *Novel Functions of IFI44L as a Feedback Regulator of Host Antiviral Responses.* J Virol, 2019. **93**(21).

100. Holden, M.T., et al., *Complete genomes of two clinical Staphylococcus aureus strains: evidence for the rapid evolution of virulence and drug resistance.* Proc Natl Acad Sci U S A, 2004. **101**(26): p. 9786-91.

101. Highlander, S.K., et al., *Subtle genetic changes enhance virulence of methicillin resistant and sensitive Staphylococcus aureus.* BMC Microbiol, 2007. **7**: p. 99.

102. Lee, H.S., et al., *Comparative Study of the Difference in Behavior of the Accessory Gene Regulator (Agr) in USA300 and USA400 Community-Associated Methicillin-Resistant Staphylococcus aureus (CA-MRSA).* J Microbiol Biotechnol, 2021. **31**(8): p. 1060-1068.

103. Jones, M.B., et al., *Genomic and transcriptomic differences in community acquired methicillin resistant Staphylococcus aureus USA300 and USA400 strains.* BMC Genomics, 2014. **15**: p. 1145.

104. Bor, D.H., et al., *Infective Endocarditis in the U.S., 1998–2009: A Nationwide Study.* PLOS ONE, 2013. **8**(3): p. e60033.

105. Federspiel, J.J., et al., *Increasing US rates of endocarditis with Staphylococcus aureus: 1999-2008.* Archives of internal medicine, 2012. **172**(4): p. 363-365.

106. Gibson, G.W., et al., *Development of a mouse model of induced Staphylococcus aureus infective endocarditis.* Comparative medicine, 2007. **57**(6): p. 563-569.

107. Liesenborghs, L., et al., *Staphylococcus aureus endocarditis: distinct mechanisms of bacterial adhesion to damaged and inflamed heart valves.* Eur Heart J, 2019. **40**(39): p. 3248-3259.

108. Wasi, F. and J. Shuter, *Primary bacterial infection of the myocardium.* Front Biosci, 2003. **8**: p. s228-31.

109. LeLeiko, R.M., D.J. Bower, and C.P. Larsen, *MRSA-associated bacterial myocarditis causing ruptured ventricle and tamponade.* Cardiology, 2008. **111**(3): p. 188-90.

110. Jariwala, P., et al., *Myocardial abscess secondary to staphylococcal septicemia: diagnosis with 3D echocardiography.* Indian Heart J, 2013. **65**(1): p. 124-5.

111. Lee, L.-N., et al., *Characteristics and local risk factors of community-acquired and health-care-associated Staphylococcus aureus pneumonia.* Scientific Reports, 2022. **12**(1): p. 18670.

112. Redi, D., et al., *Staphylococcus aureus vaccine preclinical and clinical development: current state of the art.* New Microbiol, 2018. **41**(3): p. 208-213.

113. Miller, L.S., et al., *Development of a vaccine against Staphylococcus aureus invasive infections: Evidence based on human immunity, genetics and bacterial evasion mechanisms.* FEMS Microbiol Rev, 2020. **44**(1): p. 123-153.

114. Collin, R., et al., *Common Heritable Immunological Variations Revealed in Genetically Diverse Inbred Mouse Strains of the Collaborative Cross.* The Journal of Immunology, 2019. **202**(3): p. 777-786.

115. Visscher, P.M., W.G. Hill, and N.R. Wray, *Heritability in the genomics era-- concepts and misconceptions.* Nat Rev Genet, 2008. **9**(4): p. 255-66.

116. Nagarajan, A., et al., *Using the collaborative cross to identify the role of host genetics in defining the murine gut microbiome.* Microbiome, 2023. **11**(1): p. 149.

117. Churchill, G.A. and R.W. Doerge, *Empirical threshold values for quantitative trait mapping.* Genetics, 1994. **138**(3): p. 963-71.

118. Bogue, M.A., et al., *Mouse Phenome Database: towards a more FAIR-compliant and TRUST-worthy data repository and tool suite for phenotypes and genotypes.* Nucleic Acids Res, 2023. **51**(D1): p. D1067-d1074.

119. Cunningham, F., et al., *Ensembl 2022.* Nucleic Acids Res, 2022. **50**(D1): p. D988-d995.

120. Krueger, F., *Taking appropriate QC measures for RRBS-type or other -Seq applications with Trim Galore!* Github, 2019.

121. Kechin, A., et al., *cutPrimers: A New Tool for Accurate Cutting of Primers from Reads of Targeted Next Generation Sequencing.* J Comput Biol, 2017. **24**(11): p. 1138-1143.

122. *FastQC.* 2015.

123. Andrews, S., *FastQC: a quality control tool for high throughput sequence data.* 2010, Babraham Bioinformatics, Babraham Institute, Cambridge, United Kingdom.

124. Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1): p. 15-21.

125.   Ge, S.X., E.W. Son, and R. Yao, *iDEP: an integrated web application for*
differential expression and pathway analysis of RNA-Seq data. BMC
Bioinformatics, 2018. **19**(1): p. 534.
126.   Ge, X., *iDEP Web Application for RNA-Seq Data Analysis.* Methods Mol Biol,
2021. **2284**: p. 417-443.
127.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and*
dispersion for RNA-seq data with DESeq2. Genome Biol, 2014. **15**(12): p. 550.

1064

Fig 1

Fig 2

Fig 3

Fig 4

A. Early Susceptibility

B. Late Survival

Fig 5

A.

C.

B.

GENE	SNP_ID	A/J	NZO	WSB	Cons
Npc1	rs30985329	A	A	T	missense
Lama3	rs51262081	C	C	G	missense
Cabyr	rs49718484	A	A	G	missense
Osbpl1a	rs48171054	T	T	C	missense
Psma8	rs31135735	A	A	G	missense
Kctd1	rs226935597	G	G	T	missense
Cdh2	rs243156099	C	C	T	splice region

Fig 6

A.

C.

B.

GENE	SNP_ID	A/J	CAST	NOD	NZO	WSB	Cons
Uox	rs31217970	T	C	C	C	C	synonymous
Adgrl4	rs30505737	T	A	A	A	A	splice region
Ifi44	rs48866215	A.	G	G	G	G	missense
Ifi44l	rs31149902	G	A	A	A	A	synonymous
Miga1	rs31134017	G	T	T	T	T	missense