Gene expression variation underlying tissue-specific responses to copper
stress in *Drosophila melanogaster*

Elizabeth R Everman[1,2*] and Stuart J Macdonald[1,3]

[1]1200 Sunnyside Ave, University of Kansas, Molecular Biosciences, Lawrence, KS 66045, USA
[2]730 Van Vleet Oval, University of Oklahoma, Biology, Norman, OK 73019, USA
[3]1200 Sunnyside Ave, University of Kansas, Center for Computational Biology, Lawrence, KS
66045, USA

* Correspondence:
Elizabeth R. Everman
e.everman@ou.edu

Keywords: heavy metals, gene expression, eQTL mapping, DSPR

Running head: Tissue-specific copper response

Abstract

Copper is one of a handful of biologically necessary heavy metals that is also a common
environmental pollutant. Under normal conditions, copper ions are required for many key
physiological processes. However, in excess, copper quickly results in cell and tissue damage that
can range in severity from temporary injury to permanent neurological damage. Because of its
biological relevance, and because many conserved copper-responsive genes also respond to
other non-essential heavy metal pollutants, copper resistance in *Drosophila melanogaster* is a
useful model system with which to investigate the genetic control of the response to heavy metal
stress. Because heavy metal toxicity has the potential to differently impact specific tissues, we
genetically characterized the control of the gene expression response to copper stress in a tissue-
specific manner in this study. We assessed the copper stress response in head and gut tissue of
96 inbred strains from the *Drosophila* Synthetic Population Resource (DSPR) using a combination
of differential expression analysis and expression quantitative trait locus (eQTL) mapping.
Differential expression analysis revealed clear patterns of tissue-specific expression, primarily
driven by a more pronounced gene expression response in gut tissue. eQTL mapping of gene
expression under control and copper conditions as well as for the change in gene expression
following copper exposure (copper response eQTL) revealed hundreds of genes with tissue-
specific local *cis*-eQTL and many distant *trans*-eQTL. eQTL associated with *MtnA*, *Mdr49*, *Mdr50*,
and *Sod3* exhibited genotype by environment effects on gene expression under copper stress,
illuminating several tissue- and treatment-specific patterns of gene expression control. Together,

our data build a nuanced description of the roles and interactions between allelic and expression variation in copper-responsive genes, provide valuable insight into the genomic architecture of susceptibility to metal toxicity, and highlight many candidate genes for future functional characterization.

Introduction

Many forms of stress resistance contribute to the overall health of the individual, both through immediate consequences brought on by direct exposure to the stressor, and indirectly by increasing the risk of experiencing deleterious consequences in the future. Heavy metals are one such type of stressor. Broadly, exposure to heavy metals results in oxidative stress due to the reactive state of metal ions (Stohs and Bagchi 1995; Ercal *et al.* 2001; Jomova *et al.* 2010). Acute exposure can cause gastrointestinal distress and vomiting as well as damage to intestinal lining (Taylor *et al.* 2020; Gerhardsson 2022). Chronic heavy metal exposure has been linked to neurodegenerative diseases in humans including Alzheimer's and Parkinson's Disease in adults (Bonilla-Ramirez *et al.* 2011; Chen *et al.* 2016b; Bagheri *et al.* 2018; Liddell and White 2018; Bisaglia and Bubacco 2020) and learning and behavioral disorders in developing individuals (Jomova *et al.* 2010; Blaurock-Busch *et al.* 2011; Hsueh *et al.* 2017; Lee *et al.* 2018). Additionally, exposure can increase morbidity associated with health conditions including multiple sclerosis and osteoporosis (Åkesson *et al.* 2006) as well as anemia (Turgut *et al.* 2007). Heavy metal exposure risk is often related to occupation (Castilla and Nealler 1978; Neuberger *et al.* 1990; World Health Organization *et al.* 1996; Khan *et al.* 2008; Karnchanawong and Limpiteeprakan 2009; He *et al.* 2015; Knoblauch *et al.* 2017), and like many diseases ((e.g., type 2 diabetes, Crohn's disease, heart disease, IBD (http://www.genome.gov/gwasstudies) (Hindorff *et al.* 2009; Plomin *et al.* 2009)), susceptibility to metal stress is a genetically complex trait (Zhou *et al.* 2016; Everman *et al.* 2021, 2023).

Quantitative trait locus (QTL) mapping has provided a powerful means of characterizing allelic variation that contributes to lead, cadmium, copper, and zinc resistance in several model systems including worms, flies, yeast, and plants (MacNair 1983; Jones *et al.* 2006; Bálint *et al.* 2007; Courbot *et al.* 2007; Ruden *et al.* 2009; Ehrenreich *et al.* 2012; Evans *et al.* 2018; Everman

et al. 2021). The *D. melanogaster* model system has been leveraged in several of these studies to examine the genetic architecture of metal stress response. It is an excellent model because all of the major genes that are involved in metal detoxification play similar roles in humans, allowing for the insight gained from work in *Drosophila* to have broader applications for understanding the impact of heavy metals on human health (Balamurugan *et al.* 2004; Egli *et al.* 2006a; b; Turski and Thiele 2007; Burke *et al.* 2008; Hatori and Lutsenko 2013; Calap-Quintana *et al.* 2017; Zhou *et al.* 2017). Genome-wide association mapping in flies revealed multiple SNPs that are associated with resistance to lead and cadmium toxicity (Zhou *et al.* 2016, 2017), and we previously demonstrated that resistance to copper toxicity is influenced by multiple QTL that included several conserved genes involved in copper metabolism and homeostasis (Everman *et al.* 2021). Collectively these studies have provided valuable insight into naturally occurring genetic variants that contribute to variation in resistance to several common heavy metal pollutants. However, most studies that examine the genetic control of heavy metal resistance have focused either on the whole-animal response or on single tissues. Furthermore, work over the last decade has clearly demonstrated that the majority of sites that influence trait variation are more likely to have impacts on gene regulation rather than to result in protein coding changes (Pickrell 2014; Zhang and Lupski 2015; reviewed in Boyle *et al.* 2017; Alsheikh *et al.* 2022). Therefore, an important opportunity remains to characterize the genetic control of the gene expression response to heavy metal stress and to determine the degree to which this control varies across tissues.

Transcriptomics studies have repeatedly demonstrated that exposure to heavy metal stressors increases expression of several gene families involved in metal detoxification, metabolism, and transport as well as oxidative stress response (González *et al.* 2008; Catalán *et al.* 2016; Calap-Quintana *et al.* 2017; Qu *et al.* 2018; Everman *et al.* 2021; Felmlee *et al.* 2022; Green *et al.* 2022). However, the gene expression response to metal stress is itself also subject to genetic variation in the control of the transcriptional response, i.e., different genotypes can vary in their response to the same exposure (Findley *et al.* 2021). The loci that contribute to variation in the gene expression response can be identified using expression QTL (eQTL) mapping (Jansen 2001; De Koning and Haley 2005; Gilad *et al.* 2008). By combining RNA sequencing and

eQTL mapping, we examined the tissue-specific genetic control of the gene expression response to the common metal pollutant copper using the *D. melanogaster* model system.

Although a broad oxidative stress response is expected across tissues in response to metal stress, copper is one of several heavy metals that have the potential to elicit different gene expression responses in neurological and gut tissue. This expected tissue-specific difference is in part due to the spatial distribution of specialized copper accumulation cells found in the gut of most animals including flies and humans (Calap-Quintana *et al.* 2017; Miguel-Aliaga *et al.* 2018). Genes that are responsible for maintaining normal homeostasis of the set of essential heavy metals (e.g., copper and zinc), and that play a role in the expulsion of toxic metals such as cadmium and lead, are primarily expressed in these specialized cells where they play an important protective role against excess metal ions (Calap-Quintana *et al.* 2017). Here we compared the gut response with that of head tissue due to the link between metal exposure and neurological disease in humans (Chen *et al.* 2016b), and for the opportunity to characterize differences in the tissue-specific response to oxidative stress using copper as a model metal.

Our study leverages the *Drosophila* Synthetic Population Resource (DSPR (King *et al.* 2012a)), a panel of multiparental, advanced intercross strains exhibiting dramatic variation in copper resistance that we have previously employed to map numerous QTL for resistance (Everman *et al.* 2021). Here we build on this study and use a subset of 96 strains to characterize the regulatory control of copper stress. Since the DSPR is composed of genetically stable inbred strains, it is especially well suited to eQTL mapping studies that quantify expression across tissues and treatments. We identified thousands of eQTL in both gut and head tissue using animals exposed to either control or copper-exposure treatments. Tissue-specific genetic control of gene expression was extensive, suggesting that regulatory variation has the potential to have profound, yet context-specific effects on susceptibility to heavy metal stress. Ultimately, our characterization of the genetic control of gene expression response to copper stress provides nuanced insight into the genetic control of heavy metal resistance and its potential health consequences.

1 Methods

2 Fly stocks

We used the *Drosophila* Synthetic Population Resource (DSPR), a large multiparental

mapping panel with >1500 recombinant inbred lines (RILs) (see King *et al.* 2012a, b for additional

details). In a previous study, we exposed 60 adult females from each of 1556 RILs (767 and 789

RILs from the A and B populations, respectively) to 50mM $CuSO_4$, expressing resistance as the

percentage of adults alive at 48 hours (Everman *et al.* 2021). For the present study, we selected

48 of the "resistant" RILs from the B population that were in the top 25% of the distribution of

adult copper resistance and 48 of the "sensitive" RILs from the bottom 25% of the distribution

(Figure S1).

12 Rearing and assay conditions

Flies were reared and assayed across multiple batches in the same incubator at 25°C and

50% humidity with a 12:12hr light:dark photoperiod. To maintain consistency with prior work,

experimental females for the present study were obtained from each of the 96 DSPR RILs in the

same manner used to measure adult copper resistance (Everman *et al.* 2021). Briefly, adults were

placed on cornmeal-molasses-yeast media and were allowed to oviposit for 2 days before being

discarded. Experimental females from the following generation were sorted over CO_2 into groups

of 20 and allowed to recover for 24 hours on fresh cornmeal-molasses-yeast media. Following

recovery, 3-5 day old females were transferred to Instant *Drosophila* Media (Carolina Biological

Supply Company 173200) hydrated with either water (control) or 50mM $CuSO_4$ (Copper(II)

sulfate, Sigma-Aldrich C1297) and exposed to treatment conditions for 8 hours, after which

tissues were harvested (see below). The 8-hour exposure period lasted from lights on (0800hrs)

to 4 hours before lights off. No flies died during the 8-hour exposure period.

Strains were assayed in a series of small batches to accommodate the time required for

tissue processing following the exposure period, with sensitive and resistant RILs evenly

distributed across each batch. Given the different processing techniques for heads and gut, we

obtained these tissues from different individuals of the same genotypes in different batches. The

average batch size for head collection was 25 DSPR RILs and average batch size for gut dissection

was 6 DSPR strains.

Tissue collection, RNA isolation, and sequencing library preparation

To obtain head tissue, we exposed 60 females per DSPR RIL to control and copper

conditions (three vials of 20 females per treatment) for 8 hours. At the end of the exposure

period, flies from each treatment and RIL were moved to a labeled screw-top 50mL tube, flash

frozen in liquid nitrogen, immediately vortexed to separate heads from bodies, and then stored

at -80°C for up to five days. We used a series of three-inch diameter stacking brass sieves – chilled

on dry ice for at least five minutes to prevent tissue thawing – to isolate the heads from each

sample (see Supplementary Methods). Heads for each RIL/treatment were dispensed into a

screw-top microcentrifuge tube containing 3-4 glass beads and held on dry ice. Subsequently, we

added 300μL TRIzol Reagent (Invitrogen, 15596018) to each sample and stored samples at -80°C

until RNA extraction.

To obtain gut tissue, we exposed 20 females per DSPR RIL to control and copper

conditions for 8 hours. Guts, including the fore-, mid-, and hindgut, were dissected from 10 live

individuals per RIL/treatment combination in 1X PBS and placed in screw-top microcentrifuge

tubes containing 3-4 glass beads and 300μL TRIzol. Samples were chilled on ice during dissection

and stored at -80°C until RNA extraction.

RNA was extracted from each of the 384 samples in batches of 30 samples using the

Direct-zol RNA Miniprep kit (Zymo Research, R2050). Batches consisted of samples from the same

tissue and were processed in the order individual samples were collected. RNA was eluted in

50μL water, and concentrations were determined via a NanoDrop spectrophotometer and then

standardized to 20ng/μL in 96-well plates. Because head and gut tissues were not collected

contemporaneously, two plates contained samples from heads and two contained samples from

gut tissue. The order of the samples from each RIL and treatment was largely consistent across

the head and gut sets of plates; all plates included both copper and control samples for a given

RIL and held an even representation of sensitive and resistant RILs (Table S1).

Half-reaction, unique dual indexed TruSeq Stranded mRNA sequencing libraries (Illumina,

20020595) were generated and sequenced by the University of Kansas Genome Sequencing Core.

Concentrations for every library were obtained using the Qubit dsDNA HS assay kit

(ThermoFisher, Q32854), and successful libraries were pooled based on concentration within

each of the four tissue-specific plates described above (Table S1). This resulted in two pools of

head libraries (one 94- and one 96-plex), and two pools of gut libraries (one 93- and one 96-plex).

Paired-end 37bp reads were obtained by sequencing each pool separately on NextSeq550 High-

Output flowcells. Raw read counts for the 94-plex Head, 93-plex Gut, and 96-plex Gut libraries

were 5.1 – 5.6 million PE37 read pairs per sample (Figure S2). The initial sequencing run of the

96-plex Head library pool under-clustered, so the pool was sequenced a second time on a

NextSeq550 Mid-Output flowcell. Assessment of the reads revealed no difference in our ability

to map reads from either run (Figure S3), so we combined reads across the two runs, resulting in

an average of 3.8 million PE37 read pairs per sample for this library pool (Figure S2).

Sequencing data preprocessing

Read alignment and gene filtering

Quality assessment, adapter removal and read trimming (retaining reads with a minimum

of 15 bp, Figure S4) were performed with fastp (v. 0.20.1) (Chen *et al.* 2018). Only paired reads

were retained. Filtered read pairs were aligned in a variant-aware manner to the *Drosophila*

reference genome (Release 6.33) using HISAT2 (v. 2.1.0) (Kim *et al.* 2019). SNP variants identified

in the DSPR founders

(http://wfitch.bio.uci.edu/~tdlong/SantaCruzTracks/DSPR_R6/dm6/variation/DSPR.r6.SNPs.vcf.

gz) were included in the HISAT2 genome index following instructions provided with the software.

Aligned reads were sorted with SAMtools (v. 1.9) (Li *et al.* 2009) and quantified with

featureCounts (v. 2.0.1) (Liao *et al.* 2014). Post alignment and quantification analyses were all

performed in R (v. 4.1.3) (R Core Team 2017). Average library size prior to additional filtering was

4.03 million reads. In R, genes with low expression were filtered out by first removing genes with

zero counts across all samples and then by removing genes with fewer than 10 counts in 47

samples (the number of samples unique to each of the three-way interaction categories, e.g. low

copper resistance head tissue samples exposed to copper stress (Chen *et al.* 2016a)). Following

filtering, 9842 genes were retained for downstream analyses.

Comparison between alignment pipelines

We also aligned fastp-filtered reads using the kallisto (v. 0.46.2 (Bray *et al.* 2016))

pseudoalignment pipeline and the Ensembl *Drosophila* transcriptome release 90. Average library

size was 3.96 million reads following kallisto alignment. We applied the same post-alignment

filtering parameters described above to the kallisto generated counts. Because kallisto alignment

generates output quantified by transcript, filtering resulted in retention of a slightly different set

of genes totaling 9913 genes, 9434 of which were shared with the filtered HISAT2 gene set. To

compare gene expression results from the HISAT2 and kallisto pipelines, we calculated gene-wise

correlations in expression following log2 transformation and quantile normalization across the

379 libraries. Sample and tissue-specific read counts for the 9434 genes retained in both the

kallisto and HISAT2/SAMtools/featureCounts pipeline were highly correlated (mean R = 98.2%,

Figure S5). Correlations for normalized gene counts within each tissue/resistance class/treatment

set of samples were also high (96% - 97%). The 267 genes with counts that were less than 90%

correlated between the two pipelines were not enriched for any functional or molecular

categories following gene ontology analysis (Flymine, (Lyne *et al.* 2007)), suggesting that

differences in gene expression counts were not due to systematic discrepancies between

pipelines. Given the high consistency between pipelines, all downstream differential expression

(DE) analyses and eQTL mapping presented below are derived from the

HISAT2/SAMtools/featureCounts pipeline.

Effect of tissue, treatment, and resistance class on gene expression

Gene expression was normalized using the weighted trimmed mean of M-values (mean

of log expression ratios (Robinson and Oshlack 2010)), and DE analysis was performed with limma

(Ritchie *et al.* 2015) by fitting the full-factorial linear model (Expression ~ Tissue * Treatment *

Resistance class + Sample Pooling). The "Sample Pooling" model term accounts for technical

variation due to plate, library pool and/or sequencing flowcell (these 3 factors cannot be

distinguished in our design). Genes with expression variation significantly influenced by each

term in the model were identified by fitting contrasts to the full model in limma with the contrasts.fit function. This step was followed by the eBayes function, which uses an empirical Bayes method to calculate log fold change in normalized gene expression and to determine significance between each group identified by the contrast (contrast order: expression in gut relative to head tissue, copper relative to control treated samples, sensitive relative to resistant strains). Significant DE was determined using adjusted P values to account for multiple comparisons (Benjamini-Hochberg multiple test correction, FDR threshold = 5%).

Tissue and treatment-specific eQTL analysis

We performed eQTL mapping for 6 separate datasets, testing different numbers of gene expression traits in each due to some genes having tissue- and/or treatment-specific expression (Head-Control: 9783, Head-Copper: 9842, Head-Response: 9830, Gut-Control: 9842, Gut-Copper: 9842, Gut-Response: 9841 genes). The "Response" to copper treatment was calculated as gene expression under copper conditions minus expression under control conditions for all paired samples. Positive Response values indicate genes with higher expression under copper conditions (copper treatment-induced), while negative Response values indicate genes that are repressed by copper treatment. Each of the six datasets was separately quantile normalized following log2 transformation. To preserve the direction of gene expression change in both Response datasets, we performed log2 transformation and quantile normalization on the absolute values of the copper-response expression and reassigned the sign following normalization.

We corrected expression measures for variation due to technical and environmental factors using principal components analysis (PCA; described in further detail below). Separately for each dataset, gene expression was corrected by regressing out the effects of PCs that explained more than 2% of the variance in gene expression, or that were correlated with known technical factors. eQTL mapping was performed on quantile normalized residual gene expression for each of the six datasets using R/qtl2 (Broman *et al.* 2019). eQTL mapping treats each gene expression measure as a trait, and we test for associations between genotype and variation in gene expression at each genome marker (every 10kb in the DSPR (King *et al.* 2012a)). For each gene, we regressed expression on additive probabilities indicating the likelihood that a particular

region of the genome was inherited from one of the 8 DSPR founders. Gene-specific significance

thresholds were assigned by permuting gene expression estimates among the DSPR strains 1000

times.

Given the modest sample size for each eQTL analysis (93-96 DSPR strains), we defined

QTL intervals using a 3-LOD drop, since a 2-LOD drop can give overly narrow intervals when fewer

RILs are assayed (King *et al.* 2012a). *cis* eQTL were defined as above-threshold peaks for which

the upper or lower boundary of the 3-LOD drop was within 1.5cM of the target gene, while peaks

outside this interval were classified as distant or *trans* eQTL. We note that broader peak intervals

result in more peaks being designated as *cis* eQTL and fewer peaks as *trans* eQTL (King *et al.* 2014;

King and Long 2017). Peaks were removed if they consisted of a single, above-threshold marker,

or if the peak position was outside the lower and upper peak boundary (such phenomena are

more common in experiments with limited sample size and are often found near telomere and

centromere regions where the impact of low power is exacerbated). eQTL peaks, estimated

founder effects, and percent variance were identified and calculated using custom code derived

from the DSPRqtl (King *et al.* 2012b) and R/qtl2 mapping programs.

We examined the six datasets for evidence of eQTL that were shared among datasets.

Comparisons were made between tissues within treatment (Head-Control vs Gut-Control, Head-

Copper vs. Gut-Copper, Head-Response vs. Gut-Response) and within tissue between treatments

(Heads-Control vs Heads-Copper and Gut-Control vs Gut-Copper). Our ability to identify shared

eQTL across multiple datasets is influenced by power to detect eQTL and the effect size of the

eQTL. In an attempt to account for increased uncertainty, eQTL were considered shared if the

peak positions were within 1.5cM of each other and/or if the peak intervals overlapped.

PCA correction of gene expression for eQTL analysis

A common strategy to reduce spurious gene-eQTL associations in high-dimensional data

is to use PCA to account for known and unknown technical or environmental factors (Leek and

Storey 2007; Pickrell *et al.* 2010; Gaffney *et al.* 2012; King *et al.* 2014). To determine the most

appropriate PCA correction for our data, we compared three versions of the eQTL mapping

analysis using 1) uncorrected data, 2) data corrected for PCs that explained more than 2% of the

variance and/or were correlated with tissue collection batch or sample pool (general linear model

test of the effect of technical variation on PC, uncorrected $\alpha = 0.05$) and 3) data corrected for PCs

that explained more than 1% of variance and/or were correlated with technical variation. The set

of PCs accounted for in each case are listed in Table S2. For brevity, we will simply refer to the

two PCA-corrected versions of the dataset as 2% and 1% corrected. Corrections were made by

regressing out PCs separately for each of the six expression datasets. Residuals were quantile

normalized once more and treated as gene expression values for each of the eQTL analyses and

for all downstream analyses.

We performed the three eQTL analyses (no correction, 2% correction, 1% correction) for

each of the six datasets as described above. In Head-Control, Head-Copper, Gut-Control, and Gut-

Copper eQTL analyses, failing to correct for technical variation via PCA resulted in many more

peaks far from the target gene compared to the 2% and 1% cutoffs. Because not correcting for

technical factors increases noise, it is likely the large number of distant peaks in the uncorrected

data includes many more false positives than true *trans* eQTL (Figure S6). The relationship

between peak number and data correction strategy was more pronounced for head samples,

likely due to the known technical effect of sequencing pool (Figure S6) that we account for in the

PCA-corrected versions. Head- and Gut-Response data were less sensitive to the PCA correction

(Figure S6, S7). Removing PCs explaining 1% of the variance further reduced the total number of

peaks in both head and gut analyses relative to the 2% variance correction (Figure S7). However,

this difference was slight, and overall levels of noise observed in individual LOD plots were

visually and statistically similar between the 2% and 1% data correction (average correlation in

genome-wide LOD scores between 1% and 2% corrected data for genes with eQTL ranged from

70% - 83%; Figure S8). To avoid removing true positives through use of a strict 1% cutoff, all

subsequent analyses/results employ the 2% PCA-corrected version of the dataset.

Identification of eQTL hotspots

Identification of eQTL hotspots in each of the six eQTL mapping datasets entailed two

initial steps. First, we used a chi-square goodness of fit test, assessing each dataset separately,

to determine if *trans* eQTL were overrepresented in any of the 2cM intervals distributed along

each chromosome arm. Second, because the chi-square test only indicates whether any interval

on the chromosome arm had an observed number of eQTL that deviated from expected number

of eQTL, we calculated the percent contribution of each interval to the overall chi-square value.

Percent contribution was calculated for each interval as

5
$$\% \: Contribution = \left(\mathrm{X}^2 \: value / \sum \mathrm{X}^2 \: value \right) \times 100$$

Intervals with a percent contribution value greater than 10% were considered potential hotspots.

We chose 10% as a permissive threshold for detecting hotspots based on a visual assessment of

percent contribution values as they varied across the chromosome arm. In most cases, our

approach highlighted one interval per chromosome arm with enrichment for *trans* eQTL.

We took a three-step approach to further characterize *trans* eQTL hotspots and to identify

potential regulatory candidate genes responsible for hotspots. Each step was carried out

independently for each hotspot. First, we created a composite variable using PCA to summarize

expression variation for all genes with a *trans* eQTL in that hotspot. We then used QTL mapping

in the same manner as described above to isolate QTL for this composite variable (PC1), and if a

peak was detected that overlapped the location of the hotspot, the hotspot was retained in

subsequent steps.

Second, we repeated QTL mapping of the composite variable, but now iteratively tested

whether each gene that fell within the hotspot interval had a mediating effect on the composite

QTL peak by including the expression level of each gene as a covariate in the QTL analysis. If

adding a gene expression covariate eliminated the composite variable QTL (or dramatically

reduces the LOD score at the peak), the gene was a candidate mediating gene of the *trans* eQTL

hotspot. The degree to which the LOD score was reduced by potential mediating genes varied

across all the hotspots, making it difficult to establish a consistent cutoff value. We therefore

assessed sets of mediating genes separately for each hotspot, and identified candidates that had

the most pronounced mediating effect at the composite peak position. If several genes had

similar mediating effects on the composite peak, we included them as candidates. This

permissive approach produced a list of genes that should be considered for future follow up

studies to further distinguish between true mediating genes versus those that have correlated

expression with mediating genes. Because the 2cM intervals used to initially identify hotspots

were created arbitrarily, we used the boundaries of the *trans* eQTL peak intervals to define the genomic region associated with the hotspot by using the most minimum and most maximum interval boundary for all *trans* eQTL in each hotspot. All the genes that fell within these expanded regions were included in this second step except for those genes that did not appear in our gene expression dataset. Some genes with very low levels of expression were removed from our expression datasets during filtering and so were not included in this analysis (see above). Covariate data used in each analysis corresponded to the dataset in which the hotspot was identified. For example, gene expression in gut tissue under copper conditions was used as covariate data to account for a composite variable derived for a hotspot detected in the Gut-Copper dataset.

Third, we further examined potential mediating genes by testing the correlation between estimated founder haplotype effects at any *cis* eQTL that happened to be associated with the potential mediating gene that fell within the hotspot interval. As in the second step, we used the *cis* eQTL that corresponded to the dataset in which the *trans* eQTL hotspot was detected.

Gene annotations and ontology analyses

We obtained annotation and ontology information for focal sets of genes highlighted by differential expression analysis, eQTL mapping, and assessment of eQTL hotspots using the *Drosophila melanogaster* annotation tool available from biomaRt (v. 2.50.3 (Durinck *et al.* 2005)) via Ensembl (Cunningham *et al.* 2022) and the orb.DM.eg.db R package (v. 3.14.0 (Carlson 2019)). Gene enrichment analyses were performed using the R package GOstats (v. 2.60.0 (Falcon and Gentleman 2007)), which uses hypergeometric tests for overrepresentation of GO terms with a correction for multiple tests.

Data availability

RNAseq reads are available from NCBI SRA BioProject PRJNA993605. All data including DE results, quantile-normalized expression data used for eQTL analyses, and code used to perform eQTL analyses are available from FigShare. Supplementary material contains Supplementary Methods that provide more detail on the isolation of head tissue as well as the alignment

1 pipeline, code to carry out eQTL mapping analysis, and all input and output data generated for

2 each analytical step.

Results

Effect of tissue, treatment, and resistance class on gene expression

A principal goal of our study was to characterize the gut- and head-specific gene

expression responses to copper exposure in genetically diverse copper-sensitive and copper-

resistant DSPR strains. Our differential expression model tested the three-way interaction

between tissue (gut, head), treatment (control, 8-hour exposure to 50mM $CuSO_4$), and DSPR

strain resistance class (resistant, sensitive). The primary model parameters that influenced

expression were tissue and treatment. Gene expression was strikingly distinct between gut and

head samples resulting in differential expression (DE) of 8915 of the 9842 genes tested (91% of

genes, 5% FDR; Figure 1A). Treatment alone influenced gene expression in 1156 genes (12%,

Figure 1B), while resistance class alone influenced expression of three genes: *asRNA:CR44107*,

CG18563 (involved in serine-type endopeptidase activity (Thurmond *et al.* 2019)), and *CG6023*.

Figure 1. Gene expression varied between tissues and treatments. Here we employ PCA to depict variation in gene expression in the top differentially-expressed genes with FDR < 0.05 and log fold change > 1 for four significant model terms. Each PCA was carried out with a different set of genes: A. 5031 genes, B. 59 genes, C. 113 genes, D. 341 genes. The effect of tissue on gene expression is shown in (A) but is also evident in all other plots. B. Gene expression shifted in both head and gut tissue in response to copper exposure. C. Gene expression shifted in response to copper treatment in both head and gut tissue. Shifts were typically in the same direction in both tissues with a slightly higher magnitude response observed for genes in gut tissue relative to head, leading to the interaction. D. Tissue and resistance class interacted to influence gene expression, although the effect of resistance class was slight. The significant interaction was driven by a larger magnitude difference in expression between sensitive and resistant strains in the gut tissue in response to copper. In all plots, triangle symbols indicate head samples, circles indicate gut samples. Log2, quantile-normalized gene expression data were corrected for the effect of sequencing pool prior to clustering analysis.

We also found that for 70% of genes, expression was influenced by an interaction

between tissue and treatment (Figure 1C). On average, this tissue x treatment interaction was

driven by a response to copper exposure that was in the same direction in head and gut tissue,

but that was more pronounced in gut tissue (Figure 1C). This pattern is consistent with the slightly

larger effect of treatment alone in gut tissue (Figure 1B) and might be expected because flies

were exposed to copper via ingestion in our assay. The effect of resistance class varied subtly

between gut and head tissue, influencing 32% of genes (Figure 1D), with the interaction driven

by a slightly greater effect of resistance class in gut tissue versus head tissue. The treatment and

resistance class interaction did not influence the expression of any gene tested, and the three-

way interaction between tissue, treatment, and resistance class influenced 13 genes (0.13%), of

which 8 had known or predicted functions. None of the genes influenced by the three-way

interaction were known copper response genes or had clear connections to metal toxicity

response. Information on all DE genes is available in the Supplementary data.

Differential expression of copper-related genes

Seven genes that have been previously linked to binding, metabolism, and detoxification

of copper were among the DE genes influenced by treatment (Figure 2A) and 28 genes in these

categories were influenced by the tissue by treatment interaction (Figure 2B). For several genes

that have been previously investigated in the context of copper toxicity, the change in expression

across treatments/tissues was in the expected direction. For instance, *Syx5* is a critical copper

homeostasis gene that is required for proper accumulation of copper ions under normal (copper-

scarce) conditions and is hypothesized to aid in proper localization of copper import proteins

(Norgate *et al.* 2010). Norgate *et al.* (2010) demonstrated reduced expression of *Syx5* increased

resistance to excess copper. Complementing this previous work, we found exposure to copper

stress significantly reduced expression of *Syx5* in both head and gut tissues (Figure 2A),

suggesting that downregulation of *Syx5* in the whole organism may result from copper stress.

Expression of the metallothionein family genes in response to copper stress was also

consistent with expectations. Copper exposure was expected to increase expression of the

metallothionein genes, which are involved in the sequestration of excess copper ions as a first

defense against copper toxicity (Egli *et al.* 2006a; Calap-Quintana *et al.* 2017). Further, expression

was expected to be higher in gut compared to head tissue because primary expression of

metallothioneins has been shown to occur in specialized copper accumulating cells that line the

midgut in flies (Calap-Quintana *et al.* 2017) As expected, we found that expression of four

metallothioneins (*MtnA*, *MtnB*, *MtnD*, and *MtnE*) was significantly increased in response to

copper exposure in both head and gut tissue but the increase in expression was more

pronounced in gut tissue (Figure 2).

Figure 2. Treatment and the interaction between treatment and tissue influenced expression of several genes previously shown to
be involved in detoxification, homeostasis, or binding of copper and other heavy metals. A. Heatmap of genes differentially

Three key copper transporters (*Ctr1A*, *Ctr1B*, and *ATP7*) were also among the genes

influenced by an interaction between tissue and treatment that followed expected expression

patterns based on previous reports. The ATP7 and Ctr1B copper transporters function primarily

in specialized copper-accumulating cells that line the intestine in mammals and the midgut in

flies (Zhou *et al.* 2003; Turski and Thiele 2007; Calap-Quintana *et al.* 2017), whereas the Ctr1A

copper transporter is ubiquitously expressed in both mammals and flies (Turski and Thiele 2007;

Calap-Quintana *et al.* 2017). Under control and copper conditions, we observed higher

expression of *ATP7* and *Ctr1B* in gut tissue relative to head tissue and higher expression of *Ctr1A*

in head tissue compared to gut tissue (Figure 2B). Our data are also consistent with previous

reports that *Ctr1A/B* copper importers are downregulated in *D. melanogaster* in response to

copper overexposure (Zhou *et al.* 2003; Calap-Quintana *et al.* 2017) as we observed a decrease

in expression of both *Ctr1A* and *Ctr1B* following copper exposure. The pattern of decreased

expression in response to copper was consistent across tissue but was more pronounced in head

tissue (*Ctr1A*) or gut tissue (*Ctr1B*) depending on the gene. Although *ATP7* expression under

control conditions followed tissue-specific expectations, we observed that exposure to copper

stress significantly increased *ATP7* expression, a pattern that was more pronounced in gut tissue

(Figure 2B). *ATP7* is a major copper exporter (Zhou *et al.* 2003), facilitating the transfer of copper

from copper accumulating cells in the gut to other tissues. Increased copper export under

stressful conditions may play an important role in the response to copper toxicity.

Expression of the copper chaperone *Atox1* was also differentially affected by treatment

and tissue. As proteins, Atox1 transports copper to ATP7 (Calap-Quintana *et al.* 2017; Kamiya *et*

al. 2018) and to the extracellular antioxidant enzyme SOD3 (Itoh *et al.* 2009; Kamiya *et al.* 2018).

Both *Atox1* and *Sod3* were more highly expressed in gut tissue than in head tissue (Figure 2B),

and exposure to copper stress led to a decrease in expression of both genes. Work presented by

Itoh *et al.* (2009) suggests that this shift in expression may be mechanistically linked because

Atox1 functions both to transfer copper to SOD3 and to regulate its expression as a copper-

dependent transcription factor (Itoh *et al.* 2009). Under normal conditions, null mutations in

Atox1 have been shown to lead to copper deficiency, and excess copper led to decreased Atox1

protein expression in wild-type flies (Hua *et al.* 2011).

To characterize the tissue-specific stress responses more broadly we used gene ontology

analysis of those genes showing significant tissue by treatment interactions in our differential

expression analysis. Of the 6362 genes with tissue- and treatment-specific patterns in expression,

778 could be categorized into one or more stress response categories (Table 1). As expected,

when examined on a gene-by-gene basis within each category we often saw different responses

to copper stress in head and gut tissue, indicating a range of tissue-by-treatment interactions

(Figure 3A). However, when the average response of genes in each GO category was considered,

we found that the overall expression response to copper treatment was often more pronounced

in gut tissue compared to head tissue (Figure 3B). This pattern was observed for regulation of

and response to stress (both cellular and general), response to endoplasmic reticulum stress, and

categories related to stress-activation of the MAPK and protein kinase signaling cascades (Figure

3B). In contrast, copper-induced shifts in expression of genes involved in the response to

oxidative stress was more pronounced in head tissue compared to gut tissue (Figure 3B).

Importantly, each of the oxidative stress response genes also contribute to each of the categories

that showed a more pronounced response in gut tissue. This implies that oxidative stress

response is also elevated in gut tissue, but it is one stress response pathway among many that

are activated in the gut, whereas in head tissue oxidative stress response is singularly

pronounced.

Table 1. Enrichment of stress-related GO categories (5% FDR) in genes with a tissue by treatment interaction.

Term	GO ID	N Genes in GO Category	N Genes (%) with sig Interaction	P Value
Response to stress	GO:0006950	1321	735 (56%)	$<2.47e^{-24}$
Regulation of response to stress	GO:0080134	346	218 (63%)	$<5.97e^{-15}$
Cellular response to stress	GO:0033554	646	370 (57%)	$<6.83e^{-15}$
Response to oxidative stress	GO:0006979	140	91 (65%)	$<5.83e^{-8}$
Regulation of metal ion transport	GO:0010959	57	40 (70%)	$<2.16e^{-5}$
Response to endoplasmic reticulum stress	GO:0034976	89	57 (64%)	$<3.27e^{-5}$
Stress-activated protein kinase signaling cascade	GO:0031098	88	56 (64%)	$<4.99e^{-5}$
Regulation of cellular response to stress	GO:0080135	131	78 (60%)	$<5.95e^{-5}$
Stress-activated MAPK cascade	GO:0051403	87	55 (63%)	$<7.58e^{-5}$
Cellular response to metal ion	GO:0071248	15	13 (87%)	$<5.68e^{-4}$

Figure 3. 778 genes with a significant tissue by treatment interaction effect could be classified by their involvement in stress response pathways. A. At the gene level, mean expression across DSPR strains under copper and control conditions in gut and head tissue demonstrated many distinct tissue and treatment specific patterns of expression response to copper. B. When expression was summarized at the level of GO category, we found that most genes that are involved in cellular response to stress, regulation of stress response, response to endoplasmic reticulum stress, and involvement in stress-activated MAPK and protein kinase signaling had a more pronounced response in gut tissue compared to head tissue. Conversely, genes involved in the response to oxidative stress had a more consistently pronounced response in head tissue compared to gut tissue. Data shown are mean expression response for all genes in each treatment and tissue combination +/- SE. In both plots, gold indicates gut samples; red indicates head samples.

Overall, we found pervasive differences in expression between tissues, with many genes

– including candidate metal responsive genes – showing an expression change in response to

copper treatment as well as variation in copper response among tissues. We previously reported

a minor effect of DSPR strain-specific copper resistance on the gene expression response to

copper (Everman *et al.* 2021). Classifying DSPR strains into copper resistant and susceptible

classes in the present study explained relatively little of the expression variation, and resistance

class influenced only a small number of genes, none of which are members of known metal

response pathways. Many of the key copper response genes highlighted above and in Figure 2

and Figure 3 have essentially equal expression in resistant and sensitive strains, suggesting that

the base level toxicity response to our copper treatment is similar between resistant and sensitive

strains assayed in this study.

Expression QTL mapping

Properties of mapped eQTL

eQTL mapping was executed for 6 datasets: Head-Copper, Head-Control, Head-Response,

Gut-Copper, Gut-Control, Gut-Response (see Methods for additional detail). In total over all

datasets, 4377 genes were associated with at least one eQTL and 98% of these 4377 genes were

differentially expressed due to one or more of the DE model terms and interactions (described

above). For most genes (83-98%; Table 2), we detected one eQTL (either *cis* or *trans*) per gene

per dataset (Figure S9). Genes with more than one eQTL per dataset included a multidrug

resistance gene *Mdr65* and two cytochrome p450 genes *Cyp6t3* and *Cyp12d1-d*, which have been

linked to response to insecticides (Daborn *et al.* 2007; Esteves *et al.* 2021). Two genes linked to

copper ion binding and homeostasis (*Mco1* and *CG6908* (Lang *et al.* 2012; Thurmond *et al.* 2019))

were also among the genes with multiple eQTL per dataset. However, no gene enrichment

related to metal detoxification or response was evident in genes with more than one eQTL peak

per dataset. Gene annotation and GO analysis for genes with multiple eQTL is available in the

Supplementary data.

Table 2. Summary statistics for eQTL mapping analyses.

Analysis	eQTL Type	N eQTL Peaks	N genes with eQTL	Total N unique genes with eQTL	% Genes with 1 eQTL peak	Mean Percent Variance (range)
Gut-Control	*cis*	1784	1779	1993	96%	43.1 (25.0-74.4)
Gut-Control	*trans*	295	286			33.7 (10.4-63.0)

Gut-Copper	*cis*	1677	1674	1860	97%	42.4 (22.0-76.0)
Gut-Copper	*trans*	248	237			33.4 (21.2-68.2)
Gut-Response	*cis*	139	138	316	98%	40.0 (25.5-72.5)
Gut-Response	*trans*	183	180			32.3 (16.6-41.9)
Head-Control	*cis*	1875	1869	2125	96%	43.6 (22.8-76.8)
Head-Control	*trans*	334	320			33.4 (18.3-66.4)
Head-Copper	*cis*	1879	1870	2152	97%	43.6 (23.9-78.1)
Head-Copper	*trans*	351	343			34.2 (20.5-67.4)
Head-Response	*cis*	34	34	197	83%	34.3 (21.3-64.3)
Head-Response	*trans*	235	167			30.0 (18.8-37.5)

For each of the Control and Copper datasets, *cis* eQTL outnumbered *trans* eQTL (Table 2,

Figure S7). In both Response datasets, we identified fewer eQTL overall but *trans* eQTL were more

common than *cis* eQTL (Table 2, Figure S7). This pattern is anticipated because the Response

datasets are based on the difference in gene expression between control and copper treatments,

which should eliminate genetic effects on expression that are consistent in both treatments.

Since the bulk of treatment-specific eQTL are *cis* eQTL, the Response data set is expected to have

fewer eQTL with local effects.

Across all datasets, the percent variance in gene expression explained by individual eQTL

ranged from 10.4% - 78.1% (Figure S10, Table 2). Consistent with previous work (Dixon *et al.*

2007; Emilsson *et al.* 2008; King *et al.* 2014; Albert *et al.* 2018; Keele *et al.* 2020), *cis* eQTL tended

to have higher percent variance estimates compared to *trans* eQTL in each of the six datasets

(Figure S10, Table 2), suggesting that *cis* eQTL tend to have a larger effect on transcriptional

variation compared to *trans* eQTL (Gibson and Weir 2005). However, percent variance estimates

are likely overestimated due to Beavis effects so should be assessed with care (Beavis 1995; King

and Long 2017).

Tissue specificity of mapped eQTL

Although most genes had only one eQTL per dataset (Table 2), 53.7% of genes had one or

more eQTL in at least two datasets. For instance, expression of the extracellular Copper/Zinc

superoxide dismutase *Sod3* was associated with a single eQTL per dataset but a similarly-located

Sod3 cis eQTL was detected in five of the six datasets (Figure 4A, B). For other genes, eQTL in

different datasets were clearly distinct. For example, expression of the metallothionein gene

MtnA was associated with a *trans* eQTL in the Head-Copper dataset and a *cis* eQTL in the Gut-

Copper dataset (Figure 4C, D). This implies that tissue-specific variants influence *MtnA* expression

under copper stress.

Figure 4. Many genes (53.7%) were associated with eQTL in multiple datasets, but these eQTL were often localized to distinct
intervals. A. and B. Expression of the gene Sod3 *was associated with* cis *eQTL in all datasets except Head-Response. C. and D. The*
gene MtnA *was associated with a* trans *eQTL on chromosome arm 2L in the Head-Copper dataset (C) and was associated with a*
cis eQTL *in the Gut-Control dataset (D). In all plots, the dashed horizontal line indicates the significance threshold based on*
permutation and the red triangle point indicates the position of the gene. LOD curves are colored based on treatment: gray =
Control, blue = Copper, green = Response.

To determine whether a similarly localized eQTL was detected in multiple datasets, we

examined eQTL overlap on a per gene basis between tissues within treatment (Head-Control vs

Gut-Control, Head-Copper vs. Gut-Copper, Head-Response vs. Gut-Response) and within tissue

between treatments (Heads-Control vs Heads-Copper and Gut-Control vs Gut-Copper). eQTL

were classified as overlapping if the peak positions were within 1.5cM or if the peak intervals

overlapped (see Methods). Overall, *cis* eQTL were more likely than *trans* eQTL to be detected in

multiple datasets regardless of which datasets were compared (Figure S11). The number of eQTL

that were detected in more than one dataset was highest in within-tissue comparisons (Figure

S11A, B), whereas more eQTL were distinct between tissues within treatment (Figure S11C, D).

eQTL detected in the Response datasets were most distinct with only 7 *cis* eQTL and no *trans*

eQTL shared between Head- and Gut-Response datasets (Figure S11E). Together, these results

are consistent with our DE analysis, which identified tissue as having the greatest impact on

expression variation (Figure 1).

Within each tissue the majority of *cis* Response eQTL (64-85%) were also detected in the

Control and Copper datasets while the majority of *trans* Response eQTL were unique to the

Response dataset (Figure S11F-I). The overlapping *cis* Response eQTL were typically detected as

cis eQTL in both Control and Copper datasets, suggesting that these *cis* eQTL are either associated

with different variants in the Control and Copper datasets or that the variant has different

additive or magnitude effects on the expression of genes under control and copper conditions.

Thorough tests of this hypothesis are beyond the scope of our data; however, correlations

between Response and Control or Copper founder haplotype effects at each eQTL suggest both

patterns may contribute (Figure S12).

Overall, our eQTL mapping results suggest that regulatory variation influences gene

expression in a tissue-specific manner for 2581 genes with distinct eQTL (detected in only one

dataset) that were involved in a broad range of processes spanning response to stimulus to

cellular metabolism (Figure S13). In contrast, regulatory variants influenced expression of 705

genes with shared eQTL that were detected in both head and gut tissues following copper

exposure (Figure S13). These 705 genes were enriched for a smaller number of GO categories

including metal response, detoxification, oxidative stress response, and similar stress response

pathways (Figure S13). This observation suggests that the control of some major components of

metal stress response is potentially consistent across tissues. This said, our data provide some

evidence that founder haplotype effects at shared eQTL are not always consistent, suggesting

that physically-overlapping eQTL do not necessarily represent identical genetic effects. For

example, founder haplotype effects at shared *cis* eQTL in Head and Gut tissue associated with

Mdr49, *Mdr65*, and *Sid* (involved in insecticide resistance (Sun *et al.* 2017; Denecke *et al.* 2017)

and oxidative stress response (Seong *et al.* 2014)) were negatively correlated, implying the

regulatory variant may have opposite effects on expression of these genes in head and gut tissue,

or that the eQTL actually represent the effects of distinct variants (Figure S14).

We also assessed tissue-specific eQTL patterns using our differential expression analysis

results as a guide. Of the 778 stress-related genes (defined using GO terms available from

FlyBase) with a tissue by treatment interaction (Figure 3), 349 were associated with a mapped

eQTL. Of the 349 genes, 269 had eQTL that were detected in up to three of the four main datasets

(for instance, detected in head tissue under copper and control conditions but not in either gut

dataset), indicating that most of the genes that were highlighted by DE analysis also had eQTL

that followed a tissue- and treatment-specific pattern. Combining our DE and eQTL mapping

results provides an additional layer of insight into how gene expression varies across tissue and

treatment. For the 269 genes with differential expression and eQTL, allelic variation that

influences gene expression variation likely contributes to overall tissue-specific responses to

copper stress.

Treatment specificity of eQTL

We directly compared the Control and Copper eQTL mapping results within each tissue

to identify genetic variants with treatment-dependent regulatory effects on gene expression as

well as eQTL with regulatory effects on expression that are consistent over treatments. Overall,

we found that eQTL detected under both control and copper conditions were common in head

and gut tissue (Figure S11A, B). These overlapping eQTL were associated with expression of genes

that included those related to metal response, toxicity response, and copper stress, which

resulted in enrichment of gene ontology terms related to stress responses that are consistent

with the effects of heavy metal stress (Figure S15). Interestingly, while copper treatment changed

the level of expression of metal response genes (based on GO annotation from FlyBase) with

eQTL shared by both treatments (Figure 5A, D), the estimated founder haplotype effects at eQTL

were generally consistent under both copper and control conditions (Figure 5B, E), i.e., the

genetic effects on expression are largely preserved over treatment. Despite this, eQTL had larger

effects on gene expression under copper conditions relative to control conditions in both tissues (Head Tissue: $F_{3,1496}$ = 529.2, P < 0.0001, Figure 5C; Gut Tissue: $F_{3,1258}$ = 311.2, P < 0.0001, Figure 5F), and this pattern was most pronounced for eQTL associated with metal response genes in gut tissue (Head Tissue, metal responsive vs non-metal responsive interaction: P = 0.4, Figure 5C; Gut Tissue, metal responsive vs non-metal responsive: P < 0.05, Figure 5F). Therefore, when genes have eQTL in both copper and control treatments, the eQTL contributes to expression variation in each environment in a similar way, but with the magnitude of the eQTL effect varying over tissue.

Figure 5. eQTL shared between control and copper conditions are influenced by regulatory variants in a consistent manner across treatments. A. and D. Genes previously linked to copper response, toxicity response, and oxidative stress (metal response genes) shifted in expression level in response to copper treatment in head (A) and gut (D) tissue. B. and E. Founder haplotype estimated effects for shared eQTL were similar under copper and control conditions for both metal response (blue) and non-metal response genes (grey) in head (B) and gut (E) tissues. C. and F. Percent variance explained by shared eQTL estimated in control and copper conditions was positively correlated but indicated larger effects on expression under copper conditions (C, Head Tissue: $F_{3,1496}$ = 529, r^2 = 51%, P < 0.0001; F, Gut Tissue: $F_{3,1258}$ = 312.2, r^2 = 43%, P < 0.0001). We found an interaction between percent variance estimates under control and copper conditions for metal response vs non-metal response gene in gut tissue, suggesting that the additive shift in effect size was greater for metal response gene expression variation under copper conditions. In both C and F the dashed line indicates the 1:1 slope to illustrate the reduced slopes of the regressions. Solid lines indicate the best fit using glm estimation and shading indicates the 95% confidence interval of the regression.

eQTL detected in just one of the two treatment conditions per tissue represent potential

loci with genotype by environment effects on gene expression. Compared to shared eQTL, there

were fewer distinct eQTL between the copper and control treatments for head and gut tissue

(Figure S11A, B). In head tissue, we found copper-specific eQTL associated with genes enriched

for functions related to ion, cation, and metal ion transport (Figure S16A). Copper-specific head

eQTL were associated with several copper-related genes including *MtnA*, *Mco1*, *Mco4*, *Loxl2*, and

CG17996 (Thurmond *et al.* 2019; The UniProt Consortium *et al.* 2021). Of note are *MtnA* and

Mco1. *MtnA* encodes one of five metal scavenger proteins that reduce the levels of free copper

ions available in the cell (Calap-Quintana *et al.* 2017). Exposure to copper increases *MtnA*

expression in head tissue (Figure 2B), and the *trans* eQTL associated with *MtnA* (Figure 4C) may

indicate a distant regulatory element that influences expression under copper stress. *Mco1* uses

copper as a cofactor for iron metabolism (Lang *et al.* 2012) and was associated with a *cis* and a

trans eQTL. The *trans* eQTL associated with *Mco1* falls near the metal-dependent transcription

binding factor *MTF-1*, which is one of the primary transcription factors involved in metal stress

response (Egli *et al.* 2003; Balamurugan *et al.* 2004). Control-specific eQTL in head tissue

highlighted a set of genes primarily enriched for functions related to broad categories including

cellular process and response to stimulus (Figure S16B). Despite the lack of enrichment for metal-

related GO terms, control-specific eQTL were associated with several genes involved in copper

ion homeostasis including *Mvl*, *CG10505*, *CG7194*, Grx1, and *Ctr1B* (Thurmond *et al.* 2019; The

UniProt Consortium *et al.* 2021). Maintaining proper copper homeostasis is critical for normal

physiological function, and these eQTL with effects on expression of copper homeostasis genes

may provide insight into regulation of copper transport under non-stressful conditions.

In gut tissue, we found enrichment for GO categories related to lipid metabolism,

localization, and homeostasis among genes with copper-specific eQTL (Figure S17A) and

enrichment for GO categories related to localization, transport, and regulation among genes with

control-specific eQTL (Figure S17B). Two copper-related genes (*olf413* and *CG6908*) had eQTL

under copper conditions in gut tissue (Thurmond *et al.* 2019; The UniProt Consortium *et al.* 2021).

Both eQTL were far from the gene positions (*trans*) and neither were near known metal-

responsive transcription factors. Copper-specific eQTL also implicated several genes from the

glutathione s transferase and cytochrome p450 families of genes, which are involved in the response to toxic substances (Kim and Yim 2013; Esteves *et al.* 2021), and we observed a copper-specific eQTL for the metallothionein gene *MtnA*. Similar to head tissue, several copper homeostasis genes had eQTL that were only detected under control conditions including *Ccs*, *CG11590*, *CG11825*, and *Atox1* (Culotta *et al.* 1997; Hatori and Lutsenko 2013), suggesting that allelic variation in the DSPR contributes to variation in regulation of expression of these genes under non-stressful conditions.

Treatment-dependent eQTL were also identified using the Head- and Gut-Response datasets. Response eQTL reveal loci that impact the difference in gene expression between copper and control conditions in a genotype-specific manner and are thus inherently genotype by environment eQTL. Response eQTL detected in both Gut- and Head-Response datasets were associated with several potential candidate genes that may play a role in the response to copper toxicity. GO analysis of genes with Gut-Response eQTL highlighted categories related to detoxification pathways and response to toxins including glutathione s transferase family genes (glutathione metabolic process; hypergeometric test for overrepresentation of GO terms, adj p < 0.001; Figure 6) and cytochrome p450 genes (response to toxic substances and insecticides, adj p < 0.001; Figure 6) (Kim and Yim 2013; Esteves *et al.* 2021). We also detected Response eQTL for two ABC transporter multidrug resistance protein genes (*Mdr49* and *Mdr69*) as well as two genes that are known to play a role in copper metabolism or binding (*Sod3* and *Tbh*) (Itoh *et al.* 2009; The UniProt Consortium *et al.* 2021). More generally, 50 out of 316 annotated genes with Response eQTL in the Gut-Response dataset are known or predicted to be involved in metal ion binding (Thurmond *et al.* 2019; The UniProt Consortium *et al.* 2021).

Head-Response eQTL included fewer potential candidate genes. Top GO enrichment categories included proteolysis and metabolic process (adj p < 0.0001) followed by response to insecticide (adj p < 0.001; Figure 6). We detected Head-Response eQTL for ABC transporter gene *Mdr50* as well as a different *trans* eQTL associated with *Mco1* that was distinct from those identified by comparing the copper and control-specific head eQTL. Of the 197 annotated Response eQTL genes, 26 play or are predicted to play a role in metal ion binding. There were 19 eQTL genes that were shared between the Head- and Gut-Response datasets, which included

several cytochrome p450 family genes. The shared set of Response eQTL were associated with

expression of genes that contribute to response to insecticide (adj p < 0.0001), response to toxic

substance (adj p < 0.001) and response to DDT (adj p < 0.001) (Figure 6).

Figure 6. Enrichment of GO categories for Response eQTL genes by tissue (head tissue = 197 genes; gut tissue = 316) and for the
set of eQTL genes that were shared between tissues (Shared-Response, 19 genes).

Identification of eQTL hotspots

Using a chi-square goodness of fit test, we identified between two and seven

chromosomal regions that encompassed more *trans* eQTL peaks than expected by chance per

dataset; these are potential *trans* eQTL hotspots (Table 3). Head- and Gut-Response datasets had

the most hotspots, and overall, more hotspots were detected in head tissue compared to gut

tissue datasets (Table 3). Several hotspots flanked the centromeric regions of chromosomes 2

and 3 (Figure 7) and were detected in multiple tissues and datasets (Table 3; Figure 7). This is

likely because we identified hotspot locations using genetic distance, and since crossing over is

low and gene density is higher in centromeric regions, the chance of detecting a hotspot at

centromeres is likely inflated.

Table 3. Chi-square analysis of trans eQTL distribution along chromosome arms and characteristics of hotspots.

Dataset	Chr	P Value	Hotspot	Hotspot Region* (cM)	N *trans* eQTL	N Genes/ Hotspot**	Mean Cor***	Possible Mediating Genes
Head-Control	2L	4.24e-07	H1	53.3-58.5	15	669	0.16	*CG31793, CG10366*
	2R	9.31e-07	H2	53.7-61.0	15	737	0.19	*CG14464, Cyp6a2*
	3L	6.42e-08	H3	42.9-46.4	14	835	0.16	*CG4962, CG4306*
	3R	7.91e-46	H4	45.0-48.8	43	813	0.15	*lncRNA:CR44317, CG2846*
Head-Copper	2L	3.95e-08	H5	18.2-22.9	12	175	0.56	*homer, Galt*
	2R	1.04e-13	H6	53.5-60.9	18	726	0.17	*TpnC41C, Tsp42Eg*
	3R	2.07e-53	H7	45.9-48.8	47	828	0.16	*lncRNA:CR44317, CG2846*
Head-Response	2L	0.0003	H8	3.42-6.68	8	184	0.48	*Npc2a, Auxd1*
	2R	1.42e-05	H9	84.5-88.7	9	256	0.26	*Zda, Eip55E*
			H10	86.7-90.2	8	281	0.49	*Fak*
	3L	2.27e-07	H11	10.8-18.0	10	294	0.54	*Jon65Aiv*
	3R	1.70e-35	H12	46.0-48.6	19	765	0.65	*CG14606, CG18748*
			H13	77.0-79.0	32	61	0.78	*N/A*
	X	1.44e-27	H14	14.1-20.9	22	206	0.72	*CG32751*
Gut-Control	2L	3.93e-05	G1	53.3-57.8	11	657	0.20	*Tsp39D*
	3R	3.17e-21	G2	46.0-48.8	25	813	0.15	*stck*
Gut-Copper	2L	0.002	G3	51.3-55.7	8	1046	0.13	*Ncm, CG15170, lectin-37Da*
	2R	0.002	G4	54.0-58.2	8	486	0.33	*CG14464, Ars2, Tsp42Eq, TspEr, Cyp6a2*
	3R	2.03e-24	G5	45.6-48.9	28	831	0.15	*DhpD, CH42564, Syt4, alpha-Est5*
	X	0.26	G6	0.006-1.32	5	202	0.19	*CG11384*
			G7	42.5-47.8	5	155	0.13	*inaE*
Gut-Response	2L	0.03	G8	51.0-53.7	6	580	0.10	*Ku80, CG10341*
	2R	0.0009	G9	82.4-85.8	8	251	0.14	*CG5742*
	3L	8.46e-07	G10	-0.004-2.5	8	353	0.15	*LysB, LysC, LysD*
			G11	42.9-47.0	8	1108	0.11	*Jon74E*
	3R	5.97e-05	G12	50.3-53.1	10	369	0.13	*CG12224, fabp Dip-C*

*Defined by the most minimum and most maximum boundaries of the *trans* eQTL that compose each hotspot
**Number of genes that fell within the boundary of the hotspot region
***Mean absolute pairwise correlations of gene expression (or response) values for *trans* eQTL genes corresponding to a given hotspot

If all genes with a *trans* eQTL in a given hotspot are co-regulated by a genetic element in the hotspot region, we anticipate that the expression of these genes (or their expression response to copper treatment) will be correlated to some extent. Mean absolute pairwise

correlations of gene expression or gene response values for *trans* eQTL genes corresponding to

a given hotspot ranged from 6-78% in Head datasets and from 10-60% in Gut datasets (Figure

S18, Figure S19; Table 3). We performed PCA on the matrix of per-RIL expression or response

measures for all the genes with a *trans* eQTL in each hotspot, with PC1 representing a composite

variable of the shared expression (or response) variation of genes with *trans* eQTL in that hotspot.

All composite hotspot variables mapped a QTL at the location of the hotspot, and genes with high

pairwise correlations (Figure S18, Figure S19) had consistent loadings on PC1 in each hotspot

(Figure S20).

To identify genes that may modulate expression of the set of genes with *trans* eQTL at a

hotspot, we used a mediation analysis approach that treated the expression level of each gene

that was encompassed by a hotspot interval (Table 3) as a covariate in QTL analysis of the

composite hotspot variable (see Methods). Including a true mediator as a covariate in the analysis

is expected to eliminate the presence of the composite variable QTL, resulting in a drop in the

LOD score at the peak position. With this approach we were able to identify between one and

five genes that had complete or partial mediating effects on the composite variable for all but

one hotspot (Figure S21, Figure S22; Table 3).

Our mediation analysis revealed three candidate genes that completely accounted for the

H11 composite hotspot QTL (Figure 8A). H11 is composed of 10 *trans* eQTL peaks that were

detected in the Head-Response dataset (Table 2). The three candidate mediating genes

(*Jon65Aiv*, *Jon65Aiii*, and *yip7*) are serine proteases expressed in the adult head and are involved

in proteolytic activity and protein metabolism (Kumar *et al.* 2021) (Figure 8A). We did not identify

any eQTL associated with *Jon65Aiii* or *yip7* but did detect one Head-Response *cis* eQTL near

Jon65Aiv and compared the estimated haplotype effects at this QTL with those at the H11

composite peak. Estimated founder haplotype effects were significantly negatively correlated (r

= -99%, p < 0.001, Figure 8B), suggesting that the haplotypes at this location have similar effects

on the response of *Jon65Aiv* to copper and on the response at genes that have a response *trans*

eQTL at the H11 hotspot. *Jon65Aiv* was recently identified as a novel candidate gene that may

contribute to the gene expression response to copper stress in *D. melanogaster* strains derived

from wild populations (Green *et al.* 2022). Copper exposure in wild-collected flies from that study

resulted in decreased expression of *Jon65Aiv*; we did not observe a significant change in expression of any of the three H11 candidate mediating genes in response to copper exposure.

We leveraged the STRING database to examine potential interactions between candidate mediating genes and genes with *trans* eQTL at the H11 hotspot (Szklarczyk *et al.* 2023). Using information gathered on predicted and known protein interactions and gene co-expression, STRING analysis showed that *Jon65Aiv*, *Jon65Aiii*, and *yip7* are directly or indirectly co-expressed with five of the genes with *trans* eQTL at the H11 hotspot, and several of the H11 *trans* eQTL genes have serine-type endopeptidase activity (GO enrichment analysis; p < 0.001), protein catalytic activity (p < 0.001), or hydrolase activity (p < 0.001) (Figure 8C). Together, these results suggest that *Jon65Aiv*, *Jon65Aiii*, and *yip7* are not regulatory mediating genes but rather in a shared pathway with the other H11 *trans* eQTL genes.

Figure 8. Three genes had complete mediating effects on the composite eQTL peak for hotspot H11. A. Including Jon65Aiv *(bottom purple line),* Jon65Aiii*, and* yip7 *(top purple lines) individually as covariates accounted for the composite QTL peak for H11 (black line). No other gene encompassed by the H11 composite peak had a strong mediating effect on the composite peak (green lines). B. Estimated founder haplotype effects at the composite QTL peak and at a cis eQTL associated with* Jon65Aiv *were significantly negatively correlated (Pearson r = -99%; p < 0.001). C. STRING analysis of the three candidate mediating genes and genes with a* trans *eQTL peak at the hotspot suggests that the candidate mediating genes are co-expressed with* trans *eQTL genes rather than having a regulatory role. Black edges indicate co-expression, green edges indicate similarity through textmining, light blue edges indicate protein homology, and dark blue edges indicate gene co-occurrence.*

We also identified a serine protease as a candidate gene with a partial mediating effect on the Gut-Response G11 *trans* eQTL hotspot (Figure 9A). *Jon74E* response to copper was associated with a *cis* eQTL, and similar to *Jon65Aiv*, estimated founder haplotype effects of the *cis* eQTL were negatively correlated with the estimated haplotype effects at the composite G11 hotspot peak (Pearson's r = -98%, p < 0.001, Figure 9B). Unlike the previous candidate mediating

genes, *Jon74E* did not have any connections to any other gene with a *trans* eQTL at the G11 hotspot (Figure 9C).

Figure 9. One gene had a partial mediating effect on the composite eQTL peak for hotspot G11. A. Including Jon74E *(purple line) as a covariate in QTL analysis of the G11 composite variable accounted for the composite QTL peak for H11 (black line). No other gene encompassed by the G11 composite peak had a strong mediating effect on the composite peak (green lines). B. Estimated founder haplotype effects at the composite QTL peak and at a cis eQTL associated with* Jon74E *were significantly negatively correlated (Pearson r = -98%; p < 0.001). C. STRING analysis of the candidate mediating gene and genes with a* trans *eQTL peak at the hotspot did not provide any insight into the link between the candidate mediator and the genes with* trans *eQTL in the hotspot.*

Overall, while we identified several genes with expression patterns that appeared to mediate a *trans* eQTL hotspot effect (Figures S21 and S22), we found limited evidence of functional links between expression of candidate mediating genes and the genes with *trans* eQTL at the corresponding hotspot. Our hotspot analysis did not identify any candidate transcription factors or genes that have previously reported regulatory effects on *trans* eQTL genes in either head or gut tissue.

Discussion

Copper exposure results in tissue- and treatment-specific expression

The genetic architectures of complex traits such as stress resistance and disease are highly context-dependent and can vary across scales as broad as between species and populations (Whitehead and Crawford 2005; Breschi *et al.* 2016; Findley *et al.* 2021) and as small as between individuals, regions of a particular tissue (Fournier and Schacherer 2017; Çelik and Akdaş 2019; Munro *et al.* 2022), or even within individuals across age or development (Everman and Morgan 2017; Huang *et al.* 2020; Everman *et al.* 2021). Given anticipated tissue-specific gene expression

patterns resulting from the spatial distribution of specialized copper-accumulating cells (Calap-Quintana *et al.* 2017; Miguel-Aliaga *et al.* 2018) and the potential for copper toxicity to result in acute damage to digestive tissues as well as neurological tissues (e.g., Tchounwou *et al.* 2008; Jomova *et al.* 2010), one of our principal goals was to characterize the tissue-specific transcriptomic response to copper toxicity. We demonstrate striking patterns of tissue- and treatment-specific genetic response to copper exposure using a combination of differential expression analysis and eQTL mapping. More than 90% of all genes had tissue-specific patterns of expression, and a significant tissue by treatment interaction affected 70% of the genome (Figure 1). Our data suggest that this interaction may be due to greater magnitude and diversity of the gene expression response in Gut tissue compared to Head tissue (Figure 3). Particularly, when stress-related genes were considered as a group we found that the stress response in gut tissue involved multiple stress pathways, whereas oxidative stress response was the primary stress response pathway activated in head tissue (Figure 3). Our expression QTL (eQTL) mapping results provide similar insight; the majority of eQTL were tissue-specific within treatment (Figure S11), suggesting that the genetic control of gene expression response to stress is highly context-dependent.

In general, overexposure to heavy metals results in the accumulation of reactive oxygen species (ROS) and activation of oxidative stress response pathways (e.g. Ercal *et al.* 2001; Gaetke 2003; Uriu-Adams and Keen 2005). Especially for heavy metals that are biologically necessary, there are also metal-specific metabolism pathways that can contribute to the metal toxicity response (Calap-Quintana *et al.* 2017). In the case of copper, specialized cells that are involved in uptake, metabolism, and detoxification of copper ions from the diet line the middle midgut of the fly and the acidic region of the digestive system in vertebrates (Dubreuil 2004; Calap-Quintana *et al.* 2017). Many of the genes that have been linked to copper response in previous studies were significantly differentially expressed under copper conditions relative to control conditions in our study (Figure 2), and our eQTL mapping results suggest that many of these copper-responsive genes are influenced by genetic variation in regulatory elements. Furthermore, we found that genes with functions related to metal response followed expected tissue-specific expression patterns (e.g. *Mtn* family genes; *Ctr1A* and *Ctr1B*; *Atox1* and *SOD3*;

ATP7; discussed above (Zhou *et al.* 2003; Turski and Thiele 2007; Itoh *et al.* 2009; Calap-Quintana

et al. 2017; Kamiya *et al.* 2018)) (Figure 2). For instance, and consistent with previous reports, we

found that *MtnA* expression was strongly induced by copper exposure (Figure 2B); however, the

level of induction was much more pronounced in gut tissue compared with head tissue, leading

to a significant tissue by treatment interaction. We also found variation in the genetic control of

MtnA expression under copper conditions that followed a tissue specific pattern. In gut tissue,

MtnA expression in response to copper is influenced by a *cis* eQTL (Figure 4C), whereas in head

tissue *MtnA* expression under copper conditions is influenced by a *trans* eQTL (Figure 4D).

Although the genetic variants that contribute to *MtnA* expression in either tissue are present in

both tissues, each variant appears to only influence gene expression in a particular tissue.

While *D. melanogaster* is commonly used to characterize the genetic control of heavy

metal response (e.g. Egli *et al.* 2003, 2006a; Balamurugan *et al.* 2004; Norgate *et al.* 2010; Hua *et*

al. 2011), tissue-specific comparisons are not common (Fasae and Abolaji 2022). One of the few

examples of tissue-specific response to heavy metal toxicity in *D. melanogaster* demonstrated

that variation in *CncC* pathway activity (involved in autophagy regulation) across muscle and

neurological tissue contributed to the toxicity response to methylmercury (Gunderson *et al.*

2020). Our differential expression analysis provides novel insight and suggests that there are

tissue-specific pattens in broad responses to stress through the differential activation of

pathways. Although all the stress response categories represented by our subset of genes can be

linked to heavy metal stress response (Darling and Cook 1843; Kefaloyianni *et al.* 2005), our data

suggest that oxidative stress response is the primary stress response mechanism to copper

exposure in neurological tissue (proxied with head tissue in our study) while the primary stress

response mechanisms in gut tissue are more diverse. Because previous studies that offer insight

into tissue-specific gene expression response often focus on sets of tissues that infrequently

overlap, it is challenging to determine whether the patterns we observe in our study are unique

or reflective of a general pattern. However, our data do strongly suggest that there is a tissue-

specific response to copper toxicity over the eight-hour exposure period used in this study.

Whether this difference is due to tissue-specific vulnerability, to mode of exposure, to temporal

progression of copper ions throughout the organism, or a combination of these and other factors

remains to be fully determined. Additional follow up with tissue-specific knockdown or ablation

of key genes involved in oxidative stress and other stress response pathways would be needed

to strengthen these observations.

Transcriptomic response to copper stress is influenced by regulatory allelic variation

Allelic variation and its contribution to trait variation has been repeatedly dissected,

characterized, and functionally tested in the context of complex traits ranging from those

associated with human disease (e.g., Granhall *et al.* 2006; Hardy *et al.* 2018) to various stress

responses including heavy metal resistance (e.g., Zhou *et al.* 2017; Evans *et al.* 2018; Everman *et*

al. 2019, 2021). In addition to the potential to influence the functional gene product, allelic

variation can also influence the response to stress via effects on gene expression patterns. The

majority of allelic variation is found in non-coding regions of the genome (Umans *et al.* 2021),

leading to the expectation that these non-coding variants play a regulatory role. By treating

expression of individual genes as traits measured in multiple treatments and tissues, we were

able to gain insight into context-dependent genetic control of gene expression using expression

eQTL mapping. Approximately half of the annotated *Drosophila* genome was associated with one

or more eQTL in our study, providing ample evidence of genetic variation in the control of gene

expression (Table 2). Nearly all (98%) of genes that were associated with at least one eQTL were

also differentially expressed, suggesting that the variants associated with these genes likely

influence expression, although functional validation would be necessary to verify this hypothesis.

This observation is consistent with work by Boyle *et al.* (2017) demonstrating that SNPs which

contribute to trait variation are enriched near genes that are actively transcribed under disease

states; our observations suggest this pattern extends to stress conditions as well.

Although eQTL mapping provides detailed insight into the genetic control of a given trait

by examining elements that contribute to trait variation on a gene-by-gene basis, there are

several challenges with this approach. Power to detect eQTL with modest effects is directly

related to the number of strains in which the trait is measured (Ruden *et al.* 2009; King *et al.*

2012a; Qu *et al.* 2018; Arvanitis *et al.* 2022). By using 93-96 genotypes in our eQTL analyses, lack

of power to detect modest-effect eQTL likely contributes to our estimates of tissue-specificity.

For example, power estimates to detect a QTL with an effect size of 10 with 100 DSPR strains is

approximately 20% (King *et al.* 2012a). Detection of tissue-specific eQTL is also more sensitive to

false negatives, for example due to low levels of expression of a particular gene in a given tissue.

However, despite these challenges our data suggest that allelic variation that contributes to the

regulation of gene expression is tissue-specific for approximately 65% of eQTL detected in the

control and copper datasets.

Similar studies of tissue-specific eQTL mapping are relatively rare in model organisms but

have provided examples of tissue-specific eQTL. Using a mouse multiparental mapping

population (Collaborative Cross (Aylor *et al.* 2011)) that is similar in concept to the DSPR, Keele

et al. (2020) demonstrated that among lung, liver, and kidney tissue genetic control of several

genes varied across tissues. Spatially discrete genetic control of gene expression can also vary at

the sub-tissue level. Munro *et al.* (2022) demonstrated gene expression patterns that are specific

to five regions of the rat brain are influenced by a combination of shared and sub-tissue specific

eQTL. Shared eQTL were most common in similar brain tissue types, but sub-tissue specific eQTL

were common between more distinct types of brain tissue (Munro *et al.* 2022). eQTL mapping

studies of human disease and native state gene expression have also provided evidence of tissue-

and cell type-specific genetic control of gene expression (Dimas *et al.* 2009; Nica *et al.* 2011;

Fairfax *et al.* 2012; Peters *et al.* 2016; GTEx Consortium 2017; GTEx Consortium *et al.* 2018). A

recent study presenting a novel analytical approach to identify colocalized eQTL suggests that

tissue-specific eQTL may be more common than previously thought and key for identifying

regulatory variants associated with complex disease traits (Arvanitis *et al.* 2022). The existence

of other examples of tissue-specific eQTL combined with our evidence strongly suggest that

tissue-specific eQTL contribute to the stress response to copper toxicity in *D. melanogaster*.

Consistent versus tissue-specific effects of eQTL

In addition to identifying tissue-specific eQTL, our study also provides insight into eQTL

that have consistent versus environment-dependent effects within tissue type. Complex traits

are dependent on both genotype and environmental variation, and the interaction between

genotype and environment can contribute to overall susceptibility of individuals to stress and

disease (Li *et al.* 2006; Duveau and Félix 2012; Moyerbrailean *et al.* 2016; Lea *et al.* 2022). The

importance of genotype by environment interactions and the detection of quantitative loci with

context-dependent effects was demonstrated by Lea *et al*. (2022) in a study using 544 human cell

lines exposed to 12 environments. Using a high-powered experimental design, they

demonstrated that the complex traits measured were influenced by a combination of eQTL with

consistent effects in multiple environments as well as eQTL with environment-dependent effects

(Lea *et al.* 2022). Similarly, Umans *et al*. (2021) offer a review and perspective on the importance

of examining regulatory variants in multiple contexts. Recognizing the dynamic nature of context-

dependent effects of alleles associated with disease in humans, Umans *et al.* (2021) propose that

a critical approach to characterizing allelic variation in regulatory elements is to use multiple

treatment conditions to detect important disease associated variants.

Overall, the majority of the eQTL we detected were near the position of the gene they

were associated with (*cis* eQTL), suggesting that genetic variation in local regulatory elements

plays an important role in the gene expression response to copper stress. *cis* eQTL generally have

larger effects on trait variation and are thus easier to detect with modestly powered designs (Hill

et al. 2021), and *trans* eQTL are additionally difficult to detect because they may not act at the

level of mRNA regulation but instead have larger detectable effects at the protein level (Boyle *et*

al. 2017). *cis* regulatory variation is widely appreciated to be an important contributor to

phenotypic variation and plays a decisive role in the evolution of traits in natural populations and

in human disease (Wray 2007; Savinkova *et al.* 2009; Wittkopp and Kalay 2011; Hill *et al.* 2021).

At the treatment level and consistent with previous studies in humans, flies, and worms (Ruden

et al. 2009; Snoek *et al.* 2017; Qu *et al.* 2018; Sterken *et al.* 2020; Lea *et al.* 2022), we found eQTL

consistently in both treatments as well as eQTL that were only detected in one of the two

treatments (Figure S11). The number of eQTL that were consistently detected was higher than

treatment-specific eQTL, similar to a pattern previously reported for human cell lines (Lea *et al.*

2022). Similarly, in *D. melanogaster*, Qu *et al.* (2018) and Ruden *et al.* (2009) demonstrated that

a small but non-negligeable number of eQTL were only detected following exposure to lead in

head tissue or whole animals, respectively. Our analysis provides an additional layer of insight by

examining the correlation between the estimated haplotype effects at each of the peaks that

were consistently detected under control and copper treatment conditions in head and gut tissue

(Figure 5). We found that estimated effects of founder alleles at shared eQTL often had similar

effects on expression of the gene under different conditions. Despite this apparent consistency,

an effect of treatment was still present in the eQTL detected under copper and control conditions

in gut tissue; eQTL associated with metal-associated genes had larger effects on gene expression

variation under copper conditions compared to control conditions (Figure 5F). While our results

highlight the potential for even eQTL that are repeatedly detected across treatments to have

treatment-specific effects on the expression of a given gene, adding a deeper dimension to gene

by environment interaction eQTL, it is important to consider that the QTL intervals may include

more than one variant that influences gene expression. One of these variants may influence

expression under control conditions while the other influenced expression under copper

conditions. Without additional follow-up studies, our analyses do not provide sufficient

resolution to distinguish this case from one in which the same variant has treatment-specific

effects on gene expression.

In addition to characterizing genotype by environment eQTL by examining the difference

between the control and copper datasets for each tissue, we also examined eQTL that were

associated with a summary metric of the gene expression response to copper stress. The majority

of eQTL detected in the Head- and Gut-Response datasets were *trans* eQTL. Our study design

partially accounts for the deficit in *cis* eQTL in the Response datasets, as any effects of *cis* eQTL

that were detected with similar effects in control and copper datasets would be reduced in the

Response eQTL mapping analyses. However, an enrichment of *trans* eQTL associated with specific

environments has been previously observed in *C. elegans* in response to temperature stress (Li

et al. 2006; Snoek *et al.* 2017). *trans* eQTL that are associated with the dynamic shift in gene

expression response to the environment call attention to allelic variants that may play a role in

the regulation of stress-dependent expression change that may otherwise not be detected if the

treatment conditions are considered independently. We found several Response *trans* eQTL near

genes with a wide range of functions related to response to toxic conditions (Figure 6) that

highlight potential candidate regulatory QTL. Additional follow up would be necessary to

corroborate our observations and to fully characterize the role that these potential QTL sites may

play in the regulation of the response to copper stress.

Conclusions

Exposure to heavy metal stress results in a profound change in transcriptional activity
across multiple tissues, and the genetic control of the gene expression response varies depending
on the environmental conditions and tissue. By taking a combined approach of differential
expression analysis and eQTL mapping, we were able to dissect and characterize more subtle
differences in tissue-specific response to copper toxicity. Our work provides a novel set of
candidate loci that may have context-dependent effects on gene expression and plasticity. From
the patterns that differentiate the genetic response observed in head and gut tissue, we gain
deeper insight into the level of toxicity response that is activated by short exposure to copper
stress.

Acknowledgements

12   We thank Kristen Cloud-Richardson for assistance with collection of head tissue. We also thank
13   the anonymous reviewers for their suggestions that improved this manuscript.

Funding

This project was supported by funding from NIH: ERE was supported by two postdoctoral
fellowships (F32 GM133111 and K99 ES033257) and by the Kansas INBRE (Kansas Idea Network
of Biomedical Research Excellence) project (P30 GM145499). Additional support for experiments
was provided by NIH R01 ES029922 and R01 OD034064 awarded to SJM. Sequencing was
performed by the KU Genome Sequencing Core which is supported by the Center for Molecular
Analysis of Disease Pathways and NIH Centers of Biomedical Research Excellence (P30
GM145499).

Conflict of Interest

23   The authors declare no conflicts of interest.

References

Åkesson A., P. Bjellerup, T. Lundh, J. Lidfeldt, C. Nerbrand, *et al.*, 2006 Cadmium-induced effects on bone in a population-based study of women. Environ. Health Perspect. 114: 830–834. https://doi.org/10.1289/ehp.8763

Albert F. W., J. S. Bloom, J. Siegel, L. Day, and L. Kruglyak, 2018 Genetics of trans-regulatory variation in gene expression. eLife 7: e35471. https://doi.org/10.7554/eLife.35471

Alsheikh A. J., S. Wollenhaupt, E. A. King, J. Reeb, S. Ghosh, *et al.*, 2022 The landscape of GWAS validation; systematic review identifying 309 validated non-coding variants across 130 human diseases. BMC Med. Genomics 15: 74. https://doi.org/10.1186/s12920-022-01216-w

Arvanitis M., K. Tayeb, B. J. Strober, and A. Battle, 2022 Redefining tissue specificity of genetic regulation of gene expression in the presence of allelic heterogeneity. Am. J. Hum. Genet. 109: 223–239. https://doi.org/10.1016/j.ajhg.2022.01.002

Aylor D. L., W. Valdar, W. Foulds-Mathes, R. J. Buus, R. A. Verdugo, *et al.*, 2011 Genetic analysis of complex traits in the emerging Collaborative Cross. Genome Res. 21: 1213–1222. https://doi.org/10.1101/gr.111310.110

Bagheri S., R. Squitti, T. Haertlé, M. Siotto, and A. A. Saboury, 2018 Role of copper in the onset of Alzheimer's Disease compared to other metals. Front. Aging Neurosci. 9: 446. https://doi.org/10.3389/fnagi.2017.00446

Balamurugan K., D. Egli, A. Selvaraj, B. Zhang, O. Georgiev, *et al.*, 2004 Metal-responsive transcription factor (MTF-1) and heavy metal stress response in *Drosophila* and mammalian cells: a functional comparison. Biol. Chem. 385: 597–603. https://doi.org/10.1515/BC.2004.074

Bálint A. F., M. S. Röder, R. Hell, G. Galiba, and A. Börner, 2007 Mapping of QTLs affecting copper tolerance and the Cu, Fe, Mn and Zn contents in the shoots of wheat seedlings. Biol. Plant. 51: 129–134. https://doi.org/10.1007/s10535-007-0025-9

Beavis W. D., 1995 The power and deceit of qtl experiments: Lessons from comparative qtl studies. Proc. Forth-Ninth Annu. Corn Sorghum Ind. Res. Conf. 252–268.

Bisaglia M., and L. Bubacco, 2020 Copper ions and Parkinson's Disease: Why is homeostasis so relevant? Biomolecules 10: 195. https://doi.org/10.3390/biom10020195

Blaurock-Busch E., O. R. Amin, and T. Rabah, 2011 Heavy metals and trace elements in hair and urine of a sample of Arab children with Autistic Spectrum Disorder. Mædica - J. Clin. Med. 6: 247–257.

Bonilla-Ramirez L., M. Jimenez-Del-Rio, and C. Velez-Pardo, 2011 Acute and chronic metal exposure impairs locomotion activity in *Drosophila melanogaster*: a model to study Parkinsonism. BioMetals 24: 1045–1057. https://doi.org/10.1007/s10534-011-9463-0

Boyle E. A., Y. I. Li, and J. K. Pritchard, 2017 An expanded view of complex traits: from polygenic to omnigenic. Cell 169: 1177–1186. https://doi.org/10.1016/j.cell.2017.05.038

Bray N. L., H. Pimentel, P. Melsted, and L. Pachter, 2016 Near-optimal probabilistic RNA-seq quantification. Nat. Biotechnol. 34: 525–527. https://doi.org/10.1038/nbt.3519

Breschi A., S. Djebali, J. Gillis, D. D. Pervouchine, A. Dobin, *et al.*, 2016 Gene-specific patterns of expression variation across organs and species. Genome Biol. 17: 151. https://doi.org/10.1186/s13059-016-1008-y

Broman K. W., D. M. Gatti, P. Simecek, N. A. Furlotte, P. Prins, *et al.*, 2019 R/qtl2: Software for mapping quantitative trait loci with high-dimensional data and multiparent populations. Genetics 211: 495–502. https://doi.org/10.1534/genetics.118.301595

Burke R., E. Commons, and J. Camakaris, 2008 Expression and localisation of the essential copper transporter DmATP7 in *Drosophila* neuronal and intestinal tissues. Int. J. Biochem. Cell Biol. 40: 1850–1860. https://doi.org/10.1016/j.biocel.2008.01.021

Calap-Quintana P., J. González-Fernández, N. Sebastiá-Ortega, J. Llorens, and M. Moltó, 2017 *Drosophila melanogaster* models of metal-related human diseases and metal toxicity. Int. J. Mol. Sci. 18: 1456. https://doi.org/10.3390/ijms18071456

Carlson M., 2019 org.Dm.eg.db: Genome wide annotation for fly. R Package Version 382. https://doi.org/10.18129/B9.bioc.org.Dm.eg.db

Castilla J. C., and E. Nealler, 1978 Marine environmental impact due to mining activities of El Salvador copper mine, Chile. Mar. Pollut. Bull. 9: 67–70. https://doi.org/10.1016/0025-326X(78)90451-4

Catalán A., A. Glaser-Schmitt, E. Argyridou, P. Duchen, and J. Parsch, 2016 An indel polymorphism in the *MtnA* 3' untranslated region is associated with gene expression variation and local adaptation in *Drosophila melanogaster*, (D. J. Begun, Ed.). PLOS Genet. 12: e1005987. https://doi.org/10.1371/journal.pgen.1005987

Çelik Ö., and E. Y. Akdaş, 2019 Tissue-specific transcriptional regulation of seven heavy metal stress-responsive miRNAs and their putative targets in nickel indicator castor bean (*R. communis L.*) plants. Ecotoxicol. Environ. Saf. 170: 682–690. https://doi.org/10.1016/j.ecoenv.2018.12.006

Chen Y., A. T. Lun, and G. K. Smyth, 2016a From reads to genes to pathways: differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR qyasi-likelihood pipeline. F1000Research 5: 1–51. https://doi.org/10.12688/f1000research.8987.2

Chen P., M. R. Miah, and M. Aschner, 2016b Metals and neurodegeneration. F1000Research 5: 366. https://doi.org/10.12688/f1000research.7431.1

Chen S., Y. Zhou, Y. Chen, and J. Gu, 2018 fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics 34: i884–i890. https://doi.org/10.1093/bioinformatics/bty560

Courbot M., G. Willems, P. Motte, S. Arvidsson, N. Roosens, *et al.*, 2007 A major quantitative trait locus for cadmium tolerance in *Arabidopsis halleri* colocalizes with *HMA4*, a gene encoding a heavy metal ATPase. Plant Physiol. 144: 1052–1065. https://doi.org/10.1104/pp.106.095133

Culotta V. C., L. W. J. Klomp, J. Strain, R. L. B. Casareno, B. Krems, *et al.*, 1997 The copper chaperone for superoxide dismutase. J. Biol. Chem. 272: 23469–23472. https://doi.org/10.1074/jbc.272.38.23469

Cunningham F., J. E. Allen, J. Allen, J. Alvarez-Jarreta, M. R. Amode, *et al.*, 2022 Ensembl 2022. Nucleic Acids Res. 50: D988–D995. https://doi.org/10.1093/nar/gkab1049

Daborn P. J., C. Lumb, A. Boey, W. Wong, R. H. ffrench-Constant, *et al.*, 2007 Evaluating the insecticide resistance potential of eight *Drosophila melanogaster* cytochrome P450 genes by transgenic over-expression. Insect Biochem. Mol. Biol. 37: 512–519. https://doi.org/10.1016/j.ibmb.2007.02.008

Darling N. J., and S. J. Cook, 1843 The role of MAPK signalling pathways in the response to endoplasmic reticulum stress. Biochim. Biophys. Acta.

De Koning D.-J., and C. S. Haley, 2005 Genetical genomics in humans and model organisms. Trends Genet. 21: 377–381. https://doi.org/10.1016/j.tig.2005.05.004

Denecke S., R. Fusetto, and P. Batterham, 2017 Describing the role of *Drosophila melanogaster* ABC transporters in insecticide biology using CRISPR-Cas9 knockouts. Insect Biochem. Mol. Biol. 91: 1–9. https://doi.org/10.1016/j.ibmb.2017.09.017

Dimas A. S., S. Deutsch, B. E. Stranger, S. B. Montgomery, C. Borel, *et al.*, 2009 Common regulatory variation impacts gene expression in a cell type–dependent manner. Science 325: 1246–1250. https://doi.org/10.1126/science.1174148

Dixon A. L., L. Liang, M. F. Moffatt, W. Chen, S. Heath, *et al.*, 2007 A genome-wide association study of global gene expression. Nat. Genet. 39: 1202–1207. https://doi.org/10.1038/ng2109

Dubreuil R. R., 2004 Copper cells and stomach acid secretion in the *Drosophila* midgut. Int. J. Biochem. Cell Biol. 36: 742–752. https://doi.org/10.1016/j.biocel.2003.07.004

Durinck S., Y. Moreau, A. Kasprzyk, S. Davis, B. De Moor, *et al.*, 2005 BioMart and Bioconductor: a powerful link between biological databases and microarray data analysis. Bioinformatics 21: 3439–3440. https://doi.org/10.1093/bioinformatics/bti525

Duveau F., and M.-A. Félix, 2012 Role of pleiotropy in the evolution of a cryptic developmental variation in *Caenorhabditis elegans*, (M. A. F. Noor, Ed.). PLoS Biol. 10: e1001230. https://doi.org/10.1371/journal.pbio.1001230

Egli D., A. Selvaraj, H. Yepiskoposyan, B. Zhang, E. Hafen, *et al.*, 2003 Knockout of 'metal-responsive transcription factor' MTF-1 in *Drosophila* by homologous recombination reveals its central role in heavy metal homeostasis. EMBO J. 22: 100–108. https://doi.org/10.1093/emboj/cdg012

Egli D., H. Yepiskoposyan, A. Selvaraj, K. Balamurugan, R. Rajaram, *et al.*, 2006a A family knockout of all four *Drosophila* metallothioneins reveals a central role in copper homeostasis and detoxification. Mol. Cell. Biol. 26: 2286–2296. https://doi.org/10.1128/MCB.26.6.2286-2296.2006

Egli D., J. Domènech, A. Selvaraj, K. Balamurugan, H. Hua, *et al.*, 2006b The four members of the *Drosophila* metallothionein family exhibit distinct yet overlapping roles in heavy metal homeostasis and detoxification. Genes Cells 11: 647–658. https://doi.org/10.1111/j.1365-2443.2006.00971.x

Ehrenreich I. M., J. Bloom, N. Torabi, X. Wang, Y. Jia, *et al.*, 2012 Genetic architecture of highly complex chemical resistance traits across four yeast strains, (A. P. Gasch, Ed.). PLoS Genet. 8: e1002570. https://doi.org/10.1371/journal.pgen.1002570

Emilsson V., G. Thorleifsson, B. Zhang, A. S. Leonardson, F. Zink, *et al.*, 2008 Genetics of gene expression and its effect on disease. Nature 452: 423–428. https://doi.org/10.1038/nature06758

Ercal N., H. Gurer-Orhan, and N. Aykin-Burns, 2001 Toxic metals and oxidative stress part I: Mechanisms involved in metal induced oxidative damage. Curr. Top. Med. Chem. 1: 529–539.

Esteves F., J. Rueff, and M. Kranendonk, 2021 The central role of cytochrome P450 in xenobiotic metabolism—a brief review on a fascinating enzyme family. J. Xenobiotics 11: 94–114. https://doi.org/10.3390/jox11030007

Evans K. S., S. C. Brady, J. S. Bloom, R. E. Tanny, D. E. Cook, *et al.*, 2018 Shared genomic regions underlie natural variation in diverse toxin responses. Genetics 210: 1509–1525. https://doi.org/10.1534/genetics.118.301311

Everman E. R., and T. J. Morgan, 2017 Evolution of age-specific decline in stress phenotypes is driven by both antagonistic pleiotropy and mutation accumulation. bioRxiv 115931.

Everman E. R., C. L. McNeil, J. L. Hackett, C. L. Bain, and S. J. Macdonald, 2019 Dissection of complex, fitness-related traits in multiple *Drosophila* mapping populations offers insight into the genetic control of stress resistance. Genetics 211: 19. https://doi.org/10.1534/genetics.119.301930

Everman E. R., K. M. Cloud-Richardson, and S. J. Macdonald, 2021 Characterizing the genetic basis of copper toxicity in *Drosophila* reveals a complex pattern of allelic, regulatory, and behavioral variation. Genetics 217: 1–20. https://doi.org/10.1093/genetics/iyaa020

Everman E. R., S. J. Macdonald, and J. K. Kelly, 2023 The genetic basis of adaptation to copper pollution in *Drosophila melanogaster*. Front. Genet. 14: 1144221. https://doi.org/10.3389/fgene.2023.1144221

Fairfax B. P., S. Makino, J. Radhakrishnan, K. Plant, S. Leslie, *et al.*, 2012 Genetics of gene expression in primary immune cells identifies cell type–specific master regulators and roles of HLA alleles. Nat. Genet. 44: 502–510. https://doi.org/10.1038/ng.2205

Falcon S., and R. Gentleman, 2007 Using GOstats to test gene lists for GO term association. Bioinformatics 23: 257–258. https://doi.org/10.18129/B9.bioc.GOstats

Fasae K. D., and A. O. Abolaji, 2022 Interactions and toxicity of non-essential heavy metals (Cd, Pb and Hg): lessons from *Drosophila melanogaster*. Curr. Opin. Insect Sci. 51: 100900. https://doi.org/10.1016/j.cois.2022.100900

Felmlee K. R., S. J. Macdonald, and E. R. Everman, 2022 Pre-adult exposure to three heavy metals leads to changes in the head transcriptome of adult flies. MicroPublication Biol. 1–8.

Findley A. S., A. Monziani, A. L. Richards, K. Rhodes, M. C. Ward, *et al.*, 2021 Functional dynamic genetic effects on gene regulation are specific to particular cell types and environmental conditions. eLife 10: e67077. https://doi.org/10.7554/eLife.67077

Fournier T., and J. Schacherer, 2017 Genetic backgrounds and hidden trait complexity in natural populations. Curr. Opin. Genet. Dev. 47: 48–53. https://doi.org/10.1016/j.gde.2017.08.009

Gaetke L., 2003 Copper toxicity, oxidative stress, and antioxidant nutrients. Toxicology 189: 147–163. https://doi.org/10.1016/S0300-483X(03)00159-8

Gaffney D. J., J.-B. Veyrieras, J. F. Degner, R. Pique-Regi, A. A. Pai, *et al.*, 2012 Dissecting the regulatory architecture of gene expression QTLs. Genome Biol. 13: R7. https://doi.org/10.1186/gb-2012-13-1-r7

Gerhardsson L., 2022 Diagnosis and treatment of metal poisoning general aspects, pp. 663–684 in *Handbook on the Toxicology of Metals: general considerations*, Academic Press.

Gibson G., and B. Weir, 2005 The quantitative genetics of transcription. Trends Genet. 21: 616–623. https://doi.org/10.1016/j.tig.2005.08.010

Gilad Y., S. A. Rifkin, and J. K. Pritchard, 2008 Revealing the architecture of gene regulation: the promise of eQTL studies. Trends Genet. 24: 408–415. https://doi.org/10.1016/j.tig.2008.06.001

González M., A. Reyes-Jara, M. Suazo, W. J. Jo, and C. Vulpe, 2008 Expression of copper-related genes in response to copper load. Am. J. Clin. Nutr. 88: 830S-834S. https://doi.org/10.1093/ajcn/88.3.830S

Granhall C., H.-B. Park, H. Fakhrai-Rad, and H. Luthman, 2006 High-resolution quantitative trait locus analysis reveals multiple diabetes susceptibility loci mapped to intervals <800 kb in the species-conserved *Niddm1i* of the GK rat. Genetics 174: 1565–1572. https://doi.org/10.1534/genetics.106.062208

Green L., M. Coronado-Zamora, S. Radío, G. E. Rech, J. Salces-Ortiz, *et al.*, 2022 The genomic basis of copper tolerance in *Drosophila* is shaped by a complex interplay of regulatory and environmental factors. BMC Biol. 20: 275. https://doi.org/10.1186/s12915-022-01479-w

GTEx Consortium, 2017 Genetic effects on gene expression across human tissues. Nature 550: 204–213. https://doi.org/10.1038/nature24277

GTEx Consortium, E. R. Gamazon, A. V. Segrè, M. van de Bunt, X. Wen, *et al.*, 2018 Using an atlas of gene regulation across 44 human tissues to inform complex disease- and trait-associated variation. Nat. Genet. 50: 956–967. https://doi.org/10.1038/s41588-018-0154-4

Gunderson J. T., A. E. Peppriell, D. Vorojeikina, and M. D. Rand, 2020 Tissue-specific Nrf2 signaling protects against methylmercury toxicity in *Drosophila* neuromuscular development. Arch. Toxicol. 94: 4007–4022. https://doi.org/10.1007/s00204-020-02879-z

Hardy C. M., M. K. Burke, L. J. Everett, M. V. Han, K. M. Lantz, *et al.*, 2018 Genome-wide analysis of starvation-selected *Drosophila melanogaster*—a genetic model of obesity. Mol. Biol. Evol. 35: 50–65. https://doi.org/10.1093/molbev/msx254

Hatori Y., and S. Lutsenko, 2013 An expanding range of functions for the copper chaperone/antioxidant protein Atox1. Antioxid. Redox Signal. 19: 945–957. https://doi.org/10.1089/ars.2012.5086

He Z., J. Shentu, X. Yang, V. C. Baligar, T. Zhang, *et al.*, 2015 Heavy metal contamination of soils: sources, indicators, and assessment. J. Environ. Indic. 9: 17–18.

Hill M. S., P. Vande Zande, and P. J. Wittkopp, 2021 Molecular and evolutionary processes generating variation in gene expression. Nat. Rev. Genet. 22: 203–215. https://doi.org/10.1038/s41576-020-00304-w

Hindorff L. A., P. Sethupathy, H. A. Junkins, E. M. Ramos, J. P. Mehta, *et al.*, 2009 Potential etiologic and functional implications of genome-wide association loci for human diseases and traits. Proc. Natl. Acad. Sci. 106: 9362–9367. https://doi.org/10.1073/pnas.0903103106

Hsueh Y.-M., C.-Y. Lee, S.-N. Chien, W.-J. Chen, H.-S. Shiue, *et al.*, 2017 Association of blood heavy metals with developmental delays and health status in children. Sci. Rep. 7: 43608. https://doi.org/10.1038/srep43608

Hua H., V. Günther, O. Georgiev, and W. Schaffner, 2011 Distorted copper homeostasis with decreased sensitivity to cisplatin upon chaperone *Atox1* deletion in *Drosophila*. BioMetals 24: 445–453. https://doi.org/10.1007/s10534-011-9438-1

Huang W., T. Campbell, M. A. Carbone, W. E. Jones, D. Unselt, *et al.*, 2020 Context-dependent genetic architecture of *Drosophila* life span, (N. H. Barton, Ed.). PLOS Biol. 18: e3000645. https://doi.org/10.1371/journal.pbio.3000645

Itoh S., K. Ozumi, H. Kim, O. Nakagawa, R. Mckinney, *et al.*, 2009 Novel mechanism for regulation of extracellular SOD transcription and activity by copper: Role of antioxidant-1. Free Radic. Biol. Med. 46: 95–104. https://doi.org/10.1016/j.freeradbiomed.2008.09.039

Jansen R., 2001 Genetical genomics: the added value from segregation. Trends Genet. 17: 388–391. https://doi.org/10.1016/S0168-9525(01)02310-1

Jomova K., D. Vondrakova, M. Lawson, and M. Valko, 2010 Metals, oxidative stress and neurodegenerative disorders. Mol. Cell. Biochem. 345: 91–104. https://doi.org/10.1007/s11010-010-0563-x

Jones L. C., K. A. McCarthy, J. L. Beard, C. L. Keen, and B. C. Jones, 2006 Quantitative genetic analysis of brain copper and zinc in BXD recombinant inbred mice. Nutr. Neurosci. 9: 81–92. https://doi.org/10.1080/00268970600691365

Kamiya T., K. Takeuchi, S. Fukudome, H. Hara, and T. Adachi, 2018 Copper chaperone antioxidant-1, Atox-1, is involved in the induction of SOD3 in THP-1 cells. BioMetals 31: 61–68. https://doi.org/10.1007/s10534-017-0067-1

Karnchanawong S., and P. Limpiteeprakan, 2009 Evaluation of heavy metal leaching from spent household batteries disposed in municipal solid waste. Waste Manag. 29: 550–558. https://doi.org/10.1016/j.wasman.2008.03.018

Keele G. R., B. C. Quach, J. W. Israel, G. A. Chappell, L. Lewis, *et al.*, 2020 Integrative QTL analysis of gene expression and chromatin accessibility identifies multi-tissue patterns of genetic regulation, (G. S. Barsh, Ed.). PLOS Genet. 16: e1008537. https://doi.org/10.1371/journal.pgen.1008537

Kefaloyianni E., E. Gourgou, V. Ferle, E. Kotsakis, C. Gaitanaki, *et al.*, 2005 Acute thermal stress and various heavy metals induce tissue-specific pro-or anti-apoptotic events *via* the p38-MAPK signal transduction pathway in *Mytilus galloprovincialis* (Lam.). J. Exp. Biol. 208: 4427–4436. https://doi.org/10.1242/jeb.01924

Khan S., Q. Cao, Y. M. Zheng, Y. Z. Huang, and Y. G. Zhu, 2008 Health risks of heavy metals in contaminated soils and food crops irrigated with wastewater in Beijing, China. Environ. Pollut. 152: 686–692. https://doi.org/10.1016/j.envpol.2007.06.056

Kim K., and J. Yim, 2013 The diverse biological functions of glutathione S-transferase omega in *Drosophila*. Pteridines 24: 117–120. https://doi.org/10.1515/pterid-2013-0017

Kim D., J. M. Paggi, C. Park, C. Bennett, and S. L. Salzberg, 2019 Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat. Biotechnol. 37: 907–915. https://doi.org/10.1038/s41587-019-0201-4

King E. G., S. J. Macdonald, and A. D. Long, 2012a Properties and power of the *Drosophila* Synthetic Population Resource for the routine dissection of complex traits. Genetics 191: 935–949. https://doi.org/10.1534/genetics.112.138537

King E. G., C. M. Merkes, C. L. McNeil, S. R. Hoofer, S. Sen, *et al.*, 2012b Genetic dissection of a model complex trait using the *Drosophila* Synthetic Population Resource. Genome Res. 22: 1558–1566. https://doi.org/10.1101/gr.134031.111

King E. G., B. J. Sanderson, C. L. McNeil, A. D. Long, and S. J. Macdonald, 2014 Genetic dissection of the *Drosophila melanogaster* female head transcriptome reveals widespread allelic heterogeneity, (G. Gibson, Ed.). PLoS Genet. 10: e1004322. https://doi.org/10.1371/journal.pgen.1004322

King E. G., and A. D. Long, 2017 The Beavis effect in next-generation mapping panels in *Drosophila melanogaster*. G3 Genes Genomes Genet. 7: 1643–1652.

Knoblauch A., M. Divall, M. Owuor, C. Archer, K. Nduna, *et al.*, 2017 Monitoring of selected health indicators in children living in a copper mine development area in northwestern Zambia. Int. J. Environ. Res. Public. Health 14: 315. https://doi.org/10.3390/ijerph14030315

Kumar K., A. Mhetre, G. S. Ratnaparkhi, and S. S. Kamat, 2021 A superfamily-wide activity atlas of serine hydrolases in *Drosophila melanogaster*. Biochemistry 60: 1312–1324. https://doi.org/10.1021/acs.biochem.1c00171

Lang M., C. L. Braun, M. R. Kanost, and M. J. Gorman, 2012 Multicopper oxidase-1 is a ferroxidase essential for iron homeostasis in *Drosophila melanogaster*. Proc. Natl. Acad. Sci. 109: 13337–13342. https://doi.org/10.1073/pnas.1208703109

Lea A. J., J. Peng, and J. F. Ayroles, 2022 Diverse environmental perturbations reveal the evolution and context-dependency of genetic effects on gene expression levels. Genome Res. genome;gr.276430.121v1. https://doi.org/10.1101/gr.276430.121

Lee M.-J., M.-C. Chou, W.-J. Chou, C.-W. Huang, H.-C. Kuo, *et al.*, 2018 Heavy metals' effect on susceptibility to Attention-Deficit/Hyperactivity Disorder: Implication of lead, cadmium, and antimony. Int. J. Environ. Res. Public. Health 15: 1221. https://doi.org/10.3390/ijerph15061221

Leek J. T., and J. D. Storey, 2007 Capturing heterogeneity in gene expression studies by surrogate variable analysis. PLOS Genet. 3: e161. https://doi.org/doi:10.1371/journal.pgen. 0030161

Li Y., E. Hazendonk, P. Prins, R. H. A. Plasterk, R. C. Jansen, *et al.*, 2006 Mapping determinants of gene expression plasticity by genetical genomics in *C. elegans*. PLoS Genet. 2.

Li H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, *et al.*, 2009 The Sequence Alignment/Map format and SAMtools. Bioinformatics 25: 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Liao Y., G. K. Smyth, and W. Shi, 2014 featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics 30: 923–930. https://doi.org/10.1093/bioinformatics/btt656

Liddell J. R., and A. R. White, 2018 Nexus between mitochondrial function, iron, copper and glutathione in Parkinson's disease. Neurochem. Int. 117: 126–138. https://doi.org/10.1016/j.neuint.2017.05.016

Lyne R., R. Smith, K. Rutherford, M. Wakeling, A. Varley, *et al.*, 2007 FlyMine: an integrated database for *Drosophila* and *Anopheles* genomics. Genome Biol. 8: R129. https://doi.org/10.1186/gb-2007-8-7-r129

MacNair M. R., 1983 The genetic control of copper tolerance in the yellow monkey flower, *Mimulus guttatus*. Heredity 53: 283–293.

Miguel-Aliaga I., H. Jasper, and B. Lemaitre, 2018 Anatomy and physiology of the digestive tract of *Drosophila melanogaster*. Genetics 210: 357–396. https://doi.org/10.1534/genetics.118.300224

Moyerbrailean G. A., A. L. Richards, D. Kurtz, C. A. Kalita, G. O. Davis, *et al.*, 2016 High-throughput allele-specific expression across 250 environmental conditions. Genome Res. 26: 1627–1638. https://doi.org/10.1101/gr.209759.116

Munro D., T. Wang, A. S. Chitre, O. Polesskaya, N. Ehsan, *et al.*, 2022 The regulatory landscape of multiple brain regions in outbred heterogeneous stock rats. Nucleic Acids Res. 50: 10882–10895. https://doi.org/10.1093/nar/gkac912

Neuberger J. S., M. Mulhall, M. C. Pomatto, J. Sheverbush, and R. S. Hassanein, 1990 Health problems in Galena, Kansas: a heavy metal superfund site. Sci. Total Environ. 94: 261–272.

Nica A. C., L. Parts, D. Glass, J. Nisbet, A. Barrett, *et al.*, 2011 The architecture of gene regulatory variation across multiple human tissues: The MuTHER Study, (G. Barsh, Ed.). PLoS Genet. 7: e1002003. https://doi.org/10.1371/journal.pgen.1002003

Norgate M., A. Southon, M. Greenough, M. Cater, A. Farlow, *et al.*, 2010 Syntaxin 5 is required for copper homeostasis in *Drosophila* and mammals, (J. M. Bridger, Ed.). PLoS ONE 5: e14303. https://doi.org/10.1371/journal.pone.0014303

Peters J. E., P. A. Lyons, J. C. Lee, A. C. Richard, M. D. Fortune, *et al.*, 2016 Insight into genotype-phenotype associations through eQTL mapping in multiple cell types in health and immune-mediated disease, (V. Plagnol, Ed.). PLOS Genet. 12: e1005908. https://doi.org/10.1371/journal.pgen.1005908

Pickrell J. K., J. C. Marioni, A. A. Pai, J. F. Degner, B. E. Engelhardt, *et al.*, 2010 Understanding mechanisms underlying human gene expression variation with RNA sequencing. Nature 464: 768–772. https://doi.org/10.1038/nature08872

Pickrell J. K., 2014 Joint analysis of functional genomic data and genome-wide association studies of 18 human traits. Am. J. Hum. Genet. 94: 559–573. https://doi.org/10.1016/j.ajhg.2014.03.004

Plomin R., C. M. A. Haworth, and O. S. P. Davis, 2009 Common disorders are quantitative traits. Nat. Rev. Genet. 10: 872–878. https://doi.org/10.1038/nrg2670

Qu W., K. Gurdziel, R. Pique-Regi, and D. M. Ruden, 2018 Lead modulates trans- and cis-expression quantitative trait loci (eQTLs) in *Drosophila melanogaster* heads. Front. Genet. 9: 395. https://doi.org/10.3389/fgene.2018.00395

R Core Team, 2017 R: A language and environment for statistical computing

Ritchie M. E., B. Phipson, D. Wu, Y. Hu, C. W. Law, *et al.*, 2015 *limma* powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res. 43: e47–e47. https://doi.org/10.1093/nar/gkv007

Robinson M. D., and A. Oshlack, 2010 A scaling normalization method for differential expression analysis of RNA-seq data. Genome Biol. 11: R25. https://doi.org/10.1186/gb-2010-11-3-r25

Ruden D. M., L. Chen, D. Possidente, B. Possidente, P. Rasouli, *et al.*, 2009 Genetical toxicogenomics in *Drosophila* identifies master-modulatory loci that are regulated by

developmental exposure to lead. NeuroToxicology 30: 898–914.
https://doi.org/10.1016/j.neuro.2009.08.011

Savinkova L. K., M. P. Ponomarenko, P. M. Ponomarenko, I. A. Drachkova, M. V. Lysova, *et al.*, 2009 TATA box polymorphisms in human gene promoters and associated hereditary pathologies. Biochem. Mosc. 74: 117–129. https://doi.org/10.1134/S0006297909020011

Seong C.-S., A. Varela-Ramirez, X. Tang, B. Anchondo, D. Magallanes, *et al.*, 2014 Cloning and characterization of a novel *Drosophila* stress induced DNase, (S. Cotterill, Ed.). PLoS ONE 9: e103564. https://doi.org/10.1371/journal.pone.0103564

Snoek B. L., M. G. Sterken, R. P. J. Bevers, R. J. M. Volkers, A. van't Hof, *et al.*, 2017 Contribution of trans regulatory eQTL to cryptic genetic variation in *C. elegans*. BMC Genomics 18: 500. https://doi.org/10.1186/s12864-017-3899-8

Sterken M. G., R. P. J. Bevers, R. J. M. Volkers, J. A. G. Riksen, J. E. Kammenga, *et al.*, 2020 Dissecting the eQTL micro-architecture in *Caenorhabditis elegans*. Front. Genet. 11: 501376. https://doi.org/10.3389/fgene.2020.501376

Stohs S. J., and D. Bagchi, 1995 Oxidative mechanisms in the toxicity of metal ions. Free Radic. Biol. Med. 18: 321–336.

Sun H., N. Buchon, and J. G. Scott, 2017 Mdr65 decreases toxicity of multiple insecticides in *Drosophila melanogaster*. Insect Biochem. Mol. Biol. 89: 11–16. https://doi.org/10.1016/j.ibmb.2017.08.002

Szklarczyk D., R. Kirsch, M. Koutrouli, K. Nastou, F. Mehryary, *et al.*, 2023 The STRING database in 2023: protein–protein association networks and functional enrichment analyses for any sequenced genome of interest. Nucleic Acids Res. 51: D638–D646. https://doi.org/10.1093/nar/gkac1000

Taylor A. A., J. S. Tsuji, M. R. Garry, M. E. McArdle, W. L. Goodfellow, *et al.*, 2020 Critical review of exposure and effects: implications for setting regulatory health criteria for ingested copper. Environ. Manage. 65: 131–159. https://doi.org/10.1007/s00267-019-01234-y

Tchounwou P. B., C. Newsome, J. Williams, and K. Glass, 2008 Copper-induced cytotoxicity and transcriptional activation of stress genes in human liver carcinoma (HepG2) cells. Met. Ions Biol. Med. 10: 285–290.

The UniProt Consortium, A. Bateman, M.-J. Martin, S. Orchard, M. Magrane, *et al.*, 2021 UniProt: the universal protein knowledgebase in 2021. Nucleic Acids Res. 49: D480–D489. https://doi.org/10.1093/nar/gkaa1100

Thurmond J., J. L. Goodman, V. B. Strelets, H. Attrill, L. S. Gramates, *et al.*, 2019 FlyBase 2.0: the next generation. Nucleic Acids Res. 47: D759–D765. https://doi.org/10.1093/nar/gky1003

Turgut S., A. Polat, G. Turgut, G. Emmungil, M. Bican, *et al.*, 2007 Interaction betwen anemia and blood leves of iron, zinc, copper, cadmium and lead in children. Indian J. Pediatr. 74: 827–830. https://doi.org/DOI: 10.1007/s12098-007-0147-2

Turski M. L., and D. J. Thiele, 2007 *Drosophila* Ctr1A functions as a copper transporter essential for development. J. Biol. Chem. 282: 24017–24026. https://doi.org/10.1074/jbc.M703792200

Umans B. D., A. Battle, and Y. Gilad, 2021 Where are the disease-associated eQTLs? Trends Genet. 37: 109–124. https://doi.org/10.1016/j.tig.2020.08.009

Uriu-Adams J. Y., and C. L. Keen, 2005 Copper, oxidative stress, and human health. Mol. Aspects Med. 26: 268–298. https://doi.org/10.1016/j.mam.2005.07.015

Whitehead A., and D. L. Crawford, 2005 Variation in tissue-specific gene expression among natural populations. Genome Biol. 6:R13

Wittkopp P. J., and G. Kalay, 2011 Cis-regulatory elements: molecular mechanisms and evolutionary processes underlying divergence. Nat. Rev. Genet. 13: 59-69. https://doi.org/10.1038/nrg3095

World Health Organization, Food and Agriculture Organization of the United Nations, and International Atomic Energy Agency (Eds.), 1996 *Trace elements in human nutrition and health*. World Health Organization, Geneva.

Wray G. A., 2007 The evolutionary significance of cis-regulatory mutations. Nat. Rev. Genet. 8: 206–216. https://doi.org/10.1038/nrg2063

Zhang F., and J. R. Lupski, 2015 Non-coding genetic variants in human disease. Hum. Mol. Genet. 24: R102–R110. https://doi.org/10.1093/hmg/ddv259

Zhou H., K. M. Cadigan, and D. J. Thiele, 2003 A copper-regulated transporter required for copper acquisition, pigmentation, and specific stages of development in *Drosophila melanogaster*. J. Biol. Chem. 278: 48210–48218. https://doi.org/10.1074/jbc.M309820200

Zhou S., T. V. Morozova, Y. N. Hussain, S. E. Luoma, L. McCoy, *et al.*, 2016 The genetic basis for variation in sensitivity to lead toxicity in *Drosophila melanogaster*. Environ. Health Perspect. 124: 1062–1070. https://doi.org/10.1289/ehp.1510513

Zhou S., S. E. Luoma, G. E. St. Armour, E. Thakkar, T. F. C. Mackay, *et al.*, 2017 A *Drosophila* model for toxicogenomics: Genetic variation in susceptibility to heavy metal exposure, (G. P. Copenhaver, Ed.). PLOS Genet. 13: e1006907. https://doi.org/10.1371/journal.pgen.1006907

Supplemental Figures

Figure S1. Adult copper resistance of B panel RILs used in the current study. We randomly sampled 48 resistant and 48 sensitive DSPR B panel RILs from the top and bottom 25% of the distribution of adult copper survival response following 48-hour exposure to 50mM $CuSO_4$ reported in (Everman et al. 2021).

Figure S2. Distributions of raw read pair counts by library pool. Each tissue-specific pool included copper and control sample pairs, and an even representation of resistant and sensitive RILs. HO stands for "high-output" NextSeq500 flowcell, MO stands for "mid-output" flowcell. Raw reads from the two sequencing runs of the 96-plex Heads library (green shading) were combined prior to read processing.

Figure S3. Comparisons of the number of raw read pairs, the number retained following trimming, the percentage aligned, and the percentage assigned to genes following the HISAT2 pipeline from the high-output (HO) and mid-output (MO) sequencing runs for the 96-plex Head

library. Apart from raw pair count, which varies due to under-clustering of the 96-plex Head library, data are consistent between the HO and MO sequencing of the same 96-plex library and are comparable to the separate 94-plex Head library that was only sequenced on one HO flowcell.

Figure S4. Variation in read length across all 379 samples. After trimming low quality bases and filtering short reads, only read pairs with >15 bp per read were retained.

Figure S5. Gene expression estimates from the HISAT2 and kallisto pipelines were consistent. After alignment and filtering out of lowly-expressed genes, expression estimates for 9842 and 9913 genes from the HISAT2 and kallisto pipelines, respectively, were retained. Gene-wise correlations in read counts across all samples for shared retained genes (9434) were highly correlated (mean R = 98.2%).

Figure S6. Correcting for known and unknown sources of technical variation reduced noise in eQTL mapping. Head-Control (top row) and Head-Copper (middle row) eQTL maps for *Sod1* are noisy when technical variation is not accounted for (left column). Correcting for 2% (middle column) and 1% PCs (right column) along with those associated with known technical factors appears to limit noise. Head-Response data (bottom row) were less sensitive to data correction. In each plot, orange vertical lines indicate the location of the gene. Horizontal lines indicate the gene- and dataset-specific significance threshold.

Figure S7. Correcting for PCs that explained 2% or 1% of the variation in gene expression resulted in similar numbers of eQTL. Head samples are shown in A; Gut samples are shown in B. In both panels, the left column shows the 2% correction, the right column shows the 1% correction.

Figure S8. Genome-wide LOD scores were similar between the 1% and 2% corrected datasets for each tissue and treatment. Head-Copper, -Control, and -Response correlations are shown in A-C. Gut-Copper, -Control, and -Response correlations are shown in D-F. In each panel, orange highlights genome-wide LOD score correlations for genes that are associated with at least one eQTL in the 2% corrected data; grey highlights genes that were not associated with eQTL. Vertical color-coded lines indicate mean correlations for each group of genes.

Figure S9. For most genes (>80%), mapping yielded a single eQTL. Data shown are counts of distinct eQTL peaks per gene for each of the 6 datasets. Gut tissue is shown in orange, head tissue is shown in red. Numbers above bars provide the actual number of genes in each category.

Figure S10. Percent variance explained by mapped *cis* and *trans* eQTL. *cis* eQTL (blue) tended to have higher estimates compared to *trans* eQTL (red). Estimates were slightly lower in both response datasets for both *cis* and *trans* eQTL. Numbers at the base of each plot report the total number of *cis* and *trans* eQTL detected per dataset.

Figure S11. Many eQTL were detected in multiple datasets. Overlap was highest within tissue between treatments (A and B) but was still substantial in comparisons between tissues within treatment (C, D). Comparisons between the Head- and Gut-Response datasets revealed that

most eQTL were distinct between the tissue-specific characterization of copper response. Overlap was also high in comparisons of the Response datasets against the tissue- and treatment-specific datasets (F – I). The majority (64 – 81%) of *cis* Response eQTL were shared with either the Control or Copper datasets for the corresponding tissue. For Head tissue, all but 9 *cis* Response eQTL were shared amongst Head-Control or Head-Copper *cis* eQTL; for gut tissue, all but 14 *cis* Response eQTL were shared amongst Gut-Control or Gut-Copper cis eQTL (data not shown).

Figure S12. Founder haplotype effects at *cis* eQTL peak positions that were detected in Response, Copper, and Control datasets for a given tissue ranged in strength of correlation for each comparison: A. Head-Response vs Head-Control, B. Head-Response vs Head-Copper, C. Gut-Response vs Gut-Control, D. Gut-Response vs Gut-Copper. Founder effect estimates are impacted by low replication per DSPR strain and should be interpreted with care; however, patterns suggest the detection of Response eQTL in our study may be influenced by a combination of genetic variants with different magnitude effects and treatment-specific additive effects.

Figure S13. Enrichment of GO categories for genes with eQTL in Head-Copper, Gut-Copper, or both datasets.

Figure S14. Correlations of estimated founder haplotype effects at peak positions of eQTL detected in both the Head-Copper and Gut-Copper datasets skewed positive for the majority of eQTL. A. Estimated founder haplotype effect correlations at eQTL associated with genes with functions unrelated to metal response based on FlyBase annotations were generally continuous and ranged from a small number of strong negative correlations to strong positive correlations. B. Estimated founder haplotype effect correlations at eQTL associated with genes with functions that are related to metal response (oxidative stress response, response to copper, response to xenobiotics, detoxification) generally fell into two groups with a small number of strong negative correlations and a larger number of strong positive correlations.

Figure S15. Gene ontology enrichment for potential genotype by environment eQTL that were detected in both control and copper conditions. A. Genes with eQTL that were detected in head tissue under copper and control conditions were enriched for categories related to metal ion transport and stress. B. Genes with eQTL that were detected in gut tissue under copper and control conditions were also enriched for metal-related processes. The size of plotted circles represents the relative number of genes per category. Z-score represents the relative proportion of genes with normalized expression higher or lower than the average (0) under copper conditions.

Figure S16. Gene ontology enrichment for potential genotype by environment eQTL. A. Genes with eQTL that were detected in head tissue only under copper conditions were enriched for categories related to metal ion transport and stress. B. Genes with eQTL that were detected in head tissue only under control conditions were enriched for categories related to cellular processes, response to stimulus, and biological regulation. The size of plotted circles represents

the relative number of genes per category. Z-score represents the relative proportion of genes with normalized expression higher or lower than the average (0).

Figure S17. Gene ontology enrichment for potential genotype by environment eQTL. A. Genes with eQTL that were detected in gut tissue only under copper conditions were enriched for categories related to lipid metabolism. B. Genes with eQTL that were detected in gut tissue only under control conditions were enriched for categories related localization and transport. The size of plotted circles represents the relative number of genes per category. Z-score represents the relative proportion of genes with normalized expression higher or lower than the average (0).

Figure S18. Pairwise correlations of gene expression or expression response of *trans* eQTL genes detected in the three Head datasets. The bottom lollipop plot summarizes mean absolute Pearson correlations for each set of *trans* eQTL genes.

Figure S19. Pairwise correlations of gene expression or expression response of *trans* eQTL genes detected in the three Gut datasets. The bottom lollipop plot summarizes mean absolute Pearson correlations for each set of *trans* eQTL genes.

Figure S20. Loadings for gene included in each *trans* eQTL hotspot that were highly correlated loaded evenly on PC1, which was used as a composite variable in assessment of potential mediators of *trans* eQTL hotspots.

Figure S21. Mediation analysis of Head hotspots implicated between zero and three potential candidate genes that may influence the expression or response of the *trans* eQTL genes corresponding to most hotspots. In all QTL maps above, the peak associated with variation in the composite variable representing the *trans* eQTL genes is shown in black. Genes that fell within hotspot intervals are highlighted in green; genes with partial or complete mediating effects on the composite QTL are shown in purple. In the regression plots, estimated haplotype effects for the composite peak and for the *cis* eQTL associated with the potential candidate genes are shown. The purple lines represent the best fit based on a linear model with 95% confidence intervals shaded in grey; statistics provide Pearson correlations and p values are presented without multiple test correction. The DSPR has eight founder strains. Haplotype estimates based on low representation of a founder strain at a given position were not included in the correlation analyses.

Figure S22. Mediation analysis of Gut hotspots implicated between one and five potential candidate genes that may influence the expression or response of the *trans* eQTL genes corresponding to most hotspots. In all QTL maps above, the peak associated with variation in the composite variable representing the *trans* eQTL genes is shown in black. Genes that fell within hotspot intervals are highlighted in green; genes with partial or complete mediating effects on the composite QTL are shown in purple. In the regression plots, estimated haplotype effects for the composite peak and for the *cis* eQTL associated with the potential candidate genes are shown. The purple lines represent the best fit based on a linear model with 95% confidence intervals shaded in grey; statistics provide Pearson correlations and p values are presented

without multiple test correction. The DSPR has eight founder strains. Haplotype estimates based on low representation of a founder strain at a given position were not included in the correlation analyses.

Supplemental Tables

Table S1. Samples from head and gut tissue were arrayed across four 96-well plates. Sample names are written as "AB_00000_CC" where "A" is tissue (H = head, G = gut), "B" is the level of resistance exhibited by the RIL in Everman et al. (2021) (R = resistant, S = susceptible), "00000" is the five digit RIL ID and "CC" is the treatment (Cu = exposed to copper sulfate, NA = water control). Plates 1 and 2 contained only head samples; plates 3 and 4 contained only gut samples. Copper and control-treated RIL pairs were kept together on tissue-specific plates. Each plate had an even representation of resistant and sensitive RILs. Copper samples (Cu) are shaded in blue; control samples (NA) are not shaded. Black text indicates resistant RILs; red text indicates sensitive RILs. Samples for which library prep failed are indicated with strikethrough text (HR_21180_Cu, HR_21180_NA, GR_21180_Cu, GR_21232_NA, GS_21281_NA).

Table S2. Principal components used in correction of datasets in preparation for eQTL mapping.