1
FoxP3 recognizes microsatellites and bridges DNA through multimerization

Wenxiang Zhang[1,2,5], Fangwei Leng[1,2,5], Xi Wang[1,2], Ricardo N. Ramirez[3,4], Jinseok Park[3], Christophe Benoist[3,4], and Sun Hur[1,2]*

[1]Howard Hughes Medical Institute and Program in Cellular and Molecular Medicine,

Boston Children's Hospital, MA 02115, USA

[2]Department of Biological Chemistry and Molecular Pharmacology

[3]Department of Immunology, Harvard Medical School, Boston, MA 02115, USA

[4]Evergrande Center for Immunologic Diseases, Harvard Medical School and

Brigham and Women's Hospital, Boston, MA 02115, USA

[5]These authors contributed equally.

*Correspondence: Sun.Hur@crystal.harvard.edu

Abstract

FoxP3 is a transcription factor (TF) essential for development of regulatory T cells (Tregs), a branch of T cells that suppress excessive inflammation and autoimmunity[1-5]. Molecular mechanisms of FoxP3, however, remain elusive. We here show that FoxP3 utilizes the Forkhead domain—a DNA binding domain (DBD) that is commonly thought to function as a monomer or dimer—to form a higher-order multimer upon binding to T_nG repeat microsatellites. A cryo-electron microscopy structure of FoxP3 in complex with T_3G repeats reveals a ladder-like architecture, where two double-stranded DNA molecules form the two "side rails" bridged by five pairs of FoxP3 molecules, with each pair forming a "rung". Each FoxP3 subunit occupies TGTTTGT within the repeats in the manner indistinguishable from that of FoxP3 bound to the Forkhead consensus motif (FKHM; TGTTTAC). Mutations in the "intra-rung" interface impair T_nG repeat recognition, DNA bridging and cellular functions of FoxP3, all without affecting FKHM binding. FoxP3 can tolerate variable "inter-rung" spacings, explaining its broad specificity for T_nG repeat-like sequences *in vivo* and *in vitro*. Both FoxP3 orthologs and paralogs show similar T_nG repeat recognition and DNA bridging. These findings thus reveal a new mode of DNA recognition that involves TF homo-multimerization and DNA bridging, and further implicates microsatellites in transcriptional regulation and diseases.

Introduction

How TFs utilize a limited repertoire of DBDs to orchestrate complex gene regulatory networks is a central and yet unresolved question[6-9]. While certain TFs, such as those with zinc-finger (ZF) DBDs, can expand the complexity of their sequence specificity by forming an array of DBDs, the vast majority of TFs utilize a single DBD with narrow sequence specificity shared with other members of the DBD family[7]. One prominent model to rationalize this apparent paradox is that cooperative actions of multiple distinct TFs with distinct DBDs give rise to combinatorial complexity[10,11]. However, whether a single TF with a single DBD can also recognize distinct sequences on its own and perform divergent transcriptional functions, depending on the conformation or multimerization state, has not been fully addressed.

FoxP3 is an essential TF in Tregs development, of which loss-of-function mutations cause a severe multiorgan autoimmune disease, immune dysregulation, polyendocrinopathy, enteropathy, X-linked (IPEX) syndrome[1-5]. Previous studies showed that FoxP3 remodels the global transcriptome and three-dimensional genome organization in the late stage of Treg development[12-15]. However, molecular mechanisms of FoxP3, including its direct target genes and *in vivo* sequence specificity, remain unclear[13-16].

FoxP3 DNA binding is primarily mediated by a Forkhead domain, which is shared among ~50 TFs of the Forkhead family[17,18]. Most Forkhead domains form a conserved winged-helix conformation and recognize the consensus FKHM sequence (TGTTTAC)[19]. While isolated Forkhead domain of FoxP3 was originally crystallized as an unusual domain-swap dimer[20,21], a recent study showed that FoxP3 does not form a domain-swap dimer but instead folds into the canonical winged-helix conformation in the presence of the adjacent Runx1-binding region (RBR)[22]. It was further shown that FoxP3 has a strong preference for inverted repeat FKHM (IR-FKHM) over a single FKHM *in vitro* by forming a head-to-head dimer[22]. However, previous chromatin immunoprecipitation sequencing (ChIP-seq)[14,23,24] and Cleavage Under Targets & Release Using Nuclease sequencing (CNR-seq)[14] analyses did not reveal enrichment of IR-FKHM in FoxP3-occupied genomic regions within cells[22]. While individual FKHM is present in ~10% of the FoxP3 ChIP peaks, they too may not be the FoxP3 binding sites as DNase I protection patterns at these sites were unaffected by FoxP3 deletion[24]. These observations raised the question of what sequences FoxP3 in fact recognizes in cells and whether FoxP3 can employ a previously unknown mode of binding to recognize new sequence motifs distinct from FKHM.

Results

FoxP3 binds T_nG repeat microsatellites

To re-evaluate FoxP3 sequence specificity, we performed an unbiased pull-down of genomic DNA with recombinant FoxP3 protein. Use of genomic DNA, as opposed to synthetic DNA oligos, enables testing sequence specificity in the context of naturally existing repertoire of sequences. It can also allow identification of longer motifs by using genomic DNA fragments longer than ~20-40 bp—the typical lengths used in previous biochemical studies of FoxP3[22,25,26]. We isolated genomic DNA from mouse EL4 cells, fragmented to ~100-200 bp, incubated with purified, MBP-tagged mouse FoxP3 and performed MBP pull-down, followed by next-

generation sequencing (NGS) of the co-purified DNA (FoxP3 PD-seq, Fig. 1a). We used
recombinant FoxP3 protein (FoxP3$^{\Delta N}$) harboring zinc finger (ZF), coiled coil (CC), RBR and
Forkhead domains but lacking the N-terminal Pro-rich region (Fig. 1a). FoxP3$^{\Delta N}$ was previously
shown to display the same DNA specificity as full-length FoxP3 among the test set[22]. *De novo*
motif analysis showed a strong enrichment of T_nG repeats (n=2-5) by FoxP3 pull-down, using
either MBP alone pull-down or input as a control (Fig. 1b, Supplementary Table 1a). T_3G repeat
sequence was the highest-ranking motif, which accounted for 49.8% of the peaks. No other
motifs, including the canonical FKHM or other repeats, were similarly enriched (Supplementary
Table 1a). FoxP3 pull-down using nucleosomal DNA from EL4 cells revealed similar
enrichment of T_nG repeat-like sequences (Supplementary Table 1a).

De novo motif analysis of previously published FoxP3 CNR-seq[12,14] and ChIP-seq data[14,23] also
identified T_nG repeat-like motifs as one of the most significant motifs in all four datasets (Fig.
1b, Supplementary Table 1b). The enrichment score for T_nG repeat-like motifs (E-value) was
more significant than that of FKHM in all cases (Fig. 1b, Supplementary Table 1b). Note that
T_nG repeat-like motifs have not been reported from these original studies, likely reflecting the
common practice of discarding simple repeats in motif analysis. T_nG repeat-like motifs were not
identified from open chromatin regions (as measured by ATAC-seq[27]) in Treg cells that were not
occupied by FoxP3 (Supplementary Table 1b).

To examine whether T_nG repeat-like sequences indeed contribute to FoxP3–DNA interaction in
Treg cells, we analyzed published FoxP3 CNR-seq data generated using F1 hybrids of B6 and
Cast mice strains[14]. Because of the wide divergence between B6 and Cast genomes, such data
allow evaluation of the impact of sequence variations on TF binding. Out of 196 sites showing
allelic imbalance (fold change >=4) in FoxP3 CNR-seq, 78 sites harbored T_nG repeat-like
elements in at least one allele, the frequency (39.3%) significantly higher than that in the mouse
genome (~0.06%, $p<$1e-8, Extended Data Fig. 1a). Furthermore, all but 4 sites showed T_nG
repeat length mirroring the allelic bias in FoxP3 occupancy (Fig. 1c, 1d). Of the 78 sites, we
randomly chose 50 sites, 25 each from B6- and Cast-biased peaks, and tested FoxP3 binding
efficiency by FoxP3$^{\Delta N}$ pull-down. Out of the 50 pairs of sequences tested, pull-down efficiency
of 47 pairs recapitulated differential binding in CNR-seq (Figs. 1c, 1e). All 47 sites showed
significant truncations in the T_nG repeats in the less-preferred allele (see Supplementary Table 2a
for the full list of sequences). Note that the pull-down preference for longer T_nG repeats was not
due to the different DNA lengths used—an extension of the less-preferred allele sequences with
a random sequence at a DNA end (B6* and Cast* in Fig. 1c, see Supplementary Table 2b for
sequence) did not alter the allele bias. Altogether, these results suggest that T_nG repeat-like
elements play an important role in FoxP3–DNA interaction *in vitro* and *in vivo*.

Genome-wide analysis showed that there are 46,574 loci in the *M. musculus* genome with T_nG
repeat-like sequences and that they are predominantly located distal to annotated transcription
start sites (TSSs) with 9.5% residing within 3 kb of the annotated TSSs (Extended Data Figs.
1a,1b). In comparison, among the T_nG repeat-containing FoxP3 CNR peaks[12,14] (n=3,301 out of
the 9,062 CNR peaks), 38.4% were found within 3 kb of TSSs (Extended Data Fig. 1c). T_nG
repeat-containing FoxP3 CNR peaks also displayed higher levels of trimethylated H3K4
(H3K4me3), acetylated H3K27 (H3K27ac) and chromatin accessibility than the genome-wide
T_nG repeats (Extended Data Figs. 1d, 1e, 1f). These results suggest that while T_nG repeat-like

sequences are common in the *M. musculus* genome, FoxP3 utilizes a small fraction of T_nG
repeat-like sequences in accessible, functional sites for transcriptional regulation.
133
FoxP3 multimerizes on T_nG repeats

To examine T_nG repeat enrichment in PD-seq and CNR/ChIP-seq indeed represent previously
unrecognized sequence specificity of FoxP3, we compared FoxP3 binding to DNA with T_nG
repeats (n=1 to 6) vs. those containing IR-FKHM, the highest affinity sequence reported for
FoxP3 to date[22]. All DNAs were 45 bp-long (see Supplementary Table 2b for sequences). We
found that T_3G repeat was comparable to IR-FKHM in FoxP3 binding and was the tightest
binder among the T_nG repeats (Fig. 1f), consistent with it being the most significant motif in PD-
seq (Fig. 1b). T_2G, T_4G and T_5G repeats also showed more efficient binding than a single FKHM
(1XFKHM) or random sequence (no FKHM). No other simple repeats showed FoxP3 binding
comparable to T_3G repeats (Fig. 1g). FoxP3 affinity increased with the copy number of T_3G,
when compared among DNAs of the same length (Fig. 1h). The preference for T_3G repeats was
also seen using full-length FoxP3 expressed in HEK293T cells (Extended Data Fig. 1g) or when
pull-down bait was switched from FoxP3 to DNA (Extended Data Fig. 1h). Finally, FoxP3 can
bind T_3G repeats even in the presence of nucleosomes (Extended Data Fig. 1i), suggesting that
similar interaction can occur in the context of chromatinized DNA.

We next investigated how FoxP3 recognizes T_3G repeats. Unlike IR-FKHM, T_3G-repeat DNA
induced FoxP3 multimerization as indicated by slowly migrating species in native gel-shift assay
(Fig. 1i). Protein-protein crosslinking also suggested higher-order multimerization in the
presence of T_3G repeats, but not with IR-FKHM or 1XFKHM (Extended Data Fig. 1j). In
support of T_3G repeat-induced multimerization, MBP-tagged FoxP3 was co-purified with GST-
tagged FoxP3 only in the presence of T_3G repeat, but not with IR-FKHM (Extended Data Fig.
1k). Finally, negative electron microscopy revealed a filamentous multimeric architecture of
FoxP3 on 36 tandem repeats of T_3G (Fig. 1j), the copy number chosen to aid clear visualization.
Other DNAs of the same length, such as $(A_3G)_{36}$, $(TGTG)_{36}$ or $(IR\text{-}FKHM)_5$, did not show
similar multimeric architectures under equivalent conditions (Fig. 1j, Extended Data Fig. 1l).
These results suggest that FoxP3 forms distinct multimers on T_3G repeats.

Structure of FoxP3 bound to T_3G repeats

To understand how FoxP3 forms multimers on T_3G repeats, we determined the cryo-electron
microscopy (cryo-EM) structure of FoxP3$^{\Delta N}$ in complex with $(T_3G)_{18}$. Single particle
reconstruction led to a 3.6 Å resolution map after global refinement and a 3.3 Å resolution map
after focused refinement of the central region (Extended Data Figs. 2a-2f, Extended Data Table
1). The density map revealed two continuous double-stranded DNA molecules spanning ~50 bp
(Fig. 2a). Both DNA molecules adopted the classic B-form DNA with the average twist angle of
33.5°/bp and the average rise of 3.19 Å/bp. The density map could also be fitted with the crystal
structure of DNA-bound FoxP3 monomer containing part of RBR and Forkhead (residue 326-
412), allowing placement of ten FoxP3 subunits without ZF, CC and RBR residues 188-325.
Only the non-swap, winged-helix conformation was compatible with the density map (Extended
Data Fig. 2g). Consistent with this, R337Q, a loss-of-function IPEX mutation that induces

domain-swap dimerization[22], showed significantly reduced affinity for T_3G repeats (Extended
Data Fig. 2h).
178
DNA sequence assignment (see Extended Data Fig. 3a) revealed that all ten FoxP3 subunits
interacted with T_3G repeat DNA in the manner indistinguishable from that of FoxP3 bound to the
canonical FKHM, recognizing TGTTTGT in place of TGTTTAC (Fig. 2b). This FoxP3-DNA
register was further confirmed by FoxP3 footprint analysis using DNA mutagenesis and NFAT–
FoxP3 cooperativity (Extended Data Figs. 3b, 3c). Note that NFAT is a known interaction
partner of FoxP3 and assists FoxP3 binding to DNA only when their binding sites are 3 bp apart,
the property used for inferring FoxP3–DNA registers (see Extended Data Fig. 3b for details).
186
The overall architecture resembled a ladder where the two double-stranded DNA molecules
formed side rails bridged by five "rungs", each of which consisted of two FoxP3 subunits bound
to different DNA and joined by direct protein-protein interactions ("intra-rung" interactions)
(Fig. 2a, 2c). These rungs were separated by 8 bp or 12 bp in an alternating fashion, forming two
different types of "inter-rung" interactions (inter-rung^{8bp} and inter-rung^{12bp}) with divergent
significance (to be discussed later). Given that both DNA molecules had the helical periodicity
of 10.7 bp per turn, this alternating spacing pattern allowed FoxP3 molecules to occupy
consecutive major grooves on one side of each DNA. This geometry, in turn, enabled the FoxP3
molecules on opposing DNA to face each other and form the "rungs" of the ladder. None of the
intra- and inter-rung interactions resembled the previously reported head-to-head dimerization
interaction (Extended Data Fig. 3d)[22], revealing a new mode of molecular assembly for FoxP3.
198
The two DNA molecules are skew to each other (i.e. non-parallel, non-intersecting). When
projected onto the x-y plane as in Fig. 2a, they appeared parallel, but projection onto the x-z
plane as in Fig. 2d suggested that they approached each other at an angle of 35˚. The divergence
of the two DNA molecules can explain why the multimeric assembly was limited to the decamer
spanning ~50 bp near the projected intersection point (Fig. 2d), even though the DNA sample in
cryo-EM was 72 bp-long and had many more T_3G repeats to accommodate additional FoxP3
molecules. The lack of cryo-EM density for FoxP3 molecules bound to DNA without forming
the "rung" suggests that the intra-rung interaction is critical for stable protein-DNA interaction.
In other words, DNA bridging may be an integral part of the assembly.
208
To test whether DNA bridging indeed occurs in solution, we examined co-purification of non-
biotinylated DNA (prey) with biotinylated DNA (bait) in the presence and absence of FoxP3. We
observed DNA bridging between biotinylated and non-biotinylated T_3G repeats only in the
presence of FoxP3$^{\Delta N}$ (Fig. 2e). DNA bridging was not observed between IR-FKHM and IR-
FKHM DNAs or between $(T_3G)_{12}$ and IR-FKHM DNAs. Similar T_nG repeat-dependent bridging
was observed with full-length FoxP3 expressed in 293T cells (Extended Data Fig. 3e).
Additionally, T_3G repeat DNA bridging occurred more efficiently with an increasing
concentration of FoxP3 (Extended Data Fig. 3f), suggesting that DNA bridging is not an artificial
consequence of saturating multimeric FoxP3 with DNA.
218
To further examine whether FoxP3 binding to T_nG repeats mediates long-distance chromatin
contacts in Tregs, we analyzed the available HiC-seq, PLAC-seq and HiChIP-seq data[12,13]. The
limited resolution of these data (5-10kb) precluded direct motif analysis of the chromatin contact

anchors. Instead, we asked how frequently contacts are made between anchors containing FoxP3
CNR peaks with T_nG repeats (T_nG anchors) vs. those lacking T_nG repeats (non-T_nG anchors).
Among the high-frequency contacts (HiC frequency>5, PLAC frequency>5-75) between FoxP3-
bound anchors, we found that those between two T_nG anchors (30-53%) were more prevalent
than expected by chance (13.7%) and that such T_nG–T_nG contacts were more enriched among
the stronger contacts (Fig. 2f, Supplementary Table 3, tabs 1-6). In contrast, non-T_nG–non-T_nG
contacts were more depleted among the stronger contacts. This is despite the fact that non-T_nG
CNR peaks have higher levels of chromatin accessibility and H3K4me3 than T_nG CNR peaks,
while displaying similar H3K27ac levels (Extended Data Figs. 4a-4c). Most of the T_nG–T_nG
contacts showed increased frequency in WT Tregs relative to in FoxP3 knock-out Treg-like cells
(Extended Data Fig. 4d). Furthermore, many of the anchors for the T_nG–T_nG contacts were
nearby Treg signature genes (e.g. *Il2ra, Cd28, Tnfaip3, Ets1* in Supplementary Table 3, tab 7),
and overlapped with previously characterized enhancer-promoter loop anchors in Tregs
(Extended Data Fig. 4e), implicating their transcriptional functions. These results together
support that FoxP3 multimerization on T_nG repeats contributes to long-distance chromatin
contacts in Tregs.
238
239 **Intra-rung interaction is essential**
240
Examination of the intra-rung interaction showed that multiple distinct parts of the protein are
involved; wing1 (W1), a loop between helix 2 and 4 (H2/H4 loop) and Helix6 (H6) of one
subunit interacted with RBR and H2/H4 of the other subunit within the rung (Fig. 3a). While the
resolution at the interface was insufficient to assign precise side chain conformations, the
structure identified R356 in the H2/H4 loop, V396 and V398 in W1, D409/E410/F411 in H6 as
residues at the interface (Fig. 3a). We also chose V408 in H6, which was adjacent to the interface
residues and is mutated to Met in a subset of IPEX patients[15,28,29]. Mutations of these interface
residues, including V408M, disrupted T_3G repeat binding (Fig. 3b, right) and DNA bridging
(Fig. 3c). The same mutations had a minimal impact on IR-FKHM binding (Fig. 3b, left), which
requires head-to-head dimerization of FoxP3[22]. This is consistent with the previous structure
showing that these residues are far from either the DNA binding or the head-to-head
dimerization interface[22]. The negative effect of the intra-rung mutations on T_3G repeat binding as
well as DNA bridging further supports that DNA bridging is required for FoxP3 multimerization
on T_3G repeats, rather than a simple consequence of FoxP3 multimerization.
255
These intra-rung mutations disrupted cellular functions of FoxP3, as measured by FoxP3-induced
gene expression (*e.g. CTLA4* and *CD25* protein levels in Fig 3d and RNA-seq in Extended Data
Fig. 5a), target loci binding (as measured by FoxP3 ChIP-seq in Extended Data Fig. 5b) and T
cell-suppressive functions (Fig. 3e). None of these mutations affected nuclear localization, the
level of FoxP3 (Extended Data Figs. 5c, 5d) or FoxP3's interaction with NFAT (Extended Data
Fig. 5e), although slight reduction in NFAT binding was seen for V398E. These results suggest
that the ladder-like assembly is important for the transcriptional functions of FoxP3.
263
264 **Relaxed sequence specificity of multimer**
265
We next examined the potential role of the inter-rung interactions. The inter-rung[8bp] interaction
was mediated by RBR–RBR contacts, which displayed continuous EM density indicative of a

strong or ordered interaction (Extended Data Fig. 2f, Fig. 4a). Unlike the intra-rung interface
mutations, mutations in RBR, for example F331D, disrupted FoxP3 binding to both T_3G repeats
and IR-FKHM[22] (Fig. 4b, Extended Data Fig. 6a), suggesting that RBR plays an important role
in both ladder-like multimerization and head-to-head dimerization[22]. Consistent with the
importance of the inter-rung^{8bp} interaction, changes in the inter-rung^{8bp} spacing from 8 bp (1 bp
gap) to 9 bp (2 bp gap) or 7 bp (no gap) resulted in a significant impairment in FoxP3 binding to
T_3G repeats (Fig. 4c).
275
In contrast to the inter-rung^{8bp} interaction, cryo-EM density for the inter-rung^{12bp} interaction was
difficult to interpret, which could reflect a weak or less-ordered interaction. In keeping with this,
FoxP3 binding tolerated a wide range of inter-rung^{12bp} spacings, with equivalent affinity
observed for spacings of 11-13 bp (Fig. 4d). Interestingly, while 14-19 bp spacings were not
tolerated, DNA with 21-22 bp spacings showed moderate binding. Given that 11-13 bp, 14-19 bp
and 21-22 bp spacings would place FoxP3 one, one and a half and two helical turns away from
the upstream FoxP3 molecule, respectively, this cyclical pattern suggests that the precise
positions of FoxP3 are not essential for multimeric assembly, so far as the DNA sequence allows
FoxP3 molecules to line up on one side of DNA and form the "rungs". Consistent with the idea,
DNA bridging activity showed a similar cyclical pattern (Extended Data Fig. 6b).
286
This architectural flexibility may explain our observations in Fig. 1, which showed that FoxP3
could bind to a broad range of T_nG repeat-like sequences besides perfect T_3G repeats. These
include tandem repeats of T_2G, T_4G, T_5G and their various mixtures found in the CNR-seq peaks
with allelic imbalance (Supplementary Table 2a). To examine whether a similar ladder-like
architecture forms with T_nG repeat-like sequences that are not perfect T_3G repeats, we used
DNA-bridging activity as a measure of the ladder-like assembly. All 47 pairs of the DNA
sequences showing allelic bias in FoxP3 binding *in vivo* and *in vitro* displayed the same allelic
bias in DNA bridging (Fig. 4e, Supplementary Table 2a). The multimerization-specific IPEX
mutation V408 abrogated bridging of T_2G, T_4G, T_5G repeat DNAs (Fig. 4f), suggesting a similar
multimeric architecture for FoxP3 regardless of the exact T_nG repeat sequences. Interestingly,
suboptimal T_nG repeats (n=2, 4, 5) were bridged with T_3G repeats more efficiently than with
themselves (Fig. 4g, Extended Data Fig. 6c), suggesting that having a strong DNA as a bridging
partner helps FoxP3 binding to suboptimal sequences. These results reveal yet another layer of
complexity that can broaden the sequence specificity of FoxP3.
301
302 **T_nG repeat binding is conserved in FoxPs**
303
Studies above were done with FoxP3 and T_nG repeat-like elements from *M. musculus*. We next
asked whether T_nG repeat recognition by FoxP3 is conserved in other species besides *M.*
musculus. Inspection of T_nG repeat-like elements in the *H. sapiens* and *D. rerio* genomes
revealed 18,164 and 5,517 distinct sites harboring T_nG repeats (>29 nt), respectively, in
comparison to the 46,574 sites in the *M. musculus* genome (Extended Data Fig. 1a). While T_nG-
like repeats are more frequently located distal to TSSs in all three genomes of *H. sapiens*, *M.*
musculus and *D. rerio*, greater fractions are located within ~3 kb of TSS in higher eukaryotes
(12.66%, 9.50% and 5.72% for *H. sapiens*, *M. musculus* and *D. rerio*, respectively) (Extended
Data Fig. 1b), even though all three species have similar genes-to-genome size ratios (see Table

in Extended Data Fig. 1a). This observation suggests that T_nG repeats may have been coopted for
transcriptional functions in higher eukaryotes.
315
We examined FoxP3 from *H. sapiens*, *O. anatinus* and *D. rerio*. All three FoxP3 orthologs
showed preferential binding to T_3G repeats and IR-FKHM in comparison to a single FKHM or
no FKHM (Extended Data Fig. 6d). They also bridged T_3G repeats (Extended Data Fig. 6e),
suggesting a ladder-like assembly similar to that of *M. musculus* FoxP3. This is in keeping with
the fact that the key residues for multimerization were broadly conserved or interchanged with
similar amino acids in FoxP3 orthologs (Extended Data Fig. 6f). Given that *D. rerio* FoxP3
represents one of the most distant orthologs from mammalian FoxP3, these results suggest that
T_nG repeat recognition and ladder-like assembly may be ancient properties of FoxP3.
324
Inspection of sequence alignment of Forkhead TFs revealed that the key residues for
multimerization are also well-conserved within the FoxP family, but not outside (Fig. 5a).
Biochemical analysis of *M. musculus* FoxP1, FoxP2 and FoxP4 in the FoxP family showed that
they preferentially bound T_3G repeats and bridged T_3G repeat DNA as with FoxP3 (Figs. 5b, 5c
and Extended Data Fig. 6g). *De novo* motif analysis of previously published ChIP-seq data
showed that T_nG repeat-like motifs were indeed enriched in FoxP1- and FoxP4-occupied sites
(Fig. 5d, the full list and references included in Supplementary Table 1c). This feature was
particularly strong for FoxP1 in lymphoma cell lines (SU-DHL-6 and U-2932) and mouse neural
stem cells—T_nG repeat-like motif was the most significant motif, while FKHM ranked far lower
(Fig. 5d). However, in VCap and K-562 cell lines, FoxP1 ChIP-seq peaks did not show T_nG-like
elements, although FKHM was identified as one of the most significant motifs in these cells
(Supplementary Table 1c). Similar context-dependent enrichment of T_nG repeat-like elements
was seen with FoxP4, although the motif enrichment was not as strong as with FoxP1 or FoxP3
(Fig. 5d, Supplementary Table 1c). In contrast, long (>10 nt) T_nG repeat-like elements were not
identified from any of the 48 distinct sets of ChIP-seq data for FoxA1, FoxM1, FoxJ2, FoxJ3,
FoxQ1 and FoxS1, while FKHM ranked as one of the strongest motifs in many (Supplementary
Table 1c). These results suggest that preference for T_nG repeat-like sequence and ladder-like
assembly are conserved properties of FoxP3 paralogs and orthologs, but may not be shared
among all Forkhead TFs.
344
345 **Discussion**
346
In summary, our findings showed a new mode of TF–DNA interaction that involves TF homo-
multimerization and DNA bridging. Upon binding to T_nG repeats, FoxP3 forms a ladder-like
multimer, in which FoxP3 utilizes two DNA molecules as scaffolds to facilitate cooperative
multimeric assembly. That is, the first set of FoxP3 molecules (possibly a dimer or two dimers
with an 8-bp spacing) that bridge DNA would help recruit additional FoxP3 "rungs", which
would in turn stabilize the bridged DNA architecture and subsequent rounds of FoxP3
recruitment. Such cooperative assembly enables FoxP3 to preferentially target long repeats of
T_nG rather than spurious sequences containing a few copies of T_nG. The DNA bridging activity
also implicates FoxP3 as a previously unrecognized class of TF that can directly mediate
architectural functions, which may explain the recently observed role of FoxP3 in chromatin loop
formation[12,13].
358

How do we reconcile the ladder-like assembly of FoxP3 on T_nG repeats and the previously reported head-to-head dimeric structure on IR-FKHM or related sequences[22]? Unlike the ladder-like multimerization, cellular evidence for the head-to-head dimerization is currently limited based on the available FoxP3 ChIP or CNR-seq data. Additionally, our new data showed that previously reported mutations that disrupt the head-to-head dimerization also affected the ladder-like multimerization, further limiting the ability to probe the physiological functions of the head-to-head dimerization. Nevertheless, given that head-to-head dimerization is unique to FoxP3, while the ladder-like multimerization is shared among all four FoxP TFs, we speculate that both forms exist in cells and carry out distinct functions depending on the bound DNA sequence. For example, DNA bridging would be a unique consequence of the ladder-like assembly, not shared with the head-to-head dimer, while the head-to-head dimerization may allow recruitment of certain cofactors using the unique surface created by the dimerization. This fits the previous microscopic analysis where FoxP3 were found in two distinct types of nuclear clusters associated with different cofactors[16]. Altogether, these findings suggest FoxP3 as a versatile TF that can interpret a wide range of sequences by assembling at least two distinct homo-multimeric structures.

Our findings also implicate functional roles of microsatellites in FoxP TF-mediated transcription regulation. While widely used as genetic tracing markers due to their high degrees of polymorphism, reports of biological functions of microsatellites[30,31], besides its well-known pathogenic roles[32-35], remain sparse[36,37]. Our finding of the T_nG repeat recognition by FoxP3 and other members of the FoxP family, raises the question of whether microsatellites play greater and more direct roles in transcriptional regulation than previously thought. This also prompts speculation that microsatellite polymorphism may contribute to a broad spectrum of diseases through FOXP TF dysregulation, such as autoimmunity via FOXP3, neurodevelopmental disorders via FOXP1, speech and language impairments via FOXP2, and heart and hearing defects via FOXP4.

Main text references

Brunkow, M. E. *et al.* Disruption of a new forkhead/winged-helix protein, scurfin, results in the fatal lymphoproliferative disorder of the scurfy mouse. *Nat Genet* **27**, 68-73, doi:10.1038/83784 (2001).

Bennett, C. L. *et al.* The immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome (IPEX) is caused by mutations of FOXP3. *Nat Genet* **27**, 20-21, doi:10.1038/83713 (2001).

Fontenot, J. D., Gavin, M. A. & Rudensky, A. Y. Foxp3 programs the development and function of CD4+CD25+ regulatory T cells. *Nat Immunol* **4**, 330-336, doi:10.1038/ni904 (2003).

Hori, S., Nomura, T. & Sakaguchi, S. Control of regulatory T cell development by the transcription factor Foxp3. *Science* **299**, 1057-1061, doi:10.1126/science.1079490 (2003).

Chatila, T. A. *et al.* JM2, encoding a fork head-related protein, is mutated in X-linked autoimmunity-allergic disregulation syndrome. *J Clin Invest* **106**, R75-81, doi:10.1172/JCI11679 (2000).

Badis, G. *et al.* Diversity and complexity in DNA recognition by transcription factors. *Science* **324**, 1720-1723, doi:10.1126/science.1162327 (2009).

7   Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* **172**, 650-665,
doi:10.1016/j.cell.2018.01.029 (2018).
8   Inukai, S., Kock, K. H. & Bulyk, M. L. Transcription factor-DNA binding: beyond
binding site motifs. *Curr Opin Genet Dev* **43**, 110-119, doi:10.1016/j.gde.2017.02.007
(2017).
9   Kribelbauer, J. F., Rastogi, C., Bussemaker, H. J. & Mann, R. S. Low-Affinity Binding
Sites and the Transcription Factor Specificity Paradox in Eukaryotes. *Annu Rev Cell Dev*
Biol **35**, 357-379, doi:10.1146/annurev-cellbio-100617-062719 (2019).
10   Reiter, F., Wienerroither, S. & Stark, A. Combinatorial function of transcription factors
and cofactors. *Curr Opin Genet Dev* **43**, 73-81, doi:10.1016/j.gde.2016.12.007 (2017).
11   Jolma, A. *et al.* DNA-dependent formation of transcription factor pairs alters their
binding specificity. *Nature* **527**, 384-388, doi:10.1038/nature15518 (2015).
12   Liu, Z., Lee, D. S., Liang, Y., Zheng, Y. & Dixon, J. Foxp3 Orchestrates Reorganization
of Chromatin Architecture to Establish Regulatory T Cell Identity. *bioRxiv*,
doi:10.1101/2023.02.22.529589 (2023).
13   Ramirez, R. N., Chowdhary, K., Leon, J., Mathis, D. & Benoist, C. FoxP3 associates with
enhancer-promoter loops to regulate T(reg)-specific gene expression. *Sci Immunol* **7**,
eabj9836, doi:10.1126/sciimmunol.abj9836 (2022).
14   van der Veeken, J. *et al.* The Transcription Factor Foxp3 Shapes Regulatory T Cell
Identity by Tuning the Activity of trans-Acting Intermediaries. *Immunity* **53**, 971-984
e975, doi:10.1016/j.immuni.2020.10.010 (2020).
15   Zemmour, D. *et al.* Single-cell analysis of FOXP3 deficiencies in humans and mice
unmasks intrinsic and extrinsic CD4(+) T cell perturbations. *Nat Immunol* **22**, 607-619,
doi:10.1038/s41590-021-00910-8 (2021).
16   Kwon, H.-K., Chen, H.-M., Mathis, D. & Benoist, C. Different molecular complexes that
mediate transcriptional induction and repression by FoxP3. *Nature Immunology* **18**, 1238-
1248, doi:10.1038/ni.3835 (2017).
17   Hannenhalli, S. & Kaestner, K. H. The evolution of Fox genes and their role in
development and disease. *Nat Rev Genet* **10**, 233-240, doi:10.1038/nrg2523 (2009).
18   Benayoun, B. A., Caburet, S. & Veitia, R. A. Forkhead transcription factors: key players
in health and disease. *Trends Genet* **27**, 224-232, doi:10.1016/j.tig.2011.03.003 (2011).
19   Dai, S., Qu, L., Li, J. & Chen, Y. Toward a mechanistic understanding of DNA binding
by forkhead transcription factors and its perturbation by pathogenic mutations. *Nucleic*
Acids Res **49**, 10235-10249, doi:10.1093/nar/gkab807 (2021).
20   Bandukwala, Hozefa S. *et al.* Structure of a Domain-Swapped FOXP3 Dimer on DNA
and Its Function in Regulatory T Cells. *Immunity* **34**, 479-491,
doi:10.1016/j.immuni.2011.02.017 (2011).
21   Chen, Y. *et al.* DNA binding by FOXP3 domain-swapped dimer suggests mechanisms of
long-range chromosomal interactions. *Nucleic Acids Research* **43**, 1268-1282,
doi:10.1093/nar/gku1373 (2015).
22   Leng, F. *et al.* The transcription factor FoxP3 can fold into two dimerization states with
divergent implications for regulatory T cell function and immune homeostasis. *Immunity*
55, 1354-1369 e1358, doi:10.1016/j.immuni.2022.07.002 (2022).
23   Kitagawa, Y. *et al.* Guidance of regulatory T cell development by Satb1-dependent
super-enhancer establishment. *Nat Immunol* **18**, 173-183, doi:10.1038/ni.3646 (2017).

24   Samstein, R. M. *et al.* Foxp3 exploits a pre-existent enhancer landscape for regulatory T
cell lineage specification. *Cell* **151**, 153-166, doi:10.1016/j.cell.2012.06.053 (2012).
25   Jolma, A. *et al.* DNA-binding specificities of human transcription factors. *Cell* **152**, 327-
339, doi:10.1016/j.cell.2012.12.009 (2013).
26   Koh, K. P., Sundrud, M. S. & Rao, A. Domain requirements and sequence specificity of
DNA binding for the forkhead transcription factor FOXP3. *PLoS One* **4**, e8109,
doi:10.1371/journal.pone.0008109 (2009).
27   Yoshida, H. *et al.* The cis-Regulatory Atlas of the Mouse Immune System. *Cell* **176**, 897-
912 e820, doi:10.1016/j.cell.2018.12.036 (2019).
28   Rubio-Cabezas, O. *et al.* Clinical heterogeneity in patients with FOXP3 mutations
presenting with permanent neonatal diabetes. *Diabetes Care* **32**, 111-116,
doi:10.2337/dc08-1188 (2009).
29   Consonni, F., Ciullini Mannurita, S. & Gambineri, E. Atypical Presentations of IPEX:
Expect the Unexpected. *Front Pediatr* **9**, 643094, doi:10.3389/fped.2021.643094 (2021).
30   Ibrahim, A. *et al.* MeCP2 is a microsatellite binding protein that protects CA repeats from
nucleosome invasion. *Science* **372**, doi:10.1126/science.abd5581 (2021).
31   Contente, A., Dittmer, A., Koch, M. C., Roth, J. & Dobbelstein, M. A polymorphic
microsatellite that mediates induction of PIG3 by p53. *Nat Genet* **30**, 315-320,
doi:10.1038/ng836 (2002).
32   Li, K., Luo, H., Huang, L., Luo, H. & Zhu, X. Microsatellite instability: a review of what
the oncologist should know. *Cancer Cell Int* **20**, 16, doi:10.1186/s12935-019-1091-8
(2020).
33   Pecina-Slaus, N., Kafka, A., Salamon, I. & Bukovac, A. Mismatch Repair Pathway,
Genome Stability and Cancer. *Front Mol Biosci* **7**, 122, doi:10.3389/fmolb.2020.00122
(2020).
34   Bonneville, R. *et al.* Landscape of Microsatellite Instability Across 39 Cancer Types.
JCO Precis Oncol **2017**, doi:10.1200/PO.17.00073 (2017).
35   Kloor, M. & von Knebel Doeberitz, M. The Immune Biology of Microsatellite-Unstable
Cancer. *Trends Cancer* **2**, 121-133, doi:10.1016/j.trecan.2016.02.004 (2016).
36   Bagshaw, A. T. M. Functional Mechanisms of Microsatellite DNA in Eukaryotic
Genomes. *Genome Biol Evol* **9**, 2428-2443, doi:10.1093/gbe/evx164 (2017).
37   Gharesouran, J., Hosseinzadeh, H., Ghafouri-Fard, S., Taheri, M. & Rezazadeh, M.
STRs: Ancient Architectures of the Genome beyond the Sequence. *J Mol Neurosci* **71**,
2441-2455, doi:10.1007/s12031-021-01850-6 (2021).
38   Braccioli, L. *et al.* FOXP1 Promotes Embryonic Neural Stem Cell Differentiation by
Repressing Jagged1 Expression. *Stem Cell Reports* **9**, 1530-1545,
doi:10.1016/j.stemcr.2017.10.012 (2017).
39   Consortium, E. P. An integrated encyclopedia of DNA elements in the human genome.
Nature **489**, 57-74, doi:10.1038/nature11247 (2012).

489
490

Main figure legends
Figure 1. FoxP3 recognizes T_nG repeat microsatellites.
    a. FoxP3 domain architecture and schematic of FoxP3 pull-down (PD)-seq. ZF: zinc finger;
        CC: coiled-coil; RBR: Runx1-binding region.

b. *De novo* motif analysis of FoxP3 PD-seq peaks (n=21,605) and CNR-seq peaks[14]
(n=6,655) by MEME-ChIP and STREME. E-score and percentage of peaks containing
the given motif are shown on the right. See also Supplementary Table 1a and 1b for the
comprehensive list of motifs for PD-seq, CNR-seq[12,14] and ChIP-seq data[14,23].
c. Allelic imbalance in FoxP3 binding *in vivo*. Left: genome browser view of CNR-seq[14]
showing B6-biased (top) and Cast-biased (bottom) peaks. B6 genomic coordinates are
shown on the upper left corner. Right: B6 and Cast DNA oligos were mixed 1:1 and
subjected to FoxP3$^{\Delta N}$ pull-down and gel analysis. Cast* and B6* represent oligos
extended with the random sequence (Supplementary Table 2b) to reverse their length
bias.
d. T_nG repeat-length comparison between Cast and B6 at 78 loci showing allelic bias in (c).
Repeat lengths were measured in nucleotide. n = 25 for Cast-biased loci and n = 25 for
B6-biased loci were used for this comparison. Two-tailed unpaired t-tests. ****, $p <$
0.0001.
e. Allelic imbalance in FoxP3 binding *in vitro*. 50 pairs of Cast and B6 sequences
(Supplementary Table 2a) were chosen from the 78 pairs in (d) and subjected to FoxP3$^{\Delta N}$
pull-down. For each pair, the recovery rate of the Cast and B6 DNA were measured and
their ratios were plotted. Each data point represents an average of the two pull-downs.
Two-tailed unpaired t-tests. ****, $p < 0.0001$.
f-h. FoxP3–DNA interaction as measured by FoxP3$^{\Delta N}$ pull-down. DNA harboring a random
sequence (no FKHM), a single FKHM (1XFKHM), IR-FKHM or tandem repeats of T_nG
(n=1-6) were used. All DNAs were 45 bp-long in (f) and (h), and 30 bp-long in (g).
i. Native gel shift assay of MBP-tagged FoxP3$^{\Delta N}$ (0-0.4 μM) with DNA (30 bp, 0.05 μM)
containing IR-FKHM or $(T_3G)_6$.
j. Representative negative-stain EM images of FoxP3$^{\Delta N}$ in complex with $(T_3G)_{36}$ and (IR-
FKHM)$_5$. Both DNAs were 144 bp-long.
521

522 Figure 2. FoxP3 forms a ladder-like multimer upon binding to T_3G repeat DNA.
a. Cryo-EM structure of FoxP3$^{\Delta N}$ decamer in complex with two DNA molecules (grey)
harboring $(T_3G)_{18}$. Each of the ten FoxP3 subunits are colored differently.
b. Comparison of a representative FoxP3$^{\Delta N}$ subunit from (a) (orange) with a FoxP3$^{\Delta N}$
subunit from the head-to-head dimeric structure (grey, PDB: 7TDX). Helix3 recognizes
DNA sequence (TGTTTAC in the head-to-head dimer, TGTTTGT in the ladder-like
multimer) by inserting it into the major groove.
c. Schematic of the ladder-like architecture of FoxP3 on T_3G repeat DNA.
d. Skew relationship between the two DNA molecules, which is evident when looking down
the y-axis of (a).
e. DNA bridging assay. Biotinylated DNA (bio-DNA, 82 bp) and non-biotinylated DNA
(non-bio-DNA, 60 bp) were mixed at 1:1 ratio (0.1 μM each), incubated with FoxP3$^{\Delta N}$
(0.4 μM), and subjected to streptavidin pull-down prior to gel analysis. Non-biotinylated
DNA in the eluate was visualized by SybrGold staining.
f. Chromatin contacts at FoxP3-bound anchors identified from HiC- and PLAC-seq[12].
Contacts with frequency>5 in WT Treg HiC and connected by two FoxP3-bound anchors
were analyzed with an increasing FoxP3 PLAC-seq count threshold. Percentage of the
unique contacts mediated by two T_nG anchors (out of all unique contacts between two
FoxP3-bound anchors) was indicated. All T_nG-T_nG contacts were between two distinct

10 kb anchor bins. **** indicates $p<0.0001$ using the null hypothesis that the T_nG-T_nG
contacts occur by random chance.

Figure 3. Intra-rung interaction is essential for T_nG repeat recognition, DNA bridging, and
cellular functions of FoxP3.

    a. Intra-rung interface. Ca of R356, V396, V398, V408 and D409/E410/F411 are shown in
545   spheres. These residues on yellow subunit interact with RBR and helix2-helix4 (H2/4) of
546   the orange subunit. Subunit colors are as in Fig. 2a.
547

b. Effect of intra-rung interface mutations on DNA binding. MBP-tagged FoxP3$^{\Delta N}$ (0.4 μM)
was incubated with IR-FKHM or $(T_3G)_{12}$ (60 bp for both) and analyzed by native gel
shift assay.

c. Effect of intra-rung interface mutations on DNA bridging. FoxP3 (or empty vector, EV)
was expressed in 293T cells and lysate was incubated with a mixture of biotinylated and
non-biotinylated DNA (1:1 ratio), followed by streptavidin pull-down and gel analysis.
Relative level of non-biotinylated DNA co-purified with biotinylated DNA was
quantitated from three independent pull-downs. The difference was compared with WT in
the presence of bio-DNA. Two-tailed paired t-tests. $p<0.001$ for *** and $p<0.005$ for **.

d. Transcriptional activity of FoxP3. CD4+ T cells were retrovirally transduced to express
FoxP3, and its transcriptional activity was analyzed by measuring the level of the known
target genes CTLA4 and CD25 using FACS. FoxP3 level was measured using Thy 1.1,
which is under the control of IRES from the bicistronic mRNA for FoxP3. MFI indicates
mean fluorescence intensity.

e. T cell suppression assay of intra-rung interface mutations. FoxP3-transduced T cells
(suppressors) were mixed with naïve T cells (responders) at a 1:2 ratio and the effect of
the suppressor cells on proliferation of the responder cells were measured by the CFSE
dilution profile of the responder T cells.

Figure 4. Architectural flexibility of the ladder-like assembly broadens sequence specificity
of FoxP3.

a. Structure highlighting the inter-rung^{8bp} interaction between orange and red subunits (in
surface representation). The interaction is primarily between RBRs, where F331 resides.

b. Effect of inter-rung^{8bp} mutation (F331D) on FoxP3–DNA interaction by FoxP3$^{\Delta N}$ pull-
down.

c. Effect of inter-rung^{8bp} spacings on FoxP3 binding by FoxP3$^{\Delta N}$ pull-down. Both inter-
rung^{8bp} gap nucleotides were changed from T in $(T_3G)_{10}$ to A (8-bp spacing), to AC (9-bp
spacing) or to no nucleotide (7-bp spacing). Black arrows indicate FoxP3 footprints. Gray
arrow indicates the NFAT binding site. NFAT interacts with FoxP3 and helps fixing the
FoxP3-DNA register, which was necessary to examine the effect of DNA sequence
variations at or between the FoxP3 footprints.

d. Effect of inter-rung^{12bp} spacings on FoxP3 binding by FoxP3$^{\Delta N}$ pull-down. The inter-
rung^{12bp} spacing was changed from 12 bp in $(T_3G)_{10}$ to 10-23 bp (see Supplementary
Table 2b for sequences). Averaged recovery rate of DNA from three independent pull-
downs were plotted. Two-tailed paired t-tests in comparison to $(T_3G)_{10}$. $p<0.005$ for **,
$p<0.05$ for * and $p>0.05$ for ns.

e. Comparison of Cast and B6 sequences in FoxP3 binding (left) and DNA bridging (right).
Three pairs of sequences at loci CN53, CN118 and CN16 with Cast-bias in CNR-seq
were compared (Supplementary Table 2a) by FoxP3$^{\Delta N}$ pull-down.

f. DNA bridging between $(T_2G)_{14}$ and $(T_2G)_{14}$, between $(T_4G)_9$ and $(T_4G)_9$, and between
$(T_5G)_7$ and $(T_5G)_7$ in the presence of WT FoxP3 or the IPEX mutant V408M.
Biotinylated and non-biotinylated DNA were colored red and blue, respectively.
g. DNA bridging between $(T_2G)_{14}$ and $(T_2G)_{14}$, and between $(T_2G)_{14}$ and $(T_3G)_{11}$ by FoxP3
(0-0.4 μM).
592

Figure 5. T_nG microsatellite recognition is conserved among FoxP3 orthologs and paralogs.
a. Sequence alignment of forkhead TFs. Residues equivalent to the key interface residues in
mouse FoxP3 (arrow on top) were highlighted yellow (when similar to the mouse
residues) or green (when dissimilar).
b. DNA-binding activity of FoxP1, FoxP2 and FoxP4. HA-tagged FoxP1, FoxP2 and FoxP4
were transiently expressed in HEK293T cells and purified by anti-HA IP. Equivalent
amounts of indicated DNAs (all 45 bp) were added to FoxP1/2/4-bound beads and further
purified prior to gel analysis (SybrGold).
c. DNA-bridging activity of FoxP1, FoxP2 and FoxP4. Experiments were performed as in
Fig. 3c using 293T lysate expressing HA-tagged FoxP TFs.
d. *De novo* motif analysis of FoxP1 and FoxP4 ChIP-seq peaks from published database
[38,39]. See Supplementary Table 1c for the comprehensive list and their references.
605

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mice

C57BL/6N mice, sourced from Taconic Biosciences and overseen by Harvard Medical Area
(HMA) Standing Committee on Animals, were housed in an individually ventilated cage system
at the specific-pathogen-free New Research Building facility of Harvard Medical School. The
mice were maintained at a controlled environment with a temperature of 20-22°C, humidity
ranging from 40-55%, and a 12-hour light-dark cycle. The spleens of 12~14 weeks old female
C57BL/6 mice were isolated for the study.

Naive CD4+ T Cells

Cells were isolated by using Naive $CD4^+$ T Cell Isolation Kit (Miltenyi Biotec, Cat#130-104-
453) according to the manufacturer's instructions and maintained in complete RPMI medium
(10% FBS heat-inactivated, 2mM L-Glutamine, 1mM Sodium Pyruvate, 100μM NEAA, 5mM
HEPES, 0.05mM 2-ME).

HEK293T & A549 cells

HEK293T cells (were purchased from ATCC (CRL-11268)) and A549 cells (kind gift from Dr.
Susan Weiss (U Penn)) were maintained in DMEM (High glucose, L-glutamine, Pyruvate) with
10% fetal bovine serum, 1% penicillin/streptomycin.

EL4 cells

Cells (gift from Dr. Christophe Benoist lab (Harvard Medical School)) were cultured in DMEM
(High glucose, L-glutamine, Pyruvate) supplemented with 10% fetal bovine serum, ranging from
$1X10^5$ to $1X10^6$ cells/mL.

METHOD DETAILS

Plasmids

Mouse FoxP3 plasmids were made as previously described[22]. For Mammalian expression
plasmids, HA-tagged mouse FoxP3 CDS was inserted into pcDNA3.1+ vector between KpnI and
BamHI sites. All FoxP3 mutations including R356E, V396E, V398E, V408M and 409-411AAA
were generated by site-directed mutagenesis using Phusion High Fidelity (New England Biolabs)
DNA polymerases. For retroviral packaging plasmids, HA-tagged mouse FOXP3 CDS was
inserted into MSCV-IRES-Thy1.1 vector.

For mammalian expression plasmids of FoxP3 orthologs from *H. sapiens, O. anatinus and D.*
rerio, respective FoxP3 CDS with overhangs of pcDNA vector was synthesized by IDTDNA and
then assembled with NEBuilder® HiFi DNA Assembly Cloning Kit (NEB, #5520G). FoxP3
paralogs (FoxP1, FoxP2 and FoxP4) mammalian expression plasmids were made in the same
way. Other forkhead TFs, such as FoxA1, FoxM1, FoxQ1 and FoxS1, were gifts from Stefan
Koch lab[40] through Addgene.

DNA oligos

Single-stranded DNA (ssDNA) oligos were synthesized by IDTDNA. Double-stranded DNA
(dsDNA) oligos for EMSA assay, pulldown assay and DNA bridging assay were annealed from
single-stranded, complementary oligos. After briefly spinning down each oligonucleotide pellet,
ssDNAs were dissolved in the annealing buffer (10 mM Tris-HCl pH 7.5, 50 mM NaCl).
Complementary ssDNAs were then mixed together in equal molar amounts, heated to 94°C for 2
minutes and gradually cooled down to room temperature. For dsDNA in cryo-EM analysis,
HPLC purified single-stranded, complementary oligos were purchased from IDTDNA. After
annealing, dsDNA was further purified by size-exclusion chromatography on Superdex 75
Increase 10/300 (GE Healthcare) columns in 20 mM Tris-HCl pH 7.5, 150 mM NaCl. Biotin
labeled ssDNA oligos were synthesized by IDTDNA and then dissolved in annealing buffer (10
mM Tris-HCl pH 7.5, 50 mM NaCl). Complementary, biotin-labeled ssDNAs were then mixed
together in equal molar amounts, heated to 94°C for 2 minutes and gradually cooled down to
room temperature. Sequence of all the DNA oligos used are shown in Supplementary Table 2b.

Protein expression and purification

All recombinant proteins in this paper were expressed in BL21(DE3) at 18°C for 16-20 hrs
following induction with 0.2 mM IPTG. Cells were lysed by high-pressure homogenization using
an Emulsiflex C3 (Avestin). All proteins are from the *Mus. musculus* sequence, unless mentioned
otherwise. FoxP3$^{\Delta N}$ (residues 188-423) was expressed as a fusion protein with an N-terminal
His$_6$-NusA tag. After purification using Ni-NTA agarose, the protein was treated with HRV3C

protease to cleave the His$_6$-NusA-tag and were further purified by a series of chromatography
purification using HiTrap Heparin (GE Healthcare), Hitrip SP (GE Healthcare) and Superdex
200 Increase 10/300 (GE Healthcare) columns. The final size-exclusion chromatography (SEC)
was done in 20 mM Tris-HCl pH 7.5, 500 mM NaCl, 2 mM DTT. NFAT1 protein (residues 394-
680) was also expressed as a fusion protein with an N-terminal His$_6$-NusA tag. After purification
using Ni-NTA agarose, the His$_6$-NusA-tag was removed using the HRV3C protease and was
further purified by SEC on Superdex 75 Increase 10/300 (GE Healthcare) column in 20 mM
Tris-HCl pH 7.5, 500 mM NaCl, 5% Glycerol, 2 mM DTT. His$_6$-MBP fused FoxP3$^{\Delta N}$ variants
were purified using Ni-NTA affinity column and Superdex 200 Increase 10/300 (GE Healthcare)
SEC column in 20 mM Tris-HCl pH 7.5, 500 mM NaCl, 2 mM DTT.

MBP-FoxP3$^{\Delta N}$ pulldown-seq (FoxP3 PD-seq)

Mouse EL4 genomic DNA was isolated by QIAGEN Blood & Cell Culture DNA Kit (QIAGEN,
#13343). The purified genomic DNA was then fragmented to 100~200 bp using DNase I
(ZYMO RESEARCH, #E1010) in the digestion buffer (50 mM NaCl, 20 mM Tris-HCl PH 7.5,
1.5 mM MgCl$_2$) (for 200μL system, 50μg genomic DNA was treated with 8μL DNase I for
3~4mins to obtain 100~200 bp DNA fragments). The digested genomic DNA was then purified
by QIAquick Nucleotide Removal Kit (QIAGEN, #28306) and used as an input for the
pulldown-seq.

Purified MBP-tag or MBP-FoxP3$^{\Delta N}$ protein was incubated with the input DNA fragments in the
incubation buffer (20 mM Tris-HCl pH 7.5, 100 mM NaCl, 1.5 mM MgCl$_2$) for 20 mins at room
temperature and then subjected to MBP pulldown using Amylose Resin (New England Biolabs).
The bound DNA was recovered using proteinase K (New England Biolabs) and purified using
QIAquick Nucleotide Removal kit (QIAGEN). The sequencing libraries were made using
NEBNext® Ultra™ II DNA Library Prep Kit (Illumina) according to the manufacturer's
instructions and submitted to Novogene for NGS with PE150.

Nucleosome pulldown-seq

Mouse EL4 cells were lysed using a hypotonic buffer (20mM Bis-Tris pH 7.5, 0.05% NP-40,
1.5mM MgCl$_2$, 10mM KCL, 5mM EDTA, 1Xmammalian protease inhibitor) and the nuclear
fraction was isolated by spinning down at 4° and 2,500 RPM for 10 mins. The isolated nuclear
fraction was then digested with Micrococcal Nuclease (Thermo Scientific, #88216) for 1 hour at
4°C to fragment the chromatin into individual nucleosomes. Lysate was then spun down at 4°
and 13,000 RPM for 10 mins. Cleared lysate containing the nucleosomes was incubated with
purified MBP-tag or MBP-FoxP3$^{\Delta N}$ protein (1uM) for 1 hour at 4°C and then subjected to MBP
pulldown using Amylose Resin (New England Biolabs). After proteinase K (New England
Biolabs) treatment, the final nucleosomal DNAs were recovered by QIAquick Nucleotide
Removal kit (QIAGEN) and used for library preparation. The libraries were made by NEBNext®
Ultra™ II DNA Library Prep Kit (Illumina) according to the manufacturer's instructions and
submitted to Novogene for NGS with PE150.

MBP-FoxP3$^{\Delta N}$ pulldown assay

0.4 μM purified MBP-mFoxP3$^{\Delta N}$ protein was incubated with 0.1 μM DNA in the incubation
buffer for 20 mins. The FoxP3-DNA mixture was then incubated with 25μL Amylose Resin
(New England Biolabs) for 30 mins with rotation at room temperature. The bound DNA was
recovered using proteinase K (New England Biolabs), purified using QIAquick Nucleotide
Removal kit (QIAGEN) and analyzed on 10% Novex TBE Gels (Invitrogen). DNA was
visualized by Sybr Gold staining. The expression of MBP-FoxP3$^{\Delta N}$ was validated by Western
blot using Mouse MBP Tag antibody(8G1) (Cell Signaling Technology, Cat#2396, 1:2000
dilution).

HA-FoxP3 pulldown assay

HEK293T cells were transfected with pcDNA encoding HA-tagged FoxP3 (wild-type or
mutants). After 48 hours, cells were lysed using RIPA buffer (10mM Tris-HCl pH 8.0, 1mM
EDTA, 1% Triton X-100, 0.1% Sodium Deoxycholate, 0.1% SDS, 140 mM NaCl and 1X
proteinase inhibitor) and treated with Benzonase (Millipore) for 30 mins. The lysate was then
incubated with Anti-HA Magnetic Beads (Thermo Fisher) for 1 hour. Beads were washed three
times using RIPA buffer and incubated with DNA oligos for 20 mins at room temperature.
Bound DNA was recovered using proteinase K (New England Biolabs), purified using QIAquick
Nucleotide Removal kit (QIAGEN) and analyzed on 10% Novex TBE Gels (Invitrogen). DNA
was visualized by Sybr Gold staining.

Nucleosome reconstitution and EMSA analysis

Nucleosome core particles (NCP) were reconstituted with recombinant Histone Octamer H3.1
(Active motif) and DNAs as described previously[41]. Briefly, 1 μM of TTTG repeats (144bp),
AAAG repeats (144bp), TGTG repeats (144bp) and DNA harboring 601 sequence (181bp) were
incubated with 1 μM of the histone octamer and were dialyzed against 10mM Tris-HCl PH 7.5,
1mM EDTA, 2mM DTT for 24 hrs. Nucleosomes (0.05 μM) were incubated with the indicated
amount of FoxP3$^{\Delta N}$ in the buffer (10mM Tris-HCl pH 7.5, 50mM NaCl, 1mM EDTA, and 2mM
DTT) for 30 min at 4 °C and analyzed on 6% TBE gels (Life Technologies) at 4 °C. After
staining with Sybr Gold stain (Life Technologies), Sybr Gold fluorescence was recorded using
iBright FL1000 (Invitrogen) and analyzed with iBright Analysis Software.

Biotin-DNA pulldown assay

HA-FoxP3 was transiently expressed in HEK293T cells as above. Cells were lysed using RIPA
buffer. The lysate was incubated with biotin-dsDNA (1μM) for 1 hour, and then with
streptavidin agarose beads (Thermo Fisher, 25μL) for additional 30mins. Beads were spun down
and washed three times with RIPA buffer. Bead-bound protein was extracted using the SDS
loading buffer and analyzed by SDS-PAGE and western blotting using anti-HA (Primary)
antibody (Cell Signaling, Cat#3724S, 1:3000 dilution) and anti-rabbit IgG-HRP (Secondary)
antibody (Cell Signaling, Cat#7074, 1:5000 dilution).

Electrophoretic Mobility Shift Assay (EMSA)

738 DNA (0.05μM) was mixed with the indicated amount of FoxP3 in buffer A (20mM HEPES pH
739 7.5, 150mM NaCl, 1.5mM $MgCl_2$ and 2mM DTT), incubated for 30 min at 4 °C and analyzed on
740 3-12% gradient Bis-Tris native gels (Life Technologies) at 4 °C. After staining with Sybr Gold
741 stain (Life Technologies), Sybr Gold fluorescence was recorded using iBright FL1000
742 (Invitrogen) and analyzed with iBright Analysis Software.

Crosslinking analysis

744 Protein-protein crosslinking using BMOE (Thermo Scientific) was carried out according to the
745 product manual. Briefly, 0.4μM FoxP3 $^{\Delta N}$ was incubated with 0.05μM DNAs at 25°C for 10
746 mins in 1X PBS, then BMOE was added to the final concentration of 100 μM. After 1 hour
747 incubation at 25°C, DTT (10mM) was added to quench the crosslinking reaction. Samples were
748 then analyzed by SDS-PAGE and Krypton staining (Thermo Scientific).

DNA bridging assay

Biotin-DNA (bait, 0.1μM) was incubated with Streptavidin Agarose (25μL, Thermo Fisher) in
buffer B (20mM Tris-HCl pH 7.5, 100mM NaCl, 1.5mM $MgCl_2$, 5mM DTT) for 30 mins by
rotating the mixture at room temperature. Agarose beads were washed three times with buffer B
and incubated with non-biotinylated DNA (prey, 0.1μM) and purified FoxP3 protein (or 293T
lysate expressing FoxP3). After incubation for 30 mins with rotation, bead-bound DNA was
recovered using proteinase K (New England Biolabs), purified by QIAquick Nucleotide
Removal kit (QIAGEN) and analyzed on 10% Novex TBE Gels (Invitrogen). DNA was
visualized by Sybr Gold staining.

Cryo-EM sample preparation and data collection

FoxP3$^{\Delta N}$ was incubated with $(T_3G)_{18}$ DNA at a molar ratio of 8:1 in buffer B at RT for 10 mins.
The complex was then crosslinked using 0.5 % glutaraldehyde for 10 mins at RT prior to
quenching with 1/10 volume of 1M Tris-HCl pH 7.5 (for a final Tris concentration of 0.1M). The
FoxP3$^{\Delta N}$-DNA complex was then purified by Superose 6 Increase 10/300 GL (GE Healthcare)
column in 20mM Tris-HCl pH 7.5, 100mM NaCl, 2mM DTT. The sample was concentrated to 1
mg/ml (final for protein) and applied to freshly glow-discharged C-flat 300 mesh copper grids
(CF-1.2/1.3, Electron Microscopy Sciences) at 4°C. The grids were plunged into liquid ethane
after blotting for 5 s using Vitrobot Mark IV (FEI) at the humidity setting of 100%. The grids
were screened at the Harvard Cryo-EM Center and UMass Cryo-EM core facility using Talos
Arctica microscope (FEI). The grids that showed a good sample distribution and ice thickness
were used for data collection on Titan Krios (Janelia Cryo-EM facility) operated at 300 kV and
equipped with Gatan K3 camera. A total of 11624 micrographs were taken at a magnification of
81,000x with a pixel size of 0.844 Å. Each movie comprised of 60 frames at total dose of 60 e$^-$
/Å^2. The data were collected in a desired defocus range of -0.7 to -2.1 mm.

Cryo-EM data processing and structure refinement

Data were processed using cryoSPARC V4.2.0[42] and RELION V4.0.1[43,44]. The dose-fractionated
movies were motion corrected using MotionCor2[45]. The contrast transfer function (CTF) was

estimated with CTFFIND 4.1[46]. Particles were picked using the Auto pick function in
RELION[47]. 4,201,166 raw particles were transferred to cryoSPARC for 2D classification.
1,009,168 particles from selected 2D classes were used for Ab-initio reconstruction, where they
were divided into 6 Ab-initio classes. 317,175 particles from class 1 were then refined to a final
resolution of 3.7 Å with non-uniform refinement. To improve local resolution, we performed
local refinement using a mask covering the central FoxP3 tetramer, and obtained a 3.3 Å-
resolution map. For structure refinement, a previous crystal structure of a FoxP3$^{\Delta N}$ monomer
bound to DNA (PDB: 7TDX) was docked into the EM density map from global refinement using
UCSF Chimera[48]. The total of ten copies of FoxP3$^{\Delta N}$ monomers were located for the global
refinement map. For the mask-focused local refinement map, four copies of FoxP3$^{\Delta N}$ monomers
in complex with DNA were docked. Subsequently, the decamer and tetramer models were built
manually against the respective density map using COOT[49], and refined using
phenix.real_space_refine[50]. The structure validation was performed using MolProbity[51] from the
PHENIX package. The curve representing model versus full map was calculated, based on the
final model and the full map. The statistics of the 3D reconstruction and model refinement are
summarized in Extended Data Table 1. All molecular graphics figures were prepared with Pymol
(Schrödinger, LLC) and UCSF Chimera[48]. All softwares used for cryo-EM data processing and
model building were installed and managed by SBGrid[52].

Negative stain EM

795   FoxP3$^{\Delta N}$ (0.4μM) was incubated with DNA (0.05μM) in buffer B at RT for 10 mins. The
796   samples were diluted 10-fold with buffer A, immediately adsorbed to freshly glow-discharged
797   carbon-coated grids (Ted Pella) and stained with 0.75% uranyl formate as described
798   before[53].sImages were collected using a JEM-1400 transmission electron microscope (JEOL) at
799   50,000X magnification.

De novo **motif analysis of FoxP3-occupied sites** *in vitro* **and** *in vivo*

FoxP PD-seq data were mapped to mm10 using Bowtie2[54] and sorted by samtools[55]. Peaks were
called using MACS2[56] with either input or MBP pull-down as controls. Default setting was used
for peak calling. *De novo* motif analysis was done using MEME-ChIP[57] and STREAM[58] with
minimum and maximum motif lengths set at 6 and 30 nt, respectively.
FoxP3 CUT&RUN-seq and ChIP-seq data[14] were mapped to mm10 using Bowtie2[54]. Peaks were
called using MACS2[56]. Bedtools was used to obtain CNR -seq consensus (n=1,372) and union
(n=9,062) peaksbetween Rudensky and Dixon CNR peaks. Motif analysis was done as above. To
independently validate the results, similar motif analysis was repeated using different ChIP-seq
data[23,24], which were mapped to mm10 using Bowtie2. Peaks were called using HOMER with an
input control[22] and ranked them based on the signal intensity using samtools[55]. Top 5,000
overlapping FoxP3 ChIP-seq peaks were calculated by bedtools using a 50% reciprocal overlap
criterion. FoxP3-negative open chromatin regions (OCRs) were derived from all observed Treg
OCRs[27]. Intersections and non-overlapping genomic features were extracted using bedtools[59]
intersect functionality and were subjected to the motif analysis as above. The versions and
parameters for softwares used above have been uploaded to GitHub
(https://github.com/DylannnWX/Hurlab/tree/main/Foxp3_manuscript).

Genome-wide analysis of TnG repeat-like elements

FIMO[60] was used to identify TnG repeat-like elements. The TnG repeat-like motif identified
from the MEME-ChIP analysis of the overlap of Rudensky and Dixon CNR peaks
(Supplementary Table 1b) was used as a query motif, and search was done against the human
(GrCh38), mouse (GrCm38), and Zebrafish (GrCz11) genomes. The default p-value cutoff
(p=0.05) was used. FIMO outputs of all regions that match the query motif were converted to
bed file format, and the overlapping TnG regions from FIMO outputs were combined into a
single region by bedtools merge function.

Comparison between FoxP3 CNR union peaks with and without T_nG repeat-like elements

FIMO[60] was used as above to identify TnG repeat-containing peaks from the Rudensky/Dixon
CNR union peaks (n=9,062). Out of the 9,062 peaks, 3,301 peaks showed at least one T_nG region
lower than the default p-value cutoff (p=0.05), and were classified as T_nG-containing peaks. The
non-TnG-containing peaks were then calculated by bedtools peak subtraction with intersect -v.
Genomic feature analysis was done using ChIPseeker[61]. To compare H3K4me3, H3K27ac and
ATAC signal intensity, H3K4me3 and H3K27ac ChIP-seq and ATAC-seq data[23] were mapped
to mm10 using Bowtie2[54] and the intensity was calculated within 2 kb upstream and downstream
of the FoxP3 CNR peak summits using Deeptools[62] bamCoverage and Deeptools
computeMatrix. The versions and parameters for softwares used above have been uploaded to
GitHub (https://github.com/DylannnWX/Hurlab/tree/main/Foxp3_manuscript).

Motif analysis of other Forkhead TFs

837 Peak bed files for Foxp1, Foxp2, Foxp4, Foxj2, Foxj3, FoxA1, FoxM1, FoxS1 and FoxQ1, were
838 from downloaded from ChIP-Atlas (http://chip-atlas.org/) and converted to fasta files using
839 bedtools[59] getfasta. The individual fasta file was then subjected to *de novo* motif analysis using
840 MEME-ChIP[57] with the minimum and maximum motif lengths set at 6 and 30 nt, respectively.
841 The results are summarized in Supplementary Table 1c.

CD4+ T Cell isolation and retroviral transduction

Naïve CD4+ T cells were isolated by negative selection from mouse spleens using the isolation
kit (Miltenyi Biotec) according to the manufacturer's instruction. The purity was estimated to be
>90% as measured by PE anti-CD4 (Biolegend, Cat#100408, 1:1000 dilution) staining and
FACS analysis. Naïve CD4+ T cells were then activated with anti-CD3 (Biolegend, Cat#100340,
1:500 dilution to 5μg/mL), anti-CD28 (Biolegend, Cat#102116, 1:500 dilution to 5μg/mL) and
50 U/mL of IL2 (Peprotech) in complete RPMI medium (10% FBS heat-inactivated, 2 mM L-
Glutamine, 1 mM Sodium Pyruvate, 100 μM NEAA, 5 mM HEPES, 0.05 mM 2-ME). The
activation state of T cells was confirmed with increased cell size and CD44 (BioLegend)
expression by FACS. After 48 hours, cells were spin-infected with retrovirus containing
supernatant from HEK293T cells transfected with retroviral expression plasmids (Empty MSCV-
IRES-Thy1.1 vector, wildtype-FoxP3 and mutations encoding vectors) and cultured for 2~3 days
in complete RPMI medium with 100 U/mL of IL2.

FoxP3 transcriptional activity assay in CD4⁺T cells

FoxP3 transcriptional activity was measured by levels of two known target genes, CD25 and CTLA4, and the FoxP3 expression marker Thy1.1. FoxP3-transduced CD4$^+$ T cells were stained with antibodies targeting the cell surface antigens CD25 (Biolegend, Cat#102022, 1:1000 dilution) and Thy1.1 (Biolegend, Cat#202520, 1:1000 dilution) on day 2 post-retroviral infection. The level of CTLA4 was measure by intracellular staining using anti-CTLA4 (Biolegend, Cat#106311, 1:1000 dilution) and the Transcription Factor Staining Buffer Set (eBioscience) on day 3 post retroviral infection. Flow cytometry data were analyzed with FlowJo software and presented as plots of mean fluorescence intensity (MFI) of CD25 and CTLA4 in cells grouped into bins of Thy1.1 intensity, which is the expression marker for FoxP3. Each result is representative of 3 independent experiments.

FoxP3 ChIP-seq analysis

FoxP3 ChIP-seq was conducted using CD4$^+$ T cells following a published procedure[63]. Activated CD4$^+$ T cells that had been transduced with wild-type or mutant FoxP3 were sorted based on Thy1.1 reporter expression. For each sample (5×10^6 cells), cross-linking was achieved with 1% formaldehyde for 10 minutes. Subsequently, the cells were lysed on ice using RIPA buffer (10mM Tris-HCl pH 8.0, 1mM EDTA, 1% Triton X-100, 0.1% Sodium Deoxycholate, 0.1% SDS, 140 mM NaCl, and 1X proteinase inhibitor). Chromatin fragmentation was achieved using an AFA Focused-ultrasonicator (Covaris M220) for 30 minutes (5% duty cycle, 140W max power, 200 cycle/burst), resulting in DNA fragments ranging from 100 to 200 bp. The sheared material underwent a 10-minute centrifugation at 13,000 rpm at 4°C to clear the solution. The cleared material was then subjected to immunoprecipitation overnight with HA-tag antibody (Cell Signaling, #3724) at 4°C, and Protein G beads (Active motif, #53014) were added for an additional 2 hours. The beads were sequentially washed with various buffers: RIPA Wash Buffer (0.1% SDS, 0.1% Sodium Deoxycholate, 1% Triton X-100, 1mM EDTA, 10mM Tris-HCl pH 8.0, 150mM NaCl), RIPA 500 Wash Buffer (0.1% SDS, 0.1% Sodium Deoxycholate, 1% Triton X-100, 1mM EDTA, 10mM Tris-HCl pH 8.0, 500mM NaCl), LiCl wash buffer (10mM Tris-HCl, pH 8.0, 250mM LiCl, 0.5% Triton X-100, 0.5% Sodium Deoxycholate), and Tris buffer (10mM Tris-HCl, pH 8.5). The chromatin was eluted from the beads using Elution buffer (1X TE, pH 8.0, 0.1% SDS, 150mM NaCl, 5 mM DTT). After elution, the DNA was treated with 1 µg DNase-free RNase (Roche) for 30 minutes at 37°C, followed by treatment with Proteinase K (Roche) for at least 4 hours at 63°C to reverse the crosslinks. The reverse-crosslinked DNA was then purified using SPRI beads (Beckman, #B23318). Subsequent steps, including end repair, A-base addition, adaptor-ligation, and PCR amplification, were carried out to prepare the ChIP-seq library for each sample. The libraries were made by NEBNext® Ultra™ II DNA Library Prep Kit (Illumina) according to the manufacturer's instructions and submitted to Novogene for NGS with PE150.

mRNA-seq analysis

mRNA-seq was conducted using CD4$^+$ T cells. Activated CD4$^+$ T cells that had been transduced with wild-type or mutant FoxP3 were sorted based on Thy1.1 reporter expression. For each sample, 1×10^6 cells were sorted and subjected to total RNA extraction using TRIzol reagent and

Direct-zol RNA Miniprep Kits. Quality control and the construction of mRNA-seq libraries were
undertaken by Novogene Co. Ltd.. The NEB Next Ultra II kit and the non-directional mRNA
approach with the polyA pipeline were used. The libraries were subsequently sequenced using
the Illumina NovaSeq 6000 instrument, generating paired-end reads with a length of 2X150 bp,
resulting in ~30 M reads per sample. Raw sequence files were subjected to pre-processing using
Trimmomatic v.0.36 to remove Illumina adaptor sequences and low-quality bases. Trimmed
reads were then aligned to the mouse genome (UCSC mm10) using bowtie2/2.3.4.3. For gene
read counting, HTseq-count (v. 0.12.4) was employed. Normalization of gene counts and
differential analysis were carried out using DESeq2 (v. 5). The creation of heatmaps was
accomplished using Pheatmap.

Chromatin contact analysis

HiC- and PLAC-seq datasets were downloaded from GSE217147[12], and the list of Treg
enhancer-promoter loops (EPLs) was obtained from[13]. All .hic files were converted to .cool files
to hic2cool, and all .cool files were decompressed into .txt files by cooler dump –join function.
These decompressed files were loaded as python pandas dataframes. All possible bins in .cool
files were converted to bed file formats, and intersected with TnG-containing or TnG-absent
CNR union peaks using bedtools intersect -wa function to acquire the bins that contain TnG bins
and non-TnG (NTnG) bins. These bins were used as anchors to filter raw .cool files for contact
pairs between TnG–TnG (2TnG), TnG–NTnG (TnGNTnG), and NTnG–NTnG (2NTnG). These
contact pairs were then filtered by [more than 5 in WT Treg HiC-seq] and [more than indicated
threshold in FoxP3 PLAC-seq]. Contact counts in Fig. 2f were listed in Supplementary Table 3.

The p-value of 2TnG pair enrichment was first calculated by getting the expected 2TnG pair
counts in a given list of pairs assuming random distribution (number of contact pairs *
proportion of all potential TnG bins^2). Then, this number was compared with the observed
2TnG pair counts using binomial distribution. The proportion of all potential TnG bins is 0.37,
which matches the proportion of TnG CNR peaks out of all CNR peaks (3,301 out of 9,062). The
p-value was the cumulated probability that the observed 2TnG pair counts happen by chance,
and the alternative hypothesis, if the p-value is low, indicates the probability that in 2TnG pair is
enriched in the given list of contact.

To compare HiC/PLAC-seq anchors (in mm9) to EPL anchors (in mm10), the reference genomes
of mm9 were lifted to mm10 using UCSC genome browser to acquire the correlating bin
coordinates in mm10, and their overlaps were analyzed using bedtools intersect function.

T cell suppression assay

Isolated naïve $CD4^+$ T cells were activated with anti-CD3 (Biolegend), anti-CD28 (Biolegend)
and 50 U/mL of IL2 (Peprotech) in complete RPMI medium. After 48 hours, activated $CD4^+$ T
cells were retrovirally transduced to express FoxP3 and were used as "suppressors". In parallel,
freshly isolated naïve $CD4^+$ T cells were labelled with CellTrace™ CFSE (Invitrogen) and used
as "responders". $CD3^-$ T cells representing APC cells were also isolated using the isolation kit
(Miltenyi Biotec) according to the manufacturer's instructions. For suppression assay, the CFSE-
labeled responder cells ($5X10^4$ cells) were stimulated with APC cells (10^4 cells) and anti-CD3 (1

μg/mL) in 96-well round-bottom plates for 3 days, in the presence or absence of FoxP3-
transduced suppressor cells (at a responder-to-suppressor ratio of 2:1). Proliferation ratio of the
responders were calculated as a function of CFSE dye dilution by FACS analysis.

Statistics and Reproducibility

940 Data in Figures 1f-j, 2e, 3b-e, 4b, 4d, 4e, 4f, 4g, 5b, 5c and extended data figures 1g-l, 2a, 2h, 3b,
941 3c, 3e, 3f, 5c-e, 6a-e, 6g are representative of at least three independent experiments and each
942 experiment was repeated independently with similar results.

Data availability

Naked genomic DNA pulldown-seq, nucleosome pulldown-seq, FoxP3 mRNA-seq and FoxP3
ChIP-seq data have been deposited to the GEO database with the accession code of GSE243606.
The structures and cryo-EM maps have been deposited to the PDB and the EMDB under the
accession codes of 8SRP and EMD-40737 for the decameric FoxP3 in complex with DNA, and
8SRO and EMD-40736 for the central FoxP3 tetramer in complex with DNA (focused
refinement). Other research materials reported here are available on request.
950

Code availability

952 All custom codes used in this project are deposited to GitHub (link:
953 https://github.com/DylannnWX/Hurlab/tree/main/Foxp3_manuscript). These include the
954 processing of Deeptools matrix outputs, FIMO region to peak bed files, and HiC/Cool data
955 processing. All are standalone Jupyter Notebook instances. In each instance, detailed user
956 instructions, example inputs and expected outputs were also included in this GitHub Repo.

Additional references associated with Methods

40 Moparthi, L. & Koch, S. A uniform expression library for the exploration of FOX
transcription factor biology. *Differentiation* **115**, 30-36, doi:10.1016/j.diff.2020.08.002
(2020).
41 Dyer, P. N. *et al.* Reconstitution of nucleosome core particles from recombinant histones
and DNA. *Methods Enzymol* **375**, 23-44, doi:10.1016/s0076-6879(03)75002-2 (2004).
42 Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for
rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296,
doi:10.1038/nmeth.4169 (2017).
43 Scheres, S. H. A Bayesian view on cryo-EM structure determination. *J Mol Biol* **415**,
406-418, doi:10.1016/j.jmb.2011.11.010 (2012).
44 Kimanius, D., Dong, L., Sharov, G., Nakane, T. & Scheres, S. H. W. New tools for
automated cryo-EM single-particle analysis in RELION-4.0. *Biochem J* **478**, 4169-4185,
doi:10.1042/BCJ20210708 (2021).
45 Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat Methods* **14**, 331-332, doi:10.1038/nmeth.4193
(2017).

46 Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from
electron micrographs. *J Struct Biol* **192**, 216-221, doi:10.1016/j.jsb.2015.08.008 (2015).
47 Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure
determination in RELION-3. *Elife* **7**, doi:10.7554/eLife.42166 (2018).
48 Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and
analysis. *J Comput Chem* **25**, 1605-1612, doi:10.1002/jcc.20084 (2004).
49 Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot.
Acta Crystallogr D Biol Crystallogr **66**, 486-501, doi:10.1107/S0907444910007493
(2010).
50 Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and
electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* **75**, 861-877,
doi:10.1107/S2059798319011471 (2019).
51 Williams, C. J. *et al.* MolProbity: More and better reference data for improved all-atom
structure validation. *Protein Sci* **27**, 293-315, doi:10.1002/pro.3330 (2018).
52 Morin, A. *et al.* Collaboration gets the most out of software. *Elife* **2**, e01456,
doi:10.7554/eLife.01456 (2013).
53 Ohi, M., Li, Y., Cheng, Y. & Walz, T. Negative Staining and Image Classification -
Powerful Tools in Modern Electron Microscopy. *Biol Proced Online* **6**, 23-34,
doi:10.1251/bpo70 (2004).
54 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*
9, 357-359, doi:10.1038/nmeth.1923 (2012).
55 Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**,
doi:10.1093/gigascience/giab008 (2021).
56 Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137,
doi:10.1186/gb-2008-9-9-r137 (2008).
57 Bailey, T. L., Johnson, J., Grant, C. E. & Noble, W. S. The MEME Suite. *Nucleic Acids*
Res **43**, W39-49, doi:10.1093/nar/gkv416 (2015).
58 Bailey, T. L. STREME: accurate and versatile sequence motif discovery. *Bioinformatics*
37, 2834-2840, doi:10.1093/bioinformatics/btab203 (2021).
59 Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**, 841-842, doi:10.1093/bioinformatics/btq033
(2010).
60 Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given
motif. *Bioinformatics* **27**, 1017-1018, doi:10.1093/bioinformatics/btr064 (2011).
61 Yu, G., Wang, L. G. & He, Q. Y. ChIPseeker: an R/Bioconductor package for ChIP peak
annotation, comparison and visualization. *Bioinformatics* **31**, 2382-2383,
doi:10.1093/bioinformatics/btv145 (2015).
62 Ramirez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data
analysis. *Nucleic Acids Res* **44**, W160-165, doi:10.1093/nar/gkw257 (2016).
63 Kwon, H. K., Chen, H. M., Mathis, D. & Benoist, C. Different molecular complexes that
mediate transcriptional induction and repression by FoxP3. *Nat Immunol* **18**, 1238-1248,
doi:10.1038/ni.3835 (2017).
1016

Acknowledgements

We thank all members of the Hur lab, A. Rudensky (MSKCC) and Y. Zhong (Shanghai Jiao
Tong University) for their helpful discussion and feedback. This study was supported by Modell

fellowship to W.Z., NIH grants (R01AI180137, R01AI154653 and R01AI111784 to S.H;
AI150686 and AI165697 to C.B.) and Howard Hughes Medical Institute (S.H.). Cryo-EM data
were collected at Cryo-EM facilities at Janelia, Harvard Medical School and University of
Massachusetts Worcester.
1024
1025   **Author contributions**
1026   W.Z., F.L., S.H. conceived and designed the project. W.Z., F.L. performed all the experiments.
1027   F.L. determined the structure. W.Z., X.W, R.R. performed bioinformatic analysis. J.P. assisted
1028   experiments. C.B. and S.H. supervised bioinformatic analysis. S.H. supervised overall project.
1029
1030   **Competing Interests**
1031   The authors declare no competing interests.
1032

1033   **Extended figure legends**

1034

Extended Data Figure 1. Analysis of T_3G repeats in the genome and FoxP3 multimerization
on T_3G repeats

a. T_nG repeat-like sequences in the genomes of *H. sapiens*, *M. musculus* and *D. rerio*.
Sequences that match the T_nG repeat-like motif (29 nt motif from FoxP3 CNR overlap
peaks, see Supplementary Table 1b) were identified using FIMO ($p = 0.05$, see Methods).
Genomic percentage of T_nG repeat-like sequences (in parenthesis) was the number of
T_nG repeat regions multiplied by the average size of the repeats (31, 33 and 38 bp for *H.*
sapiens, *M. musculus* and *D. rerio*, respectively), divided by the genome size (3.2, 2.7
and 1.4 billion bp, respectively). Below: length distribution of the TnG repeat-like
sequences. Genes-to-genome size ratio was calculated by dividing the number of genes
used in the feature annotation (31,074, 24,528 and 13,576 in *H. sapiens*, *M. musculus* and
D. rerio) by the genome size.
b. Distribution of T_nG repeat-like sequences in the genomes of *H. sapiens*, *M. musculus* and
D. rerio relative to Transcription Start Sites (TSSs).
c. Distribution of CNR union peaks (union of Rudensky CNR peaks and Dixon CNR peaks,
n=9,062) relative to TSSs. CNR union peaks with and without T_nG repeat-like sequences
(n=3,301 and 5,761, respectively) were identified using FIMO ($p = 0.05$) as in (a).
d-f. Comparison of (d) H3K4me3-ChIP, (e) H3K27ac-ChIP and (f) ATAC signal [23] around
the T_nG repeat-like sequences that overlap with FoxP3 CNR union peaks vs. those
genome-wide in thymic Tregs (n = 4,837 peaks and 41,889 peaks respectively). See
Extended Data Figs. 4a-4c for pre-thymic Tregs, which showed that high levels of
H3K4me3, H3K27ac and ATAC signals were maintained prior to FoxP3 expression. T_nG
repeats in the blacklist were removed. Right: ChIP/ATAC signal was averaged over +/-
500 bp around the T_nG repeat-like sequences. Two-tailed unpaired t-tests. ****, $p <$
0.0001.
g. DNA sequence specificity of FoxP3 as measured by FoxP3 pull-down. HA-tagged, full-
length FoxP3 was transiently expressed in HEK293T cells and purified by anti-HA IP.
Equivalent amounts of indicated DNAs (30-31 bp) were added to FoxP3-bound beads
and further purified by anti-HA IP prior to gel analysis.

h.   DNA sequence specificity of FoxP3 as measured by DNA pull-down. Equivalent
amounts of biotinylated DNAs were mixed with FoxP3-expressing 293T lysate and were
subjected to streptavidin pull-down. Co-purified FoxP3 was analyzed by anti-HA WB.
i.   FoxP3 binding to nucleosomal DNA as measured by native gel-shift assay. Indicated
DNA was incubated with the histone octamer at 1:1 molar ratio (black circle), followed
by incubation with FoxP3 (0.2 or 0.4 μM for light and dark green circles, respectively).
Empty dotted circles indicate no histone or FoxP3. Sybrgold stain was used for
visualization. With an increasing concentration of FoxP3, the intensity of the
nucleosomal TTTG repeat decreased, while the signal in the gel well (red arrow)
increased. Such changes were not observed with other DNAs.
j.   BMOE crosslinking of FoxP3$^{\Delta N}$ with and without DNA. FoxP3$^{\Delta N}$ can only form
multimers on $(T_3G)_6$ DNA.
k.   Multimerization analysis of FoxP3, as measured by co-purification of FoxP3 with
different tags. GST- and MBP-tagged FoxP3 were incubated together in the presence and
absence of indicated DNA and were subjected to MBP pull-down, followed by WB
analysis of GST-FoxP3 in eluate. Note that GST replaced the CC domain in FoxP3,
disallowing hetero-dimerization between MBP-FoxP3 and GST-FoxP3. Thus, co-
purification of these two proteins in the presence of T_3G repeats suggests DNA sequence-
dependent multimerization of the FoxP3 homodimer.
l.   Representative negative-stain EM images of FoxP3$^{\Delta N}$ in complex with $(AAAG)_{36}$ (left)
and $(TGTG)_{36}$ (right).
1085

Extended Data Figure 2. Cryo-EM structure of the FoxP3$^{\Delta N}$–$(T_3G)_{18}$ complex.

1086

a.   Representative negative-stain EM (left) and cryo-EM images (right) of FoxP3$^{\Delta N}$
multimers on $(T_3G)_{18}$ DNA.
b.   2D classes chosen for 3D reconstruction.
c.   Cryo-EM image processing workflow. See details in Methods.
d.   Local resolution for the maps of global refinement (left) and local refinement with a mask
covering the central four subunits of FoxP3 (right). Local resolution was calculated by
CryoSPARC. Resolution range was indicated according to the color bar.
e.   Fourier shell correlation (FSC) curve for global refinement (left) and local refinement
(right). Map-to-Map FSC curve was calculated between the two independently refined
half-maps after masking (blue line), and the overall resolution was determined by gold
standard FSC=0.143 criterion. Map-to-Model FSC was calculated between the refined
atomic models and maps (red line).
f.   Cryo-EM map and ribbon model of FoxP3$^{\Delta N}$ decamer in complex with two $(T_3G)_{18}$
DNAs (PDB: 8SRP, EMDB: 40737). DNA molecules are colored grey. Individual FoxP3
monomers are colored differently.
g.   Superposition of the domain-swap dimeric structure of FoxP3 (cyan, PDB:4WK8) onto
any subunit of the FoxP3 multimeric structure (represented here by the orange subunit)
by aligning the common portions of FoxP3 reveals that the domain-swap dimer is
incompatible with the density map.
h.   Native gel shift analysis of MBP-tagged FoxP3$^{\Delta N}$ (WT or R337Q, 0.4 μM) with $(T_3G)_{12}$
DNA (0.05 μM). Note that R337Q induces domain-swap dimerization.
1108

Extended Data Figure 3. DNA sequence and conformational analysis.

1109

a. DNA sequence assignment by Q-score analysis using mapQ. For each of the four DNA strands, eight possible sequence alignments (right) were tested against the local refinement map. The sequence alignment showing the highest overall score was highlighted with a thick red line. The best sequence alignments for all four DNA strands were consistent with each other and were used for cryo-EM reconstruction.

b. Experimental validation of FoxP3–DNA registers using NFAT–FoxP3 cooperativity analysis. This assay utilizes the FoxP3 interaction partner NFAT, which assists FoxP3 binding to DNA only when their binding sites are 3 bp apart in one particular orientation (as in the schematic). To investigate FoxP3 footprints on T_3G repeat DNA, we varied the position and orientation of NFAT consensus sequence (GGAAA, green) relative to T_3G repeats (Var1-6), and performed FoxP3 pull-down. An internal control DNA (cntrl, harboring the NFAT motif followed by FKHM with 3 nt gap) was used to normalize the test DNA (Var1-6) pull-down efficiency. Only DNA with a single nucleotide gap between GGAAA and TTTG (Var2) showed a positive effect of NFAT on the FoxP3–DNA interaction. This suggests that the most upstream FoxP3 subunit recognizes TGTTTGT. Two-tailed paired t-tests, comparing with and without NFAT. $p<0.005$ for **, $p<0.05$ for * and $p>0.05$ for ns.

c. FoxP3 interaction with $(T_3G)_{10}$ variants. Variations in DNA sequence outside the FoxP3 footprints were tolerated (Var7 and Var8), but those within the footprints (Var9 and Var10) were not.

d. Comparison of the inter-subunit interactions in FoxP3 decamer on T_nG repeats vs head-to-head (H-H) dimer on IR-FKHM (PDB:7TDX). Superposition of the H-H dimer (dark grey) onto any of the ten subunits in the decamer structure showed distinct modes of inter-subunit interactions. Shown are two examples where subunit 1 of the H-H dimer was aligned to orange (top) or magenta (bottom) subunits of the decamer, showing that subunit 2 of the H-H dimer did not align with any of the decamer subunits.

e. DNA bridging assay using FoxP3 expressed in 293T cells. Biotinylated and non-biotinylated DNA (82 and 60 bp, respectively) were mixed at 1:1 ratio and further incubated with 293T lysates expressing HA-tagged FoxP3, followed by streptavidin pull-down and gel analysis of non-biotinylated DNA by SybrGold staining.

f. DNA bridging assay using an increasing concentration of purified FoxP3$^{\Delta N}$. 0.1 μM each of biotinylated and non-biotinylated $(T_3G)_{12}$ DNAs were used. FoxP3 concentrations are indicated at the bottom.

Extended Data Figure 4. FoxP3 can bridge T_nG repeat-containing sites in vivo.

a-c. Comparison of (a) H3K4me3-ChIP, (b) ATAC and (c) H3K27ac-ChIP signal [23] around the CNR union peaks with and without T_nG repeat-like sequences in pre-thymic Tregs (pre-tTregs) and thymic Tregs (tTregs) (n=3,301 peaks and 5,761 peaks, respectively). Right: ChIP/ATAC signal was averaged over +/- 500 bp around the CNR peak summits. Two-tailed unpaired t-tests. ****, $p < 0.0001$.

d. FoxP3-dependence of the chromatin contacts at FoxP3-bound T_nG anchors in Fig. 2f. FoxP3-bound T_nG anchors were defined as anchors that overlap with FoxP3 CNR peaks with T_nG repeat-like sequences. Contacts with frequency>5 in WT Treg HiC and connected by two T_nG anchors were analyzed with an increasing FoxP3 PLAC-seq count threshold. For each contact, log2 foldchange of HiC counts from WT to FoxP3 knock-out Treg were plotted. Contacts with FDR<10 were colored red. The majority of the T_nG–

1156	T_nG contacts were less frequent in FoxP3 knock-out than in WT Tregs, although smaller
1157	fractions (10-15%) showed statistically significant FoxP3 dependence (FDR<10), as
1158	previously reported[12]. n=4365, 2559, 813, 204 and 60 anchors respectively. Mean ± SD
1159	were shown in black and blue lines. See also Supplementary Table 3.

1160   e. Fraction of the FoxP3-bound T_nG anchors from Fig. 2f that overlap with previously
1161   published Treg EPL anchors [13].

1162

Extended Data Figure 5. Characterization of intra-rung interface mutant FoxP3

a. mRNA-seq heatmap analysis. CD4$^+$ T cells were transduced and sorted to express FoxP3
as in Fig. 3d and were subjected to mRNA-seq. Top 100 genes showing the most
significant difference between WT FoxP3 and EV were chosen for the evaluation of
individual mutants. All four mutants were impaired in transcriptional functions, albeit to
varying extents. The level of FoxP3 was equivalent for WT and all mutants. A few genes
previously reported to be FoxP3-dependent were indicated in larger fonts. Note that
V398E was not tested due to its negative effect on NFAT binding in (e).

b. ChIP-seq of HA-tagged FoxP3. Cells were transduced as in Fig. 3d and were subjected to
anti-HA ChIP-seq. WT FoxP3 bound peaks were identified using MACS2 (n=8,607,
$p<0.01$), and heatmaps of the ChIP signal were generated for each mutant at the WT peak
locations. Below: averaged intensity of ChIP signal within 0.5 kb of the WT peak
summit. Peaks with and without T_nG repeats (n=1,900 peaks and 6,707 peaks,
respectively) were compared. Two-tailed paired t-tests, comparing mutants to WT. $p <$
0.0001 for ****, $p<0.001$ for *** and $p<0.005$ for **.

c. Nuclear localization of WT FoxP3 and intra-rung interface mutants. HA-tagged FoxP3
was transiently expressed in A549 cells and was subjected to anti-HA immunofluorescent
(yellow) analysis. Nuclei were shown with DAPI (blue) staining.

d. Expression levels of WT FoxP3 and intra-rung mutants in A549 cells.

e. Effect of the intra-rung mutations on the NFAT–FoxP3 interaction, as measured by
native gel shift assay. FoxP3 (0.1 μM) was incubated with DNA harboring IR-FKHM
and the NFAT site (with a 3-bp gap as in Extended Data Fig. 3b, 0.05 μM). NFAT (0.1
μM) was added to the mixture to monitor formation of the ternary complex NFAT–
FoxP3–DNA. Note that V398E showed slight but reproducible reduction in NFAT
binding.

1188

Extended Data Figure 6. Multimerization on T_nG repeats is conserved in FoxP3 orthologs
and paralogs.

a. T_nG repeats DNA-binding activity of FoxP3 with mutations in RBR. All seven RBR
mutations previously shown to disrupt the head-to-head dimerization [22] disrupted $(T_3G)_{12}$
binding.

b. Effect of inter-rung^{12bp} spacing variations on FoxP3-mediated DNA bridging. Non-
biotinylated DNAs in Fig. 4d were mixed with biotinylated $(T_3G)_{10}$ and FoxP3$^{\Delta N}$ (0.2
μM) prior to streptavidin pull-down and gel analysis. Relative level of non-biotinylated
DNA co-purified with biotinylated DNA was quantitated from three independent pull-
downs. Two-tailed paired t-tests, in comparison to $(T_3G)_{10}$. $p<0.001$ for ***, $p<0.05$ for *
and $p>0.05$ for ns.

c. DNA-bridging activity of FoxP3 with different combinations of T_nG repeats.
Biotinylated-DNA (red) and non-biotinylated DNA (blue) were mixed at 1:1 ratio and

were incubated with FoxP3$^{\Delta N}$ (0.2 μM) prior to streptavidin pull-down and gel analysis of non-biotinylated DNA. T$_2$G, T$_4$G and T$_5$G repeats bridged better with T$_3$G repeats than with themselves.

d. DNA-binding activity of FoxP3 orthologs with indicated DNA. Experiments were performed as in Fig. 5b.

e. DNA-bridging activity of FoxP3 orthologs. Experiments were performed as in Fig. 5c.

f. Sequence alignment of FoxP3 orthologs from different species, showing conservation of the key interface residues (yellow highlight, arrows on top with the residue identities in *M. musculus* FoxP3).

g. DNA-binding activity of FoxP3 paralogs with T$_n$G repeats (n=1-4). Experiments were performed as in Fig. 5b.

Extended data table 1

ED table 1. Table for Cryo-EM data collection, refinement and validation statistics. The statistics of the 3D reconstruction and model refinement are summarized.

a

F331

b

IR-FKHM (T₃G)₁₂

H-H dimer multimer

DNA DNA

− WT F331D − WT F331D

c

inter-rung^8bp spacing

(T₃G)₁₀

8bp   8bp

TAGGAAA GTT TGTTTGTT TGTTTGTTTGTT TGTTTGTT TGTTTGTTT

variants 8bp A   A
          9bp AC   AC
          7bp −   −

1XFKHM (T₃G)₁₀ | inter-rung^8bp spacing
                 8   9   7

IP: FoxP3
DNA
input

d

(T₃G)₁₀   12bp

variants 10-23bp

1XFKHM (T₃G)₁₀ | inter-rung^12bp spacing
10 11 12 13 14 17 19 20 21 22 23

DNA IP: FoxP3
input

Relative intensity

p=0.015 *ns,p*=0.117 *p*=0.037 **p*=0.002 ***p*=0.004 ****p*=0.002 *****p*=0.003 ******p*=0.013 *******p*=0.033 *ns,p*=0.073 *ns,p*=0.085

1XFKHM (T₃G)₁₀ 10 11 12 13 14 17 19 20 21 22 23
inter-rung^12bp spacing

e

DNA binding | DNA bridging

input FoxP3 PD | input biotin PD input biotin PD

CN53
Cast (T₂G)₁₆
B6 (T₂G)₆

CN118
Cast (T₄G)₁₀
B6 (T₄G)₆

CN16
Cast (T₅G)₁₅
B6 (T₅G)₅

f

T₂G–T₂G   T₄G–T₄G   T₅G–T₅G

FoxP3 − WT V408M − WT V408M − WT V408M

non-bio-DNA IP: biotin
input

g

T₂G–T₂G   T₃G–T₂G

FoxP3   FoxP3

non-bio-DNA IP: biotin
input