A simulation framework for modeling the within-patient evolutionary dynamics of SARS-CoV-2

John W Terbot II[1,2*], Brandon S. Cooper[2], Jeffrey M. Good[2], Jeffrey D. Jensen[1*]

[1] Arizona State University, School of Life Sciences, Center for Evolution & Medicine, Tempe,

Arizona, United States of America

[2] University of Montana, Division of Biological Sciences, Missoula, Montana, United States of

America

* jterbot@asu.edu (JWT); jeffrey.d.jensen@asu.edu (JDJ)

Abstract

The global impact of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) has led

to considerable interest in detecting novel beneficial mutations and other genomic changes that

may signal the development of variants of concern (VOCs). The ability to accurately detect these

changes within individual patient samples is important in enabling early detection of VOCs.

Such genomic scans for positive selection are best performed via comparison of empirical data to

simulated data wherein evolutionary factors, including mutation and recombination rates,

reproductive and infection dynamics, and purifying and background selection, can be carefully

accounted for and parameterized. While there has been work to quantify these factors in SARS-

CoV-2, they have yet to be integrated into a baseline model describing intra-host evolutionary

dynamics. To construct such a baseline model, we develop a simulation framework that enables

one to establish expectations for underlying levels and patterns of patient-level variation. By

varying eight key parameters, we evaluated 12,096 different model-parameter combinations and

compared them to existing empirical data. Of these, 592 models (~5%) were plausible based on

the resulting mean expected number of segregating variants. These plausible models shared

several commonalities shedding light on intra-host SARS-CoV-2 evolutionary dynamics: severe

infection bottlenecks, low levels of reproductive skew, and a distribution of fitness effects

skewed towards strongly deleterious mutations. We also describe important areas of model

uncertainty and highlight additional sequence data that may help to further refine a baseline

model. This study lays the groundwork for the improved analysis of existing and future SARS-

CoV-2 within-patient data.

Key Words: evolutionary genomics, population genetics, SARS-CoV-2, viral evolution

Significance Statement

Despite its tremendous impact on human health, a comprehensive evolutionary baseline model

has yet to be developed for studying the within-host population genomics of SARS-CoV-2.

Importantly, such modeling would enable improved analysis and provide insights into the key

evolutionary dynamics governing SARS-CoV-2 evolution. Given this need, we have here

quantified a set of plausible baseline models via large-scale simulation. The commonly shared

features of these relevant models - including severe infection bottlenecks, low levels of progeny

skew, and a high rate of strongly deleterious mutations - lay the foundation for sophisticated

analyses of SARS-CoV-2 evolution within patients using these baseline models.

Introduction

The emergence of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) in late 2019 is the most impactful human pathogen to arise thus far in the 21st century. Since its emergence, SARS-CoV-2 is directly responsible for nearly 8 million deaths as of December 2022 (Institute for Health Metrics and Evaluation 2022, Hay et al. 2023). However, the true impact of SARS-CoV-2—considering under-reporting, late reporting, and indirect deaths (*e.g.*, via strained health care systems)— is likely much greater, with worldwide excess mortality estimated to exceed 14 million as of December 2021 (Wang et al. 2022, Msemburi et al. 2023). Moreover, SARS-CoV-2 continues to persist worldwide and appears likely to become an endemic virus going forward, as observed with other human coronaviruses (HCoV-229E, -NL63, -OC43, and -HKU1; Corman et al. 2018).

Understanding the evolutionary dynamics of SARS-CoV-2 and predicting new Variants of Concern (VOCs) that may result in waves of increased infection and mortality remains vital. While the tools for studying these questions as framed via inter-host spread of SARS-CoV-2 have been well-developed (Rambaut et al. 2020, O'Toole et al. 2021), the ability to study intra-host evolution of SARS-CoV-2 is considerably less established. Transmission between hosts is an obviously important stage in viral spread; however, inter-host spread is a brief portion of the viral life cycle, with the entirety of viral reproductive activity occurring within a host. Thus, dissecting the intra-host evolutionary dynamics are of key importance in monitoring contemporary and future SARS-CoV-2 spread. In particular, mutations that influence evasion of the host immune system, increase success in cell invasion, and otherwise improve the successful completion of metabolic and reproductive tasks within a host cell could all be of clinical consequence. Given complete information, such mutations would first be detectable within a

single host. While strongly beneficial mutations can eventually become identifiable when

observing inter-host data through their increased prevalence within the meta-population (should

they escape stochastic loss), the evolutionary dynamics that ultimately dictate their spread will be

determined at the intra-host level.

For the viral population within a single patient, these episodic, beneficial mutations may

be expected to modify patterns of genomic variation in a manner that would deviate from

background patterns produced under constantly operating evolutionary processes (*e.g.*, via a

selective sweep of the beneficial mutation (Charlesworth and Jensen 2021). As such, as with any

natural population, the study of viral intra-host evolution requires the construction of an

evolutionary 'null' model to quantify expected baseline levels and patterns of genomic variation

(Jensen 2021, Johri et al. 2020, Terbot et al. 2023). At a minimum, a viral baseline model should

include mutation, recombination, reproductive dynamics, purifying and background selection,

and the history of bottlenecks and growth characterizing patient infection (Irwin et al. 2016).

Without comparison to such a baseline model, it is not possible to determine if observed within-

patient allele frequencies are attributable to these common evolutionary processes or to the

comparatively rare action of positive selection (Barton 1998, Johri et al. 2022b).

In this study, we present a first attempt to construct such a model and narrow the range of

key parameter values governing the intra-host evolutionary dynamics of SARS-CoV-2. Using

forward simulations and comparison to existing patient data summarizing intra-host variation, we

identify more and less likely areas of parameter space. We find that transmission and infection

bottlenecks appear to be strongly constrained (*i.e.,* on the order of <5 virions) in plausible

models, consistent with other recent results for SARS-CoV-2 (Lythgoe et al. 2021, Martin and

Koelle 2021, Bendall et al. 2023) and the transmission of other airborne viruses like seasonal

influenza (McCrone et al. 2018, Valesano et al. 2019). We also describe important areas of

uncertainty and correlations between parameter values. For example, if the distribution of new

mutational effects is heavily skewed towards strongly deleterious mutations, the range of

uncertainty in other parameter values is inflated. Furthermore, we highlight additional data that

may help to further narrow this parameter space. For example, the commonly employed 2%

minor allele frequency threshold cut-off greatly limits model resolution, and may be improved by

higher-coverage sequencing of individual patient samples that increases the confidence in

individual single nucleotide polymorphism (SNP) calls. Thus, the presented exploration of the

SARS-CoV-2 evolutionary parameter space will be valuable in informing future modeling

studies, in interpreting newly emerging patient data, and in guiding future data collection.

Results

A total of 12096 model-parameter combinations were analyzed (Table 1, Table 2, and Figure 1).

The great majority were rejected for generating too few or too many segregating SNPs relative to

available empirical data, leaving 592 models remaining (Supplemental Table 1). Due to the

nested nature of the models, the feasible model-parameter sets can be readily categorized

according to the 108 possible combinations of the required parameters. Of these, 18 specific

parameter combinations produced viable models, highlighting which parameters are likely to

have the strongest influence of intra-host evolution of SARS-CoV-2. We found that all plausible

models included a strong infection bottleneck *(i.e.*, a bottleneck under 5 virions appears most

consistent with the data, represented here by a bottleneck of 1; Figure 2, Figure 3). Most retained

models also had low or moderate mutation rates. Only three models with the highest mutation

rate were retained. All three of these models were full models using a distribution of fitness

effects (DFE) with the largest proportion of strongly deleterious mutations, the highest carrying

capacity, and briefest infection duration; the parameter values for recombination and progeny

skew varied between these models. In contrast, all 12 required parameter sets using the lowest

mutation rate and a severe bottleneck of 1 cleared the SNP threshold, and 5 of the 12 sets with

the midpoint mutation rate resulted in plausible models. Generally, larger carrying capacities

resulted in better fitting models (4/12 parameter sets had plausible models with the lowest value

of carrying capacity and a bottleneck of 1, 6/12 with the midpoint value for carrying capacity,

and 8/12 for the highest carrying capacity).

Within these required parameter sets, there are a total of 112 nested models (1 bottleneck

model, 3 with the addition of recombination, 27 with the additions of recombination and progeny

skew, and 81 with the additions of recombination, progeny skew, and a DFE); the number of

models that cleared the filtering requirement varied considerably between the required parameter

sets. Briefly, 9 of the 18 required parameter sets had bottleneck models consistent with the

empirical data (of which 8 had examples of all nested models also clear the threshold), 11 of the

18 had consistent recombination-only models, 10 out of 18 had consistent recombination +

progeny skew models, and all 18 had consistent full models. Generally, if a simpler model could

be accepted, then more complicated models nested within that model also tended to produce

plausible results. The distribution of these non-rejected models are more fully detailed in Table 3

and Figure 2. In terms of general patterns, models with strong infection bottlenecks, lower

progeny skews, and DFEs containing a greater proportion of strongly deleterious mutations were

more likely to be accepted (Figure 3, and see Discussion).

Discussion

The evaluation of intra-host population genomic data from SARS-CoV-2 patient samples has the

potential to detect signatures of positive selection and allow for the early identification of VOCs.

However, it is important to recognize that levels and patterns of genetic variation in a population

are the result of a variety of factors including mutation, recombination, reproductive dynamics,

purifying and background selection, and the history of bottlenecks and growth characterizing

patient infection. One of the key methods of differentiating between these alternative

explanations is through comparisons between empirical data and plausible simulated data (Irwin

et al. 2016, Johri et al. 2020, Jensen 2021). However, given that the possible model and

parameter space is essentially infinite, a key first step in developing an evolutionary baseline

model is determining the relevant parameter bounds.

Our study supports three main conclusions that help define a feasible parameter space

governing intra-host SARS-CoV-2 evolutionary dynamics. First, our results support recent

conclusions that SARS-CoV-2 transmission is generally associated with strong infection

bottlenecks of one or a few virions (Bendall et al. 2023). Specifically, we found a bottleneck of 1

to be the only bottleneck size tested that produced plausible models. However, the unexamined

space between 1 and 5 (the midpoint value for bottlenecks) may also produce plausible models

as they were not tested in this study. Regardless, a bottleneck of 1 to 4 virions conforms with the

recent empirical findings based on patterns of intra-host SARS-CoV-2 variation between likely

transmissions pairs (Bendall et al. 2023).

Secondly, our results highlight the importance of considering the pervasive effects of

purifying and background selection, via a consideration of the DFE, in constraining levels of

variation segregating above the 2% frequency threshold. The wide-spread production of strongly

deleterious mutations appears necessary to explain the apparent contradiction between the high

mutation rates of RNA viruses (Drake and Holland 1999, Elena and Sanjuán 2005, Jensen et al.

2020) and the generally low number of SNPs identified in patients ((Lythgoe et al. 2021,

Valesano et al. 2021, Wang et al. 2021, Bendall et al. 2023, Gu et al. 2023). Given that coding

regions comprise most of the SARS-CoV-2 genome, this result is not particularly surprising.

This importance may be observed in the simulated parameter sets: among the models that

accumulated only neutral mutations, 11.8% were plausible; among the models with primarily

neutral mutations, 23.9% were plausible; among the models with equivalent rates of neutral and

strongly deleterious mutations, 28.4% were plausible; and among the models with primarily

strongly deleterious mutations, 42.8% were plausible. Current best estimates of the true DFE

underlying SARS-CoV-2 intra-host evolution suggest a bimodal distribution of fitness effects

(Flynn et al. 2022, Terbot et al. 2023) with peaks centered around strongly deleterious and

neutral fitness impacts. At a minimum, between 15-20% of sites in the SARS-CoV-2 genome

seem to be entirely invariant (Neher 2022), providing a minimum estimate for the proportion of

strongly deleterious mutations in the DFE. However, other studies indicate this value is more

likely in the range of 40-50% (Flynn et al. 2022, Terbot et al. 2023). Therefore, while the DFE

most severely skewed towards strongly deleterious mutations produced the most plausible

models, it is unlikely to reflect the true proportion of strongly deleterious mutations. However, it

is notable that the two other DFE distributions reflecting current best understandings of the true

DFE both produced around twice as many viable models as models including only neutral

mutations, emphasizing the central importance of purifying selection in the evolution of SARS-

CoV-2 within patients.

Finally, lower progeny skew values produce more plausible models. Specifically, models

with progeny skew and containing either lower probabilities of skewed offspring events and/or

lower burst sizes were over-represented amongst acceptances (397/555 plausible models or

71.5%). Reproduction of SARS-CoV-2 within a host cell is generally non-lytic and instead

involves release of new virions through continuous budding (Bar-On et al. 2020; Park et al.

2020). The prominence of lower levels of progeny skew in plausible models is consistent with a

lack of a single large burst of reproduction (*i.e*, lower values of burst size). Alternatively, it may

be related to the production of sub-genomic RNA which are not packaged directly into offspring

virions, but through recombination, mutations present in their sequences can be incorporated into

offspring virions (*i.e.*, lower values of Ξ; Langsjoen et al. 2020, Gribble et al. 2021).

Further model differentiation is limited by current data available for intra-host variation

of SARS-CoV-2. The 2% allele frequency cutoff used in this study mirrors the cutoff used in

several previous studies (Table 4). While this criterion successfully narrowed the relevant

evolutionary parameter space in our simulations, the scarce number of SNPs are insufficient to

explore more sophisticated statistics related to the site frequency spectrum or linkage

disequilibrium. However, new mutations arising during a typical patient infection will be rare

and thus may be missed when applying a frequency-based filter. There is of course an inherent

trade-off between including lower frequency SNPs and reducing the confidence in individual

SNP calls (Jacot et al. 2021). This trade-off can be partially ameliorated via the generation of

high-quality and higher-depth sequencing. A rule of thumb proposed by Lauring (Lauring 2020)

suggests that coverage should be ten times the inverse of the variant's frequency. So, a coverage

depth of 1000 would be required to confidently detect variants at 1% of the population, a depth

of 2000 for a 0.5% variant, and so on.

These general bioinformatic recommendations assume that sequencing data will

effectively sample intra-host variation without bias. However, most SARS-CoV-2 sequencing

protocols rely on targeted PCR or probe-based enrichment of the SARS-CoV-2 genome to

reduce background contamination of host nucleotides. Targeted enrichment approaches have

been widely used in biology and are usually sensitive to any standing genetic variation that

impacts the binding efficacy of enrichment probes or primers (Mamanova et al. 2010, Jones and

Good 2015). Indeed, assay-dependent effects have been a constant concern during the

development SARS-CoV-2 sequence protocols as demonstrated by the recurrent need to redesign

enrichment primers (*e.g.*, the ARCTIC protocol) to fully sequence the genomes of newly

circulating VOCs (Ulhuq et al. 2023). SARS-CoV-2 targeted enrichment is also sensitive to low

viral loads (Lam et al. 2021), which could further limit understanding of changes in intra-host

diversity through the time course of an infection. Given these concerns, technical replication of

individual patient samples may be required to reliably detect and quantify the frequency of rare

variants.

In addition to limitations relating to the detection of rare alleles, our study was also

unable to consider the impacts of host compartmentalization (*i.e.*, localized subpopulations

within different organs and areas of organ systems) on intra-host genetic diversity.

Compartmentalization is likely an important factor for SARS-CoV-2 evolution in regard to the

production of viral reservoirs in prolonged infections or infection of immune-compromised

patients (Gonzalez-Reiche et al. 2023, Normandin et al. 2023). However, there is currently

insufficient information regarding the number and connectivity of compartments used by SARS-

CoV-2 to allow for this aspect of model complexity to be reasonably parameterized. As more

empirical evidence regarding compartmentalization becomes available, future simulation studies

may be able to incorporate compartmentalization to determine the impacts—specifically the

influence of gene flow and recombination among compartments—on the intra-host population

genetics of SARS-CoV-2.

With that said, based on the currently available data, our study has successfully

quantified areas of the SARS-CoV-2 evolutionary parameter space that are most plausible. In

addition, the simulation framework presented here may be utilized to compare against future

sequencing studies, which will likely enable a further narrowing of likely models. However, even

the relatively broad parameter space here identified may be utilized to more effectively screen

for newly emerging positively selected mutations (*e.g.,* those contributing to the rapid spread of

VOCs), and in so doing reduce the traditionally high false-positive rates traditionally associated

with such selection scans (Johri et al 2020, Johri et al. 2022a). This work will also likely be

useful in providing a baseline simulation for studies looking at intra-host population genetics of

SARS-CoV-2 over time within a single host. Such studies may also allow for greater

discrimination between baseline models while retaining the use of a robust, minimum minor

allele frequency by providing information on how genetic variation accumulates (or persists)

over the course of an infection within a single patient.

Materials and Methods

Simulations

We used the SLiM software package (v4.0.1, Haller and Messer 2023) to conduct forward-in-

time simulations. We performed all simulations using SLiM's non-Wright-Fisher tick cycle. To

represent the single-stranded, haploid genome of one metabolically-active virion, we simulated

genomes consisting of 30kb. Each tick of the simulation represented approximately one hour,

during which all virions in the population produce one 'child virion' excluding cases of progeny

skew (described further below). We ran four different, nested models to gain insights into the

importance of various population genetic factors. The simplest model, hereafter referred to as the

bottleneck model, consisted of four parameters: the mutation rate, the initial number of virions

drawn from the burn-in period (or bottleneck size), the carrying capacity of the host, and the

number of ticks over which the simulation runs *i.e.*, the time between infection and sampling

(infection duration). The other three models added key factors: recombination, recombination

plus progeny skew, and a full model adding recombination, progeny skew, and a DFE. The latter

describes the proportion new mutations that are strongly deleterious or neutral. The model and

parameter space are summarized in Table 1 and Table 2. We based the tested parameter ranges

upon the current literature, as recently described in Terbot et al. 2023 unless otherwise stated.

Progeny genomes added mutations according to a single, genome-wide mutation rate.

Contrary to SLiM's default behavior for mutations arising at the same site, the simulation

retained only the most recent mutation occurring in a genome. The initial bottleneck was

performed by drawing virions from a common burn-in simulation, and the size of this bottleneck

reflected the range reported in the literature (Popa et al. 2020, Braun et al. 2021, Lythgoe et al.

2021, Martin and Koelle 2021, Bendall et al. 2023). We based the levels used for carrying

capacity on the number of virions estimated at peak infection of 10^9 to 10^{11} (Bar-On et al. 2020,

Sender et al. 2021). Due to computational constraints, and because this figure represents a census

population size as opposed to an effective population size, we scaled the carrying capacity

downward. Owing to this scaling of population size, we scaled up the range of mutation and

recombination rates to maintain a constant population-scaled product. The duration of infection

was chosen to represent a range of potential times to sampling. Given most empirical studies

report sampling in terms of days from symptom onset (Table 4), and there is considerable

uncertainty regarding the incubation time of SARS-CoV-2 from initial infection to symptom

presentation (Bar-On et al. 2020, Lauer et al. 2020, Li et al. 2020, Du et al. 2022, Wu et al.

2022), this range extends from brief (7 days) to extended (42 days).

To model recombination, we used a single parameter for the genome-wide recombination

rate. For each virion, the simulation chose another random virion to serve as the recombination

partner; and the simulation used this recombination partner for all progeny produced by the focal

virion in that tick cycle, including those experiencing progeny skew. We used a multiple-merger

coalescent model of progeny skew (Irwin et al. 2016) which required two parameters: Ξ

corresponding to the probability of a virion having multiple reproduction events in a single tick,

and the size of this reproductive burst. Values of Ξ represent the eclipse time of a virion (10

hours, *i.e.*, a value of $\Xi = 0.1$) as the max rate (Bar-On et al. 2020) or the product of this eclipse

time and the number of virions present in a cell that is actively budding virions (*i.e.*, 100000

virions multiplied by 10 hours, or a value of $\Xi = 0.000001$) as the minimum rate (Sender et al.

2021). The geometric mean of the minimum and maximum value (0.001) was used as the

midpoint value. We selected the highest level of burst size by determining the largest possible

burst that could still be completed with the maximum levels of Ξ and carrying capacity and the

requested computing resources; we selected lower levels as proportions of that maximum value.

Finally, we parameterized the DFE as the ratio of strongly deleterious relative to neutral

mutations.

Simulations used a common, burn-in source population which was created using a full-

model simulation run for 25,000 ticks at the highest value of carrying capacity and the mid-level

value for all other parameters. We replicated each parameter combination 5 times. Therefore, we

ran a total of 12,096 model-parameter combinations (represented visually in Figure 1), resulting

in 60,480 replicates. To simulate variation in sequencing depth of empirical studies, we

independently sampled 100 and 1000 genomes and stored them as .ms files. A total of 26

replicates failed to complete due to requiring more than the allotted computational resources and

were excluded from further analysis (Supplemental Table 2).

Simulation Assessment

Using the scikit-allel package (Miles et al. 2021) and custom python script (Rossum and Drake

2009), we calculated a set of summary statistics for each .ms file. The primary statistic we used

to identify models resulting in levels of variation similar to that observed in empirical patient

samples was the total number of SNPs (often referred to as intra-host single nucleotide variations

or iSNVs in the SARS-CoV-2 literature). We compared the simulated data to multiple recent

studies that have sought to quantify the amount of intra-host variation in SARS-CoV-2 (Lythgoe

et al. 2021, Valesano et al. 2021, Wang et al. 2021, Bendall et al. 2023, Gu et al. 2023). To make

this comparison, we applied filtering criteria to the simulated data reflecting that implemented in

the empirical data (*i.e.,* removing variants segregating below 2% frequency). From this existing

literature, the number of SNPs found within an intra-host population of SARS-CoV-2 clearing

this frequency threshold tends to be 5 or fewer (Table 4). Additionally, we required each

parameter combination to clear this threshold for both the 100 and 1000 genome sampling

schemes. We performed model filtering, and figure creation was performed using a custom R

script (R Core Team 2022) and the packages ggplot2 (Wickham 2016) and gridExtra (Auguie

2017). All eidos, bash, python, and R scripts have been deposited on github

(https://github.com/jwterbot2/SARS-CoV-2_InitialBaselineModel)

Supplementary Material

Supplementary tables and figures are available in a linked file for the purposes of review.

Acknowledgements

Research reported in this manuscript was supported by the National Institute of General Medical

Sciences of the National Institutes of Health (NIH) under Award Number P30GM140963 (to

JMG and BSC). NIH awards R35GM124701 (BSC) and R35GM139383 (JDJ) also supported

this work. The funders had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript. We are grateful to Parul Johri and Vivak Soni for their

advice regarding simulation design. We would also like to thank the computing resources used

during this study: the common burn-in was run on the Agave computing cluster at ASU and

simulation replicates were run using the Open Science Pool resources (OSG 2006, Pordes et al.

2007, Sfiligoi et al. 2009).

Data Availability

Scripts and code used in this study along with a copy of data used to generate figures are

available online via github (https://github.com/jwterbot2/SARS-CoV-2_InitialBaselineModel).

Literature Cited

Auguie B. 2017. gridExtra: Miscellaneous Functions for 'Grid' Graphics. https://cran.r-

project.org/package=gridExtra.

Bar-On YM, Flamholz A, Phillips R, Milo R. 2020. SARS-CoV-2 (COVID-19) by the numbers.

eLife. 9. doi: 10.7554/eLife.57309.

Barton NH. 1998. The effect of hitch-hiking on neutral genealogies. Genetical Research. 72:123–

133. doi: 10.1017/S0016672398003462.

Bendall EE et al. 2023. Rapid transmission and tight bottlenecks constrain the evolution of

highly transmissible SARS-CoV-2 variants. Nat Commun. 14:272. doi: 10.1038/s41467-

023-36001-5.

Braun KM et al. 2021. Acute SARS-CoV-2 infections harbor limited within-host diversity and

transmit via tight transmission bottlenecks. PLoS Pathogens. 17:e1009849. doi:

10.1371/journal.ppat.1009849.

Charlesworth B, Jensen JD. 2021. Effects of selection at linked sites on patterns of genetic

variability. Annu Rev of Ecol Evol and Systematics. 52:177–197. doi: 10.1146/annurev-

ecolsys-010621-044528.

Corman VM, Muth D, Niemeyer D, Drosten C. 2018. Hosts and sources of endemic human

coronaviruses. In: *Advances in Viral Research*. Academic Press: Cambridge, MA pp.

163–188. doi: 10.1016/bs.aivir.2018.01.001.

Drake JW, Holland JJ. 1999. Mutation rates among RNA viruses. Proc Natl Acad Sci U S A.

96:13910–13913. doi: 10.1073/pnas.96.24.13910.

Du Z et al. 2022. Shorter serial intervals and incubation periods in SARS-CoV-2 variants than

the SARS-CoV-2 ancestral strain. J Travel Medicine. 29:1–3. doi: 10.1093/jtm/taac052.

Elena SF, Sanjuán R. 2005. Adaptive value of high mutation rates of RNA viruses: separating

causes from consequences. J Virology. 79:11555–11558. doi: 10.1128/JVI.79.18.11555-

11558.2005.

Flynn JM et al. 2022. Comprehensive fitness landscape of SARS-CoV-2 M^{pro} reveals insights

into viral resistance mechanisms. eLife. 11:1–27. doi: 10.7554/eLife.77433.

Gonzalez-Reiche AS et al. 2023. Sequential intrahost evolution and onward transmission of

SARS-CoV-2 variants. Nat Commun. 14:3235. doi: 10.1038/s41467-023-38867-x

Gribble J et al. 2021. The coronavirus proofreading exoribonuclease mediates extensive viral

recombination. PLoS Pathogens. 17:e1009226. doi: 10.1371/journal.ppat.1009226.

Gu H et al. 2023. Within-host genetic diversity of SARS-CoV-2 lineages in unvaccinated and

vaccinated individuals. Nat Commun. 14:1793. doi: 10.1038/s41467-023-37468-y.

Haller BC, Messer PW. 2023. SLiM 4: Multispecies eco-evolutionary modeling. The American

Naturalist. 201:E000–E000. doi: 10.1086/723601.

Hay SI, Murray CJL. 2023. Conflicting COVID-19 excess mortality estimates – Authors' reply.

The Lancet. 401:433–434. doi: 10.1016/S0140-6736(23)00227-1.

Institute for Health Metrics and Evaluation. 2022. Covid-19 Projections.

https://covid19.healthdata.org/global?view=cumulative-deaths&tab=trend. (July 13,

2023)

Irwin KK et al. 2016. On the importance of skewed offspring distributions and background

selection in virus population genetics. Heredity. 117:393–399. doi: 10.1038/hdy.2016.58.

Jacot D, Pillonel T, Greub G, Bertelli C. 2021. Assessment of SARS-CoV-2 genome sequencing:

quality criteria and low-frequency variants. J Clinical Microbiology. 59:1–10. doi:

10.1128/JCM.00944-21.

Jensen JD. 2021. Studying population genetic processes in viruses: From drug-resistance

evolution to patient infection dynamics. In: *Encyclopedia of Virology*. Elsevier pp. 227–

232. doi: 10.1016/B978-0-12-814515-9.00113-2.

Jensen JD, Stikeleather RA, Kowalik TF, Lynch M. 2020. Imposed mutational meltdown as an

antiviral strategy. Evolution. 74:2549–2559. doi: 10.1111/evo.14107.

Johri P et al. 2022a. Recommendations for improving statistical inference in population

genomics. PLoS Biology. 20:e3001669. doi: 10.1371/journal.pbio.3001669.

Johri P, Charlesworth B, Jensen JD. 2020. Toward an evolutionarily appropriate null model:

jointly inferring demography and purifying selection. Genetics. 215:173–192. doi:

10.1534/genetics.119.303002.

Johri P, Stephan W, Jensen JD. 2022b. Soft selective sweeps: Addressing new definitions,

evaluating competing models, and interpreting empirical outliers. PLoS Genetics.

18:e1010022. doi: 10.1371/journal.pgen.1010022.

Jones MR, Good JM. 2016. Targeted capture in evolutionary and ecological genomics. Mol Ecol.

25:185–202. doi: 10.1111/mec.13304.

Lam C et al. 2021. SARS-CoV-2 Genome sequencing methods differ in their abilities to detect

variants from low-viral-load samples. J Clinical Microbiology. 59. doi:

10.1128/JCM.01046-21.

Langsjoen RM, Muruato AE, Kunkel SR, Jaworski E, Routh A. 2020. Differential Alphavirus

defective RNA diversity between intracellular and extracellular compartments is driven

by subgenomic recombination events. mBio. 11:1–20. doi: 10.1128/mBio.00731-20.

Lauer SA et al. 2020. The incubation period of coronavirus disease 2019 (CoVID-19) from

publicly reported confirmed cases: Estimation and application. Annals of Internal

Medicine. 172:577–582. doi: 10.7326/M20-0504.

Lauring AS. 2020. Within-host viral diversity: A window into viral evolution. Annu Rev

Virology. 7:63–81. doi: 10.1146/annurev-virology-010320-061642.

Li Q et al. 2020. Early transmission dynamics in Wuhan, China, of novel coronavirus–infected

pneumonia. New England Journal of Medicine. 382:1199–1207. doi:

10.1056/nejmoa2001316.

Lythgoe KA et al. 2021. SARS-CoV-2 within-host diversity and transmission. Science. 372. doi:

10.1126/science.abg0821.

Mamanova L et al. 2010. Target-enrichment strategies for next-generation sequencing. Nat

Meth. 7:111–118. doi: 10.1038/nmeth.1419.

Martin MA, Koelle K. 2021. Comment on "Genomic epidemiology of superspreading events in

Austria reveals mutational dynamics and transmission properties of SARS-CoV-2". Sci

Transl Med. 13:eabh1803. doi: 10.1126/scitranslmed.abh1803.

McCrone JT et al. 2018. Stochastic processes constrain the within and between host evolution of

influenza virus. eLife. 7. doi: 10.7554/eLife.35962.

Miles A et al. 2021. cggh/scikit-allel: v1.3.3. doi: 10.5281/zenodo.4759368.

Msemburi W et al. 2023. The WHO estimates of excess mortality associated with the COVID-19

pandemic. Nature. 613:130–137. doi: 10.1038/s41586-022-05522-2.

Neher RA. 2022. Contributions of adaptation and purifying selection to SARS-CoV-2 evolution.

Virus Evol. 8:2022.08.22.504731. doi: 10.1093/ve/veac113.

Normandin E et al. 2023. High-depth sequencing characterization of viral dynamics across tissues in fatal COVID-19 reveals compartmentalized infection. Nat Commun. 14:574. doi: 10.1038/s41467-022-34256-y.

O'Toole Á et al. 2021. Assignment of epidemiological lineages in an emerging pandemic using the pangolin tool. Virus Evol. 7:1–9. doi: 10.1093/ve/veab064.

OSG. 2006. OSPool. doi: 10.21231/906P-4D78.

Park WB et al. 2020. Virus isolation from the first patient with sars-cov-2 in Korea. J Korean Medical Science. 35:10–14. doi: 10.3346/jkms.2020.35.e84.

Popa A et al. 2020. Genomic epidemiology of superspreading events in Austria reveals mutational dynamics and transmission properties of SARS-CoV-2. Science Translational Medicine. 12:2555. doi: 10.1126/scitranslmed.abe2555.

Pordes R et al. 2007. The open science grid. J Physics: Conference Series. 78:012057. doi: 10.1088/1742-6596/78/1/012057.

R Core Team. 2018. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing: Vienna, Austria https://www.R-project.org/.

Rambaut A et al. 2020. A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. Nat Microbiol. 5:1403–1407. doi: 10.1038/s41564-020-0770-5.

Sender R et al. 2021. The total number and mass of SARS-CoV-2 virions. Proc Natl Acad Sci U S A. 118:1–9. doi: 10.1073/pnas.2024815118.

Sfiligoi I et al. 2009. The pilot way to grid resources using glideinWMS. In: *2009 WRI World Congress on Computer Science and Information Engineering*. IEEE pp. 428–432. doi: 10.1109/CSIE.2009.950.

Terbot JW et al. 2023. Developing an appropriate evolutionary baseline model for the study of

SARS-CoV-2 patient samples. PLoS Pathogens. 19:e1011265. doi:

10.1371/journal.ppat.1011265.

Ulhuq FR et al. 2023. Analysis of the ARTIC V4 and V4.1 SARS-CoV-2 primers and their

impact on the detection of Omicron BA.1 and BA.2 lineage-defining mutations.

Microbial Genomics. 9:2022.12.01.22282842. doi: 10.1099/mgen.0.000991.

Valesano AL et al. 2020. Influenza B viruses exhibit lower within-host diversity than influenza

A viruses in human hosts. J Virology. 94. doi: 10.1128/JVI.01710-19.

Valesano AL et al. 2021. Temporal dynamics of SARS-CoV-2 mutation accumulation within

and across infected hosts. PLoS Pathogens. 17:e1009499. doi:

10.1371/journal.ppat.1009499.Van

Rossum G, Drake FL. 2009. Python 3 Reference Manual. doi: 10.5555/1593511.

Wang H et al. 2022. Estimating excess mortality due to the COVID-19 pandemic: a systematic

analysis of COVID-19-related mortality, 2020–21. The Lancet. 399:1513–1536. doi:

10.1016/S0140-6736(21)02796-3.

Wang Y et al. 2021. Intra-host variation and evolutionary dynamics of SARS-CoV-2 populations

in COVID-19 patients. Genome Medicine. 13:30. doi: 10.1186/s13073-021-00847-5.

Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. doi: 10.1007/978-3-319-24277-

4.

Wu Y et al. 2022. Incubation Period of COVID-19 Caused by Unique SARS-CoV-2 Strains.

JAMA Network Open. 5:e2228008. doi: 10.1001/jamanetworkopen.2022.28008.

Tables and Figures

Table 1. Outline of Model Complexity

Each row represents a nested model which includes all parameters in that row and from rows above it.

Number of parameters added by each stage is detailed in column "+k" and the specific parameters added

are described in the rightmost column.

Model	+k	Parameters
Bottleneck	4	Bottleneck, Mutation Rate (μ), Infection Duration, Carrying Capacity (K)
+ Recombination	1	Recombination Rate (R)
+ Progeny Skew	2	Probability of Multiple Coalescent Event (Ξ), Burst Size
+ DFE	1	Ratio of neutral to strongly deleterious mutations

Table 2. Parameter Values Used in Models

Each parameter, other than infection duration, had three possible levels: a low point estimate, a high point

estimate, and a value representing a midpoint value. The column labeled "Burn-In" details the parameter

levels used during the common burn-in; note that aside from infection duration and carrying capacity, all

parameter levels used were the midpoint level. The highest carrying capacity and an extended infection

duration were used to allow mutations to accumulate and begin to reach equilibrium as may be expected

across the entire metapopulation of a pathogen during a prolonged pandemic.

	Burn-In	Lowest	Low	Midpoint	High
Initial Bottleneck	N/A	N/A	1	5	100
Infection Duration	25e3	168 (7 days)	336 (14 days)	672 (28 Days)	1008 (42 Days)
Mutation Rate	2.135e-6	N/A	2.135e-7	2.135e-6	2.135e-5
Carrying Capacity	1e5	N/A	5e3	5e4	1e5
Recombination Rate	5.5e-5	N/A	1e-5	5.5e-5	1e-4
Progeny Skew Probability	0.003	N/A	0.0001	0.003	0.1
Progeny Skew Amount	100	N/A	20	100	200
DFE (Neutral:Deleterious)	1:1	N/A	4:1	1:1	1:4

Table 3: Number of Plausible Models Given Required Parameters and Model Complexity

Counts and percentage of nested models within a given set of values for the required parameters carrying capacity, mutation rate, and infection

duration detailed in the first three columns (the initial bottleneck for all accepted models was the low value or a single virion). Each set of required

parameters had a single bottleneck model (columns 4 and 5), 3 recombination models (columns 6 and 7), 27 progeny skew models (columns 8 and

9), and 81 full models (columns 10 and 11). The final two columns represent the count and percentage of accepted models (out of a possible 112)

given a set of required parameters.

Infection Duration	Mutation Rate (μ)	Carrying Capacity (K)	Bottleneck Count	Bottleneck Percent	Recomb. Count	Recomb. Percent	Prog. Skew Count	Prog. Skew Percent	Full Count	Full Percent	All Count	All Percent
Lowest	Low	Low	1	100.00%	2	66.70%	6	22.20%	55	67.90%	64	57.10%
Lowest	Low	Midpoint	1	100.00%	3	100.00%	14	51.90%	56	69.10%	74	66.10%
Lowest	Low	High	1	100.00%	2	66.70%	19	70.40%	39	48.10%	61	54.50%
Lowest	Midpoint	Midpoint	1	100.00%	3	100.00%	8	29.60%	37	45.70%	49	43.80%
Lowest	Midpoint	High	1	100.00%	3	100.00%	11	40.70%	43	53.10%	58	51.80%
Lowest	High	High	0	0.00%	0	0.00%	0	0.00%	3	3.70%	3	2.70%
Low	Low	Low	0	0.00%	0	0.00%	0	0.00%	28	34.60%	28	25.00%
Low	Low	Midpoint	1	100.00%	3	100.00%	10	37.00%	33	40.70%	47	42.00%
Low	Low	High	0	0.00%	3	100.00%	12	44.40%	40	49.40%	52	46.40%
Low	Midpoint	Midpoint	0	0.00%	0	0.00%	0	0.00%	9	11.10%	9	8.00%
Low	Midpoint	High	0	0.00%	3	100.00%	3	11.10%	21	25.90%	27	24.10%
Midpoint	Low	Low	0	0.00%	0	0.00%	0	0.00%	9	11.10%	9	8.00%
Midpoint	Low	Midpoint	1	100.00%	2	66.70%	1	3.70%	23	28.40%	27	24.10%
Midpoint	Low	High	1	100.00%	3	100.00%	9	33.30%	25	30.90%	38	33.90%
Midpoint	Midpoint	High	0	0.00%	0	0.00%	0	0.00%	3	3.70%	3	2.70%
High	Low	Low	0	0.00%	0	0.00%	0	0.00%	5	6.20%	5	4.50%
High	Low	Midpoint	0	0.00%	0	0.00%	0	0.00%	10	12.30%	10	8.90%
High	Low	High	1	100.00%	1	33.30%	0	0.00%	23	28.40%	25	22.30%

478   Table 4. Summary of Empirical Studies Referenced

479   Sampling schema and results from empirical studies that reported the number of intra-host SNPs identified in patient samples. Note that all studies

480   apart from Lythgoe et al. 2021 utilized serial sampling within at least one patient during their data collection; this did not impact their reported

481   number of intra-host SNPs, but did allow for confirmation of SNPs as true positives.

Paper	Time to Sample	MAF Filter	Intra-host SNPs
Lythgoe et al. 2021	"symptomatic individuals on admission to the hospital"	3%	1.4 (mean)
Valesano et al. 2021	-7 to 20 days post symptom onset	2%	1 (median); 0-2 (IQR)
Wang et al. 2021	10 to 37 days post-symptom onset (18.09 days (mean))	5%	7.33 (mean); 1-23 (Range)
Bendall et al. 2023	Symptoms and positive test within 7 days	2%*	~0.82 (mean); 0-5 (Range)
Gu et al. 2023	With 5 days of symptom onset (2.3 days (mean))	2.50%	10.05 (mean); 5 (median)

486   * Intra-host SNP had to clear MAF filter in two technical replicates

Figure 1. Representations of the total parameter space for models' required parameters. Each required parameter [*i.e.*, bottleneck size, infection duration, mutation rate (μ), and carrying capacity (K)] is represented by a node. Line color and shape correspond to parameter value: dotted, red is lowest; solid, orange is low (the lowest value for bottleneck size, μ, and K); dashed, green is the mid-point value; and dotted-dashed, blue is the high value. The asterisk (*) denotes the only area of parameter space that had plausible models: those with bottlenecks of 1 (see Figure 2).

Figure 2. Parameter space for all plausible models. Line color and shape correspond to the parameter

values as in Figure 1. For each set of required parameters, a box is included that contains the fraction of

plausible models out of the total number of models using that set. Further information on the specific

models which were plausible is detailed in Table 3.

Figure 3. Each line represents the range of filtered SNPs for a particular model using a sampling of 1000

genomes, and each point is the mean of that model's replicates. All models in this figure used the lowest

bottleneck size (*i.e.*, 1); the carrying capacity used increases in panels from left to right, and the value of

the mutation rate used increases in panels from top to bottom. Within each panel, dashed lines separate

models into subpanels with different infection durations, increasing from left to right. Within each

subpanel the order of the models is the same and is detailed in Supplemental Table 3. The blue, horizontal

line represents the threshold of 5 SNPs used in this study to accept or reject a potential model (note that

Y-axes differ by row). Models with a mean value lower than this line (but not zero) were accepted as

plausible. Similar figures for models using larger bottlenecks are available as Supplemental Figures 1 and

2.

 Supplemental Table 1. List of parameter levels for all plausible models.

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Low	Low	Low	Lowest	n/a	n/a	n/a	n/a
Low	Low	Low	Lowest	Low	n/a	n/a	n/a
Low	Low	Low	Lowest	Low	Low	Low	n/a
Low	Low	Low	Lowest	Low	Low	Mid	n/a
Low	Low	Low	Lowest	Low	Mid	Low	n/a
Low	Low	Low	Lowest	Low	Low	Low	4:1
Low	Low	Low	Lowest	Low	Low	Mid	4:1
Low	Low	Low	Lowest	Low	Low	High	4:1
Low	Low	Low	Lowest	Low	Mid	Low	4:1
Low	Low	Low	Lowest	Low	Mid	High	4:1
Low	Low	Low	Lowest	Low	Low	Low	1:1
Low	Low	Low	Lowest	Low	Low	Mid	1:1
Low	Low	Low	Lowest	Low	Low	High	1:1
Low	Low	Low	Lowest	Low	Mid	Low	1:1
Low	Low	Low	Lowest	Low	Mid	Mid	1:1
Low	Low	Low	Lowest	Low	Mid	High	1:1
Low	Low	Low	Lowest	Low	Low	Low	1:4
Low	Low	Low	Lowest	Low	Low	Mid	1:4
Low	Low	Low	Lowest	Low	Low	High	1:4
Low	Low	Low	Lowest	Low	Mid	Mid	1:4
Low	Low	Low	Lowest	Low	Mid	High	1:4
Low	Low	Low	Lowest	Low	High	Mid	1:4
Low	Low	Low	Lowest	Low	High	High	1:4
Low	Low	Low	Lowest	Mid	n/a	n/a	n/a
Low	Low	Low	Lowest	Mid	Low	Low	n/a
Low	Low	Low	Lowest	Mid	Mid	Low	n/a
Low	Low	Low	Lowest	Mid	Low	Low	4:1
Low	Low	Low	Lowest	Mid	Mid	Low	4:1
Low	Low	Low	Lowest	Mid	Mid	High	4:1
Low	Low	Low	Lowest	Mid	Low	Low	1:1
Low	Low	Low	Lowest	Mid	Low	Mid	1:1
Low	Low	Low	Lowest	Mid	Low	High	1:1
Low	Low	Low	Lowest	Mid	Mid	Low	1:1
Low	Low	Low	Lowest	Mid	Mid	High	1:1
Low	Low	Low	Lowest	Mid	Low	Low	1:4
Low	Low	Low	Lowest	Mid	Low	Mid	1:4
Low	Low	Low	Lowest	Mid	Low	High	1:4
Low	Low	Low	Lowest	Mid	Mid	Low	1:4
Low	Low	Low	Lowest	Mid	Mid	Mid	1:4

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Low	Low	Low	Lowest	Mid	Mid	High	1:4
Low	Low	Low	Lowest	Mid	High	Low	1:4
Low	Low	Low	Lowest	Mid	High	Mid	1:4
Low	Low	Low	Lowest	Mid	High	High	1:4
Low	Low	Low	Lowest	High	Low	Low	n/a
Low	Low	Low	Lowest	High	Low	Low	4:1
Low	Low	Low	Lowest	High	Low	High	4:1
Low	Low	Low	Lowest	High	Mid	Low	4:1
Low	Low	Low	Lowest	High	Mid	High	4:1
Low	Low	Low	Lowest	High	Low	Low	1:1
Low	Low	Low	Lowest	High	Low	Mid	1:1
Low	Low	Low	Lowest	High	Low	High	1:1
Low	Low	Low	Lowest	High	Mid	Low	1:1
Low	Low	Low	Lowest	High	Mid	Mid	1:1
Low	Low	Low	Lowest	High	Mid	High	1:1
Low	Low	Low	Lowest	High	High	Low	1:1
Low	Low	Low	Lowest	High	Low	Low	1:4
Low	Low	Low	Lowest	High	Low	Mid	1:4
Low	Low	Low	Lowest	High	Low	High	1:4
Low	Low	Low	Lowest	High	Mid	Low	1:4
Low	Low	Low	Lowest	High	Mid	Mid	1:4
Low	Low	Low	Lowest	High	Mid	High	1:4
Low	Low	Low	Lowest	High	High	Low	1:4
Low	Low	Low	Lowest	High	High	Mid	1:4
Low	Low	Low	Lowest	High	High	High	1:4
Low	Low	Mid	Lowest	n/a	n/a	n/a	n/a
Low	Low	Mid	Lowest	Low	n/a	n/a	n/a
Low	Low	Mid	Lowest	Low	Low	Mid	n/a
Low	Low	Mid	Lowest	Low	Mid	Mid	n/a
Low	Low	Mid	Lowest	Low	High	Low	n/a
Low	Low	Mid	Lowest	Low	Low	Mid	4:1
Low	Low	Mid	Lowest	Low	Low	High	4:1
Low	Low	Mid	Lowest	Low	Mid	Low	4:1
Low	Low	Mid	Lowest	Low	Mid	Mid	4:1
Low	Low	Mid	Lowest	Low	High	Low	4:1
Low	Low	Mid	Lowest	Low	High	Mid	4:1
Low	Low	Mid	Lowest	Low	High	High	4:1
Low	Low	Mid	Lowest	Low	Low	Low	1:1
Low	Low	Mid	Lowest	Low	Low	Mid	1:1
Low	Low	Mid	Lowest	Low	Mid	Low	1:1
Low	Low	Mid	Lowest	Low	Mid	Mid	1:1

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Low	Low	Mid	Lowest	Low	Mid	High	1:1
Low	Low	Mid	Lowest	Low	High	Low	1:1
Low	Low	Mid	Lowest	Low	High	Mid	1:1
Low	Low	Mid	Lowest	Low	High	High	1:1
Low	Low	Mid	Lowest	Low	Mid	Low	1:4
Low	Low	Mid	Lowest	Low	Mid	Mid	1:4
Low	Low	Mid	Lowest	Low	Mid	High	1:4
Low	Low	Mid	Lowest	Low	High	Mid	1:4
Low	Low	Mid	Lowest	Low	High	High	1:4
Low	Low	Mid	Lowest	Mid	n/a	n/a	n/a
Low	Low	Mid	Lowest	Mid	Low	Low	n/a
Low	Low	Mid	Lowest	Mid	Low	Mid	n/a
Low	Low	Mid	Lowest	Mid	Mid	Mid	n/a
Low	Low	Mid	Lowest	Mid	High	Low	n/a
Low	Low	Mid	Lowest	Mid	High	Mid	n/a
Low	Low	Mid	Lowest	Mid	Low	Low	4:1
Low	Low	Mid	Lowest	Mid	Low	Mid	4:1
Low	Low	Mid	Lowest	Mid	Low	High	4:1
Low	Low	Mid	Lowest	Mid	Mid	Mid	4:1
Low	Low	Mid	Lowest	Mid	Mid	High	4:1
Low	Low	Mid	Lowest	Mid	High	Low	4:1
Low	Low	Mid	Lowest	Mid	High	Mid	4:1
Low	Low	Mid	Lowest	Mid	High	High	4:1
Low	Low	Mid	Lowest	Mid	Low	Low	1:1
Low	Low	Mid	Lowest	Mid	Low	High	1:1
Low	Low	Mid	Lowest	Mid	Mid	Mid	1:1
Low	Low	Mid	Lowest	Mid	Mid	High	1:1
Low	Low	Mid	Lowest	Mid	High	Mid	1:1
Low	Low	Mid	Lowest	Mid	High	High	1:1
Low	Low	Mid	Lowest	Mid	Low	Low	1:4
Low	Low	Mid	Lowest	Mid	Mid	Mid	1:4
Low	Low	Mid	Lowest	Mid	Mid	High	1:4
Low	Low	Mid	Lowest	Mid	High	Mid	1:4
Low	Low	Mid	Lowest	Mid	High	High	1:4
Low	Low	Mid	Lowest	High	n/a	n/a	n/a
Low	Low	Mid	Lowest	High	Low	Low	n/a
Low	Low	Mid	Lowest	High	Low	Mid	n/a
Low	Low	Mid	Lowest	High	Mid	Low	n/a
Low	Low	Mid	Lowest	High	Mid	Mid	n/a
Low	Low	Mid	Lowest	High	High	Low	n/a
Low	Low	Mid	Lowest	High	High	High	n/a

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Low	Low	Mid	Lowest	High	Low	Low	4:1
Low	Low	Mid	Lowest	High	Low	Mid	4:1
Low	Low	Mid	Lowest	High	Low	High	4:1
Low	Low	Mid	Lowest	High	Mid	Mid	4:1
Low	Low	Mid	Lowest	High	High	Mid	4:1
Low	Low	Mid	Lowest	High	High	High	4:1
Low	Low	Mid	Lowest	High	Low	Mid	1:1
Low	Low	Mid	Lowest	High	Low	High	1:1
Low	Low	Mid	Lowest	High	Mid	High	1:1
Low	Low	Mid	Lowest	High	High	Low	1:1
Low	Low	Mid	Lowest	High	High	Mid	1:1
Low	Low	Mid	Lowest	High	High	High	1:1
Low	Low	Mid	Lowest	High	Low	Mid	1:4
Low	Low	Mid	Lowest	High	Mid	Mid	1:4
Low	Low	Mid	Lowest	High	Mid	High	1:4
Low	Low	Mid	Lowest	High	High	Mid	1:4
Low	Low	Mid	Lowest	High	High	High	1:4
Low	Low	High	Lowest	n/a	n/a	n/a	n/a
Low	Low	High	Lowest	Low	Low	Mid	n/a
Low	Low	High	Lowest	Low	Low	High	n/a
Low	Low	High	Lowest	Low	Mid	Low	n/a
Low	Low	High	Lowest	Low	Mid	Mid	n/a
Low	Low	High	Lowest	Low	High	Low	n/a
Low	Low	High	Lowest	Low	High	High	n/a
Low	Low	High	Lowest	Low	Low	Low	4:1
Low	Low	High	Lowest	Low	Mid	Low	4:1
Low	Low	High	Lowest	Low	Mid	High	4:1
Low	Low	High	Lowest	Low	High	Mid	4:1
Low	Low	High	Lowest	Low	High	High	4:1
Low	Low	High	Lowest	Low	Mid	Low	1:1
Low	Low	High	Lowest	Low	Mid	Mid	1:1
Low	Low	High	Lowest	Low	Mid	High	1:1
Low	Low	High	Lowest	Low	High	Mid	1:1
Low	Low	High	Lowest	Low	High	High	1:1
Low	Low	High	Lowest	Low	Mid	High	1:4
Low	Low	High	Lowest	Low	High	High	1:4
Low	Low	High	Lowest	Mid	n/a	n/a	n/a
Low	Low	High	Lowest	Mid	Low	Low	n/a
Low	Low	High	Lowest	Mid	Low	Mid	n/a
Low	Low	High	Lowest	Mid	Low	High	n/a
Low	Low	High	Lowest	Mid	Mid	Low	n/a

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Low	Low	High	Lowest	Mid	Mid	Mid	n/a
Low	Low	High	Lowest	Mid	Mid	High	n/a
Low	Low	High	Lowest	Mid	High	Mid	n/a
Low	Low	High	Lowest	Mid	High	High	n/a
Low	Low	High	Lowest	Mid	Low	Low	4:1
Low	Low	High	Lowest	Mid	Low	High	4:1
Low	Low	High	Lowest	Mid	Mid	Mid	4:1
Low	Low	High	Lowest	Mid	Mid	High	4:1
Low	Low	High	Lowest	Mid	High	Low	4:1
Low	Low	High	Lowest	Mid	High	Mid	4:1
Low	Low	High	Lowest	Mid	High	High	4:1
Low	Low	High	Lowest	Mid	Low	Low	1:1
Low	Low	High	Lowest	Mid	Low	Mid	1:1
Low	Low	High	Lowest	Mid	Low	High	1:1
Low	Low	High	Lowest	Mid	Mid	Mid	1:1
Low	Low	High	Lowest	Mid	Mid	High	1:1
Low	Low	High	Lowest	Mid	High	Mid	1:1
Low	Low	High	Lowest	Mid	Mid	High	1:4
Low	Low	High	Lowest	Mid	High	Mid	1:4
Low	Low	High	Lowest	High	n/a	n/a	n/a
Low	Low	High	Lowest	High	Low	Mid	n/a
Low	Low	High	Lowest	High	Mid	Mid	n/a
Low	Low	High	Lowest	High	Mid	High	n/a
Low	Low	High	Lowest	High	High	Low	n/a
Low	Low	High	Lowest	High	High	High	n/a
Low	Low	High	Lowest	High	Low	Low	4:1
Low	Low	High	Lowest	High	Low	Mid	4:1
Low	Low	High	Lowest	High	Mid	Mid	4:1
Low	Low	High	Lowest	High	Mid	High	4:1
Low	Low	High	Lowest	High	High	Low	4:1
Low	Low	High	Lowest	High	High	Mid	4:1
Low	Low	High	Lowest	High	High	High	4:1
Low	Low	High	Lowest	High	Low	Mid	1:1
Low	Low	High	Lowest	High	Mid	Mid	1:1
Low	Low	High	Lowest	High	Mid	High	1:1
Low	Low	High	Lowest	High	High	Mid	1:1
Low	Low	High	Lowest	High	High	High	1:1
Mid	Low	Mid	Lowest	n/a	n/a	n/a	n/a
Mid	Low	Mid	Lowest	Low	n/a	n/a	n/a
Mid	Low	Mid	Lowest	Low	Low	Low	n/a
Mid	Low	Mid	Lowest	Low	Mid	Low	n/a

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Mid	Low	Mid	Lowest	Low	Low	Low	4:1
Mid	Low	Mid	Lowest	Low	Low	Mid	4:1
Mid	Low	Mid	Lowest	Low	Low	High	4:1
Mid	Low	Mid	Lowest	Low	Mid	Low	4:1
Mid	Low	Mid	Lowest	Low	Low	Low	1:1
Mid	Low	Mid	Lowest	Low	Low	Mid	1:1
Mid	Low	Mid	Lowest	Low	Low	High	1:1
Mid	Low	Mid	Lowest	Low	Mid	Low	1:1
Mid	Low	Mid	Lowest	Low	Low	Low	1:4
Mid	Low	Mid	Lowest	Low	Low	Mid	1:4
Mid	Low	Mid	Lowest	Low	Mid	Low	1:4
Mid	Low	Mid	Lowest	Low	High	Low	1:4
Mid	Low	Mid	Lowest	Mid	n/a	n/a	n/a
Mid	Low	Mid	Lowest	Mid	Low	Low	n/a
Mid	Low	Mid	Lowest	Mid	Low	Mid	n/a
Mid	Low	Mid	Lowest	Mid	Mid	Low	n/a
Mid	Low	Mid	Lowest	Mid	Low	Low	4:1
Mid	Low	Mid	Lowest	Mid	Low	Mid	4:1
Mid	Low	Mid	Lowest	Mid	Mid	Low	4:1
Mid	Low	Mid	Lowest	Mid	Low	Low	1:1
Mid	Low	Mid	Lowest	Mid	Low	Mid	1:1
Mid	Low	Mid	Lowest	Mid	Low	High	1:1
Mid	Low	Mid	Lowest	Mid	Mid	Low	1:1
Mid	Low	Mid	Lowest	Mid	Low	Low	1:4
Mid	Low	Mid	Lowest	Mid	Low	Mid	1:4
Mid	Low	Mid	Lowest	Mid	Low	High	1:4
Mid	Low	Mid	Lowest	Mid	Mid	Low	1:4
Mid	Low	Mid	Lowest	Mid	Mid	Mid	1:4
Mid	Low	Mid	Lowest	Mid	High	Low	1:4
Mid	Low	Mid	Lowest	High	n/a	n/a	n/a
Mid	Low	Mid	Lowest	High	Low	Low	n/a
Mid	Low	Mid	Lowest	High	Low	Mid	n/a
Mid	Low	Mid	Lowest	High	Mid	Low	n/a
Mid	Low	Mid	Lowest	High	Low	Low	4:1
Mid	Low	Mid	Lowest	High	Low	Mid	4:1
Mid	Low	Mid	Lowest	High	Low	High	4:1
Mid	Low	Mid	Lowest	High	Mid	Low	4:1
Mid	Low	Mid	Lowest	High	Low	Low	1:1
Mid	Low	Mid	Lowest	High	Low	Mid	1:1
Mid	Low	Mid	Lowest	High	Mid	Low	1:1
Mid	Low	Mid	Lowest	High	Low	Low	1:4

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Mid	Low	Mid	Lowest	High	Low	Mid	1:4
Mid	Low	Mid	Lowest	High	Low	High	1:4
Mid	Low	Mid	Lowest	High	Mid	Low	1:4
Mid	Low	Mid	Lowest	High	High	Low	1:4
Mid	Low	High	Lowest	n/a	n/a	n/a	n/a
Mid	Low	High	Lowest	Low	n/a	n/a	n/a
Mid	Low	High	Lowest	Low	Low	Low	n/a
Mid	Low	High	Lowest	Low	Low	Mid	n/a
Mid	Low	High	Lowest	Low	Low	High	n/a
Mid	Low	High	Lowest	Low	Mid	Low	n/a
Mid	Low	High	Lowest	Low	Low	Low	4:1
Mid	Low	High	Lowest	Low	Low	Mid	4:1
Mid	Low	High	Lowest	Low	Low	High	4:1
Mid	Low	High	Lowest	Low	Mid	Low	4:1
Mid	Low	High	Lowest	Low	High	Low	4:1
Mid	Low	High	Lowest	Low	Low	Low	1:1
Mid	Low	High	Lowest	Low	Low	Mid	1:1
Mid	Low	High	Lowest	Low	Low	High	1:1
Mid	Low	High	Lowest	Low	Mid	Low	1:1
Mid	Low	High	Lowest	Low	High	Low	1:1
Mid	Low	High	Lowest	Low	Low	Low	1:4
Mid	Low	High	Lowest	Low	Low	Mid	1:4
Mid	Low	High	Lowest	Low	Low	High	1:4
Mid	Low	High	Lowest	Low	Mid	Low	1:4
Mid	Low	High	Lowest	Low	Mid	Mid	1:4
Mid	Low	High	Lowest	Low	High	Low	1:4
Mid	Low	High	Lowest	Mid	n/a	n/a	n/a
Mid	Low	High	Lowest	Mid	Low	Low	n/a
Mid	Low	High	Lowest	Mid	Low	Mid	n/a
Mid	Low	High	Lowest	Mid	Mid	Low	n/a
Mid	Low	High	Lowest	Mid	Low	Low	4:1
Mid	Low	High	Lowest	Mid	Low	Mid	4:1
Mid	Low	High	Lowest	Mid	Low	High	4:1
Mid	Low	High	Lowest	Mid	Mid	Low	4:1
Mid	Low	High	Lowest	Mid	Low	Low	1:1
Mid	Low	High	Lowest	Mid	Low	Mid	1:1
Mid	Low	High	Lowest	Mid	Low	High	1:1
Mid	Low	High	Lowest	Mid	Mid	Low	1:1
Mid	Low	High	Lowest	Mid	Low	Mid	1:4
Mid	Low	High	Lowest	Mid	Low	High	1:4
Mid	Low	High	Lowest	Mid	Mid	Low	1:4

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Mid	Low	High	Lowest	Mid	Mid	Mid	1:4
Mid	Low	High	Lowest	Mid	High	Mid	1:4
Mid	Low	High	Lowest	High	n/a	n/a	n/a
Mid	Low	High	Lowest	High	Low	Low	n/a
Mid	Low	High	Lowest	High	Low	Mid	n/a
Mid	Low	High	Lowest	High	Low	High	n/a
Mid	Low	High	Lowest	High	Mid	Low	n/a
Mid	Low	High	Lowest	High	Low	Low	4:1
Mid	Low	High	Lowest	High	Low	Mid	4:1
Mid	Low	High	Lowest	High	Low	High	4:1
Mid	Low	High	Lowest	High	Mid	Low	4:1
Mid	Low	High	Lowest	High	Low	Low	1:1
Mid	Low	High	Lowest	High	Low	Mid	1:1
Mid	Low	High	Lowest	High	Low	High	1:1
Mid	Low	High	Lowest	High	Mid	Low	1:1
Mid	Low	High	Lowest	High	High	Low	1:1
Mid	Low	High	Lowest	High	Low	Low	1:4
Mid	Low	High	Lowest	High	Low	Mid	1:4
Mid	Low	High	Lowest	High	Low	High	1:4
Mid	Low	High	Lowest	High	Mid	Mid	1:4
Mid	Low	High	Lowest	High	High	Low	1:4
High	Low	High	Lowest	Low	Low	Low	1:4
High	Low	High	Lowest	Mid	Low	Low	1:4
High	Low	High	Lowest	High	Low	Mid	1:4
Low	Low	Low	Low	Low	Mid	High	1:1
Low	Low	Low	Low	Low	Low	Low	1:4
Low	Low	Low	Low	Low	Low	Mid	1:4
Low	Low	Low	Low	Low	Low	High	1:4
Low	Low	Low	Low	Low	Mid	Low	1:4
Low	Low	Low	Low	Low	Mid	Mid	1:4
Low	Low	Low	Low	Low	Mid	High	1:4
Low	Low	Low	Low	Low	High	Low	1:4
Low	Low	Low	Low	Low	High	Mid	1:4
Low	Low	Low	Low	Low	High	High	1:4
Low	Low	Low	Low	Mid	Mid	High	1:1
Low	Low	Low	Low	Mid	Low	Low	1:4
Low	Low	Low	Low	Mid	Low	Mid	1:4
Low	Low	Low	Low	Mid	Low	High	1:4
Low	Low	Low	Low	Mid	Mid	Low	1:4
Low	Low	Low	Low	Mid	Mid	Mid	1:4
Low	Low	Low	Low	Mid	Mid	High	1:4

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Low	Low	Low	Low	Mid	High	Low	1:4
Low	Low	Low	Low	Mid	High	Mid	1:4
Low	Low	Low	Low	Mid	High	High	1:4
Low	Low	Low	Low	High	Mid	High	1:1
Low	Low	Low	Low	High	Low	Low	1:4
Low	Low	Low	Low	High	Low	Mid	1:4
Low	Low	Low	Low	High	Low	High	1:4
Low	Low	Low	Low	High	Mid	Low	1:4
Low	Low	Low	Low	High	Mid	High	1:4
Low	Low	Low	Low	High	High	Low	1:4
Low	Low	Low	Low	High	High	High	1:4
Low	Low	Mid	Low	n/a	n/a	n/a	n/a
Low	Low	Mid	Low	Low	n/a	n/a	n/a
Low	Low	Mid	Low	Low	Low	Low	n/a
Low	Low	Mid	Low	Low	Low	Mid	n/a
Low	Low	Mid	Low	Low	Low	High	n/a
Low	Low	Mid	Low	Low	Mid	Low	n/a
Low	Low	Mid	Low	Low	Low	Mid	4:1
Low	Low	Mid	Low	Low	Low	High	4:1
Low	Low	Mid	Low	Low	Mid	Low	4:1
Low	Low	Mid	Low	Low	Low	Mid	1:1
Low	Low	Mid	Low	Low	Low	High	1:1
Low	Low	Mid	Low	Low	Mid	Low	1:1
Low	Low	Mid	Low	Low	High	Low	1:1
Low	Low	Mid	Low	Low	Low	High	1:4
Low	Low	Mid	Low	Low	Mid	Mid	1:4
Low	Low	Mid	Low	Low	Mid	High	1:4
Low	Low	Mid	Low	Low	High	Low	1:4
Low	Low	Mid	Low	Mid	n/a	n/a	n/a
Low	Low	Mid	Low	Mid	Low	Low	n/a
Low	Low	Mid	Low	Mid	Low	Mid	n/a
Low	Low	Mid	Low	Mid	Low	High	n/a
Low	Low	Mid	Low	Mid	Low	Low	4:1
Low	Low	Mid	Low	Mid	Low	Mid	4:1
Low	Low	Mid	Low	Mid	Low	High	4:1
Low	Low	Mid	Low	Mid	Mid	Low	4:1
Low	Low	Mid	Low	Mid	High	Low	1:1
Low	Low	Mid	Low	Mid	Low	High	1:4
Low	Low	Mid	Low	Mid	Mid	Mid	1:4
Low	Low	Mid	Low	Mid	Mid	High	1:4
Low	Low	Mid	Low	Mid	High	Low	1:4

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Low	Low	Mid	Low	Mid	High	Mid	1:4
Low	Low	Mid	Low	Mid	High	High	1:4
Low	Low	Mid	Low	High	n/a	n/a	n/a
Low	Low	Mid	Low	High	Low	Low	n/a
Low	Low	Mid	Low	High	Low	Mid	n/a
Low	Low	Mid	Low	High	Low	High	n/a
Low	Low	Mid	Low	High	Low	Mid	4:1
Low	Low	Mid	Low	High	Low	High	4:1
Low	Low	Mid	Low	High	Mid	Low	4:1
Low	Low	Mid	Low	High	Low	Low	1:1
Low	Low	Mid	Low	High	Mid	Low	1:1
Low	Low	Mid	Low	High	Mid	Mid	1:1
Low	Low	Mid	Low	High	Mid	Mid	1:4
Low	Low	Mid	Low	High	Mid	High	1:4
Low	Low	Mid	Low	High	High	Low	1:4
Low	Low	Mid	Low	High	High	Mid	1:4
Low	Low	Mid	Low	High	High	High	1:4
Low	Low	High	Low	Low	n/a	n/a	n/a
Low	Low	High	Low	Low	Low	Low	n/a
Low	Low	High	Low	Low	Low	Mid	n/a
Low	Low	High	Low	Low	Mid	Low	n/a
Low	Low	High	Low	Low	High	Low	n/a
Low	Low	High	Low	Low	Low	Low	4:1
Low	Low	High	Low	Low	Mid	Mid	4:1
Low	Low	High	Low	Low	High	Low	4:1
Low	Low	High	Low	Low	Low	High	1:1
Low	Low	High	Low	Low	Mid	Low	1:1
Low	Low	High	Low	Low	Mid	Mid	1:1
Low	Low	High	Low	Low	High	Low	1:1
Low	Low	High	Low	Low	Mid	Mid	1:4
Low	Low	High	Low	Low	Mid	High	1:4
Low	Low	High	Low	Low	High	Low	1:4
Low	Low	High	Low	Low	High	Mid	1:4
Low	Low	High	Low	Low	High	High	1:4
Low	Low	High	Low	Mid	n/a	n/a	n/a
Low	Low	High	Low	Mid	Low	Low	n/a
Low	Low	High	Low	Mid	Low	High	n/a
Low	Low	High	Low	Mid	Mid	Low	n/a
Low	Low	High	Low	Mid	High	Low	n/a
Low	Low	High	Low	Mid	Low	Low	4:1
Low	Low	High	Low	Mid	Low	Mid	4:1

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Low	Low	High	Low	Mid	Low	High	4:1
Low	Low	High	Low	Mid	Mid	Low	4:1
Low	Low	High	Low	Mid	Mid	Mid	4:1
Low	Low	High	Low	Mid	High	Low	4:1
Low	Low	High	Low	Mid	Low	Mid	1:1
Low	Low	High	Low	Mid	Mid	Low	1:1
Low	Low	High	Low	Mid	Mid	Mid	1:1
Low	Low	High	Low	Mid	High	Low	1:1
Low	Low	High	Low	Mid	High	Mid	1:1
Low	Low	High	Low	Mid	Low	Low	1:4
Low	Low	High	Low	Mid	Mid	Mid	1:4
Low	Low	High	Low	Mid	Mid	High	1:4
Low	Low	High	Low	Mid	High	Mid	1:4
Low	Low	High	Low	Mid	High	High	1:4
Low	Low	High	Low	High	n/a	n/a	n/a
Low	Low	High	Low	High	Low	Low	n/a
Low	Low	High	Low	High	Low	Mid	n/a
Low	Low	High	Low	High	Mid	Low	n/a
Low	Low	High	Low	High	High	Low	n/a
Low	Low	High	Low	High	Low	Mid	4:1
Low	Low	High	Low	High	Mid	Mid	4:1
Low	Low	High	Low	High	High	Low	4:1
Low	Low	High	Low	High	Low	Low	1:1
Low	Low	High	Low	High	Low	Mid	1:1
Low	Low	High	Low	High	Mid	Mid	1:1
Low	Low	High	Low	High	High	High	1:1
Low	Low	High	Low	High	Low	High	1:4
Low	Low	High	Low	High	Mid	Mid	1:4
Low	Low	High	Low	High	Mid	High	1:4
Low	Low	High	Low	High	High	Mid	1:4
Low	Low	High	Low	High	High	High	1:4
Mid	Low	Mid	Low	Low	Low	Low	1:4
Mid	Low	Mid	Low	Low	Low	Mid	1:4
Mid	Low	Mid	Low	Low	Mid	Low	1:4
Mid	Low	Mid	Low	Mid	Low	Low	1:4
Mid	Low	Mid	Low	Mid	Low	Mid	1:4
Mid	Low	Mid	Low	Mid	Mid	Low	1:4
Mid	Low	Mid	Low	High	Low	Low	1:4
Mid	Low	Mid	Low	High	Low	Mid	1:4
Mid	Low	Mid	Low	High	Mid	Low	1:4
Mid	Low	High	Low	Low	n/a	n/a	n/a

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Mid	Low	High	Low	Low	Low	Low	n/a
Mid	Low	High	Low	Low	Low	Mid	n/a
Mid	Low	High	Low	Low	Low	Low	4:1
Mid	Low	High	Low	Low	Low	Low	1:1
Mid	Low	High	Low	Low	Low	Low	1:4
Mid	Low	High	Low	Mid	n/a	n/a	n/a
Mid	Low	High	Low	Mid	Low	Low	n/a
Mid	Low	High	Low	Mid	Low	Low	4:1
Mid	Low	High	Low	Mid	Low	Mid	4:1
Mid	Low	High	Low	Mid	Low	Low	1:1
Mid	Low	High	Low	Mid	Low	Mid	1:1
Mid	Low	High	Low	Mid	Mid	Low	1:1
Mid	Low	High	Low	Mid	Low	Low	1:4
Mid	Low	High	Low	Mid	Low	Mid	1:4
Mid	Low	High	Low	Mid	Low	High	1:4
Mid	Low	High	Low	Mid	Mid	Low	1:4
Mid	Low	High	Low	High	n/a	n/a	n/a
Mid	Low	High	Low	High	Low	Low	4:1
Mid	Low	High	Low	High	Low	Low	1:1
Mid	Low	High	Low	High	Low	Mid	1:1
Mid	Low	High	Low	High	Mid	Low	1:1
Mid	Low	High	Low	High	Low	Low	1:4
Mid	Low	High	Low	High	Low	Mid	1:4
Mid	Low	High	Low	High	Low	High	1:4
Mid	Low	High	Low	High	Mid	Low	1:4
Mid	Low	High	Low	High	High	Low	1:4
Low	Low	Low	Mid	Low	Mid	High	1:1
Low	Low	Low	Mid	Low	Mid	Mid	1:4
Low	Low	Low	Mid	Low	Mid	High	1:4
Low	Low	Low	Mid	Mid	Mid	High	1:1
Low	Low	Low	Mid	Mid	Mid	High	1:4
Low	Low	Low	Mid	High	Mid	High	1:1
Low	Low	Low	Mid	High	Low	High	1:4
Low	Low	Low	Mid	High	Mid	High	1:4
Low	Low	Low	Mid	High	High	Mid	1:4
Low	Low	Mid	Mid	n/a	n/a	n/a	n/a
Low	Low	Mid	Mid	Low	n/a	n/a	n/a
Low	Low	Mid	Mid	Low	Low	Low	n/a
Low	Low	Mid	Mid	Low	Low	Low	4:1
Low	Low	Mid	Mid	Low	Mid	Low	4:1
Low	Low	Mid	Mid	Low	Low	Low	1:1

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Low	Low	Mid	Mid	Low	Low	Mid	1:1
Low	Low	Mid	Mid	Low	Low	High	1:1
Low	Low	Mid	Mid	Low	Mid	Low	1:1
Low	Low	Mid	Mid	Low	Low	Low	1:4
Low	Low	Mid	Mid	Low	Low	Mid	1:4
Low	Low	Mid	Mid	Low	Low	High	1:4
Low	Low	Mid	Mid	Low	Mid	Low	1:4
Low	Low	Mid	Mid	Mid	Low	Low	4:1
Low	Low	Mid	Mid	Mid	Mid	Low	4:1
Low	Low	Mid	Mid	Mid	Low	Low	1:1
Low	Low	Mid	Mid	Mid	Low	Mid	1:1
Low	Low	Mid	Mid	Mid	Mid	Low	1:1
Low	Low	Mid	Mid	Mid	Low	Mid	1:4
Low	Low	Mid	Mid	Mid	Low	High	1:4
Low	Low	Mid	Mid	Mid	Mid	Low	1:4
Low	Low	Mid	Mid	High	n/a	n/a	n/a
Low	Low	Mid	Mid	High	Low	Mid	1:1
Low	Low	Mid	Mid	High	Low	Low	1:4
Low	Low	Mid	Mid	High	Low	Mid	1:4
Low	Low	Mid	Mid	High	Low	High	1:4
Low	Low	Mid	Mid	High	Mid	Low	1:4
Low	Low	High	Mid	n/a	n/a	n/a	n/a
Low	Low	High	Mid	Low	n/a	n/a	n/a
Low	Low	High	Mid	Low	Low	Mid	n/a
Low	Low	High	Mid	Low	Mid	Low	n/a
Low	Low	High	Mid	Low	Low	Mid	4:1
Low	Low	High	Mid	Low	Low	High	4:1
Low	Low	High	Mid	Low	Low	Low	1:1
Low	Low	High	Mid	Low	Low	Mid	1:1
Low	Low	High	Mid	Low	Low	High	1:1
Low	Low	High	Mid	Low	Mid	Low	1:1
Low	Low	High	Mid	Low	Low	Mid	1:4
Low	Low	High	Mid	Low	Mid	Mid	1:4
Low	Low	High	Mid	Low	High	Low	1:4
Low	Low	High	Mid	Mid	n/a	n/a	n/a
Low	Low	High	Mid	Mid	Low	Low	n/a
Low	Low	High	Mid	Mid	Low	Mid	n/a
Low	Low	High	Mid	Mid	Mid	Low	n/a
Low	Low	High	Mid	Mid	Low	Low	4:1
Low	Low	High	Mid	Mid	Low	Low	1:1
Low	Low	High	Mid	Mid	Low	Mid	1:1

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Low	Low	High	Mid	Mid	Low	High	1:4
Low	Low	High	Mid	Mid	Mid	Mid	1:4
Low	Low	High	Mid	Mid	High	Low	1:4
Low	Low	High	Mid	High	n/a	n/a	n/a
Low	Low	High	Mid	High	Low	Low	n/a
Low	Low	High	Mid	High	Low	Mid	n/a
Low	Low	High	Mid	High	Low	High	n/a
Low	Low	High	Mid	High	Mid	Low	n/a
Low	Low	High	Mid	High	Low	Mid	4:1
Low	Low	High	Mid	High	Low	High	4:1
Low	Low	High	Mid	High	Mid	Low	4:1
Low	Low	High	Mid	High	Low	Low	1:1
Low	Low	High	Mid	High	Low	High	1:1
Low	Low	High	Mid	High	High	Low	1:1
Low	Low	High	Mid	High	Low	High	1:4
Low	Low	High	Mid	High	Mid	Low	1:4
Low	Low	High	Mid	High	Mid	Mid	1:4
Low	Low	High	Mid	High	High	Low	1:4
Mid	Low	High	Mid	Low	Low	Mid	1:4
Mid	Low	High	Mid	Mid	Low	Mid	1:4
Mid	Low	High	Mid	High	Low	Mid	1:4
Low	Low	Low	High	Low	Mid	Mid	1:4
Low	Low	Low	High	Low	Mid	High	1:4
Low	Low	Low	High	Mid	Mid	Mid	1:4
Low	Low	Low	High	Mid	Mid	High	1:4
Low	Low	Low	High	High	Mid	High	1:4
Low	Low	Mid	High	Low	Low	Low	1:4
Low	Low	Mid	High	Low	Low	Mid	1:4
Low	Low	Mid	High	Low	Mid	Low	1:4
Low	Low	Mid	High	Mid	Low	Low	1:4
Low	Low	Mid	High	Mid	Low	Mid	1:4
Low	Low	Mid	High	Mid	Low	High	1:4
Low	Low	Mid	High	Mid	Mid	Low	1:4
Low	Low	Mid	High	High	Low	Low	1:4
Low	Low	Mid	High	High	Low	Mid	1:4
Low	Low	Mid	High	High	Mid	Low	1:4
Low	Low	High	High	n/a	n/a	n/a	n/a
Low	Low	High	High	Low	Low	Low	4:1
Low	Low	High	High	Low	Low	Mid	4:1
Low	Low	High	High	Low	Low	Low	1:1
Low	Low	High	High	Low	Low	Mid	1:1

μ	Bottleneck Size	K	Infection Duration	Recombination	Ξ	Burst Size	DFE
Low	Low	High	High	Low	Mid	Low	1:1
Low	Low	High	High	Low	Low	Mid	1:4
Low	Low	High	High	Low	Low	High	1:4
Low	Low	High	High	Low	Mid	Low	1:4
Low	Low	High	High	Mid	Low	Low	4:1
Low	Low	High	High	Mid	Low	Low	1:1
Low	Low	High	High	Mid	Low	Mid	1:1
Low	Low	High	High	Mid	Mid	Low	1:1
Low	Low	High	High	Mid	Low	Low	1:4
Low	Low	High	High	Mid	Low	Mid	1:4
Low	Low	High	High	Mid	Low	High	1:4
Low	Low	High	High	Mid	Mid	Low	1:4
Low	Low	High	High	High	n/a	n/a	n/a
Low	Low	High	High	High	Low	Low	4:1
Low	Low	High	High	High	Low	Mid	1:1
Low	Low	High	High	High	Mid	Low	1:1
Low	Low	High	High	High	Low	Low	1:4
Low	Low	High	High	High	Low	Mid	1:4
Low	Low	High	High	High	Low	High	1:4
Low	Low	High	High	High	Mid	Low	1:4

515   Supplemental Table 2. List of models with replicates that failed to complete due to exceeding allocated

516   memory.

Mutation Rate (μ)	Carrying Capacity (K)	Bottleneck Size	Infection Duration	Recombination Rate (R)	Progeny Skew Probability (Ξ)	Burst Size	DFE	Missing Reps
High	High	Mid	High	Mid	High	High	n/a	1
High	High	Mid	Mid	High	High	High	n/a	4
High	High	High	Mid	High	High	High	n/a	5
High	High	Mid	High	High	High	High	n/a	2
High	High	High	High	High	High	High	n/a	5
High	High	High	Mid	High	High	High	4,1	1
High	High	Mid	High	High	High	High	4,1	5
High	High	High	High	High	High	High	4,1	3

518   Supplemental Table 3. Order of models presented in Figure 3, Supplemental Figure 1, and Supplemental

519   Figure 2 as lines (range) and points (mean value). Note that for each figure the mutation rate (μ), initial

520   bottleneck size, and carrying capacity (K) are the same for all models. The infection duration varies

521   between models as described in figure captions. Models with no completed replicates (Supplemental

522   Table 2) appear as gaps between neighboring models.

Order	R	Ξ	Burst Size	DFE
1	n/a	n/a	n/a	n/a
2	Low	n/a	n/a	n/a
3	Mid	n/a	n/a	n/a
4	High	n/a	n/a	n/a
5	Low	Low	Low	n/a
6	Mid	Low	Low	n/a
7	High	Low	Low	n/a
8	Low	Mid	Low	n/a
9	Mid	Mid	Low	n/a
10	High	Mid	Low	n/a
11	Low	High	Low	n/a
12	Mid	High	Low	n/a
13	High	High	Low	n/a
14	Low	Low	Mid	n/a
15	Mid	Low	Mid	n/a
16	High	Low	Mid	n/a
17	Low	Mid	Mid	n/a
18	Mid	Mid	Mid	n/a
19	High	Mid	Mid	n/a
20	Low	High	Mid	n/a
21	Mid	High	Mid	n/a
22	High	High	Mid	n/a
23	Low	Low	High	n/a
24	Mid	Low	High	n/a
25	High	Low	High	n/a
26	Low	Mid	High	n/a
27	Mid	Mid	High	n/a
28	High	Mid	High	n/a
29	Low	High	High	n/a
30	Mid	High	High	n/a
31	High	High	High	n/a
32	Low	Low	Low	4:1

Order	R	Ξ	Burst Size	DFE
33	Mid	Low	Low	4:1
34	High	Low	Low	4:1
35	Low	Mid	Low	4:1
36	Mid	Mid	Low	4:1
37	High	Mid	Low	4:1
38	Low	High	Low	4:1
39	Mid	High	Low	4:1
40	High	High	Low	4:1
41	Low	Low	Mid	4:1
42	Mid	Low	Mid	4:1
43	High	Low	Mid	4:1
44	Low	Mid	Mid	4:1
45	Mid	Mid	Mid	4:1
46	High	Mid	Mid	4:1
47	Low	High	Mid	4:1
48	Mid	High	Mid	4:1
49	High	High	Mid	4:1
50	Low	Low	High	4:1
51	Mid	Low	High	4:1
52	High	Low	High	4:1
53	Low	Mid	High	4:1
54	Mid	Mid	High	4:1
55	High	Mid	High	4:1
56	Low	High	High	4:1
57	Mid	High	High	4:1
58	High	High	High	4:1
59	Low	Low	Low	1:1
60	Mid	Low	Low	1:1
61	High	Low	Low	1:1
62	Low	Mid	Low	1:1
63	Mid	Mid	Low	1:1
64	High	Mid	Low	1:1
65	Low	High	Low	1:1
66	Mid	High	Low	1:1
67	High	High	Low	1:1
68	Low	Low	Mid	1:1
69	Mid	Low	Mid	1:1
70	High	Low	Mid	1:1
71	Low	Mid	Mid	1:1
72	Mid	Mid	Mid	1:1
73	High	Mid	Mid	1:1

Order	R	Ξ	Burst Size	DFE
74	Low	High	Mid	1:1
75	Mid	High	Mid	1:1
76	High	High	Mid	1:1
77	Low	Low	High	1:1
78	Mid	Low	High	1:1
79	High	Low	High	1:1
80	Low	Mid	High	1:1
81	Mid	Mid	High	1:1
82	High	Mid	High	1:1
83	Low	High	High	1:1
84	Mid	High	High	1:1
85	High	High	High	1:1
86	Low	Low	Low	1:4
87	Mid	Low	Low	1:4
88	High	Low	Low	1:4
89	Low	Mid	Low	1:4
90	Mid	Mid	Low	1:4
91	High	Mid	Low	1:4
92	Low	High	Low	1:4
93	Mid	High	Low	1:4
94	High	High	Low	1:4
95	Low	Low	Mid	1:4
96	Mid	Low	Mid	1:4
97	High	Low	Mid	1:4
98	Low	Mid	Mid	1:4
99	Mid	Mid	Mid	1:4
100	High	Mid	Mid	1:4
101	Low	High	Mid	1:4
102	Mid	High	Mid	1:4
103	High	High	Mid	1:4
104	Low	Low	High	1:4
105	Mid	Low	High	1:4
106	High	Low	High	1:4
107	Low	Mid	High	1:4
108	Mid	Mid	High	1:4
109	High	Mid	High	1:4
110	Low	High	High	1:4
111	Mid	High	High	1:4
112	High	High	High	1:4

524
Supplemental Figure 1. Each line represents the range of filtered SNPs for a particular model using a
sampling of 1000 genomes; each point is the mean of that model's replicates. All models in this figure
used the lowest bottleneck size (*i.e*., 5); panels increase from left to right in the value of carrying capacity
used and increase from top to bottom in the value of mutation rate used. Within each panel, dashed lines
separate models into subpanels with different infection durations, increasing from left to right. Within
each subpanel the order of the models is the same and is detailed in Supplemental Table 3. The blue,
horizontal line represents the threshold of 5 SNPs used in this study to accept or reject a potential model
(note that Y-axes differ by row).

533
Supplemental Figure 2. Each line represents the range of filtered SNPs for a particular model using a
sampling of 1000 genomes; each point is the mean of that model's replicates. All models in this figure
used the high bottleneck size (*i.e.*, 100); panels increase from left to right in the value of carrying capacity
used and increase from top to bottom in the value of mutation rate used. Within each panel, dashed lines
separate models into subpanels with different infection durations, increasing from left to right. Within
each subpanel the order of the models is the same and is detailed in Supplemental Table 3. The blue,
horizontal line represents the threshold of 5 SNPs used in this study to accept or reject a potential model
(note that Y-axes differ by row).

542