Bidirectional Regulation of Motor Circuits Using Magnetogenetic Gene Therapy

Authors: Santiago R. Unda[1], Roberta Marongiu[1], Lisa E. Pomeranz[2], Jonathan P. Dyke[3], Edward

K. Fung[3], Logan Grosenick[4], Rick Zirkel[5], Aldana M. Antoniazzi[1], Sofya Norman[1], Conor M.

Liston[4], Chris Schaffer[5], Nozomi Nishimura[5], Sarah A. Stanley[6,7], Jeffrey M. Friedman[2], Michael

G. Kaplitt[1*].

1. Laboratory of Molecular Neurosurgery, Department of Neurological Surgery, Weill
7   Cornell Medical College, Cornell University; New York, NY, USA.
8   2. Laboratory of Molecular Genetics, Rockefeller University; New York, NY, USA.
9   3. Citigroup Bioimaging Center, Weill Cornell Medical College, Cornell University; New
10   York, NY, USA.
11   4. Department of Psychiatry, Weill Cornell Medical College, Cornell University; New York,
12   NY, USA.
13   5. Meining School of Biomedical Engineering, Cornell University, Ithaca, NY, USA.
14   6. Diabetes, Obesity and Metabolism Institute, Icahn School of Medicine at Mount Sinai,
15   New York, NY, USA
16   7. Nash Family Department of Neuroscience and Friedman Brain Institute, Icahn School of
17   Medicine at Mount Sinai, New York, NY, USA
18
19   ***Corresponding author:** Michael G. Kaplitt (mik2002@med.cornell.edu)

Abstract

Regulating the activity of discrete neuronal populations in living mammals after delivery of modified ion channels can be used to map functional circuits and potentially treat neurological diseases. Here we report a novel suite of magnetogenetic tools, based on a single anti-ferritin nanobody-TRPV1 receptor fusion protein, which regulated neuronal activity in motor circuits when exposed to magnetic fields. AAV-mediated delivery of a cre-dependent nanobody-TRPV1 calcium channel into the striatum of adenosine 2a (A2a) receptor-cre driver mice led to restricted expression within D2 neurons, resulting in motor freezing when placed in a 3T MRI or adjacent to a transcranial magnetic stimulation (TMS) device. Functional imaging and fiber photometry both confirmed focal activation of the target region in response to the magnetic fields. Expression of the same construct in the striatum of wild-type mice along with a second injection of an AAVretro expressing cre into the globus pallidus led to similar circuit specificity and motor responses. Finally, a mutation was generated to gate chloride and inhibit neuronal activity. Expression of this variant in subthalamic nucleus (STN) projection neurons in PitX2-cre parkinsonian mice resulted in reduced local c-fos expression and a corresponding improvement in motor rotational behavior during magnetic field exposure. These data demonstrate that AAV delivery of magnetogenetic constructs can bidirectionally regulate activity of specific neuronal circuits non-invasively *in vivo* using clinically available devices for both preclinical analysis of circuit effects on behavior and potential human clinical translation.

Introduction

Neuromodulation has become a critical tool for both delineating functional brain circuits underlying behaviors in animal studies and treating human patients with circuit disorders [1]. Electrical stimulation has long been used in pre-clinical studies and deep brain stimulation (DBS) is a standard of care for patients with advanced tremors and Parkinson's disease (PD) [2]. Despite the focal effects of electrical stimulation, the need for greater precision in modulating specific circuits or neuronal populations has promoted the development of novel technologies. Optogenetics combines viral delivery of light-sensitive ion channels with regulated light probes for precise millisecond-time scale control of neural activity [3,4], but the microbial origin of the channels and the need for a fiber optic implant can limit applications in animal studies and provide challenges to human translation [5]. In contrast, chemogenetics delivers genes for modified channels or receptors to allow drug-regulated control of neural activity without an implant [6], but the response is dependent on drug diffusion to targeted regions and the time course and longevity of responses rely on the pharmacokinetics of the drug.

Precise spatial and temporal control of neural circuitry without implanted devices could be transformative for both pre-clinical studies and for human therapeutic applications [7]. *In vivo* magnetic-field-based stimulation to control neural activity has shown promise in mammalian [5,7-11] and non-mammalian systems [7,12]. Electromagnetic fields can penetrate tissue and transmit energy to metal/metal oxide particles which in turn gate receptors and ion channels [13]. Taking advantage of these properties, regulation of several brain structures has been achieved using endogenous or genetically encoded members of the transient receptor potential cation channel subfamily V (TRPV). Tethering of endogenous ferritin or external iron oxide nanoparticles (MNP)

[5] [7,9] has been shown to lead to alternations of neural activity by magnetic fields *in vitro* and *in*

vivo. Previously the use of this system required the use of adenovirus vectors to deliver TRPV1

and ferritin cDNA clones which were too large for a single AAV vector.

Here, we report highly cell-specific, temporally precise, remote, and reversible

modulation of deep brain structures in awake, freely moving mice using a novel nanobody (Nb)

to endogenous ferritin (Ft) fused to the TRPV1 channel (Nb-Ft-TRPV1) encoded by a construct

packaged into a single AAV vector[14]. We show that cre-dependent magnetogenetic activation of

striatopallidal neurons elicits robust motor freezing in Adora-2a (A2a)-cre mice, similar to that

reported using opto- and chemogenetics [15-17]. The magnetic fields were delivered using standard

MRI machines or a commercially available Transcranial Magnetic Stimulation (TMS) device used

for human therapy, with a threshold of 180 mT to induce behavior alterations. Neural activation

was further confirmed using expression of *c-fos*, positron emission tomography (PET), and *in vivo*

fiber photometry to validate magnetogenetic-induced changes of neural activation. Circuit-

specific activation of striatopallidal neurons was also achieved in wild type (wt) mice using a

retrograde AAV encoding cre recombinase in combination with the cre-conditional

magnetogenetic construct. Finally, we achieved significant motor improvements in parkinsonian

mice using non-invasive, magnetogenetic inhibition of subthalamic nucleus (STN) neurons with a

nanobody-TRPV1 construct modified to block neuronal activity. The approach that is presented

provides for a means for inducing bidirectional, real-time regulation of neuronal circuits in freely

moving animals in a magnetic field and may have potential for a variety of human therapeutic

applications.

Results

Magnetogenetic stimulation of striatal indirect pathway alters motor processing *in vivo.*

We targeted the indirect pathway *in vivo* using an AAV1/2 vector expressing an HA-tagged, cre-

dependent excitatory magnetogenetic Nb-Ft-TRPV1^{Ca2+} cassette flanked by a double floxed

inverted open reading frame (DIO) under the control of the JET promoter (AAV1/2-JET-DIO-Nb-

Ft-TRPV1^{Ca2+}-HA). The nanobody expressed from this construct has been shown previously to be

gated in the presence of a magnetic field via interactions between the anti-ferritin nanobody (Nb-

Ft) and endogenous ferritin with resulting neuronal activation, thus obviating the need for

delivering exogenous ferritin cDNAs[14].

92   The cre-dependent Nb-Ft-TRPV1^{Ca2+} AAV was injected bilaterally into the dorsal striatum

of transgenic mice expressing cre-recombinase in indirect spiny projection neurons (iSPN)

expressing dopamine type 2 (D2) receptors (A2a-cre mice) **(Fig. 1A upper panel)**. Immunostaining

for HA or RFP (for the AAV1/2-JET-DIO-mCherry control group) demonstrated efficient gene

expression in the dorsal striatum **(Fig. 1A lower panel)**. Histological analysis of transcript

expression using RNAScope confirmed that AAV transcripts co-localized only in D2 neurons **(Fig.**

1B). To determine if this new magnetogenetic construct could functionally activate iSPNs *in vivo*,

we first examined motor activity after placing animals in a direct magnetic field (DMF) **(Fig 1C)**.

Parkinson's disease is associated with increased iSPN activity and optogenetic activation of these

neurons in normal mice results in rapid freezing [15]. Bilateral DMF stimulation (500 mT-1.3 T) using

a 3T MRI magnet significantly decreased activity compared to baseline, with activity returning to

normal after removal from the DMF **(Fig. 1D)**. There was also a marked increase in the time spent

freezing (n=10, 20.3 ± 2.8 vs 1.2 ± 0.69 sec, p<0.001), and a decrease of ambulation (n=10, 23.0

± 5.1 vs 44.7 ± 1.9 sec, p<0.001) during DMF application in the Nb-Ft-TRPV1^{Ca2+} group placed in a

magnetic field compared to the mCherry group **(Fig. 1E & F)**.

To determine the threshold for magnetic field strength required to regulate striatal

cellular activity, we compared two different field gradients, 500 mT-1.3 T (High DMF titration)

(Supplementary Fig. 1A) and 100 mT-270 mT (Low DMF titration) as determined by placing a

gaussmeter in different locations in the cage at different distances from the 3T magnet **(Fig. 2A)**.

After mapping the magnetic field strength in the behavioral boxes **(Fig. 2B & Supplementary Fig.**

1B), we then recorded the locomotor activity in animals expressing the magnetogenetic

construct. The activity heatmap in the presence of the Low DMF revealed a consistent decrease

of activity only when the mice receiving Nb-Ft-TRPV1^{Ca2+} were in proximal sections of the

behavior box with a field strength at or above 180 mT **(Fig. 2B)**. This contrasted with animals

exposed to the High DMF which showed reduced activity across the entire box **(Supplementary**

Fig. 1C). Thus, while the freezing time was consistent throughout the box in the mice in the High

DMF titration **(Supplementary Fig. 1D)**, in the Low DMF titration there was a significant increase

in freezing time (n=5, p<0.05) only in the regions with >180 mT field strength **(Fig. 2B)**. This

threshold is consistent with a prior [10] report which used an earlier generation magnetogenetic

system consisting of separate TRPV1 and ferritin cDNAs to demonstrate that a DMF source of

~0.2 T was sufficient to regulate deep brain circuits involved in food intake and glucose

homeostasis. To further evaluate whether a magnetic field gradient over 180 mT is sufficient to

induce a robust motor phenotype, we repeated our behavioral assessment at low DMF and high

DMF **(Fig. 2C)**. Consistent with the prior results, we found that mice expressing Nb-Ft-TRPV1^{Ca2+}

stayed at a relatively fixed location during both Low and High DMF application while control mice

expressing mCherry continued to move normally during DMF conditions **(Fig. 2 D & E)**. Indirect

pathway activation with both High and Low DMF application decreased the time spent

ambulating (n=5-8, p<0.001), and increased time spent freezing (n=5-8, p<0.001). **(Fig. 2 F & G)**.

Magnetogenetic stimulation increases striatal neuronal activity *in vivo.*

We next assessed whether the changes in locomotor activity were correlated with changes in

neural activity. Previously the immediate early gene *c-fos* was shown to be increased in *vitro* and

in vivo in cells expressing TRPV1 and ferritin in a magnetic field [8,10,11]. Consistent with this, a

significantly higher proportion of striatal neurons expressing Nb-Ft-TRPV1^{Ca2+} showed c-*fos+*

expression compared to mCherry controls following exposure exposed to the MRI magnet (500

mT-1.3 T) (n=5, 23.65 ± 3.49 vs 5.79 ± 0.77 %, p<0.05) **(Fig. 3A & B)**. There was a correlation

between *c-fos* expression and freezing (n=4-5, Pearson's r=0.39, two-tailed n.s.) with the Nb-Ft-

TRPV1^{Ca2+} and mCherry mice clustering at opposite extremes of the correlation matrix **(Fig. 3C)**.

The data are consistent with there being a robust activation of the indirect pathway neurons,

with relatively low variability between the expression of gene activity-induced makers, and the

freezing phenotype.

To further assess striatal activity after magnetogenetic activation living animals, we used

positron emission tomography (PET) with [^{18}F] fluorodeoxyglucose (^{18}F-FDG), which measures

glucose utilization as a reflection of underlying neural activity [18]. Micro-PET-computed

tomography (CT) was performed in A2a mice expressing Nb-Ft-TRPV1^{Ca2+} fasted for 12 hrs. vs

mice receiving an mCherry in the striatum construct. ^{18}F-FDG was administered to the two groups

first without DMF exposure, followed 3 days later by PET imaging following the application of

DMF **(Fig. 3D upper panel)**. DMF exposure significantly increased striatal uptake of ^{18}F-FDG,

expressed as the ratio of uptake at baseline prior to DMF exposure to the magnetic field to the

signal after placement in a magnetic field. The ratio of the PET signal was significantly greater in

Nb-Ft-TRPV1^{Ca2+} group compared to mCherry controls (n=9-10, 1.14 ± 0.06 vs 0.98 ± 0.001,

p<0.05) **(Fig. 3D lower panel & E)**. To account for individual variability, we also quantified the

percentage of the ^{18}F-FDG dose that was injected dose gram (%ID/g) in the striatum compared

to uptake in cerebellum, with and without DMF treatment. DMF exposure significantly increased

^{18}F-FDG uptake in the striatum in mice with Nb-Ft-TRPV1^{Ca2+} expression in D2 neurons relative to

mCherry controls (n=10, 1.37 ± 0.06 vs 1.19 ± 0.03 %ID/g, p<0.001) **(Fig. 3F)**. No change in the

PET signal was seen in cerebellum. Since clinically ^{18}F-FDG PET-CT is used to monitor Parkinson's

disease progression and can be correlated with motor features [18], we reasoned that the levels of

striatal glucose uptake might correlate with the time mice spent freezing during DMF application.

Consistent with this, the data showed a significant positive correlation (n=7-9, Pearson's r=0.69,

two-tailed p<0.05) between striatal glucose uptake and freezing time in the Nb-Ft-TRPV1^{Ca2+}

expressing mice **(Fig. 3G)**, with the most marked increase in the time spent freezing observed in

animals with a ^{18}F-FDG uptake over 1.4 using the cerebellum as a reference.

Finally, we used genetically encoded calcium indicators to measure magnetogenetic-

induced neural activity [5,7,11,12]. To selectively record D2 iSPN calcium transients during pre-DMF,

DMF, and post-DMF periods, we bilaterally injected a cre-dependent excitatory magnetogenetic

vector (AAV1/2-JET-DIO- Nb-Ft-TRPV1^{Ca2+} -HA) together with a cre-dependent calcium indicator

(AAV9-Syn-DIO-GCaMP6s) and implanted an optical fiber in the dorsal striatum of A2a-cre mice

(Fig. 4A). This required the use of optical fibers that did not contain any metal which would heat

in the presence of a magnetic field. Nb-Ft-TRPV1^{Ca2+} expressing mice again showed motor

freezing in response to DMF that was correlated with an increased ΔF/F striatal GCaMP

fluorescence signal **(Fig. 4B)**, which was not seen in mCherry expressing mice **(Fig. 4D)**. Mean

GCaMP6 fluorescence (ΔF/F) percentage aligned to Pre-DMF, DMF, and Post-DMF revealed a

marked increase of fluorescence during DMF stimulation in the Nb-Ft-TRPV1^{Ca2+} group **(Fig. 4C)**

but not in the mCherry group **(Fig. 4E)**. Calcium transients in D2 iSPNs demonstrated by an

increase in GCaMP6 fluorescence (ΔF/F) were significantly increased by DMF treatment in the

Nb-Ft-TRPV1^{Ca2+} but not in the mCherry expressing mice (n=3-4, 27.91 ± 8.24 vs -1.37 ± 1.05%,

p<0.05) **(Fig. 4G)**. In parallel, DMF treatment significantly increased time spent freezing in Nb-Ft-

TRPV1^{Ca2+} compared to mCherry controls (n=3-4, p<0.05) confirming the functional efficacy of

magnetogenetic activation **(Fig. 4B & D)**. Consistent with the *c-fos* expression level and the PET

studies, there was a positive correlation between the change in fluorescent signal and the extent

of the motor inhibition (n=3-4, Pearson's r=0.60, two-tailed n.s.), with the experimental and

control groups mostly clustering at opposite ends of the correlation matrix **(Fig. 4H)**. This

combined data demonstrates that the fused ferritin nanobody-TRPV1 channel, expressed from a

single AAV in striatal D2 neurons, resulted in activation of transduced iSPNs which then caused

the resulting motor freezing.

Transcranial magnetic stimulation (TMS) increases Nb-Ft-TRPV1^{Ca2+} mediated calcium

transients and elicits parkinsonian phenotype.

While the MRI magnet was highly effective for demonstrating and characterizing the

magnetogenetic effect of the new construct on neurons within the brain, the feasibility of

translating into human application a technology which requires proximity to an MRI for

therapeutic efficacy is limited. To evaluate the potential of a more clinically relevant technology,

we repeated the studies above using a commercial transcranial magnetic stimulation (TMS)

system which is FDA approved for depression and other applications [19] **(Fig. 5A)**. Similar to the

effect of an MRI machine, TMS application at 20% output capacity using 50 twin pulses per

second with 1 second inter-stimulus intervals decreased motor ambulation in Nb-Ft-TRPV1^{Ca2+}

expressing mice compared with mCherry controls **(Fig. 5B)**. Analysis of motor parameters

including distance moved **(Fig. 5C & D)**, activity **(Supplementary Fig. 2A & B)**, and change of

location **(Supplementary Fig. 2D & E)** was consistent with activation of D2 iSPN in Nb-Ft-

TRPV1^{Ca2+} expressing mice compared to mCherry controls. Average distance moved and activity

percentage during TMS application was significantly decreased in the Nb-Ft-TRPV1^{Ca2+} expressing

mice (n=4-5, p<0.05) **(Fig. 5E, Supplementary Fig. 2C)**. Again, using targeted GCAMP expression

and fiber photometry, TMS treatment produced rapid generation of fluorescent positive spikes

in the striatum of Nb-Ft-TRPV1^{Ca2+} expressing mice while TMS treatment showed a sustained

decrease in GCaMP6 fluorescence in mCherry-expressing controls **(Fig. 5F & G)** [20]. Fluorescence

(ΔF/F) quantification showed a significant increase in the Nb-Ft-TRPV1^{Ca2+} expressing mice

compared to their own baseline (n=3, 14.41 ± 4.15 vs -2.38 ± 0.81%, p<0.05) and compared with

mCherry mice during TMS application (n=3-4, 14.41 ± 4.15 vs -15.81 ± 5.61%, p<0.05) **(Fig. 5H)**.

These studies confirm that, as was seen with MRI generated magnetic fields, a clinically approved

TMS device is capable of activating transduced neurons and eliciting motor freezing in Nb-Ft-

TRPV1^{Ca2+} expressing mice at well below the maximal output of the device.

Magnetogenetic stimulation of striatopallidal neurons in wild type mice reproduces a robust

motor phenotype.

Cre-driver mouse lines provide cell specificity for functional mapping of neural circuits underlying

numerous behaviors and many diseases, but this approach is limited to animals. In addition,

expression is based on cellular identity and not on functional anatomical connections [21]. In order

to determine if magnetogenetics could be restricted to specific circuits in normal mice, we

utilized a dual vector system to restrict expression to the striatopallidal iSPN circuitry. AAV1/2-

JET-DIO-Nb-Ft-TRPV1^{Ca2+}-HA was first injected bilaterally into the dorsal striatum of normal mice,

which would not lead to expression of the viral transgene due to the absence of cre. This was

followed by a second injection of a retrograde AAV vector carrying a cre recombinase fused to

GFP (RetroAAV2-CMV-Cre-GFP) into the globus pallidus (Gp). This approach allows retrograde

uptake of AAV into afferents to the Gp, with resulting cre expression in neurons which project to

Gp, including D2 iSPNs. Since the only pallidal afferents containing a floxed transgene with this

approach would be those from the dorsal striatum, this should lead to restricted expression of

the magnetogenetic construct only in striatopallidal neurons. As expected, immunostaining for

HA or RFP (for AAV1/2-JET-DIO and the mCherry control group, respectively) demonstrated co-

localization of each with GFP from the transduced SPNs **(Fig. 6 A)**. The same behavioral tests were

then performed as were used to previously characterize the transgenic A2a-cre mice (**Fig.1**).

Consistent with the prior results, there was a robust decrease in locomotor activity when the

animals receiving the AAV1/2-JET-DIO-Nb-Ft-TRPV1^{Ca2+}-HA injections were placed in a magnetic

field compared to controls (**Fig. 6B)**. There was a significant decrease in ambulation during DMF

application (n=8, 16.13 ± 3.02 vs 39.38 ± 3.28 sec, p<0.001), and significant increase in time spent

freezing (n=8, 23.75 ± 4.63 vs 4.13 ± 1.34 sec, p<0.001) **(Fig. 6C & D)**. *c-fos* expression was again

significantly higher in the transduced iSPNs of Nb-Ft-TRPV1^{Ca2+} expressing mice compared to

mCherry controls (n=4, 26.34 ± 3.13 vs 10.76 ± 2.41 %, p<0.001) **(Fig. 6E & F).** To control for the

possibility that striatal overexpression of TRPV1 ion channels alone could mediate activation of

the indirect pathway in the absence of ferritin tethered to the TRPV1 channel, we compared the

effects of DMF treatment on wt mice expressing either Nb-Ft-TRPV1^{Ca2+} or TRPV1^{Ca2+} alone **(Fig.**

6G). DMF application had no observable motor effect on TRPV1^{Ca2+} expressing mice but again

suppressed ambulation and increased freezing in mice expressing Nb-Ft-TRPV1^{Ca2+}, confirming

that fusion of the ferritin nanobody to the TRPV1 channel was required for neuronal activation

in the presence of a magnetic field **(Fig. 6H & I)**.

Magnetogenetic inhibition of subthalamic nucleus neurons improves motor impairment in

parkinsonian mice.

One advantage of chemogenetic and optogenetic systems is the availability of a suite of modified

channels which can inhibit neurons in addition to those which can activate neuronal firing [22].

Developing an inhibitory magnetogenetic system would not only expand the opportunities for

scientific research using this technology but would also provide translational opportunities for

diseases where neuronal inhibition would be desirable. We first validated *in vitro* the utility of

the anti-ferritin nanobody fused to a mutant TRPV1 channel modified to allow permeability of

Chloride (Cl-) ions (Nb-Ft-TRPV1^{Cl-}), which should increase chloride flux to inhibit neuronal activity

(Fig. 7A). Neuro2A cells were transfected with plasmids expressing mCherry, TRPV1^{Ca2+} or Nb-Ft-

TRPV1^{Cl-} for fluorescent chloride imaging using MQAE, which is quenched by intracellular chloride

(Supplementary Fig. 3A). Application of a low intensity magnetic field (MF, 230-250mT) showed

a significant decrease in the ΔF/F0 in the Nb-Ft-TRPV1^{Cl-} expressing cells compared to both

mCherry and TRPV1^{Ca2+} controls **(Fig. 7B)** (ΔF/F0 at 150s: Nb-Ft-TRPV^{Cl-} -0.16 ± 0.01,

TRPV1^{Ca2+}0.06 ± 0.01, mCherry -0.002 ± 0.01, p<0.001) consistent with increased intracellular

chloride concentrations.

The use of an inhibitory magnetogenetic system may be of potential use for symptomatic

treatment of Parkinson's disease (PD), where chemical or gene therapy-based inhibition of

subthalamic nucleus (STN) projection neurons can improve motor function in rodents, non-

human primates, and humans [23-27]. To test this, we generated unilateral 6-hydroxydopamine (6-

OHDA) lesions of nigral dopaminergic neurons in transgenic mice expressing the cre recombinase

in PitX-2-positive neurons (PitX-2-cre mice) **(Supplementary Fig. 3B)**, which restricts cre

expression to STN glutamatergic projection neurons with virtually no expression within other

populations in the vicinity [28]. Stable and functionally significant lesions were documented

through quantification of apomorphine-induced rotations, followed by randomization of animals

to balanced experimental and control groups and confirmation that DMF alone does not affect

rotations **(Supplementary Fig. 3C-D)**. A DIO cre-dependent AAV1/2 vector encoding the Nb-Ft-

TRPV1^{Cl-} cassette under the control of the JET promoter (AAV1/2-JET-DIO- Nb-Ft-TRPV1^{Cl-}-HA)

was then injected into the ipsilateral STN of 6OHDA-lesioned PitX2 mice, with another group

injected with a DIO cre-dependent mCherry vector as a negative control **(Fig. 7D)**. Mice

expressing Nb-Ft-TRPV1^{Cl-} in the STN showed a significant reduction of contralateral rotations

induced by the dopaminergic agonist, apomorphine, in the presence of a DMF application relative

to baseline and controls baseline (n=9-10, -46.62 ± 6.5 vs -2.67 ± 14.69%, p<0.05), **(Fig. 7E,**

Supplementary Fig. 3E & F). The number of *c-fos* positive transduced neurons in the STN of PitX2-

cre mice was significantly decreased following DMF exposure in the group which received the

inhibitory magnetogenetic Nb-Ft-TRPV1^{Cl-}vector compared to mCherry controls **(Fig. 7F & G)**

(n=4, 15.01 ± 2.83 vs 48.89 ± 6.11%, p<0.05). This confirms that AAV-mediated expression of the

magnetogenetic inhibitory construct exclusively within PitX2-positive STN projection neurons

resulted in reduction of neuronal activity *in vivo* and a consequent improvement in motor

function following 6OHDA lesioning of the substantia nigra.

Discussion

The demonstration that viral vectors such as AAV can be used for gene transfer in the

mammalian brain [29] has transformed pre-clinical neuroscience research. Furthermore, many

completed and active clinical trials of gene therapy in humans have shown promise for treating

circuit disorders [27,30]. While initially genes cloned into AAV were expressed constitutively, more

recently techniques such as optogenetics and chemogenetics have been used to regulate neural

activity in an inducible manner, using light or drugs respectively. Optogenetics provides

temporally rapid and precise control of neuronal function but utilizes microbial-derived channels

that can be immunogenic and require permanent implantation of light-generating devices that

potentially tether small animals and present engineering and safety challenges for human

applications [4,31]. Chemogenetics uses drugs to regulate channels, which obviates the need for a

device implant, but chemogenetic regulation occurs in a more delayed fashion based upon the

pharmacokinetics of the activator molecules [32,33]. In this report, we validate magnetogenetics as

an alternative approach for modulating neural activity and show that magnetic field-based

regulation of neuronal function can be used for bidirectional control of neuronal activity. This

method provides both anatomical and temporal precision in awake, freely moving animals

without the need for an implanted device and uses a viral vector potentially applicable for clinical

studies and can respond to clinically approved devices which deliver sufficient strength magnetic

fields.

We validated this system using variants of TRPV1, which are endogenous mammalian

channels, together with a nano-body that binds to mouse and human ferritin[14]. We tested this

approach by eliciting a parkinsonian phenotype in freely moving mice to demonstrate the

functional efficacy of magnetogenetic activation of striatal indirect medium spinal neurons.

Activation of these neurons in a magnetic field was evident only in mice receiving the

magnetogenetic construct compared with controls, as confirmed by *c-fos* expression post-

mortem as well as both FDG-PET and fiber photometry in living animals. Optogenetic control of

basal ganglia circuitry has previously demonstrated similar changes in motor behavior after

selective activation of indirect pathways *in vivo* using an implanted light device to control neurons

expressing Channelrhodopsin-2 (ChR2) [15]. More recently, Munshi and colleagues [5] reported

successful magnetogenetic modulation of motor circuits targeting motor cortex and dorsal

striatum. Consistent with the classical model of basal ganglia circuitry, this study reported

decrease in voluntary movement by activation of striatal D2 iSPN using a combination of metal

nanoparticles (MNPs) and AAVs encoding TRPV1 channels stereotactically injected 4 weeks apart

in the ridge between dorsal and ventral striatum [5]. However, synthetic magnetic nanoparticles

would not allow for stable, chronic magnetogenetic activation and the use of MNPs could

potentially tether to other thermo-sensitive channels causing off target activation and safety

issues [7,34]. In addition, the intrinsic variability of separate injections and non-cell specific

transgene transduction can limit conclusions regarding the relationship between phenotype and

cells modulated by the exogenous particle-based magnetogenetic system. This is in contrast to

the genetically encoded system which we employed, which only requires injection of a single AAV

to improve both dosing consistency between subjects and longevity of neuromodulation due to

chronic AAV-mediated expression. For human translation, the ability to deliver a single construct

with a single infusion has both safety and regulatory advantages, and here we demonstrate the

viability of AAV-mediated delivery of a single magnetogenetic construct encoding a fused Nb

capable of tethering endogenous ferritin to an excitatory TRPV1 channel for inducible modulation

of neuronal activity leading to definable motor phenotypes.

The ability use of optogenetics, through a combination of focal delivery of viral constructs

expressing opsins and implantation of light emitting devices in a target brain region, has led to a

wealth of knowledge regarding the relevance of neuronal circuits to particular behaviors [35]. The

potential benefit of the magnetogenetic system described here is the use of a non-invasive

magnetic field to regulate specific circuits in freely moving animals, rather than a permanent

optical fiber implant which is tethered to an external device. We first demonstrated that a cre-

dependent magnetogenetic system can target particular neuronal populations in cre-driver mice.

However, this approach is limited to animals and may also not provide the anatomical specificity

required to examine anatomical circuits. To address this, we also developed a modified approach

combining delivery of the cre-dependent magnetogenetic activating construct into the striatum

with injection of a retrograde viral vector (AAVretro) into globus pallidus, a known target of

striatal iSPNs, to restrict expression of the activating construct exclusively within the same

striatopallidal iSPNs targeted in the A2a-cre driver mice. Histological analysis confirmed that

expression was restricted to striatal D2 neurons and the behavioral and physiological responses

to the magnetic field were similarly robust compared with data generated in the A2a-cre mice.

However, AAVretro may not efficiently transduce all neurons in a retrograde fashion and must

be individually tested for particular applications, though there are alternative retrograde viral

vectors that could also be used in such an approach if the pattern of retrograde uptake differs

between vectors [36]. Nonetheless, our data provide a novel means for non-invasive regulation of

specific neuronal circuits without the need for transgenic animals, which could not only be useful

for pre-clinical research but even have potential for circuit-specific human translation.

One potential limitation of the application of magnetogenetics in a clinical setting is the

availability of a device for generating a magnetic field that is suitable for human therapeutic

applications. Stanley and colleagues [10] previously reported successful bidirectional

magnetogenetic modulation of ventromedial hypothalamus (VMH) neurons using the earlier dual

construct system and an MRI to deliver the magnetic field, and we similarly observed a robust

motor phenotype following exposure to a 3T MRI in mice expressing Nb-Ft-TRPV1^{Ca2+} in iSPNs at

similar field strength thresholds. While this is feasible for experimental studies, any requirement

of proximity to a 3T MRI for efficacy could limit the translational potential of this technology. We

addressed this by also testing a TMS device as a source of AMF to induce magnetic-field-based

stimulation of D2 iSPNs. As with the MRI machine, we observed a sustained effect from the TMS

generated magnetic field in mice expressing Nb-Ft-TRPV1^{Ca2+} in striatal D2 iSPNs, with decreased

locomotion and corresponding increase in GCaMP fluorescence. Interestingly, *in vivo* fiber

photometry analysis also revealed that TMS reduced neural activity in control mice, which could

potentially be attributed to TMS frequency-dependent neural activation-inhibition. This is not

surprising, as TMS does influence neuronal activity in isolation, which is the basis for current

clinical applications. Human [37] and animal [20] reports that have shown suppression of neural

activity in motor-related regions with different protocols of TMS treatment, especially low-

frequency TMS. Nonetheless, this reduced activity had no behavioral effect and if anything

suggests that we may even have underestimated the magnitude of magnetogenetic neuronal

activation. Given the widespread and increasing use of TMS as a clinical device for non-invasive

neuromodulation in a variety of conditions [19], it is possible that magnetogenetics could be used

to either augment TMS or allow TMS to target deeper neuronal populations where the strength

of the magnetic field weakens but is still above the threshold that we have observed for

functional activation of the recombinant engineered channel. The ability to generate low field

strength at-home devices for intermittent modulation similar to TMS, while maintaining

anatomical specificity through focal delivery of the gene therapeutic, could also allow broader

application of this technology. Finally, the ability to regulate peripheral or autonomic neurons is

plausible using this technology for indications where either activation or inhibition of neural

activity is desirable, potentially including the control of pain, neuropathy symptoms or organ

function. Our data establishing the threshold for magnetic field strength also raises the possibility

that other more simple or even wearable magnetic devices may enable regulation of more

accessible neuronal populations.

To expand the utility of this system, we have also developed a novel construct with a

mutation that converted the sodium channel into a chloride channel, allowing magnetogenetic

inhibition of neuronal activity. Our *in vitro* data demonstrated the functionality of this new

construct, while AAV-mediated delivery of the construct to a population of glutamatergic

projection neurons within the STN of hemiparkinsonian PitX2-cre mice confirmed the *in vivo*

efficacy of magnetogenetic inhibition and consequent motor improvements in this setting. Pitx2

expression is restricted to neurons within the general region of the STN [38,39], allowing very limited

expression of our cre-conditional magnetogenetic system to this population of STN neurons.

Altered activity of the STN is widely believed to be one of the pathophysiological mechanisms

leading to PD motor deficits [40], hence traditional high frequency DBS in the STN has emerged as

a standard therapy for advance stages of PD [2]. Nonetheless, there is equipoise regarding the

mechanism of DBS action and the role of inhibition of STN neurons as a PD therapeutic. Gradinaru

and colleagues [40] found that optical inhibition of glutamatergic STN neurons in hemi-parkinsonian

rats was insufficient to affect motor behavior using early generation inhibitory opsins, yet STN

lesioning, infusion of GABA agonists into the STN and gene therapy to promote GABA production

within the STN have all demonstrated improved motor function in both pre-clinical studies and

human clinical trials [23-25,27,41]. While DBS may still function through mechanisms other than simple

neuronal inhibition [42], our results support a significant body of evidence suggesting that inhibition

of STN projection neurons can improve motor function in Parkinson's disease [23,27,43-46]. The

availability of an inhibitory magnetogenetic system, which can also be combined with retrograde

cre vectors to target specific populations and can be applied to peripheral neurons as well, could

vastly expand both the experimental and clinical potential of this technology.

In summary, the present study demonstrates that magnetogenetic gene therapy

efficiently regulates basal ganglia circuitry. A single AAV encoding the activating magnetogenetic

construct restricted to iSPNs elicited freezing phenotype in transgenic mice. Circuit-specific

magnetogenetic activation of iSPNs was replicated in wt mice using AAVretro to deliver cre to a

specific neuronal population, while expression of a novel inhibitory magnetogenetic construct

within the STN reduced activity with consequent motor improvement in hemiparkinsonian mice.

Our results provide a bidirectional toolkit for magnetogenetic regulation to study specific central

or peripheral neural circuits, while further development of the technology may facilitate

translation into amenable human conditions where non-invasive and immediate modulation of

neuronal activity could have therapeutic benefit.

Materials and Methods

Generation of AAV Constructs

pAAV-JET-DIO-MCS/pLP362 is the backbone for the AAV vectors used to make **pAAV-JET-DIO-**

mCherry, pAAV-JET-DIO-Nb-FT-TRPV1^{Ca2+}, pAAV-JET-DIO-Nb-Ft-TRPV1^{Cl-} and pAAV-JET-DIO-

TRPV1^{Ca2+} and was made by replacing the BsrGI/NheI region of pAAV-hSyn-DIO-hM3D(Gq)-

mCherry (Addgene Cat No. 44361) with a multiple cloning site

(TGTACAACTAGTACCGGTTCGCGAGCATGCCCTAGGGCTAGC) and inserting a 242-bp PCR product

containing the JET promoter[47] amplified using primers 5'-

CTGCGGCCGCACGCGTGTACCATTGACGAATTCGGGCG and 5'-

ACTCTAGAGGATCCGGTACCTGTCAAGTGACGATCACAGGG into the MluI/KpnI sites.

To make **pAAV-JET-DIO-mCherry/**pLP388**,** the mCherry open reading frame was amplified using

primers 5'-ATTACCGGTGTTAACATGGTGAGCAAGGGCGAGGA

and 5'-TAAACCGGTCTTAAGTTACTTGTACAGCTCGTCCATG and cloned into AgeI-digested pAAV-

JET-DIO-MCS with the mCherry ORF in the reverse orientation to the JET promoter.

pAAV-JET-DIO-Nb-FT-TRPV1^{Ca2+}/pLP363 was made by ligating a 2943bp SpeI/EcoRV fragment

containing NbFT-TRPV1^{Ca2+} [14] to NruI/AvrII digested pAAV-JET-DIO-MCS.

The 865-bp HindIII/Tth111I fragment of pAAV-JET-DIO-Nb-FT-TRPV1^{Ca2+} was replaced with a 445-

bp HindIII-Tth111I fragment of synthesized DNA (Azenta Life Sciences, NJ) lacking the Nb-Ft

sequence to make **pAAV-JET-DIO-TRPV1^{Ca2+}**/pLP452.

To make **pAAV-JET-DIO-Nb-Ft-TRPV1^{Cl-}/**pLP375, primers 5'-

TCCATGGTGTTCTCCCTGGCAATGGGCTGGACCAACATGCTCT and

5'-AGACTAGTGTTATTTCTCCCCTGGGACCA were used to remove one NcoI site and amplify an

876-bp NcoI fragment of TRPV1^{mutant} encoding the I679K mutation[10] which was cloned into NcoI-

digested pAAV-JET-DIO-NbFT-TRPV1^{Ca2+}.

pAAV-CMV-Cre-GFP was obtained from Addgene Cat No. 68544.

Cell Culture and Adeno-Associated Virus (AAV) Preparation

Human embryonic kidney (HEK293, ATCC #CRL-1573) cells were cultured in DMEM (Gibco),

supplemented with 10% (vol/vol) FBS (Sigma-Aldrich) and 1% (vol/vol) penicillin-streptomycin

(Gibco), at 37 °C in 95% humidified air and 5% CO2. Vector stocks were prepared by packaging

the plasmids into rgAAV2 or mixed serotype AAV1/2 particles using a calcium phosphate

transfection system as described previously[48]. Cells were harvested and lysed at 72 h after

transfection. The vectors were purified using iodixanol gradient and dialyzed against PBS with

2mM MgCl2. AAV titers were determined by qPCR using Syber Green chemistry and with primers

to the WPRE fragment of the AAV backbone.

Quantitative Real Time PCR (qPCR)

To titer our purified viral vectors, AAV were processed as previously described[48] and viral

genomes quantified via qPCR using Fast SYBR green master mix (Applied Biosystems Cat. No

4385612) on the AB 7500 FAST Real Time PCR platform (Applied Biosystems) and primers to the

WPRE element: WPRE-Fw: 5'-GGCTGTTGGGCACTGACAAT-3'; WPRE-Rev: 5'-

CTTCTGCTACGTCCCTTCGG-3'. The relative number of full viral particles were calculated using the

standard curve method by normalizing to known standard samples.

Animals

Male mice were housed two to five per cage and kept at 22°C on a reverse 11 am-light/11 pm-

dark cycle, with standard mouse chow and water provided ad libitum throughout the duration of

the study. All animal procedures were approved by the Institutional Animal Care and Use

Committee of Weill Cornell Medicine and were in accordance with National Institutes of Health

guidelines.

The following mice were used in these studies: i) wt C57BL/6J mice, obtained from Jackson

laboratory (Jax.org; strain #:000664); ii) 129S-Pitx2tm4(cre)Jfm/Mmucd (Pitx2-Cre; MMRRC stock

#:000126-UCD) and *B6.FVB(Cg)-Tg (Adora2a-cre) KG139Gsat/Mmucd* (A2a-Cre; MMRRC stock

#:036158-UCD).

Stereotactic Surgery

All stereotactic surgical procedures were performed on 8–12-week-old mice, weighing 20g to

25g, under a mixture of ketamine/xylazine anesthesia. Ketamine (Butler Animal Health Supply)

and xylazine (Lloyd Laboratories) were administered at concentrations of 110 and 10 mg/kg body

weight i.p., respectively. After the induction of anesthesia, the animals were placed into a

stereotactic frame (David Kopf Instruments). All infusions were performed using a 10-µL

stereotactic syringe attached to a micro-infusion pump (World Precision Instruments) at a rate

of 0.1-0.4 µL/min. To prevent reflux, after each infusion, the injection needle was left in place for

5 min, withdrawn a short distance (0.3–0.5 mm), and then left in the new position for an

additional 2 min before removal. with a 33 G needle

To generate 6OHDA lesioned mice, animals were injected with a total volume of 0.6 µL of 6OHDA

hydrobromide (Sigma-Aldrich) in PBS with 0.1% ascorbate unilaterally into the medial forebrain

bundle (MFB) at a concentration of 2.5 mg/mL and an infusion rate of 0.1 µL/min with a 10-µL

Hamilton syringe with a 30 G needle. The coordinates for the injection were AP −1.1 mm, ML -

1.1 mm, DV −5.0 mm relative to bregma and the dural surface. Before lesion surgery, the

norepinephrine reuptake inhibitor desipramine (25 mg/kg, i.p.) was administered via i.p.

injection at least 30 min before 6OHDA infusion, to protect neostriatal and cerebellar

noradrenergic neurons from the toxin-induced damage. Mice were allowed 4-6-week recovery

before being subjected to the apomorphine-induced a behavioral test to estimate the extent of

dopamine depletion in the substantia nigra (see following section).

For experiments in Pitx2-cre mice, AAV1/2 vectors were injected into the subthalamic nucleus

(STN) unilateral to the lesioned side (AP -2.0 mm, ML -1.7 mm, DV −4.7 mm from bregma) of

6OHDA mice. For experiments in A2a-cre mice, AAV1/2 vectors were injected bilaterally into the

dorsal striatum (AP +0.5 mm, ML ±2.3 mm, DV −3.5 mm from bregma). For experiments in wt

mice, AAV1/2 vectors were injected bilaterally into the dorsal striatum (AP +0.5 mm, ML ±2.3

mm, DV −3.5 mm from bregma) and AAV2retro into the globus pallidus (AP +0.5 mm, ML ±2.3

mm, DV −3.5 mm from bregma). All craniotomies to inject AAV vectors were performed at a rate

of 0.4 µL /min using a 10-µL WPI syringe with a 33 G needle. A total of 2×10^{11} genomic particles/µL

(2 µL in PBS) of vectors were injected.

The mice were allowed a 6–8-week recovery before being subjected to the behavioral tests

during which maximal transgene expression from AAV vectors is achieved. At the conclusion of

the experiments, injection site accuracy within the targeted brain region was determined by

immunohistochemistry for the specific transgene expressed by the AAV vectors, and mice with

mistargeted injections were excluded from analysis before their data were unblinded.

Sources of Magnetic field

Direct magnetic field (DMF)

The magnetic field for *in vivo* studies was generated by the superconducting electromagnetic MRI

field from a Siemens 3.0 Tesla PRISMA MRI Scanner (Siemens Healthcare). The magnetic field

strength was measured and mapped using a gaussmeter (F.W. BELL MODEL 5080). High DMF was

defined as a region with a strength of 0.5-1.3 T, and Low DMF as 0.1-0.27 T.

Transcranial magnetic stimulation (TMS)

A butterfly C-B60 coil (165 x 85 x 19mm/inner diameter 35mm/outer diameter 75mm) was used

to deliver magnetic field during *in vivo* studies. Biphasic stimulation pulses were delivered by a

waveform generator (MagPro100x with MagOption) with the following settings: 20% machine

capacity (~600 mT), twin/dual mode 50 Hz stimulation, 500μs rise time, 1ms inter pulse interval,

10 pulses in train, number of trains 20, and inter train interval 1 s for 20 s.

Behavioral Assessments

All behavioral tests were run during the dark phase of the mouse daily cycle. The experiments

were performed by an examiner blinded to treatment groups.

Apomorphine induced rotations

To test for 6OHDA lesion efficiency in Pitx2-cre mice, rotational behaviors were performed. In

brief, mice were placed in a body harness connected to a transducer/swivel in a bowl-shaped

testing arena (rotometer). Full 360° clockwise and counter-clockwise rotations were measured

over a 45-min period after drug injection for apomorphine-induced rotations. Apomorphine

(Sigma-Aldrich) was dissolved in saline solution, 0.9% NaCl, and 0.1% ascorbate, and

administered at a dose of 0.25 mg/Kg via s.c. injection. Mice we randomized based on the net

number of contralateral rotations before receiving STN injections of the inhibitory

magnetogenetic construct.

To test the effect of our AAV.Nb-Ft-TRPV1^{Cl-} vectors on inhibiting STN neuronal activity, rotational

behaviors were performed inside a 3T MRI room in a sound insulated, ventilated Styrofoam

chamber that contained a plastic arena of 26.5 x 26.5 x 26.5 cm. Video recordings of apomorphine

induced rotations were obtained beginning 10 minutes after 0.25 mg/Kg via s.c. injections for 5

minutes outside the magnetic field followed by 5 minutes inside the magnetic field [1.3T -

500mT], and another 5 minutes outside the magnetic field in the start position using a magnetic

field compatible camera (MRC HiSpeed camera, https://www.mrc-systems.de/en/products/mr-

compatible-cameras#12m-camera). Upon the conclusion of behavioral studies, nigral lesion

efficiency by 6OHDA was determined by immunohistochemistry for the tyrosine hydroxylase (TH)

protein as a marker for dopaminergic neurons in the substantia nigra. Mice with mistargeted

injections were excluded from analysis before their data were unblinded.

Open-Field Arena

Locomotor behavior in an open-field arena was performed inside a 3T MRI room using the same

chambers and video recording system as described above for apomorphine induced rotations.

For 2 days prior to testing, each mouse was habituated in the open field box for 10 minutes to

reduce any possible confounding effect of a new environment. On the day of testing, each mouse

was placed in the center of the open field, and spontaneous horizontal and vertical behaviors

were recorded during 5-min sessions of pre-DMF, during DMF, and post-DMF.

Noldus EthoVision

EthoVision XT14 was used to analyze offline rotational behaviors and open field videos. The

automatic animal detection with deep learning settings was used to track center to-nose and

center to-tail points in defined arenas. Video recordings were used to analyze the following

locomotion and rotational parameters: (1) distance travelled, (2) activity percentage (%), (3)

rotation >90 degrees clockwise, and counterclockwise. Tracking data and heatmaps were

graphed with EthoVision analysis tools.

TRACKER

TRACKER video analysis and modeling tool (TRACKER 6 free software,

https://physlets.org/tracker/) was used to track precise motor responses, including freezing and

ambulation time. Each second of the recordings was divided into 5 frames to track nose, center,

fore and hind paws, and tail every 2 frames resulting in 150 positions measured per minute.

Spatial motion in the Y axis, referred as "change of location", was also analyzed as a function of

time.

Immunohistochemistry

On completion of all assessments, mice were deeply anesthetized with sodium pentobarbital

(150 mg/kg, i.p.) and transcardially perfused with 4% paraformaldehyde (PFA). Brains were

extracted and post-fixed overnight in 4% PFA, cryoprotected in 30% sucrose, and cut into 30-μm

sections using a vibratome (Leica Microsystems). Free-floating sections were treated with various

antibodies to visualize proteins of interest using immunofluorescence labeling, including tyrosine

hydroxylase (TH) to quantify the extent of nigral neurodegeneration, and mCherry or HA to

estimate the transduction efficiency from our AAV vectors. The following primary antibodies

were used: rabbit anti-RFP to detect mCherry (Rockland Cat. No 600-401-379; dil. 1:1000), sheep

anti-TH (Millipore Cat. No AB1542; dil. 1:1000), rabbit anti-HA (Sigma-Aldrich Cat. No H6908,

dil.1:500), and rat anti-cfos (Synaptic systems Cat. No 226-017, dil. 1:1000). Alexa Fluor-

conjugated fluorescent secondary antibodies (Molecular probes) were obtained from Life

Technologies and used at a 1:1000 dilution.

RNAScope in situ hybridization

Thirty μm free floating brain sections were mounted in PBS onto charged Superfrost slides (Fisher

Scientific) and were allowed to air dry for ~ 2h. The experiments were performed as per

manufacturer's instructions (ACDbio). The incubations at 40°C and 60°C took place in the HybEZ

oven using the EZ-Batch system (ACDbio). Sections were first baked at 60°C for 1h and post-fixed

in 4% PFA for 15 min at 4°C. After the dehydration steps, slides were dried at 60°C for 15 min,

treated with hydrogen peroxidase (Cat No. 322335, ACDbio) at room temperature for 10 min,

and dried again at 60°C for 15 min before performing the retrieval step. Sections were incubated

in target retrieval buffer (Cat No. 322000, ACDbio) for 10 min at 100°C and washed two times

with double deionized water (ddH20). After 3 min incubation in 100% Ethanol and drying at 60°C

for 10 min, sections were then incubated in Protease III (Cat No. 322337, ACDbio) at 40°C for

30min, washed five times with ddH20, and incubated for 2h at 40°C with probes including Mm-

Drd2 – C1, Mm-Drd1 – C3, mCherry-C4. A slide for either positive and negative control probes

were also included (Cat No. 321811 and 321831 respectively, ACDbio). Two washes occurred in

between the following 40°C incubations with wash buffer (Cat No. 310091). Sections were

incubated for 30min with AMP1, 30 min with AMP2, and 15min with AMP3 (kit Cat No.323110).

Next, sections were incubated in HRP C1 for 15min and Opal 520 (1:1500; Cat No. OP-001001) in

TSA buffer (Cat No. 322809). Sections were next incubated in an HRP blocker for 15 min.

Following sections were incubated with HRP C2 in combination with Opal 650 (1:1500; Cat No.

OP-001003), and HRP C4 in combination with Opal 570 (1:1500; Cat No. OP-001005). Slides were

cover slipped with Prolong Gold containing DAPI (ACDbio).

PET/CT animal imaging studies

All animal imaging experiments were performed in the Citigroup Biomedical Imaging Center

(CBIC, Weill Cornell Medicine). For baseline experiments, mice were fasted for 12 hours prior

imaging followed by. Intraperitoneal (i.p.) injection of ~500μCi ^{18}F-fluorodeoxyglucose (FDG), and

residual dose was measured. 1 hour after injection, mice were anesthetized with isoflurane and

scanned for 1 hour in a micro-PET/CT scanner (Siemens Inveon). For DMF experiments, mice were

exposed to a 3T MRI (DMF=500 mT − 1.3 T) for a total of 25 minutes after ^{18}F-FDG i.p.,

administration, and scanned following the same protocol as for baseline experiments. Following

imaging, scans were histogrammed and reconstructed using Siemens Inveon Acquisition

Workplace (Siemens Medical Solutions, Knoxville, TN, USA) and analyzed in DICOM format.

MRICron and MATLAB were used to do a PET-CT co-registration, and PMOD Fusion tool (version

3.6, PMOD Technologies Ltd., Zurich, Switzerland) was used for rigid transformations (manually

co-registered via rigid transformations [translations and rotations]), matching with MRI mouse

template, normalization with Mirrione mouse brain atlas, and regions of interest (ROI) analysis.

All analyses were conducted in a blinded fashion and were adjusted for mouse weight, ^{18}F-FDG

final dose (^{18}F-FDG$_{initial\ dose}$-^{18}F-FDG$_{residual\ dose}$), as well as relative to regions that do not express

dopamine 2-type receptors such as cerebellum. Right and left dorsal striatum were averaged for

each animal and MBq/ml or Percent injected dose per gram (%ID/g) were calculated to plot data.

In vivo fiber photometry

Fiber photometry was performed to measure *in vivo* calcium dependent activity dynamics during

DMF/TMS stimulation. Cre-dependent AAVs expressing GCaMP6s (AAV9-Syn-DIO-GCaMP6s,

Addgene Cat. No 100845-AAV9) and Nb-Ft-TRPV1^{Ca2+}-HA/mCherry were injected bilaterally into

the dorsal striatum of A2a-Cre transgenic mice to restrict expression of GCaMP6s and NbFt-

TRPV1^{Ca2+}-HA or mCherry to D2 iSPNs. Two weeks following AAV injection, a 400-µm-diameter

optical fiber (Doric, MFC_400/430-0.48_5.5mm_ZF2.5(G)_FLT, Cat. No B280-4418-5) was

implanted in the dorsal striatum 0.5 mm above the AAV injection site (AP: +2.00mm, ML: −0.25

mm, DV: −2.15mm) and was secured with Metabond. Prior to behavioral testing, mice were

habituated for 2 days to a patch cord attached to the implanted optical fiber for 3 min sessions.

During behavioral testing, fiber photometry recordings were taken for 2 min before, during, and

post magnetic stimulation. Behaviors were performed and recorded using the same settings as

described above for open-field arena locomotor quantification. Following behavioral testing,

fluorescence microscopy was utilized to confirm GCaMP6s expression and optical fiber

placement. To excite GCaMP6s, light from a 470nm LED (Thorlabs, M470F3) modulated at a

frequency of 521Hz was passed through a filter (Semrock, FF02-472/30), reflected by a dichroic

(Semrock, FF495-Di03) and coupled to a 0.48 NA, 400µm core optical fiber patch cord (Doric).

Emitted fluorescence traveled back through the patch cord, passed through the dichroic, a filter

(Semrock, FF01-535/50), and was focused onto a photodetector (Newport, Model 2151). The

modulated signal passed from the photodetector to a RP2.1 real-time processor (Tucker Davis

Technologies) where it was demodulated, and low pass filtered using a corner frequency of 15Hz.

TTL pulses denoting the start of behavioral trials were passed to the processor in real time for

alignment of calcium recordings to behavioral measures.

In vitro chloride imaging

Neuro 2A cells (ATCC ccl-131, mycoplasma testing and STR profiling performed by ATCC) were

grown in Eagles' minimum essential medium with 10% fetal bovine serum (Gibco) at 37°C and 5%

CO_2.

For immunocytochemistry studies, cells were cultured on 12-mm cover glass (Fisher Scientific,

Pittsburgh, PA) coated with fibronectin (Sigma). Cells were transfected with constructs

expressing Nb-Ft-TRPV1^{Cl-} or TRPV1^{Ca2+} 24 h after plating. Cells were stained 48-72hrs after

transfection.

For imaging studies using the chloride-dependent reporter MQAE, Neuro2A cells were grown on

12-mm cover glass and transfected with plasmids expressing mCherry alone, co-transfected with

NbFt-TRPV1^{Cl-} and mCherry or co-transfected with TRPV1^{Ca2+} and mCherry. Holotransferrin (2

mg/ml, Sigma) was added to cells 24 hours after transfection. Cells were placed at 32°C 24 hours

before imaging. Cells were washed 3 times with Krebs-HEPES buffer (NaCl 110 mM, KCl 5.5 mM,

$CaCl_2$ 2.5 mM, $MgCl_2$ 1.25 mM, HEPES 5mM and glucose 10mM) then incubated with MQAE 5mM

(Invitrogen) in Krebs-HEPES buffer for 1 hour at 37°C. Cells were then washed in Krebs-HEPES

buffer three times and placed in glass-bottom dishes with Krebs-HEPES buffer for 15 minutes

before imaging. Imaging was performed using an inverted fluorescent microscope Zeiss Axio

Observer Z1 microscope (Carl Zeiss, Oberkochen GER). Cells were imaged every 6 seconds for a

total of 3 mins with imaging before and during low AMF treatment with a neodymium magnet (K

and J magnetics) to produce a peak field 230-250mT. Image analysis was performed using FIJI.

Briefly, regions of interest (ROI) were selected based on mCherry expression and mean intensity

calculated for each ROI in each image after background subtraction. Chloride responses were

quantified as change in fluorescence intensity normalized to baseline fluorescence (ΔF/F0).

Imaging was performed on at least three occasions for each condition and chloride responses

were measured in 142 cells (Nb-Ft-TRPV1^{Cl-}), 76 cells (TRPV1^{Ca2+}) and 181 cells (mCherry).

Immunocytochemistry (ICC) was used to confirm cell surface expression of TRPV1. Live cells were

incubated with rabbit anti-TRPV1 (extracellular) polyclonal antibody (Thermo Fisher Scientific

cat#PA5-77361) (1:50) in culture medium for 10 min at 37 °C followed by 5 washes with RT culture

medium. Cells were then incubated with goat anti-rabbit Alexa Fluor 633 (1:1000) for 10 min at

32 °C followed by a further 5 washes with RT culture medium. Cells were then fixed in 4%

paraformaldehyde in Hank's balanced salt solution (HBSS) for 30 min at room temperature and

washed 5 times in HBSS before mounting using Fluoromount with DAPI (Southern Biotech,

Birmingham, AL).

Imaging acquisition and analysis

Images were taken by an epi-fluorescent microscope (Olympus BX61 microscope fluorescent

microscope fitted with an Olympus DP71 digital camera) or a confocal microscope (Zeiss LSM

900) and analyzed with Image J software (version 1.52p, NIH, US). For quantification of *c-fos*+

cells, coronal sections were sampled at intervals of 120-160μm for immunostaining. 3 dorsal

striatal or nigral sections per mouse were identified using the mouse brain atlas of Paxinos (2007)

and scanned bilaterally using a 20x objective. For each selected section, three randomly chosen

regions of interest (ROIs) with fixed areas were selected for quantification. A Macro Plugin was

applied to each section obtained, with a Gaussian blur filter threshold set to 30 and size filter to

20 to remove small objects and low signal cells. *c-fos*+ neurons were determined only when clear

co-localization with DAPI staining was observed. Automatized detection using Image J software

in 4-5 mice per condition (controls vs treated).

Statistics

To make valid comparisons among experimental groups, mice were randomized based on

baseline motor behavior to achieve balanced, unbiased groups. Pitx2-cre mice were randomized

based upon contralateral rotational counts following systemic apomorphine injection (0.25

mg/Kg), measured at 4 weeks after 6OHDA lesion surgery. The sample size for our experiments

was determined by power analysis[49], published work in the field, and previous studies performed

by the PIs. Exact sample sizes for each experiment are indicated in figure legends. The power was

set at 95% probability, effect size to 0.3-0.5 depending on the experimental outcome, and

significance at $p < 0.05$.

Statistical analyses were conducted using GraphPad Prism 7.0. The statistical test used in each

experiment depended on the type of data collected (reported in figure legends). We tested for

normality using the Kolmogorov-Smirnov test and Q-Q plots. Two-way ANOVA or two-tailed t test

for statistical comparisons were performed as appropriate.

For fiber photometry experiments, the change in fluorescent intensity (ΔF/F) percentage was

calculated by subtracting the linear fit of the isosbestic signal to the raw GCaMP signal from the

raw GCaMP signal, median for the entire recording session was then subtracted from each

GCaMP event value and divided by the median; that is, $(465_{raw} - 405_{fitted}) = 465_{net}$; $(465_{net} -$

$465_{median})/(465_{median})$ x 100. Events were computed comparing means in pre-DMF/DMF/post-

DMF or baseline/TMS conditions.

Pearson's correlation coefficients were used to quantify associations between different readouts

(e.g., c-fos+ immunoreactivity vs freezing of gait). When comparing two sets of normally

distributed data, unpaired two-tailed t-tests were utilized.

References

1.   Krauss JK, Lipsman N, Aziz T, et al. Technology of deep brain stimulation: current status
700   and future directions. *Nat Rev Neurol.* 2021;17(2):75-87.

701   2.   Benabid AL, Chabardes S, Mitrofanis J, Pollak P. Deep brain stimulation of the subthalamic
702   nucleus for the treatment of Parkinson's disease. *Lancet Neurol.* 2009;8(1):67-81.

703   3.   Boyden ES, Zhang F, Bamberg E, Nagel G, Deisseroth K. Millisecond-timescale, genetically
704   targeted optical control of neural activity. *Nat Neurosci.* 2005;8(9):1263-1268.

705   4.   Deisseroth K. Optogenetics. *Nat Methods.* 2011;8(1):26-29.

706   5.   Munshi R, Qadri SM, Zhang Q, Castellanos Rubio I, Del Pino P, Pralle A. Magnetothermal
707   genetic deep brain stimulation of motor behaviors in awake, freely moving mice. *Elife.*
708   2017;6.

709   6.   Keifer O, Kambara K, Lau A, Makinson S, Bertrand D. Chemogenetics a robust approach to
710   pharmacology and gene therapy. *Biochem Pharmacol.* 2020;175:113889.

711   7.   Wheeler MA, Smith CJ, Ottolini M, et al. Genetically targeted magnetic control of the
712   nervous system. *Nat Neurosci.* 2016;19(5):756-761.

713   8.   Stanley SA, Gagner JE, Damanpour S, Yoshida M, Dordick JS, Friedman JM. Radio-wave
714   heating of iron oxide nanoparticles can regulate plasma glucose in mice. *Science.*
715   2012;336(6081):604-608.

716   9.   Stanley SA, Sauer J, Kane RS, Dordick JS, Friedman JM. Remote regulation of glucose
717   homeostasis in mice using genetically encoded nanoparticles. *Nat Med.* 2015;21(1):92-
718   98.

719   10.   Stanley SA, Kelly L, Latcha KN, et al. Bidirectional electromagnetic control of the
720   hypothalamus regulates feeding and metabolism. *Nature.* 2016;531(7596):647-650.

721   11.   Chen R, Romero G, Christiansen MG, Mohr A, Anikeeva P. Wireless magnetothermal deep
722   brain stimulation. *Science.* 2015;347(6229):1477-1480.

723   12.   Long X, Ye J, Zhao D, Zhang SJ. Magnetogenetics: remote non-invasive magnetic activation
724   of neuronal activity with a magnetoreceptor. *Sci Bull (Beijing).* 2015;60:2107-2119.

725   13.   Stanley SA, Friedman JM. Electromagnetic Regulation of Cell Activity. *Cold Spring Harb
726   Perspect Med.* 2019;9(5).

727   14.   Stanley SA, Pomeranz LE, Li R, et al. Magnetogenetic cell activation using endogenous
728   ferritin. *bioRxiv.* 2023:2023.2006.2020.545120.

15.   Kravitz AV, Freeze BS, Parker PR, et al. Regulation of parkinsonian motor behaviours by
optogenetic control of basal ganglia circuitry. *Nature.* 2010;466(7306):622-626.

16.   Yu C, Cassar IR, Sambangi J, Grill WM. Frequency-Specific Optogenetic Deep Brain
Stimulation of Subthalamic Nucleus Improves Parkinsonian Motor Behaviors. *J Neurosci.*
2020;40(22):4323-4334.

17.   Alcacer C, Andreoli L, Sebastianutto I, Jakobsson J, Fieblinger T, Cenci MA. Chemogenetic
stimulation of striatal projection neurons modulates responses to Parkinson's disease
therapy. *J Clin Invest.* 2017;127(2):720-734.

18.   Berti V, Pupi A, Mosconi L. PET/CT in diagnosis of movement disorders. *Ann N Y Acad Sci.*
2011;1228:93-108.

19.   Cohen SL, Bikson M, Badran BW, George MS. A visual and narrative timeline of US FDA
milestones for Transcranial Magnetic Stimulation (TMS) devices. *Brain Stimul.*
2022;15(1):73-75.

20.   Muller PA, Dhamne SC, Vahabzadeh-Hagh AM, Pascual-Leone A, Jensen FE, Rotenberg A.
Suppression of motor cortical excitability in anesthetized rats by low frequency repetitive
transcranial magnetic stimulation. *PLoS One.* 2014;9(3):e91065.

21.   Harris JA, Hirokawa KE, Sorensen SA, et al. Anatomical characterization of Cre driver mice
for neural circuit mapping and manipulation. *Front Neural Circuits.* 2014;8:76.

22.   Vlasov K, Van Dort CJ, Solt K. Optogenetics and Chemogenetics. *Methods Enzymol.*
2018;603:181-196.

23.   Luo J, Kaplitt MG, Fitzsimons HL, et al. Subthalamic GAD gene therapy in a Parkinson's
disease rat model. *Science.* 2002;298(5592):425-429.

24.   LeWitt PA, Rezai AR, Leehey MA, et al. AAV2-GAD gene therapy for advanced Parkinson's
disease: a double-blind, sham-surgery controlled, randomised trial. *Lancet Neurol.*
2011;10(4):309-319.

25.   Kaplitt MG, Feigin A, Tang C, et al. Safety and tolerability of gene therapy with an adeno-
associated virus (AAV) borne GAD gene for Parkinson's disease: an open label, phase I
trial. *Lancet.* 2007;369(9579):2097-2105.

26.   Hasegawa T, Chiken S, Kobayashi K, Nambu A. Subthalamic nucleus stabilizes movements
by reducing neural spike variability in monkey basal ganglia. *Nat Commun.*
2022;13(1):2233.

27.   Niethammer M, Tang CC, LeWitt PA, et al. Long-term follow-up of a randomized AAV2-
GAD gene therapy trial for Parkinson's disease. *JCI Insight.* 2017;2(7):e90133.

28.   Guillaumin A, Pietro Serra G, Georges F, Wallén-Mackenzie Å. Optogenetic investigation
into the role of the subthalamic nucleus in motor control. *bioRxiv.* 2020.

29.   Kaplitt MG, Leone P, Samulski RJ, et al. Long-term gene expression and phenotypic
correction using adeno-associated virus vectors in the mammalian brain. *Nat Genet.*
1994;8(2):148-154.

30.   Deverman BE, Ravina BM, Bankiewicz KS, Paul SM, Sah DWY. Gene therapy for
neurological disorders: progress and prospects. *Nat Rev Drug Discov.* 2018;17(10):767.

31.   Shen Y, Campbell RE, Cote DC, Paquet ME. Challenges for Therapeutic Applications of
Opsin-Based Optogenetic Tools in Humans. *Front Neural Circuits.* 2020;14:41.

32.   Poth KM, Texakalidis P, Boulis NM. Chemogenetics: Beyond Lesions and Electrodes.
Neurosurgery. 2021;89(2):185-195.

33.   Jendryka M, Palchaudhuri M, Ursu D, et al. Pharmacokinetic and pharmacodynamic
774   actions of clozapine-N-oxide, clozapine, and compound 21 in DREADD-based
775   chemogenetics in mice. *Sci Rep.* 2019;9(1):4522.
776   34.   Hernandez-Morales M, Han V, Kramer RH, Liu C. Evaluating methods and protocols of
777   ferritin-based magnetogenetics. *iScience.* 2021;24(10):103094.
778   35.   Tye KM, Deisseroth K. Optogenetic investigation of neural circuits underlying brain
779   disease in animal models. *Nat Rev Neurosci.* 2012;13(4):251-266.
780   36.   Sarno E, Robison AJ. Emerging role of viral vectors for circuit-specific gene interrogation
781   and manipulation in rodent brain. *Pharmacol Biochem Behav.* 2018;174:2-8.
782   37.   Wang WC, Wing EA, Murphy DLK, et al. Excitatory TMS modulates memory
783   representations. *Cogn Neurosci.* 2018;9(3-4):151-166.
784   38.   Parolari L, Schneeberger M, Heintz N, Friedman JM. Functional analysis of distinct
785   populations of subthalamic nucleus neurons on Parkinson's disease and OCD-like
786   behaviors in mice. *Mol Psychiatry.* 2021;26(11):7029-7046.
787   39.   Wallen-Mackenzie A, Dumas S, Papathanou M, et al. Spatio-molecular domains identified
788   in the mouse subthalamic nucleus and neighboring glutamatergic and GABAergic brain
789   structures. *Commun Biol.* 2020;3(1):338.
790   40.   Gradinaru V, Mogri M, Thompson KR, Henderson JM, Deisseroth K. Optical deconstruction
791   of parkinsonian neural circuitry. *Science.* 2009;324(5925):354-359.
792   41.   During MJ, Kaplitt MG, Stern MB, Eidelberg D. Subthalamic GAD gene transfer in
793   Parkinson disease patients who are candidates for deep brain stimulation. *Hum Gene*
794   *Ther.* 2001;12(12):1589-1591.
795   42.   Ashkan K, Rogers P, Bergman H, Ughratdar I. Insights into the mechanisms of deep brain
796   stimulation. *Nat Rev Neurol.* 2017;13(9):548-554.
797   43.   Ricardo Y, Pavon N, Alvarez L, et al. Long-term effect of unilateral subthalamotomy for
798   Parkinson's disease. *J Neurol Neurosurg Psychiatry.* 2019;90(12):1380-1381.
799   44.   Martinez-Fernandez R, Manez-Miro JU, Rodriguez-Rojas R, et al. Randomized Trial of
800   Focused Ultrasound Subthalamotomy for Parkinson's Disease. *N Engl J Med.*
801   2020;383(26):2501-2513.
802   45.   Ji YW, Zhang X, Fan JP, et al. Differential remodeling of subthalamic projections to basal
803   ganglia output nuclei and locomotor deficits in 6-OHDA-induced hemiparkinsonian mice.
804   *Cell Rep.* 2023;42(3):112178.
805   46.   Milosevic L, Kalia SK, Hodaie M, Lozano A, Popovic MR, Hutchison W. Subthalamic
806   suppression defines therapeutic threshold of deep brain stimulation in Parkinson's
807   disease. *J Neurol Neurosurg Psychiatry.* 2019;90(10):1105-1108.
808   47.   Tornoe J, Kusk P, Johansen TE, Jensen PR. Generation of a synthetic mammalian promoter
809   library by modification of sequences spacing transcription factor binding sites. *Gene.*
810   2002;297(1-2):21-32.
811   48.   Morgenstern PF, Marongiu R, Musatov SA, Kaplitt MG. Adeno-associated viral gene
812   delivery in neurodegenerative disease. *Methods Mol Biol.* 2011;793:443-455.
813   49.   Charan J, Kantharia ND. How to calculate sample size in animal studies? *J Pharmacol*
814   *Pharmacother.* 2013;4(4):303-306.
815

Acknowledgments

This study was supported by the National Institute of Health R01 NS097184 (JMF, MGK) and the

JPB Foundation (JMF, MGK).

We acknowledge Henning U. Voss and Immanuel Elbau for assistance with the video-recording

systems during MRI and TMS, William Tower for assistance with the EthoVision software,

Neranjan de Silva for assistance breeding mice, Violetta Manusajyan for technical support during

MRI recordings, Leah Kelly for helpful discussions during experiments and Muc Chieu Du and

Kamaar Lawson for assistance with use of the Citigroup Biomedical Imaging Center.

Author contributions:

Conceptualization: JMF, MGK

Methodology: SRU, RM, JMF, MGK

Investigation: SRU, RM, LEP, AMA, SN, JPD, EKF, LG, RZ, SAS, MGK

Visualization: JMF, MGK.

Funding acquisition: MGK, JMF

Supervision: JMF, MGK

Writing – original draft: SRU, MGK

Writing – review & editing: SAS, JMF, MGK

Competing interest:

The authors declare no competing interests.

Materials & Correspondence:

Michael G. Kaplitt (mik2002@med.cornell.edu)

Figures

Fig. 1. Selective viral-mediated expression of Nb-Ft-TRVP1^{Ca2+} in striatal iSPNs elicit parkinsonian motor behavior. a. Schema of the double-floxed Cre-dependent AAV vector expressing the Nb-Ft-TRVP1^{Ca2+} or mCherry under the control of the JET promoter, and

immunostaining for HA in Adora-2A (A2a)-Cre mice demonstrates selective expression of the AAV- Nb-Ft-TRVP1^{Ca2+} in dorsal striatum. **b.** Selective transduction of D2-type iSPN with RNA in-situ hybridization (ISH). Co-localization of D2-type (green) iSPN with mCherry (red). **c.** Protocol and Schema for direct magnetic field (DMF) neural activation to assess motor behavior. **d.** Example of altered motor activity during bilateral striatal pre-DMF (green), DMF (red) and post-DMF (blue) stimulation; individual lines represent the path of a single mouse. Effect of DMF stimulation on **e.** Ambulation bout duration, and **f.** Freezing of gait in mCherry (blue bars/dots, n=10) and Nb-Ft-TRVP1^{Ca2+} (red bars/dots, n=10) mice. Error bars show SEM. ** represents p value < 0.001 with Student's t test.

Fig. 2. Nb-Ft-TRPV1^{Ca2+} in D2 iSPNs stimulated with high DMF and low DMF treatment produce freezing of gait. a. Schema for low direct magnetic field (DMF) titration. **b.** Magnetic field strength and individual activity heatmap example of motor activity during bilateral striatal Low DMF titration in Nb-Ft-TRPV1^{Ca2+} expressing mice. Freezing of gait in different magnetic

field gradients in the Low DMF titration ranges (blue bars, n=5) in Nb-Ft-TRPV1^{Ca2+} expressing mice. **c.** Protocol and schema for High DMF and Low DMF stimulation in A2a-Cre mice injected with the double-floxed Cre-dependent AAV vector expressing the Nb-Ft-TRPV1^{Ca2+} or mCherry under the control of the JET promoter. Representative tracking data of change of location in **d.** Nb-Ft-TRPV1^{Ca2+} and **e.** mCherry A2a-Cre mice during High DMF (red line), Low DMF (blue line), and No DMF (green line) treatment. Effect of High DMF and Low DMF stimulation on **f.** Ambulation and **g.** Freezing of gait in Nb-Ft-TRPV1^{Ca2+} (red dot/bars, n= 8) and mCherry (blue dots/bars, n=5). Error bars show SEM. * represents p value < 0.05, ** represents p value < 0.001 with Student's t test.

Fig. 3. Nb-Ft-TRPV1^{Ca2+} increases *c-fos* expression and cerebral metabolism upon DMF treatment. a. Immunostaining for *c-fos* (cyan) and HA or RFP (red) of dorsal striatum of A2a-Cre mice euthanized 1 hour following 15 minutes of DMF exposure. **b.** Percentage (%) of *c-fos* positive (+) neurons over total number of iSPNs expressing HA or RFP (n=5 mice per group/3

sections per mice). **c.** Correlation between *c-fos* expression and freezing of gait during DMF exposure (n=5 mice per group). **d.** Protocol and schema for 18-FDG micro-PET/CT scans procedure and analysis with representative micro-PET/CT imaging of A2a-Cre mouse brains expressing the Nb-Ft-TRPV1^{Ca2+} or the mCherry control virus following DMF exposure. 18-FDG uptake in dorsal striatum calculated as the percentage of injected dose/weight (%ID/g), **e.** DMF over baseline ratio. **f.** DMF over baseline normalized to cerebellum, in mCherry (blue bars/dots, n=8) and Nb-Ft-TRPV1^{Ca2+} (red bars/dots, n=8) mice. **g.** Correlation between DMF over baseline 18-FDG uptake ratio and freezing of gait during DMF exposure (n=10). Error bars show SEM. * represents p value < 0.05, ** represents p value < 0.001 with Student's t test. Anti-HA staining to detect Nb-Ft-TRVP1^{Ca2+} expression, anti-RFP staining to detect mCherry expression.

Fig. 4. Nb-Ft-TRPV1^{Ca2+} increases calcium transients in dorsal striatum. a. Schema of the double-floxed Cre-dependent AAV vector expressing the Nb-Ft-TRPV1^{Ca2+} or mCherry under the control of the JET promoter (red) combined with double floxed Cre-dependent AAV vector expressing GCaMP6s under the control of the Syn promoter (green). Behavioral methodology to

assess calcium transients during pre-DMF (green), DMF (red), and post-DMF (blue) treatment. **b.** Activity trace example of altered motor activity during bilateral striatal pre-DMF (green), DMF (red), and post-DMF (blue) stimulation in an Nb-Ft-TRPV1^{Ca2+} mouse with heatmap of GCaMP fluorescence aligned to pre-DMF (left), DMF (middle), and post-DMF (right) of the same animal shown in activity trace example expressing the Nb-Ft-TRPV1^{Ca2+} vector. **c.** Mean fluorescence aligned to pre-DMF, DMF, and post-DMF in Nb-Ft-TRPV1^{Ca2+} group (n=3). **d.** Activity trace example of altered motor activity during bilateral striatal pre-DMF (green), DMF (red), and post-DMF (blue) stimulation in an mCherry mouse with heatmap of GCaMP fluorescence aligned to pre-DMF (left), DMF (middle), and post-DMF (right) for the same animal shown in activity trace example expressing the mCherry vector. **e.** Mean fluorescence aligned to pre-DMF, DMF, and post-DMF in the mCherry group (n=4) **f.** Freezing of gait (FOG), **g.** Average delta (Δ) F/F (%) during pre-DMF, DMF, and post-DMF stimulation. **h.** Correlation between ΔF/F (%) calcium transients and FOG during DMF treatment. Error bars show SEM. * represents p value < 0.05 with Student's t test.

Fig. 5. Transcranial magnetic stimulation (TMS) treatment increases Nb-Ft-TRPV1^{Ca2+} mediated motor freezing and calcium transients in dorsal striatum. a. Schema of the TMS set up and A2a-Cre mice with bilateral injection of the double-floxed Cre-dependent AAV vector expressing the Nb-Ft-TRPV1^{Ca2+} or mCherry under the control of the JET promoter (red) combined with double floxed Cre-dependent AAV vector expressing GCaMP6s under the control of the Syn promoter (green) in dorsal striatum. **b.** Activity trace example of altered motor activity during baseline (left) and TMS (right) treatment in an mCherry mouse (blue), and Nb-Ft-TRPV1^{Ca2+} mouse (red). Distance ambulated during baseline (left) and TMS (right)

treatment in the **c.** mCherry group (blue), and **d.** Nb-Ft-TRPV1^{Ca2+} group (red). (n=4-5; data presented as the mean ± SEM in grey.). **e.** Average distance during baseline and TMS treatment in mCherry (blue) and Nb-Ft-TRPV1^{Ca2+} groups. **f.** Mean fluorescence during baseline in Nb-Ft-TRPV1^{Ca2+} (red) and mCherry (blue) mice (n=3; data presented as the mean ± SEM). **g.** Mean fluorescence during 20 seconds of TMS triggers every 2 seconds in Nb-Ft-TRPV1^{Ca2+} (red) and mCherry (blue) mice (n=3-4; data presented as the mean ± SEM). **h.** Grouped bar graph of Nb-Ft-TRPV1^{Ca2+} (red) and mCherry (blue) mice during baseline, and 20 seconds of TMS triggers every 2 seconds (n=3-4). Error bars show SEM. * represents p value < 0.05 with Student's t test.

Fig. 6. Selective viral-mediated expression of Nb-Ft-TRPV1^{Ca2+} in the striatopallidal pathway elicit parkinsonian motor behavior in wild type mice. a. Schema of the double-floxed Cre-

dependent AAV vector expressing the Nb-Ft-TRPV1^{Ca2+} or mCherry under the control of the JET promoter in striatal iSPN projecting to the Gp achieved by Gp injection of a retrograde AAV2 vector carrying Cre-GFP, and immunostaining for GFP (green), HA (red) in wt mice demonstrates selective expression of the AAV- Nb-Ft-TRPV1^{Ca2+} in dorsal striatum iSPN projecting to the Gpe, white arrows show colocalization of GFP with HA. **b.** Example of altered motor activity during bilateral striatopallidal pre-DMF (green), DMF (red) and post-DMF (blue) stimulation, lines represent the mouse's path. Effect of DMF stimulation on **c.** Ambulation bout duration, and **d.** Freezing of gait in mCherry (blue bars/dots, n=8) and Nb-Ft-TRPV1^{Ca2+} (red bars/dots, n=8) mice. **e.** Immunostaining for *c-fos* (cyan) and HA (Nb-Ft-TRVP1^{Ca2+}; red) or RFP (mCherry; red) of dorsal striatum of wt mice euthanized 1 hour following 15 minutes of DMF exposure. White arrows show co-localization. **f.** Percentage (%) of *c-fos* positive (+) neurons over total number of iSPNs expressing HA or RFP (n=3 mice per group/3 sections per mice). **g.** Schema of the double-floxed Cre-dependent AAV vector expressing the Nb-Ft-TRPV1^{Ca2+} or TRPV1^{Ca2+} under the control of the JET promoter in striatal iSPNs projecting to the Gp achieved by Gp injection of a retrograde AAV2 vector carrying Cre-GFP. Effect of DMF stimulation on **h.** Ambulation bout duration, and **i.** Freezing of gait in TRPV1^{Ca2+} (blue bars/dots, n=5) and Nb-Ft-TRPV1^{Ca2+} (red bars/dots, n=7) mice. Error bars show SEM. ** represents p value < 0.001 with Student's t test.

Fig. 7. Mutant Nb-Ft-TRPV1^{Cl-} inhibits neuronal activity of subthalamic nucleus projection neurons and rescues motor impairment in parkinsonian mice. a. Schema of inhibition system

with mutant Nb-Ft-TRPV1^{Cl-} membrane channel. **b.** Normalized change in MQAE fluorescence Intensity (ΔF/F0) of Neuro2A cells expressing mCherry, TRPV1^{Ca2+} or Nb-Ft-TRPV1^{Cl-} with low AMF application *in vitro*. **c.** AUC analysis of chloride imaging for Nb-Ft-TRPV1^{Cl-} channel. **d.** Protocol and schema for generation of parkinsonian PitX-2-Cre mice with unilateral injection of 6 hydroxydopamine (6-OHDA) in the medial forebrain bundle (MFB), followed by randomization based contralateral rotations induced by i.p. administration of apomorphine (0.25mg/kg) and ipsilateral intracranial injection of the double-floxed Cre-dependent AAV vector expressing the mutant Nb-Ft-TRPV1^{Cl-} under the control of the JET promoter in the subthalamic nucleus (STN). Representative immunostaining of the selective expression of Nb-Ft-TRPV1^{Cl-} (red) in STN neurons expressing Cre recombinase (Green). **e.** Percentage of change from pre-DMF treatment in contralateral rotations induced with apomorphine (0.25mg/kg, i.p.) in parkinsonian PitX-2-Cre mice expressing the Nb-Ft-TRPV1^{Cl-} or the mCherry control viruses in the STN during DMF, and post-DMF. **f.** Immunostaining for *c-fos* (cyan) and HA or mCherry (red) of STN of PiX-2-Cre mice euthanized 1 hour following 15 minutes of DMF exposure with apomorphine induced rotations (0.25mg/kg). White arrows show co-localization. **g.** Percentage (%) of *c-fos* positive (+) neurons over total number of iSPNs expressing HA or RFP (n=4 mice per group/3 sections per mice). Error bars show SEM. * represents p value < 0.05 with Student's t test. Anti-HA staining to detect Nb-Ft-TRVP1^{Ca2+} expression, anti-RFP staining to detect mCherry expression.