# Interchromosomal contacts between regulatory regions trigger stable transgenerational epigenetic inheritance in Drosophila

Maximilian H. Fitz-James[1], Gonzalo Sabarís[1], Peter Sarkies[2], Frédéric Bantignies[1], Giacomo Cavalli[1#]

[1]Institute of Human Genetics, CNRS and University of Montpellier, 141 Rue de la Cardonille, 34094 Montpellier, France

[2]Department of Biochemistry, University of Oxford, South Parks Road, Oxford OX1 3QU, UK

#Corresponding Author: giacomo.cavalli@igh.cnrs.fr

**Abstract**

Non-genetic information can be inherited through the germline and maintained across generations by the process of Transgenerational Epigenetic Inheritance (TEI). TEI can be established by various triggering events, including transient genetic perturbations. In *Drosophila*, hemizygosity of the *Fab-7* regulatory element leads to establishment of TEI at a homologous locus on another chromosome, resulting in heritable epigenetic differences in eye colour and in levels of the histone mark H3K27me3. Here, we investigate the mechanism by which hemizygosity of *Fab-7* leads to establishment of TEI. By mutating transcription factor binding sites within two key regions of *Fab-7*, we demonstrate the importance of two proteins in the establishment and maintenance of TEI: GAGA-factor and Pleiohomeotic. We show that these proteins function by recruiting the Polycomb Repressive Complex 2 and by mediating interchromosomal chromatin contacts between *Fab-7* and its homologous locus. Finally, we recapitulate these chromatin contacts using an *in vivo* synthetic biology system and show that chromatin contacts alone can establish TEI in the absence of hemizygosity. This work provides a mechanism by which hemizygosity of one locus can establish an epigenetic memory at another *in trans* through long-distance chromatin contacts.

## Introduction

Epigenetic information has long been known to be a major factor in the regulation of gene expression[1]. Whether such information can be transmitted across generations in various organisms has been a more elusive question, made difficult by the potential for genetic factors to confound experiments on heredity[2,3]. Through careful experimentation in model organisms, recent work has now demonstrated that such transgenerational epigenetic inheritance (TEI) does occur in a variety of organisms[4,5]. In addition, although much more work is required, the molecular mechanisms underlying these instances of inheritance have begun to be described[6]. Many of the most well-known epigenetic regulators of gene expression have been implicated in different cases of TEI, including non-coding RNAs[7–12], DNA methylation[13–15] and histone modifications[12,16–19], while less typical sources of non-genetic information, such as 3D chromatin organisation[19] and transcription factor binding[20], have been suggested as secondary signals in some cases.

Just as genes and their allelic variants are the basis of genetic variation, so are "epialleles" the basic units of heritable epigenetic change. Similarly, while mutation is the means by which genetic variation arises, "epimutation" describes the appearance of a heritable change in epigenetic information that gives rise to an epiallele. Epimutation provides an alternative source of heritable variation which differs from genetic mutation in that it has the potential to be more rapid, targeted and reversible, allowing for fast adaptation to a fluctuating environment[21]. Nonetheless, the underlying causes of epimutation and the mechanism by which they arise remain unclear and likely vary between organisms.

Given the complexity of mechanisms involved in TEI in higher eukaryotes, model systems that can be easily manipulated and allow one to both effectively track heritable phenotypes and analyse the underlying molecular events at play are critically needed in the field. One clear example of TEI occurs in *Drosophila melanogaster*, in a transgenic line called "Fab2L", involving a transgene that drives the expression of the eye pigmentation gene *mini-white*[22]. Flies of the Fab2L line exhibit a stochastic phenotype, manifesting as mosaicism of pigmentation in the adult eye. A memory of this phenotype can be established by a transient genetic perturbation[19] which can then be maintained epigenetically for countless generations following the initial trigger. The Fab2L line thus represents a valuable model system to investigate the underlying mechanism of epigenetic variability and the means by which it is triggered to become heritable. Here, we investigate the mechanistic basis for the establishment of transgenerational epigenetic inheritance at the Fab2L locus. We show that it is mediated by two key regulatory regions of the

transgene through the binding of the transcription factors GAGA-factor (GAF) and
Pleiohomeotic (Pho). We show that these transcription factors recruit the Polycomb Repressive
Complex 2 (PRC2) to the transgene, leading to deposition of H3K27me3, and promote
interchromosomal chromatin contacts between the transgene and a homologous region
elsewhere in the genome. Using an *in vivo* synthetic biology system, we show that chromatin
contacts alone are sufficient to induce TEI in *Drosophila*, providing a mechanism whereby
genetic perturbation of one locus can trigger TEI at another through long-distance chromatin
interactions.

**Results**

**Binding of GAGA-Factor and Pho is responsible for epigenetic variability at the Fab2L**
**transgene**

The *Drosophila* Fab2L line carries a single copy 12.4kb transgene inserted into chromosome
arm 2L at cytogenetic position 37B[22,23]. This transgene contains the reporter genes *LacZ* and
*mini-white* under the control of the *Fab-7* element. *Fab-7* is a well-studied regulatory region of
the bithorax complex on chromosome 3, where it regulates the expression of the Hox gene *Abd-*
*b*[24–26]. Importantly, the Fab2L line therefore contains two versions of the *Fab-7* element in its
genome, one at its endogenous location on chromosome 3 and one inserted ectopically within
the transgene on chromosome 2 (**Fig. 1A**).

The *mini-white* reporter gene, which controls red pigment deposition in the eye, is not expressed
uniformly in Fab2L flies but shows a mosaic pattern of eye pigmentation, with some ommatidia
showing strong *mini-white* expression and others strong repression, within the same individual
(**Supp. Fig. 1A,B**). Stochastic binding of the Polycomb Repressive Complex 2 (PRC2), resulting
in deposition of the repressive histone mark H3K27me3, has been suggested as an explanation
for the variability of mini-white expression in transgenes carrying Polycomb-bound elements.
This mosaic eye pattern therefore represents a very evident and visible instance of epigenetic
variation in the absence of any underlying genetic change[19].

To further investigate the mechanism behind this epigenetic variation, we generated transgenic
versions of the Fab2L transgene with mutations in key sequence motifs in the *Fab-7* element.
*Fab-7* contains within it two important subdomains: an insulator region, which in its endogenous
state prevents mis-regulation of *Abd-B* by adjacent regulatory regions in the wrong body

segments, and a Polycomb Response Element (PRE), which recruits either Polycomb Group

(PcG) or Trithorax Group (TrxG) proteins to maintain the pattern of *Abd-B* expression

established early in development. Both of these regions contain several consensus sequence

motifs for the DNA-binding proteins GAF and Pho (**Fig. 1B**). Among other functions, both of

these proteins are known to be recruiters of PRC2, strongly suggesting their potential

involvement in the variable H3K27me3 levels, and thus the eye colour phenotype, in Fab2L.

We generated three transgenic lines mutating all GAF and Pho sequence motifs within either

the insulator (Fab2L-INS), the PRE (Fab2L-PRE) or both (Fab2L-INS-PRE) in the Fab2L

transgene (**Fig. 1B**). Fab2L-INS showed a significant decrease in GAF binding to the insulator,

although Pho binding was little affected as was GAF binding to the PRE. Interestingly, Fab2L-

PRE had the opposite effect, with GAF binding at both sites unaffected, while Pho binding was

significantly decreased at the PRE only (**Fig. 1C,D**). These results extended to the recruitment

of PRC2 to Fab2L, which was largely unaffected in Fab2L-INS but significantly decreased in

Fab2L-PRE, not only at the PRE itself but also exterior to the *Fab-7* at the downstream LacZ

region (**Fig. 1E**). Mutation of both regions together shows clear additive effects. Indeed, in the

Fab2L-INS-PRE line binding of both GAF and Pho are decreased at both the insulator and PRE,

and in most cases to a significantly greater extent that in either of the single mutants (**Fig.**

**1C,D**), although the effects on PRC2 binding were not decreased to a significantly greater

extent than in Fab2L-PRE (**Fig. 1E**). Taken together, these results suggest that the insulator

and PRE regions of Fab-7 act cooperatively to recruit GAF and Pho to the Fab2L transgene,

although the PRE may play the greater role in the subsequent recruitment of PRC2.

We then analysed the downstream effects of this altered GAF and Pho recruitment on the

chromatin and phenotype of the mutant lines. All three mutant lines had significantly decreased

levels of H3K27me3 across the transgene, with the decrease being much more pronounced in

the double mutant Fab2L-INS-PRE line (**Fig. 1F**). These changes in chromatin translated to

phenotypic effects on the eye colour of the adult flies. Indeed, all three mutant lines displayed

shifts towards red eye colour compared to naïve Fab2L (**Fig. 1G-J, Supp. Fig. 1**). While it

remained significant, the shift in Fab2L-INS was milder than for the other two lines, with

approximately 16% of females and 61% of males exhibiting fully red eyes (**Fig. 1H, Supp. Fig.**

**1C,D**). In contrast, almost all Fab2L-PRE and Fab2L-INS-PRE flies of both sexes exhibited fully

red eyes. It is interesting to note, however, that while the shift towards red was complete for

Fab2L-INS-PRE, with all individuals having uniform red eyes (**Fig. 1J, Supp. Fig. 1G,H**),

Fab2L-PRE retained some stochasticity, with around 7% of females possessing at least some

white ommatidia (**Fig. 1I, Supp. Fig. 1E**). These results reinforce the idea that the insulator and

PRE together are responsible for the epigenetic and phenotypic variability of the Fab2L fly line.

**The insulator and PRE regions of *Fab-7* are individually sufficient to mediate epigenetic**

**inheritance**

In WT flies carrying the Fab2L transgene, epigenetic differences in expression between

individuals are not inherited transgenerationally. However, a transient genetic manipulation in

the form of a single generation of hemizygosity of the endogenous *Fab-7* results in

establishment of transgenerational epigenetic inheritance (TEI) which is subsequently

maintained[19]. Indeed, when we applied repeated selection and crossing of the most extreme

individuals in the population of an unmanipulated Fab2L line over ten generations, we did not

obtain any significant differences in eye colour across the population (**Supp. Fig. 2A,B**).

However, crossing this "naïve" Fab2L line with another Fab2L line bearing a homozygous

deletion of the endogenous *Fab-7* gives F1 individuals carrying the transgene in a homozygous

state, but the endogenous *Fab-7* in a hemizygous state (**Supp. Fig. 2C**). This hemizygosity

establishes transgenerational epigenetic memory at the transgene, such that reconstituting the

Fab2L genotype in the F2 results in a line which is genetically identical to the P0 Fab2L, but in

which TEI is now possible. Indeed, in this line selection over 10 generations results in either red

or white "epilines": populations of flies with a significant proportion of individuals with

monochrome eye colour, i.e. with 100% of their ommatidia either pigmented or unpigmented

(**Supp. Fig. 2D**). We also found that other crossing schemes which induced *Fab-7* hemizygosity

for one or two generations, were able to trigger TEI in Fab2L (**Supp. Fig. 2E-J**).

Given the altered phenotype of the mutant Fab2L transgenic lines, we asked whether these

mutations also interfered with the ability of the Fab2L transgene to maintain a memory of its

epigenetic state across generations by TEI. Due to the shift towards red eyes in the naïve

Fab2L-INS, Fab2L-PRE and Fab2L-INS-PRE lines, selection towards red eyes would not be

informative. We therefore performed a transgenerational epigenetic selection experiment to

determine if these mutant lines could be selected towards a more repressed, white-eyed

phenotype than the naïve population. Just as with wild-type Fab2L (**Supp. Fig. 2C**), we

introduced a single generation of *Fab-7* hemizygosity while leaving the mutant versions of the

Fab2L transgene unmanipulated (**Fig. 2A**). We then reconstituted the parental genotype,

homozygous for the endogenous *Fab-7*, and selectively bred the most white-eyed individuals

over subsequent generations. As a control, we used a wild-type Fab2L line which had previously been selected for a red-eyed phenotype. The Fab2L, Fab2L-INS and Fab2L-PRE, which all showed a greater or lesser degree of variability in their starting populations, were receptive to selection, showing a clear and gradual shift towards whiter eyes in both females and males over the generations (**Fig. 2 B-D, Supp. Fig. 3A-C**). In the case of Fab2L-INS, the appearance of some individuals with fully white eyes was even observed (**Fig. 2C, Supp. Fig. 3B**), consistent with the less extreme de-repression observed in this line compared to the other mutants. In contrast, the Fab2L-INS-PRE line displayed no variation at any point during the experiment, with all individuals of both sexes maintaining a uniform red eye colour (**Fig. 2D, Supp. Fig. 3C**). As expected, some form of epigenetic variation at the Fab2L transgene is therefore prerequisite for TEI. However, these results demonstrate that the insulator or PRE regions alone are still able to mediate TEI, indicating that they act together to maintain an epigenetic memory at the transgenic *Fab-7* element in the Fab2L line.

**The insulator and PRE regions of *Fab-7* are required for transmission of a repressed epigenetic state through paramutation**

The Fab2L transgene is not only able to acquire an altered epigenetic state by selection over generations, but can do so in a single generation by the process of "paramutation"[19]. Paramutation denotes the horizontal transfer of an epigenetic state *in trans* between two homologous alleles, and has been described in many organisms including *Drosophila*[27–30]. In the Fab2L line, crossing a naïve Fab2L with a Fab2L epiline (either white or red-eyed) results in the acquisition by the naïve allele of the epigenetic state of the epiline allele. This phenomenon can be tracked by the use of a *black[1]* marker allele, closely linked to the Fab2L transgene, such that F2 individuals that have inherited both copies of Fab2L from the naïve parent can be determined with high probability (**Fig. 3A, Supp. Fig. 4A**). Although these F2 flies possess the genetic material of the naïve P0 population, their epigenetic state resembles that of the epiline with which it was crossed, attesting to the acquisition over this genomic region of a new epigenetic state (**Fig. 3B, Supp. Fig. 4B**).

We performed paramutation crosses with the Fab2L-INS, Fab2L-PRE and Fab2L-INS-PRE lines to determine if these mutant versions of the Fab2L transgene could acquire a repressed epigenetic state by horizontal transmission in addition to, or instead of, selection over generations. Again, as control we used a wild-type Fab2L epiline which had previously acquired

a de-repressed, red-eyed epigenetic state. Just as with the selection, Fab2L, Fab2L-INS and

Fab2L-PRE were able to acquire a more white-eyed phenotype than the naïve parental lines,

with Fab2L-INS again being the most receptive to repression (**Fig. 3C-E**). Conversely, all

Fab2L-INS-PRE individuals maintained their uniform red coloration of the eyes even after

exposure in the F1 of the cross to a repressed epiallele (**Fig. 3F**). While lines bearing a wild-

type version of the insulator or PRE of *Fab-7* alone were therefore able to acquire an altered

epigenetic state by both selection over generations and paramutation, mutation of both regions

together completely prevents acquisition of a repressed epigenetic state by either method.

These results therefore suggest that both the insulator and PRE work together to mediate not

only epigenetic variation at the transgenic *Fab-7*, but are also key in maintaining an epigenetic

memory across generations.

**GAF mediates long-range chromatin contacts through the insulator region of *Fab-7***

Our results point to PRC2 and GAF as key factors mediating the epigenetic variability, and its

inheritance across generations, at the Fab2L transgene. PRC2 has a direct role in regulating the

expression of the Fab2L transgene, as the differences in *mini-white* expression and phenotype

of the Fab2L epilines correlate with differences in PRC2-deposited H3K27me3 across the

transgene[19]. The role of GAF is less clear as mutation of the GAF sites in the insulator ultimately

had little effect on PRC2 binding (**Fig. 1E**), suggesting that GAF's primary role at the *Fab-7*

element may not be the recruitment of PRC2. Intriguingly, among its many other functions, GAF

has been shown to mediate long-range chromatin contacts between its target genes[31–33]. The

three-dimensional organization of chromatin is a major factor in the regulation of gene

expression, and polycomb target genes in particular are frequently found to colocalize in the

nucleus at so-called "Polycomb bodies"[34]. Moreover, the transgenic and endogenous copies of

*Fab-7* were previously found to form chromatin contacts in the Fab2L line[19].

To investigate the effect of Fab2L mutation on these chromatin contacts, we performed FISH in

embryos of Fab2L and our mutated transgenic lines in order to visualise the regions surrounding

the *Fab-7* elements. These regions showed significant colocalization in the nuclei of Fab2L

embryos, but not in Fab2L ; *Fab7[1]* embryos which lack the endogenous *Fab-7*. This is

measured both in a decrease in average distance between the two loci (**Fig. 4A**) and in an

increase in the proportion of nuclei in which the loci are in close proximity (**Fig. 4B**). This shows

that chromatin contacts do occur between these loci dependent on the presence of both *Fab-7*

elements. Fab2L-INS-PRE mutants also showed no colocalization, similar to Fab2L ; *Fab7[1]*, indicating that the functionality of the *Fab-7* element is required for these contacts. In Fab2L-PRE mutants, the average distance between the two loci was not significantly different from wild-type Fab2L, suggesting that the insulator region alone can mediate contacts. Accordingly, contacts were significantly reduced in a Fab2L-INS mutant. It is worth noting, however, that inter-loci average distance in Fab2L-INS was not increased to the same extent as in Fab2L-INS-PRE when compared to Fab2L (**Fig. 4A**). This strongly suggests that contacts between the *Fab-7* elements are primarily mediated through the insulator region, and that the PRE may also play a stabilizing role.

**Artificially-induced chromatin contacts are sufficient to induce TEI at the Fab2L locus**

To build on these results, we sought to investigate the role of chromatin contacts in TEI at Fab2L. Inter-loci Fab2L-*Fab-7* distance does not differ significantly between white- or red-eyed epilines of Fab2L[19], arguing against a contribution of chromatin contacts to the epigenetic differences between these epilines. We also found that while these contacts are robust in late stage embryos (stage 14-15) they are not observable in early stages (stage 4-5) (**Supp. Fig. 5**), indicating that chromatin contacts themselves are unlikely to be the transgenerationally inherited signal of TEI. However, the Fab2L-*Fab-7* chromatin contacts observable in Fab2L increase even further in Fab2L ; *Fab7[1]*/+ individuals hemizygous for the endogenous *Fab-7* (**Supp. Fig. 5, Fig. 5D,E**). This increase in contacts therefore correlates with the triggering of TEI in the genetic crosses previously discussed (**Supp. Fig. 2**), making gain of chromatin contacts a prime candidate for the molecular trigger that establishes TEI. We therefore wished to explore this correlation in greater detail.

To directly investigate the role of chromatin contacts in the triggering of TEI, we therefore sought to induce contacts rather than abrogate them. To achieve this, we developed an *in vivo* system to induce interchromosomal contacts between the two regions of interest without recourse to any genetic perturbation. We dubbed this system "Three-Dimensional Contact Induction System" or "3D-CIS", for its ability to bring two distant loci in proximity (**Fig. 5A**). To create this system, we inserted arrays of Lac or Tet operators adjacent to the transgenic and endogenous *Fab-7* elements, respectively. Aside from the addition of these arrays, this line has the same genotype as the Fab2L line, and has a similar average distance and contact frequency between the two loci (**Fig. 5B,D,E**). However, activation of the system by introducing

a TetR-LacI fusion protein which binds to both arrays (**Supp. Fig. 6E,F**) resulted in anchoring of

the two *Fab-7* elements to each other (**Fig. 5C**). This anchoring leads to a decrease in the

average Fab2L-*Fab-7* distance and increase in the frequency of close contacts, both to levels

comparable to the hemizygous Fab2L ; *Fab7[1]* / + in which TEI is established (**Fig. 5D,E**).

Importantly, at no point are either the transgenic or endogenous *Fab-7* in a hemizygous state

(**Supp. Fig. 6A,B**). The 3D-CIS system therefore allows us to investigate the effect of increasing

chromatin contacts between the two *Fab-7* elements in the absence of any genetic perturbation.

Just as with Fab2L in the absence of genetic perturbation (**Supp. Fig. 2A,B**), selection of the

3D-CIS line in the "OFF" state over several generations did not result in any change in eye

colour across the population, towards either white or red eyes (**Fig. 5B,F**). After ten generations

of selection, these lines also exhibited no difference in H3K27me3 levels between each other

(**Fig. 5I**). However, activation of the 3D-CIS system, by introduction of the TetR-LacI fusion

protein and thus increase in contacts between the Fab2L transgene and endogenous *Fab-7*,

was able to establish TEI, such that selection over subsequent generations resulted in both

white and red epilines (**Fig. 5C, G**). These epilines also had significant differences in

H3K27me3 levels between them (**Fig. 5J**), reminiscent of the differences between Fab2L

epilines obtained by selection after transient hemizygosity of *Fab-7* (**Fig. 5H**). These results

demonstrate that chromatin contacts alone, in the absence of any genetic perturbation, are

sufficient to induce TEI at the Fab2L transgene. As further controls, we generated two more

lines expressing either a LacI-LacI or a TetR-TetR fusion protein as part of the 3D-CIS system

(**Supp. Fig. 6C,D**). These lines were also unable to trigger TEI (**Supp. Fig. 7**), showing that

triggering is not due to expression of a fusion protein or its binding to either array singly, but

conclusively results from the binding of the fusion protein to both arrays in tandem.

**The *Fab-7* element is required for stable chromatin contacts**

The ability of the 3D-CIS system to induce TEI in Fab2L suggests that the primary role of the

*Fab-7* element in the establishment of TEI is to mediate long-range chromatin contacts between

the transgenic and endogenous *Fab-7* elements. To determine whether induced chromatin

contacts can trigger TEI at the Fab2L transgene even in the absence of the *Fab-7* element, we

generated a new version of the 3D-CIS system in which the transgenic *Fab-7* was deleted

(**Supp. Fig. 8A**). As expected, phenotypically, this line resembled Fab2L-INS-PRE, with all

individuals possessing uniform red eyes (**Fig. 1J, Supp. Fig. 8B**). Similarly, this LacO-Fab2L-

Fab7Δ line was unable to acquire a repressed epigenetic state by either selection or paramutation (**Supp. Fig. 8C-E**). Activation of the 3D-CIS system by introduction of the LacI-TetR fusion protein was also unable to trigger TEI in this line (**Supp. Fig. 8F-I**). However, FISH analysis revealed that chromatin contacts between the transgene and the endogenous *Fab-7* were not increased in this line. Indeed in both the "OFF" and "ON" state, 3D-CIS-Fab7Δ flies did not show any significant contacts between the two loci, comparable to Fab2L ; *Fab7[1]* (**Supp. Fig. 8J**). This suggests that 3D-CIS is insufficient to mediate long-range chromatin contacts on its own, but rather acts to stabilise or reinforce contacts already established between the two *Fab-7* elements.

**Altered epigenetic states remain stable in the absence of artificially-induced chromatin contacts**

These results demonstrate a clear role for chromatin contacts in the initial triggering of TEI in Fab2L. However, due to experimental constraints, the 3D-CIS system remains active throughout the selection towards epilines. To determine whether the altered epigenetic states triggered by the 3D-CIS system can be maintained even in the absence of induced chromatin contacts, we crossed the 3D-CIS epilines with a naïve LacO-Fab2L. In the F2, flies lacking the TetR-LacI fusion protein (as determined by a GFP marker, see Methods) were selected and counted. Even in the absence of the TetR-LacI inducing chromatin contacts, this F2 generation had a majority of individuals with primarily white eyes, in the case of the white epiline (**Fig. 5K**), or primarily red eyes, in the case of the red epilines (**Supp. Fig. 9A**). The distribution of the population was comparable with those derived from a cross with Fab2L epilines triggered by transient hemizygosity rather than 3D-CIS (**Supp. Fig. 9B,C**). The LacO-Fab2L was therefore able to maintain the memory of its altered epigenetic state, even in the absence of artificially-induced chromatin contacts with the endogenous *Fab-7*, demonstrating that chromatin contacts are required to establish, but not maintain, transgenerational epigenetic inheritance.

## Discussion

**GAF-mediated chromatin contacts and PRC2-mediated epigenetic variability together account for TEI at the Fab2L transgene**

Our results highlight the crucial role played by two subdomains of the *Fab-7* element in the

establishment of epigenetic variation at the Fab2L transgene, and the maintenance of its

memory across generations. These subdomains are an insulator and a PRE, which act through

the recruitment of the transcription factors GAF and Pho. Alone, one of these regions remains

sufficient to maintain a certain degree of variation and epigenetic memory at the Fab2L

transgene, albeit in a manner skewed towards de-repression. Mutation of all GAF and Pho sites

across both regions however completely abrogates all variation, demonstrating that at least

some binding of these proteins is essential (**Fig. 1-2**).

Our findings suggest that the insulator and PRE cooperate to control epigenetic regulation of the

*Fab-7* element in two ways. The first is the recruitment of PRC2, which deposits H3K27me3 in a

stochastic manner, leading to the observed variable eye colour phenotype. The second is to

mediate long-range chromatin contacts between the two distant *Fab-7* elements in the genome.

However, mutation of these subdomains suggests that the PRE is the comparatively more

important of the two regions for PRC2 recruitment (and thus epigenetic variability) (**Fig. 1E**),

while the insulator is much more involved in the establishment of chromatin contacts (**Fig. 4**).

Nevertheless, both elements contribute to some extent to both aspects of Fab2L regulation.

The distribution of GAF and Pho bindings sites between these subdomains suggests differing

roles for these proteins, in agreement with what is known of their function. Indeed, the majority

of GAF binding sites (6 out of 9) are located in the insulator, while the majority of Pho sites (3

out of 4) are in the PRE (**Fig. 1B**).  Based on these observations, we propose that GAF is the

primary mediator of chromatin contacts, whereas Pho is the primary recruiter of PRC2, and thus

responsible for the epigenetic variability at the Fab2L transgene. Together, these proteins thus

mediate the dual functions of the *Fab-7* element, both of which are essential to TEI in this model

system.

**PRC2-dependent epigenetic memory at the Fab2L locus**

Our results clearly point to a central role for chromatin contacts in the establishment of TEI at

Fab2L, but not in the inheritance of the alternative gene expression at Fab2L, as contacts are

not present in early development (**Supp. Fig. 5**). PRC2-deposited H3K27me3 is thus the

primary epigenetic signal underpinning the variability at the Fab2L transgene and must be

inherited by another mechanism independent of chromatin contacts, either by direct inheritance

or by reconstruction in each generation based on an epigenetic memory maintained by some

other signal[6]. Alternatively, it is worth considering that a combination of direct and indirect

inheritance mechanisms could serve to reinforce each other, providing a more stable epigenetic

memory that either pathway alone. Previous work in *Drosophila* has provided evidence of

germline inheritance of H3K27me3, arguing for its ability to be transmitted through

gametogenesis at least[35]. Moreover, a recent study found that a DNA-binding protein, in this

case CTCF, can maintain its association with chromatin through development and serve as an

epigenetic signal itself[20]. That this could extend to a complex like PRC2, which both binds to

and deposits H3K27me3, is an interesting prospect, as inheritance of both these signals could

provide redundancy and reinforcement through a positive feedback loop. While our study is

primarily concerned with explaining how TEI is triggered at the Fab2L locus, future

investigations into the mechanism by which this epigenetic state is transmitted across

generations will provide a more complete picture of this instance of TEI.

**Chromatin contacts trigger PRC2-dependent TEI at the Fab2L locus**

Our results indicate that the primary mechanism for the triggering of TEI at the Fab2L transgene

is the promotion of physical contact within the nucleus between Fab2L and the endogenous

*Fab-7*. Increased contact frequency can either be induced by hemizygosity (**Supp. Fig. 2**) –

probably because the remaining copy of *Fab-7* forms contacts more efficiently with its distant

homolog at another locus in the absence of its homologous allele – or in a synthetic manner,

such as in our transgenic 3D-CIS system (**Fig. 5**). This demonstrates that chromatin contacts

can establish transgenerational epigenetic memory in the absence of any genetic perturbation.

Nevertheless, we note that this system is unable to mimic the effects of hemizygosity in the

absence of an adjacent *Fab-7* element (**Supp. Fig. 8**). This is explained by the fact that Fab2L-

*Fab-7* contacts are already observed to a lesser extent in Fab2L ; + individuals, but are

increased in Fab2L ; *Fab[7]*/+ hemizygotes (**Fig. 5D,E**). Thus, 3D-CIS acts to increase or

stabilise the contacts already occurring between the two loci, rather than driving the contacts *de*

*novo*.

As this trigger can lead to inheritance of epigenetic state in both directions (repression and de-

repression), any mechanism explaining how this trigger occurs must take into account this

plasticity. We therefore propose a model whereby stabilisation of Fab2L-*Fab-7* chromatin

contacts allows for the exchange of PRC2 between the endogenous and transgenic *Fab-7*

elements, thereby triggering an epigenetic memory that can be selected towards extremes over

generations (**Fig. 6**). In this model, naïve Fab2L flies have stochastic recruitment of PRC2 to the

transgene by Pho, leading to a random mosaic eye colour pattern (**Fig. 6A**). Chromatin contacts

are mediated by GAF, but in the absence of manipulation these contacts primarily occur

between homologous alleles, i.e. Fab2L to Fab2L or endogenous *Fab-7* to endogenous *Fab-7*

(**Fig. 6B**). While interchromosomal contacts between Fab2L and *Fab-7* do occur, they are

transient and outcompeted by the preferential interaction between homologous alleles that is

common in dipteran species[36] (**Fig. 6C**). Stabilisation of these contacts is achieved upon *Fab-7*

hemizygosity (**Fig. 6D**), because the remaining endogenous *Fab-7,* having lost its preferred

interaction partner, is free to form more stable contacts with its imperfect transgenic partner

without being outcompeted by its homologous allele. This situation can be mimicked in a

homozygous state, and in the absence of genetic perturbation, thanks to the 3D-CIS transgenic

system, which artificially stimulates chromatin contacts between Fab2L and *Fab-7*, making them

interact preferentially with each other rather than with their homologous alleles (**Fig. 6E**). When

these trans interactions are stable, exchange of PRC2 can occur between the PREs of the *Fab-*

*7* elements. Pho sites within the PRE can act as either donors or acceptors of PRC2, leading to

small changes in expression of the transgenic *mini-white* reporter. The memory of this

expression is maintained across generations, meaning that over time, these differences can be

selected to extremes, leading to either fully repressed or fully de-repressed transgene

expression, and monochrome eye colour (**Fig. 6F**).

A few aspects of this model are worth highlighting. First, it would predict that a Fab2L/+ ;

*Fab7[1]*/+ double hemizygote, in which both loci have lost their preferential interaction partner

on the homologue, should be an even more effective trigger of TEI than single hemizygosity.

While we have not examined this in detail, comparison of the speed of selection towards

epilines in our different TEI triggering crosses supports this (**Supp. Fig. 2**). Second, exchange

of PRC2 binding between Fab2L alleles, rather than between Fab2L and *Fab-7*, could also

explain how paramutation is able to transfer epigenetic state, albeit imperfectly, between

homologous alleles in this line. Our model thus accounts for both the triggering of epiallelic

identity, and its horizontal transfer by paramutation.

## A broader role for hemizygosity and chromatin contacts in triggering TEI

410  One major question in the field of epigenetic inheritance is how heritable epigenetic variability,

411  or epimutation, arises in the first place. Studies in plants suggest that heritable changes in DNA

methylation can occur apparently spontaneously in these organisms, leading to long-term epigenetic differences between lines[37–39]. Recent studies in *Caenorhabditis elegans* have extended these observations to metazoans and to RNA and chromatin-based epigenetic changes[40,41]. Other studies have sought to identify environmental triggers for TEI, directly linking epigenetic variation to an external stress to which it is intended to respond[2,42,43]. The final prominent candidate for sources of epigenetic variation is genetic perturbation, different types of which have been shown to trigger TEI in a variety of organisms[15,16,44,45].

In the *Drosophila* Fab2L line, this genetic perturbation takes the form of transient hemizygosity of the endogenous *Fab-7* region for at least one generation. It is interesting to note that unlike some cases of genetically-triggered TEI, in Fab2L the epigenetic memory is triggered and maintained not at the locus which is perturbed (the endogenous *Fab-7*) but elsewhere in the genome (the Fab2L transgene), testifying to the ability of trans-interactions to induce TEI. *Fab-7* has been found to have a similar effect on other loci. Indeed, hemizygosity of *Fab-7* has been shown to affect the expression of another PRE-containing gene in the distant Antennapedia cluster, with a phenotype that persisted for several generations after restoration of *Fab-7* homozygosity[19]. This raises the question of whether similar mechanisms could be acting to trigger TEI at other loci in natural populations.

Recent sequencing of wild *Drosophila melanogaster* lines has revealed the incredible genomic variation between populations of this single species. This includes numerous and large-scale deletions, duplications and translocations across the genome[46]. Mixing of two such genomically disparate populations would lead to a number of hemizygosity or heterozygosity events, as well as homology between very distant loci reminiscent of what is observed in Fab2L. Breeding in the wild thus has the potential to lead to many instances of naturally occurring genetic perturbation as potential triggers for TEI. Just as in Fab2L, it could be that establishment of TEI might be possible only between certain regions which are already prone to contact each other. In this respect, PRC2 targets may be particularly interesting candidates for naturally occurring TEI. Indeed, as previously mentioned, many PRC2 targets are frequently clustered within the nucleus in polycomb bodies, forming a large domain of silenced chromatin[47]. Interestingly, genes regulated in this manner are more likely to possess both an insulator region and a PRE, just like *Fab-7*. Our study provides insight into the mechanisms by which this type of epimutation could occur, but it is only by extending this insight to a broader context that we will be able to determine the role of TEI in the phenotypic variation, and thus potentially adaptation, of natural populations.

## Methods

### Fly stocks and culture

Flies were raised in standard cornmeal yeast extract media. Standard temperature was 21°C, with the exception of P0 and F1 crosses in the experiments of Fab2L epiallele establishment by hemizygosity or 3D-CIS (**Fig. 2, Fig. 5F,G, Sup. Fig. 2, Supp. Fig. 6A-D, Supp. Fig. 7D,E, Sup. Fig. 8D-I**), for which temperature was 18°C. The Fab2L and Fab2L ; *Fab7[1]* lines were described in Bantignies *et al.*, 2003[22]. The Fab2L, *black[1]* line and pre-established Fab2L epilines (Fab2L-R* and Fab2L-W*) were described in Ciabrelli *et al.*, 2017[19].

For generation of the Fab2L mutant lines, a transgenic line was made containing an AttP insertion site at cytogenetic position 37B, by CRISPR-Cas9 of a w[1118] line (Bloomington Drosophila Stock Center) to cut at the exact site of Fab2L transgene insertion in the Fab2L line. Fab2L-INS was then generated by Phi-recombination of an AttB-containing plasmid containing the entirety of the Fab2L transgene, with directed mutations of the insulator GAF and Pho sites, into this 37B-AttP line. Injection services for these two lines were provided by BestGene Inc.. Fab2L-PRE and Fab2L-INS-PRE were then generated by CRISPR-Cas9 editing of Fab2L-INS, with a two-guide RNA strategy designed and implemented by Rainbowgene Transgenic Flies Inc.

To create the 3D-CIS system, arrays of 7 Tet operators and 21 Lac operators were inserted adjacent to the endogenous or transgenic *Fab-7*, respectively, using a single guide RNA to cut immediately to their 3'. Cassettes encoding recombinant proteins combining the Lac and/or Tet repressors with a GFP marker (TetR-GFP-LacI, TetR-GFP-TetR and LacI-GFP-LacI) under expression of an *Actin-5C* promoter were inserted into chromosome arm 3L separately by Phi recombination into an established AttP containing line (Bloomington 24480). The transgenes encoding these proteins were then recombined with the TetO-*Fab-7*, and introduced into a Fab2L background, ready to be crossed with the LacO-Fab2L as described in **Supp. Fig. 6**. All injections for these lines were provided by BestGene Inc. The Fab2L-Fab-7Δ line was derived from the LacO-Fab2L line by CRISPR-Cas9 targeted deletion of the transgenic *Fab-7*, designed and implemented by Rainbowgene Transgenic Flies Inc.

474 Fab2L-INS, *black[1]*, Fab2L-PRE, *black[1]*, Fab2L-INS-PRE, *black[1]* and Fab2L-Fab7Δ, *black[1]*

475 were generated by recombining the Fab2L transgene with the *black[1]* allele from the w[1118];

476 *black[1]* line (Bloomington Drosophila Stock Center).

**Chromatin Immunoprecipitation and antibodies**

0 to 16 hour old embryos were collected in Embryo Wash Buffer (0.03% Triton X-100, 140mM

NaCl) and dechorionated with bleach. Samples were crosslinked in 1 ml A1 buffer (60 mM KCl,

15 mM NaCl, 15 mM HEPES [pH 7.6], 4 mM MgCl2, 0.5% Triton X-100, 0.5 mM dithiothreitol

(DTT), 10 mM sodium butyrate and complete EDTA-free protease inhibitor cocktail [Roche]), in

the presence of 1.8% formaldehyde. Samples were homogenized with a micropestle and

incubated for a total time of 15 minutes at room temperature. Crosslinking was stopped by

adding 350 mM glycine followed by incubation for 5 min. The homogenate was transferred to a 2

ml tube and centrifuged for 5 minutes, 4,000g at 4°C. The supernatant was discarded, and the

nuclear pellet was washed three times in 2 ml A1 buffer and once in 2 ml of Lysis buffer (140

mM NaCl, 15 mM HEPES [pH 7.6], 1 mM EDTA, 0.5mM EGTA, 1%Triton X-100, 0.5mMDTT,

0.1% sodium deoxycholate, 10 mM sodium butyrate and complete EDTA-free protease inhibitor

cocktail [Roche]) at 4°C. Nuclei were than resuspended in 1.5 ml Lysis buffer in the presence of

0.1% SDS and 0.5% N-Laurosylsarcosine, transferred to a 15 ml falcon tube and incubated for 2

hours with agitation at 4°C. Samples were adjusted to 3 ml and chromatin was sonicated using a

Q700 sonicator with microtip (QSonica) for a total of 6 minutes and 30 seconds at amplitude 50

(settings: 30 s on, 1min 30 s off x 13 cycles) in an ice bucket. Sheared chromatin had size range

of 100 to 300 base pairs. After sonication and 5 minutes high-speed centrifugation at 4°C,

fragmented chromatin was recovered in the supernatant and aliquoted in 5 µg (for H3K27me3

ChIP) or 20 µg (for non-histone protein ChIP) aliquots adjusted to a volume of 500 µl in Lysis

Buffer with 0.1% SDS and 0.5% N-Laurosylsarcosine for storage at -20°C.

To perform the ChIP, samples were thawed on ice and chromatin was precleared by addition of

15 µl of Protein A Dynabeads (Invitrogen 10002D) followed by incubation for at least 1 hour at

4°C. Dynabeads were removed on a magnetic rack and antibodies were added at a dilution of

1:100 (a mock control in the presence of rabbit IgG was performed at the same time, while an

input of the same size was set aside). Samples were incubated for overnight at 4°C on a rotating

wheel. 30 µl of Protein A Dynabeads were added and incubation was continued for at least 2

hours at 4°C. Antibody-protein complexes bound to beads were washed 4 times in Lysis Buffer

with 0.05% SDS and twice in TE Buffer (0.1 mM EDTA, 10 mM Tris (pH 8)) in 1 ml each time.

Chromatin was eluted from beads in 100 µl of 10 mM EDTA, 1% SDS, 50 mM Tris (pH 8) at

65°C for 15 minutes and eluted again in 150 µl of 10 mM EDTA, 0.67% SDS, 50 mM Tris (pH 8)

at 65°C for 15 minutes, with the eluate collected on a magnetic rack each time. The 250 µl

eluates and 250 µl of the Input DNA samples (1:2 input) were incubated overnight at 65°C to

reverse crosslinks and treated with Proteinase K for 3 hours at 56°C. DNA was isolated by

addition of an equal volume of phenol-chloroform, supernatants collected and then ethanol

precipitated for 2 hours at -20°C in the presence of 20 µg glycogen by addition of 25 µl 3M

sodium acetate and 625 µl ethanol. Samples were centrifuged at high speed for 1 hour and

washed in 500 µl of 70% ethanol before resuspension in 200 µl H2O. Immunoprecipitated DNA

was used to analyze the enrichment of specific DNA fragments by real-time PCR (qPCR), using

a Roche Light Cycler 480 and the Light Cycler 480 SYBR green I Master mix. For each

amplicon, IP DNA was normalized to Input DNA. The ChIP/Input ratio was further normalized to

a positive control region (engrailed). ChIP amplicons for the insulator or PRE regions were

specific to either the WT or mutated transgenic sequence, depending on the genotype analysed.

Antibodies used in this study were as follows: anti-GAF polyclonal antibody[48]; anti-Pho

polyclonal antibody[48,49]; anti-E(z) polyclonal antibody[31]; anti-H3K27me3 polyclonal antibody

(Active Motif 39155), anti-GFP polyclonal antibody (Abcam ab290), normal rabbit IgG (Cell

Signalling 2729).

**Fluorescence in situ hybridization**

Two-color 3D FISH was performed as previously described[50]. For a detailed protocol, see

Bantignies and Cavalli, 2014[51]. Briefly, embryos were dechorionated with bleach and fixed in

buffer A (60 mM KCl; 15 mM NaCl; 0.5 mM spermidine; 0.15 mM spermine; 2 mM EDTA; 0.5

mM EGTA; 15 mM PIPES, pH 7.4) with 4% paraformaldehyde for 25 min in the presence of

heptane. Embryos were then devitellinized by adding methanol to the heptane phase, extracted

and washed three times in methanol. Embryos were kept for a maximum of 4 months in

methanol at 4C before proceeding to FISH. Fixed embryos were sequentially re-hydrated in PBT

(PBS, 0.1% Tween 20) before being treated with 100–200 µg/ml RNaseA in PBT for 2 hours at

room temperature. Embryos were then sequentially transferred into a pre-Hybridization Mixture

(pHM: 50% formamide; 4XSSC; 100 mM NaH2PO4, pH 7.0; 0.1% Tween 20). Embryonic DNA

was denatured in pHM at 80°C for 15 minutes. The pHM was removed, and denatured probes

diluted in the FISH Hybridization Buffer (FHB: 10% dextransulfat; 50% deionized formamide;

2XSSC; 0.5 mg/ml Salmon Sperm DNA) were added to the tissues without prior cooling.

Hybridization was performed at 37°C overnight with gentle agitation. Post-hybridization washes
were performed, starting with 50% formamide, 2XSSC, 0.3% CHAPS and sequentially returning
to PBT. After an additional wash in PBS-Tr, DNA was counterstained with DAPI (at a final
concentration of 0.1 ng/µl) in PBT and embryos were mounted with ProLong Gold Antifade
(Invitrogen).

FISH probes for the 37B and 89E regions were made from a previous design described in
Ciabrelli et al. 2017[19]. For each region, 6 non-overlapping probes of between 1.2 and 1.7kb
covering an area of approximately 12kb were generated using the FISH Tag DNA kit with Alexa
Fluor 555 or Alexa Fluor 647 dyes (Invitrogen Life Technologies). 100ng of each probe were
added to the 30µL of FHB for hybridization.

**Microscopy and image analysis**

For the FISH, the 3D distances between 37B and 89E loci were acquired and measured as
follows: due to somatic pairing of homologous chromosomes in *Drosophila*, the majority of the
nuclei in embryos show a single FISH spot for each probe. In the cases of non-overlap FISH
signals between homologues, the closest distance between the centres of the two probes was
considered. To measure distances, 3D stacks were collected from 3-5 different embryos.
Optical sections were collected at 0.5 µm intervals along Z-axis using a Leica SP8-UV
microscope, Montpellier Resources Imaging (MRI) facility. Relative 3D distances between FISH
signals were analyzed in approximately 80 to 120 nuclei per 3D stack using the Imaris software
(Oxford Instruments). The distance distribution between the two probes was obtained by pooling
replicates for each condition.

**Acknowledgements**

We thank Montpellier Resources Imaging facility as well as the *Drosophila* facility (both affiliated
to BioCampus University of Montpellier, CNRS, INSERM, Montpellier, France). We thank Yuki
Ogiyama for assistance with the cloning of the plasmids for *Drosophila* injection and Judith
Kassis for the gift of the anti-Pho antibody. M.F-J. and G.S. were supported by the European
Research Council Advanced Grant 3DEpi with additional support for M. F.-J from the MSD
Avenir Foundation Grant GENE-IGH. F.B. and G.C. were supported by CNRS. Research in the
P.S. lab was supported by the John Fell Fund (Grant No 0011417). Research in the G.C.

laboratory was supported by grants from the European Research Council (Advanced Grant
3DEpi), the European CHROMDESIGN ITN project (Marie Skłodowska-Curie grant agreement
No 813327), the European E-RARE NEURO DISEASES grant "IMPACT", by the Agence
Nationale de la Recherche (PLASMADIFF3D, grant N. ANR-18-CE15-0010, LIUVCHROM,
grant N. ANR-21-CE45-0011), by the Fondation ARC (EpiMM3D), by the MSD Avenir
Foundation ((Project GENE-IGH), and by the French National Cancer Institute (INCa, PIT-MM
grant N. INCA-PLBIO18-362).

**Author contributions**

578 M. F-J. and G.C. conceived of and led the project. M. F-J. designed and performed the
579 experiments. M. F-J. and G.C. interpreted the data. M. F-J., G.S. and F.B. performed the
580 *Drosophila* transgenerational selection experiments. M. F-J. composed the manuscript with
581 editorial input from G.C. and P.S. All authors reviewed and commented on the manuscript.

**Competing interest declaration**

583 The authors declare no competing interests.

**References**

1. Allis, C. D. & Jenuwein, T. The molecular hallmarks of epigenetic control. *Nat. Rev.*
*Genet.* **17**, 487–500 (2016).

2. Cavalli, G. & Heard, E. Advances in epigenetics link genetics to the environment and
disease. *Nature* **571**, 489–499 (2019).

3. Bohacek, J. & Mansuy, I. M. A guide to designing germline-dependent epigenetic
inheritance experiments in mammals. *Nature Methods* **14** 243–249 (2017).

4. Liberman, N., Wang, S. Y. & Greer, E. L. Transgenerational Epigenetic Inheritance: From
Phenomena to Molecular Mechanisms. *Curr. Opin. Neurobiol.* **59** 189-206 (2019).

5. Bošković, A. & Rando, O. J. Transgenerational epigenetic inheritance. *Annu. Rev. Genet.*
**52**, 21–41 (2018).

6. Fitz-James, M. H. & Cavalli, G. Molecular mechanisms of transgenerational epigenetic
inheritance. *Nat. Rev. Genet.* **23**, 325–341 (2022).

7. Rechavi, O. *et al.* Starvation-induced transgenerational inheritance of small RNAs in C.

elegans. *Cell* **158**, 277–287 (2014).

8.    Schott, D., Yanai, I. & Hunter, C. P. Natural RNA interference directs a heritable response
to the environment. *Sci. Rep.* **4**, 7387 (2015).

9.    Teixeira, F. K. *et al.* PiRNA-mediated regulation of transposon alternative splicing in the
soma and germ line. *Nature* **552**, 268–272 (2017).

10.   Duempelmann, L., Skribbe, M. & Bühler, M. Small RNAs in the Transgenerational
Inheritance of Epigenetic Information. *Trends Genet.* **36**, 203–214 (2020).

11.   Cecere, G. Small RNAs in epigenetic inheritance: from mechanisms to trait transmission.
*FEBS Lett.* **595**, 2953–2977 (2021).

12.   Yu, R., Wang, X. & Moazed, D. Epigenetic inheritance mediated by coupling of RNAi and
histone H3K9 methylation. *Nature* **558**, 615–619 (2018).

13.   Furci, L. *et al.* Identification and characterisation of hypomethylated DNA loci controlling
quantitative resistance in Arabidopsis. *Elife* **8**, e40655 (2019).

14.   Silveira, A. B. *et al.* Extensive Natural Epigenetic Variation at a De Novo Originated
Gene. *PLoS Genet.* **9**, e1003437 (2013).

15.   Takahashi, Y. *et al.* Transgenerational inheritance of acquired epigenetic signatures at
CpG islands in mice. *Cell* **186**, 715–731 (2023).

16.   Greer, E. L. *et al.* Transgenerational epigenetic inheritance of longevity in Caenorhabditis
elegans. *Nature* **479**, 365–371 (2011).

17.   Torres-Garcia, S. *et al.* Epigenetic gene silencing by heterochromatin primes fungal
resistance. *Nature* **585**, 453–458 (2020).

18.   Seong, K. H., Li, D., Shimizu, H., Nakamura, R. & Ishii, S. Inheritance of stress-induced,
ATF-2-dependent epigenetic change. *Cell* **145**, 1049–1061 (2011).

19.   Ciabrelli, F. *et al.* Stable Polycomb-dependent transgenerational inheritance of chromatin
states in Drosophila. *Nat. Genet.* **49**, 876–886 (2017).

20.   Jung, Y. H. *et al.* Recruitment of CTCF to an *Fto* enhancer is responsible for
transgenerational inheritance of BPA-induced obesity. *Proc. Natl. Acad. Sci.* **119**, (2022).

21.   Sabarís, G., Fitz-James, M. H. & Cavalli, G. Epigenetic inheritance in adaptive evolution.

*Ann. N. Y. Acad. Sci.* **1524** 22-29 (2023).

22.     Bantignies, F., Grimaud, C., Lavrov, S., Gabut, M. & Cavalli, G. Inheritance of polycomb-
629             dependent chromosomal interactions in Drosophila. *Genes Dev.* **17**, 2406–2420 (2003).

630     23.     Zink, D. & Paro, R. Drosophila Polycomb-group regulated chromatin inhibits the
631             accessibility of a trans-activator to its target DNA. *EMBO J.* **14**, 5660–5671 (1995).

632     24.     Maeda, R. K. & Karch, F. The ABC of the BX-C: the bithorax complex explained.
633             *Development* **133**, 1413–1422 (2006).

634     25.     Kyrchanova, O. *et al.* Functional Dissection of the Blocking and Bypass Activities of the
635             Fab-8 Boundary in the Drosophila Bithorax Complex. *PLoS Genet.* **12**, 1–22 (2016).

636     26.     Kyrchanova, O. *et al.* Complete reconstitution of bypass and blocking functions in a
637             minimal artificial *Fab-7* insulator from *Drosophila bithorax* complex. *Proc. Natl. Acad. Sci.*
638             **116,** 13462-13467 (2019)

639     27.     Hollick, J. B. Paramutation and related phenomena in diverse species. *Nat. Rev. Genet.*
640             **18**, 5–23 (2016).

641     28.     Dorer, D. R. & Henikoff, S. Transgene repeat arrays interact with distant heterochromatin
642             and cause silencing in cis and trans. *Genetics* **147**, 1181–1190 (1997).

643     29.     De Vanssay, A. *et al.* Paramutation in Drosophila linked to emergence of a piRNA-
644             producing locus. *Nature* **490**, 112–115 (2012).

645     30.     Capovilla, M., Robichon, A. & Rassoulzadegan, M. A new paramutation-like example at
646             the Delta gene of Drosophila. *PLoS One* **12**, 1–16 (2017).

647     31.     Ogiyama, Y., Schuettengruber, B., Papadopoulos, G. L., Chang, J. M. & Cavalli, G.
648             Polycomb-Dependent Chromatin Looping Contributes to Gene Silencing during
649             Drosophila Development. *Mol. Cell* **71**, 73-88.e5 (2018).

650     32.     Mahmoudi, T., Katsani, K. R. & Verrijzer, C. P. GAGA can mediate enhancer function in
651             trans by linking two separate DNA molecules. *EMBO J.* **21**, 1775–1781 (2002).

652     33.     Chetverina, D., Erokhin, M. & Schedl, P. GAGA factor – a multifunctional pioneering
653             chromatin protein. *Cell Mol. Life Sci.* **78**, 4125–4141 (2021).

654     34.     Entrevan, M., Schuettengruber, B. & Cavalli, G. Regulation of Genome Architecture and
655             Function by Polycomb Proteins. *Trends Cell Biol.* **26**, 511–525 (2016).

35.  Zenk, F. *et al.* Germ line–inherited H3K27me3 restricts enhancer function during
maternal-to-zygotic transition. *Science* **357**, 212–216 (2017).

36.  McKee, B. D. Homologous pairing and chromosome dynamics in meiosis and mitosis.
*Biochim. Biophys. Acta - Gene Struct. Expr.* **1677**, 165–180 (2004).

37.  Johannes, F. & Schmitz, R. J. Spontaneous epimutations in plants. *New Phytol.* **221**,
1253–1259 (2019).

38.  Johannes, F. *et al.* Assessing the Impact of Transgenerational Epigenetic Variation on
Complex Traits. *PLoS Genet.* **5**, e1000530 (2009).

39.  Kawakatsu, T. *et al.* Epigenomic Diversity in a Global Collection of Arabidopsis thaliana
Accessions. *Cell* **166**, 492–505 (2016).

40.  Beltran, T., Shahrezaei, V., Katju, V. & Sarkies, P. Epimutations driven by small RNAs
arise frequently but most have limited duration in Caenorhabditis elegans. *Nat. Ecol.*
*Evol.* **4**, 1539–1548 (2020).

41.  Wilson, R., Le Bourgeois, M., Perez, M. & Sarkies, P. *Fluctuations in chromatin state at*
*regulatory loci occur spontaneously under relaxed selection and are associated with*
*epigenetically inherited variation in C. elegans gene expression. PLOS Genetics* **19**
(2023).

42.  Klosin, A., Casas, E., Hidalgo-Carcedo, C., Vavouri, T. & Lehner, B. Transgenerational
transmission of environmental information in C. elegans. *Science* **356**, 320–323 (2017).

43.  Casier, K., Boivin, A. & Teysset, L. Environmentally-Induced Transgenerational
Epigenetic Inheritance◻: Implication of PIWI Interacting RNAs. *Cells* **8** 1108 (2019).

44.  Reinders,  a., Burckert, N., Boller, T., Wiemken,  a. & De Virgilio, C. Saccharomyces
cerevisiae cAMP-dependent protein kinase controls entry into stationary phase through
the Rim15p protein kinase. *Genes Dev.* **12**, 2943–2955 (1998).

45.  Rassoulzadegan, M. *et al.* RNA-mediated non-mendelian inheritance of an epigenetic
change in the mouse. *Nature* **441**, 469–474 (2006).

46.  MacKay, T. F. C. *et al.* The Drosophila melanogaster Genetic Reference Panel. *Nature*
**482**, 173–178 (2012).

47.  Entrevan, M., Schuettengruber, B. & Cavalli, G. Regulation of Genome Architecture and

Function by Polycomb Proteins. *Trends Cell Biol.* **26**, 511–525 (2016).

48.    Schuettengruber, B. *et al.* Functional anatomy of polycomb and trithorax chromatin

landscapes in Drosophila embryos. *PLoS Biol.* **7**, (2009).

49.    Brown, J. L., Fritsch, C., Mueller, J. & Kassis, J. A. The Drosophila pho-like gene

encodes a YY1-related DNA binding protein that is redundant with pleiohomeotic in

homeotic gene silencing. *Development* **130**, 285–294 (2003).

50.    Bantignies, F. *et al.* Polycomb-dependent regulatory contacts between distant Hox loci in

Drosophila. *Cell* **144**, 214–26 (2011).

51.    Bantignies, F. & Cavalli, G. Topological Organization of Drosophila Hox Genes Using

DNA Fluorescent In Situ Hybridization. *Methods Mol. Biol.* **1196**, 103–120 (2014).

**Figure 1.** Mutation of GAF and Pho binding sites decreases epigenetic variability of the Fab2L transgene. (**A**) Schematic representation of the Fab2L transgene at its insertion site at cytological position 37B on chromosome 2, alongside the homologous *Fab-7* region on chromosome 3. (**B**) Illustration of the *Fab-7* element with important subdomains and transcription factor binding sites in wild-type and mutated versions of the Fab2L transgene. (**C-F**) ChIP-qPCR assays performed in embryos of the indicated genotypes at regions within the Fab2L transgene. Error bars represent +/- standard error from the mean (SEM) of three independent repeats. Samples were normalised to engrailed as a positive control and compared to wild-type Fab2L or between each other by the t-test (*p<0.05, **p<0.01, n.s. = not significant). (**G-J**) Phenotypic classification of eye colour in female and male adults of the indicated genotypes. Flies were sorted into five classes on the basis of eye colour, representing the number of pigmented ommatidia: Class 1 = 0% ; Class 2 = 1-10% ; Class 3 = 10-75%  ; Class 4 = 75-99% ; Class 5 = 100%.

**Figure 2.** Epigenetic inheritance of eye colour is abrogated in the absence of GAF and Pho binding to the Fab2L transgene. (**A**) Crossing scheme for the triggering of TEI at wild-type and mutant versions of Fab2L, with diagrammatic representation of the copy number of the *Fab-7* element on chromosomes 2 and 3. (**B-E**) Results of selection for the most white-eyed flies in each generation beyond the F2 of the crossing scheme. At top, curves represent the percentage of Class 5 females in the population across generations. Error bars are +/- standard deviation (SD) of 3 independent repeats. At bottom, pie charts represent the phenotypic distribution of the eye colour within the population in the first and last generations of selection.

**Figure 3.** The insulator and PRE regions are required for Fab2L to acquire a repressed epigenetic state through paramutation. (**A**) Illustration of the paramutation crossing scheme for the acquisition of a repressed epigenetic state by a naive Fab2L allele from an established epiallele *in trans*. The presence of a *black[1]* marker linked to Fab2L allows for the identification of F2 individuals carrying two copies of the transgene from the naive parent (grey chromosomes) rather than from the epiline parent (blue chromosomes). (**B-F**) Paramutation crossing schemes and phenotypic distribution of the populations with the indicated genotypes and epiline identities.

**Figure 4.** Long-distance chromatin contacts between Fab2L and *Fab-7* depend on the functionality of the insulator region. (**A**) Violin plots representing the distance distributions of the 37B and 89E regions surrounding the two *Fab-7* elements as determined by FISH in the indicated genotypes. Distances were measured in stage 14-15 embryos in T1 and T2 segments. Distributions were compared using the t-test (*p<0.05, **p<0.01, n.s. = not significant). (**B**) Bar graphs representing the percentage of cells from the same FISH assays in which the inter-loci distance was less than 1um.

**Figure 5.** An *in vivo* synthetic biology system promotes interchromosomal contacts between *Fab-7* elements and is sufficient to induce TEI without genetic perturbation. (**A-C**) Schematic representation of the 3D-CIS system. Arrays of Lac and Tet operons are inserted next to the transgenic and endogenous *Fab-7* elements. Expression of a TetR-LacI fusion protein binding to both arrays promotes contacts between the two loci. Illustrative micrographs represent nuclei from embryos with the 3D-CIS system in the "OFF" or "ON" state with FISH highlighting of the regions surrounding the *Fab-7* elements. (**D**) Violin plots representing the distance distributions of the 37B and 89E regions surrounding the two *Fab-7* elements as determined by FISH in the indicated genotypes. Distances were measured in stage 14-15 embryos in T1 and T2 segments. Distributions were compared using the t-test (*p<0.05, **p<0.01, n.s. = not significant). (**E**) Bar graphs representing the percentage of cells from the same FISH assays in which the inter-loci distance was less than 1um. (**F,G**) Results of selection for the most white or red-eyed flies in each generation of 3D-CIS flies in either the "OFF" or "ON" state. Curves represent the percentage of males of Class 1 or Class 4+5 in the population across generations. Error bars are +/- SD of 3 independent repeats. Pie charts represent the phenotypic distribution of the eye colour within the population in the first and last generations of selection. (**H-J**) ChIP-qPCR assays agasint H3K27me3 performed in embryos of the indicated genotypes after at least 10 generations of selection towards white or red epiallele identity, at regions within the Fab2L transgene. Error bars represent +/- SEM of three independent repeats. Samples were normalised to engrailed as a positive control and compared to each other by the t-test (*p<0.05, **p<0.01, n.s. = not significant). (**K**) Paramutation crossing scheme and phenotypic distribution of the populations with the indicated genotypes and epiline identities.

**Figure 6.** Model: Stabilisation of interchromosomal contacts triggers an epigenetic memory of PRC2 binding. (**A**) The *Fab-7* element recruits PRC2 by Pho binding to its PRE, leading to stochastic silencing of a mini-white transgene and a mosaic eye colour. (**B**) This PRC2 recruitment is coupled with long-range chromatin contacts with other *Fab-7* elements mediated by GAF through an insulator region. (**C**) When more than one copy of *Fab-7* is present in the genome contacts can be initiated between distant *Fab-7* elements, but these contacts are outcompeted by inter-allelic contacts and remain transient. Stabilisation of these contacts can be achieved through hemizygosity of one *Fab-7* copy (**D**) or through synthetic biology tools (**E**). This stabilisation leads to exchange of PRC2 between the PREs, resulting in either increased PRC2 association and silencing, or decreased PRC2 association and de-repression, and triggering an epigenetic memory of this altered association. (**F**) Over generations these slight differences can be selected to extremes, resulting in either very strong or very weak repression and strikingly different phenotypes.

**Supplemental Figure 1.** Images showing randomly selected samples of 10 females and males of the indicated genotypes.

**Supplemental Figure 2.** Crossing schemes and results of subsequent selection towards epilines of the Fab2L transgene. Selection was performed after either no cross (**A,B**), transient hemizygosity of the endogenous *Fab-7* element (**C,D**) or both the endogenous and transgeic *Fab-7* elements, achieved from crosses with different parental genotypes (**E-J**). In each case, Fab2L ; + flies derived from the indicated cross were split into two independent lines and subjected to repeated selection of the most extreme white or red-eyed flies, respectively, in each generation. Curves represent the percentage of Class 1 or Class 5 males in the population across generations. Pie charts represent the phenotypic distribution of the eye colour within the population in the first and last generations of selection.

**Supplemental Figure 3.** Results of selection for the most white-eyed flies in each generation beyond the F2 of the crossing scheme. At top, curves represent the percentage of Class 5 males in the population across generations. Error bars are +/- standard deviation (SD) of 3 independent repeats. At bottom, pie charts represent the phenotypic distribution of the eye colour within the population in the first and last generations of selection.

**Supplemental Figure 4.** Illustration, crossing scheme and phenotypic distribution of a paramutation cross for the acquisition of a repressed epigenetic state by a naive Fab2L allele from an established epiallele *in trans*. The presence of a *black[1]* marker linked to Fab2L allows for the identification of F2 individuals carrying two copies of the transgene from the naive parent (grey chromosomes) rather than from the epiline parent (blue chromosomes).

**Supplemental Figure 5.** Violin plots representing the distance distributions of the 37B and 89E regions surrounding the two *Fab-7* elements as determined by FISH in the indicated genotypes and embryo stages. Distances were measured. Distributions were compared using the t-test (*p<0.05, **p<0.01, n.s. = not significant).

**Supplemental Figure 6. (A-D)** Crossing schemes used to generate the 3D-CIS lines and associated controls. Chromosomes 2 and 3 for each genotype of the crosses are illustrated with the different versions of the *Fab-7* element highlighted. Red bars indicate a wild-type *Fab-7* at either the endogenous (chromosome 3) or transgenic Fab2L (chromosome 2) locus. Purple and orange bars indicate the array-containing LacO-Fab2L and TetO-Fab7, respectively. As illustrated, none of the crosses introduce hemizygosity of either *Fab-7* element at any point. **(E,F)** ChIP-qPCR assays in embryos of the indicated genotypes against LacI-TetR, TetR-TetR or LacI-LacI using a GFP tag. Approximate positions of the loci analysed by each primer pair within or adjacent to the *Fab-7* elements are indicated.

**Supplemental Figure 7.** (**A-C**) Schematic representation of the 3D-CIS Tet and 3D-CIS Lac systems, variations on the 3D-CIS used as controls. Arrays of Lac and Tet operons are inserted next to the transgenic and endogenous *Fab-7* elements. Expression of a TetR-TetR or LacI-LacI fusion protein binding to only one array controls for the potential induction of contacts between homologous alleles rather than interchromosomal loci.(**F,G**) Results of selection for the most white or red-eyed flies in each generation of 3D-CIS Tet and 3D-CIS Lac flies in the "ON" state. Curves represent the percentage of males of Class 1 or Class 4+5 in the population across generations. Error bars are +/- SD of 3 independent repeats. Pie charts represent the phenotypic distribution of the eye colour within the population in the first and last generations of selection. (**F,G**) ChIP-qPCR assays agasint H3K27me3 performed in embryos of the indicated genotypes after selection towards white or red epiallele identity, at regions within the Fab2L transgene. Error bars represent +/- SEM of three independent repeats. Samples were normalised to engrailed as a positive control and compared to each other by the t-test (*$p<0.05$, **$p<0.01$, n.s. = not significant).

**Supplemental Figure 8.** (**A,F,H**) Schematic representation of the 3D-CIS-Fab7 Δ line in which the *Fab-7* element was deleted from the Fab2L transgene of the 3D-CIS line (**B**) Phenotypic classification of eye colour of the 3D-CIS-Fab7Δ flies. (**C**) Paramutation crossing scheme and phenotypic distribution of the populations with the indicated genotypes and epiline identities. (**D**) Crossing scheme for the triggering of TEI in Fab2L-Fab7Δ, with diagrammatic representation of the copy number of the *Fab-7* element on chromosomes 2 and 3. (**E,G,I**) Results of selection for the most white-eyed flies in each generation after the cross or in the 3D-CIS system. Curves represent the percentage of Class 5 males in the population across generations. Error bars are +/- standard deviation (SD) of 3 independent repeats. (**J**) Violin plots representing the distance distributions of the 37B and 89E regions surrounding the two *Fab-7* elements as determined by FISH in the indicated genotypes. Distances were measured in stage 14-15 embryos in T1 and T2 segments. Distributions were compared using the t-test (n.s. = not significant, i.e. p>0.05).

**Supplemental Figure 9.** Paramutation crossing schemes and phenotypic distribution of the populations with the indicated genotypes and epiline identities.