ABSTRACT

Several methods of ecological network inference have been proposed, but their consistency

and applicability for use across ecologically relevant scales require further investigation.

Here, we infer ecological networks using two data sets (YFDP, FIA) describing distributional

and attribute information at local, regional, and continental scales for woody species across

North America. We accomplish this inference using four different methodologies

(COOCCUR, NETASSOC, HMSC, NDD-RIM), incorporating biological data along an

occurrence-performance spectrum while accounting (or not) for various confounding

parameters. We contrast 1-1 associations at each evaluated scale to quantify consistency

amongst inference approaches. We also assess consistency across scales within each

inference approach. Ultimately, we find that inferred networks are inconsistent across scales

and methodologies, particularly at continental scales. We highlight how such inconsistencies

between network inference methods may be linked to using occurrence or performance

information and incorporating or not confounding factors. Finally, we argue that identifying

the "best" inference method is non-trivial. Thus, to facilitate the choice of inference methods

for a given purpose, we suggest aligning specific research questions and the scale

applicability of the method when interpreting the inferred links, network topology, and

ecological processes governing network assembly.

INTRODUCTION

Climate change, habitat loss, and biological invasions are reorganising Earth's biosphere

(Blowes et al. 2019), altering species interactions (Estes et al. 2011) and, consequently,

ecosystem functioning (Woodward et al. 2010), resulting in a call for ecological research to

consider species interactions when predicting the emergence of novel communities and

changes in ecosystem behaviour and functioning (Gilman et al. 2010). This goal can be

achieved by assessing how ecological networks, which encode species identities as nodes and

pairwise species interactions as links (Morales-Castilla et al. 2015), would be altered due to

changes in the underlying composition of ecosystems and biological communities.

Traditionally, ecological network research efforts have focused on characterising species-

species networks at local scales (Bimler et al. 2022). Recent works have taken a broader

perspective by characterising networks at regional scales (Hackett et al. 2019) and integrating

multiple local networks into global meta-networks (Fricke et al. 2022). Various models to

quantify links between species in the absence of prior information on their likely interactions

have also been proposed. These models exist on a continuum defined by two axes: (1) the

used biological input data and (2) the capability to integrate additional data sources (Figure

1).

Biological inputs, which describe the first axis, can be classified along an occurrence-

performance spectrum. Here, some network inference methodologies focus on occurrence

(Veech 2013) or performance data either implicitly through abundance (Ulrich and Gotelli

2010) or explicitly through biomass or fitness measurements (Bimler et al. 2022). This

occurrence-performance axis arguably represents one of the most important philosophical

and methodological discriminating characteristics between network inference approaches.

The second axis encapsulates the capacity to account for different covariates such as

environmental gradients, phylogenetic relatedness, and functional traits of species (Morales-

Castilla et al. 2015). Approaches along this spectrum can be sorted based on covariates, like

environmental conditions, being either absent or present (again, either implicitly or explicitly)

when inferring the associations in a network. The importance of including this information

has been demonstrated through climatological conditions altering the persistence of

competing species (Usinowicz and Levine 2021) and the role of functional traits in governing

interactions between forest tree species (Kunstler et al. 2012). Still, most ecological network

inference approaches do not incorporate such information (Blanchet et al. 2020), most likely

due to data availability constraints, concerns of collinearity, or complexity of analyses.

Lastly, ecological networks are affected by the species richness of their corresponding

ecosystems (Poisot et al. 2012), a property determined by the assessment scale (e.g., larger

areas contain larger species pools, Conor & McCoy, 2013). The occurrence and magnitude of

a link between species may thus differ across geographic scales due to changes in species

pool. While such effects of species turnover have been studied for in-situ quantified networks

and demonstrated to alter 1-1 links, no such assessment exists for inferred ecological

networks.

Practitioners of ecological network inference are faced with many decisions when selecting

an appropriate approach and data sources for their respective study. The philosophical and

methodological distinctions between approaches raise an important question: *How*

comparable are different methods of ecological network inference? The necessary

comparisons of modern ecological network inference methods at a scale relevant to

quantifying the broader impacts of global environmental change are missing. Here, we focus

on the comparability of inferred networks rather than their accuracy for three reasons. First,

locally in-situ measured networks often fail to describe all existing links and hence cannot be

used to benchmark interference accuracy. Second, existing frameworks to generate "virtual"

networks fail to incorporate the two dimensions that differentiate between ecological network

inference approaches and variable scales. Last, we aim to showcase the need to align the

inference approach with the research questions and the scale applicability of the method.

Doing this will give necessary context to the interpretation of the inferred links, network

topology, and ecological processes governing network assembly.

In this study, we compare and quantify similarities between four methods of ecological

network inference. By doing so, we investigate how consistent inferred ecological networks

are when quantifying links (1) at the same scale using different inference approaches and (2)

across ecologically relevant scales using the same inference approach. We expect networks

corresponding to opposing sides of the occurrence-performance spectrum to differ notably in

their structure, with lower dissimilarities between approaches leveraging similar input types.

We also expect that the largest differences between approaches will be between those that

account for abiotic factors and those that do not. We postulate that such differences will be

amplified at regional and continental spatial scales where environmental gradients are the

main determinants of species distributional patterns (Stephan et al. 2021). Lastly, we expect

ecological networks inferred with the same approach, but varying species-pools, to differ

across geographic scales. Within this comparison across scales, we expect changes in species

pools to contribute to overall network dissimilarity but not sufficiently to explain total

dissimilarities of networks.

MATERIAL & METHODS

ECOLOGICAL NETWORK INFERENCE METHODOLOGY

We inferred horizontal (i.e., non-trophic) ecological networks using four methods that

encapsulate the variability in statistical principles, data requirements, and ecological

interpretation (Table 1). A verbose description of each inference method is provided in the

supplementary material. These methods have been used primarily to infer associations

between bird species (Griffith et al. 2016, Tikhonov et al. 2020) and plant species (Morueta-

Holme et al. 2016, Bimler et al. 2022) at scales ranging from local to continental applications.

While other inference methods have been established (e.g.: Damgaard et al., 2018; Lamonica

et al., 2021; Swain et al., 2021), we have selected this set of four methods as distinct

examples of the occurrence-performance axis, as well as their differences in considering

environmental conditions (Figure 1).

Data handling, network inferences, and analyses were carried out in R (R Core Team 2021).

The packages cooccur (Griffith et al. 2016), netassoc (Blonder and Morueta-Holme 2017),

and hmsc (Tikhonov et al. 2020) were used for the COOCCUR, NETASSOC, and HMSC

methods, respectively. NDD-RIM was implemented via custom R code. The workflow

diagrams of data handling and network inference are presented in Figures S1 and S2.

DATA SOURCES & DATA HANDLING

We inferred networks based on two datasets of North American woody plants: the Forest

Inventory Analysis Database (FIA, Burrill *et al.*, 2018) and the Yosemite Forest Dynamics

Plot (YFDP, Lutz *et al.*, 2012). These were aggregated to describe a pool of possibly

interacting species at three scales: local (0.256 km^2), regional (3,030.17 km^2), and macro

(1,508,451 km^2).

The local scale pool of interacting species was defined using the YFDP data, which is a 25.6

ha study area within which all trees ≥1 cm diameter at breast height (dbh) are measured

according to the protocols of the Smithsonian ForestGEO network (Davies et al. 2021). We

used a subset of the YFDP data set containing 34,589 records of dbh of 11 woody plant

species measured in a grid of 640, 20 m × 20 m quadrats which we subsequently treated as

individual plots. For consistency with the FIA, we considered only those stems ≥2.5 cm dbh

(Gray et al. 2012), retaining the records of 34,444 individuals and their dbh, as a

measurement of resource allocation to growth, for network inference. These data were further

aggregated to express species' abundances and occurrence in the 640 quadrants of the YFDP.

Species pools at the regional and macro scales were defined by aggregating FIA-plot data.

This dataset summarizes information on tree species' presence, abundance, and size/biomass

across the forested portion of the United States of America (Gray et al. 2012). To remove

possible biases in inferred association matrices introduced by rare species, we removed all

species present in less than 10% of all locations for our analyses. We used the FIA-contained

records of biomass per species per plot as proxies for species' performances. The regional

scale species pool was defined by limiting FIA-plot data to the region encompassing

Yosemite National Park as defined by the U.S. National Park Service (National Park Service

2009). This selection resulted in 291 observations (i.e., biomass and abundance of a species

at an FIA-plot) at 101 plots containing 13 species. The macro scale species pool was defined

by limiting FIA-plot data to the Temperate Conifer Forests Biome polygon (World Wildlife

Fund 2020). This masking resulted in 96,169 observations across 46,328 plots containing 15

species.

FUNCTIONAL TRAIT DATA

The YFDP dataset also includes species-specific measurements of multiple functional traits.

We selected specific leaf area, leaf carbon content, and leaf nitrogen content for our analyses.

These traits describe important trade-offs in plant investment in photosynthetic organs and

storage of resources (Wright et al. 2004). The FIA dataset does not record functional trait

measurements for its respective scales. While functional trait data is available for our target

species from functional trait repositories like BIEN (Enquist et al. 2009), functional leaf trait

expression is modulated by the environment (Wright et al. 2005), potentially leading to an

inaccurate representation of local realities at large scales (Violle et al. 2012). We therefor

chose to use functional trait means exclusively when implementing HMSC at local scale

(Table 1).

ENVIRONMENTAL DATA

To quantify plot-level environmental conditions experienced by species/individuals, we

matched air temperature, soil moisture in a depth of 0-7 cm, total precipitation, and potential

evaporation to our data using the KrigR package (Kusch and Davy 2022). We chose these

climate parameters due to their link to vegetation performance (Kusch et al. 2022). To reduce

biases introduced by climate regime changes, we aggregated environmental conditions to 10-

year climatologies leading up to the sampling date of each plot.

PHYLOGENETIC RELATEDNESS

We established phylogenetic relatedness by pruning a phylogenetic mega tree supplied with

the V.PhyloMaker package (Jin and Qian 2019). All species in our data not represented in

this backbone phylogeny were removed from our analyses which resulted in the above-

mentioned 11, 13, and 15 species at plot-, regional-, and macro-scale respectively.

SPECIES DISTRIBUTIONS

We obtained species range maps (for use with NETASSOC, Table 1) using the BIEN

package (Maitner et al. 2018) to inform null expectations of species co-occurrences at FIA

scales (i.e., regional and macro). At the YFDP scale, null models for NETASSOC were

established by resampling the observed species-by-site matrix.

NETWORK CONSISTENCY QUANTIFICATION

To quantify dissimilarities of inferred ecological networks at the level of individual links

(interactions/associations) and node sets (species composition), we calculated interaction

beta-diversity (a relative measure of how many species-species links and/or species identities

are shared between two networks) using the betalink (Poisot et al. 2012) R package. This

comparison evaluates whether link patterns or species turnover drive changes in the identity

and sign of links. For this analysis, we reduced link sets to statistically significant links (p-

value < 0.05 for COOCCUR and NETASSOC; 90% credible interval not overlapping 0 for

HMSC and NDD-RIM) in the inferred weighted ecological networks. To ensure

comparability across methods, we collapsed opposing links in NDD-RIM networks into mean

links when computing topological metrics and assigned statistical significance only when

both underlying directed links were identified as statistically significant. Additionally, we

changed link weights to -1 and 1 based on their original sign. The pairwise associations

common to all inferred networks owing to the shared species are highlighted with black

outlines in Figures 2 and 3.

RESULTS

To assess the effects of ecological network inference method choice and specification along

the occurrence-performance axis and with explicit or implicit adjustments for climate

gradients, we focus on plot scale YFDP data (Figure 2) because most data types are available

at this scale (Table 1). Furthermore, the trends observed at plot-scale are mirrored at regional

and macro-scales (Figure S3 & Figure S4).

Neither association strength nor sign of 1-1 links showed consistent patterns across methods

for the YFDP dataset (Figure 2). This pattern holds even for different specifications of the

HMSC approach (Figure 2D to F). Inference methods focused on performance data (NDD-

RIM and HMSC in Figure 2F) identify more statistically significant links between species

pairs than those focused only on occurrence or abundance data (Figure 2A, B, D, E).

As expected, the network matrices for the eight-species pairings shared across all scales

(black outlined tiles in Figure 3) show no scale consistency within the evaluated methods.

The magnitude and statistical significance of links change across scales for all methods. In

the case of HMSC (Figure 3C), we infer a heterogenous network matrix at the plot scale

(containing negative and positive statistically significant links) which changes to an

exclusively positive, statistically significant link matrix at the macro scale. This scale-

inconsistency can also be seen for COOCCUR (Figure 3A) and NETASSOC (Figure 3B),

where we fail to detect any statistically significant associations at the regional scale. Lastly,

using NDD-RIM, there is little consistency in the sign and magnitude of statistically

significantly identified directed links at the plot scale (Figure 1C), regional scale (Figure

S3C) and macro scale (Figure S4C). These inconsistencies remain when collapsing opposing

directed links inferred by NDD-RIM into undirected links (Figure 3D).

Quantifying network dissimilarity via decomposition of network-dissimilarities into node-set

and link-pattern changes reflects the overall lack of inference consistency across scales and

approaches. Network dissimilarities within scales (Figure 4, Table S2) are high and cannot be

traced back to species turnover (by definition, each network for the same scale consists of the

same node set). Instead, these must be attributed to each method input data or statistical

philosophy. In line with our expected differences due to the use of occurrence and

performance data, the lowest dissimilarity scores within-scales can be found between HMSC

and NDD-RIM outputs. As for abiotic factors governing ecological interactions and their

inference, we find comparatively low link dissimilarity of individual links between HMSC

and NETASSOC at the macro scale, for which we hypothesised climatological gradients to

play crucial roles. When investigating cross-scale consistencies, we find that network

dissimilarity is driven by changes in links between species present across the plot, regional,

and macro scale instead of added or removed link structures introduced by species turnover

(Table S3). This dissimilarity may be reduced by large amounts of data (Figure S5). This

finding is consistent for all inference methods assessed here. Finally, these inconsistencies

extend beyond comparisons of individual link structures and lead to a lack of consistency in

network topology as described by modularity and node centrality (see supplementary material

and Figures S6 & S7).

<u>DISCUSSION</u>

We have identified a lack of consensus among networks inferred using COOCCUR,

NETASSOC, HMSC, and NDD-RIM. This discrepancy could be attributed to (1) differences

in philosophy of methods (e.g., occurrence vs. performance input data, accounting for abiotic

conditions and niche preferences), (2) scale-dependant applicability of different methods and

data availability/richness, (3) realised biological interactions change across scales due to

differences in species pools, and (4) changes in the ecological processes driving community

assembly across scales.

CHOICE OF NETWORK INFERENCE APPROACH ACROSS SCALES

Choice of network inference method should be driven by method applicability given study's

spatial extent and data availability. Our results highlight two factors that require careful

consideration when choosing which method to implement: (1) biological inputs being

analysed and (2) data integration and availability.

First, the traditional reliance on occurrence data for network inference has been criticised as

being too simplistic to measure ecological process outcomes (Blanchet et al. 2020). As seen

in Figure 2 and Figure 3, methods of network inference that leverage abundance or

performance data identify more statistically significant associations than their occurrence-

driven counterparts. We postulate that difference in identified statistically significant

ecological associations is partially driven by the amount of biologically relevant information in abundance and performance data compared to occurrence datasets (i.e., quantification of species performances implicitly and explicitly vs. qualitative information). Indeed, performance-based networks reported the lowest dissimilarity scores (Figure 4). Such differences in information content have already resulted in null model analyses to discriminate community assembly rules being augmented with abundance data (Ulrich and Gotelli 2010). We suggest that network inference approaches also be subjected to this paradigm shift, thus leveraging performance or abundance data wherever available despite the considerable data collection efforts this entails, particularly at large geographic scales.

Second, macroecological research has long recognised the relevance of environmental conditions on species presence (Wilkinson et al. 2019), abundance (Johnson and Sinclair 2017), and performance (Coutts et al. 2016). Thus, the scale applicability of ecological network inference depends on whether or not climatological gradients ought to be considered in a given study. Of the four evaluated methods, COOCCUR and NDD-RIM do not account for climatological gradients, while NETASSOC does so implicitly through range expectations, and HMSC does so explicitly through direct use of climate data to quantify contributions of bioclimatic niches and environmental conditions to species performance. By accounting for environmental gradients, which alter species co-occurrence potential, NETASSOC and HMSC address recurring criticisms of ecological network inference (Blanchet et al. 2020). The effect of climatological data integration into ecological network inference manifests in lower network dissimilarity scores for networks inferred with NETASSOC and HMSC at macro scales.

In addition to including climatological patterns, there are calls to account for the shared evolutionary history of species pairs as well as their functional trait expressions (Ovaskainen et al. 2017, Blanchet et al. 2020). Meta-analyses of networks across geographic scales, regions, link types, and trophic levels have revealed that a combination of these information criteria is required to uncover mechanisms explaining network structure (Xing & Fayle,

2021). However, this information is unavailable at some geographic scales (Table 1). Our

results clearly show that data sparsity can also affect network inference at the regional scale

(Figure 3), where NDD-RIM was the only capable method. This capability is likely due to the

response-impact model component of the method, designed to infer rare/unobserved links.

While complex methods like HMSC are seemingly the only ones capable of inferring

ecological networks at macroecological scales with respect to integration of additional

confounding variables, this scale is also where selective data sparsity problem is the

strongest, and inference may still be limited as the sampling of relevant information is costly

at such scales.

Lastly, the methods of ecological network inference assessed here focus on representing

networks emerging from different ecological processes. While COOCCUR and NETASSOC

aim to identify species associations and their outcome on space-utilisation by association

partners, HMSC is primarily aimed at disentangling community assembly processes with

link-inference as a side-product of residuals. In contrast, NDD-RIM considers asymmetries in

species associations and aims to identify the effect of one association partner on the

performance of the other. In addition, the response-impact model component of NDD-RIM

may prove helpful in predicting novel interactions as communities reshuffle, thus giving a

valuable set of testable hypotheses for experimental macroecology (Alexander et al. 2016).

The ecological mechanism supporting a given methodology and the biological interpretation

of network-links must thus align with the research questions being investigated to ensure the

results' reliability and interpretation. This consideration should be prioritised over the other

factors mentioned above.

SCALE-CONSISTENCY & BIOLOGICAL PROCESSES

Biological processes revealed by ecological networks are conditional on the choice and scale

of the inference method. Figure 3 and Figure S5 illustrate that 1-1 associations are

inconsistent across geographic scales when inferred using the same approach. These scale-

inconsistencies are not necessarily a flaw in the methods of network inference but instead a

feature of (1) the inference method applicability at varying scales, (2) the scale-dependence

of species-interactions, and (3) which biological processes are being captured at geographic

scales.

Inferred species links can be considered data-driven hypotheses about realised biological

interactions (Ovaskainen et al. 2016), which have been demonstrated to change in strength

and sign when additional species are introduced/removed (Bartomeus et al. 2021). Our

analyses of network link beta diversity suggest that scale-inconsistencies of 1-1 associations

are partially driven by variation in realised interactions but cannot be explained by changes in

species pools alone. Therefore, while there is a precedent for using ecological networks

measured/inferred at one scale to draw conclusion at other scales (Guimarães 2020), our

findings indicate that neither 1-1 associations nor whole-network topology measures should

be transferred between scales. We recommend ecological networks be inferred and

interpreted at the scales at which they are to be used for further analyses.

CONCLUSION

Methods of network inference provide a tool for ecological studies. They allow identifying

biological interactions which are extremely difficult to observe in nature (e.g., large

communities with many potential and realised interactions). We demonstrate however, that

scale inconsistencies are considerable for pairwise species links. We show how focusing on

the "best" or "most" consistent ecological network inference is a disingenuous goal. Instead,

the choice of method needs to consider: (1) alignment of link-interpretation with research

questions, (2) applicability of the network inference method at the geographic scale of study,

(3) data availability and information content at the geographic scale of study, and (4)

biological mechanisms which are identifiable using network inference at the scale of study.

ACKNOWLEDGEMENTS

This research was supported by the Aarhus University Research Foundation Strat-up Grant

(grant no. AUFF-2018-7-8) to AO. AO considers this a contribution to the Center for

Ecological Dynamics in a Novel Biosphere (ECONOVO), funded by Danish National

Research Foundation (grant DNRF173).

AUTHOR CONTRIBUTIONS

EK and AO conceptualised this study. EK led the analysis. EK created all R scripts necessary

for the analyses. NDD-RIM integration was aided by MB. Results were interpreted by EK

and AO. YFDP data was provided by JAL. Manuscript writing was led by EK. All authors

contributed critically to the drafts and gave final approval for publication.

REFERENCES

Alexander, J. M. et al. 2016. When Climate Reshuffles Competitors: A Call for Experimental
Macroecology. Trends in Ecology and Evolution 31:831–841.
Bartomeus, I. et al. 2021. Experimental evidence of the importance of multitrophic structure
for species persistence. Proceedings of the National Academy of Sciences
118:e2023872118.
Bimler, M. D. et al. 2022. Estimating interaction matrices from performance data for diverse
systems.
Blanchet, F. G. et al. 2020. Co□occurrence is not evidence of ecological interactions.
Ecology Letters:ele.13525.
Blonder, B., and N. Morueta-Holme. 2017. netassoc: Inference of Species Associations from
Co-Occurrence Data.
Blowes, S. A. et al. 2019. The geography of biodiversity change in marine and terrestrial
assemblages. Science 366:339–345.
Burrill, E. A. et al. 2018. The Forest Inventory and Analysis Database: database description
and user guide version 8.0 for Phase 2. Page U.S. Department of Agriculture, Forest
Service.
Conor, E. F., and E. D. McCoy. 2013. Species-Area Relationships. Pages 640–650
Encyclopedia of Biodiversity: Second Edition. Academic Press.
Coutts, S. R. et al. 2016. Extrapolating demography with climate, proximity and phylogeny:
approach with caution. Ecology Letters 19:1429–1438.
Damgaard, C. et al. 2018. Insights on plant interaction between dominating species from
patterns of plant association: expected covariance of pin-point cover measurements of
two species. Environmental and Ecological Statistics 25:221–235.
Davies, S. J. et al. 2021. ForestGEO: Understanding forest diversity and dynamics through a
global observatory network. Biological Conservation 253:108907.
Enquist, B. J. et al. 2009. The Botanical Information and Ecology Network (BIEN):
Cyberinfrastructure for an integrated botanical information network to investigate the
ecological impacts of global climate change on plant biodiversity. Page The iPlant
Collaborative.
Estes, J. A. et al. 2011. Trophic Downgrading of Planet Earth. Science 333:301–306.
Fricke, E. C. et al. 2022. The effects of defaunation on plants' capacity to track climate
change. Science 375:210–214.
Gilman, S. E. et al. 2010. A framework for community interactions under climate change.

Trends in Ecology and Evolution 25:325–331.

Gray, A. et al. 2012. Forest Inventory and Analysis Database of the United States of America (FIA). Biodiversity and Ecology 4:225–231.

Griffith, D. M. et al. 2016. cooccur: Probabilistic species co-occurrence analysis in R. Journal of Statistical Software 69:1–17.

Guimarães, P. R. 2020. The Structure of Ecological Networks Across Levels of Organization. Annual Review of Ecology, Evolution, and Systematics 51:433–460.

Hackett, T. D. et al. 2019. Reshaping our understanding of species' roles in landscape-scale networks. Ecology Letters 22:1367–1377.

Jin, Y., and H. Qian. 2019. V.PhyloMaker: an R package that can generate very large phylogenies for vascular plants. Ecography 42:1353–1359.

Johnson, D. S., and E. H. Sinclair. 2017. Modeling joint abundance of multiple species using Dirichlet process mixtures. Environmetrics 28.

Kunstler, G. et al. 2012. Competitive interactions between forest trees are driven by species' trait hierarchy, not phylogenetic or functional similarity: Implications for forest community assembly. Ecology Letters 15:831–840.

Kusch, E. et al. 2022. Vegetation☐memory effects and their association with vegetation resilience in global drylands. Journal of Ecology 110:1561–1574.

Kusch, E., and R. Davy. 2022. KrigR—a tool for downloading and statistically downscaling climate reanalysis data. Environmental Research Letters 17:024005.

Lamonica, D. et al. 2021. Predicting the dynamics of establishing tree populations: a framework for statistical inference and lessons for data collection. Methods in Ecology and Evolution:2041–210X.13656.

Lutz, J. A. et al. 2012. Ecological importance of large-diameter trees in a temperate mixed-conifer forest. PLoS ONE 7.

Maitner, B. S. et al. 2018. The bien r package: A tool to access the Botanical Information and Ecology Network (BIEN) database. Methods in Ecology and Evolution 9:373–379.

Morales-Castilla, I. et al. 2015. Inferring biotic interactions from proxies. Trends in Ecology and Evolution 30:347–356.

Morueta-Holme, N. et al. 2016. A network approach for inferring species associations from co-occurrence data. Ecography 39:1139–1150.

National Park Service. 2009. Boundary for Yosemite National Park. https://databasin.org/datasets/7ae51cf6-aac8-4636-b1db-eb206ed012d0/.

Ovaskainen, O. et al. 2016. Using latent variable models to identify large networks of species-to-species associations at different spatial scales. Methods in Ecology and Evolution 7:549–555.

Ovaskainen, O. et al. 2017. How to make more out of community data? A conceptual framework and its implementation as models and software. Ecology Letters 20:561–576.

Poisot, T. et al. 2012. The dissimilarity of species interaction networks.

R Core Team. 2021. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

Stephan, P. et al. 2021. Positive species interactions shape species' range limits. Oikos:oik.08146.

Swain, A. et al. 2021. Inferring, comparing and exploring ecological networks from time-series data through R packages constructnet, disgraph and dynet:1–15.

Tikhonov, G. et al. 2020. Joint species distribution modelling with the R-package Hmsc. Methods in Ecology and Evolution 11:442–447.

Tikhonov, G. et al. 2021. hmsc: Hierarchical Model of Species Communities.

Ulrich, W., and N. J. Gotelli. 2010. Null model analysis of species associations using abundance data. Ecology 91:3384–3397.

406   Usinowicz, J., and J. M. Levine. 2021. Climate-driven range shifts reduce persistence of
407   competitors in a perennial plant community. Global Change Biology:1890–1903.
408   Veech, J. A. 2013. A probabilistic model for analysing species co-occurrence. Global
409   Ecology and Biogeography 22:252–260.
410   Violle, C. et al. 2012. The return of the variance: intraspecific variability in community
411   ecology. Trends in Ecology and Evolution 27:244–252.
412   Wilkinson, D. P. et al. 2019. A comparison of joint species distribution models for presence–
413   absence data. Methods in Ecology and Evolution 10:198–211.
414   Woodward, G. et al. 2010. Ecological Networks in a Changing Climate. Page Advances in
415   Ecological Research.
416   World Wildlife Fund. 2020. Terrestrial Ecoregions of the World.
417   http://omap.africanmarineatlas.org/BIOSPHERE/data/note_areas_sp/Ecoregions_Ecosys
418   tems/WWF_Ecoregions/WWFecoregions.htm.
419   Wright, I. J. et al. 2004. The worldwide leaf economics spectrum. Nature 428:821–827.
420   Wright, I. J. et al. 2005. Modulation of leaf economic traits and trait relationships by climate.
421   Global Ecology and Biogeography 14:411–421.
422
423

FIGURE 1 – VARIATION ACROSS APPROACHES OF ECOLOGICAL NETWORK INFERENCE –

Network inference approaches can be distinguished with regards to their biodiversity input data type

(corresponding to the occurrence-performance spectrum, see Figure S1) and their consideration of

environmental conditions. The network inference approaches we selected for our study were chosen due to their

comprehensive coverage of the resulting spectrum of data type use.

TABLE 1 - METHODS OF NETWORK INFERENCE - Description of statistical principles, data
requirements, and ecological interpretation of outputs of the four methods of network inference used in this
study. The evaluated methods are: **Probabilistic Species Co-Occurrence Analysis in R** (COOCCUR**,** Griffith
et al., 2016), **Inference of Species Associations from Co-Occurrence Data** (NETASSOC**,** Blonder &
Morueta-Holme, 2017), **Hierarchical Model of Species Communities** (HMSC**,** Tikhonov et al., 2021), and
Neighbour-Density Dependent - Response Impact Model (NDD-RIM**,** Bimler et al., 2022). In our
implementation of NETASSOC, range estimates for local-scale data were not used in our analyses as they did
not represent informative null models. In our implementation of HMSC**,** trait means for each species were only
used locally for our analyses as functional trait expressions become too diverse at regional scales to be
expressed with a mean value.

Method	Principle	Input	Output	Interpretation of Links
COOCCUR	Probability of species-pair co-occurrence being higher or lower than random.	• Species-by-Site occurrence		Spatial aggregation/disaggregation of species-pairs
NETASSOC	Frequency of species-pair co-occurrence compared to null model.	• Species-by-Site occurrence or abundance • Species-by-Site expectation of range estimates as occurrence	Association matrix	
HMSC	Hierarchical modelling of community assembly processes were residual species-species correlations are treated as association metrics.	• Species-by-site occurrence, abundance, or performance • Environmental conditions at each site • Phylogeny of species • Trait means for each species		Aggregation/disaggregation (in space, abundances, or performance) of species-pairs that aren't explained through other community assembly processes
NDD-RIM	Neighbour-density dependent regression and deconstruction of species interactions into response and effect coefficients for inference of rare interactions.	• Individual/species performance at each site • Species-by-Site matrix of abundance	Interaction matrix	Change in performance of interaction subject with increase in abundance of interaction actor

FIGURE 2 - MODEL SPECIFICATION DRIVES NETWORK INFERENCE – Ecological network adjacency matrices inferred at plot scale using the four methods of network inference with differing specifications of spatial products (i.e., occurrence, abundance, performance). COOCCUR (A), NETASSOC (B), NDD-RIM (C) represent use-cases for occurrence, abundance, and performance data. HMSC (D-F) can incorporate occurrence (D), abundance (E), or performance (F) data. Black outlines highlight species pairings shared across all scales of assessment. Black squares in the network matrices signify statistically significant inferred connections between species. See table S1 for an overview of four-letter abbreviations and corresponding species identities.

449
FIGURE 3 - NETWORK INFERENCE IS NOT CONSISTENT ACROSS METHODS OR SCALES –

Contrasting the ecological networks inferred using four separate methods (A-D) reveals a lack of scale-

consistency when assessing species-species links across geographic scales (i.e., plot scale – left column, region

scale – centre column, and macro scale – right column). Black outlines highlight species pairings shared across

all scales of assessment. Black outlines highlight species pairings shared across all scales of assessment. Black

squares in the network matrices signify statistically significant inferred connections between species. White

squares indicate in NDD-RIM visualisation (D) indicate that only one of the underlying directed links was

identified statistically significantly. See table S1 for an overview of four-letter abbreviations and corresponding

species identities.

FIGURE 4 – NETWORK BETADIVERSITY ACROSS APPROACHES WITHIN SCALES OF

ASSESSMENT – There exists considerably dissimilarity between ecological networks made up of statistically

significant links inferred by the four target approaches within each scale of assessment. Notably, inference at

region scale (centre panel) was held back by low data availability.