Distinct genetic underpinnings of inter-individual differences in the sensorimotor-association axis of cortical organisation

Giacomo Bignardi[1,2*], Michel G. Nivard[3], H. Lina Schaare[4,5], Boris C. Bernhardt[6], Richard A.I. Bethlehem[7,8], Simon E. Fisher[1,9,†], & Sofie L. Valk[4,5,10,† *]

[1] *Language and Genetics Department, Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands;*
[2] *Max Planck School of Cognition, Leipzig, Germany;*
[3] *Department of Biological Psychology, Faculty of Behaviour and Movement Sciences, VU University, Amsterdam, The Netherlands;*
[4] *Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany;*
[5] *Research Center Jülich, Institute for Neuroscience and Medicine (INM-7), Germany;*
[6] *McGill University, Montreal, Canada;*
[7] *Autism Research Centre, Department of Psychiatry, University of Cambridge, Cambridge, United Kingdom;*
[8] *Department of Psychology, University of Cambridge, Cambridge, United Kingdom;*
[9] *Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen 6500 HB, The Netherlands;*
[10] *Heinrich-Heine-Universitaet Duesseldorf, Germany*

* Giacomo Bignardi
* Sofie L Valk.
† These authors share last authorship.
Email: giacomo.bignardi@mpi.nl; valk@cbs.mpg.de

Author Contributions: S.L.V. and G.B. conceptualised the work; S.L.V. & S.E.F. supervised the research; G.B. performed primary analyses; S.L.V., S.E.F, M.G.N, B.C.B., and R.A.I. gave input on analysis and Figures; S.L.V. performed functional connectomics, microstructural and geodesic distance analysis; H.L.S. provided input in diffusion map embedding analysis; G.B. computed Figures; G.B, & S.L.V. drafted the manuscript; S.L.V., S.E.F, M.G.N, B.C.B., H.L.S., and R.A.I. contributed to revising the manuscript.

Competing Interest Statement: The authors declare no competing interest.

Keywords: Neuroscience, Sensorimotor-Association axis, Behaviour Genetics, Structural Equation Modelling, Measurement Error.
This file includes:
>   Main Text
>   Figures 1 to 5

Bignardi et al.

Abstract

The topological differentiation of sensorimotor and association cortical regions along a sensorimotor-
association (S-A) axis has undergone profound evolutionary change along the mammalian lineage. In
humans, patterns of gene expression, neurotransmitter receptors, cell-type markers, microstructure,
and functional connectivity have been shown to vary systematically along such S-A axis. Despite such
a notable convergence, it is not yet known whether patterns of S-A axis inter-individual differences
mirror the robust spatial relationships across neurobiological properties observed at the group-level.
Here, we examined genetic variation associated with inter-individual differences in the S-A axis and
assessed whether structural and functional properties that are highly correlated at the group-level also
share genetic underpinnings. Using an augmented multivariate twin design, we quantified the expected
genetic overlap between properties of the S-A axis, accounting for measurement error heterogeneity
across the cortex and its impact on structure-function relationships. Despite robust spatial relationships
and highly heritable inter-individual differences in S-A axis microstructure and functional organisation,
and contrary to group-level findings, our results indicate partly distinct genetic effects across the
different brain features that co-vary along the S-A axis. Our approach highlights the complexity of
genetic contributions to brain organisation and their potential implications for cognitive variability.

Introduction

The human brain supports perception and action but also abstract cognition (1,2). This diversity of functions is thought to be reflected by the gradual dissociation between unimodal sensory and transmodal association cortical areas along a sensorimotor-association (S-A) axis (3). The S-A axis spans a vast array of neuroanatomical properties, including microstructural variation (myelination and cytoarchitecture) and inter-areal connectivity distance (1,3–6). Here, sensory areas show increased layer differentiation, myelination, and, predominantly, short-range connections. In contrast, association areas show less differentiated microstructural profiles, reduced myelination, and a combination of short- and long-range connectivity profiles (5,7,8). The S-A axis underwent profound evolutionary changes (3), with an expansion of cortical association areas paralleled by a marked laminarisation of sensory areas in human primates (9,10). Such structural re-organisation and evolutionary changes along the S-A axis (1,6,11,12) may have provided the scaffold for functional differentiation (13,14), allowing in turn for human-specific cognitive and behavioural flexibility (1,13).

In recent years, several discoveries have enhanced our understanding of the link between structural and functional features of the S-A axis. These findings have highlighted spatial associations between microarchitectonic differentiation (15) and cortical geometry (16) with intrinsic functional organisation (13,17). For example, T1w/T2w maps derived from non-invasive Magnetic Resonance Imaging (MRI)— indexing cortical microstructural differences—, and histological markers based on cell staining have been shown to relate strongly to gene transcriptional profiles and functional dissociation along the S-A axis. This suggests that a canonical genetic architecture may shape S-A axis structural organisation, in turn allowing for the differentiation of cortical function (6).

Recent studies noted that inter-individual variability of the S-A axis is associated with a host of trait differences, from sex (18) and age (19) to human neuropsychiatric traits, such as autism (20), schizophrenia (21), and depression (22), highlighting the importance of the S-A axis for human complex trait variation. However, despite well-documented strong associations at the group-level (3,4,6,23–25), it is still unclear whether patterns of inter-individual differences in S-A axis properties mirror group-level findings. Knowing whether such patterns converge or diverge is crucial because it can directly inform the study of the impact of S-A axis alterations on complex human traits. For example, differences between individual- and group-level findings could imply that the impact of any given neurobiological S-A property could extend well beyond the effects of other seemingly related features.

Here, we asked: do individual differences in structural properties of the S-A axis relate to differences in functional properties? In other words, we studied whether previously widely reported group averages can inform S-A axis associations at the individual level. Specifically, we tested whether individual regional cortical microstructure (6) and structural cortico-cortical network proximity (26), captured by the geodesic distance of inter-connected regions across the cortical mantle (1,26), relate to the well-known functional dissociations between sensory and transmodal association areas (1,3). To account for the known issue of measurement error heterogeneity across the cortex (27) and its impact on

association estimates (28,29), we adapted and applied measurement error models in the form of
structural equation models (30,31). This allowed us to tease apart unreliable intra-individual from
reliable inter-individual variation in the functional organisation of the S-A axis.

We then focused on upstream sources of inter-individual differences in the S-A axis in structure and
function: genetic variation. We asked: Are genetic effects on the S-A axis shared across structural and
functional properties? Here, we analysed a genetic informative sample and quantified the extent of
overlap across genetic effects on structural and functional properties of the S-A axis. Specifically, we
used an augmented twin-informed design to quantify and tease out genetic overlaps between the S-A
axis's structural and functional properties while accounting for the impact of reliability on classic
heritability estimates (31,32). Last, we evaluated the robustness of our results, both across subsamples,
between regional and global cortical metrics, and between and within individuals' S-A axis properties.

Results

To quantify structural and functional S-A axis properties, we combined microstructural and resting-state
functional MRI (rsfMRI) data from the Human Connectome Project (HCP (33); N = 992 adults; 529
women, mean age 28 y; 22-37 y). We computed two structural metrics and one functional metric
indexing the S-A axis:

- Regional microstructure: we quantified regional microstructure indexing the differentiation
between sensorimotor and association areas using the individuals' mean intensity of regional
T1w/T2w (T1w/T2w$_{mi}$) in 400 parcels (6,34)
- Geodesic distance: we quantified regional cortico-cortical network proximity (26) by computing
the Geodesic Distance (GD) between every cortical region and its corresponding functional
network, averaging within each region to get parcel-wise estimates (20)
- Functional gradient loadings: we quantified individuals' functional S-A axis by obtaining the first
component of the individual functional connectomes (FC$_{G1}$) using diffusion map embedding (1)

We started our analysis by testing whether associations between group-level averaged maps of
structural S-A axis properties correlated with the functional S-A axis (Fig. 1A-C). By using a subsample
of n = 482 adults (229 women, mean age 28 y; 22-37 y; a subsample obtained by excluding respective
all twins included in the full HCP sample), we were able to replicate group-level findings between
averaged T1w/T2w$_{mi}$ and FC$_{G1}$, extending the results to GD and FC$_{G1}$ (Fig. 1D-E).

Figure 1. Structural and functional S-A axes strongly correlate at the group-level. Structural (**A-B**) and functional (**C**) indices of Sensorimotor-Association (S-A) axes plotted on inflated cortical surfaces (57). Values represent averages of individual T1w/T2w mean intensity profiles (A; T1w/T2w$_{mi}$), averages of individual Geodesic Distances (B; GD), and functional gradients loadings (C; FC$_{G1}$) extracted from the average of individual functional connectomes across 400 cortical regions. (**D**) Structural indices are strongly associated with functional indices of the S-A axis; Spearman ρ = -.61 and ρ = .75 between T1w/T2w$_{mi}$ and FC$_{G1}$, and GD and FC$_{G1}$, respectively; all p<.05). Each dot represents a regional value; the colour represents Yeo-Krienen 7 network membership. (**E**) Conceptual representation of group-level analysis. Note that individual and regional information is lost in favour of group-level results.

Pervasive inter-individual differences in the S-A axis of cortical organisation. Having estimated the extent of overlap between structural and functional S-A axis properties at the group-level, we shifted the focus to the individual level. Since group-level S-A axis can mask substantial individual variability (Fig. 2A), we asked: does individual variability in structural S-A axis properties relate to variability in S-A functional properties, as for group-level analysis (Fig. 2B)? By shifting analysis from group-level summary statistics to individual variability, we harnessed the fundamental distinction between intra- and inter-individual differences (35). The first is known to index unreliable and fluctuating variability within individuals over time, while the second indexes the reliable and stable part of the overall variability between individuals (Fig. 2C) (35). This distinction is crucial, as intra-individual variability can downward bias effect sizes, reduce statistical power (35), additionally downward biasing genetic estimates (36), all of the above heterogeneously across the whole cortex (27), and can, therefore, increase reproducibility issues (see (28) for details). We were able to make such a distinction by exploiting one of the strengths of the HCP design, which emphasises multiple resting-state fMRI sessions (across two

days of scanning sessions, ~30 min each). This feature of the HCP design allowed us to partly discard

intra-individual fluctuations in rsfMRI data from inter-individual differences in the functional gradient.

Precisely, we partition the inter-individual variance ($\sigma_{inter}FC_{G1i}^2$) from the overall observed variance in

the functional gradient (σFC_{G1i}^2 for any parcel i) by applying a measurement error model ((30,31) Fig.

2D, see Methods).

Estimates obtained from the measurement error model indicate that 33% of the total variability in the

functional gradient was, on average, accounted for by intra-individual variance (Fig. 2E) even when

using individual functional gradients extracted from functional connectomes averaged across two days

of rsfMRI sessions (totalling ~60 min of scanning session). In other words, estimates for the association

between the functional gradient and other S-A axis properties (or any other variable) would be, on

average, biased downward by a factor of bias(r-observed, r-true) = .82 (a lower bound calculated

assuming perfect reliability for the other S-A axis property (28)). Second, we observed systematic

differences in estimates obtained from the measurement models across functional cortical networks,

$F(6, 393) = 33.21$, $p < .001$; $\eta^2 = 0.34$, 95% CI [0.27, 1.00]), with estimates for parcel-wise inter-individual

variances ranging from $\sigma_{inter-114}^2 = .39$ to $\sigma_{inter-294}^2 = .89$ (Figure 2D, SI Appendix, Fig. S1). In other

words, bias is heterogeneous and expected to influence estimates across the cortex systematically.

187
Figure 2. Pervasive inter-individual differences in the S-A axis of functional connectivity. (**A**)
group-level estimates (black contour) overshadow pervasive individual differences in S-A axis
properties. (**B**) The shift between levels of analyses: from group-level (grey square) to between-
individuals (coloured squares); The gradient square conceptually captures panel A. (**C**) Measurement
error model to partition, for any parcel i, variance in the functional gradient loadings into intra- ($\sigma_{d\text{-}intra(i)}^2$,
for regional values measured at day 1 or 2 of the testing session, i.e., rectangles) and inter- ($\sigma_{inter(i)}^2$, for
the latent component, i.e., circle) individual variance. Parameter estimates for any parcel i can be found

in Supporting File 1. (**D**) The proportion of intra- and inter-individual variance in the functional network
across Yeo-Krienen functional networks: the horizontal line displays the median; lower and upper
hinges correspond to the first and third quartile; the whisker extends from the hinge to the largest/lower
value no further than 1.5 * interquartile range from the hinge. Note that across all parcels, observed
variance includes substantial inter-individual variation. *Notes on measurement model: Rectangles*
represent the measured phenotypes; The circle is the latent component; the double-headed arrows
within the circle represent the variance associated with the latent components; one-headed arrows are
the paths (here all set to 1).
203
204

Individual differences in regional cortico-cortical network proximity, rather than microstructure,
relate to the functional gradient of the S-A axis of cortical organisation. To simultaneously
deattenuate the heterogenous downward biases and handle structural and functional S-A metrics, we
used a Structural Equation Modelling (SEM) approach. Precisely, we specified a model in which the
inter-individual differences in the functional gradient estimated via the measurement error model were
directly tested for associations with microstructural profiles and geodesic distances parcel-wise data
(Fig. 3A). Here, we note that we avoided making assumptions about the causal structure generating
the possible correlations between structural and functional metrics. We simply limited ourselves to
estimating the association between regional properties of the S-A axis. On the one hand, contrary to
group-level topographies, we found less than 2% of the 400 parcels to display a significant association
between individuals' microstructural profiles and functional gradient loadings. These significant
associations were all negative, weak ($-.20 > r > -.27$), and spread across both hemispheres and the
dorsal, ventral, and default-mode functional networks. Conversely, we found large overlaps between
individual geodesic distances and functional gradient loadings (Fig. 3B), with 57% of the 400 parcels
showing significant associations after Bonferroni correction. The directionality of the estimates for the
association between individual regional geodesic distances and functional gradient loadings highlighted
systematic differences across functional networks. Significant positive associations were preferentially
clustered within the visual and the default mode (one sample t-test, two-sided, $t(21) = 5.82$, $p < .001$,
average $r = .51$, and $t(53) = 4.04$, $p = .001$, average $r = .25$), while negative associations were
preferentially clustered within the somatomotor and ventral attention networks (one sample t-test, two-
sided, $t(59) = -13.32$, $p < .001$, average $r = -.48$, and $t(27) = -8.14$, $p < .001$, average $r = -.40$,
respectively, all test corrected for Bonferroni). Estimates obtained from standard correlation analysis
further confirmed that the SEM approach successfully deattenuated measurement error bias (SI
Appendix, Fig. S2).
229

Bignardi et al.

A Between-individual SEM

B Phenotypic correlations with FC$_{G1}$

Figure 3. Structural and functional S-A axes selectively correlate between individuals. (A) Simplified structural measurement error model to estimate the correlation between S-A axis properties considering the distinction between intra- and inter-individual differences in functional gradient loading variability. Observed parcel-wise values and latent components are standardised before model fitting. **(B)** Summary for the standardised estimates on the inflated cortical surface (57) from the structural measurement error model indicates little and weak phenotypic correlations (r_p) between microstructural intensity (T1w/T2w$_{mi}$) and functional gradient loadings (FC$_{G1}$) inter-individual differences but large and highly significant ($p<.05$ corrected for Bonferroni comparisons) overlaps between functional gradient loadings and geodesic distances. All parameter estimates for any parcel i, including the covariances between T1w/T2w$_{mi}$ and GD that were not the focus of the current study, can be found in Supporting File 2. *Notes on structural equation models: double-headed arrows between circles represent covariances; here, path coefficients for the measurement error model are estimated (λ). Other abbreviations and symbols are as in Figure 2.*

Genetic effects on regional S-A axis variability are substantial yet mostly distinct. After having related inter-individual differences in structural and functional S-A axis properties, we asked whether genetic effects were mostly common or distinct across S-A axis properties. Here, we further exploited the HCP's family structure to partition the relationship between structural and functional properties of the S-A axis in common and distinct genetic sources. Precisely, by applying a multivariate twin design, we were able to partition genetic (σ_A^2; A: additive) and unsystematic environmental (σ_E^2; E: Unique-Environmental) sources of variability in microstructure, geodesic distances, and functional gradients loadings. (Here, we note that since intra-individual variability is partitioned in the E component of the model (37), this strategy made it also possible to discard further intra-individual sources of variance in

structural metrics, even in the absence of repeated measures (36)). To do so, we focused our analysis

on the twin HCP subsample, which includes both monozygotic (MZ) and dizygotic (DZ) twins (n = 328,

195 MZ and 133 DZ individual twins, 124 and 88 women, respectively; mean age 29 y, range = 22-35

y; see Methods for details on inclusion criteria).

To partition variability (σ_P^2; p: S-A axis phenotypic property) within S-A modalities in σ_A^2 and σ_E^2 sources

and further unpack genetic and environmental structure-function associations, we specified a

multigroup multivariate model with only A and E components (Fig. 4A). Microstructural profiles,

geodesic distances, and functional gradients loadings all displayed substantial heritability (h_{twin}^2, Fig.

4B), with mean h_{twin}^2 = .43, SD = 0.11, mean h_{twin}^2 = .34, SD = 0.11, and mean h_{twin}^2 = .57, SD = 0.14,

respectively. Consistent with previous work (31), a comparison of univariate models fitted to functional

gradient loadings not accounting for measurement error confirmed that the inclusion of the

measurement error model substantially boosted h_{twin}^2 estimates of 54% relative to h_{twin}^2 not accounting

for intra-individual variance (univariate h_{twin}^2 = .37, SD = 0.11 when not including a measurement error

model, see SI Appendix, Fig. S3). However, notwithstanding such relatively high h_{twin}^2 for both

microstructural profiles and functional gradient loadings, we found no significant additive genetic

correlation between the two (all p > .05, Bonferroni corrected; Fig. 4B). This suggested little room for

possible common genetic causes between microstructural intensity and functional gradient loadings S-

A axis properties. Conversely, 14% of the parcels displayed significant additive genetic correlations

between geodesic distances and functional gradient loadings (7% negative and 6% positive in

directionality, respectively p < .05, Bonferroni corrected, Fig. 4C). The average magnitude of the genetic

correlation (r_A) was r_A = -.67, SD = 0.16, and r_A = .64, SD = 0.16, for the negative and positive

association, respectively. Furthermore, we found that for 30% of the parcels, complementary

environmental effects mostly correlated between geodesic distances and functional gradient loadings.

280
Figure 4. Genetic sources of the S-A axis' structural and functional properties are selectively
distinct. (**A**) Simplified graphical representation of the multivariate twin-informed SEM. All parameter
estimates for any parcel *i* can be found in Supporting File 3**.** (**B**) Variance explained by the additive
genetic (A) and non-common environmental (E) components. Note that there is genetic variation
associated with individual differences across all parcels. (**C**) Summary for the significant additive genetic
and environmental correlations, r_A and r_E, respectively, across S-A axis properties on the inflated cortical
surface (46). Note only significant genetic correlations between cortico-cortical network proximity, as
measured by geodesic distances and functional gradients loadings (FC_{G1}). The r_A and r_E estimates can
be found in Supporting Files 4-5 (panels B and C, respectively). *Notes on structural equation models:*
additional circles represent latent additive genetic and environmental components; Double-headed
arrows between circles represent genetic and environmental covariances. Where not noted, path
coefficients are set to 1. Other abbreviations and symbols are as in Figure 2.
293

Associations between structural and functional properties of the S-A axis are robust across

samples. To test for the robustness of the results discussed so far, we estimated the overlap of the

significant regional genetic or environmental association in the genetically informative subsample with

the significant regional associations obtained from the first subsample. Of the 104 parcels that displayed

either or both significant genetic and environmental correlations between geodesic distances and

functional gradient loadings in the genetically informative sample, 99 also displayed a significant

correlation in the first subsample. In other words, we found a 95% overlap between subsamples. These

results show that regional results were robust across two subsamples drawn from the HCP.

Genetic and environmental associations extend beyond regional S-A axis variability. As a final

analysis, we asked whether associations between geodesic distances and functional gradients were

generalisable beyond regional differences. First, we quantified global S-A axis properties variability as

the overall Median Absolute Deviation (MAD) across all parcels within individuals. Within an individual,

higher MAD scores indicate a larger dispersion in S-A axis values across the cortex. Once more, we

found no significant genetic or environmental associations between microstructural profile intensity and

functional gradients. Yet, we found a substantial negative genetic correlation between the geodesic

distances and functional gradient between individuals' S-A axis MAD scores (r_A = -.78, 95% CI [-1.19,

-.34], CFI = .93, RMSEA = .04; Fig. 5A). Additionally, to obtain a complementary estimate of global S-

A axis variability, we quantified microstructural profile intensity, geodesic distance, and functional

gradient ρ similarity indices. These indices assessed how similar S-A axis properties in one individual

are compared to the average. Consistent with regional and global variances differences, ρ similarity

indices in geodesic distance, rather than microstructure, showed strong and positive genetic

correlations with global differences in the functional gradient (r_A = .61, 95% CI [.43, .79], CFI = .95,

RMSEA = .04; Fig. 5B). Findings were robust to intra-cranial volume as a possible common cause of

S-A axis structure-function covariance (SI Appendix, Fig. S4)

A Multivariate model for overall within-individual S-A axis MAD   **B** Multivariate model for S-A axis ρ similarity indices

Figure 5. Findings extend beyond regional S-A axis associations. (A) Simplified graphical representation of the multivariate twin-informed SEM for overall within-individual Median Absolute Deviations (MAD). Note the strong but negative significant associations between the latent additive genetic components underlying geodesic distances (GD; centre) and functional gradient loadings (FC$_{G1}$; right). **(B)** Simplified graphical representation of the multivariate twin-informed SEM for ρ similarity indices (Fisher-z transformed). As for the model reported in panel A, the only significant associations are found between GD and FC$_{G1}$. * $p < .05$. *Notes on structural equation models: abbreviations and symbols are as in Fig. 4. Here, double-headed arrows between latent variables indicate correlations (since we report standardised solutions). Dashed arrows represent nonsignificant correlations.*

Discussion

Our results collectively reveal that group-level estimates of spatial associations between structural and functional gradients differentiating sensory from association areas in the human brain might mask pervasive inter-individual differences. These inter-individual differences, in turn, might display different patterns of associations to the one depicted at the group level. Notwithstanding the comparable heritability of the S-A axis properties as shown in this study, theoretical models of S-A axis development and evolution (3), and group-level relationships between patterns of gene expression, cortical microstructure, and functional differentiation of sensorimotor to transmodal-association areas (6), we found little evidence overlap between microstructural and functional S-A axis similarities at the level of the individual. Precisely, we found little evidence of phenotypic and an absence of evidence for genetic or environmental associations between cortical microstructure (as measured by T1w/T2w) and S-A function (as measured by the principal gradient of functional connectivity) of the cortex. At the same time, our results showed substantial genetic and environmental associations between individual-level differences in regional and global cortico-cortical network proximity (as measured by the geodesic distance of inter-connected regions across the cortical mantle) and function. These latter results align with theories emphasising geometric constraints of brain function, yet do not fully align with previous and current group-level estimates. While group-level associations indicate a positive relationship between cortico-cortical network proximity, our results uncover a mixture of positive and negative relationships at the individual level of analysis (the former preferentially clustered within the visual and default mode network, the latter with the somatomotor and the ventral attention network). Moreover, we found negative, not positive, genetic correlations when shifting from local to global association, as we did when analysing overall within-individual S-A axis dispersion. This suggests that genetic differences between people that tend to co-occur with decreased variation in geodesic distances across the cortex also tend to co-occur with more dispersed functional gradients.

Fundamental principles of brain organisation can appear to be highly conserved across neurobiological
properties if based on group-based analyses of topological co-variation, yet our results suggest this
conservation breaks down when assessed at the level of the individual. Based on group-level
associations, previous work suggested that cortical maturation of diverse neurobiological properties
proceeds along an evolutionary conserved and developmentally rooted S-A axis of cortical organisation
(3,4). However, our results indicate that genetic variation within a population is expected to be
selectively associated with some properties (e.g., function and cortico-cortical network proximity) but
not others (e.g., microstructure). A possible explanation may lie in the physiological and cognitive
implications of the resting state signal and downstream effects on its functional gradient. It is possible
that, although in general terms resting state networks are topologically organised along the S-A axes,
their individual fluctuations reflect physiological variability not captured by its microstructural scaffold
(38). Indeed, various studies have reported that network integration and segregation align with neural
gain, which could be explained by alterations in neuromodulatory systems (39,40).

Regardless of the possible explanation, these findings are important for understanding the origins of
differences between individuals, such as in various neuropsychiatric disorders or work on brain-
behaviour associations in general. Complementing studies of S-A axis variability with informative
genetic models of structural and functional S-A axis variability would allow us to capture how genetic
effects on S-A variability contribute to variation in human neuropsychiatric traits, facilitating novel
mechanistic insights into these notoriously complex phenotypes. We also note that such models can
be easily applied to many neurobiological properties (e.g., resting-state fMRI, see (31)) to enhance
current brain-behaviour mapping efforts (41). To facilitate this, we have made all the code available and
provided all SEM functions in R and lavaan syntax.

The measurement error modelling approach can successfully tease apart unstable intra-individual
differences from stable inter-individual differences, and this effect can have a substantial downstream
impact on estimates. For example, applying the measurement error modelling approach, in line with
previous results (31), resulted in a nearly 50% increase in estimates of heritability. We foresee that this
approach could have further direct application in ongoing research on the origins of psychiatric disorders
and brain-behaviour studies, and in the analysis of the genomic architecture of principles of brain
organisation, for example, by mitigating the impact of measurement error heterogeneity on estimates.
Indeed, when individual variability in the S-A axis is the predictor of interest, such as in brain-behaviour
studies, applying any measurement error model is expected to deattenuate downwardly biased
estimates (28,35,42,43). Moreover, genome-wide association studies could easily implement genome-
based structural equation modelling (44,45) extensions of our approach to discard unstable and
unreliable variance, overcoming attenuation biases in associations between single nucleotide
polymorphisms and target phenotypes (e.g., similarly to what has been done for analyses based on
polygenic indices (43)).

Yet, it is important to note that the interpretation of deattenuated estimates rests upon assumptions of
what type of error is expected to influence the measurement of functional S-A axis properties across
days of measurement. Here, we assumed that what is measured as being shared between sessions is
the common cause of what is measured within each session. Within this framework, what is left is
unique to each session, including sources of measurement error. Such application of the measurement
error model to S-A axis functional properties is not expected to perfectly segregate S-A axis error-free
variance in the inter-individual component (35). For example, the systematic error of individual S-A axis
measurement would be indistinguishable from meaningful inter-individual variability and still be
captured by the inter-individual variance component. Additionally, this measurement error model could
confound unsystematic S-A axis error with genuine intra-individual differences, which may even be
partially heritable (46). Therefore, the actual sources of session-to-session differences may be more
nuanced than the simple measurement error model implies. Consequently, we advise caution when
interpreting inter- and intra-individual differences as exclusively "error-free" and "error" variance,
respectively (see Methods for further details).
409

Another limitation in our study is the lack of repeated structural metric measures. Although applying a
measurement error model allowed us to partly disentangle intra- and inter-individual variability in
functional gradient loadings, we could not account for the differences in structural properties within
individuals. This limitation may have attenuated the estimated relationship between structure and
function. However, the nature of the metrics and the twin design employed to elucidate differences
between individuals should mitigate the impact of such a lack of repeated structural metrics (36). By
applying the classical twin design, we were able to further partition unstable measurement error in the
environmental (E) component of the model, which minimised possible biases introduced by hypothetical
measurement error, at least for the additive genetic (A) correlations (r_A) estimates.
419

In sum, our findings reveal that group-level results can overshadow substantial inter-individual
differences within and between different neurobiological properties. By focusing on these previously
underappreciated differences, we could highlight selective associations of individual variation in S-A
axis cortical structure and function. These inter-individual differences and associations open a window
into genetic sources of S-A axis structure and function, which we reveal to be selectively distinct. These
results underscore the complex interplay between the S-A axis's structural and intrinsic functional
properties, and provide a set of tools that can be used to test their potentially differential roles in shaping
cognition.
428

Materials and Methods
Sample. We used data from the Human Connectome Project (HCP) S1200 release. The HCP includes
data from 1206 individuals (656 women) that comprise 298 Monozygotic (MZ) twins, 188 Dizygotic (DZ)
twins, and 720 individuals, with mean age ± *SD* = 28.8 ± 3.7 years (age range = 22-37 years). Informed
consent for all individuals was obtained by HCP, and our data usage was approved by HCP and
complied with all relevant ethical regulations for working with human participants (see (13,33,47)). The

435 primary participant pool comes from individuals born in Missouri to families that include twins, sampled
436 as healthy representatives of ethnic and socioeconomic diversity of US individuals, based on data from
437 the Missouri Department of Health and Senior Services Bureau of Vital Records. We followed standard
438 guidelines for inclusion criteria as described elsewhere (13). Our sample, in line with Valk et al., (13)
439 comprised 992 (529 women) individuals. The first subsample of n = 482 (229 women) was created by
440 excluding all individual twins. The second genetically informative subsample of n = 328 (212 women)
441 was created by including only individual twins with genotyped zygosity matching self-reported zygosity
442 (195 MZ and 133 DZ; 124 women and 88 women, respectively).

444 **Functional imaging.** Functional connectivity matrices were based on four 14 min 33 s of functional
445 Magnetic Resonance Imaging (fMRI) data acquired over two sessions, spaced two days apart, through
446 the HCP, which underwent HCP's minimal preprocessing. For each individual, four functional
447 connectivity matrices were computed using the minimally preprocessed, spatially normalised resting-
448 state fMRI (rsfMRI) scans, which were co-registered using MSMAll to template HCP 32k_LR surface
449 space. 32k_LR surface space consists of 32,492 total nodes per hemisphere (59,412 excluding the
450 medial wall). We computed four functional connectivity matrices per individual from the average time
451 series extracted in each of the 400 Schaefer cortical parcels. The individual functional connectomes
452 were generated by averaging preprocessed time series within nodes, Pearson correlating nodal time
453 series and converting them to Fisher-z scores. The average functional connectomes were obtained by
454 averaging functional connectomes within individuals (i.e., between sessions) and between individuals.

455 **Structural imaging.** MRI protocols of the HCP have been previously described (33,47). MRI data were
456 acquired originally on the same day on the HCP's custom 3T Siemens Skyra equipped with a 32-
457 channel head coil. T1w images with identical parameters were acquired using a 3D-MP-RAGE
458 sequence over 7 min 40 s (0.7 mm isovoxels, matrix = 320 × 320, 256 sagittal slices; TR = 2400 ms,
459 TE = 2.14 ms, TI = 1000 ms, flip angle = 8°; iPAT = 2). T2w images were acquired using a 3D T2-
460 SPACE sequence with identical geometry over 8 min and 24 s (TR = 3200 ms, TE = 565 ms, variable
461 flip angle; iPAT = 2). We followed the preprocessing steps outlined in Valk et al. (13).

462 **Parcellation and functional networks**. We used the Schaefer group-level hard-parcellation, originally
463 obtained by a gradient-weighted Markov random field model integrating local gradient and global
464 similarity approaches (34). To stratify results within established cortical functionally coupled networks,
465 we used the seven Yeo-Krienen networks (48).

466 **Microstructural profiles (T1w/T2w$_{mi}$).** We used T1w/T2w imaging myelin-sensitive contrast from the
467 HCP minimal processing pipeline, which uses the T2w to correct for inhomogeneities in the T1w image
468 to estimate mean intensity T1w/T2w microstructural profiles (T1w/T2w$_{mi}$). T1w/T2w$_{mi}$ has been shown
469 to map to model-based tract-tracing histological data in macaque, estimate intracortical myelin content,
470 and thus approximate architectural complexity and cortical hierarchy (6).

471 **Geodesic distance (GD).** Individual Geodesic Distances (GD) were computed using the Micapipe
472 toolbox (26). Briefly, we computed GD between each region and their top 10% of maximally functionally

connected regions along each individual native cortical midsurface. We further averaged within regions

to obtain a parcel-wise value and improve computation performance. Micapipe implements the Dijkstra

algorithm (49) (further details can be found in (26)).

Functional gradient loadings (FC$_{G1}$). We sequentially averaged FCs, first within days, resulting in two

FCs per individual, and then between days, resulting in one FC per individual. We then extracted the

three first components from the two sequentially averaged and one averaged FCs, using the Python

package BrainSpace (50). Extraction of the first eigenvector followed standard procedures, with the

original individual FCs set at a connection density of 10% (i.e., the FCs were made sparse by setting a

sparsity threshold of 90%). The first ten eigenvectors were then obtained by decomposing the FCs by

diffusion map embedding, a robust non-linear manifold learning technique (1). To aid comparability

across individuals, we aligned individual eigenvectors to the template eigenvector by Procrustes

rotation (51). The template functional gradient was directly extracted from the overall mean FC matrix.

Group-level associations analysis. We computed Spearman rank-order correlations (ρ) between the

structural (T1w/T2w$_{mi}$ and GD) and functional (FC$_{G1}$) S-A axis group-level properties. Group-level

properties were obtained from the average of the individual structural S-A properties (i.e., average

T1w/T2w$_{mi}$ and GD), and from the decomposition of the average FC (i.e., principal gradient obtained

via diffusion map embedding of the average FC).

Measurement model of error in individual variability of the functional S-A axis. To partition stable

inter-individual variability in functional gradient loading, we adapted previous measurement error

models to rsfMRI to gradients (30,31). The intuition behind such a modelling strategy is simple. Suppose

parcel-wise values are measured without error and are stable over a reasonable period of time (e.g.,

one day). In that case, the correlations across individuals between the values obtained across two time

points will equal 1. If the correlations deviate from 1 instead, regional values will be measured with

some error, with bigger deviations corresponding to higher error or fluctuation over time. As can be

seen in Fig. 2C, when errors or changes over time are present, it may become difficult to distinguish

regional differences between individuals from regional differences within individuals. In this case, we

can use the measurement error model to estimate what stays constant across time, indexing the "true"

regional values. Across the manuscript, for correctness, since "error" variance can include meaningful,

yet unstable, fluctuation in rsfMRI, while "true" variance can also consist of systematic measurement

error across sessions, we refer to the former term as intra-individual and the latter as inter-individual

variability (35). First, we fit a measurement model to parcel-wise functional gradient loadings averaged

within days. In line with Teeuw et al. (31), we did not constrain intra-individual variance components to

be equal across days of scanning sessions. We performed model fitting in lavaan (44) after

standardising observed variables (i.e., std.ov = T). We then used model estimates obtained for the

variances of the latent and observed components. Using Spearman-Brown correction, we computed

the averaged proportion of stable inter-individual variance in functional gradient loadings across days

as the intra-class correlation (ICC) (52). For each parcel *i,* the ICC was calculated as follows:

510
$$ICC(2,k)_{(i)} = \frac{k * ICC(2,1)_{(i)}}{1 + (k-1) * ICC(2,1)_{(i)}}$$

511 Where *k* is a constant equal to the number of measures (i.e., *k* = 2) and the $ICC(2,1)_{(i)}$ is calculated as
512 follows:

513
$$ICC(2,1)_{(i)} = \frac{\sigma^2_{inter(i)}}{\sigma^2_{inter(i)} + (\frac{\sigma^2_{1-intra(i)} + \sigma^2_{2-intra(i)}}{2})}$$

514 $ICC(2,k)_{(i)}$ estimates the proportion of inter-individual variance over the total variance, $\sigma^2_{inter(i)}$, in the
515 functional loadings as if they were obtained from the average of the two scanning sessions. The
516 proportion of intra-individual variance for a parcel *i*, $\sigma^2_{intra(i)}$, is obtained simply by subtracting the
517 $ICC(2,k)_{(i)}$ from 1.

518 The expected bias for any parcel *i* was calculated following Tiego et al. (28):

519
$$bias_{(i)} = \sqrt{R_{p,p} * R_{FCG1,FCG1(i)}}$$

Where $R_{p,p}$, the reliability for the structural S-A axis property *p* (e.g., T1w/T2w$_{mi}$), was set to be equal
to 1 across all parcels, and $R_{FCG1,FCG1(i)}$, the reliability of parcel-wise value for the functional gradient
loading, was calculated as $ICC(2,k)_{(i)}$.

Structural Measurement Error Equation Modelling. We used Structural Equation Modelling (SEM)
to estimate correlations between structural (i.e., T1w/T2w$_{mi(i)}$, GD$_{(i)}$) and functional (i.e, FC$_{G1(i)}$) S-A
modalities. Each multivariate model simultaneously accounted for intra-individual variances by including
the measurement error model. All models were fitted in lavaan (53) after standardising all observed
variables (i.e., std.ov = T). Prior to model fitting, sex and age were regressed from parcel-wise S-A axis
values using the function umx::umx_residualize() (54). Structural equation models were fit to residual
scores. We assumed missing data to be missed at random and followed parameters' estimation via full-
information Maximum Likelihood (i.e., missing = "ML").

Twin-informed Multivariate Structural Equation Modelling. We used multigroup SEM to partition
parcel-wise variability in structural ($\sigma T1w/T2w_{mi(i)}{}^2$, $\sigma GD_{(i)}{}^2$) and functional ($\sigma^2_{inter(i)}$) S-A modalities into
either additive genetic (σ_A^2) and unsystematic environmental (σ_E^2) sources of variance. Structural
equation models were fit to T1w/T2w$_{mii}$, GD$_i$, and FC$_{G1i1}$ (day 1) and FC$_{G1i2}$ (day 2) data, grouped by
zygosity (i.e., two groups). The model specification was informed by the multivariate twin design (55).
Briefly, monozygotic (MZ) twins are ~100 % genetically identical, coming from the same fertilised egg.
In contrast, dizygotic (DZ) twins are, on average, only 50% additively genetically similar regarding allelic
variants coming from two different fertilised eggs. Thus, the correlation between the additive genetic
component (A) is set to be equal to 1 for MZ while .5 for DZ. In contrast, each twin's unique environment
(E) component will be unique; therefore, their correlation will be equal to 0. In total, we fit one
multivariate AE model per parcel. Following the measurement error procedure outlined above and in
ref. (31), a common pathway measurement error model was included in the specification of the

Bignardi et al.

multigroup SEM. As such, each multivariate model simultaneously accounted for intra-individual
variance. We employed the direct symmetric approach by estimating variance components directly
while setting path coefficients to 1 (with the exception of the measurement model, for which we fixed
the variance to be equal to 1, and estimated the path coefficients, instead). We chose this approach as
it has been shown to reduce type I errors and produce asymptotically unbiased χ^2 (56).

All models were fitted in lavaan (53), with standardisation of observed variables before model fitting
(i.e., std.ov = T). Similarly to what was reported above, to control for the effect of age and sex on S-A
axis properties, we residualised parcel-wise variables prior to modelling using the function
umx::umx_residualize() (54). Residuals were used as observed variables in later twin modelling. We
estimated parameters via full-information Maximum Likelihood (i.e., missing = "ML") and evaluated the
goodness of fit for each parcel by comparative fit index (CFI) and root mean square error of
approximation (RMSEA) scores. Following standard cut-offs (31), we retained only models with a
"satisfactory" CFI>.90 and an RMSEA<.08. Narrow-sense twin heritability (h_{twin}^2) estimates for each
parcel i were defined as the ratio of the additive genetic variance over the sum of the additive genetic
and environmental variances:

558
$$h^2_{\text{twin-p}(i)} = \frac{\sigma^2_{\text{Ap}(i)}}{\sigma^2_{\text{Ap}(i)} + \sigma^2_{\text{Ep}(i)}}$$

559 Where $\sigma_{\text{Ap}(i)}^2$ is the additive genetic variance for the S-A axis parcel-wise value for the given property p
560 (e.g., GD). For functional gradient loadings, the heritability was calculated as

561
$$h^2_{\text{twin-inter}(i)} = \sigma^2_{\text{Ainter}(i)}$$

562 After imposing the equality constrain on the common factor FC$_{\text{G1}inter(i)}$

563
$$\sigma^2_{\text{Ainter}(i)} + \sigma^2_{\text{Einter}(i)} = 1$$

564 Genetic correlations (r_A) were calculated as:

565
$$r_A = \frac{\sigma_{\text{Ap1i,Ap2i}}}{\sqrt{\sigma^2_{\text{Ap1i}} * \sigma^2_{\text{Ap2i}}}}$$

Where $\sigma_{\text{Ap1,Ap2}}^2$ is the additive genetic covariance between two S-A axis properties, $p1$ and $p2$ (e.g., GD
and T1w/T2w$_{\text{mi}}$). Environmental correlations (r_E) were calculated similarly to r_A but using environmental
variance and covariances estimates.

Generalisation beyond regional associations. For each individual, we obtain two metrics for
structural and functional S-A axis properties (i.e., a total of six measures per individual):

Overall within-individual Median Absolute Deviation: we quantified the spread of the regional values
across the cortex by computing within-individual Median Absolute Deviation (MAD) of microstructure,
geodesic distances, and functional gradient loadings. MAD is a robust univariate measure of statistical
dispersion and is simply calculated as follows:

575
$$MAD_{pj} = med(|p_{pji} - med(p_{pj})|)$$

Where p_{ij} is the parcel-wise value for a property *p*, an individual *j*, and a parcel *I* and *med* is the median.

ρ similarity index: we obtained the similarity index by estimating the Spearman rank (ρ) correlations
between each individual microstructure, cortico-cortical network proximity, and functional gradient
loadings with the respective S-A group-level modality vectors. For example, the ρ similarity index for the
cortico-cortical network proximity for an individual *j* was obtained by correlating their GD with the group-
level GD. Similarly, for the same individual *j*, the similarity index for their functional gradient loading was
obtained by correlating their FC_{G1} on day 1 and on day 2 of scanning with the group-level FC_{G1}.

Similar to what is outlined above for regional analysis, we fit two multivariate AE models, one per metric.
Before model fitting, ρ similarity indices were first Fisher-z transformed. To recapitulate regional analysis
as closely as possible within the multivariate model, we also included the measurement error model to
overall within-individual MAD and ρ similarity index functional gradient loadings obtained on days 1 and
2 of scanning sessions. Note that standardised coefficients are obtained using the
lavaan::standardizedSolution() function. As a final sensitivity analysis, to discount individuals' whole
brain volume as a possible confounding effect of the relationship between SA axis structure-function
associations, we additionally included total intra-cranial volume. Precisely, we followed a two-step
procedure to discount intra-cranial volume as a possible common cause. First, we regressed out intra-
cranial volume from overall within-individual MAD and Fisher-z transformed ρ similarity indices for all S-
A axis properties. We then re-fit the exact multivariate twin models to the residuals.

Data availability
We obtained human data from the open-access Human Connectome Project (HCP) S1200 young
adult sample. HCP Young Adult data are available at https://www.humanconnectome.org/study/hcp-
young-adult. Supporting Files 1-5 with summary statistics can be found at
https://github.com/giacomobignardi/h2_SA_axis/tree/main/SI.

Code availability
All code is available and can be found at https://github.com/giacomobignardi/h2_SA_axis . SEM and
twin-based analysis were carried out using the statistical package latent variable analysis (lavaan)
https://lavaan.ugent.be/. The function to apply the measurement error model (meermo) can be found
here https://github.com/giacomobignardi/h2_SA_axis/tree/main/R/functions/meermo. The lavaan
syntax for latent variable analysis of twin data (lavaantwda) can be found in the repository
https://github.com/giacomobignardi/h2_SA_axis/tree/main/R/functions/lavaantwda. An introduction to
twin modelling using lavaan can be found at https://rpubs.com/MichelNivard/798608. Code and tutorial
for functional gradient decomposition of functional connectomes are available
at https://brainspace.readthedocs.io/en/latest/pages/install.html. The code and tutorial to generate
geodesic distances can be found at https://micapipe.readthedocs.io/en/latest/.

Bignardi et al.

Acknowledgements

We want to thank the Human Connectome Project, Washington University, the University of Minnesota,

and Oxford University Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil;

1U54MH091657) originally funded by the 16 N.I.H. Institutes and Centers that support the N.I.H.

Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at

Washington University. We would also like to thank Jitse Amelink, Meike D. Hettwer, and MacKenzie

D. Trupp for their comments on an earlier version of the manuscript. This study was supported by

the German Federal Ministry of Education and Research (BMBF), the Max Planck Society, and the Max

Planck School of Cognition.

Bignardi et al.

References

1. Margulies DS, Ghosh SS, Goulas A, Falkiewicz M, Huntenburg JM, Langs G, et al. Situating the default-mode network along a principal gradient of macroscale cortical organization. Proc Natl Acad Sci. 2016 Nov 1;113(44):12574–9.

2. Mesulam MM. From sensation to cognition. Brain. 1998 Jun 1;121(6):1013–52.

3. Sydnor VJ, Larsen B, Bassett DS, Alexander-Bloch A, Fair DA, Liston C, et al. Neurodevelopment of the association cortices: Patterns, mechanisms, and implications for psychopathology. Neuron. 2021 Sep 15;109(18):2820–46.

4. Sydnor VJ, Larsen B, Seidlitz J, Adebimpe A, Alexander-Bloch AF, Bassett DS, et al. Intrinsic activity development unfolds along a sensorimotor–association cortical axis in youth. Nat Neurosci. 2023 Apr;26(4):638–49.

5. von Economo CF, Koskinas GN. Die cytoarchitektonik der hirnrinde des erwachsenen menschen. J Springer. 1925;

6. Burt JB, Demirtaş M, Eckner WJ, Navejar NM, Ji JL, Martin WJ, et al. Hierarchy of transcriptomic specialization across human cortex captured by structural neuroimaging topography. Nat Neurosci. 2018 Sep;21(9):1251–9.

7. Brodmann K. ergleichende Lokalisationslehre der Grosshirnrinde in ihren Prinzipien dargestellt auf Grund des Zellenbaues. Barth. 1909;

8. Ercsey-Ravasz M, Markov NT, Lamy C, Van Essen DC, Knoblauch K, Toroczkai Z, et al. A predictive network model of cerebral cortical connectivity based on a distance rule. Neuron. 2013 Oct 2;80(1):184–97.

9. Buckner RL, Krienen FM. The evolution of distributed association networks in the human brain. Trends Cogn Sci. 2013 Dec 1;17(12):648–65.

10. García-Cabezas MÁ, Hacker JL, Zikopoulos B. Homology of neocortical areas in rats and primates based on cortical type analysis: an update of the Hypothesis on the Dual Origin of the Neocortex. Brain Struct Funct. 2023 Jun 1;228(5):1069–93.

11. Saberi A, Paquola C, Wagstyl K, Hettwer MD, Bernhardt BC, Eickhoff SB, et al. The regional variation of laminar thickness in the human isocortex is related to cortical hierarchy and interregional connectivity. PLoS Biol. 2023 Nov;21(11):e3002365.

12. Markov NT, Ercsey-Ravasz M, Van Essen DC, Knoblauch K, Toroczkai Z, Kennedy H. Cortical high-density counterstream architectures. Science. 2013 Nov 1;342(6158):1238406.

13. Valk SL, Xu T, Paquola C, Park B yong, Bethlehem RAI, Vos de Wael R, et al. Genetic and phylogenetic uncoupling of structure and function in human transmodal cortex. Nat Commun. 2022 May 9;13(1):2341.

14. Demirtaş M, Burt JB, Helmer M, Ji JL, Adkinson BD, Glasser MF, et al. Hierarchical Heterogeneity across Human Cortex Shapes Large-Scale Neural Dynamics. Neuron. 2019 Mar 20;101(6):1181-1194.e13.

15. Paquola C, Amunts K, Evans A, Smallwood J, Bernhardt B. Closing the mechanistic gap: the value of microarchitecture in understanding cognitive networks. Trends Cogn Sci. 2022 Oct 1;26(10):873–86.

16. Leech R, Vos De Wael R, Váša F, Xu T, Austin Benn R, Scholz R, et al. Variation in spatial dependencies across the cortical mantle discriminates the functional behaviour of primary and association cortex. Nat Commun. 2023 Sep 13;14(1):5656.

17. Suárez LE, Markello RD, Betzel RF, Misic B. Linking Structure and Function in Macroscale Brain Networks. Trends Cogn Sci. 2020 Apr 1;24(4):302–15.

18. Serio B, Hettwer MD, Wiersch L, Bignardi G, Sacher J, Weis S, et al. Sex differences in functional cortical organization reflect differences in network topology rather than cortical morphometry. Nat Commun. 2024 Sep 4;15(1):7714.

19. Dong HM, Margulies DS, Zuo XN, Holmes AJ. Shifting gradients of macroscale cortical organization mark the transition from childhood to adolescence. Proc Natl Acad Sci. 2021 Jul 13;118(28):e2024448118.

20. Hong SJ, Vos de Wael R, Bethlehem RAI, Lariviere S, Paquola C, Valk SL, et al. Atypical functional connectome hierarchy in autism. Nat Commun. 2019 Mar 4;10(1):1022.

21. Dong D, Yao D, Wang Y, Hong SJ, Genon S, Xin F, et al. Compressed sensorimotor-to-transmodal hierarchical organization in schizophrenia. Psychol Med. 2023 Feb;53(3):771–84.

22. Xia M, Liu J, Mechelli A, Sun X, Ma Q, Wang X, et al. Connectome gradient dysfunction in major depression and its association with gene expression profiles and treatment outcomes. Mol Psychiatry. 2022 Mar;27(3):1384–93.

23. Huntenburg JM, Bazin PL, Margulies DS. Large-Scale Gradients in Human Cortical Organization. Trends Cogn Sci. 2018 Jan 1;22(1):21–31.

24. Zhang XH, Anderson KM, Dong HM, Chopra S, Dhamala E, Emani PS, et al. The cell-type underpinnings of the human functional cortical connectome. Nat Neurosci. 2024 Nov 21;1–11.

25. Goulas A, Changeux JP, Wagstyl K, Amunts K, Palomero-Gallagher N, Hilgetag CC. The natural axis of transmitter receptor distribution in the human cerebral cortex. Proc Natl Acad Sci. 2021 Jan 19;118(3):e2020574118.

26. Cruces RR, Royer J, Herholz P, Larivière S, Vos de Wael R, Paquola C, et al. Micapipe: A pipeline for multimodal neuroimaging and connectome analysis. NeuroImage. 2022 Nov 1;263:119612.

27. Helwegen K, Libedinsky I, Heuvel MP van den. Statistical power in network
neuroscience. Trends Cogn Sci. 2023 Mar 1;27(3):282–301.

28. Tiego J, Martin EA, DeYoung CG, Hagan K, Cooper SE, Pasion R, et al. Precision
behavioral phenotyping as a strategy for uncovering the biological correlates of
psychopathology. Nat Ment Health. 2023 May;1(5):304–15.

29. Kang K, Seidlitz J, Bethlehem RAI, Xiong J, Jones MT, Mehta K, et al. Study design
features increase replicability in brain-wide association studies. Nature. 2024 Nov 27;1–
9.

30. Brandmaier AM, Wenger E, Bodammer NC, Kühn S, Raz N, Lindenberger U. Assessing
reliability in neuroimaging research through intra-class effect decomposition (ICED).
Johansen-Berg H, Kastner S, editors. eLife. 2018 Jul 2;7:e35718.

31. Teeuw J, Hulshoff Pol HE, Boomsma DI, Brouwer RM. Reliability modelling of resting-
state functional connectivity. NeuroImage. 2021 May 1;231:117842.

32. Ge T, Holmes AJ, Buckner RL, Smoller JW, Sabuncu MR. Heritability analysis with repeat
measurements and its application to resting-state functional connectivity. Proc Natl
Acad Sci. 2017 May 23;114(21):5521–6.

33. Van Essen DC, Ugurbil K, Auerbach E, Barch D, Behrens TEJ, Bucholz R, et al. The Human
Connectome Project: A data acquisition perspective. NeuroImage. 2012 Oct;62(4):2222–
31.

34. Schaefer A, Kong R, Gordon EM, Laumann TO, Zuo XN, Holmes AJ, et al. Local-Global
Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI.
Cereb Cortex. 2018 Sep 1;28(9):3095–114.

35. Zuo XN, Xu T, Milham MP. Harnessing reliability for neuroscience research. Nat Hum
Behav. 2019 Aug;3(8):768–71.

36. Hagenbeek FA, Hirzinger JS, Breunig S, Bruins S, Kuznetsov DV, Schut K, et al. Maximizing
the value of twin studies in health and behaviour. Nat Hum Behav. 2023 Jun;7(6):849–
60.

37. Falconer DS, Mackay TFC. Introduction to Quantitative Genetics. 4 edition. Harlow:
Pearson; 1996. 480 p.

38. Laumann TO, Snyder AZ. Brain activity is not only for thinking. Curr Opin Behav Sci. 2021
Aug 1;40:130–6.

39. Shine JM, Aburn MJ, Breakspear M, Poldrack RA. The modulation of neural gain
facilitates a transition between functional segregation and integration in the brain. Deco
G, editor. eLife. 2018 Jan 29;7:e31130.

40. Luppi AI, Hansen JY, Adapa R, Carhart-Harris RL, Roseman L, Timmermann C, et al. In
vivo mapping of pharmacologically induced functional reorganization onto the human
brain's neurotransmitter landscape. Sci Adv. 2023 Jun 14;9(24):eadf8332.

41. Marek S, Tervo-Clemmens B, Calabro FJ, Montez DF, Kay BP, Hatoum AS, et al.
Reproducible brain-wide association studies require thousands of individuals. Nature.
2022 Mar;603(7902):654–60.

42. Xu T, Kiar G, Cho JW, Bridgeford EW, Nikolaidis A, Vogelstein JT, et al. ReX: an integrative
tool for quantifying and optimizing measurement reliability for the study of individual
differences. Nat Methods. 2023 Jul;20(7):1025–8.

43. van Kippersluis H, Biroli P, Dias Pereira R, Galama TJ, von Hinke S, Meddens SFW, et al.
Overcoming attenuation bias in regressions using polygenic indices. Nat Commun. 2023
Jul 25;14(1):4473.

44. Pritikin JN, Neale MC, Prom-Wormley EC, Clark SL, Verhulst B. GW-SEM 2.0: Efficient,
Flexible, and Accessible Multivariate GWAS. Behav Genet. 2021 May;51(3):343–57.

45. Grotzinger AD, Rhemtulla M, de Vlaming R, Ritchie SJ, Mallard TT, Hill WD, et al.
Genomic structural equation modelling provides insights into the multivariate genetic
architecture of complex traits. Nat Hum Behav. 2019 May;3(5):513–25.

46. Kemper KE, Sidorenko J, Wang H, Hayes BJ, Wray NR, Yengo L, et al. Genetic influence
on within-person longitudinal change in anthropometric traits in the UK Biobank. Nat
Commun. 2024 May 6;15(1):3776.

47. Glasser MF, Sotiropoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL, et al. The
minimal preprocessing pipelines for the Human Connectome Project. NeuroImage. 2013
Oct 15;80:105–24.

48. Yeo BTT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M, et al. The
organization of the human cerebral cortex estimated by intrinsic functional connectivity.
J Neurophysiol. 2011 Sep;106(3):1125–65.

49. Dijkstra EW. A note on two problems in connexion with graphs. Numer Math. 1959 Dec
1;1(1):269–71.

50. Vos de Wael R, Benkarim O, Paquola C, Lariviere S, Royer J, Tavakol S, et al. BrainSpace:
a toolbox for the analysis of macroscale gradients in neuroimaging and connectomics
datasets. Commun Biol. 2020 Mar 5;3(1):1–10.

51. Bethlehem RAI, Paquola C, Seidlitz J, Ronan L, Bernhardt B, Consortium CC, et al.
Dispersion of functional gradients across the adult lifespan. NeuroImage. 2020 Nov
15;222:117299.

52. Koo TK, Li MY. A Guideline of Selecting and Reporting Intraclass Correlation Coefficients
for Reliability Research. J Chiropr Med. 2016 Jun;15(2):155–63.

Bignardi et al.

53. Rosseel Y. lavaan: An R Package for Structural Equation Modeling. J Stat Softw. 2012
May 24;48(1):1–36.

54. Bates TC, Maes H, Neale MC. umx: Twin and Path-Based Structural Equation Modeling in
R. Twin Res Hum Genet Off J Int Soc Twin Stud. 2019 Feb;22(1):27–41.

55. Boomsma D, Busjahn A, Peltonen L. Classical twin studies and beyond. Nat Rev Genet.
2002 Nov;3(11):872–82.

56. Verhulst B, Prom-Wormley E, Keller M, Medland S, Neale MC. Type I Error Rates and
Parameter Bias in Multivariate Behavioral Genetic Models. Behav Genet. 2019
Jan;49(1):99–111.

57. Mowinckel AM, Vidal-Piñeiro D. Visualization of Brain Statistics With R Packages ggseg
and ggseg3d. Adv Methods Pract Psychol Sci. 2020 Dec 1;3(4):466–83.

774