Distinct genetic underpinnings of inter-individual differences in the sensorimotor-association axis of cortical organisation

Giacomo Bignardi[1,2*], Michel G. Nivard[3,4], H. Lina Schaare[5,6], Boris C. Bernhardt[7], Richard A.I. Bethlehem[8,9], Simon E. Fisher[1,10,†], & Sofie L. Valk[5,6,11,† *]

[1] *Language and Genetics Department, Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands;*
[2] *Max Planck School of Cognition, Leipzig, Germany;*
[3] *Medical Research Council Integrative Epidemiology Unit, University of Bristol, Bristol, UK;*
[4] *Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK;*
[5] *Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany;*
[6] *Research Center Jülich, Institute for Neuroscience and Medicine (INM-7), Germany;*
[7] *McGill University, Montreal, Canada;*
[8] *Autism Research Centre, Department of Psychiatry, University of Cambridge, Cambridge, United Kingdom;*
[9] *Department of Psychology, University of Cambridge, Cambridge, United Kingdom;*
[10] *Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen 6500 HB, The Netherlands;*
[11] *Heinrich-Heine-Universitaet Duesseldorf, Germany.*

* Giacomo Bignardi
* Sofie L Valk.
† These authors share last authorship.
Email: giacomo.bignardi@mpi.nl; valk@cbs.mpg.de

Author Contributions: S.L.V. and G.B. conceptualised the work; S.L.V. & S.E.F. supervised the research; G.B. performed primary analyses; S.L.V., S.E.F., M.G.N., B.C.B., and R.A.I. gave input on analysis and figures; S.L.V. performed functional connectomics, microstructural and geodesic distance analysis; H.L.S. provided input in diffusion map embedding analysis; G.B. computed figures; G.B., & S.L.V. drafted the manuscript; S.L.V., S.E.F., M.G.N., B.C.B., H.L.S., and R.A.I. contributed to revising the manuscript.

Competing Interest Statement: The authors declare no competing interest.

Keywords: Neuroscience, Sensorimotor-Association axis, Behaviour Genetics, Structural Equation Modelling, Measurement Error.

Bignardi et al.

Abstract

In humans, many neurobiological features of the cortex—including gene expression patterns, microstructure, and functional connectivity—vary systematically along a sensorimotor-association (S-A) axis of brain organisation. To date, it is still poorly understood whether inter-individual differences in patterns of S-A axis capture these robust spatial relationships across neurobiological properties observed at the group-level. Here, we examine inter-individual differences in structural and functional properties of the S-A axis, namely cortical microstructure, geodesic distances, and the functional gradient, in a sample of young adults from the Human Connectome Project (N = 992, including 328 twins). We quantified heritable variation associated with inter-individual differences in the S-A axis, and assessed whether structural and functional properties that are highly spatially correlated at the group-level also share genetic underpinnings. To consider measurement errors in resting-state functional connectivity data and their impact on properties of the S-A axis, we used a multivariate twin design capable of disentangling individual-level variation in both intra- and inter-individual differences. After accounting for some of the intra-individual variation, we found average heritable individual differences in both the functional gradient (h_{twin}^2 = 57%), cortical microstructure (h_{twin}^2 = 43%), and geodesic distances (h_{twin}^2 = 34%). However, these genetic influences were mostly distinct and deviated from group-level patterns. In particular, we found no significant genetic correlation between the functional gradient and microstructure, while we found both positive and negative genetic associations between the functional gradient and geodesic distances. Our approach highlights the complexity of genetic contributions to brain organisation and may have potential implications for understanding cognitive variability within the S-A axis framework.

Introduction

The human brain supports perception and action but also abstract cognition (1,2). This diversity of functions is thought to be reflected by the gradual dissociation between unimodal sensory and transmodal association cortical areas along a sensorimotor-association (S-A) axis (3). The S-A axis spans a vast array of neuroanatomical properties, including microstructural variation (myelination and cytoarchitecture) and inter-areal connectivity distance (1,3–6). Here, sensory areas show increased layer differentiation, myelination, and, predominantly, short-range connections. In contrast, association areas show less differentiated microstructural profiles, reduced myelination, and a combination of short- and long-range connectivity profiles (5,7,8).

The differentiation between sensory and association areas underwent evolutionary changes (3), with an expansion of cortical association areas paralleled by a marked laminarisation of sensory areas in human primates (9,10). Such structural re-organisation and evolutionary changes along the S-A axis (1,6,11,12) may have provided the scaffold for functional differentiation (13,14), allowing in turn for human-specific cognitive and behavioural flexibility (1,13).

Several discoveries have enhanced our understanding of links between structural and functional features of the S-A axis. These findings have highlighted spatial associations of microarchitectonic differentiation (15) and cortical geometry (16) with functional organisation (13,17). For example, T1w/T2w maps derived from non-invasive Magnetic Resonance Imaging (MRI)—indexing cortical microstructural differences —and histological markers based on cell staining have been shown to relate strongly to gene transcriptional profiles and functional dissociation along the S-A axis. Such group-level associations suggest that a canonical genetic architecture may shape S-A axis structural organisation, providing foundations for the differentiation of cortical function (6).

Recent studies noted that various features of the S-A axis show inter-individual differences in human populations that are associated with variability in a host of traits, such as neuropsychiatric traits, including autism (18), schizophrenia (19), and depression (20), as well as sex (21) and developmental (22) differences. These findings highlight the overall importance of the S-A axis for human complex trait variation. However, it remains unclear how the different structural and functional properties of this axis relate to each other when viewed at the level of the individual, i.e. whether well-documented strong associations between such properties at the group-level (3,4,6,23–25) are also reflected in patterns of inter-individual differences. Knowing whether such patterns converge or diverge is crucial because it directly informs the study of the impact of alterations along the S-A axis on human traits. For instance, divergences between findings at the individual and group levels may suggest that associations between functional characteristics of the S-A axis and trait variability do not parallel analogous structural differences. Similarly, these differences may entail that a lack of associations between a specific neurobiological feature of the S-A axis and a particular trait does not necessarily imply the absence of relationships across other S-A axis neurobiological properties.

Here, we asked: *Do individual differences in structural properties of the S-A axis relate to differences in functional properties?* In other words, we studied whether previous widely reported group averages can inform S-A axis associations at the individual level. Specifically, we tested whether anatomical individual differences in regional cortical microstructure (6) and cortical geometry—captured by the geodesic distance of inter-connected regions across the cortical mantle (1,26)—relate to the well-known functional dissociations between sensory and transmodal association areas (1,3).

To thoroughly account for the known issue of measurement error heterogeneity across the cortex (27) and its impact on association estimates (28,29), we adapted and applied measurement error models in the form of structural equation models (30,31). This allowed us to rigorously tease apart unreliable intra-individual from reliable inter-individual variation in the functional organisation of the S-A axis.

We then went further to interrogate upstream sources of inter-individual differences in the S-A axis in
structure and function: genetic variation. We asked: *Are genetic effects on the S-A axis shared across*
structural and functional properties? Here, we analysed a genetically informative sample and quantified
the extent of overlap across genetic effects on structural and functional properties of the S-A axis.
Specifically, we used an augmented twin-informed design to quantify and tease out genetic overlaps
between the S-A axis's structural and functional properties while accounting for the impact of reliability
on classic heritability estimates (31,32). Last, we evaluated the robustness of our results, both across
subsamples, between regional and global cortical metrics, and between and within individuals' S-A axis
properties.
119

120 **Results**

121 To quantify structural and functional S-A axis properties, we combined microstructural and resting-state
122 functional MRI (rsfMRI) data from the Human Connectome Project (HCP (33); N = 992 adults; 529
123 women, mean age 28 y; 22-37 y). We computed two structural metrics and one functional metric
124 indexing the S-A axis:
125

• Regional microstructure: we quantified regional microstructure indexing the differentiation
between sensorimotor and association areas using the individuals' mean intensity of regional
T1w/T2w (T1w/T2w$_{mi}$) in 400 parcels (6,34)
• Geodesic distance: we quantified cortical geometry in each individual as regional cortico-
cortical network proximity (26) by computing the geodesic distance (GD) between every cortical
region and its corresponding functional network, averaging within each region to get parcel-
wise estimates (18). We chose geodesic distances because they align with the anatomical
layout of axonal wiring along the cortical sheet and are strongly associated with functional
connectivity gradients (1,35), thus providing a spatial framework that complements
microstructural measures
• Functional gradient loadings: we quantified the functional S-A axis in each individual by
obtaining the first component of the individual functional connectomes (FC$_{G1}$) using diffusion
map embedding (1)

We started our analysis by testing whether associations between group-level averaged maps of
structural S-A axis properties correlated with the functional S-A axis (Fig. 1A-C). By using a subsample
of n = 482 adults (229 women, mean age 28 y; 22-37 y; a subsample obtained by excluding all twins
included in the full HCP sample), we were able to replicate group-level findings between averaged
T1w/T2w$_{mi}$ and FC$_{G1}$, extending the results to GD and FC$_{G1}$ (Fig. 1D-E).

A   Microstructural intensity   B   Geodesic distances   C   Functional gradient

D   Group-level associations   E   Group level-analysis

144
Figure 1. *Structural and functional S-A axes strongly correlate at the group-level*. Structural (**A-**
B) and functional (**C**) indices of Sensorimotor-Association (S-A) axes plotted on inflated cortical
surfaces (36). Values represent averages of individual T1w/T2w mean intensity profiles (A; T1w/T2w$_{mi}$),
averages of individual geodesic distances (B; GD), and functional gradients loadings (C; FC$_{G1}$)
extracted from the average of individual functional connectomes across 400 cortical regions. (**D**)
Structural indices are strongly associated with functional indices of the S-A axis; Spearman ρ = -.61
and ρ = .75 between T1w/T2w$_{mi}$ and FC$_{G1}$, and GD and FC$_{G1}$, respectively; all p < .05. Each dot
represents a regional value; the colour represents canonical Yeo-Krienen 7 network membership. (**E**)
Conceptual representation of group-level analysis. Note that individual and regional information is lost
in favour of group-level results.
155

156   **Pervasive inter-individual differences in the S-A axis of cortical organisation.** Having estimated
157   the extent of overlap between structural and functional S-A axis properties at the group-level, we shifted
158   the focus to the individual level. Since group-level S-A axis can mask substantial individual variability
159   (Fig. 2A), we asked: does individual variability in structural S-A axis properties relate to variability in S-
160   A functional properties, as for group-level analysis (Fig. 2B)?
161

In shifting analysis from group-level summary statistics to individual variability, we harnessed the
distinction between intra- and inter-individual differences (37). The first (i.e., intra-individual) is known
to index unreliable and fluctuating variability within individuals over time, while the second (i.e., inter-
individual) indexes the reliable and stable part of the overall variability between individuals (Fig. 2C)
(37). This distinction is crucial, as intra-individual variability can downward bias effect sizes and reduce
statistical power (37), additionally downward biasing genetic estimates (38). Such factors apply
heterogeneously across the whole cortex (27), and can, therefore, increase reproducibility issues (see
(28) for details).
170

We were able to robustly distinguish between intra- and inter-individual effects by exploiting one of the
strengths of the HCP design, which emphasises multiple rsfMRI sessions (across two days of scanning
sessions, ~30 min each). This feature of the HCP design allowed us to partly discard intra-individual
fluctuations in rsfMRI data from inter-individual differences in the functional gradient. Precisely, we
partition the inter-individual variance ($\sigma_{inter(i)}^2$) from the overall observed variance in the functional
gradient ($\sigma FC_{G1(i)}^2$ for any parcel i) by applying a measurement error model ((30,31) Fig. 2D, see
Methods).

Estimates obtained from the measurement error model indicate that 33% of the total variability in the
functional gradient was, on average, accounted for by intra-individual variance (Fig. 2D) even when
using individual functional gradients extracted from functional connectomes averaged across two days
of rsfMRI sessions (totalling ~60 min of scanning session). In other words, estimates for the association
between the functional gradient and other S-A axis properties (or any other variable) would be, on
average, biased downward by a factor of bias(r-observed, r-true) = .82 (a lower bound calculated
assuming perfect reliability for the other S-A axis property (28)). Second, we observed systematic
differences in estimates obtained across functional cortical networks, $F(6, 393) = 33.21$, $p < .001$; $\eta^2 =$
0.34, 95% CI [0.27, 1.00]), with estimates for parcel-wise inter-individual variances ranging from
$\sigma_{inter(114)}^2 = .39$ to $\sigma_{inter(294)}^2 = .89$ (Fig. 2D). That is, bias is heterogeneous and expected to influence
estimates across the cortex systematically.
189

190
Figure 2. *Pervasive inter-individual differences in the S-A axis of functional connectivity.* (**A**)
group-level estimates (black contour) overshadow pervasive individual differences in S-A axis
properties. (**B**) The shift between levels of analyses: from group-level (grey square) to between-
individuals (coloured squares); the gradient square conceptually captures panel A. (**C**) Measurement
error model to partition, for any parcel i, variance in the functional gradient loadings into intra- ($\sigma_{d\text{-}intra(i)}^2$,
for regional values measured at day 1 or 2 of the testing session, i.e., rectangles) and inter- ($\sigma_{inter(i)}^2$, for
the latent component, i.e., circle) individual variance. Parameter estimates for any parcel i can be found
in Supplementary Table S1. (**D**) The proportion of intra- and inter-individual variance in the functional
network across Yeo-Krienen functional networks: the horizontal dashed line represents the mean
proportion of variance across networks; the horizontal lines display the median within network; lower
and upper hinges correspond to the first and third quartile; the whisker extends from the hinge to the
largest/lower value no further than 1.5 * interquartile range from the hinge. Note that across all parcels,
observed variance includes substantial inter-individual variation. *Notes on measurement model:*
Rectangles represent the measured phenotypes; the circle is the latent component; the double-headed
arrows within the circle represent the variance associated with the latent components; one-headed
arrows are the paths (here all set to 1).
207

Individual differences in regional cortico-cortical network proximity, rather than microstructure,
relate to the functional gradient of the S-A axis of cortical organisation. To simultaneously
deattenuate the heterogeneous downward biases and handle structural and functional S-A metrics, we
used a Structural Equation Modelling (SEM) approach. Precisely, we specified a model in which the
inter-individual differences in the functional gradient estimated via the measurement error model were
directly tested for associations with microstructural profiles and geodesic distances parcel-wise data

(see Methods). Here, we note that we avoided making assumptions about the causal structure
generating the possible correlations between structural and functional metrics. We simply limited
ourselves to estimating the association between regional properties of the S-A axis.

On the one hand, contrary to group-level topographies, we found less than 2% of the 400 parcels to
display a significant association between individuals' microstructural profiles and functional gradient
loadings. These significant associations were all negative, weak ($-.20 > r > -.27$), and spread across
both hemispheres and the dorsal, ventral, and default-mode functional networks. Conversely, we found
large overlaps between individual geodesic distances and functional gradient loadings (Fig. 3A), with
57% of the 400 parcels showing significant associations after Bonferroni correction. The directionality
of the estimates for the association between individual regional geodesic distances and functional
gradient loadings highlighted systematic differences across functional networks. Significant positive
associations were preferentially clustered within the visual and the default mode (one sample t-test,
two-sided, $t(21) = 5.82$, $p < .001$, average $r = .51$, and $t(53) = 4.04$, $p = .001$, average $r = .25$), while
negative associations were preferentially clustered within the somatomotor and ventral attention
networks (one sample t-test, two-sided, $t(59) = -13.32$, $p < .001$, average $r = -.48$, and $t(27) = -8.14$, p
$< .001$, average $r = -.40$, respectively, all tests accounting for multiple-testing via Bonferroni correction).
Estimates obtained from standard correlation analysis further confirmed that the SEM approach
successfully deattenuated measurement error bias (Fig. 3B).

Figure 3. *Structural and functional S-A axes selectively correlate between individuals.* (**A**)
Summary for the standardised estimates on the inflated cortical surface (36) from the structural
measurement error model (see Methods) indicates little and weak phenotypic correlations (r_p) between
microstructural intensity (T1w/T2w$_{mi}$) and functional gradient loadings (FC$_{G1}$) inter-individual differences
but large and highly significant ($p<.05$ after Bonferroni multiple-testing correction) overlaps between
functional gradient loadings and geodesic distances. All parameter estimates for any parcel i, including
the covariances between T1w/T2w$_{mi}$ and GD that were not the focus of the current study, can be found
in Supplementary Table S2. (**B**) The scatter plot shows how correlations between regional functional
gradient loadings and geodesic distance estimated following the classic Pearson correlation approach
(r_{gd-g1}, on the x-axis) relate to correlations estimated with the measurement error model approach
(deattenuated r_{gd-g1}, on the y-axis). The grey line represents the deviation from the expected relationship
between the two approaches under no estimated difference. As can be seen, negative and positive
downward biases tend to be deattenuated.

Genetic influences on inter-individual differences in the functional gradient of cortical
organisation. We went on to assess the extent of genetic influences on the S-A axis and the degree
to which their estimates are impacted by measurement error. To do so, we exploited the family structure
in the HCP to partition individual differences. In particular, we applied a twin design and partitioned the
variability of functional gradient loadings into genetic (σ_A^2; A: additive) and unsystematic environmental
(σ_E^2; E: unique-environmental) sources via SEM. We focused our analysis on the twin HCP subsample
including both monozygotic (MZ) and dizygotic (DZ) twins ($n = 328$, 195 MZ and 133 DZ individual twins,
124 and 88 women, respectively; mean age 29 y, range = 22-35 y; see Methods for details on inclusion
criteria), and derived twin-based heritability estimates (h_{twin}^2). These h_{twin}^2 estimates were derived from

an integrated SEM that incorporated the measurement error model. This approach made it possible to
separate intra- and inter-individual differences in the functional gradient, yielding h_{twin}^2 estimates that
robustly accounted for intra-individual variation, including measurement error.

We benchmarked our h_{twin}^2 estimates for functional gradient loadings (obtained from the model that
accounted for measurement error) by comparing them to a classic twin model (not accounting for
measurement error). Following standard cut-offs (31), we retained 395 and 398 parcel-wise
measurement error and classic twin models, respectively, with a satisfactory CFI > .90 and an RMSEA
< .08. We excluded an additional parcel-wise measurement error twin model as it returned an out-of-
bounds estimate of h_{twin}^2 > 1.The inclusion of the measurement error model substantially boosted h_{twin}^2
estimates relative to the estimates obtained from models not accounting for intra-individual variance. In
particular, accounting for intra-individual differences resulted in a much larger average heritability for
functional gradient loadings: h_{twin}^2 = .57, SD = 0.13, compared to the h_{twin}^2 = .37, SD = 0.11 obtained
from classic models (Fig. 4A-B). Fig. 4C provides an illustration of the enhanced estimates for the
heritability of S-A axis functional organisation when properly accounting for measurement error.

274
Figure 4. *Univariate twin AE models that account for measurement error boost functional*
***gradient heritability estimation*.** Parcel-wise twin heritability (h_{twin}^2) estimates for the 392 parcels with
satisfactory fit indices across classic and measurement error twin models. **A** Box plot of the h_{twin}^2
stratified per Yeo-Krienen 7 functional networks. **B** Box plot of the h_{twin}^2 stratified per Yeo-Krienen 7
functional networks obtained from the model accounting for measurement error (here, intra-individual
differences, see Methods). The dashed line displays the mean h_{twin}^2 across networks. Note that the
average heritability is 53% higher in B. Parameter estimates for any parcel *i* can be found in
Supplementary Table 3. **C** Scatter plot showing the increase of the h_{twin}^2 estimate across the cortex in
both models. Each dot represents one parcel.

Genetic effects on different properties of regional S-A axis variability are substantial yet mostly
distinct. After having related inter-individual differences in structural and functional S-A axis properties,
and showing heritable variation in functional organisation of the S-A axis, we asked whether genetic
effects were mostly common or distinct across S-A axis properties. To partition variability (σ_p^2; p: S-A
axis phenotypic property) within S-A modalities in σ_A^2 and σ_E^2 sources, and further unpack genetic and
environmental structure-function associations, we specified a multigroup multivariate model with only A
and E components (see Methods). We retained all multivariate models as they displayed satisfactory
fit indices, except a model returning an out-of-bounds estimate of h_{twin}^2 > 1. Similar to functional gradient
loadings, microstructural profiles, mean h_{twin}^2 = .43, SD = 0.11, and geodesic distances, mean h_{twin}^2 =
.34, SD = 0.11, displayed substantial h_{twin}^2 across the cortex (Fig. 5A-B).

Notwithstanding such relatively high h_{twin}^2 for both microstructural profiles and functional gradient
loadings, we found no significant additive genetic correlation between the two (all *p* > .05, Bonferroni
corrected; Fig. 5B). This suggested little room for possible shared genetic causes between the S-A axis
properties indexed by microstructural intensity and functional gradient loadings. Conversely, 14% of the
parcels displayed significant additive genetic correlations between geodesic distances and functional

gradient loadings (7% negative and 7% positive in directionality, $p < .05$, Bonferroni corrected, Fig. 5C).
The average magnitude of the genetic correlation (r_A) was $r_A = -.67$, $SD = 0.16$, and $r_A = .64$, $SD = 0.16$.
Furthermore, we found that for 30% of the parcels, complementary environmental effects mostly
correlated between geodesic distances and functional gradient loadings (Fig. 5D).
305

306
Figure 5. *Genetic sources of structural and functional properties of the S-A axis are selectively*
distinct. (**A**) Parcel-wise twin heritability (h_{twin}^2) estimates for microstructure and (**B**) geodesic
distances. (**C**) Summary for the significant additive genetic correlations r_A and (**D**) and environmental
correlations (r_E) across structural S-A axis properties with the functional gradient loadings on the inflated
cortical surface (46). Note that the only significant genetic correlations are between geodesic distances
and functional gradients loadings. The r_A estimates can be found in Supplementary Tables S4-S5.
313

Associations between structural and functional properties of the S-A axis are robust across
samples. To test for the robustness of the results discussed so far, we estimated the overlap of the
significant regional genetic or environmental associations in the genetically informative subsample with
the significant regional phenotypic associations obtained from the first subsample. Of the 104 parcels
that displayed significant genetic and/or environmental correlations between geodesic distances and
functional gradient loadings in the genetically informative sample, 99 also displayed a significant
phenotypic correlation in the first subsample. In other words, we found a 95% overlap between
subsamples in terms of the parcels implicated. These results show that regional results were robust
across two subsamples drawn from the HCP.
323

Genetic and environmental associations extend beyond regional S-A axis variability. As a final
analysis, we asked whether associations between geodesic distances and functional gradients were
generalisable beyond regional differences. First, we quantified variability in global S-A axis properties
as the overall Median Absolute Deviation (MAD) across all parcels within individuals. Within an
individual, higher MAD scores indicate a larger dispersion in S-A axis values across the cortex. Once
more, we found no significant genetic or environmental associations between microstructural profile
intensity and functional gradients MAD scores. Yet, we found a substantial negative genetic correlation
between the geodesic distances and functional gradient MAD scores ($r_A = -.78$, 95% CI [-1.19, -.34],
CFI = .93, RMSEA = .04; Fig. 6A).
333

Additionally, to obtain a complementary estimate of global S-A axis variability, we quantified
microstructural profile intensity, geodesic distance, and functional gradient ρ similarity indices. These

indices assessed how similar S-A axis properties in one individual are compared to the average.
Consistent with regional and global variance differences, ρ similarity indices in geodesic distance, rather
than microstructure, showed strong and positive genetic correlations with global differences in the
functional gradient (r_A = .61, 95% CI [.43, .79], CFI = .95, RMSEA = .04; Fig. 6B). Findings were robust
to intra-cranial volume as a possible common cause of S-A axis structure-function covariance (Fig.6C-
D)
342

343
Figure 6. *Findings extend beyond regional S-A axis associations.* **(A)** Simplified graphical
representation of the multivariate twin-informed SEM for overall within-individual Median Absolute
Deviations (MAD). Note the strong but negative significant associations between the latent additive
genetic components (*A*, circles in red) underlying geodesic distances (GD; centre) and functional
gradient loadings (FC$_{G1}$; right). The blue circles represent the latent residual environmental components
(E). **(B)** Simplified graphical representation of the multivariate twin-informed SEM for ρ similarity indices
(Fisher-*z* transformed). As for the model reported in panel A, the only significant associations are found
between GD and FC$_{G1}$. * *p* < .05. **(C)** and **(D)** panels show that the associations between MAD scores
and ρ similarity indices are robust to intra-cranial volume (ICV) as a possible confounder. Here, double-
headed arrows between latent variables indicate correlations (since we report standardised solutions).
Dashed arrows represent nonsignificant correlations.
355

356 **Discussion**
357 Our results reveal that group-level estimates of spatial associations between structural and functional
358 gradients differentiating sensory from association areas in the human brain might mask pervasive inter-
359 individual differences. These inter-individual differences, in turn, might display different patterns of
360 associations from those depicted at the group level.
361

Notwithstanding the comparable heritability of the different S-A axis properties as shown in this study,
theoretical models of S-A axis development and evolution (3), and group-level relationships between
patterns of gene expression, cortical microstructure, and functional differentiation of sensorimotor to
transmodal-association areas (6), our findings revealed minimal convergence between microstructural
and functional features of the S-A axis at the individual level. Specifically, we found little evidence of
phenotypic and an absence of evidence for genetic associations between cortical microstructure (as
measured by T1w/T2w) and S-A function (as measured by the principal gradient of functional
connectivity) of the cortex.

At the same time, our results showed substantial genetic and environmental associations between
individual-level differences in cortico-cortical network proximity (as measured by the geodesic distance
of inter-connected regions across the cortical mantle) and S-A function. These latter results align with
theories emphasising geometric constraints of brain function (1), yet do not fully align with group-level
estimates. While group-level associations indicate a positive relationship between cortico-cortical
network proximity, our results uncover a mixture of positive and negative relationships at the individual
level of analysis (the former preferentially clustered within the visual and default mode network, the
latter with the somatomotor and the ventral attention network).

Moreover, we found negative, not positive, genetic correlations when shifting from local to global
association, as we did when analysing overall within-individual S-A axis dispersion. This suggests that
genetic differences between people that tend to co-occur with decreased variation in geodesic distances
across the cortex also tend to co-occur with more dispersed functional gradients. It is worth noting that
the stronger relationship between geodesic distances and functional gradients (compared to
microstructure) likely reflects the spatial constraints imposed by cortical geometry on functional
organisation. While geodesic distances capture anatomical spatial relationships and functional
gradients reflect connectivity patterns, both are shaped by the underlying architecture of neural
connections. In particular, short-range anatomical proximity supports stronger functional coupling,
which may contribute to the observed robust association. The selective associations between geodesic
distances and functional gradients at the individual level also support the observed dissociation in S-A
axis with microstructure, rather than reflecting a broader issue with finer spatial features of gradient-
based approaches.

Fundamental principles of brain organisation can appear to be highly conserved across neurobiological
properties if based on group-based analyses of topological co-variation, yet our results suggest this
conservation may break down when assessed at the level of the individual. Based on group-level
associations, previous work suggested that cortical maturation of diverse neurobiological properties
proceeds along an evolutionary conserved and developmentally rooted S-A axis of cortical organisation
(3,4). However, our results indicate that genetic variation within a population is expected to be
selectively associated with some properties (e.g., function and cortico-cortical network proximity) but
not others (e.g., microstructure), at least within cortical regions. The apparent paradox of observing
stable group-level patterns and individual variation fits with current models of brain organisation, which
recognise both shared principles and meaningful personal differences. This view is in line with recent
perspectives (e.g., 39), which emphasise the importance of individual-specific brain features.

These findings may also be important for understanding the origins of differences between individuals,
such as in various neuropsychiatric disorders or work on brain-behaviour associations in general. First,
since we found a dissociation between structural and functional gradients, complementing studies of S-
A axis variability with both structural and functional features may result in the discovery of brain-
behaviour associations with selective neurobiological properties of this axis. Second, the addition of
informative genetic models of structural and functional S-A axis variability would provide several novel
insights into previously observed correlations. Along these lines, by disentangling brain-behaviour
associations into their genetic and environmental sources, genetically informative models could also

enhance our understanding of whether previously observed associations are caused by non-genetic
factors. We also note that our model can be further applied to many neurobiological properties (e.g.,
rsfMRI, see (31)) to enhance current brain-behaviour mapping efforts (40). To facilitate this, we have
made all the code available and provided all SEM functions in R and lavaan syntax.

We also underscore that the measurement error modelling approach can successfully tease apart
unstable intra-individual differences from stable inter-individual differences, and this effect can have a
substantial downstream impact on estimates. For example, applying the measurement error modelling
approach, in line with previous results (31), resulted in a nearly 50% increase in estimates of heritability.
These results should also work as a cautionary tale against interpreting differences in average
heritability across principles of S-A organisation, or, more generally, between functional and structural
properties of the cortex. Before accounting for intra-individual differences that occur also as a result of
measurement error, heritability estimates for functional gradient loadings may have seemed lower than
heritability estimates for structure. Yet, after accounting for such intra-individual variation, estimates
were higher. This apparent puzzle is reconciled due to heritability being a ratio, with intra-individual
variation being part of the denominator. Since functional connectivity tends to vary intra-individually,
estimates will tend to be smaller when not accounting for this intra-individual variability.

We foresee that the measurement error modelling approach could have further direct application in
ongoing research on the origins of psychiatric disorders and brain-behaviour studies, and in the analysis
of the genomic architecture of principles of brain organisation, exactly because we show that it may
mitigate the impact of measurement error heterogeneity on estimates. Indeed, when individual
variability in the S-A axis is the predictor of interest, such as in brain-behaviour studies, applying any
measurement error model is expected to deattenuate downwardly biased estimates (28,37,41,42).
Moreover, genome-wide association studies could easily implement genome-based structural equation
modelling (43,44) extensions of our approach to discard unstable and unreliable variance, overcoming
attenuation biases in associations between single nucleotide polymorphisms and target phenotypes
(e.g., similarly to what has been done for analyses based on polygenic indices (42)).

Yet, it is important to note that the interpretation of deattenuated estimates rests upon assumptions of
what type of error is expected to influence the measurement of functional S-A axis properties across
days of measurement. Here, we assumed that what is measured as being shared between sessions is
the common cause of what is measured within each session. Within this framework, what is left is
unique to each session, including sources of measurement error. Such application of the measurement
error model to S-A axis functional properties is not expected to perfectly segregate S-A axis error-free
variance in the inter-individual component (37). For example, the systematic error of individual S-A axis
measurement would be indistinguishable from meaningful inter-individual variability and still be
captured by the inter-individual variance component. Additionally, this measurement error model could
confound unsystematic S-A axis error with genuine intra-individual differences, which may even be
partially heritable (45). Therefore, the actual sources of session-to-session differences may be more
nuanced than the simple measurement error model implies. Consequently, we advise caution when
interpreting inter- and intra-individual differences as exclusively "error-free" and "error" variance,
respectively (see Methods for further details).

Another limitation of our study is the lack of repeated structural metric measures. Although applying a
measurement error model allowed us to partly disentangle intra- and inter-individual variability in
functional gradient loadings, we could not account for the differences in structural properties within
individuals. This limitation may have attenuated the estimated relationship between structure and
function. However, the nature of the metrics and the twin design employed to elucidate differences
between individuals should mitigate the impact of such a lack of repeated structural metrics (38). By
applying the classical twin design, we were able to further partition unstable measurement error in the

environmental (*E*) component of the model, which minimised possible biases introduced by hypothetical
measurement error, at least for the additive genetic (*A*) correlations (r_A) estimates.
467
In sum, our findings reveal that group-level results can overshadow substantial inter-individual
differences within and between different neurobiological properties. By focusing on these previously
underappreciated differences, we could highlight selective associations of individual variation in S-A
axis cortical structure and function. These inter-individual differences and associations open a window
into genetic sources of S-A axis structure and function, which we reveal to be selectively distinct. Our
results underscore the complex interplay between the S-A axis's structural and intrinsic functional
properties and provide a set of tools that can be used to test their potentially differential roles in shaping
cognition.
476

Materials and Methods
Sample. We used data from the Human Connectome Project (HCP) S1200 release. The HCP includes
data from 1206 individuals (656 women) that comprise 298 Monozygotic (MZ) twins, 188 Dizygotic (DZ)
twins, and 720 individuals, with mean age ± *SD* = 28.8 ± 3.7 years (age range = 22-37 years). Informed
consent for all individuals was obtained by HCP, and our data usage was approved by HCP and
complied with all relevant ethical regulations for working with human participants (see (13,33,46)). The
primary participant pool comes from individuals born in Missouri to families that include twins, sampled
as healthy representatives of ethnic and socioeconomic diversity of US individuals, based on data from
the Missouri Department of Health and Senior Services Bureau of Vital Records. We followed standard
guidelines for inclusion criteria as described elsewhere (13). Our sample, in line with Valk et al., (13)
comprised 992 (529 women) individuals. The first subsample of *n* = 482 (229 women) was created by
excluding all twins. The second genetically informative subsample of *n* = 328 (212 women) was created
by including only twins with genotyped zygosity matching self-reported zygosity (195 MZ and 133 DZ
individuals; 124 women and 88 men, respectively, forming between 150 and 152 complete pairs,
depending on data availability for the S-A axis modality).
492

Functional imaging. Functional connectivity matrices were based on four 14 min 33 s of functional
Magnetic Resonance Imaging (fMRI) data acquired over two sessions, spaced two days apart, through
the HCP, which underwent HCP's minimal preprocessing. No global signal regression was performed
in the processing of the fMRI data. For each individual, four functional connectivity matrices were
computed using the minimally preprocessed, spatially normalised resting-state fMRI (rsfMRI) scans,
which were co-registered using MSMAll to template HCP 32k_LR surface space. 32k_LR surface space
consists of 32,492 total nodes per hemisphere (59,412 excluding the medial wall). We computed four
functional connectivity matrices per individual from the average time series extracted in each of the 400
Schaefer cortical parcels. The individual functional connectomes were generated by averaging
preprocessed time series within nodes, Pearson correlating nodal time series and converting them to
Fisher-*z* scores. The average functional connectomes were obtained by averaging functional
connectomes within individuals (i.e., between sessions) and between individuals.

Structural imaging. MRI protocols of the HCP have been previously described (33,46). MRI data were
acquired originally on the same day on the HCP's custom 3T Siemens Skyra equipped with a 32-
channel head coil. T1w images with identical parameters were acquired using a 3D-MP-RAGE
sequence over 7 min 40 s (0.7 mm isovoxels, matrix = 320 × 320, 256 sagittal slices; TR = 2400 ms,
TE = 2.14 ms, TI = 1000 ms, flip angle = 8°; iPAT = 2). T2w images were acquired using a 3D T2-
SPACE sequence with identical geometry over 8 min and 24 s (TR = 3200 ms, TE = 565 ms, variable
flip angle; iPAT = 2). We followed the preprocessing steps outlined in Valk et al. (13).

Parcellation and functional networks. We used the Schaefer group-level hard-parcellation, originally
obtained by a gradient-weighted Markov random field model integrating local gradient and global
similarity approaches (34). To stratify results within canonical cortical functionally coupled networks, we
used the seven Yeo-Krienen networks (47).

Microstructural profiles (T1w/T2w$_{mi}$). We used T1w/T2w imaging myelin-sensitive contrast from the
HCP minimal processing pipeline, which uses the T2w to correct for inhomogeneities in the T1w image
to estimate mean intensity T1w/T2w microstructural profiles (T1w/T2w$_{mi}$). T1w/T2w$_{mi}$ has been shown
to map to model-based tract-tracing histological data in macaque, estimate intracortical myelin content,
and thus approximate architectural complexity and cortical hierarchy (6).

Geodesic distance (GD). Individual geodesic distances (GD) were computed using the Micapipe
toolbox (26). Briefly, we computed GD between each region and their top 10% of maximally functionally
connected regions along each individual native cortical midsurface. We further averaged within regions
to obtain a parcel-wise value and improve computation performance. Micapipe implements the Dijkstra
algorithm (48) (further details can be found in (26)).

Functional gradient loadings (FC$_{G1}$). We sequentially averaged FCs, first within days, resulting in two
FCs per individual, and then between days, resulting in one FC per individual. We then extracted the
three first components from the two sequentially averaged and one averaged FCs, using the Python
package BrainSpace (49). Extraction of the first eigenvector followed standard procedures, with the
original individual FCs set at a connection density of 10% (i.e., the FCs were made sparse by setting a
sparsity threshold of 90%). The first ten eigenvectors were then obtained by decomposing the FCs by
diffusion map embedding, a robust non-linear manifold learning technique (1). To aid comparability
across individuals, we aligned individual eigenvectors to the template eigenvector by Procrustes
rotation (50). The template functional gradient was directly extracted from the overall mean FC matrix.

Group-level associations analysis. We computed Spearman rank-order correlations (ρ) between the
structural (T1w/T2w$_{mi}$ and GD) and functional (FC$_{G1}$) S-A axis group-level properties. Group-level
properties were obtained from the average of the individual structural S-A properties (i.e., average
T1w/T2w$_{mi}$ and GD), and from the decomposition of the average FC (i.e., principal gradient obtained
via diffusion map embedding of the average FC).

Measurement model of error in individual variability of the functional S-A axis. To partition stable
inter-individual variability in functional gradient loading, we adapted previous measurement error
models to rsfMRI to the functional gradient (30,31). The intuition behind such a modelling strategy is
simple. Suppose parcel-wise values are measured without error and are stable over a reasonable
period of time (e.g., one day). In that case, the correlations across individuals between the values
obtained across two time points will equal 1. If the correlations deviate from 1 instead, regional values
will be measured with some error, with bigger deviations corresponding to higher error or fluctuation
over time. As can be seen in Fig. 2C, when errors or changes over time are present, it may become
difficult to distinguish regional differences between individuals from regional differences within
individuals. In this case, we can use the measurement error model to estimate what stays constant
across time, indexing the "true" regional values. Across the manuscript, for correctness, since "error"
variance can include meaningful, yet unstable, fluctuation in rsfMRI, while "true" variance can also
consist of systematic measurement error across sessions, we refer to the former term as intra-individual
and the latter as inter-individual variability (37). First, we fit a measurement model to parcel-wise
functional gradient loadings averaged within days. In line with Teeuw et al. (31), we did not constrain
intra-individual variance components to be equal across days of scanning sessions. We performed
model fitting in lavaan (44) after standardising observed variables (i.e., std.ov = T). We then used model
estimates obtained for the variances of the latent and observed components. Using Spearman-Brown
correction, we computed the averaged proportion of stable inter-individual variance in functional
gradient loadings across days as the intra-class correlation (ICC) (51). For each parcel *i,* the ICC was
calculated as follows:

561
$$\text{ICC}(2,k)_i = \frac{k * \text{ICC}(2,1)_i}{1 + (k-1) * \text{ICC}(2,1)_i} \tag{1}$$

562 Where k is a constant equal to the number of measures (i.e., $k = 2$) and the ICC(2,1)$_i$ is calculated as
563 follows:

$$\text{ICC}(2,1)_i = \frac{\sigma^2_{\text{inter}(i)}}{\sigma^2_{\text{inter}(i)} + \left(\frac{\sigma^2_{1-\text{intra}(i)} + \sigma^2_{2-\text{intra}(i)}}{2}\right)} \tag{2}$$

565 ICC(2,k)$_i$ estimates the proportion of inter-individual variance over the total variance, $\sigma_{\text{inter}(i)}^2$, in the
566 functional loadings as if they were obtained from the average of the two scanning sessions. The
567 proportion of intra-individual variance for a parcel i, $\sigma_{\text{intra}(i)}^2$, is obtained simply by subtracting the
568 ICC(2,k)$_i$ from 1.

569 The expected bias for any parcel i was calculated following Tiego et al. (28):

$$\text{bias}_{(i)} = \sqrt{R_{p,p} * R_{\text{FCG1,FCG1}(i)}} \tag{3}$$

Where $R_{p,p}$, the reliability for the structural S-A axis property p (e.g., T1w/T2w$_{mi}$), was set to be equal
to 1 across all parcels, and $R_{\text{FCG1,FCG1}(i)}$, the reliability of parcel-wise value for the functional gradient
loading, was calculated as ICC(2,k)$_i$.

Multivariate Measurement error SEM. We used Structural Equation Modelling (SEM) to estimate
correlations between structural (i.e., T1w/T2w$_{mi(i)}$, GD$_{(i)}$) and functional (i.e, FC$_{G1(i)}$) S-A modalities.
Each multivariate model simultaneously accounted for intra-individual variances by including the
measurement error model (Fig. 7A). All models were fitted in lavaan (52) after standardising all
observed variables (i.e., std.ov = T). Prior to model fitting, sex and age were regressed from parcel-
wise S-A axis values using the function umx::umx_residualize() (53). Structural equation models were
fit to residual scores. We assumed missing data to be missed at random and followed parameters'
estimation via full-information Maximum Likelihood (i.e., missing = "ML").

Twin-informed Multivariate Structural Equation Modelling. We used multigroup SEM to partition
parcel-wise variability in structural (σT1w/T2w$_{mi(i)}^2$, σGD$_{(i)}^2$) and functional ($\sigma_{\text{inter}(i)}^2$) S-A modalities into
either additive genetic (σ_A^2) or unsystematic environmental (σ_E^2) sources of variance. Structural
equation models were fit to T1w/T2w$_{mii}$, GD$_i$, and FC$_{G1i1}$ (day 1) and FC$_{G1i2}$ (day 2) data, grouped by
zygosity (i.e., two groups). The model specification was informed by the multivariate twin design (54).
Briefly, monozygotic (MZ) twins are ~100 % genetically identical, coming from the same fertilised egg.
In contrast, dizygotic (DZ) twins are, on average, only 50% additively genetically similar regarding allelic
variants coming from two different fertilised eggs. Thus, the correlation f between the additive genetic
component (A) is set to be equal to 1 for MZ and ½ for DZ. In contrast, each twin's unique environment
(E) component will be unique; therefore, their correlation will be equal to 0. In total, we fit one
multivariate AE model per parcel. Following the measurement error procedure outlined above and in
reference (31), a common pathway measurement error model was included in the specification of the
multigroup SEM (Fig. 7B). As such, each multivariate model simultaneously accounted for intra-
individual variance (Fig. 7C). We employed the direct symmetric approach by estimating variance
components directly while setting path coefficients to 1 (with the exception of the measurement model,
for which we fixed the variance to be equal to 1, and estimated the path coefficients, instead). We chose
this approach as it has been shown to reduce type I errors and produce asymptotically unbiased χ^2
(55).

Similarly to reference (56), twin models were fitted in lavaan (52), with standardisation of observed
variables before model fitting (i.e., std.ov = T). To control for the effect of age and sex on S-A axis
properties, we residualised parcel-wise variables prior to modelling using the function
umx::umx_residualize() (53). Residuals were used as observed variables in later twin modelling. We
estimated parameters via full-information Maximum Likelihood (i.e., missing = "ML") and evaluated the
goodness of fit for each parcel by comparative fit index (CFI) and root mean square error of

approximation (RMSEA) scores. Following standard cut-offs (31), we retained only models with a
"satisfactory" CFI>.90 and an RMSEA<.08. Narrow-sense twin heritability (h_{twin}^2) estimates for each
parcel i were defined as the ratio of the additive genetic variance over the sum of the additive genetic
and environmental variances:

610
$$h_{\text{twin}-p(i)}^2 = \frac{\sigma_{Ap(i)}^2}{\sigma_{Ap(i)}^2 + \sigma_{Ep(i)}^2} \tag{4}$$

611 Where $\sigma_{Ap(i)}^2$ is the additive genetic variance for the S-A axis parcel-wise value for the given property p
612 (e.g., GD). For functional gradient loadings, the heritability was calculated as

613
$$h_{\text{twin}-\text{inter}(i)}^2 = \sigma_{A\text{inter}(i)}^2 \tag{5}$$

614 After imposing the equality constraint on the common factor $FC_{G1\text{inter}(i)}$

615
$$\sigma_{A\text{inter}(i)}^2 + \sigma_{E\text{inter}(i)}^2 = 1 \tag{6}$$

616 Genetic correlations (r_A) were calculated as:

617
$$r_a = \frac{\sigma_{Ap1i,Ap2i}}{\sqrt{\sigma_{Ap1i}^2 * \sigma_{Ap2i}^2}} \tag{7}$$

618 Where $\sigma_{Ap1,Ap2}^2$ is the additive genetic covariance between two S-A axis properties, $p1$ and $p2$ (e.g., GD
619 and T1w/T2w$_{\text{mi}}$). Environmental correlations (r_E) were calculated similarly to r_A but using environmental
620 variance and covariance estimates.

621
Figure 7. *SEM approach*. (**A**) Structural measurement error model used to estimate deattenuated
correlation between S-A axis properties considering the distinction between intra- and inter-individual
differences in functional gradient loading variability. Observed parcel-wise values and latent
components are standardised before model fitting. (**B**) Twin model used to obtain deattenuated h_{twin}^2
estimates. Here, the covariance between the twins' parcel-wise functional gradient loadings is
estimated directly on the inter-individual component of variance (i.e., FC$_{G1}$) and set to $f * \sigma_{A\text{inter}}^2$, where
f = 1 for MZ twins and f = ½ for DZ twins. The model assumes absence of shared household and non-
additive genetic effects on functional gradients in young adults. (**C**) Simplified graphical representation
of the multivariate twin-informed SEM, which puts the models depicted in A and B together. *Notes on
SEM: Rectangles represent the measured structural or functional MRI-derived phenotypes; the circle is
the latent components; the double-headed arrows connecting circles with themselves represent the
variance associated with the latent components; double-headed arrows connected between circles
covariances; Where not noted, one-headed arrows are the paths (here all set to 1).*

Generalisation beyond regional associations. For each individual, we obtain two metrics for
structural and functional S-A axis properties (i.e., a total of six measures per individual):

Overall within-individual Median Absolute Deviation: we quantified the spread of the regional values
across the cortex by computing within-individual Median Absolute Deviation (MAD) of microstructure,
geodesic distances, and functional gradient loadings. MAD is a robust univariate measure of statistical
dispersion and is simply calculated as follows:

Bignardi et al.

641
$$MAD_{pj} = med(|p_{pji} - med(p_{pj})|) \tag{8}$$

Where p_{ij} is the parcel-wise value for a property p, an individual j, and a parcel I and *med* is the median.

ρ similarity index: we obtained the similarity index by estimating the Spearman rank ($ρ$) correlations
between each individual microstructure, cortico-cortical network proximity, and functional gradient
loadings with the respective S-A group-level modality vectors. For example, the $ρ$ similarity index for the
cortico-cortical network proximity for an individual j was obtained by correlating their GD with the group-
level GD. Similarly, for the same individual j, the similarity index for their functional gradient loading was
obtained by correlating their FC_{G1} on day 1 and on day 2 of scanning with the group-level $FC_{G1.}$

Similar to what is outlined above for regional analysis, we fit two multivariate AE models, one per metric.
Before model fitting, $ρ$ similarity indices were first Fisher-*z* transformed. To recapitulate regional analysis
as closely as possible within the multivariate model, we also included the measurement error model to
overall within-individual MAD and $ρ$ similarity index functional gradient loadings obtained on days 1 and
2 of scanning sessions. Note that standardised coefficients are obtained using the
lavaan::standardizedSolution() function. As a final sensitivity analysis, to discount individuals' whole
brain volume as a possible confounding effect of the relationship between SA axis structure-function
associations, we additionally included total intra-cranial volume. Precisely, we followed a two-step
procedure to discount intra-cranial volume as a possible common cause. First, we regressed out intra-
cranial volume from overall within-individual MAD and Fisher-*z* transformed $ρ$ similarity indices for all S-
A axis properties. We then re-fit the exact multivariate twin models to the residuals.

Data availability
We obtained human data from the open-access Human Connectome Project (HCP) S1200 young
adult sample. HCP Young Adult data are available at https://www.humanconnectome.org/study/hcp-
young-adult. The Supplementary Table with summary statistics can be found at
https://github.com/giacomobignardi/h2_SA_axis/tree/main/SI.
665
Code availability
All code is available and can be found at https://github.com/giacomobignardi/h2_SA_axis. SEM and
twin-based analysis were carried out using the statistical package latent variable analysis (lavaan)
https://lavaan.ugent.be/. The function to apply the measurement error model (meermo) can be found
here: https://github.com/giacomobignardi/h2_SA_axis/tree/main/R/functions/meermo. The lavaan
syntax for latent variable analysis of twin data (lavaantwda) can be found in the repository
https://github.com/giacomobignardi/h2_SA_axis/tree/main/R/functions/lavaantwda. An introduction to
twin modelling using lavaan can be found at https://rpubs.com/MichelNivard/798608. Code and tutorial
for functional gradient decomposition of functional connectomes are available
at https://brainspace.readthedocs.io/en/latest/pages/install.html. The code and tutorial to generate
geodesic distances can be found at https://micapipe.readthedocs.io/en/latest/.
677
Acknowledgements
We want to thank the Human Connectome Project, Washington University, the University of Minnesota,
and Oxford University Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil;
1U54MH091657) originally funded by the 16 N.I.H. Institutes and Centers that support the N.I.H.
Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at
Washington University. We would also like to thank Jitse Amelink, Meike D. Hettwer, and MacKenzie
D. Trupp for their comments on an earlier version of the manuscript. This study was supported by
the German Federal Ministry of Education and Research (BMBF), the Max Planck Society, and the Max
Planck School of Cognition.

Bignardi et al.

References

1. Margulies DS, Ghosh SS, Goulas A, Falkiewicz M, Huntenburg JM, Langs G, et al. Situating the default-mode network along a principal gradient of macroscale cortical organization. Proc Natl Acad Sci. 2016 Nov 1;113(44):12574–9.

2. Mesulam MM. From sensation to cognition. Brain. 1998 Jun 1;121(6):1013–52.

3. Sydnor VJ, Larsen B, Bassett DS, Alexander-Bloch A, Fair DA, Liston C, et al. Neurodevelopment of the association cortices: Patterns, mechanisms, and implications for psychopathology. Neuron. 2021 Sep 15;109(18):2820–46.

4. Sydnor VJ, Larsen B, Seidlitz J, Adebimpe A, Alexander-Bloch AF, Bassett DS, et al. Intrinsic activity development unfolds along a sensorimotor–association cortical axis in youth. Nat Neurosci. 2023 Apr;26(4):638–49.

5. von Economo CF, Koskinas GN. Die cytoarchitektonik der hirnrinde des erwachsenen menschen. J Springer. 1925;

6. Burt JB, Demirtaş M, Eckner WJ, Navejar NM, Ji JL, Martin WJ, et al. Hierarchy of transcriptomic specialization across human cortex captured by structural neuroimaging topography. Nat Neurosci. 2018 Sep;21(9):1251–9.

7. Brodmann K. ergleichende Lokalisationslehre der Grosshirnrinde in ihren Prinzipien dargestellt auf Grund des Zellenbaues. Barth. 1909;

8. Ercsey-Ravasz M, Markov NT, Lamy C, Van Essen DC, Knoblauch K, Toroczkai Z, et al. A predictive network model of cerebral cortical connectivity based on a distance rule. Neuron. 2013 Oct 2;80(1):184–97.

9. Buckner RL, Krienen FM. The evolution of distributed association networks in the human brain. Trends Cogn Sci. 2013 Dec 1;17(12):648–65.

10. García-Cabezas MÁ, Hacker JL, Zikopoulos B. Homology of neocortical areas in rats and primates based on cortical type analysis: an update of the Hypothesis on the Dual Origin of the Neocortex. Brain Struct Funct. 2023 Jun 1;228(5):1069–93.

11. Saberi A, Paquola C, Wagstyl K, Hettwer MD, Bernhardt BC, Eickhoff SB, et al. The regional variation of laminar thickness in the human isocortex is related to cortical hierarchy and interregional connectivity. PLoS Biol. 2023 Nov;21(11):e3002365.

12. Markov NT, Ercsey-Ravasz M, Van Essen DC, Knoblauch K, Toroczkai Z, Kennedy H. Cortical high-density counterstream architectures. Science. 2013 Nov 1;342(6158):1238406.

13. Valk SL, Xu T, Paquola C, Park B yong, Bethlehem RAI, Vos de Wael R, et al. Genetic and phylogenetic uncoupling of structure and function in human transmodal cortex. Nat Commun. 2022 May 9;13(1):2341.

14. Demirtaş M, Burt JB, Helmer M, Ji JL, Adkinson BD, Glasser MF, et al. Hierarchical Heterogeneity across Human Cortex Shapes Large-Scale Neural Dynamics. Neuron. 2019 Mar 20;101(6):1181-1194.e13.

15. Paquola C, Amunts K, Evans A, Smallwood J, Bernhardt B. Closing the mechanistic gap: the
value of microarchitecture in understanding cognitive networks. Trends Cogn Sci. 2022 Oct
1;26(10):873–86.

16. Leech R, Vos De Wael R, Váša F, Xu T, Austin Benn R, Scholz R, et al. Variation in spatial
dependencies across the cortical mantle discriminates the functional behaviour of primary and
association cortex. Nat Commun. 2023 Sep 13;14(1):5656.

17. Suárez LE, Markello RD, Betzel RF, Misic B. Linking Structure and Function in Macroscale Brain
Networks. Trends Cogn Sci. 2020 Apr 1;24(4):302–15.

18. Hong SJ, Vos de Wael R, Bethlehem RAI, Lariviere S, Paquola C, Valk SL, et al. Atypical
functional connectome hierarchy in autism. Nat Commun. 2019 Mar 4;10(1):1022.

19. Dong D, Yao D, Wang Y, Hong SJ, Genon S, Xin F, et al. Compressed sensorimotor-to-
transmodal hierarchical organization in schizophrenia. Psychol Med. 2023 Feb;53(3):771–84.

20. Xia M, Liu J, Mechelli A, Sun X, Ma Q, Wang X, et al. Connectome gradient dysfunction in major
depression and its association with gene expression profiles and treatment outcomes. Mol
Psychiatry. 2022 Mar;27(3):1384–93.

21. Serio B, Hettwer MD, Wiersch L, Bignardi G, Sacher J, Weis S, et al. Sex differences in functional
cortical organization reflect differences in network topology rather than cortical morphometry. Nat
Commun. 2024 Sep 4;15(1):7714.

22. Dong HM, Margulies DS, Zuo XN, Holmes AJ. Shifting gradients of macroscale cortical
organization mark the transition from childhood to adolescence. Proc Natl Acad Sci. 2021 Jul
13;118(28):e2024448118.

23. Huntenburg JM, Bazin PL, Margulies DS. Large-Scale Gradients in Human Cortical Organization.
Trends Cogn Sci. 2018 Jan 1;22(1):21–31.

24. Zhang XH, Anderson KM, Dong HM, Chopra S, Dhamala E, Emani PS, et al. The cell-type
underpinnings of the human functional cortical connectome. Nat Neurosci. 2024 Nov 21;1–11.

25. Goulas A, Changeux JP, Wagstyl K, Amunts K, Palomero-Gallagher N, Hilgetag CC. The natural
axis of transmitter receptor distribution in the human cerebral cortex. Proc Natl Acad Sci. 2021
Jan 19;118(3):e2020574118.

26. Cruces RR, Royer J, Herholz P, Larivière S, Vos de Wael R, Paquola C, et al. Micapipe: A
pipeline for multimodal neuroimaging and connectome analysis. NeuroImage. 2022 Nov
1;263:119612.

27. Helwegen K, Libedinsky I, Heuvel MP van den. Statistical power in network neuroscience. Trends
Cogn Sci. 2023 Mar 1;27(3):282–301.

28. Tiego J, Martin EA, DeYoung CG, Hagan K, Cooper SE, Pasion R, et al. Precision behavioral
phenotyping as a strategy for uncovering the biological correlates of psychopathology. Nat Ment
Health. 2023 May;1(5):304–15.

29. Kang K, Seidlitz J, Bethlehem RAI, Xiong J, Jones MT, Mehta K, et al. Study design features
increase replicability in brain-wide association studies. Nature. 2024 Nov 27;1–9.

30. Brandmaier AM, Wenger E, Bodammer NC, Kühn S, Raz N, Lindenberger U. Assessing reliability
in neuroimaging research through intra-class effect decomposition (ICED). Johansen-Berg H,
Kastner S, editors. eLife. 2018 Jul 2;7:e35718.

31. Teeuw J, Hulshoff Pol HE, Boomsma DI, Brouwer RM. Reliability modelling of resting-state
functional connectivity. NeuroImage. 2021 May 1;231:117842.

32. Ge T, Holmes AJ, Buckner RL, Smoller JW, Sabuncu MR. Heritability analysis with repeat
measurements and its application to resting-state functional connectivity. Proc Natl Acad Sci.
2017 May 23;114(21):5521–6.

33. Van Essen DC, Ugurbil K, Auerbach E, Barch D, Behrens TEJ, Bucholz R, et al. The Human
Connectome Project: A data acquisition perspective. NeuroImage. 2012 Oct;62(4):2222–31.

34. Schaefer A, Kong R, Gordon EM, Laumann TO, Zuo XN, Holmes AJ, et al. Local-Global
Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. Cereb
Cortex. 2018 Sep 1;28(9):3095–114.

35. Oligschläger S, Huntenburg JM, Golchert J, Lauckner ME, Bonnen T, Margulies DS. Gradients of
connectivity distance are anchored in primary cortex. Brain Struct Funct. 2017 Jul;222(5):2173–
82.

36. Mowinckel AM, Vidal-Piñeiro D. Visualization of Brain Statistics With R Packages ggseg and
ggseg3d. Adv Methods Pract Psychol Sci. 2020 Dec 1;3(4):466–83.

37. Zuo XN, Xu T, Milham MP. Harnessing reliability for neuroscience research. Nat Hum Behav.
2019 Aug;3(8):768–71.

38. Hagenbeek FA, Hirzinger JS, Breunig S, Bruins S, Kuznetsov DV, Schut K, et al. Maximizing the
value of twin studies in health and behaviour. Nat Hum Behav. 2023 Jun;7(6):849–60.

39. Petersen SE, Seitzman BA, Nelson SM, Wig GS, Gordon EM. Principles of cortical areas and
their implications for neuroimaging. Neuron. 2024 Sep 4;112(17):2837–53.

40. Marek S, Tervo-Clemmens B, Calabro FJ, Montez DF, Kay BP, Hatoum AS, et al. Reproducible
brain-wide association studies require thousands of individuals. Nature. 2022
Mar;603(7902):654–60.

41. Xu T, Kiar G, Cho JW, Bridgeford EW, Nikolaidis A, Vogelstein JT, et al. ReX: an integrative tool
for quantifying and optimizing measurement reliability for the study of individual differences. Nat
Methods. 2023 Jul;20(7):1025–8.

42. van Kippersluis H, Biroli P, Dias Pereira R, Galama TJ, von Hinke S, Meddens SFW, et al.
Overcoming attenuation bias in regressions using polygenic indices. Nat Commun. 2023 Jul
25;14(1):4473.

43. Pritikin JN, Neale MC, Prom-Wormley EC, Clark SL, Verhulst B. GW-SEM 2.0: Efficient, Flexible,
and Accessible Multivariate GWAS. Behav Genet. 2021 May;51(3):343–57.

44. Grotzinger AD, Rhemtulla M, de Vlaming R, Ritchie SJ, Mallard TT, Hill WD, et al. Genomic
structural equation modelling provides insights into the multivariate genetic architecture of
complex traits. Nat Hum Behav. 2019 May;3(5):513–25.

45. Kemper KE, Sidorenko J, Wang H, Hayes BJ, Wray NR, Yengo L, et al. Genetic influence on
within-person longitudinal change in anthropometric traits in the UK Biobank. Nat Commun. 2024
May 6;15(1):3776.

46. Glasser MF, Sotiropoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL, et al. The minimal
preprocessing pipelines for the Human Connectome Project. NeuroImage. 2013 Oct 15;80:105–
24.

47. Yeo BTT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M, et al. The
organization of the human cerebral cortex estimated by intrinsic functional connectivity. J
Neurophysiol. 2011 Sep;106(3):1125–65.

48. Dijkstra EW. A note on two problems in connexion with graphs. Numer Math. 1959 Dec
1;1(1):269–71.

49. Vos de Wael R, Benkarim O, Paquola C, Lariviere S, Royer J, Tavakol S, et al. BrainSpace: a
toolbox for the analysis of macroscale gradients in neuroimaging and connectomics datasets.
Commun Biol. 2020 Mar 5;3(1):1–10.

50. Bethlehem RAI, Paquola C, Seidlitz J, Ronan L, Bernhardt B, Consortium CC, et al. Dispersion of
functional gradients across the adult lifespan. NeuroImage. 2020 Nov 15;222:117299.

51. Koo TK, Li MY. A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for
Reliability Research. J Chiropr Med. 2016 Jun;15(2):155–63.

52. Rosseel Y. lavaan: An R Package for Structural Equation Modeling. J Stat Softw. 2012 May
24;48(1):1–36.

53. Bates TC, Maes H, Neale MC. umx: Twin and Path-Based Structural Equation Modeling in R.
Twin Res Hum Genet Off J Int Soc Twin Stud. 2019 Feb;22(1):27–41.

54. Boomsma D, Busjahn A, Peltonen L. Classical twin studies and beyond. Nat Rev Genet. 2002
Nov;3(11):872–82.

55. Verhulst B, Prom-Wormley E, Keller M, Medland S, Neale MC. Type I Error Rates and Parameter
Bias in Multivariate Behavioral Genetic Models. Behav Genet. 2019 Jan;49(1):99–111.

56. Bignardi G, Wesseldijk LW, Mas-Herrero E, Zatorre RJ, Ullén F, Fisher SE, et al. Twin modelling
reveals partly distinct genetic pathways to music enjoyment. Nat Commun. 2025 Mar
25;16(1):2904.

829