Serpine1 negatively regulates Th1 cell responses in experimental autoimmune

encephalomyelitis.

(Running title: Serpine1 negatively regulates Th1 cell responses in EAE)

Irshad Akbar[*,#], Ruihan Tang[†], Joanie Baillargeon[*], Andrée-Pascale Roy[*], Prenitha Mercy

Ignatius Arokia Doss[*], Chen Zhu[†], Vijay K. Kuchroo[†,‡,§,‖], Manu Rangachari[*,¶, ‖, **,††, ‡‡]

[*] axe Neurosciences, Centre de recherche du CHU de Québec-Université Laval, Québec QC

Canada

[†] Evergrande Center for Immunologic Diseases, Harvard Medical School, Brigham and

Women's Hospital, Boston MA USA

[‡] Ann Romney Center for Neurologic Diseases, Harvard Medical School, Brigham and Women's

Hospital, Boston MA USA

[§] Klarman Cell Observatory, Broad Institute of MIT and Harvard, Cambridge MA USA

[¶] Department of Molecular Medicine, Université Laval, Québec, QC, Canada

[‖] Address correspondence and reprint request to Dr. Manu Rangachari or Dr. Vijay K. Kuchroo.

Axe Neurosciences, Centre de recherche du CHU de Québec-Université Laval, 2705 boul

Laurier, Québec QC G1V 4G2 (M.R) or Brigham and Women's Hospital, Hale Building for

Transformative Research, 60 Fenwood Rd, HBTM 10016F, Boston MA 02115. Email addresses:

manu.rangachari@crchudequebec.ulaval.ca, vkuchroo@rics.bwh.harvard.edu

[#] funding to I.A., MS Society of Canada Doctoral Studentship

[**] funding to M.R., MS Society of Canada Discovery Grant #3781

[††] funding to M.R., Canadian Institutes of Health Research (CIHR) Project Grant #159713

[‡‡] funding to M.R., Senior scholar award, Fonds de recherche de Québec – Santé #313330

Phone (M.R) : ++1-418-525-4444 x 46461

Fax (M.R). : ++1-418-654-2298

Abstract

Th1 cells are critical in experimental autoimmune encephalomyelitis (EAE). Serpine1 has been posited as an inhibitor of IFNγ from T cells though its role in autoimmunity remains unclear. Here, we show that *Serpine1*$^{-/-}$ mice develop EAE of enhanced severity relative to wild-type controls. Serpine1 overexpression represses Th1 cell cytokine production and pathogenicity, while *Serpine1*$^{-/-}$ x 2D2 Th1 cells transfer EAE of increased severity in comparison to *Serpine1*$^{+/+}$ x 2D2 Th1 cells. Notably, polarized *Serpine1*$^{-/-}$ Th1 cells display delayed expression of the Th1-specific inhibitory receptor, Tim-3. *Serpine1*$^{-/-}$ Tim-3-Tg Th1 cells, which transgenically over-express Tim-3, showed increased expression of IFNγ and reduced expression of the checkpoint molecules Lag-3 and PD-1 relative to Tim-3-Tg counterparts. Further, Serpine1 deficiency restored the EAE phenotype of Tim-3-Tg mice that normally develop mild disease. Together, we identify Serpine1 as a negative regulator of Th1 cells.

Key points

- Serpine1 inhibits EAE in a T cell-dependent manner.

- Serpine1 is upregulated in Th1 cells and inhibits their pathogenicity.

- Serpine1 promotes expression and function of Th1-specific inhibitory receptor Tim-3.

Introduction

Effector CD4[+] Th1 cells are characterized by IFNγ production, recruit macrophages and

CD8[+] T cells to sites of infection and inflammation and are potent initiators and propagators of

autoimmune diseases such as in the T cell-driven EAE[1] model of MS[2] (1). Th1 cells express a

lineage-specific inhibitory receptor, Tim-3[3], that resolves inflammatory responses when

triggered at sites of inflammation (2-4). The Tim-3 pathway is crucial to repressing EAE

pathology (2, 5, 6). Understanding the processes regulating Tim-3 expression and function might

permit us to develop strategies to curb Th1-mediated inflammation in self-tissue.

Serpine1[4], or PAI-1[5], represses the conversion of the plasminogen proenzyme into mature

plasmin and was previously suggested to regulate IFNγ-driven T cell responses (7, 8). Here, we

show that Serpine1 restrains EAE pathogenicity via its inhibitory effects on Th1 cells, and that

Serpine1 is required for optimal expression of Tim-3 by Th1 cells.

Materials and Methods

Mice

Tim-3-Tg (5) are described. B6[6] *(Serpine1[+/+]*; stock #000664), *Serpine1[-/-]* (B6.129S2-

Serpine1[tm1Mlg]/J; #002507), 2D2-Tg (C57BL/6-Tg (Tcra2D2,Tcrb2D2)1KuchJ; #006912) and

Rag1[-/-] (B6.129S7-*Rag1[tm1Mom]*/J; #002216) mice were obtained from Jackson Labs.

Experimental and control animals were co-housed at the animal facilities of CRCHU de Québec-

Université Laval or Brigham & Women's Hospital. All procedures were authorized by the

Animal Care Committee of Université Laval (CPAUL-3) or the Institutional Animal Care and

Use Committee (IACUC) of Harvard University.

T helper cell differentiation

CD4[+] T cells were enriched from B6 spleens using anti-CD4 Microbeads (Miltenyi), purified as

CD4+CD62L[hi] using a FACSAria (BD) high-speed cell sorter and cultured in supplemented T

cell media as described (3). They were differentiated (9) for 2 days with plate bound anti-CD3

and anti-CD28 (2 μg mL[-1] each; BioXcell) into Th1 cells – 10 ng mL[-1] rmIL-12 (R&D

Biosystems) plus anti-IL-4 (10 μg mL[-1], BioXcell), or Th17 – rhTGFβ (3 ng mL[-1], Miltenyi) +

rmIL-6 (20 ng mL[-1], Miltenyi) + anti-IFNγ (10 μg mL[-1], BioXcell). Cells were then transferred

to uncoated tissue culture plates and cultured for an additional 3 days, with rmIL-2 (10 ng mL[-1],

Miltenyi) added to Th1, and rmIL-23 (20 ng mL[-1], R&D Biosystems) added to Th17. For

multiple rounds of polarization, cells were collected at d5 and then restimulated for a subsequent

5-day period as above. Tiplaxtinin (25 μM, Tocris) or equivalent volume of DMSO were

maintained for 5 days of culture where indicated. Serpine1 cDNA was cloned into pMIG[7] vector

and RV[8] gene transduction of Th1 or Th17 cells was conducted using our described spin-

infection protocol (10).

EAE

Active immunization was induced by s.c. injection of $MOG_{[35-55]}$[9] (CHU de Québec) in

incomplete Freund's adjuvant (Difco) supplemented with 5 mg mL^{-1} *Mycobacterium*

tuberculosis extract (Fisher). A dose of 25 μg $MOG_{[35-55]}$ per mouse was used to compare

Serpine1$^{+/+}$ to *Serpine1*$^{-/-}$ mice, while 100 μg per mouse was used in EAE experiments involving

Tim-3-Tg mice. In Figure 1D, passive EAE was induced by first immunizing *Serpine1*$^{+/+}$ 2D2

and *Serpine1*$^{-/-}$ 2D2 mice with $MOG_{[35-55]}$. Nine days later, splenocytes were collected and

stimulated *ex vivo* for 48 hours with $MOG_{[35-55]}$ (20 μg mL^{-1}), rmIL-12 and rmIL-23. Next,

$20x10^6$ splenocytes were injected i.p. into unimmunized B6 recipients. Adoptive transfer of 2D2,

Serpine1$^{-/-}$ 2D2 or RV-transduced 2D2 Th1 cells entailed i.v. injection ($2x10^6$) into *Rag1*$^{-/-}$

recipients at d5 of culture. In all EAE experiments, mice received 200 ng pertussis toxin (List

Biological Labs) i.p. at d0 and d2. Mice were assessed for clinical symptoms daily as previously

described (9).

Flow cytometry

Cell surface and intracellular flow cytometry were conducted as previously described (10). The

following Abs and dyes were used: *CD4*, clone RM4-5, ThermoFisher (TF) cats #45-0042-82,

48-0042-82; CD62L, MEL-14, TF #47-0621-82, *Tim-3*, RMT3-23, Biolegend #119706; *PD-1*,

J43, TF #25-9985-82; *Lag-3*, eBioC9B7W, TF #12-2231-82; *IFNγ*, XMG1.2, TF # 48-7311-82;

TNFα, MP6-XT22, TF #11-7321-82, 12-7321-41, 17-7321-82; IL-2, JES6-5H4, BD Biosciences

#560547; IL-17, TC11-18H10.1, Biolegend #506922; Fixable Viability Dye, TF #65-0865-14; 7-

aminoactinomycin D, TF #A1310. Data were collected using an LSRII flow cytometer (BD

Biosciences) and were analyzed with FlowJo (BD). Gates were set on fluorescence minus one

controls and the following global strategy was used: *i)* singlets were selected based on FSC-H vs

FSC-A; *ii)* live $CD4^+$ T cells based on $CD4^+$ Viability Dye^{neg} events, or on $CD4^+$7-

aminoactinomycin D^{neg} events. RV-transduced $CD4^+$ T cells were further gated on GFP

positivity as indicated in Figure legends. In Figures 4CD, live $CD4^+$ T cells were gated as

$CD4^+$Tim-3^{pos} or $CD4^+$Tim-3^{neg}.

Ex vivo assessment of T cell function

Splenocyte cultures from EAE mice were stimulated, or not, with 10 μg mL^{-1} $MOG_{[35-55]}$ for 48

hours. For proliferation studies, 1.25 μCi [^{3}H]-thymidine (Perkin-Elmer) was added to each

culture well for the last 16 hours. Cytokine supernatant ELISA were conducted using the

following capture/detection sets: *IFNγ*, clones RA-6A2/XMG1.2; *IL-2*, JES6-1412/JES6-5H4;

IL-17, TC11-18H10.1/TC11-8H4. Serpine1 protein was measured using Serpin E1/PAI-1

DuoSet ELISA (R&D Biosystems).

Statistics

Two-tailed parametric tests were conducted using Prism (GraphPad). Comparisons of 2 groups

were made by *t*-test while comparisons of >2 groups were made by ANOVA followed by post-

hoc test. In EAE studies, linear regression (6, 10) or area under curve (10) analyses were

conducted on mice with symptoms.

Results and Discussion

Serpine1 inhibits the severity of EAE.

Upon immunization with MOG$_{[35-55]}$, *Serpine1$^{-/-}$* mice developed EAE of significantly greater severity than *Serpine1$^{+/+}$* controls (Figure 1A). When the data were sex-disaggregated, *Serpine1$^{-/-}$* females showed a significantly worsened disease burden as compared to *Serpine1$^{+/+}$* females, while *Serpine1$^{-/-}$* males showed a trend towards exacerbated disease relative to *Serpine1$^{+/+}$* males (Supplementary Figure 1). Notably, antigen-specific proliferation responses to MOG$_{[35-55]}$ were enhanced in peripheral *Serpine1$^{-/-}$* CD4$^+$ T cells prior to clinical onset (Figure 1B), as was secretion of IFNγ and IL-2, but not IL-17 (Figure 1C).

Serpine1 regulates CNS fibrinolysis (11) and thus heightened EAE severity in *Serpine1$^{-/-}$* mice might be due to its absence in a non-T cell compartment. We thus crossed the *Serpine1$^{-/-}$* strain to 2D2-Tg strain, which bear a MOG$_{[35-55]}$ -specific TCR (12), and immunized *Serpine1$^{+/+}$* 2D2 and *Serpine1$^{-/-}$* 2D2 mice. Prior to disease onset, we isolated splenocytes and restimulated them with Ag plus IL-12 and IL-23. *Serpine1$^{-/-}$* 2D2 Ag-restimulated blasts induced EAE of significantly greater severity relative to *Serpine1$^{+/+}$* 2D2 blasts upon adoptive transfer (Figure 1D).

Previous studies have been conflicting with respect to the role of Serpine1 in EAE, possibly due to opposing effects on T cells and CNS repair. Treating MOG$_{[35-55]}$-immunized mice with a Serpine1-derived 18mer peptide reduced disease severity; however its effects were not T cell-specific (13). By contrast, a small molecule inhibitor of Serpine1, TM5484, reduced EAE severity, thereby suggesting that Serpine1 augments inflammation; however, this may have been due to CNS-intrinsic effects, as TM5484 can permeate the blood-brain barrier (14).

Notably, in the chronic/relapsing Biozzi model of EAE, *Serpine1*[-/-] mice developed EAE of

delayed onset and milder severity, due to superior CNS fibrinolytic capacity compared to

controls (11); it is possible that the fibrogenic, pathogenic, properties of Serpine1 outweigh its

anti-inflammatory function in this model. Here, we show that Serpine1 represses antigen-specific

T cell-driven CNS autoimmunity in a T cell-intrinsic manner.

Serpine1 suppresses Th1 cell function in vitro and in vivo

We next found that Serpine1 secretion was sharply upregulated in both Th1 (Figure 2A)

and Th17 (Figure 2B) cells upon a second round of *in vitro* differentiation. Serpine1-deficient T

cells can produce increased levels of IFNγ relative *in vivo* (7, 8); however, the functional role of

Serpine1 in differentiated *bonafide* Th1 and Th17 cells has never been directly assessed. Upon

RV OE[10] of Serpine1 in Th1 cells, we observed a downregulation in expression of IFNγ and a

trend towards reduced TNFα relative to control transduced cells. By contrast, overexpression of

Serpine1 in Th17 cells did not impact expression of either IL-17 or TNFα (Figure 2C).

To determine whether enforced expression of Serpine1 altered Th1 cell pathogenicity, we

transduced 2D2 Th1 cells with Serpine1-OE or control RV and adoptively transferred these cells

to *Rag1*[-/-] mice (6). Interestingly, Serpine1-OE 2D2 Th1 cells induced disease of lessened

severity compared to control cells (Figure 2D), characterized by a reduced frequency of

inflammatory IFNγ[+]IL-2[+] T cells *in vivo* (Figure 2E).

We next treated Th1 cells with tiplaxtinin, a small molecule that inhibits the

antiproteolytic activity of Serpine1 towards the plasminogen activators urokinase and tissue

plasminogen activator (15, 16). Tiplaxtinin enhanced IFNγ and IL-2 production from Th1 cells

(Figure 3A), suggesting that the Serpine1 may downregulate Th1 responses by repressing the

activation of mature plasmin. We then generated Th1 cells from *Serpine1*[+/+] 2D2 and *Serpine1*[-/-]

2D2 mice and found that *Serpine1*[-/-] 2D2 Th1 cells induced EAE of significantly greater severity

upon adoptive transfer (Figure 3B).

Increased IFNγ was previously observed from *Serpine1*[-/-] CD4[+] and CD8[+] T cells upon

LPS or staphylococcal enterotoxin B treatment *in vivo* (7). Further, *Serpine1*[-/-] mice are resistant

to nasal allergy in a Th2-mediated OVA sensitization model, with IFNγ production upregulated

by *Serpine1*[-/-] splenocytes upon Ag recall (8). Here, we show that while Serpine1 is expressed by

both Th1 and Th17 cells, it impacts Th1 cells specifically by downregulating their inflammatory

cytokine production and autoimmune potential.

Serpine1 promotes Tim-3 expression and inhibits Tim-3[+] Th1 cell inflammatory responses.

As Serpine1 inhibits Th1-driven inflammation, we next asked whether it could augment

the expression of inhibitory Tim-3. Repeatedly polarized *Serpine1*[-/-] Th1 cells expressed lower

Tim-3, and with delayed kinetics, relative to *Serpine1*[+/+] controls (Figure 4A). Loss of Serpine1

signaling did not increase expression of IFNγ; however, the frequency of IFNγ[+]Tim-3[+] Th1 cells

was strikingly lower in its absence (Figure 4B).

Tim-3-Tg mice ectopically overexpress Tim-3 cDNA in a T cell-restricted manner,

without the need for multiple rounds of polarization under Th1 conditions (17). Reasoning that

this might help us uncover Serpine1-dependent differences in IFNγ expression, we crossed

Serpine1[-/-] mice to the Tim-3-Tg strain and generated Th1 cells from these and Tim-3-Tg control

mice. Tim-3[pos] Th1 cells derived from *Serpine1*[-/-] Tim-3-Tg mice were strikingly more positive

for IFNγ than their *Serpine1*[+/+] Tim-3-Tg counterparts; notably, no differences were observed

with Tim-3[neg] Th1 cells between the strains (Figure 4C). This indicated that Serpine1 represses

IFNγ expression in a Tim-3-dependent manner. Interestingly, there was a concomitant reduction

in the expression of the T cell negative regulatory receptors PD-1 and Lag-3 in *Serpine1*$^{-/-}$ Tim-

3-Tg Th1 cells when compared to *Serpine1*$^{+/+}$ Tim-3-Tg controls (Figure 4D).

Tim-3-Tg mice develop EAE of attenuated severity (5). To examine whether loss of

Serpine1 expression could reverse this phenotype, we actively immunized *Serpine1*$^{+/+}$,

Serpine1$^{+/+}$ Tim-3-Tg and *Serpine1*$^{-/-}$ Tim-3-Tg mice. While *Serpine1*$^{+/+}$ Tim-3-Tg mice

developed disease of only mild severity, *Serpine1*$^{-/-}$ Tim-3-Tg mice displayed EAE that was of

comparable severity to that seen in wild-type animals (Fig 4E). Our data thus demonstrate that

Serpine1 is required for Tim-3-mediated repression of T cell inflammation and pathogenicity in

EAE.

Tim-3 marks exhausted T cells in chronic viral infections and cancers (18-20) and is

functionally tractable, as concomitant blockade of Tim-3 and PD-1 causes tumor regression

greater degree than that with anti-PD-1 alone (20). Further, depletion of Bat3, an intracellular

repressor of Tim-3 signaling, causes Th1 cells to adopt an exhausted-like phenotype *in vivo* (6,

21). While Serpine1-mediated upregulation of Tim-3 may be desirable in the context of

autoimmunity, it remains to be seen what role, if any, Serpine1 plays in T cell exhaustion.

Altogether, our data identify a novel Th1 cell-intrinsic regulatory mechanism. Serpine1

represses Th1 cell pathogenicity by inhibiting their production of inflammatory cytokines and by

restraining them from adopting a highly differentiated Tim-3^{+} phenotype. Strategies to augment

Serpine1 expression and function in Th1 cells could present an attractive therapeutic option in

autoimmune disease.

Acknowledgements

We thank Vincent Desrosiers for technical assistance, Kim Larose-Labrecque and Andrée

Brisson for animal care, and Ryder Whittaker Hawkins for critical reading of the manuscript.

Author Contributions[11]

Data availability statement[12]

Conflicts of interest[13]

References

1. Jäger, A., V. Dardalhon, R. A. Sobel, E. Bettelli, and V. K. Kuchroo. 2009. Th1, Th17, and
Th9 effector cells induce experimental autoimmune encephalomyelitis with different
pathological phenotypes. *J. Immunol.* 183: 7169–7177.
2. Monney, L., C. A. Sabatos, J. L. Gaglia, A. Ryu, H. Waldner, T. Chernova, S. Manning, E. A.
Greenfield, A. J. Coyle, R. A. Sobel, G. J. Freeman, and V. K. Kuchroo. 2002. Th1-specific cell
surface protein Tim-3 regulates macrophage activation and severity of an autoimmune disease.
Nature 415: 536–541.
3. Sabatos, C. A., S. Chakravarti, E. Cha, A. Schubart, A. Sánchez-Fueyo, X. X. Zheng, A. J.
Coyle, T. B. Strom, G. J. Freeman, and V. K. Kuchroo. 2003. Interaction of Tim-3 and Tim-3
ligand regulates T helper type 1 responses and induction of peripheral tolerance. *Nat. Immunol.*
4: 1102–1110.
4. Sánchez-Fueyo, A., J. Tian, D. Picarella, C. Domenig, X. X. Zheng, C. A. Sabatos, N.
Manlongat, O. Bender, T. Kamradt, V. K. Kuchroo, J.-C. Gutiérrez-Ramos, A. J. Coyle, and T.
B. Strom. 2003. Tim-3 inhibits T helper type 1-mediated auto- and alloimmune responses and
promotes immunological tolerance. *Nat. Immunol.* 4: 1093–1101.
5. Dardalhon, V., A. C. Anderson, J. Karman, L. Apetoh, R. Chandwaskar, D. H. Lee, M.
Cornejo, N. Nishi, A. Yamauchi, F. J. Quintana, R. A. Sobel, M. Hirashima, and V. K. Kuchroo.
2010. Tim-3/galectin-9 pathway: regulation of Th1 immunity through promotion of CD11b+Ly-
6G+ myeloid cells. *J. Immunol.* 185: 1383–1392.
6. Rangachari, M., C. Zhu, K. Sakuishi, S. Xiao, J. Karman, A. Chen, M. Angin, A. Wakeham,
E. A. Greenfield, R. A. Sobel, H. Okada, P. J. McKinnon, T. W. Mak, M. M. Addo, A. C.
Anderson, and V. K. Kuchroo. 2012. Bat3 promotes T cell responses and autoimmunity by
repressing Tim-3–mediated cell death and exhaustion. *Nat. Med.* 18: 1394–1400.
7. Renckens, R., J. M. Pater, and T. van der Poll. 2006. Plasminogen activator inhibitor type-1-
deficient mice have an enhanced IFN-gamma response to lipopolysaccharide and staphylococcal
enterotoxin B. *J. Immunol.* 177: 8171–8176.
8. Sejima, T., S. Madoiwa, J. Mimuro, T. Sugo, K. Okada, S. Ueshima, O. Matsuo, T. Ishida, K.
Ichimura, and Y. Sakata. 2005. Protection of plasminogen activator inhibitor-1-deficient mice
from nasal allergy. *J. Immunol.* 174: 8135–8143.
9. Pradeep Yeola, A., I. Akbar, J. Baillargeon, P. Mercy Ignatius Arokia Doss, V. O. Paavilainen,
and M. Rangachari. 2020. Protein translocation and retro-translocation across the endoplasmic
reticulum are crucial to inflammatory effector CD4+ T cell function. *Cytokine* 129: 154944.
10. Doss, P. M. I. A., M. Umair, J. Baillargeon, R. Fazazi, N. Fudge, I. Akbar, A. P. Yeola, J. B.
Williams, M. Leclercq, C. Joly Beauparlant, P. Beauchemin, G. F. Ruda, M. Alpaugh, A. C.
Anderson, P. E. Brennan, A. Droit, H. Lassmann, C. S. Moore, and M. Rangachari. 2021. Male
sex chromosomal complement exacerbates the pathogenicity of Th17 cells in a chronic model of
central nervous system autoimmunity. *Cell Rep* 34: 108833.
11. East, E., D. Gveric, D. Baker, G. Pryce, H. R. Lijnen, and M. L. Cuzner. 2008. Chronic
relapsing experimental allergic encephalomyelitis (CREAE) in plasminogen activator inhibitor-1
knockout mice: the effect of fibrinolysis during neuroinflammation. *Neuropathol. Appl.*
Neurobiol. 34: 216–230.
12. Bettelli, E., M. Pagany, H. L. Weiner, C. Linington, R. A. Sobel, and V. K. Kuchroo. 2003.
Myelin oligodendrocyte glycoprotein-specific T cell receptor transgenic mice develop
spontaneous autoimmune optic neuritis. *J. Exp. Med.* 197: 1073–1081.

13. Gur-Wahnon, D., T. Mizrachi, F.-Y. Maaravi-Pinto, A. Lourbopoulos, N. Grigoriadis, A.-A.
R. Higazi, and T. Brenner. 2013. The plasminogen activator system: involvement in central
nervous system inflammation and a potential site for therapeutic intervention. *J*
Neuroinflammation 10: 124.
14. Pelisch, N., T. Dan, A. Ichimura, H. Sekiguchi, D. E. Vaughan, C. van Ypersele de Strihou,
and T. Miyata. 2015. Plasminogen Activator Inhibitor-1 Antagonist TM5484 Attenuates
Demyelination and Axonal Degeneration in a Mice Model of Multiple Sclerosis. *PLoS ONE* 10:
e0124510.
15. Elokdah, H., M. Abou-Gharbia, J. K. Hennan, G. McFarlane, C. P. Mugford, G.
Krishnamurthy, and D. L. Crandall. 2004. Tiplaxtinin, a novel, orally efficacious inhibitor of
plasminogen activator inhibitor-1: design, synthesis, and preclinical characterization. *J. Med.*
Chem. 47: 3491–3494.
16. Gorlatova, N. V., J. M. Cale, H. Elokdah, D. Li, K. Fan, M. Warnock, D. L. Crandall, and D.
A. Lawrence. 2007. Mechanism of inactivation of plasminogen activator inhibitor-1 by a small
molecule inhibitor. *J. Biol. Chem.* 282: 9288–9296.
17. Dardalhon, V., A. S. Schubart, J. Reddy, J. H. Meyers, L. Monney, C. A. Sabatos, R. Ahuja,
K. Nguyen, G. J. Freeman, E. A. Greenfield, R. A. Sobel, and V. K. Kuchroo. 2005. CD226 is
specifically expressed on the surface of Th1 cells and regulates their expansion and effector
functions. *J. Immunol.* 175: 1558–1565.
18. Jones, R. B., L. C. Ndhlovu, J. D. Barbour, P. M. Sheth, A. R. Jha, B. R. Long, J. C. Wong,
M. Satkunarajah, M. Schweneker, J. M. Chapman, G. Gyenes, B. Vali, M. D. Hyrcza, F. Y. Yue,
C. Kovacs, A. Sassi, M. Loutfy, R. Halpenny, D. Persad, G. Spotts, F. M. Hecht, T.-W. Chun, J.
M. McCune, R. Kaul, J. M. Rini, D. F. Nixon, and M. A. Ostrowski. 2008. Tim-3 expression
defines a novel population of dysfunctional T cells with highly elevated frequencies in
progressive HIV-1 infection. *J. Exp. Med.* 205: 2763–2779.
19. Jin, H.-T., A. C. Anderson, W. G. Tan, E. E. West, S.-J. Ha, K. Araki, G. J. Freeman, V. K.
Kuchroo, and R. Ahmed. 2010. Cooperation of Tim-3 and PD-1 in CD8 T-cell exhaustion during
chronic viral infection. *Proc. Natl. Acad. Sci. U.S.A.* 107: 14733–14738.
20. Sakuishi, K., L. Apetoh, J. M. Sullivan, B. R. Blazar, V. K. Kuchroo, and A. C. Anderson.
2010. Targeting Tim-3 and PD-1 pathways to reverse T cell exhaustion and restore anti-tumor
immunity. *J. Exp. Med.* 207: 2187–2194.
21. Zhu, C., K. O. Dixon, K. Newcomer, G. Gu, S. Xiao, S. Zaghouani, M. A. Schramm, C.
Wang, H. Zhang, K. Goto, E. Christian, M. Rangachari, O. Rosenblatt-Rosen, H. Okada, T. Mak,
M. Singer, A. Regev, and V. Kuchroo. 2021. Tim-3 adaptor protein Bat3 is a molecular
checkpoint of T cell terminal differentiation and exhaustion. *Sci Adv* 7.

[1] Experimental autoimmune encephalomyelitis
[2] Multiple sclerosis
[3] T cell immunoglobulin and mucin domain-containing-3
[4] Serine protease inhibitor clade E1
[5] plasminogen activation inhibitor 1
[6] C57BL6/J
[7] pMSCV-IRES-GFP
[8] retroviral
[9] Myelin oligodendrocyte glycoprotein, amino acids 35-55
[10] overexpression

[11] I.A. conducted experiments and managed the project. R.T., J.B., A-P.R., P.M.I.A.D and C.Z. conducted experiments. V.K.K. co-supervised the project. M.R. conducted experiments, supervised the project and wrote the manuscript.

[12] The datasets generated for this study are available from the corresponding authors upon reasonable request.

[13] V.K.K. has an ownership stake and is a member of the Scientific Advisory Board for Tizona Therapeutics. Further, he is a co-founder of, and has an ownership stake in, Celsius Therapeutics. In addition, he is an inventor on patents related to Th17 cell function. His interests are reviewed and managed by Brigham & Women's Hospital and Partners Healthcare in accordance with their conflict of interest policies.

    Figure Legends

Figure 1. Serpine1 inhibits the severity of EAE. A. *Left,* Representative EAE curve of

$MOG_{[35-55]}$-immunized *Serpine1$^{+/+}$* (n=9) and *Serpine1$^{-/-}$* (n=7) female mice. *Middle*, linear

regression curves of the representative disease courses. The dashed lines indicate the 95%

confidence intervals for each curve. *Right*, area-under-curve (AUC) comparison of mice pooled

from 3 experiments. Filled circles, female mice (n=14, *Serpine1$^{+/+}$*, n=12, *Serpine1$^{-/-}$*); open

circles, male mice (n=10, *Serpine1$^{+/+}$*, n=10, *Serpine1$^{-/-}$*). **, p<0.01; two way-ANOVA

analysis of genotype as a variable. Incidence of EAE over all experiments was 24/25

Serpine1$^{+/+}$, 22/24 *Serpine1$^{-/-}$*. **B.** $MOG_{[35-55]}$-immunized female *Serpine1$^{+/+}$* and *Serpine1$^{-/-}$* mice

(n=3 each) were sacrificed 10 days post-immunization, and lymph node cells were assessed for

proliferation over 48 hours in response to $MOG_{[35-55]}$. cpm, counts per minute. **, p<0.01, *t*-test.

C. Splenocytes from immunized female *Serpine1$^{+/+}$* and *Serpine1$^{-/-}$* mice (n=3 each) were

restimulated with $MOG_{[35-55]}$ for 48 (IL-2) or 72 (IFNγ, IL-17) hours, and secretion of the

indicated cytokines was measured by ELISA. *, p<0.05; **, p<0.01, *t*-test. **D.** Splenocytes from

female *Serpine1$^{+/+}$* 2D2 and *Serpine1$^{-/-}$* 2D2 mice were restimulated with $MOG_{[35-55]}$ in the

presence of IL-12 and IL-23, prior to adoptive transfer to WT mice (n=5 each condition) that

were monitored for signs of EAE. Right graph, linear regression analysis with 95% confidence

intervals. ****, p<0.0001.

Figure 2. Serpine1 is expressed in Th1 cells and suppresses Th1-driven EAE. A, B. Naïve

$CD4^{+}CD62L^{hi}$ T cells were isolated from female B6 mouse spleen and were differentiated under

Th1 or Th17 conditions for 2 rounds of polarization. Secretion of Serpine1 protein was measured

by ELISA in supernatant from Th1 (**A**) or Th17 (**B**) by ELISA. *, p<0.05; ***, p<0.001, one

way ANOVA. Cultures derived from 4 independent mice. **C.** Female B6 Th1 or Th17 cells were

transduced with control or Serpine1-OE RV. Cells were analyzed for production of the indicated

cytokines after 5 days. Gated on GFP-positive live events. Quantitation represents paired *t*-test

analysis of 3 independent cultures each. *, p<0.05. **D.** Female 2D2 Th1 cells were transduced

with control- (n=5) or Serpine1-OE (n=4) RV. After 5 days of stimulation, cells were adoptively

transferred to *Rag1*$^{-/-}$ recipients who were monitored for signs of EAE. *Bottom*, linear regression

curves and 95% confidence intervals, for the disease courses. **E.** At disease endpoint (d35),

splenic CD4$^+$ T cells from mice in (**D**) were assessed for production of IFNγ and IL-2 by flow

cytometry. Gated on live CD4$^+$ events. **, p<0.01, *t*-test.

Figure 3. Loss of Serpine1 function or expression exacerbates Th1 responses. A. Female B6

Th1 cells were treated with DMSO (control) or 25 μM tiplaxtinin for 5 days, at which point IFNγ

and IL-2 were measured by flow cytometry. *, p<0.05, t-test; 3 independent observations. **B.**

Female *Serpine1*$^{+/+}$ 2D2 and *Serpine1*$^{-/-}$ 2D2 CD4$^+$ Th1 cells were transferred i.v. to *Rag1*$^{-/-}$

recipients (2x10^6 cells/mouse). n=4, *Serpine1*$^{+/+}$ 2D2; n=5, *Serpine1*$^{-/-}$ 2D2. Mice were

subsequently assessed for signs of EAE. Left graph, linear regression curves of the disease

courses.

Figure 4. Serpine1 promotes Tim-3 expression and inhibits Tim-3$^+$ Th1 cell inflammatory

responses. A, B. Female *Serpine1*$^{+/+}$ and *Serpine1*$^{-/-}$ T cells were subjected to 4 successive

rounds of Th1 polarization, and Tim-3 (**A,B**) and IFNγ (**B**) were assessed by flow cytometry. **(A)**

Quantification of Tim-3 expression at the end of each round. **, p<0.01, one-way ANOVA. **B.**

Expression of Tim-3 versus IFNγ after 20 days of total culture (end of round 4). Gated on live

CD4$^+$ events. Representative of 4 experiments. **C.** Splenic T CD4$^+$ T cells from male *Serpine1$^{+/+}$*

Tim-3-Tg and *Serpine1$^{-/-}$* Tim-3-Tg mice (n=6 independent cultures from each) were

differentiated under Th1 conditions for 5 days. CD4, Tim-3 and IFNγ expression were assessed

by flow cytometry, with cells first gated on live CD4$^+$Tim-3^{pos} or live CD4$^+$Tim-3^{neg}. ***,

p<0.001; *t*-test. **D.** Splenic T CD4$^+$ T cells from male *Serpine1$^{+/+}$* Tim-3-Tg and *Serpine1$^{-/-}$* Tim-

3-Tg mice (n=3 independent cultures from each) were differentiated under Th1 conditions for 5

days. Lag-3 and PD-1 expression were assessed by flow cytometry. *, p<0.05; *t*-test. Gated on

live CD4$^+$Tim-3$^+$ events. **E.** Female *Serpine1$^{+/+}$* (n=4), *Serpine1$^{-/-}$* Tim-3-Tg (n=14) and

Serpine1$^{+/+}$ Tim-3-Tg (n=5) mice were immunized with MOG$_{[35-55]}$ and were monitored for

signs of EAE. *Right graph*, linear regression analysis of the disease curves with Bonferroni's

correction applied. Representative of 3 immunizations.

Supplementary Figure 1. AUC comparison of MOG$_{[35-55]}$-immunized Serpine1$^{+/+}$ vs *Serpine1$^{-/-}$*

females (left) and males (right). *t*-test.

Figure 1

Figure 2

Figure 3

Figure 4