Main Manuscript for

Amyloid β accelerates age-related proteome-wide protein insolubility.

Edward Anderton[#1,2], Manish Chamoli[#*1], Dipa Bhaumik[1], Christina D. King[1], Xueshu Xie[1], Anna Foulger[1], Julie K. Andersen[1], Birgit Schilling[*1], Gordon J. Lithgow[*1]

Abstract

Loss of proteostasis is a highly conserved feature of aging across model organisms and typically results in the accumulation of insoluble protein aggregates. Protein insolubility is a central feature of major age-related neurodegenerative diseases including Alzheimer's Disease (AD), where hundreds of insoluble proteins associate with aggregated amyloid beta (Aβ) in senile plaques. Despite the established connection between aging and AD risk, therapeutic approaches to date have overlooked aging and proteome-wide protein insolubility as causal factors, instead focusing on Aβ and Tau. Here, using an unbiased proteomics approach, we questioned the relationship between Aβ and age-related protein insolubility. We demonstrate that, in *C. elegans*, Aβ expression is sufficient to drive proteome-wide protein insolubility. The Aβ-driven insoluble proteome bears a highly significant overlap with the aging-driven insoluble proteome, suggesting there exists a core, sub-proteome which is vulnerable to insolubility. Using human genome-wide association studies (GWAS) we show that this insoluble sub proteome is replete with biological processes implicated across not only neurodegenerative diseases but also across a broad array of chronic, age-related diseases (CARDs), providing suggestive evidence that age-related loss of proteostasis could play a role in general CARD risk. Finally, we show that the gut metabolite, Urolithin A, relieves Aβ toxicity supporting its use in clinical trials for dementia and other age-related diseases.

Introduction

Loss of proteostasis is a well-established feature of aging[1]. Insoluble protein aggregates accumulate during the normal aging process across eukaryote species, from yeast to mice[2–10]. In *Caenorhabditis elegans*, insoluble protein accumulation is strongly associated with lifespan[5]. Chemical compounds that slow the accumulation of insoluble proteins extend lifespan[11]; whereas a diet rich in iron accelerates protein insolubility and shortens lifespan[12]. In humans, protein insolubility and dysregulation of proteostasis are central phenomena of all major, age-related neurodegenerative diseases, despite the canonical protein associated with aggregation being different in each case eg. Tau & Amyloid β (Aβ) (Alzheimer's disease), α-synuclein (Parkinson's disease) and Huntingtin (Huntington's disease)[13–19]. In the context of Alzheimer's Disease (AD), senile plaques from patient brains contain hundreds of insoluble proteins in addition to Aβ. These proteins range from immune factors, such as complement proteins, to proteins involved in maintaining proteostasis including molecular chaperones, proteasomal subunits, and autophagy machinery[20–23]. Critically, many of these proteins have been found to aggregate due to normal aging in *C. elegans*[5,6,10,24,25]. While aging is the dominant risk factor for AD, therapeutic approaches have thus far failed to consider aging or generalized protein insolubility as causal factors[26–29]; perhaps contributing to fact that approved AD therapies provide modest improvements[30–32].

Molecular and genetic evidence implicate several aging hallmarks in the etiology of AD, such as inflammation and mitochondrial dysfunction[33–37]. However, the primary histopathological hallmarks of the disease are the accumulation of insoluble protein aggregates rich in Aβ and neurofibrillary tangles of aggregated, hyper-phosphorylated Tau[38–40]. Genetic, biochemical, and animal model data all support a central role for amyloid beta (Aβ), and in particular the $Aβ_{1-42}$ peptide, in the formation of aggregates and neurodegeneration[41–50]. However, Positron Emission Tomography (PET) studies measuring Aβ accumulation in the brains of cognitively healthy individuals have proven that Aβ deposition is not sufficient to cause dementia[51]. Intriguingly, insoluble protein extracts, specifically from old animals, significantly accelerate the aggregation of Aβ *in vitro*, suggesting a biophysical relationship between Aβ and the aging insoluble proteome[52]. We questioned whether the reverse is also true. What if Aβ accumulation accelerates the insolubility of proteins that typically insolubilize during aging? Could it be that the age-dependent insoluble proteome interacts with Aβ in a vicious feedback loop, leading to an acceleration of insolubility in AD? Moreover, if we understand which proteins are affected by Aβ in this way, could we uncover a novel mechanism through which Aβ exerts toxicity?

Here, using an unbiased proteomic approach[53], we tested the effect of Aβ expression on protein insolubility using *C. elegans*. We demonstrate that Aβ expression drives a dramatic increase in proteome-wide protein insolubility. We discovered that the Aβ-

driven insoluble proteome is highly similar to the insoluble proteome that forms due to normal aging, showing a fundamental link between AD and aging. By analyzing human genome-wide association studies (GWAS), we show that the proteins driven to become insoluble by aging or Aβ represent biological processes implicated across not only neurodegenerative diseases but also diverse chronic, age-related diseases (CARDs), providing suggestive evidence that protein insolubility could play a role in general CARD risk. Taken together, our findings provide new insights into Aβ toxicity in AD and highlight the importance of considering the insoluble proteome in not only AD but also in other age-related diseases.

Results

Aβ expression drives proteome-wide protein insolubility resembling normal aging.

To determine the effect of Aβ expression on proteostasis we utilized the *C. elegans* strain, GMC101, which expresses the human, pathogenic Aβ$_{1-42}$, peptide in muscle tissue, here referred to as simply Aβ[47,54]. When cultured at 20ºC, worms express Aβ at low levels and when shifted 25ºC overexpress Aβ. After 24 hours of exposure, the majority of worms paralyze[55]. We allowed the worms to develop in the absence of Aβ expression and then shifted to 25ºC at day 1 of adulthood. Worms were then collected after 24 hours of exposure to Aβ. To determine the effect of Aβ on protein aggregation, we extracted insoluble proteins by serial washing worm lysates with 1% Sodium Dodecyl Sulfate (SDS) buffer. We used mass spectrometry and data independent acquisitions (DIA) to identify and quantify proteins, as previously described[53,56,57] (Fig.1A). Background genotype-control worms (CL2122) lacking the Aβ transgene were cultured and processed in parallel for comparison (Fig. 1A).

Strikingly, Aβ expression in young adult animals caused a robust increase in the total amount of insoluble protein (Fig.1 B&C). We identified and quantified peptides representing 1704 proteins in the insoluble fraction across four biological replicates, of which 593 proteins robustly increased due to Aβ expression across independent experiments (Fig. S1A). We observed a proteome-wide increase in insolubility, impacting a wide array of essential functions (Fig S1C). Several protein homeostasis machineries were highly enriched in the insoluble proteome, including heat shock proteins (HSPs) from most major cell compartments including the mitochondria and endoplasmic reticulum, the proteasome regulatory subunit, the TriC chaperonin complex, and lysosomal proteins (Fig S1C). Furthermore, we observed an increase in protein insolubility of the ribosomes and translation accessory factors, the mitochondrial electron transport chain (ETC), the tricarboxylic acid (TCA) cycle, the nuclear laminar and core histones, Golgi COPI vesicles and endoplasmic reticulum, cytoskeleton, and extracellular matrix (ECM), as well as dozens of metabolic enzymes spanning, glycolysis, lipid transport and catabolism, one carbon, glycogen, and steroid metabolism. Gene Ontology (GO) and pathway enrichment analysis revealed that the insoluble proteome was enriched with proteins involved in oxidative phosphorylation (Ox. Phos), Ribosomes, the mitochondrial unfolded protein response (mitoUPR), and "Determination of lifespan" (Fig.1 D, Supp. Table 2). Specifically, 42 of the 70 proteins in the *C. elegans* proteome bearing the annotation "Determination of Lifespan" became insoluble due to

Aβ (Fig. 1D). Using STRING clustering, we found that these 42 proteins form a functionally related PPI network, connecting protein folding machineries with translation and oxidative phosphorylation (STRING, p<10-6) (Fig. 1E).

The Aβ insoluble proteome was highly enriched with mitochondrial proteins (109/593); particularly those involved in the ETC and TCA cycle (Fig. S1C, Supp. Table 2). Consistent with this, hsp-6 increased 9.2-fold in the insoluble fraction and the mitochondrial unfolded protein response (mitoUPR) was amongst the most significantly enriched pathways in the insoluble proteome (Fig. 1D, Supp. Table 2). 43 of the 88 ETC proteins became insoluble after Aβ expression (ETC complexes detailed in Supp. Table 3). Highly expressed, nuclear-encoded ETC complexes, were particularly vulnerable to insolubility (Fig. S1C). Notably, this was not accompanied by insolubility of the mitochondrial DNA-encoded ETC complex proteins, with only one mtDNA-encoded, complex IV protein becoming insoluble (Supp. Table 3). The skewed representation of nuclear-encoded subunits could be explained by the fact that we observed a significant increase in insolubility of mitochondrial outer membrane proteins. All major proteins of the TOM complex: TOM-20, TOM-22, TOM-40, and TOM-70 became insoluble. We also identified GOP-3, the orthologue of the outer membrane complex protein SAMM50[61], which is required for threading of mitochondrial β-barrel proteins into the outer and inner membrane bilayers[62], along with several small molecule transporters: VDAC-1, ANT-1.1 and MTCH-1 (Fig S1C). It is interesting to note that single nucleotide polymorphisms (SNPs) in several TOM complex genes have been linked with AD risk, most prominently *TOMM40*[58–60] (Fig. S1C). Taken together these data provide *in vivo* evidence in a multicellular organism that Aβ drives insolubility of dozens of mitochondrial proteins.

Normal aging causes a significant increase in the level of insoluble protein and we noted many of the same proteins that typically become insoluble during normal aging in the Aβ driven insoluble proteome. This led us to question the extent of the similarity between the two. To assess this rigorously, we generated a list of proteins that robustly insolubilize during normal aging by overlapping two published aging insoluble proteomes from different laboratories[6,53]. This resulted in an overlap of 457 proteins that reliably become insoluble during aging (p <0.0001, Fischer's exact test) (Fig. 1G). When we compared this with the Aβ insoluble proteome, we found that 66%, or 305 proteins, became insoluble under both conditions (p < 0.0001, Fischer's exact test) (Fig. 1G). Moreover, when we compared the GO Biological Processes (BPs) represented in the aging and Aβ insoluble proteomes, we uncovered an 89% overlap, indicating that aging and Aβ drive insolubility of proteins involved in almost identical biological processes (Fig. 1H). The fact that two thirds of proteins that become insoluble during normal aging also do so under Aβ expression suggested to us that we identified a core set of vulnerable proteins that become insoluble under stress conditions, which we refer to herein as the 'Core Insoluble Proteome' (CIP).

We suspected from previous work[5] that the CIP might be particularly enriched with regulators of aging. Indeed, we found that the CIP was highly enriched with modulators of lifespan. Specifically, 100 proteins, or roughly one third of the CIP, modulate lifespan according to the GenAge database (Fig.2A)[62]; with 80% of those proteins extending lifespan when their expression is reduced or eliminated (Fig. 2B). Furthermore, using published whole-genome RNA interference (RNAi) screens[63–67], we found that knock down of approximately one in six of the proteins in the CIP has been shown to improve disease pathology across Huntington Disease (HD) or Parkinson Disease (PD)

proteinopathy models, suggesting that the CIP might be enriched with regulators of neurodegeneration (Supp Table 4). This led us to directly question if the orthologues of CIP proteins had been implicated in neurodegenerative proteinopathies in humans. To do so, we performed disease pathway enrichment analysis with the human orthologues of the CIP and found a highly significant enrichment for a range of neurodegenerative diseases including prion disease (Pr), AD, PD and HD (Fig. 2C). In almost all case, the proteins were annotated against all four neurodegenerative diseases, suggesting a common mechanism might underly their common association. Our data point towards protein insolubility under proteotoxic stress conditions as the common factor (Supp. Table 5).

Protein insolubility impacts biological processes that increase human age-related disease risk.

In addition to bearing annotations for neurodegenerative disease pathways, we observed that many insoluble proteins bore pathway annotations for non-neurological, age-related diseases, such as non-alcoholic fatty liver disease and diabetic cardiomyopathy (Supp. Table 7B). This led us to question if the relationship between the insoluble proteome and chronic age-related disease (CARD) could go beyond simply neurodegeneration. Previous GWAS analysis demonstrated that common biological processes underly a whole array of diverse CARDs[68]. We therefore applied this analysis to query the commonality between biological processes impacted by protein insolubility and biological processes implicated in CARD risk[68] (Fig. 2D). We compared biological processes implicated in risk for developing 38 distinct CARDs against the biological processes impacted by insolubility in the CIP. Strikingly, we found that the average protein in the CIP was found to share biological processes with 18 of the 38 different CARDs (Fig. 2E). Moreover, 88% of the CIP proteins shared annotations with diseases spanning four or more of the following five CARD categories: neurodegenerative, metabolic, cancer, cardiovascular, and 'other' (capturing disparate CARDs such as macular degeneration, rheumatoid arthritis, and osteoporosis) (Fig. 2E&F). These associations were relatively enriched compared with the experimental background insoluble proteome (ie. all proteins that could be identified in the insoluble fraction across all experimental conditions) and highly enriched compared with proteins in the non-insoluble background proteome, which on average shared biological process annotations with zero CARDs (Fig 2E). It is worth noting that *TOMM40*, which we identified in the Aβ-driven insoluble proteome and two independent aging-driven insoluble proteome datasets[11,53], is one of only a very small fraction of SNPs (2.5%) that increase disease risk for 3 or more broad classes of CARD[68].

The biological processes common to the insoluble proteome and all 5 broad CARD categories included: immune activation and stress response pathways; growth signaling, such as fibroblast growth factor (FGF) and WNT signaling; RNA splicing and regulation of expression; and tissue homeostatic processes important for development and wound healing, such as ECM organization, cell migration and cell differentiation (Fig. S2A). Except for a few rare cases, soluble proteins lose their biological function when they become insoluble and therefore these data provide suggestive evidence that insolubility of a core set of vulnerable proteins might promote risk of not only neurodegenerative disease but CARDs more broadly.

The CIP can be used to identify therapeutic targets.

We speculated that insoluble proteins could be playing a direct role in the toxicity of Aβ in this *C. elegans* model and that reducing the expression of the insoluble proteins might be beneficial. To test this, we knocked down the expression of CIP-encoding genes using RNAi and measured paralysis. We prioritized testing known aging modifiers, as defined by the GO BP determination of lifespan. We found that, of the 23 genes knocked down, 12 significantly impacted disease pathology, eight had no impact, and three were either lethal or significantly delayed development and so could not be tested. Therefore, of the CIP genes we were able to test, 60% significantly impacted Aβ toxicity (Fig 3A), suggesting an enrichment for disease-modifying proteins in the CIP. Contrary to our initial hypothesis, however, most of these genes (7/12) resulted in exacerbation of paralysis rather than protection; suggesting that knocking down expression of some proteins could be undermining a compensatory response to their becoming insoluble due to Aβ.

Since we observed strong evidence of mitochondrial protein unfolding in the insoluble proteome, we hypothesized that directly targeting mitochondrial protein homeostasis with a small molecule might represent a good strategy to ameliorate amyloid toxicity. Urolithin A (UA) is a mitophagy-inducing, natural product derived from the bacterial metabolism of foods containing ellagitannins, such as pomegranate seeds and walnuts, in the gastrointestinal tract [69–71], and has been shown to improve muscle function in a phase II clinical trial for age-related frailty [72,73]. We treated Aβ-expressing worms with UA for 18 hours prior to inducing Aβ expression and then scored paralysis after 24 hours. We measured a robust and significant decrease in the percentage of paralyzed worms at 24 hours (Fig 3B). Taken with recent Aβ mouse model data, this supports that targeting mitochondrial protein homeostasis by inducing mitophagy represents a reasonable strategy for preventing or treating Aβ toxicity [74].

Discussion

In this study we used an unbiased proteomics approach to assess the impact of Aβ on protein insolubility. We discovered that Aβ expression is sufficient to cause proteome-wide protein insolubility and in particular unfolding of the mitochondrial ETC and membrane transporters. We discovered that the Aβ-driven insoluble proteome bears a striking overlap with the aging-driven insoluble proteome, suggesting there exists a core sub-proteome, vulnerable to insolubility under stress conditions. This core insoluble proteome was remarkably enriched for regulators of lifespan and disease pathology in *C. elegans*; with a particularly strong enrichment for regulators of neurodegenerative proteinopathies. Using analysis of GWAS data [68] we uncovered suggestive evidence that protein insolubility could link diverse CARDs by the disruption of common biological processes. This finding is compelling considering the geroscience hypothesis, which "posits that aging physiology plays a major role in many — if not all — chronic diseases" [75]. While purely correlative at this stage, it leads us to hypothesize that insolubility could be a contributing factor in the etiology of CARDs more generally.

The Aβ-driven insoluble proteome data is consistent with evidence in yeast and human cells that Aβ, and specifically the $Aβ_{1-42}$ peptide, interferes with the import of mitochondrial protein precursors into the mitochondria by directly interacting with the TOM proteins [76–79]. The present study provides the first evidence that outer membrane

proteins themselves become insoluble due to Aβ, including GOP-3 (SAMM50) of the SAM complex which has not been implicated before. This is consistent with our observation that the insoluble proteome is highly enriched with nuclear-encoded but not mitochondrial-encoded ETC complex proteins. Further, we found that targeting mitochondrial protein homeostasis with an established mitophagy-inducing compound, Urolithin A, was efficacious in preventing Aβ toxicity which may explain its observed therapeutic effects in Aβ-expressing pre-clinical mouse models[74]. These data clearly support the notion that Aβ disrupts mitochondrial function through driving loss of mitochondrial protein homeostasis which warrants further attention in the development of AD therapeutics.

Here we demonstrated that insoluble proteomics can be used to identify genetic modifiers of Aβ toxicity. We show that reducing the levels of specific proteins in the insoluble proteome can either ameliorate or exacerbate Aβ toxicity in *C. elegans*. We observed counter-intuitive effects which could point towards specific disease-modifying mechanisms for future research. Firstly, knock down of the large ribosomal protein RPL4 caused a robust suppression of paralysis. This was anticipated because we assumed this would reduce global translation and thus reduce the overall demand on the proteostasis machinery. However, knock down of the translation elongation factor EEF-2, which should have similarly decreased global translation, had the opposite effect, resulting in a profound increase in paralysis. Similarly, we also observed a counter-intuitive result when targeting insoluble mitochondrial proteins. Knocking down the mitochondrial ATP synthase peripheral stalk protein, ASB-2, for example was found to significantly delay paralysis. In contrast, knocking down the mitochondrial membrane transporter ANT-1.1 or the aconitase enzyme ACO-2 led to significantly worse paralysis. RNAi against ASB-2 and other subunits of the ATP synthase complex are known to increase lifespan[80,81], possibly through a mito-hormetic response, and this could explain the protective effects observed in our model. However, ANT-1.1 and ACO-2 knock down have also been shown to increase lifespan by several independent groups[5,81–84], demonstrating a more complex picture. While these data could point to a compensation response, we speculate that certain insoluble proteins might serve a cytoprotective role in sequestering misfolded proteins or, in this case, toxic Aβ monomers, into inert, insoluble aggregates. Knock down of other members of the insoluble proteome might initiate a protective response at the level of transcription or protein homeostasis machinery.

Over 50 biological processes with known links with age-related disease risk were shown here to be linked with insolubility during normal aging (Fig S2). These data suggest that clinicians and scientists working on CARD mechanisms could benefit from considering protein insolubility when studying the interplay between aging and their disease of interest; particularly of proteins implicated in this study. We propose that evidence exists to support this approach in thinking about therapeutic strategies for age-related diseases; for example, the small molecule, HBX, which slows aging by increasing protein homeostasis in invertebrate models had unexpected, positive effects on bone health during aging in the laboratory mouse[85]. Based on the discovery of a core insoluble sub proteome and the processes implicated in CARD risk, targeting the insoluble proteome provides an encompassing strategy for the prevention and treatment of disparate age-related diseases.

Experimental Methods

Animal Strains

The following strains were used to generate the proteomics data: GMC101 (dvIs100 [*unc-54*p::A-beta-1-42::*unc-54* 3'-UTR + *mtl-2p*::GFP]), CL2122 (dvIs15 [(pPD30.38) *unc-54*(vector) + (pCL26) *mtl-2*::GFP]). GMC101 was used to test Urolithin A for protection against paralysis.

The following *C. elegans* strain was generated in for this study, GL399 (dvIs100 [*unc-54*p::A-beta-1-42::*unc-54* 3'-UTR; *spe-9*(hc88) I; *rrf-3*(b26) II.), by crossing TJ1060(*spe-9*(hc88) I; *rrf-3*(b26) II) with GMC101 (dvIs100 [*unc-54p*::A-beta-1-42::*unc-54* 3'-UTR + *mtl-2p*::GFP]). GL399 was used for all RNAi paralysis assays.

Animal Maintenance

GMC101 and CL2122 worms were maintained at 20ºC on 60mm Nematode Growth Media (NGM) agar plates seeded with OP50 *E. coli*. Worms were maintained by transferring 30-50 eggs to a fresh plate on Monday and Friday. GL399 was maintained at 15ºC on 60mm NGM agar plates seeded with OP50 *E. coli*. Worms were maintained by transferring 30-50 eggs to a fresh plate weekly.

Urolithin A Paralysis Assay

A 20mM stock of Urolithin A (UA) was prepared in sterile DMSO and stored in aliquots at -20ºC. From the stock solution, 130μL of the working solution (50μM) was prepared by mixing 7.5μL of stock solution (or DMSO only for control plates) with 125.5μL of distilled sterile water and was added to the top 35mm NGM plates (3 mL NGM agar) pre-seeded with OP50 *E. coli*. A population of synchronized GMC101 worms was generated by allowing Day 2 adult worms to lay eggs for 2 hours at 20ºC on 60mm NGM agar plates seeded with OP50. Progeny were allowed to develop to larval stage 4 (L4) and then transferred to NGM plates overlaid with 50μM UA (or DMSO only for control plates, 0.05% DMSO). After 18 hours, these plates were moved to 25ºC to initiate Aβ expression. Worms were scored for paralysis after a further 24 hours. To score paralysis in an unbiased way, worms were moved to one quadrant of the plate and after 3-5 minutes scored as paralyzed if they were unable to moved away.

RNA Interference Paralysis Assay

A population of synchronized GL399 worms was generated by allowing Day 2 adult worms to lay eggs for 2 hours at 20ºC on 60mm NGM agar plates seeded with HT115 *E. coli* expressing the scrambled interfering RNA, or control vector RNAi. Progeny were allowed to develop to for 48 hours and then transferred to NGM plates seeded HT115 *E. coli* expressing either the scrambled control vector RNAi or RNAi against the gene of interest, and immediately shifted to 25ºC to initiate Aβ expression. Worms were scored for paralysis after 50 hours and 72 hours in order to identify RNAi conditions that advance or delay paralysis, respectively. To score paralysis in an unbiased way, worms were moved to one quadrant of the plate and after 5 minutes scored as paralyzed if they

were unable to moved away. Graphed results reflect the summarized findings of the two timepoints.

Insoluble Protein Extraction

Isolation of SDS-insoluble proteins from worms was completed as described in detail in[53]. Briefly, 200 day 2 adult worms were allowed to lay eggs for 5 hours on 100mm NGM agar plates seeded with 4x concentrated OP50 *E.coli*. After 50-52 hours worms were collected in sterile S-basal solution (5.85 g NaCl, 1 g K_2HPO_4, 6 g KH_2PO_4, 1 ml cholesterol (5 mg/ml in ethanol) per 1L sterile H_2O) and transferred to fresh 4x OP50 seeded plates. At 72 hours post egglay worm plates were transferred to 25ºC to initiate Aβ expression. 24h hours later worms were collected in sterile S-basal. The worms were allowed to settle under gravity and the supernatant solution was removed. This step ensured larval worms would not be taken forward for insoluble protein isolation. The worms were washed several times to remove all bacteria and larvae and then the supernatant was removed and the tubes flash-frozen in dry ice/ethanol.

Worm pellets were thawed in the presence of detergent-free lysis buffer containing protease inhibitor (20 mM Tris base, pH 7.4, 100 mM NaCl, 1 mM $MgCl_2$). Thawed pellets were then sonicated in a water bath kept at 4ºC on maximum intensity for 30 s ON and 30 s OFF to prevent sample overheating. After the first round of sonication for 10 cycles (10 min) pellets were checked under the light microscope to ensure worms were efficiently lysed. Lysates were clarified by centrifugation at 3,000g for 4 minutes. Supernatant was collected and total protein quantified by BCA.

Insoluble proteins were extracted by aliquoting lysate equivalent to 2mg of total protein and spinning at 20,000g for 15 minutes at 4ºC. The supernatant was collected as the aqueous-soluble fraction. The insoluble protein pellet was resolubilized in 1%(w/v) SDS lysis buffer (20 mM Tris base, pH 7.4, 100 mM NaCl, 1 mM $MgCl_2$ plus protease inhibitor) and centrifuged at 20,000g for 15 minutes at room temperature. The supernatant was collected as the 1% SDS-soluble fraction. The pellet was then resolubilized and washed twice more in 1% SDS lysis buffer and the supernatant fractions kept. The remaining SDS-insoluble pellet was resolubilized in sterile 70% (v/v) formic acid and sonication in a water bath for 30 minutes. Samples were then dried in a vacuum concentrator for 1 hour to completely remove formic acid. 1 x LDS sample gel buffer was added to the dried pellet and the samples were heated to 95ºC for 10 minutes. Samples were briefly centrifuged to collect condensation and stored at -80ºC or run onto a 4-12% NUPAGE Bis-Tris gel.

Mass Spectrometric Acquisitions

Dried pellets were dissolved in 40 µL of 1x LDS sample gel buffer, incubated at 95 °C for 10 min, vortexed, and spun down. Solubilized samples were run in pre-cast NuPAGE 4-12% gradient acrylamide Bis-Tris protein gels (Invitrogen) for 20 minutes to concentrate the proteins in a single band in the stacking gel. For in-gel digestion, the gel bands were diced, collected in tubes, and dehydrated with a dehydration buffer (25 mM ammonium bicarbonate in 50% acetonitrile (ACN) and water). The gel samples were dried in a vacuum concentrator, reduced with 10 mM dithiothreitol (DTT) in 25 mM ammonium bicarbonate (pH 7-8) and incubated for 1 hour at 56 ºC with agitation, then alkylated with 55 mM iodoacetamide (IAA) in 25 mM ammonium bicarbonate (pH 7-8), and incubated for 45

minutes at room temperature in the dark. The diced gel pieces were washed with 25 mM ammonium bicarbonate in water (pH 7-8), dehydrated again with the dehydration buffer, and dried in a vacuum concentrator. Afterwards, 250 ng of sequencing-grade trypsin in 25 mM ammonium bicarbonate (pH 7-8) was added to each sample. Gel pieces were vortexed for 10 min, briefly spun, and incubated at 4 ºC for 30 min without mixing. Gel pieces were covered with 25 mM ammonium bicarbonate (pH 7-8) and incubated overnight for 16-20 h at 37 ºC with agitation. Subsequently, the digested peptides were further extracted, as gel pieces were subjected to water and then two separate additions of a solution containing 50% ACN, 5% formic acid (FA) in water. After each addition of solution, the sample was mixed for 10 minutes, and then the aqueous digests from each sample were transferred into a new tube. These pooled peptide extractions were dried in a vacuum concentrator until completely dry. Proteolytic peptides were re-suspended in 30 µL of 0.2% FA in water and desalted using stage-tips made in-house containing a C_{18} disk, concentrated in a vacuum concentrator, and re-suspended in 15 µL of 0.2% FA in water and 1 µL of indexed Retention Time Standard (iRT, Biognosys, Schlieren, Switzerland).

Samples were then subjected to mass spectrometric analysis using a high-performance liquid chromatography (HPLC) system combined with a chip-based HPLC system (Eksigent nano-LC) directly connected to a quadrupole time-of-flight mass spectrometer (TripleTOF 5600, a QqTOF instrument) as detailed in our previous step-by-step protocol[53].

DIA/SWATH Data Processing and Statistical Analysis

Each sample was acquired in data-dependent acquisition (DDA) mode to build peptide spectral libraries, as we previously described[53]. Data-Independent Acquisition (DIA)/SWATH data was processed in Spectronaut (version 12.0.20491.3.15243) using DIA. Data extraction parameters were set as dynamic and non-linear iRT calibration with precision iRT was selected. DIA data was matched against an in-house *Caenorhabditis elegans* spectral library that provides quantitative DIA assays for 5,461 control and 10,140 GMC 101 *C*. elegans peptides corresponding to 1,223 protein groups for control and 1,839 protein groups for GMC 101 and supplemented with scrambled decoys (library size fraction of 0.01), using dynamic mass tolerances and dynamic extraction windows. DIA/SWATH data was processed for relative quantification comparing peptide peak areas from different days. Identification was performed using 1% precursor and protein q-value. Quantification was based on the peak areas of extracted ion chromatograms (XICs) of 3 – 6 MS2 fragment ions, specifically b- and y-ions, with q-value sparse data filtering and iRT profiling being applied. For this sample-set, local normalization was not implemented. Differential protein expression analyses for all comparisons were performed using a paired t-test, and p-values were corrected for multiple testing, using the Storey method. Specifically, group wise testing corrections were applied to obtain q-values. Protein groups with at least two unique peptides, q-value < 0.01, and absolute Log_2(fold-change) > 0.58 are significantly altered (Supp. Tables 1,11A,11B).

Data Accession

Raw data and complete MS data sets have been uploaded to the Mass Spectrometry Interactive Virtual Environment (MassIVE) repository, developed by the Center for Computational Mass Spectrometry at the University of California San Diego, and can be

Computational Methods

Enrichment analysis: All enrichment analyses were performed using significantly increased proteins and gProfiler[86] with Benjamini-Hochberg FDR correction. Enrichment results are presented as the product of Gene Count and FDR-corrected p value. Human orthologs were derived from Ortholist[61].

GWAS analysis: All SNP GO biological process data for the present study was taken from Johnson et al.[68]. All GO biological process data for the insoluble proteins was downloaded from DAVID[87,88] and Uniprot[89]. For comparison between GO biological processes enriched in CARDs with those represented in the insoluble proteome and CIP(the overlap between the Aβ-driven and aging-driven insoluble proteome), we determined overlaps between GO term annotations for the proteins and the CARDs. We used these overlaps then to determine the shared biological processes in the CIP and insoluble proteome with the five broad age-related disease categories as defined in Johnson et al.[68]. The "insoluble proteome" was defined as any protein which could be identified in the insoluble fraction in either of the experiments ie. the experimental background. For comparison with the proteome writ large, we generated a random list of 1600 proteins from the *C. elegans* proteome using a random number generator and downloaded their GO biological process data from DAVID and Uniprot. GO biological processes shared between the CIP and all 5 diverse categories of CARD were processed through ReviGO[90] to remove any redundant terms and create a relationship network between the processes, which we subsequently modified in Cytoscape[91] and presented in Fig.S2.

Acknowledgments

GJL, BS & JKA disclose support for the research of this work from NIA RF1AG057358 and GJL discloses support for the research of this work from The Larry L. Hillblom Medical Foundation Lithgow LLHF Ctr Supp. and NIA U01AG045844. BS acknowledges support of instrumentation for the TripleTOF 6600 from the NIH shared instrumentation grant 1S10 OD016281 (Buck Institute).

Special thanks to Dr Simon Johnson for providing source data from[68].

COI Statement

The authors declare no conflict of interest.

References

1. López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. Hallmarks of aging: An expanding universe. *Cell* **186**, 243–278 (2023).

2. Demontis, F. & Perrimon, N. FOXO/4E-BP signaling in Drosophila muscles regulates organism-wide proteostasis during aging. *Cell* **143**, 813–825 (2010).

3. Erjavec, N., Cvijovic, M., Klipp, E. & Nyström, T. Selective benefits of damage partitioning in unicellular systems and its effects on aging. *Proc Natl Acad Sci U S A* **105**, 18764–18769 (2008).

4. Saarikangas, J. & Barral, Y. Protein aggregates are associated with replicative aging without compromising protein quality control. *Elife* **4**, (2015).

5. Reis-Rodrigues, P. *et al.* Proteomic analysis of age-dependent changes in protein solubility identifies genes that modulate lifespan. *Aging Cell* **11**, 120–127 (2012).

6. David, D. C. *et al.* Widespread protein aggregation as an inherent part of aging in C. elegans. *PLoS Biol* **8**, 47–48 (2010).

7. Sacramento, E. K. *et al.* Reduced proteasome activity in the aging brain results in ribosome stoichiometry loss and aggregation. *Mol Syst Biol* **16**, 9596 (2020).

8. Tanase, M. *et al.* Role of carbonyl modifications on aging-associated protein aggregation. *Sci Rep* **6**, 1–14 (2016).

9. Vonk, W. I. M. *et al.* Differentiation Drives Widespread Rewiring of the Neural Stem Cell Chaperone Network. *Mol Cell* **78**, 329-345.e9 (2020).

10. Walther, D. M. *et al.* Widespread proteome remodeling and aggregation in aging C. elegans. *Cell* **161**, 919–932 (2015).

11. Mark, K. A., Dumas, K. J., Ramanathan, A., Gibson, B. W. & Lithgow, G. J. Vitamin D Promotes Protein Homeostasis and Longevity via the Stress Response Pathway Genes skn-1, ire-1, and xbp-1. *CellReports* **17**, 1227–1237 (2016).

12. Klang, I. M. *et al.* Iron promotes protein insolubility and aging in C. elegans. *Aging* **6**, 975–991 (2014).

13. Taylor, J. P., Hardy, J. & Fischbeck, K. H. Toxic proteins in neurodegenerative disease. *Science (1979)* **296**, 1991–1995 (2002).

14. Bates, G. Huntingtin aggregation and toxicity in Huntington's disease. *Lancet* **361**, 1642–1644 (2003).

15. Berke, S. J. S. & Paulson, H. L. Protein aggregation and the ubiquitin proteasome pathway: gaining the UPPer hand on neurodegeneration. *Curr. Opin. Genet. Dev.* **13**, 253–261 (2003).

16. Dawson, T. M. & Dawson, V. L. Rare genetic mutations shed light on the pathogenesis of Parkinson disease. *J. Clin. Invest* **111**, 145–151 (2003).

17.   Goedert, M. Tau protein and neurodegeneration. *Semin. Cell Dev. Biol.* **15**, 45–49 (2004).

18.   Ross, C. A. & Poirier, M. A. Protein aggregation and neurodegenerative disease. *Nature Medicine 2004 10:7* **10**, S10–S17 (2004).

19.   Bruijn, L. I. *et al.* Aggregation and motor neuron toxicity of an ALS-linked SOD1 mutant independent from wild-type SOD1. *Science (1979)* **281**, 1851–1854 (1998).

20.   Drummond, E. *et al.* The amyloid plaque proteome in early onset Alzheimer's disease and Down syndrome. *Acta Neuropathol Commun* **10**, (2022).

21.   Drummond, E. *et al.* Proteomic differences in amyloid plaques in rapidly progressive and sporadic Alzheimer's disease. *Acta Neuropathol* **133**, 933–954 (2017).

22.   Nijholt, D. A. T., Stingl, C. & Luider, T. M. Laser capture microdissection of fluorescently labeled amyloid plaques from alzheimer's disease brain tissue for mass spectrometric analysis. *Methods in Molecular Biology* **1243**, 165–173 (2014).

23.   Drummond, E., Nayak, S., Pires, G., Ueberheide, B. & Wisniewski, T. Isolation of Amyloid Plaques and Neurofibrillary Tangles from Archived Alzheimer's Disease Tissue Using Laser-Capture Microdissection for Downstream Proteomics. *Methods Mol Biol* **1723**, 319–334 (2018).

24.   Olzscha, H. *et al.* Amyloid-like aggregates sequester numerous metastable proteins with essential cellular functions. *Cell* **144**, 67–78 (2011).

25.   Vecchi, G. *et al.* Proteome-wide observation of the phenomenon of life on the edge of solubility. *Proc Natl Acad Sci U S A* **117**, 1015–1020 (2020).

26.   Brookmeyer, R., Johnson, E., Ziegler-Graham, K. & Arrighi, H. M. Forecasting the global burden of Alzheimer's disease. *Alzheimers Dement* **3**, 186–191 (2007).

27.   Brookmeyer, R. *et al.* National estimates of the prevalence of Alzheimer's disease in the United States. *Alzheimer's & Dementia* **7**, 61–73 (2011).

28.   Padmanabhan, P. & Götz, J. Clinical relevance of animal models in aging-related dementia research. *Nature Aging 2023 3:5* **3**, 481–493 (2023).

29.   The transforming landscape of dementia research. *Nature Aging 2023 3:5* **3**, 459–459 (2023).

30.   Alexander, G. C., Emerson, S. & Kesselheim, A. S. Evaluation of Aducanumab for Alzheimer Disease: Scientific Evidence and Regulatory Review Involving Efficacy, Safety, and Futility. *JAMA - Journal of the American Medical Association* **325**, 1717–1718 (2021).

31.   Tampi, R. R., Forester, B. P. & Agronin, M. Aducanumab: evidence from clinical trial data and controversies. *Drugs Context* **10**, (2021).

32.   van Dyck, C. H. *et al.* Lecanemab in Early Alzheimer's Disease. *New England Journal of Medicine* **388**, 9–21 (2022).

33. Heppner, F. L., Ransohoff, R. M. & Becher, B. Immune attack: the role of inflammation in Alzheimer disease. *Nature Reviews Neuroscience 2015 16:6* **16**, 358–372 (2015).

34. Selkoe, D. J. Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev* **81**, 741–766 (2001).

35. Wang, W., Zhao, F., Ma, X., Perry, G. & Zhu, X. Mitochondria dysfunction in the pathogenesis of Alzheimer's disease: recent advances. *Molecular Neurodegeneration 2020 15:1* **15**, 1–22 (2020).

36. Butterfield, D. A. & Halliwell, B. Oxidative stress, dysfunctional glucose metabolism and Alzheimer disease. *Nature Reviews Neuroscience 2019 20:3* **20**, 148–160 (2019).

37. Welikovitch, L. A. *et al.* Early intraneuronal amyloid triggers neuron-derived inflammatory signaling in APP transgenic rats and human brain. *Proc Natl Acad Sci U S A* **117**, 6844–6854 (2020).

38. Stelzmann, R. A., Norman Schnitzlein, H. & Reed Murtagh, F. An English translation of Alzheimer's 1907 paper, 'Uber eine eigenartige Erkankung der Hirnrinde'. *Clin Anat* **8**, 429–431 (1995).

39. Braak, H. & Braak, E. Neurofibrillary changes confined to the entorhinal region and an abundance of cortical amyloid in cases of presenile and senile dementia. *Acta Neuropathol* **80**, 479–486 (1990).

40. Braak, H. & Braak, E. Neuropathological stageing of Alzheimer-related changes. *Acta Neuropathol* **82**, 239–259 (1991).

41. Rovelet-Lecrux, A. *et al.* APP locus duplication causes autosomal dominant early-onset Alzheimer disease with cerebral amyloid angiopathy. *Nat Genet* **38**, 24–26 (2006).

42. Kelleher, R. J. & Shen, J. Presenilin-1 mutations and Alzheimer's disease. *Proc Natl Acad Sci U S A* **114**, 629 (2017).

43. Hardy, J. & Selkoe, D. J. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* **297**, 353–356 (2002).

44. Casas, C. *et al.* Massive CA1/2 neuronal loss with intraneuronal and N-terminal truncated Abeta42 accumulation in a novel Alzheimer transgenic model. *Am J Pathol* **165**, 1289–1300 (2004).

45. LaFerla, F. M., Green, K. N. & Oddo, S. Intracellular amyloid-beta in Alzheimer's disease. *Nat Rev Neurosci* **8**, 499–509 (2007).

46. Billings, L. M., Oddo, S., Green, K. N., McGaugh, J. L. & LaFerla, F. M. Intraneuronal Abeta causes the onset of early Alzheimer's disease-related cognitive deficits in transgenic mice. *Neuron* **45**, 675–688 (2005).

47. McGowan, E. *et al.* Abeta42 is essential for parenchymal and vascular amyloid deposition in mice. *Neuron* **47**, 191–199 (2005).

48.　Meyer-Luehmann, M. *et al.* Rapid appearance and local toxicity of amyloid-beta plaques in a mouse model of Alzheimer's disease. *Nature* **451**, 720–724 (2008).

49.　Meyer-Luehmann, M. *et al.* Extracellular amyloid formation and associated pathology in neural grafts. *Nat Neurosci* **6**, 370–377 (2003).

50.　Wang, J., Dickson, D. W., Trojanowski, J. Q. & Lee, V. M. Y. The levels of soluble versus insoluble brain Abeta distinguish Alzheimer's disease from normal and pathologic aging. *Exp Neurol* **158**, 328–337 (1999).

51.　Chételat, G. *et al.* Amyloid imaging in cognitively normal individuals, at-risk populations and preclinical Alzheimer's disease. *Neuroimage Clin* **2**, 356 (2013).

52.　Groh, N. *et al.* Age-dependent protein aggregation initiates amyloid-β aggregation. *Front Aging Neurosci* **9**, 138 (2017).

53.　Xie, X. *et al.* Quantification of insoluble protein aggregation in caenorhabditis elegans during aging with a novel data-independent acquisition workflow. *Journal of Visualized Experiments* **2020**, 1–17 (2020).

54.　McColl, G. *et al.* Utility of an improved model of amyloid-beta (Aβ1-42) toxicity in Caenorhabditis elegans for drug screening for Alzheimer's disease. *Mol Neurodegener* **7**, 57 (2012).

55.　Mccoll, G. *et al.* Utility of an improved model of amyloid-beta (Aβ1-42) toxicity in Caenorhabditis elegans for drug screening for Alzheimer's disease. *Mol Neurodegener* **7**, 1–9 (2012).

56.　Collins, B. C. *et al.* Multi-laboratory assessment of reproducibility, qualitative and quantitative performance of SWATH-mass spectrometry. *Nature Communications 2017 8:1* **8**, 1–12 (2017).

57.　Gillet, L. C. *et al.* Targeted Data Extraction of the MS/MS Spectra Generated by Data-independent Acquisition: A New Concept for Consistent and Accurate Proteome Analysis. *Molecular & Cellular Proteomics* **11**, O111.016717 (2012).

58.　Valant, V. *et al.* TOMM40 in Cerebral Amyloid Angiopathy Related Intracerebral Hemorrhage: Comparative Genetic Analysis with Alzheimer's Disease. *Transl Stroke Res* **3**, 102–112 (2012).

59.　Martin, E. R. *et al.* SNPing away at complex diseases: analysis of single-nucleotide polymorphisms around APOE in Alzheimer disease. *Am J Hum Genet* **67**, 383–394 (2000).

60.　Omoumi, A. *et al.* Evaluation of late-onset Alzheimer disease genetic susceptibility risks in a Canadian population. *Neurobiol Aging* **35**, 936.e5-936.e12 (2014).

61.　Kim, W., Underwood, R. S., Greenwald, I. & Shaye, D. D. OrthoList 2: A New Comparative Genomic Analysis of Human and Caenorhabditis elegans Genes. *Genetics* **210**, 445–461 (2018).

62. Ott, C. *et al.* Sam50 Functions in Mitochondrial Intermembrane Space Bridging and Biogenesis of Respiratory Complexes. *Mol Cell Biol* **32**, 1173–1188 (2012).

63. Kraemer, B. C., Burgess, J. K., Chen, J. H., Thomas, J. H. & Schellenberg, G. D. Molecular pathways that influence human tau-induced pathology in Caenorhabditis elegans. *Hum Mol Genet* **15**, 1483–1496 (2006).

64. Hamamichi, S. *et al.* Hypothesis-based RNAi screening identifies neuroprotective genes in a Parkinson's disease model. *Proc Natl Acad Sci U S A* **105**, 728–733 (2008).

65. Van Ham, T. J. *et al.* C. elegans Model Identifies Genetic Modifiers of α-Synuclein Inclusion Formation During Aging. *PLoS Genet* **4**, e1000027 (2008).

66. Silva, M. C. *et al.* A Genetic Screening Strategy Identifies Novel Regulators of the Proteostasis Network. *PLoS Genet* **7**, 1002438 (2011).

67. Nollen, E. A. A. *et al.* Genome-wide RNA interference screen identifies previously undescribed regulators of polyglutamine aggregation. *Proc Natl Acad Sci U S A* **101**, 6403–6408 (2004).

68. Johnson, S. C., Dong, X., Vijg, J. & Suh, Y. Genetic evidence for common pathways in human age-related diseases. *Aging Cell* **14**, 809–817 (2015).

69. Cerdá, B., Periago, P., Espín, J. C. & Tomás-Barberán, F. A. Identification of urolithin A as a metabolite produced by human colon microflora from ellagic acid and related compounds. *J Agric Food Chem* **53**, 5571–5576 (2005).

70. D'Amico, D. *et al.* Impact of the Natural Compound Urolithin A on Health, Disease, and Aging. *Trends Mol Med* **27**, 687–699 (2021).

71. D'Amico, D. *et al.* Urolithin A improves mitochondrial health, reduces cartilage degeneration, and alleviates pain in osteoarthritis. *Aging Cell* **21**, (2022).

72. Liu, S. *et al.* Effect of Urolithin A Supplementation on Muscle Endurance and Mitochondrial Health in Older Adults: A Randomized Clinical Trial. *JAMA Netw Open* **5**, e2144279–e2144279 (2022).

73. Andreux, P. A. *et al.* The mitophagy activator urolithin A is safe and induces a molecular signature of improved mitochondrial and cellular health in humans. *Nat Metab* **1**, 595–603 (2019).

74. Ballesteros-Álvarez, J., Nguyen, W., Sivapatham, R., Rane, A. & Andersen, J. K. Urolithin A reduces amyloid-beta load and improves cognitive deficits uncorrelated with plaque burden in a mouse model of Alzheimer's disease. *Geroscience* **45**, 1095–1113 (2023).

75. What is Geroscience? - American Federation for Aging Research. https://www.afar.org/what-is-geroscience.

76. Mossmann, D. *et al.* Amyloid-β peptide induces mitochondrial dysfunction by inhibition of preprotein maturation. *Cell Metab* **20**, 662–669 (2014).

77.   Cenini, G., Rub, C., Bruderek, M. & Voos, W. A Highlights from MBoC Selection: Amyloid β-peptides interfere with mitochondrial preprotein import competence by a coaggregation process. *Mol Biol Cell* **27**, 3257 (2016).

78.   Hansson Petersen, C. A. *et al.* The amyloid β-peptide is imported into mitochondria via the TOM import machinery and localized to mitochondrial cristae. *Proc Natl Acad Sci U S A* **105**, 13145–13150 (2008).

79.   Hu, W., Wang, Z. & Zheng, H. Mitochondrial accumulation of amyloid β (Aβ) peptides requires TOMM22 as a main Aβ receptor in yeast. *J Biol Chem* **293**, 12681 (2018).

80.   Xu, C. *et al.* Genetic inhibition of an ATP synthase subunit extends lifespan in C. elegans. *Scientific Reports 2018 8:1* **8**, 1–14 (2018).

81.   Lee, S. S. *et al.* A systematic RNAi screen identifies a critical role for mitochondria in C. elegans longevity. *Nat Genet* **33**, 40–48 (2003).

82.   Kim, Y. & Sun, H. Functional genomic approach to identify novel genes involved in the regulation of oxidative stress resistance and animal lifespan. *Aging Cell* **6**, 489–503 (2007).

83.   Hamilton, B. *et al.* A systematic RNAi screen for longevity genes in C. elegans. *Genes Dev* **19**, 1544–1555 (2005).

84.   Dong, M. Q. *et al.* Quantitative mass spectrometry identifies insulin signaling targets in C. elegans. *Science* **317**, 660–663 (2007).

85.   Evans, D. S. *et al.* Longitudinal Functional Study of Murine Aging: A Resource for Future Study Designs. *JBMR Plus* **5**, (2021).

86.   Raudvere, U. *et al.* g:Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic Acids Res* **47**, W191–W198 (2019).

87.   Huang, D. W. *et al.* DAVID Bioinformatics Resources: Expanded annotation database and novel algorithms to better extract biology from large gene lists. *Nucleic Acids Res* **35**, (2007).

88.   Huang, D. W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44–57 (2009).

89.   Bateman, A. *et al.* UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic Acids Res* **51**, D523–D531 (2023).

90.   Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO Summarizes and Visualizes Long Lists of Gene Ontology Terms. *PLoS One* **6**, e21800 (2011).

91.   Shannon, P. *et al.* Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. *Genome Res* **13**, 2498 (2003).

Figure 1: Aβ drives proteome-wide protein insolubility that resembles normal aging

Aβ drives proteome-wide protein insolubility that resembles normal aging. A. Schematic of experimental procedure. **B.** Total insoluble protein Intensity in worms expressing Aβ vs genotype control strain CL2122, all values normalized to the average of the control in each experiment n = 4, error bars = SEM, Mann-Whitney Test. **C.** Representative volcano plot from one experiment Absolute $\text{Log}_2(\text{Fold Change})>0.58$, Q <0.01 **D.** Barplot of selected enriched GO terms, KEGG pathways and Wikipathways in the Aβ insoluble proteome, Benjamini Hochberg FDR <5% (see supplementary tables for complete list) **E.** STRING protein-protein interaction network of insoluble proteins with annotations for "Determination of Lifespan", average fold change and Q values across both experiments, STRING p < 1x10[-16]. **F.** left: **F.** Overlap of proteins enriched in the insoluble proteome of old *C. elegans* from two publicly available datasets[43,49] p<0.0001 Fischer's Exact Test. right: Overlap of aging insoluble proteome with Aβ insoluble proteome, p<0.0001 Fischer's Exact Test. **G.** Overlap of all GO annotations from the aging insoluble proteome and Aβ insoluble proteome, p<0.0001 Fischer's Exact Test.

Figure 2: The Core Insoluble Proteome is enriched with modulators of aging and age-related disease

The Core Insoluble Proteome is enriched with modulators of aging and age-related disease A. Overlap of lifespan modulating proteins from GenAge database with the core insoluble proteome, p<0.0001 Fischer's Exact Test **B.** Barplot summary of published lifespan experiments where the 100 overlapping genes were either mutated or knocked down. **C.** Barplot highlighting enriched GO terms for neurodegenerative diseases using human orthologs of CIP proteins, Benjamini Hochberg, FDR <1%. **D.** Schematic representation of how GWAS analysis was used to compare GO Biological Processes shared across diverse chronic, age-related diseases (CARDs) with the biological processes represented in the CIP **E.** Box and whiskers plot shows distribution of the number of ARDs that share biological processes with proteins found in CIP, the insoluble proteome, and the proteome. The median value is displayed above. **F.** Frequency distribution of the number of broad CARD classes that share one or more biological processes with proteins found in the CIP, the insoluble proteome, and the proteome.

Figure 3: Targeting the insoluble proteome genetically or pharmacologically modulates Aβ toxicity

Targeting the insoluble proteome genetically or pharmacologically modulates Aβ toxicity A. Statistically significant hits from RNAi paralysis screen against CIP proteins. Each point represents the relative proportion of paralyzed worms on that plate normalized to the average paralysis on control vector for that experiment, approx. 40 worms per plate, 2-3 separate experiments, error bars = SEM, for statistical tests see Supplementary Table 9A, all p<0.05 with Bonferroni correction. **B.** Barplot of paralysis under 50μM UA treatment, 40 worms per plate, 3 experiment replicates, replicate experiments are indicated by shape, error bars = SEM, unpaired t test.

Supplementary Figure S1

A. Overlap of proteins which increased in abundance in the insoluble proteome after Aβ expression. **B.** Overlap of proteins which increased due to Aβ expression across both experiments and proteins identified in the aging-driven insoluble proteome in either of the two published datasets. **C.** Schematic overview of proteostasis and mitochondrial proteins enriched in the Aβ-driven insoluble proteome

Supplementary Figure S2

GO Biological Processes shared between the core insoluble proteome and all 5 ARD classes

A. ReviGo network representation of the non-redundant GO biological processes shared between the CIP and all 5 diverse CARD categories.