***Tbr1* autism mouse model displays altered structural and functional**

amygdalar connectivity and abnormal whole-brain synchronization

Tsan-Ting Hsu[1], Chien-Yao Wang[2] and Yi-Ping Hsueh[1]*

[1]Institute of Molecular Biology, Academia Sinica, Taipei, Taiwan, ROC

[2]Institute of Information Science, Academia Sinica, Taipei, Taiwan, ROC

* Corresponding to Dr. Yi-Ping Hsueh, Institute of Molecular Biology, Academia Sinica, 128,

Sec 2, Academia Rd, Taipei, 11529, Taiwan, ROC. Email: yph@gate.sinica.edu.tw

Running title: Whole-brain connectivity in *Tbr1*$^{+/-}$ autism model.

Keywords: Axonal projection, Contralateral long-range connectivity, Autism spectrum

disorders, Basolateral amygdala, Default mode network, Structural and functional

connectivity, Whole-brain connectivity.

Abstract

Autism spectrum disorders (ASD) are recognized as neural disconnectivity syndromes. Here, we establish a whole-brain immunostaining and quantification platform to investigate structural and functional connectivity in a $Tbr1^{+/-}$ autism mouse model, which shares a defect in the anterior commissure that is evolutionarily conserved with human patients. Since the basolateral amygdala (BLA) is particularly susceptible to $Tbr1$ haploinsufficiency and it projects to the contralateral brain hemisphere via the anterior commissure, we express a channelrhodopsin variant oChIEF fused with Citrine at the BLA to outline axonal projections of BLA neurons and to activate the BLA under blue light theta-burst stimulation (TBS). Next, we evaluate C-FOS expression to represent neural activity. We show that $Tbr1$ haploinsufficiency almost completely disrupts contralateral BLA axonal projections and results in ipsilateral mistargeting, thereby globally altering BLA functional connectivity. Based on correlated C-FOS expression among brain regions, we further report that $Tbr1$ deficiency severely disrupts whole-brain synchronization in the absence of salient stimulation. $Tbr1^{+/-}$ and wild-type mice exhibit opposing responses to TBS-induced amygdalar activation, with it reducing synchronization in wild-type mice but enhancing it in $Tbr1^{+/-}$ mice. Whole-brain modular organization and inter-module connectivity are also affected by Tbr1 deficiency and amygdalar activation. In fact, the synchronization and intra- and inter-modular connectivities of TBS-treated $Tbr1^{+/-}$ mice are more comparable to those of non-treated wild-type mice, suggesting a potential ameliorating effect of amygdalar stimulation on brain function. Moreover, $Tbr1^{+/-}$ mice exhibit attenuated connectivity between the amygdala and default mode network, a subnetwork highly relevant to social behaviors, strengthening the link between the impaired connectivity and social behavior deficits displayed by $Tbr1^{+/-}$ mice. Our high-resolution analytical platform reveals the inter- and intra-hemispheric connectopathies arising from an ASD condition and emphasizes the defective synchronization at a whole-brain scale caused by $Tbr1$ deficiency.

Introduction

Autism spectrum disorders (ASD), one of the most highly prevalent neurodevelopmental disorders, are characterized by two core behavioral deficits that appear in early childhood; one is impaired social interaction and communication, and the other is restricted, repetitive behavioral patterns and sensory abnormalities (https://dsm.psychiatryonline.org/doi/book/10.1176/appi.books.9780890425787). Seeking the biological mechanisms causing these core behavioral manifestations is crucial for providing a better understanding of ASD and uncovering potential treatments. One of the primary biological features correlated with behavioral phenotypes in ASD patients is altered brain connectivity [1-4], although the detailed features of rewiring alterations under ASD conditions remain elusive.

T-Brain-1 (TBR1), a neuron-specific T-box transcription factor [5], has been identified as a causative gene of ASD (Gene score 1, https://gene.sfari.org/database/human-gene/TBR1) [6, 7]. TBR1 regulates the expression of a set of genes controlling axonal projections of cortical and amygdalar neurons [7, 8]. Contralateral projection of basolateral amygdalae (BLA) neurons via the posterior part of the anterior commissure is particularly sensitive to *Tbr1* deficiency [7], with this latter eliciting anterior commissure impairment in both humans and mice [7, 9-11]. Moreover, *Tbr1* haploinsufficiency, a condition mimicking genetic variations observed in patients with ASD, results in many autism-like behavioral abnormalities in mice, including reduced social behaviors, defective vocalization, impaired olfactory discrimination, aversive memory and cognitive inflexibility [7, 8, 12]. Disconnection of the anterior commissure in mice either surgically or via unilateral chemogenetic inhibition of the BLA results in autism-like behaviors [13, 14]. Together, these studies have demonstrated that *Tbr1*$^{+/-}$ mice may serve as an excellent model for exploring the connectivity deficits associated with ASD.

Given that BLA axonal projection is particularly sensitive to *Tbr1* haploinsufficiency [7, 13] and that BLA activation by local infusion of D-cycloserine, a co-agonist of NMDAR, is sufficient to ameliorate the autism-like behaviors exhibited by *Tbr1*$^{+/-}$ mice [7], we speculate that BLA-derived circuits are critical to *TBR1*-linked ASD phenotypes. Here, we established a whole-brain analytical platform using serial mouse brain sections for immunostaining and high-content imaging and for performing registration and quantification based on the Allen Mouse Common Coordinate Framework version 3 (CCFv3) [15]. Using this system, we analyzed axonal projection of the BLA at the mesoscale level and assessed neuronal activity based on C-FOS expression at a cellular level. Our high-resolution analyses revealed the inter- and intra-hemispheric connectopathy in *Tbr1*$^{+/-}$ mice, providing clear evidence that an ASD-linked condition results in dyssynchronization at the whole-brain scale.

Results

Development of a mesoscopic whole-brain scale circuit analysis pipeline

To analyze structural and functional BLA connectivity in individual mice, we unilaterally expressed the ultra-fast channelrhodopsin variant oChIEF fused with the fluorescent protein Citrine in BLA neurons of both $Tbr1^{+/-}$ and wild-type mice. Given that oChIEF-Citrine targets the plasma membrane of entire neurons, in addition to opto-stimulating BLA neurons, oChIEF-Citrine was also used to outline BLA neurons, including axons. The relative fluorescent signal intensity of oChIEF-Critine outside of the BLA reflects the extent of axonal projections derived from unilateral BLA neurons. To analyze neuronal activity and functional connectivity of the BLA, $Tbr1^{+/-}$ and wild-type mice were further separated into two groups. One group received optical theta-burst stimulation (TBS) at the oChIEF-expressing BLA when they were freely moving in their home cages. The other control group was subjected to the same experimental procedure except for optical stimulation. Then, we analyzed neuronal activation by BLA stimulation based on C-FOS expression 2 hr after stimulation. Our whole-brain scale analysis and quantification basically comprise five elements, with an outline and detailed procedure being presented in **Figures 1, S1** and the **Methods**. The list of brain regions used in our study and their corresponding abbreviations can be found in **Tables S1** and **S2**. Brain ontology is identical to CCFv3.

Structure-wise comparisons reveal rewiring in $Tbr1^{+/-}$ mouse brains

First, we analyzed the whole-brain distribution pattern of oChIEF-Citrine. We found that oChIEF-Citrine signals were widely distributed ipsilaterally in wild-type brains and also extended contralaterally (**Figure 2A**; **Video S1**, upper). However, oChIEF-Citrine signals barely extended contralaterally in $Tbr1^{+/-}$ mouse brains (**Figure 2a**; **Video S1**, lower).

We compared brains from a total of 15 $Tbr1^{+/-}$ mice and 13 wild-type littermates. Since the BLA consists of heterologous projections with varying patterns of long-range axonal projection [16], first we investigated the distributions of the virus injection sites between wild-type and $Tbr1^{+/-}$ mouse brains and confirmed that they were indistinguishable and comparable (**Figure S2**). Next, we binned and summarized the average percentage of oChIEF-Citrine signals in all brain regions of $Tbr1^{+/-}$ mice and wild-type littermates. These results were subjected to 3D projection at both coronal and horizontal views, which indicated a reduction in contralateral projections of BLA neurons in $Tbr1^{+/-}$ mice (**Figure 2B**; **Video S2**).

Moreover, we used three different methods to identify brain regions susceptible to *Tbr1*

deficiency. The first method was a structure-wise comparison between *Tbr1*$^{+/-}$ mice and wild-

type littermates at the whole-brain scale. This volume-based quantification was performed on

the ipsilateral, i.e., the virus injection site, and contralateral hemispheres separately (**Figure**

S3A, Table S1). Consistent with our previous findings [7, 13], we found that oChIEF-Citrine

signals were greatly reduced in contralateral BLA-related regions (LA, BLA, BLAa, BLAp),

and in the associated fiber tracts (act, amc), of *Tbr1*$^{+/-}$ mice compared to those of wild-type

littermates (**Figures 2C-2E**), supporting the reliability of our whole-brain analysis system.

In addition to the amygdala and its associated fiber tracts, we found that additional 26

brain regions at the contralateral site had much lower oChIEF-Citrine signals in *Tbr1*$^{+/-}$ mice

relative to wild- type (**Figures 2C-2E**). We term these regions "affected contralateral projection

regions (ACPR)", which mainly reside in areas of the forebrain, such as the isocortex, olfactory

area (OLF), cortical subplate (CTXsp) and cerebral nuclei (CNU). In wild-type mice, these

regions actually receive topographically bilateral projections from the two amygdalae in the

two brain hemispheres (**Figures 2A-2B,** upper panel; also see Allen Mouse Brain Connectivity

Atlas: connectivity.brain-map.org/projection/experiment/113144533   and

connectivity.brain.map.org/projection/experiment/277710753). At the ipsilateral site, 18 brain

regions were identified as exhibiting increased oChIEF-Citrine signals in *Tbr1*$^{+/-}$ mouse brains

(**Figures 2C-2E**), which we denote "affected ipsilateral projection regions (AIPR)". Note that

these AIPR are not the counterparts of the ACPR (**Figures 2C-2E**). These findings indicate that

Tbr1 haploinsufficiency impairs the contralateral projections of BLA neurons and leads to

mistargeting to ipsilateral brain regions, and not necessarily limited to ACPR counterparts. We

also noticed that AIPR encompass several regions in the cortex, thalamus (TH), hypothalamus

(HY) and midbrain (MB) (**Figures 2C-2E**), implying that *Tbr1* haploinsufficiency may elicit

stronger BLA ipsilateral connections with those regions.

We selected 44 contralateral and 50 ipsilateral brain regions to manually verify the

regions of interest (ROI) and to correct mismatched ROI for every single brain slice image of

each brain sample, confirming our quantification results (**Figure 2E**, the regions labeled with

red font).

***Tbr1* deficiency alters the ratio of contralateral to ipsilateral projections**

The second method we applied was to measure the ratio of Citrine signals in paired regions of

the two brain hemispheres, i.e., the contralateral/ipsilateral (C/I) ratio, in individual mice. The

fold-change in C/I ratios between *Tbr1*$^{+/-}$ mice and WT littermates were then analyzed, which

revealed that *Tbr1*[+/−] mice exhibited reduced C/I ratios in all identified regions (**Figure S3B**).

Given that all ACPR exhibited reduced Citrine signals, their C/I ratios were consequently

reduced (**Figure S3C**). In addition, three AIPR presented lower C/I ratios because of their

stronger Citrine signals at the ipsilateral site (**Figure S3D**). Moreover, 11 additional brain

regions in *Tbr1*[+/−] mice exhibited a reduced C/I ratio due to differential expression of the Citrine

levels between the two brain hemispheres (**Figure S3E**). This analysis of C/I ratios further

confirms the altered bilateral axonal projections elicited by *Tbr1* deficiency.

Principal component analysis also reveals altered rewiring in *Tbr1*[+/−] mouse brains

The third method was to analyze BLA projections using principal component analysis (PCA).

For the contralateral site, the PCA analysis showed clear segregation between wild-type and

Tbr1[+/−] mice along PC2 (**Figures S4A-S4B**; PC2 KNN classification accuracy 86%, k = 4, *P* =

0) with the ACPR identified in **Fig. 1** contributing noticeably to that axis. In fact, the top 10%

of regions contributing to PC2 were all ACPR (**Figures S4C-S4D**). At the ipsilateral site,

although our PCA analysis revealed strong and significant segregation between the two

genotypes along PC1 (**Figures S4E-S4F**; PC1 KNN classification accuracy 75%, k = 5, *P* =

0.013), there were no correlations between PC1 loading and AIPR (**Figures S4G-S4H**),

implying that many BLA ipsilateral targeting regions exhibit greater inter-sample variation than

the difference between genotypes and supporting that some ipsilateral projections tend to be

random and that their innervations are less specific.

Taken together, using multiple different comparative approaches, we analyzed BLA-

derived circuits in *Tbr1*[+/−] mice at a whole-brain scale and uncovered aberrant rewiring caused

by *Tbr1* haploinsufficiency. The imbalanced topographic BLA bilateral axonal projections

account for the major rewiring differences, with some enhanced BLA ipsilateral projections to

some areas of the cortex, thalamus, hypothalamus and midbrain also being displayed by the

Tbr1[+/−] mice.

***Tbr1* deficiency changes neuronal activation**

Next, we investigated if the altered BLA projections in *Tbr1*[+/−] mice influence their neuronal

activity patterns. To do so, we compared the C-FOS expression of ACPR and AIPR across four

experimental groups, i.e., WT ctrl, *Tbr1*[+/−] ctrl, WT TBS and *Tbr1*[+/−] TBS. Using volume-based

quantification, the C-FOS expression patterns in all examined brain regions were

indistinguishable among the four groups (**Figure S5A**, **Table S1**). However, when we

visualized the distribution of C-FOS[+] cells after binning and averaging of a 3D mouse brain

map, it was clear that the distribution of C-FOS$^+$ cells varied among these four groups of mice (**Figure S5B**). Since a given brain region usually makes specific connections with multiple brain regions via specific subregions, the unchanged subregions may dilute any differences and result in indistinguishable C-FOS expression levels in a volume-based quantification. Therefore, we established slice-based quantification to analyze the numbers of C-FOS$^+$ cells along the anterior-posterior (AP) axis with 100 μm binning of both brain hemispheres (**Figure 3A; Figure S6A**), which revealed differential neuronal activation between *Tbr1*$^{+/-}$ mice and wild-type littermates (**Figure S6B**). Cumulative probability distributions indicated that the majority of brain regions indeed displayed comparable numbers of C-FOS$^+$ cells among the four experimental groups for both brain hemispheres (**Figures 3A, 3B**). However, compared to wild-type groups, *Tbr1*$^{+/-}$ mice had more C-FOS$^+$ cells in a small proportion (~15%) of regions at the contralateral site with or without TBS treatment (**Figure 3B**, upper). At the ipsilateral site, only the WT ctrl group had fewer C-FOS$^+$ cells than the other three groups (**Figure 3B**, lower). Thus, although only a few differentially activated brain regions exist, our slice-based quantification approach adequately identified those regions among our experimental groups.

Consequently, we used this slice-based quantification approach to conduct pairwise comparisons of our four experimental groups of mice to establish which brain regions exhibited differential C-FOS expression (**Figure 3C**, **Table S2**). For the comparison of *Tbr1*$^{+/-}$ ctrl vs. WT ctrl, we uncovered 105 and 93 regions having more C-FOS$^+$ cells in the contralateral and ipsilateral sides of the brains of the *Tbr1*$^{+/-}$ ctrl group, respectively. In contrast, only 9 regions displayed more C-FOS$^+$ cells in the WT ctrl (4 in the contralateral and 5 in the ipsilateral hemispheres) (**Figure 3C**, left). These differentially activated regions are widely distributed across the brain, from the isocortex to MB, suggesting that the abnormally high brain activities of *Tbr1*$^{+/-}$ mice during wakefulness (the state of our mice under experimental/control treatment) can be linked to circuits and areas distributed across the entire brain (**Figure 3C**, left).

We also analyzed the contributions of BLA activation to *Tbr1* phenotypes. In wild-type mice, TBS at the BLA increased the number of C-FOS$^+$ cells across different brain regions of both the contralateral and ipsilateral sites (**Figure 3C**, middle left). Thus, unilateral activation of the BLA in wild-type mice indeed activates many other brain regions, both contralaterally and ipsilaterally. In contrast, *Tbr1*$^{+/-}$ mice unexpectedly displayed more brain regions with fewer C-FOS$^+$ cells upon TBS (**Figure 3C**, middle right). These results indicate that *Tbr1* haploinsufficiency alters global functional connectivity between the BLA and other brain regions. We also detected differing C-FOS$^+$ cell distributions in many brain regions between

214 the *Tbr1*[+/−] TBS and WT TBS groups (**Figure 3C**, right). Taken together, these results imply

215 that *Tbr1* deficiency induces functional rewiring.

Correlations between rewired BLA-derived circuits and altered C-FOS expression patterns

To further explore the correlation between axonal projection and neural activation, we

conducted slice-based quantification of oChIEF-Citrine signals as per our analysis of C-FOS[+]

cell distribution patterns. Compared to our volume-based quantification results, we found that

more brain structures exhibited differential innervation patterns of BLA-derived axons in

Tbr1[+/−] mice according to slice-based analysis, regardless of contralateral or ipsilateral

projection and increased or decreased innervation patterns (**Figures S7A-S7C, Table S2**).

However, the general conclusion of BLA-derived circuit rewiring in *Tbr1*[+/−] mice remains, i.e.,

that impaired contralateral axonal projections and enhanced axonal innervations in the

ipsilateral hemisphere.

Next, we selected the brain regions exhibiting rewired axonal innervations and altered

C-FOS expression patterns upon TBS of the BLA in either wild-type or *Tbr1*[+/−] mice for

correlation analysis. First, we determined ΔoChIEF traces by pairwise subtraction of oChIEF-

Citrine signals between *Tbr1*[+/−] and wild-type samples (**Figures 4A**, **4C,** left, 10μm³ binning

cubes). Neural activation induced by TBS in *Tbr1*[+/−] mice and WT littermates, i.e., *Tbr1*[+/−] ΔC-

FOS traces and WT ΔC-FOS traces respectively, was also determined. Pairwise subtraction

between *Tbr1*[+/−] ΔC-FOS and WT ΔC-FOS traces was then executed to obtain all possible

differences, i.e., ΔΔC-FOS traces (**Figures 4A**, **4C,** middle, 100 μm³ binning cubes). A Pearson

correlation coefficient (r) was calculated according to the correlation of the averaged ΔoChIEF

trace and the averaged ΔΔC-FOS trace in each brain region. The brain regions exhibiting

positive correlations between rewired BLA-derived circuits and altered C-FOS expression

patterns reflect concurrent changes to BLA-derived axonal innervations and their neural activity

upon BLA activation. In the contralateral hemisphere of *Tbr1*[+/−] mice, the AUD regions (AUD,

AUDp), AIp, PIR, BLA and MBsen regions all exhibited impaired BLA contralateral axonal

projections and the boosting effects of BLA activation on C-FOS expression were inhibited,

indicating a loss of intensification effect of contralateral BLA axonal projections to these

regions in the mutant mice (**Figures 4A-4B**). The contralateral thalamus, DORsm regions

(DORsm, VENT) and ILM each displayed higher oChIEF-Citrine signals in *Tbr1*[+/−] mice, but

they exhibited negative correlations between ΔoChIEF and ΔΔC-FOS (**Figures 4A-4B**).

In the ipsilateral hemisphere, the correlation results between ΔoChIEF and ΔΔC-FOS

were the opposite of those for the contralateral hemisphere (**Figures 4A-4B** vs. **Figures 4C-**

4D). Only for the posterior part (ventral) of the HIP and NOT were ipsilateral BLA axonal

projections reduced in *Tbr1*$^{+/-}$ mice, and it exhibited impaired boosting effects of BLA

activation on C-FOS expression (**Figures 4C-4D, left panel**). A major group of regions at the

ipsilateral side exhibited a higher level of oChIEF-Citrine in *Tbr1*$^{+/-}$ mice but a negative

correlation between ΔoChIEF and ΔΔC-FOS (**Figures 4C-4D**). This group contains the

isocortical area (MOp, AUDd, ACA, ACAv, TEa), as well as the CNU (MEA, PALv, SI, PALc,

BST), TH (DORsm, DORpm, CL), HY (LZ) and MB (SCdg, VTA) areas.

Taken together, these findings indicate that structural rewiring of BLA-derived circuits

in *Tbr1*$^{+/-}$ mice triggers altered functional consequences for downstream targets. In general,

more contralateral target areas receive fewer BLA axonal projections that cause diminished

potentiation of neuronal activities, whereas more ipsilateral downstream target regions are

innervated by more BLA axons that induce net inhibitory effects on neuronal activities.

Altered functional connectivity of *Tbr1*$^{+/-}$ mice

In addition to examining the alterations to individual structures elicited by *Tbr1*

haploinsufficiency, we further assessed the functional changes at the network level. We

evaluated the functional connectivity of paired brain regions by determining their correlated C-

FOS expression levels across all subjects [17, 18]. We then computed an inter-regional

correlation matrix of C-FOS levels across all subjects for each experimental group. The inter-

regional correlation coefficient of C-FOS levels, r, was considered meaningful if its significant

correlation level *P* was <0.05, otherwise r was set to zero to reflect no correlation. For all inter-

regional pairs displaying a significant correlation, we identified large proportions of inter-

regional pairs with positive correlations and comparable positive r values among the four

experimental groups of WT ctrl, *Tbr1*$^{+/-}$ ctrl, WT TBS and *Tbr1*$^{+/-}$ TBS, yet very few but

variable inter-regional pairs exhibiting negative correlations in the WT ctrl (1.52%), *Tbr1*$^{+/-}$ ctrl

(2%), WT TBS (2.36%) and *Tbr1*$^{+/-}$ TBS (0.03%) groups (**Figure 5A**). Upon examining the

entire inter-regional correlation matrix, we observed that meaningful functional connections

between brain regions in *Tbr1*$^{+/-}$ mice in the awake state were noticeably reduced compared to

their WT littermates (**Figure 5B**, WT ctrl vs. *Tbr1*$^{+/-}$ ctrl). After TBS, inter-regional functional

connections were noticeably reduced in the wild-type group (**Figure 5B**, WT ctrl vs. WT TBS),

whereas they were dramatically increased in *Tbr1*$^{+/-}$ mice (**Figure 5B, *Tbr1*$^{+/-}$ ctrl vs. *Tbr1*$^{+/-}$**

TBS). Therefore, the opposing consequences caused by BLA activation in WT and *Tbr1*$^{+/-}$ mice

are not only reflected in neuronal recruitment of individual structures, but they are also

manifested by inter-regional functional connectivity.

Next, we quantified the degree centrality (DC) of each brain region based on our inter-

regional C-FOS correlation data (**Figure 5C**). A higher DC for a specific region indicates that

the activity of that region is synchronous with more brain regions. Our results show that the

WT ctrl group had a higher DC degree than the $Tbr1^{+/-}$ ctrl group. However, BLA activation

generally reduced DC degree in WT mice, but increased it in $Tbr1^{+/-}$ mice (**Figure 5C**). Thus,

the WT ctrl group was more synchronous than the $Tbr1^{+/-}$ ctrl group, and TBS at the BLA has

opposing effects in this respect on WT and $Tbr1^{+/-}$ mice.

In order to confirm that the differences in DC were not caused by the particular

correlation criteria we used, we evaluated the network stability of the four experimental groups

by increasing the r (correlation coefficient) or P (correlation significance) threshold used for

constructing the inter-regional correlation network (**Figure 5D**). Compared to the WT ctrl

network, the reduced mean DC of the $Tbr1^{+/-}$ ctrl network still existed across all the absolute r

and large range of P threshold conditions. Moreover, the contrasting modulatory effects of

BLA-derived circuit activation on DC between the wild-type and $Tbr1^{+/-}$ networks also existed

when all the absolute r and large range of P value thresholds were used for network construction

(**Figure 5D**). These results support that the differences in DC among the four experimental

groups are unbiased.

Differential modular organization and reorganization upon BLA activation

To understand which brain regions are more densely interconnected than others and to achieve

better modular integration, we used a Louvain community detection algorithm [19] to analyze

modular organization. Optimal communities were defined by C-FOS levels of the brain regions

being more correlated to each other within a community than between communities. We used

a spatial resolution parameter series, R, ranging from 0.4 to 0.9, in Louvain community

detection and calculated the difference in the modularity metric (Q) for communities

representing real networks or shuffled networks. The appropriate R-value for each network was

determined according to the maximal difference between Q and $Q_{shuffled}$ (**Figure 6A**).

Louvain community detection revealed distinct modular organizations for the four

experimental groups. The WT ctrl network contained 14 communities, with the majority of

brain regions occupying the first 3 communities (598/618 brain regions). The remaining

communities of the WT ctrl network comprised a few inter-correlated regions (communities 4,

6, 8, 9) or single brain regions (communities 5, 7, 10, 11, 12, 13, 14) (**Figures 6B, 6C**). In

contrast, the *Tbr1*[+/−] ctrl network only comprised four communities (**Figures 6B, 6C**). Indeed, it was clear that none of the communities identified in the WT ctrl group were identical to those of the *Tbr1*[+/−] ctrl group, no matter whether brain regions or mean correlation within or between communities was being considered (**Figures 6B, 6C, Figure S7D, Table S3**).

Under BLA activation, the communities in the WT and *Tbr1*[+/−] mice were substantially reorganized. All major communities of the WT TBS network (communities 1–3, total 613/618 regions) displayed weaker mean correlations within and between communities relative to the first three communities of the WT ctrl network (**Figures 6B, 6C**, WT TBS communities 1-3 vs. WT ctrl communities 1-3). For the *Tbr1*[+/−] TBS network, the brain regions were highly concentrated in two communities, i.e., communities 1 and 3 (**Figures 6B, 6C**), with strong mean within and between community correlations. Note that community 1 of *Tbr1*[+/−] TBS network is similar to community 2 of WT ctrl network (**Figure 6C**), suggesting that amygdala activation may reorganize neural network of *Tbr1*[+/−] mice to be more comparable to that in WT mice.

Overall, this community analysis suggests that Tbr1 haploinsufficiency reorganizes how the mouse brain is networked and also leads to differential connectivity upon BLA activation.

***Tbr1* deficiency and BLA activation influence the default mode network**

Finally, we applied our dataset to a physiologically relevant subnetwork analysis to explore any differences among experimental groups. Our control groups reflect a cognitive state where the mice were in their home cages during wakefulness and in the absence of any salient stimuli. Therefore, we adopted the default mode network (DMN) as the most appropriate for our analysis since it was initially identified by resting-state fMRI (functional magnetic resonance imaging) as a set of brain regions with correlated activities in the absence of goal-directed tasks and inactivation during task execution [20]. Since the DMN has been implied in social behaviors and ASD [21, 22], DMN analysis likely provide more understanding about the deficits in *Tbr1*[+/−] mice.

The core DMN regions in the mouse brain are located in the isocortical prefrontal (ACAd, ACAv, PL, ILA, ORBl, ORBm, ORBvl), somatomotor (SSp-tr, SSp-ll, MOs) and medial (VISa, VISam, RSPagl, RSPd, RSPv) areas of both brain hemispheres [23]. We extracted our data on oChIEF patterns, C-FOS expression and functional connectivity for these core DMN regions and compared them among our experimental groups. To aid visualization, the summarized signals in 10 and 100 mm^3 cubes were used to highlight differences in oChIEF-Citrine and C-FOS expression, respectively.

Among the DMN regions, $Tbr1^{+/-}$ mice exhibited more ipsilateral BLA projections, especially in the ventral part of the ipsilateral ACA (i_ACAv), compared to wild-type mice (**Figure 7A**, DMN ΔoChIEF), implying a role for TBR1 in controlling BLA axonal projections into the DMN. For C-FOS expression, we observed more cubes with higher C-FOS$^+$ cell numbers in bilateral DMN regions for the $Tbr1^{+/-}$ ctrl vs. WT ctrl and WT TBS vs. WT ctrl comparisons (**Figure 7B**). In contrast, there were more cubes with reduced numbers of C-FOS$^+$ cells for the $Tbr1^{+/-}$ TBS vs. $Tbr1^{+/-}$ Ctrl and $Tbr1^{+/-}$ TBS vs. WT TBS comparisons (**Figure 7B**). Thus, DMN activity is indeed sensitive to $Tbr1$ deficiency and BLA activation.

We also determined the DC and participation coefficient (PC) of all DMN regions. PC measures the relative distribution of a given brain region's (node) correlations (edges) with other communities versus the correlations within its own community. Many regions of the $Tbr1^{+/-}$ ctrl DMN exhibited reduced DC, and only the ipsilateral ACAv (i_ACAv) presented a slightly elevated DC (**Figure 7C**). This outcome is likely relevant to the greater BLA axonal innervation of the i_ACAv in the $Tbr1^{+/-}$ ctrl group (**Figure 7A**). In contrast to our findings for DC, $Tbr1^{+/-}$ ctrl DMN regions showed elevated PC values for many regions relative to the WT ctrl DMN (**Figure 7C**). These analyses again evidence that the connectivity of DMN in $Tbr1^{+/-}$ ctrl mice is abnormal, thus resembling the aberrant DMN displayed by ASD patients [24, 25].

We also observed that BLA activation elicited alterations to the DMN in both wild-type and $Tbr1^{+/-}$ mice, albeit with different consequences (**Figure 7C**). In wild-type mice, BLA activation induced the same decreasing tendency for DC in many of the DMN regions, but various changes in PC (**Figure 7C**, WT TBS vs. WT ctrl). In $Tbr1^{+/-}$ mice, the same stimulation treatment elevated DC but reduced PC in many of the DMN regions (**Figure 7C**, $Tbr1^{+/-}$ TBS vs. $Tbr1^{+/-}$ ctrl). These differences in the DMN between wild-type and $Tbr1^{+/-}$ mice upon BLA activation clearly demonstrate the differential responses of DMN regions to BLA activation (**Figure 7C,** $Tbr1^{+/-}$ TBS vs. WT TBS).

Finally, we examined the reciprocal C-FOS correlation within different DMN regions to visualize the inter-connected networks of the specific DMN regions (**Figure 7D**). Consistent with a previous study on the mesoscopic mouse DMN [23], our wild-type mice exhibited a strong inter-connected network among DMN regions (**Figure 7D**). However, the DMN of awake $Tbr1^{+/-}$ mice ($Tbr1^{+/-}$ ctrl) had a much more loosely connected network, showing overall hypoconnectivity (**Figure 7D**). TBS of the BLA again exerted opposing effects on $Tbr1^{+/-}$ mice and their wild-type littermates, with the WT TBS group having a loosely connected network, whereas the inter-connected network of the $Tbr1^{+/-}$ TBS group became stronger than that of $Tbr1^{+/-}$ ctrl mice (**Figure 7D**).

Thus, using DMN as an example, our analyses support that Tbr1 deficiency and BLA

activation regulate the connectivity of physiologically relevant circuits.

Discussion

Using a whole-brain mapping and quantification system, we investigated the structure and

functional rewiring of BLA-derived circuits in a *Tbr1* ASD mouse model based on BLA axonal

projection patterns and C-FOS expression. We show impairment of contralateral projection and

ipsilateral mistargeting of BLA neurons in our *Tbr1$^{+/-}$* mice, consequently altering global neural

activity under both control and TBS-mediated BLA activation conditions. We also observed

that whole-brain synchronization is altered by *Tbr1* deficiency, with or without BLA activation

(**Figure 8**). Our system provides a high-resolution method for analyzing functional connectivity

at the cellular level, which is a much more accurate approach compared to fMRI, even though

temporal resolution is not achievable in our system. To our knowledge, ours is the first study

to provide evidence in an ASD mouse model that particular circuit activation elicits differential

functional consequences in the mutant mice compared to wild-type mice subjected to the same

manipulation.

Altered local and long-range functional connectivity is one of the primary features of

ASD [26-29]. Mouse models have also been shown to recapitulate the connectivity deficits

displayed by human patients [27, 30-35]. *TBR1* deficiency is a particularly interesting case

because one of the outcomes, i.e., impairment of the anterior commissure, is shared by both

mouse and human [9-13]. Interamygdalar projections of BLA neurons into the two brain

hemispheres are defective in cases of *Tbr1* deficiency [7, 13]. In the current report, we have

further demonstrated that BLA axons in *Tbr1$^{+/-}$* mice barely cross the midline between brain

hemispheres and mainly mistarget to ipsilateral brain regions. Although we have identified

several ipsilateral brain regions with more axonal projections from the ipsilateral BLA, our

PCA analysis indicates that many ipsilateral target regions exhibit considerable inter-sample

variability, indicating that some ipsilateral projections tend to be random and less specific. Our

work further highlights abnormal network synchronization, both at the whole-brain scale and

within a specific subnetwork (i.e., the DMN), in *Tbr1* mutant mice, providing the role of DMN

in an ASD mouse model.

The default mode network (DMN), originally defined by resting-state fMRI in human,

is highly relevant to ASD [22, 24, 25, 36, 37]. DMN abnormality results in atypical information

integration, prompting social-cognitive deficits [22, 37]. Unlike in humans, there are much

fewer studies on DMN in ASD mouse models [38, 39]. Here, we show that *Tbr1* deficiency

alters the structural and functional connectivity between the BLA and DMN. Moreover, BLA activation results in opposing consequences for $Tbr1^{+/-}$ mice and wild-type littermates in terms of neuronal activity and inter-network connectivity of the DMN. Similar to global brain synchronization, TBS-induced BLA activation elicits stronger connectivity among regions of the DMN in $Tbr1^{+/-}$ mice, thereby likely improving DMN function.

Our previous study showed that bilateral infusion of D-cycloserine, a coagonist of NMDA receptor, into the BLA improved the social behaviors and associated memory of $Tbr1^{+/-}$ mice [7, 12], revealing a beneficial effect of BLA activation on $Tbr1^{+/-}$ mice. Consistently, here, we found that TBS-mediated BLA activation globally increased correlations of C-FOS expression among different brain regions, thus contributing to whole-brain synchronization. This outcome implies that deep brain stimulation at the BLA likely improves the deficits caused by $Tbr1$ deficiency, consistent with previous studies indicating that deep brain stimulation ameliorates autism phenotypes [40-42].

Methods

Ethics statement

All animal experiments were performed with the approval of the Academia Sinica Institutional Animal Care and Utilization Committee and in strict accordance with its guidelines (Protocol No. 14-11-759 and 18-10-1234). Human subject study is not included in the current work.

Mice

The $Tbr1^{+/-}$ mice were originally provided by Drs. J. L. Rubenstein (Department of Psychiatry, University of California, San Francisco) and R. F. Hevner (Department of Neurological Surgery, University of Washington, Seattle). $Tbr1^{+/-}$ mice and wild-type littermates were housed in the animal facility of the Institute of Molecular Biology, Academia Sinica, under controlled temperature and humidity and a 12 h light/12 h dark cycle. All animal experiments were performed at 3-5 months of age with the approval of the Academia Sinica Institutional Animal Care and Utilization Committee and in strict accordance with its guidelines (Protocol No. 14-11-759 and 18-10-1234).

Virus production

For axon tracing and *in vivo* stimulation of BLA-derived circuits, we packaged an adeno-associated virus (AAV) that expresses the ultra-fast channelrhodopsin (ChR) variant oChIEF-Citrine under the control of the human Synapsin promoter (AAV-Syn-oChIEF-Citrine) as

described in our previous study [13]. Briefly, a transgenic plasmid encoding oChIEF-Citrine,

AAV helper plasmid serotype AAV8 capsid and AAV2 replication proteins, together with an

adenovirus helper plasmid encoding essential proteins for AAV packaging were transfected

into adenovirus E1-expressing cells. Crude virus soup was obtained from transfected E1-

expressing cells and purified by iodixanol gradient centrifugation. The physical vector titers

were quantified by measuring the number of packaged vector genomes by real-time PCR using

SYBR Green reaction mix (Roche Diagnostics, Mannheim, Germany).

Experimental procedure of the mesoscopic whole-brain scale circuit analysis pipeline

In order to analyze the structural and functional connectivity of BLA-derived circuits, we

developed a mesoscopic whole-brain scale circuit analysis pipeline (**Figure 1 and Figure S1**).

Detailed methods are described in the following sections.

Step1: Brain sample preparation

To label the BLA-derived circuits and manipulate their activity for examining functional

consequences, we performed stereotaxic surgeries for virus infection and optic fiber

implantation. Mice were deeply anesthetized and their heads were secured on a Lab Standard™

Stereotaxic Instrument (Stoelting, Wood Dale, IL USA). AAV-Syn-oChIEF-Citrine (0.1 μl,

10^{10} genomes/μl) was slowly infused over 10 min unilaterally into the BLA (virus injection

coordinates: 1.3 mm posterior, 3.3 mm lateral and 4.95 mm ventral to the bregma). Upon

completing virus injection, an optic fiber was slowly inserted into the site to 0.05 mm above

the virus injection site (optic fiber implantation coordinates: 1.3 mm posterior, 3.3 mm lateral

and 4.9 mm ventral to the bregma).

After recovering for at least 3 weeks, the mice were habituated to being hand-held, to

having a patch cable connected to the optic fiber, and to the lid of their home cage being opened

for 10 minutes when the patch cable was being connected. Habituation sessions were conducted

once per day for 3 days. *In vivo* optogenetic simulation and brain sample fixation were carried

out on the day following the last habituation session. We used a 473 nm laser with 10 mW light

intensity (emitted from the tip of the optic fiber) for optogenetic stimulation. After 10 min of

freely exploring their home cage, mice received five repetitive theta burst frequency

stimulations (TBS: 10 trains/TBS, inter-train interval, 200 ms; 5 pulses/train, inter-pulse

interval, 10 ms) with a 30 s inter-TBS interval to a unilateral BLA. This experimental

stimulation paradigm mimics physiological theta oscillations and has been demonstrated to

intensify responses to cortical and thalamic inputs through commissural BLA projections [13].

TBS was also used to induce changes in neuronal plasticity of the BLA [43]. Therefore, the
functional consequences of TBS on BLA-derived circuits can persist and prevail over the
random activation of BLA neurons during wakefulness. After photostimulation or sham
manipulation (without light emission), mice were kept in their home cage for 2 hr until sacrifice
by perfusing them with 4% paraformaldehyde in PBS. The brains were then dissected out and
post-fixed overnight at 4 °C.

Step2: High-content immunofluorescence imaging
Brains were cryopreserved in 30% sucrose at 4 °C for 2 days and then embedded in OCT (4583,
Tissue-TeK). For whole-brain analysis, we used a cryostat microtome (CM1900, Leica) to
cryosection each brain from the posterior to anterior cerebrum into at least 120 consecutive
coronal brain slices (60 μm/slice), with the cerebellum and olfactory bulb being excluded from
our analysis. Traditional immunohistochemical methods were deployed, as described
previously [13]. Primary anti-GFP (1:5000, Abcam) and anti-C-FOS (1:200, Cell Signaling)
antibodies were used to reveal Citrine-containing BLA axonal projections and C-FOS
immunoreactivities, respectively. The stained coronal brain sections (at least 120 sections) with
intensified oChIEF-Citrine (stained with GFP antibody) and C-FOS (stained with C-FOS
antibody) signals were imaged with a high-content imaging system (Molecular Devices)
hosting a 4x objective. Images from one brain were stitched together using MetaXpress
6.2.3.733 (Molecular Devices) and manually aligned in Amira 6.4 (ThermoFisher Scientific)
or Etomo [44].

Step3: Registration and transformation to Allen Mouse Common Coordinate Framework
version 3 (CCFv3)
After acquiring the aligned consecutive slice raw images, we registered image series to CCFv3
[15]. We downloaded high-resolution coronal templates with a 10 μm resolution along the
anterior-posterior (AP) axis using Allen Software Development Kit Python code, and then
manually matched the corresponding template to each raw image according to certain landmark
brain structures, such as the beginning and end of the hippocampus (HIP), the location of the
anterior commissure (act and aco), and the disconnections of the corpus callosum (cc). After
determining the matched template for each raw image, binary experimental images were
registered to the matched binary templates by means of the medical image registration library
SimpleElastix [45], first with 12 global (translation and affine) and then one local (B-spline)

transforming parameters. Finally, the stored multiple transforming parameters were applied to

merge the experimental images of oChIEF-Citrine and C-FOS signals into registered images.

*Step4: Image segmentation for oChIEF and C-FOS signals (also see **Figure S1**)*

Before quantification of oChIEF and C-FOS signals and comparing the differences between

wild-type and $Tbr1^{+/-}$ mice, we needed to eliminate fluorescence signals due to experimental

variation and background interference. The purpose of image segmentation for oChIEF signals

was to determine which pixels are oChIEF-positive (**Figure S1A**). Therefore, first we deployed

the pixel classification algorithm ilastik1.3.3 [46] to predict the probability that every pixel in

a registered image displayed oChIEF signals. The oChIEF probability maps of each image were

cross-referenced against the registered images pixel by pixel. The resulting filtered images (raw

signal intensity x probability) were further processed to intensify edges by subtracting the

Gaussian blurred version of the filtered images. The edge-enhanced filtered images were then

subjected to thresholding/binarization (>6x S.D. of the Z-stacked image with filtering and edge

enhancement) to isolate oChIEF-positive pixels.

532   The aim of image segmentation for C-FOS signals was to determine the locations of

C-FOS$^+$ cells within each image (**Figure S1B**). We preprocessed the raw C-FOS images by

applying rolling-ball background subtraction (ImageJ, radius 50 pixels). The spot detection

algorithm of Imaris 9.3 (Bitplane) was then applied to detect C-FOS$^+$ cells from background-

subtracted images using consistent criteria (Detection spot diameter: 10 μm; Detection

threshold: 2 x S.D. of the Z-stacked image with background subtraction). The locations of C-

FOS$^+$ cells in each image could then be exported. Finally, the binarized oChIEF signals and C-

FOS$^+$ cell locations of each registered image were subjected to quantification.

Step5: Application of brain structure masks and quantification of oChIEF and C-FOS signals

We accessed the Reference Space of CCFv3 to obtain the coronal indicator masks of each brain

region by using Allen Software Development Kit Python code. The coronal indicator masks

could separately compartmentalize oChIEF-positive pixels and C-FOS$^+$ cells into different

brain regions, which served as regions of interest (ROI) to quantify the binarized oChIEF

percentage or C-FOS$^+$ cell number for each brain region along the AP axis. All computations

were carried out using the Numpy, Pandas, OpenCV, scikit-image and GeoPandas Python

packages. The data compartmentalizing oChIEF-positive pixels and C-FOS$^+$ cells into different

brain regions were subsequently used for 3D visualization. Data on quantified oChIEF

percentage, C-FOS$^+$ cell numbers, and brain region area were used for the analysis. Some brain

regions were excluded from the analysis of oChIEF signals, such as ipsilateral BLA-related regions (LA, BLA, BLAa, BLAp) and bilateral VISam, which contained artificial signals caused by experimental surgery for virus injections. Moreover, the bilateral somatosensory cortex (SSp, SSp-n, SSp-bfd, SSp-ll, SSp-tr, SSp-un, SSs) and corpus callosum-related fiber tracts (cc, fa, ec, ee, ccb, ccs) were also excluded from the analysis due to exiguous oChIEF-Citrine-expressing AAV infections during surgery.

k-nearest neighbors (KNN) algorithm-based classification

To examine the spatial separability of virus injection sites or PC locations in a PCA analysis, we conducted a KNN-based classification with five-fold cross-validation using scikit-learn Python package. A series of numbers of nearest neighbors was used to identify the one achieving the best KNN classification accuracy. Furthermore, a randomized dataset of virus-injecting locations or PC locations, shuffled 1000 times, between experimental groups was used to validate the significance of spatial separability. The spatial separability was considered significant when the KNN classification accuracy of the real dataset deviated from the same metric distribution calculated for the randomized dataset ($P < 0.05$).

Volume-based quantification and related analysis

Volume-based quantification of oChIEF signals allowed us to calculate the percentage of oChIEF signals (%) in a given brain region volume. To do this, total numbers of oChIEF pixels within a brain region were calculated first by summing over all oChIEF pixels in single slices (products of quantified oChIEF percentage and region area) belonging to the same region along the AP axis and then dividing by the sum of the region area along the AP axis. Volume-based quantification of C-FOS signals revealed the density of C-FOS$^+$ cells in a given brain region volume (cells/mm^3). To calculate these values, total C-FOS$^+$ cells within a brain region were calculated first by summing over all C-FOS$^+$ cells in single slices and then dividing by the volume of the brain region (converted to mm^3). Volume-based quantification data for oChIEF and C-FOS signals were used in structure-wise comparisons (both oChIEF and C-FOS data), principal component analysis (PCA; oChIEF data), and network analysis of inter-regional C-FOS correlations (C-FOS data), all described further below.

*Structure-wise comparison of volume-based quantification data (also see **Figure S3A**)*

Before undertaking structure-wise comparisons of oChIEF percentages or C-FOS cell densities between different experimental groups (oChIEF data: WT and *Tbr1*$^{+/-}$; C-FOS data: WT ctrl,

Tbr1$^{+/-}$ ctrl, WT TBS and *Tbr1*$^{+/-}$ TBS), we set a threshold for selecting brain regions for the

comparisons. To compare oChIEF percentages, brain regions with a oChIEF percentage greater

than the 20th percentile in all *Tbr1*$^{+/-}$ regional quantifications were selected (257 contralateral

and 357 ipsilateral brain regions). For C-FOS$^+$ cell density comparison, brain regions with a

summed C-FOS cell density for all experimental groups (WT ctrl, *Tbr1*$^{+/-}$ ctrl, WT TBS and

Tbr1$^{+/-}$ TBS) greater than 800 cells/mm^3 were selected (311 contralateral and 307 ipsilateral

brain regions). All the data on each brain region were normalized to the respective averaged

WT (oChIEF data) or WT ctrl (C-FOS data) values before being subjected to statistical analysis.

For the structure-wise oChIEF comparison, a two-sided Mann-Whitney U test was conducted

in the SciPy sub-package scipy.stats. For the structure-wise C-FOS comparison, a Kruskal-

Wallis test was conducted (using scipy.stats), followed by a post-hoc Dunn test with Bonferroni

adjustment of *P* value (using the scikit-posthocs Python package) if the Kruskal-Wallis test was

significant (P <0.05).

Structure-wise comparison of the ratio of contralateral Citrine signal to ipsilateral Citrine

signal (the C/I ratio) from volume-based quantification data

The brain regions selected for C/I ratio comparison were those displaying a significant oChIEF

percentage (above the defined threshold) in both brain hemispheres in wild-type samples. The

C/I ratio data were normalized to the averaged C/I ratio of wild-type samples. The structure-

wise C/I ratio comparison was also conducted by two-sided Mann-Whitney U test using

scipy.stats.

PCA analysis of oChIEF signals

To visualize differences in oChIEF percentages across all brain regions and between wild-type

and *Tbr1*$^{+/-}$ mouse brains, we conducted a PCA analysis on the oChIEF percentage data using

the scikit-learn. Per-sample data across all brain regions were projected onto the first two

principal components.

Network analysis of inter-regional C-FOS correlations

Functional connectivity, as measured in our study, was based on synchronization of C-FOS

expression levels between two brain regions in individual subjects. Therefore, we used relative

C-FOS density (normalized to the averaged value for WT ctrl) obtained from our volume-based

quantification data as a single metric to represent the C-FOS level of a given brain region of

one sample. To measure all possible synchronizations, we evaluated inter-regional C-FOS

correlations between all possible brain region pairs by means of two-sided Kendall's rank

correlation (using scipy.stats) across all the samples in the same experimental group. A network

of inter-regional C-FOS correlations was constructed using the significant C-FOS correlations

(edges, significance level $P < 0.05$) between brain region pairs (nodes) by using the NetworkX

Python package. Self-to-self correlations were removed. The degree centrality (DC) and

participation coefficient (PC) of individual nodes (brain regions) were calculated using

NetworkX.

To dissect the modular organization of networks, we adopted a Louvain community detection

algorithm and the modularity metric (Q) calculation provided by NetworkX. To identify the

optimal resolution parameter R value used for Louvain community detection, we generated a

shuffled network by maintaining the same correlation weights (edges) while randomizing the

source and target regions (nodes). An R value series (0.4 to 0.9 in steps of 0.1) was used for

Louvain community detection of real and shuffled networks. For each R value, the difference

in Q between the modularity of the real network and the shuffled network was calculated 1000

times. The most appropriate R value that generated the maximal mean difference of Q (Q–

$Q_{shuffled}$) was selected for Louvain community detection. These computations were achieved

using Numpy, Pandas and NetworkX.

Slice-based quantification and related analysis

Slice-based quantifications of oChIEF and C-FOS signals involved calculating the oChIEF

pixel (products of quantified oChIEF percentage and region area) and C-FOS$^+$ cell distributions

along the AP axis, respectively. To achieve fair comparisons, we carried out 100 μm binning

of oChIEF pixel and C-FOS$^+$ cell distributions along the AP axis. The binned oChIEF pixel and

C-FOS$^+$ cell distributions were then used for structure-wise comparisons and correlation

analysis between rewired BLA-derived circuits and their C-FOS modulatory ability.

*Structure-wise comparison of the data from slice-based quantification (also see **Figure S6A**)*

Before structure-wise comparison of oChIEF pixel or C-FOS$^+$ cell distributions between

different groups (oChIEF data, WT and *Tbr1*$^{+/−}$; C-FOS data, WT ctrl, *Tbr1*$^{+/−}$ ctrl, WT TBS

and *Tbr1*$^{+/−}$ TBS), we set a threshold to select brain regions for comparison. To compare

oChIEF pixel distributions, brain regions were selected with a summed total oChIEF pixels for

both groups (WT + *Tbr1*$^{+/−}$) greater than 13-fold the standard deviation of all oChIEF pixels in

all contralateral brain regions (207 contralateral and 344 ipsilateral brain regions). To compare

C-FOS$^+$ cell distributions, brain regions with a summed total C-FOS$^+$ cells for all compared

653 groups (WT ctrl, $Tbr1^{+/-}$ ctrl, WT TBS and $Tbr1^{+/-}$ TBS) greater than 100 cells were selected

654 (313 contralateral and 311 ipsilateral brain regions). For our structure-wise comparisons of

655 oChIEF pixel and C-FOS$^+$ cell distributions along the AP axis, Friedman's one-way repeated

656 measure analyses were conducted using the Pingouin Python package. A post-hoc

657 nonparametric pairwise test with one-step Bonferroni correction of P value (using Pingouin)

658 was conducted for the C-FOS$^+$ cell distributions if the outcome of the Friedman one-way

659 repeated measure analysis was significant (P <0.05). To express the extent of distributional

660 differences between compared groups, we conducted pair-wise subtractions of the oChIEF

661 pixels or C-FOS$^+$ cell numbers for the same AP coordinates between all possible pairs. The

662 differences along the AP axis of all sample pairs were then averaged and summed into single

663 metrics (sum of ΔoChIEF or sum of ΔC-FOS) to represent the extent of any differences.

665 *Correlation analysis of rewired BLA-derived circuits and their C-FOS modulatory ability*

666 Brain regions selected for correlation analysis were those receiving differential BLA axonal

667 innervations between wild-type and $Tbr1^{+/-}$ mouse brains (Friedman test P value < 0.05) and

668 altered C-FOS$^+$ cell distributions upon TBS-mediated BLA activation in either wild-type or

669 $Tbr1^{+/-}$ mice (adjusted P value <0.05 in WT TBS vs. WT ctrl or $Tbr1^{+/-}$ TBS vs. $Tbr1^{+/-}$ ctrl).

670 First, we determined ΔoChIEF traces by pairwise subtraction of oChIEF-Citrine signals

671 between $Tbr1^{+/-}$ and wild-type samples. The neural activation induced by TBS in $Tbr1^{+/-}$ mice

672 and WT littermates, i.e., WT ΔC-FOS traces and $Tbr1^{+/-}$ ΔC-FOS traces, was also determined.

673 Pairwise subtraction between $Tbr1^{+/-}$ ΔC-FOS traces and WT ΔC-FOS traces was then executed

674 to obtain all possible differences, i.e., ΔΔC-FOS traces. A Pearson correlation coefficient (r)

675 was calculated between the averaged ΔoChIEF trace and the averaged ΔΔC-FOS trace for each

676 brain region. To determine if any correlation was meaningful, i.e., distinct from non-correlated

677 pairs, the order of ΔoChIEF traces, WT ΔC-FOS traces and $Tbr1^{+/-}$ ΔC-FOS traces along the

678 AP axis was independently reshuffled 1000 times to create 1000 putatively non-correlated mean

679 ΔoChIEF and mean ΔΔC-FOS shuffled pairs. We defined that a meaningful positive or negative

680 correlation between mean ΔΔoChIEF and ΔΔC-FOS values should fit three criteria. First, the r

681 value should deviate from the same metric obtained from the shuffled dataset ($r_{shuffled}$) (positive

682 correlation, above 95% $r_{shuffled}$ distribution; negative correlation, below 95% $r_{shuffled}$ distribution).

683 Second, the r value should be >0.1 (positive correlation) or <–0.1 (negative correlation). Third,

684 the tendency thresholds of ΔoChIEF and ΔΔC-FOS traces were determined according to the

685 mean 99th percentiles of scaled differences, which were calculated from the pair-subtraction

population dataset with 1000 times AP axis reshuffling (**Figures 4B, 4D**, dashed line in middle

panel). A positive or negative correlation between mean ΔoChIEF and $\Delta\Delta$C-FOS depended on

the product of ΔoChIEF and $\Delta\Delta$C-FOS tendency thresholds (positive correlation, > 0; negative

correlation, <0). The computations of all correlation analyses were carried out using Pandas

and Numpy.

ROI corrections of brain region masks

We selected 44 contralateral and 50 ipsilateral brain regions for manual correction of the

mismatched ROI in every single brain slice image of each brain sample. To do this, the original

coronal indicator masks of each brain region were separated into contralateral and ipsilateral

masks and converted to ImageJ ROIs by sequentially applying the ImageJ Create Selection

function and adding it to ImageJ ROI Manager. Using the C-FOS or oChIEF raw images

(registered to CCFv3) as references, ROIs were verified and corrected by manually adjusting

their shapes and locations. After ROI corrections, the ROIs of each brain region in both

hemispheres were converted back to image masks by applying the ImageJ Create Mask function.

The new coronal indicator masks were then used for re-quantifications.

3D visualization

Summarized 3D stack of binarized oChIEF images

To visualize differences in BLA-derived circuits between wild-type and *Tbr1*$^{+/-}$ mice, the

binarized oChIEF image series of individual brains was binned on the dorsal-ventral and

medial-lateral axis into 10 μm/pixel. Each binned pixel intensity represented the oChIEF

probability. The AP coordinates of the binned oChIEF image series were converted into the

respective CCFv3 coordinates (10 μm/pixel) and all binned oChIEF image series of individual

brains were then merged. If the same AP coordinate contained multiple binned oChIEF image

series from different brain samples, we conducted averaging on the intensity of those binned

oChIEF image series. If an AP coordinate contained only one binned oChIEF image, then that

binned image represented the oChIEF signals for that particular AP coordinate. The 3D stack

of binned and averaged oChIEF image series was adopted to illustrate BLA-derived circuits at

a whole-brain scale. Image processing was carried out using ImageJ, Numpy and scikit-image.

3D mapping of binned BLA-derived circuits or C-FOS$^+$ cell distributions

To visualize the innervations of BLA-derived circuits or the distributions of C-FOS$^+$ cells in

particular brain regions, we used the datasets on compartmentalized oChIEF-positive pixels and

C-FOS$^+$ cells for different brain regions. The CCFv3 whole brain shell was obtained from the

image stacks of CCFv3 "root" coronal indicator masks and converted to an isosurface mesh

with smoothing using the vedo Python package. For BLA-derived circuit visualization, binned

and averaged oChIEF image stacks within particular regions were employed and binned pixels

(10 μm^3 cubes) were converted into vectorized points using Numpy, Pandas and vedo. Each

vectorized point expressed different information, such as the brain region to which the point

belonged (discontinuous color map), scaled metrics expressing the difference between

compared groups (continuous color map), and oChIEF probability (alpha level). Vectorized

oChIEF points were mapped onto the 3D whole brain shell using vedo. To visualize C-FOS$^+$

cell distributions, the locations of C-FOS$^+$ cells were first binned by 100 μm^3 cubes in every

brain sample, and then the averaged C-FOS density of every 100 μm^3 cube was calculated

across brain samples of the same experimental group. Every binned and averaged C-FOS cube

(100 μm^3) within particular regions was converted into a vectorized sphere using Numpy,

Pandas and vedo. Each vectorized sphere expressed different information, such as the brain

region to which the sphere belonged (discontinuous color map), scaled metrics expressing the

difference between compared groups (continuous color map), and the normalized C-FOS

density within the cube (size of the sphere). Vectorized C-FOS spheres were mapped onto the

3D whole brain shell using vedo.

3D mapping of C-FOS correlation networks among main brain areas

To visualize the C-FOS correlation networks shown in **Figure 8**, the central locations of the

main cortical (prefrontal, somatomotor, medial, lateral, auditory and visual) and subcortical

areas (OLF, HPF, CTXsp, CNU, TH, HY, MB and HB) in both hemispheres were first

determined by averaging the locations of all the binned 100-μm^3 cubes belonging to particular

brain areas. Mean correlations within and between brain areas were then calculated. Centroids

representing particular brain areas in the ipsilateral or contralateral hemisphere were converted

to vectorized spheres using vedo. The color of each sphere represented brain area identity and

the size reflected mean correlation extent. To visualize inter-area mean correlations, correlation

edges were generated using vedo. For each edge, color represented correlational polarity and

thickness reflected mean correlation extent. Vectorized centroid spheres (brain areas) and inter-

area edges (inter-area mean correlations) were mapped onto a 3D whole brain shell again using

vedo.

Quantification and statistical analysis

Sample size n represents the number of brain samples from individual animals in each

experimental group. We did not use statistical methods to predetermine sample sizes, but our

sample size is similar to those brain connectome studies in previous publication [47]. Volume-

based quantification and its-associated analysis of oChIEF signals (KNN-based classification

of virus injection sites and PC locations, structure-wise comparisons, PCA analysis and C/I

ratio analysis) are based on n = 13 of wild-type (6 from WT ctrl and 7 from WT TBS groups)

and n = 15 of Tbr1$^{+/-}$ brain samples (7 from Tbr1$^{+/-}$ ctrl and 8 from Tbr1$^{+/-}$ TBS groups).

Volume-based quantification of relative C-FOS density and inter-regional C-FOS correlation

analysis was based on the brain samples largely overlapped with volume-based quantification

and analysis of oChIEF signals. We included one additional brain sample for both genotypes

which has mistargeted virus expression in the control condition and excluded one sample from

Tbr1$^{+/-}$ TBS due to inappropriate image acquisition of C-FOS signals. Brain samples used for

slice-based analysis (structure-wise comparisons and correlation analysis) are almost the same

as those used in the volume-based analysis of oChIEF signals, with the exclusion of one brain

sample from the Tbr1$^{+/-}$ TBS group which had inappropriate image acquisition of C-FOS

signals. The exact n numbers of experimental groups for every analysis can be found in figure

legends. For the data distribution, we did not use statistical methods to test data normality but

we assumed most of the quantified data is not normally distributed. Therefore, non-parametric

tests were used for statistical analysis. Statistical details can be found in the method sections

describing each analysis. Data representing statistical parameters such as mean ± SEM, median

and interquartile range are described in figure legends. All test statistics (e.g. exact *P* value,

Mann-Whitney U statistic, Friedman chi-square statistic, and degrees of freedom) of structure-

wise comparisons (volume-based or slice-based analysis) can be found in **Tables S1 and S2.**

Declarations

Ethics approval and consent to participate

Only mice were used in this study. All animal experiments were performed with the approval

of the Academia Sinica Institutional Animal Care and Utilization Committee (Protocol No. 14-

11-759 and 18-10-1234) and in strict accordance with its guidelines and those of the Council

of Agriculture Guidebook for the Care and Use of Laboratory Animals.

Consent for publication

Not applicable.

Availability of data and materials

All sample data will be shared by the lead contact upon request. Custom code used for volume-

based quantification, slice-based quantification, network analysis and 3D visualization has been

uploaded to GitHub (https://github.com/HsuehYiPing/Tbr1Amyg).

Competing interests

The authors report no competing interests.

Funding

This work was supported by grants from Academia Sinica (AS-IA-106-L04 and AS-TP-110-

L10 to Y.-P.H.), and the National Science and Technology Council (NSTC 112-2326-B-001-

008 to Y.-P.H.). The funders had no role in study design, data collection and analysis, the

decision to publish or the preparation of the manuscript.

Author contributions

Conceptualization, T.-T.H. and Y.-P.H.; Methodology and investigation, T.-T.H.; Coding, T.-

T.H. and C.-Y.W.; Writing – original draft, T.-T.H. and Y.-P.H.; Writing – review and editing,

T.-T.H., C.-Y.W. and Y.-P.H.; Funding acquisition, Y.-P.H.; Supervision and project

administration, Y.-P.H. All authors contributed to the article and approved the submitted

version.

Acknowledgments

We thank the Imaging Core and Animal Facility of the Institute of Molecular Biology,

Academia Sinica, for technical assistance, Dr. John O'Brien for English editing and members

of Y.-P.H.'s laboratory for technical assistance and discussion. This work was supported by

grants from Academia Sinica (AS-TP-110-L10 and AS-IA-111-L01 to Y.-P.H.).

References

1. Geschwind DH, Levitt P. Autism spectrum disorders: developmental disconnection syndromes. Curr Opin Neurobiol. 2007;17(1):103-11. Epub 2007/02/06. doi: 10.1016/j.conb.2007.01.009.
2. Uddin L, Supekar K, Menon V. Reconceptualizing functional brain connectivity in autism from a developmental perspective. Frontiers in Human Neuroscience. 2013;7:458. doi: 10.3389/fnhum.2013.00458.
3. Di Martino A, Yan CG, Li Q, Denio E, Castellanos FX, Alaerts K, et al. The autism brain imaging data exchange: towards a large-scale evaluation of the intrinsic brain architecture in autism. Molecular Psychiatry. 2014;19(6):659-67. doi: 10.1038/mp.2013.78.
4. Ameis SH, Catani M. Altered white matter connectivity as a neural substrate for social impairment in Autism Spectrum Disorder. Cortex. 2015;62:158-81. doi: https://doi.org/10.1016/j.cortex.2014.10.014.
5. Bulfone A, Smiga SM, Shimamura K, Peterson A, Puelles L, Rubenstein JL. T-brain-1: a homolog of Brachyury whose expression defines molecularly distinct domains within the cerebral cortex. Neuron. 1995;15(1):63-78. PubMed PMID: 7619531.
6. De Rubeis S, He X, Goldberg AP, Poultney CS, Samocha K, Cicek AE, et al. Synaptic, transcriptional and chromatin genes disrupted in autism. Nature. 2014;515(7526):209-15. Epub 2014/11/05. doi: 10.1038/nature13772.
7. Huang TN, Chuang HC, Chou WH, Chen CY, Wang HF, Chou SJ, et al. Tbr1 haploinsufficiency impairs amygdalar axonal projections and results in cognitive abnormality. Nat Neurosci. 2014;17(2):240-7. Epub 2014/01/21. doi: 10.1038/nn.3626.
8. Chuang HC, Huang TN, Hsueh YP. T-Brain-1--A Potential Master Regulator in Autism Spectrum Disorders. Autism Res. 2015;8(4):412-26. Epub 2015/01/21. doi: 10.1002/aur.1456. PubMed PMID: 25600067.
9. Nambot S, Faivre L, Mirzaa G, Thevenon J, Bruel AL, Mosca-Boidron AL, et al. De novo TBR1 variants cause a neurocognitive phenotype with ID and autistic traits: report of 25 new individuals and review of the literature. Eur J Hum Genet. 2020. Epub 2020/02/02. doi: 10.1038/s41431-020-0571-6. PubMed PMID: 32005960.
10. Hsueh YP, Hsu TT, Huang TN. The evolutionarily conserved function of TBR1 in controlling the size of anterior commissure in human and mouse brains. Eur J Hum Genet. 2020;28(8):997-8. Epub 2020/04/11. doi: 10.1038/s41431-020-0621-0. PubMed PMID: 32273580.
11. Co M, Barnard RA, Jahncke JN, Grindstaff S, Fedorov LM, Adey AC, et al. Shared and Distinct Functional Effects of Patient-Specific Tbr1 Mutations on Cortical Development. J Neurosci. 2022;42(37):7166-81. Epub 2022/08/10. doi: 10.1523/jneurosci.0409-22.2022. PubMed PMID: 35944998; PubMed Central PMCID: PMCPMC9480892.
12. Huang TN, Yen TL, Qiu LR, Chuang HC, Lerch JP, Hsueh YP. Haploinsufficiency of autism causative gene Tbr1 impairs olfactory discrimination and neuronal activation of the olfactory system in mice. Mol Autism. 2019;10:5. Epub 2019/02/23. doi: 10.1186/s13229-019-0257-5. PubMed Central PMCID: PMCPMC6371489.
13. Huang TN, Hsu TT, Lin MH, Chuang HC, Hu HT, Sun CP, et al. Interhemispheric Connectivity Potentiates the Basolateral Amygdalae and Regulates Social Interaction and Memory. Cell Rep. 2019;29(1):34-48.e4. Epub 2019/10/03. doi: 10.1016/j.celrep.2019.08.082. PubMed PMID: 31577954.
14. Hsu T-T, Huang T-N, Hsueh Y-P. Anterior Commissure Regulates Neuronal Activity of Amygdalae and Influences Locomotor Activity, Social Interaction and Fear Memory in Mice. Frontiers in Molecular Neuroscience. 2020;13:47. doi: 10.3389/fnmol.2020.00047.

15. Wang Q, Ding S-L, Li Y, Royall J, Feng D, Lesnar P, et al. The Allen Mouse Brain
Common Coordinate Framework: A 3D Reference Atlas. Cell. 2020;181(4):936-53.e20.
doi: https://doi.org/10.1016/j.cell.2020.04.007.
16. Hintiryan H, Bowman I, Johnson DL, Korobkova L, Zhu M, Khanjani N, et al.
Connectivity characterization of the mouse basolateral amygdalar complex. Nature
Communications. 2021;12(1):2859. doi: 10.1038/s41467-021-22915-5.
17. Wheeler AL, Teixeira CM, Wang AH, Xiong X, Kovacevic N, Lerch JP, et al.
Identification of a Functional Connectome for Long-Term Fear Memory in Mice. PLOS
Computational Biology. 2013;9(1):e1002853. doi: 10.1371/journal.pcbi.1002853.
18. Rogers-Carter MM, Varela JA, Gribbons KB, Pierce AF, McGoey MT, Ritchey M, et al.
Insular cortex mediates approach and avoidance responses to social affective stimuli.
Nature Neuroscience. 2018;21(3):404-14. doi: 10.1038/s41593-018-0071-y.
19. Rubinov M, Sporns O. Complex network measures of brain connectivity: Uses and
interpretations.   NeuroImage.   2010;52(3):1059-69.   doi:
https://doi.org/10.1016/j.neuroimage.2009.10.003.
20. Raichle ME. The Brain's Default Mode Network. Annual Review of Neuroscience.
2015;38(1):433-47. doi: 10.1146/annurev-neuro-071013-014030. PubMed PMID:
25938726.
21. Nair A, Jolliffe M, Lograsso YSS, Bearden CE. A Review of Default Mode Network
Connectivity and Its Association With Social Cognition in Adolescents With Autism
Spectrum Disorder and Early-Onset Psychosis. Front Psychiatry. 2020;11:614. Epub
2020/07/17. doi: 10.3389/fpsyt.2020.00614. PubMed PMID: 32670121; PubMed Central
PMCID: PMCPMC7330632.
22. Harikumar A, Evans DW, Dougherty CC, Carpenter KLH, Michael AM. A Review of the
Default Mode Network in Autism Spectrum Disorders and Attention Deficit Hyperactivity
Disorder. Brain Connect. 2021;11(4):253-63. Epub 2021/01/07. doi:
10.1089/brain.2020.0865. PubMed PMID: 33403915; PubMed Central PMCID:
PMCPMC8112713.
23. Whitesell JD, Liska A, Coletta L, Hirokawa KE, Bohn P, Williford A, et al. Regional,
Layer, and Cell-Type-Specific Connectivity of the Mouse Default Mode Network. Neuron.
2021;109(3):545-59. doi: https://doi.org/10.1016/j.neuron.2020.11.011.
24. Starck T, Nikkinen J, Rahko J, Remes J, Hurtig T, Haapsamo H, et al. Resting state fMRI
reveals a default mode dissociation between retrosplenial and medial prefrontal
subnetworks in ASD despite motion scrubbing. Frontiers in Human Neuroscience.
2013;7:802. doi: 10.3389/fnhum.2013.00802.
25. Chen L, Chen Y, Zheng H, Zhang B, Wang F, Fang J, et al. Changes in the topological
organization of the default mode network in autism spectrum disorder. Brain Imaging and
Behavior. 2021;15(2):1058-67. doi: 10.1007/s11682-020-00312-8.
26. Khan S, Gramfort A, Shetty NR, Kitzbichler MG, Ganesan S, Moran JM, et al. Local and
long-range functional connectivity is reduced in concert in autism spectrum disorders. Proc
Natl Acad Sci U S A. 2013;110(8):3107-12. Epub 2013/01/16. doi:
10.1073/pnas.1214533110. PubMed Central PMCID: PMC3581984.
27. Stoodley CJ, D'Mello AM, Ellegood J, Jakkamsetti V, Liu P, Nebel MB, et al. Altered
cerebellar connectivity in autism and cerebellar-mediated rescue of autism-related
behaviors in mice. Nat Neurosci. 2017;20(12):1744-51. Epub 2017/12/01. doi:
10.1038/s41593-017-0004-1. PubMed PMID: 29184200; PubMed Central PMCID:
PMCPMC5867894.
28. Hong SJ, Vos de Wael R, Bethlehem RAI, Lariviere S, Paquola C, Valk SL, et al. Atypical
functional connectome hierarchy in autism. Nat Commun. 2019;10(1):1022. Epub

2019/03/06. doi: 10.1038/s41467-019-08944-1. PubMed PMID: 30833582; PubMed
Central PMCID: PMCPMC6399265.
29. Buch AM, Vértes PE, Seidlitz J, Kim SH, Grosenick L, Liston C. Molecular and network-
level mechanisms explaining individual differences in autism spectrum disorder. Nat
Neurosci. 2023;26(4):650-63. Epub 2023/03/10. doi: 10.1038/s41593-023-01259-x.
PubMed PMID: 36894656.
30. Haberl MG, Zerbi V, Veltien A, Ginger M, Heerschap A, Frick A. Structural-functional
connectivity deficits of neocortical circuits in the Fmr1 (-/y) mouse model of autism. Sci
Adv. 2015;1(10):e1500775. Epub 2015/12/25. doi: 10.1126/sciadv.1500775. PubMed
Central PMCID: PMC4681325.
31. Gabrielsen TP, Anderson JS, Stephenson KG, Beck J, King JB, Kellems R, et al. Functional
MRI connectivity of children with autism and low verbal and cognitive performance.
Molecular Autism. 2018;9(1):67. doi: 10.1186/s13229-018-0248-y.
32. Shofty B, Bergmann E, Zur G, Asleh J, Bosak N, Kavushansky A, et al. Autism-associated
Nf1 deficiency disrupts corticocortical and corticostriatal functional connectivity in human
and mouse. Neurobiol Dis. 2019;130:104479. Epub 2019/05/28. doi:
10.1016/j.nbd.2019.104479. PubMed PMID: 31128207; PubMed Central PMCID:
PMCPMC6689441.
33. Zhang Z, Gibson JR, Huber KM. Experience-dependent weakening of callosal synaptic
connections in the absence of postsynaptic FMRP. Elife. 2021;10:e71555. Epub
2021/10/08. doi: 10.7554/eLife.71555. PubMed PMID: 34617509; PubMed Central
PMCID: PMCPMC8526058.
34. Zerbi V, Pagani M, Markicevic M, Matteoli M, Pozzi D, Fagiolini M, et al. Brain mapping
across 16 autism mouse models reveals a spectrum of functional connectivity subtypes.
Mol Psychiatry. 2021;26(12):7610-20. Epub 2021/08/13. doi: 10.1038/s41380-021-01245-
4. PubMed PMID: 34381171; PubMed Central PMCID: PMCPMC8873017.
35. Awad PN, Zerbi V, Johnson-Venkatesh EM, Damiani F, Pagani M, Markicevic M, et al.
CDKL5 sculpts functional callosal connectivity to promote cognitive flexibility. Mol
Psychiatry. 2023. Epub 2023/02/04. doi: 10.1038/s41380-023-01962-y. PubMed PMID:
36737483.
36. Jung M, Kosaka H, Saito DN, Ishitobi M, Morita T, Inohara K, et al. Default mode network
in young male adults with autism spectrum disorder: relationship with autism spectrum
traits. Molecular Autism. 2014;5(1):35. doi: 10.1186/2040-2392-5-35.
37. Padmanabhan A, Lynch CJ, Schaer M, Menon V. The Default Mode Network in Autism.
Biol Psychiatry Cogn Neurosci Neuroimaging. 2017;2(6):476-86. Epub 2017/10/17. doi:
10.1016/j.bpsc.2017.04.004. PubMed PMID: 29034353; PubMed Central PMCID:
PMCPMC5635856.
38. Kron M, Howell CJ, Adams IT, Ransbottom M, Christian D, Ogier M, et al. Brain activity
mapping in Mecp2 mutant mice reveals functional deficits in forebrain circuits, including
key nodes in the default mode network, that are reversed with ketamine treatment. J
Neurosci. 2012;32(40):13860-72. Epub 2012/10/05. doi: 10.1523/jneurosci.2159-12.2012.
PubMed PMID: 23035095; PubMed Central PMCID: PMCPMC3500840.
39. Liska A, Bertero A, Gomolka R, Sabbioni M, Galbusera A, Barsotti N, et al. Homozygous
Loss of Autism-Risk Gene CNTNAP2 Results in Reduced Local and Long-Range
Prefrontal Functional Connectivity. Cereb Cortex. 2018;28(4):1141-53. Epub 2017/02/12.
doi: 10.1093/cercor/bhx022. PubMed PMID: 28184409.
40. Sinha S, McGovern RA, Sheth SA. Deep brain stimulation for severe autism: from
pathophysiology to procedure. Neurosurg Focus. 2015;38(6):E3. Epub 2015/06/02. doi:
10.3171/2015.3.Focus1548. PubMed PMID: 26030703.

41. Penton T, Catmur C, Banissy MJ, Bird G, Walsh V. Non-invasive stimulation of the social
brain: the methodological challenges. Soc Cogn Affect Neurosci. 2022;17(1):15-25. Epub
2020/08/01. doi: 10.1093/scan/nsaa102. PubMed PMID: 32734295; PubMed Central
PMCID: PMCPMC9083106.
42. Yan H, Elkaim LM, Venetucci Gouveia F, Huber JF, Germann J, Loh A, et al. Deep brain
stimulation for extreme behaviors associated with autism spectrum disorder converges on
a common pathway: a systematic review and connectomic analysis. J Neurosurg.
2022;137:699-708. Epub 2022/01/22. doi: 10.3171/2021.11.Jns21928. PubMed PMID:
35061980.
43. Bazelot M, Bocchio M, Kasugai Y, Fischer D, Dodson PD, Ferraguti F, et al. Hippocampal
Theta Input to the Amygdala Shapes Feedforward Inhibition to Gate Heterosynaptic
Plasticity.   Neuron.   2015;87(6):1290-303.   Epub   2015/09/25.   doi:
10.1016/j.neuron.2015.08.024. PubMed Central PMCID: PMCPMC4590554.
44. Kremer JR, Mastronarde DN, McIntosh JR. Computer Visualization of Three-Dimensional
Image Data Using IMOD. Journal of Structural Biology. 1996;116(1):71-6. doi:
https://doi.org/10.1006/jsbi.1996.0013.
45. Marstal K, Berendsen F, Staring M, Klein S, editors. SimpleElastix: A User-Friendly,
Multi-lingual Library for Medical Image Registration. 2016 IEEE Conference on
Computer Vision and Pattern Recognition Workshops (CVPRW); 2016 26 June-1 July
2016.
46. Berg S, Kutra D, Kroeger T, Straehle CN, Kausler BX, Haubold C, et al. ilastik: interactive
machine learning for (bio)image analysis. Nature Methods. 2019;16(12):1226-32. doi:
10.1038/s41592-019-0582-9.
47. DeNardo LA, Liu CD, Allen WE, Adams EL, Friedmann D, Fu L, et al. Temporal evolution
of cortical ensembles promoting remote memory retrieval. Nature Neuroscience.
2019;22(3):460-9. doi: 10.1038/s41593-018-0318-7.
987
988

Figures and figure legends

Figure 1. Workflow of whole-brain-scale analysis of BLA-derived circuits. There are five steps in the analysis pipeline. Step 1: Brain sample preparation, involving unilateral BLA infection of oChIEF-Citrine-expressing AAV, *in vivo* theta-burst stimulation (TBS) of the AAV-infected BLA while the awake mice are in their home cage, and brain fixation by PFA 2 hours after optogenetic stimulation. Step 2: High-content immunofluorescence imaging. At least 120 coronal cryosections (60 μm/section) of a single mouse brain were subjected to immunostaining to reveal C-FOS+ cells and oChIEF-Citrine signals, and then imaged by a high-content imaging system. Step 3: Registration and transformation to the Allen brain atlas. Raw image series were registered to the templates of Allen common coordinate framework version 3 (CCFv3) by sequentially applying 12 global and 1 local transform parameters. Step 4: Image

segmentation for oChIEF and C-FOS signals. The oChIEF signals were isolated by pixel

classification and binarized. C-FOS$^+$ cell locations were detected by the spot detection function

of Imaris (also see **Figure S1**). Step 5: Application of brain structure masks and quantification

of oChIEF and C-FOS signals. Examples of choropleth maps show the quantification of

binarized oChIEF signals (% of pixels/slice, upper) and C-FOS$^+$ cell number (number/slice,

lower) of different brain regions in a single coronal section. Scale bar in Step 4, 1 mm.

1007

Figure 2. Rewiring of BLA-derived circuits.

(A) Coronal image series of binarized oChIEF-Citrine signals (green) superimposed on a CCFv3 template (gray) of wild-type (WT, upper) and *Tbr1*[+/−] mouse brain (lower) (also see **Video S1**).

(B) Coronal (left) and horizontal (right) views of summarized 3D stacks with binarized and binned (10 μm^3) oChIEF signals from WT (upper) and *Tbr1*[+/−] (lower) mouse brains. Color map indicates different probabilities of oChIEF signal in a 10 μm^3 cube (also see **Video S2**).

(C) Structure-wise comparison of binarized oChIEF signals between wild-type and *Tbr1*[+/−] mouse brains after volume-based quantification (also see **Table S1**). Scatterplots showing the fold-change (*Tbr1*[+/−]/WT) versus significance (Mann-Whitney U test *P* value) of each contralateral (upper) and ipsilateral (lower) brain region are presented. Vertical and horizontal

dotted lines represent fold change = 1 and U test *P* value = 0.05, respectively. Colors represent

brain region identities. Open and filled circles represent ROI non-corrected and corrected brain

regions, respectively.

(D) Coronal (left) and horizontal (right) views of 3D stacks with binarized and binned (10 μm^3)

oChIEF pixels in the brain regions displaying significantly different BLA-circuit rewiring

(*P*<0.05) between wild-type (upper) and *Tbr1*$^{+/-}$ mice (lower). Colors represent brain regions

as shown in (C). Alpha levels (0–1) of the brain region color represent scaled oChIEF

probabilities.

(E) Box plots showing comparisons of relative oChIEF signal density between WT and *Tbr1*$^{+/-}$

mouse brains. Only the regions displaying a significant difference (*P* <0.05) are shown. Left,

contralateral; right, ipsilateral. Boxes show median and interquartile ranges, and whiskers

extend to the rest of the distribution except for outliers. Spots represent individual samples.

Asterisks label the parental brain regions of subsequent offspring regions grouped by the

horizontal lines. Red font indicates ROI-corrected regions. Color bars under the charts represent

brain region identities, as shown in (C). The horizontal dotted lines represent the WT average,

i.e., 100%.

Scale bar in (B), 1 mm. Sample sizes of mice are the same for (B)-(E) (WT, n=13; *Tbr1*$^{+/-}$, n=

15).

1038

Figure 3. Differential C-FOS expression in WT and *Tbr1⁺/⁻* mouse brains with or without activation of BLA-derived circuits.

(A) Slice-based quantification of C-FOS⁺ cell distributions in WT ctrl, *Tbr1⁺/⁻* ctrl, WT TBS and *Tbr1⁺/⁻* TBS groups (left to right). Heatmaps show binned (100 μm) and averaged C-FOS⁺ cell numbers along the anterior-poster axis (AP distance, rows) of every brain region (columns) in the contralateral (Contra., upper panel) and ipsilateral (Ipsi., lower panel) hemispheres.

(B) Cumulative plots of the numbers of C-FOS$^+$ cells along the AP for all brain regions in the

four experimental groups (WT ctrl, *Tbr1$^{+/-}$* ctrl, WT TBS and *Tbr1$^{+/-}$* TBS).

(C) Structure-wise comparison of C-FOS$^+$ cell numbers in the contralateral (upper) and

ipsilateral (lower) brain hemispheres of the four pairwise-compared groups (left to right, *Tbr1$^{+/-}$*

ctrl vs. WT ctrl, WT TBS vs. WT ctrl, *Tbr1$^{+/-}$* TBS vs. *Tbr1$^{+/-}$* ctrl, *Tbr1$^{+/-}$* TBS vs. WT TBS).

Each scatterplot shows the summed C-FOS$^+$ cell number difference along the AP axis (Sum of

ΔC-FOS) versus statistical significance (Adjusted *P* value) for each brain region of the

pairwise-compared groups (also see **Table S2, Figure S7b**). Friedman one-way repeated

measure analyses, followed by post-hoc pairwise tests with Bonferroni correction, were used

for statistical analysis. Vertical and horizontal dotted lines represent the Sum of ΔC-FOS=0 and

Adjusted *P* value=0.05, respectively. Colors represent brain region identities. Open and filled

circles represent ROI-non-corrected and -corrected brain regions, respectively. Each pie chart

above the scatterplot illustrates the proportions of brain regions that fit each condition annotated

in the right panel.

Sample sizes of mouse numbers are identical in (A)-(C) (WT ctrl, n=6; *Tbr1$^{+/-}$* ctrl, n=7; WT

TBS, n=7; *Tbr1$^{+/-}$* TBS, n=7).

1062

Figure 4. Correlations of BLA axonal projection patterns and C-FOS expression of *Tbr1*⁺/⁻ mice.

(A) 3D maps of rewired BLA-derived circuits (left, 10 μm³ binned cubes with binarized oChIEF signals) and differentially TBS-derived C-FOS expression levels (middle, 100 μm³ binned C-FOS⁺ cell numbers) in affected brain regions (right) with the Allen brain region color code. Only the results of contralateral hemisphere are shown herer. Color maps of the scaled difference show the differential oChIEF signal probabilities (left 3D map, ΔoChIEF of *Tbr1*⁺/⁻ − WT) and the difference between the C-FOS modulatory effects of BLA-derived circuits in *Tbr1*⁺/⁻ (ΔC-FOS of *Tbr1*⁺/⁻ TBS − *Tbr1*⁺/⁻ ctrl) and WT (ΔC-FOS of WT TBS − WT ctrl) mouse brains (middle left 3D map, ΔΔC-FOS). Scatterplots on the right panels show the sum of ΔoChIEF (*Tbr1*⁺/⁻ − WT) versus the correlation between ΔoChIEF and ΔΔC-FOS for the

brain regions of contralateral hemispheres. Colors indicate brain region identities. Open and

filled circles represent ROI-non-corrected and -corrected brain regions, respectively.

(B) Examples of brain regions that have positive or negative correlations between rewired BLA-

derived circuits of $Tbr1^{+/-}$ mice and differential TBS-derived C-FOS expression levels. The

oChIEF distributions (WT and $Tbr1^{+/-}$, upper panel), C-FOS$^+$ cell distributions (WT ctrl,

$Tbr1^{+/-}$ ctrl, WT TBS and $Tbr1^{+/-}$ TBS, lower panel), and scaled differences (ΔoChIEF and

$\Delta\Delta$C-FOS, middle panel) along the AP axis are shown. Dashed lines in scaled differences plots

represent the average 99th percentile of scaled differences calculated from the pair-subtraction

population data with 1000 times AP axis reshuffling.

(C) The 3D maps of the results of ipsilateral hemisphere.

(D) The correlation between oChIEF and C-FOS signals at the ipsilateral hemisphere.

Data represent mean ± SEM. Contralateral (A)-(B) and ipsilateral (C)-(D) hemispheres were

analyzed separately. Sample sizes were the same in (A)-(D) (oChIEF data: WT, n=13; $Tbr1^{+/-}$,

n= 14. C-FOS data: WT ctrl, n=6; $Tbr1^{+/-}$ ctrl, n=7; WT TBS, n=7; $Tbr1^{+/-}$ TBS, n=7).

1089

Figure 5. Distinct inter-regional correlations of C-FOS expression in WT and *Tbr1*^{+/−} mouse brains with or without BLA-derived activation.

(A) Cumulative plot showing the probability of all significant inter-regional correlations ($P <$ 0.05, Kendall's rank correlation) of C-FOS expression levels for the four experimental groups.

(B) Matrices showing whole-brain inter-regional correlations of C-FOS expression in the WT ctrl, *Tbr1*^{+/−} ctrl, WT TBS and *Tbr1*^{+/−} TBS groups (left to right).

(C) Plots showing Degree Centrality (DC, y-axis) of each brain region (annotated in the bottom row, x-axis).

(D) Plots showing mean DC of networks (y-axis) with different thresholds for the correlation coefficient r (x-axis, upper panel) or correlation *P* values (x-axis, lower panel).

Brain regions are annotated with color codes in (B)-(C). The sample sizes of mice are the same in (A)-(D) (WT ctrl, n=7; *Tbr1*^{+/−} ctrl, n=8, WT TBS, n=7; *Tbr1*^{+/−} TBS, n=7).

1104

Figure 6. Modular organization of C-FOS expression correlation networks.

(A) Plot showing the differences in modularity (y-axis) between real networks (Q) and shuffled networks ($Q_{shuffled}$) when a different resolution parameter R (x-axis) was used for community detection in the four experimental groups.

(B) The same datasets of inter-regional correlations for C-FOS expression are shown in **Figure 5B**, except the inter-regional pair positions were sorted by their communities. The R values that generate the maximum difference in Q between real and shuffled networks for each experimental group are indicated and they were used to analyze network communities.

(C) 3D maps of binned and averaged C-FOS$^+$ cubes for each community (rows) of the four experimental groups (columns). C-FOS$^+$ cubes are shown in brain region color codes, with their sizes representing the relative normalized density of C-FOS$^+$ cells within binned cubes (100 μm^3) (also see **Figure S7D**). Mean correlations within (the bars at the top of 3D maps) and between communities (the lines under the 3D maps) are annotated according to differential extents (thickness) and polarity (color). Thicker bars or lines represent a higher degree of correlation, whether positive (red) or negative (blue). Sample sizes are the same as in **Figure 5.**

1122

Figure 7. Comparison of default mode networks (DMN) in WT ctrl, *Tbr1*[+/−] ctrl, WT TBS and *Tbr1*[+/−] TBS groups.

(A) The differences of BLA axonal projections into DMN regions for *Tbr1*[+/−] and WT mice, as revealed by ΔoChIEF for 10 μm³ binned cubes.

(B) 3D maps showing differences in density of C-FOS⁺ cells (ΔC-FOS, 100 μm³ bins) for the DMN of experimental group pairs. The heat maps indicate the scaled differences for ΔoChIEF (A) and ΔC-FOS (B).

(C) Structure-wise comparison of DMN properties for four experimental group pairs. Each scatterplot shows the difference in DC versus the difference in participation coefficient (PC) in each DMN brain region for each group pair.

(D) Visualization in spring layout of the DMN networks for the four experimental groups. Brain

regions (nodes) having many reciprocal C-FOS correlations (edges) are closer together.

The color of each node represents brain region identity. The size of the node represents the

normalized C-FOS density of the brain region. Labels "i_" and "c_" represent ipsilateral or

contralateral brain regions, respectively. DMN region color codes are identical for (A), (C) and

(D). In (C), DMN regions labeled in black indicate regions showing a significant difference in

C-FOS$^+$ cell numbers between pairwise-compared groups. Blue-labeled i_ACAv exhibited a

negative correlation between the amount of BLA-derived axonal innervations and its

modulatory effects on C-FOS expression patterns. Thus, the i_ACAv received enhanced BLA

axonal projections that induce net inhibitory effects on neuronal activities in the *Tbr1*$^{+/-}$ mouse

brain. All DMN regions are ROI-corrected. Sample sizes for the DMN network analysis are

identical to those described in **Figure 5A, 5B**. Sample sizes for the DMN ΔC-FOS and DMN

ΔoChIEF analyses are the same as those described in **Figure 4**.

1146

Figure 8. Whole-brain alteration of the amygdalar mesoscopic connectome and synchronization in the *Tbr1⁺ᐟ⁻* mouse brain.

(A) Differences in the amygdalar mesoscopic connectome between wild-type and Tbr1⁺ᐟ⁻ mice are revealed by oChIEF-Citrine expression in a unilateral BLA. *Tbr1⁺ᐟ⁻* mice massively lose contralateral axonal projections of the BLA and exhibit mistargeting of ipsilateral BLA axons.

(B) The global neuronal activities revealed by C-FOS expression are different between wild-type and *Tbr1⁺ᐟ⁻* mice. In general, *Tbr1⁺ᐟ⁻* mice exhibit higher C-FOS expression during wakefulness.

(C) BLA activation elicits opposing effects on C-FOS levels when comparing wild-type to *Tbr1⁺ᐟ⁻* mice. In general, BLA activation potentiates C-FOS levels in wild-type mice, whereas it reduces C-FOS expression in *Tbr1⁺ᐟ⁻* mice.

(D) The neuronal synchronization patterns revealed by C-FOS correlation networks differ between wild-type and *Tbr1⁺ᐟ⁻* mice and are differentially modulated by BLA activation. Neuronal activities are generally less synchronized in the *Tbr1⁺ᐟ⁻* mouse brain. However, after BLA activation, neuronal synchronization is reduced in wild-type mice but increased in *Tbr1⁺ᐟ⁻*

1163   mice. The C-FOS correlation networks shown here are simplified in that brain regions

1164   belonging to the same primary brain areas (colored spots) have been combined. Mean

1165   correlations within and between primary brain areas were calculated and are expressed as spot

1166   size and edge thickness, respectively.

1167

Supporting Information

Figure S1. Workflow for image segmentation of oChIEF and C-FOS signals. (A) oChIEF signal segmentation. Raw images of oChIEF-Citrine signals were subjected to an ilastik pixel classification algorithm to obtain the oChIEF signal probability map. Filtered images were created by applying the oChIEF signal probability map to the raw images. Subtracted images were created by subtracting the filtered image from a Gaussian filter blurred version of the same image. Finally, the subtracted image was subjected to thresholding and binarization to get binarized oChIEF signals.

1178   (B) Detection of C-FOS$^+$ nuclei. Raw images of C-FOS signals were subjected to rolling-ball

1179   background subtraction and then the Imaris spot detection algorithm was used to detect C-FOS

1180   spot signals. Spot locations were then exported for quantification.

1181

Figure S2. Spatial distribution of oChIEF-Citrine-expressing AAV injection sites in wild-type and *Tbr1*⁺/⁻ mouse brains.

(A) Images showing oChIEF-Citrine expression (green) in the AAV injection sites (red spots) of wild-type (left panels) and *Tbr1*⁺/⁻ (right panels) mouse brains. BLA and LA regions are outlined.

(B) Distribution of virus injection sites in CCFv3 space.

(C) Differentiation of virus injection site locations between wild-type and *Tbr1*⁺/⁻ mouse brains by k-nearest neighbors (KNN) algorithm-based classification. Plot showing the parameter number of KNN neighbors (k) versus KNN classification accuracies.

(D) Three k parameters that presented relatively high classification accuracy (arrows in (C)) were used to compare (wild-type versus *Tbr1*⁺/⁻) the real classification accuracy (indicated by the red line) with the classification accuracy distributions of virus injection locations shuffled 1000-times. The results support that the spatial distributions of injection sites are indistinguishable between wild-type and *Tbr1*⁺/⁻ mice.

Figure S3. Volume-based quantification and rewiring of bilateral BLA axon projections in *Tbr1*⁺ᐟ⁻ mouse brain.

(A) Binarized oChIEF-Citrine percentage and C-FOS cell density in each brain region volume were calculated and compared between experimental groups.

(B)-(E) Bilateral BLA axon projections to the same regions of both brain hemispheres were

investigated by determining the ratio of contralateral Citrine signal to ipsilateral Citrine signal

(the C/I ratio) for each brain region.

(A) Scatterplot showing the fold-change ($Tbr1^{+/-}$/WT) in C/I ratio versus significance (Mann-

Whitney U test P value) in each brain region receiving bilateral BLA axon projections in wild-

type mouse brains. Vertical and horizontal dotted lines represent fold change = 1 and U test P

value = 0.05, respectively. Colors represent brain region identities. Open and filled circles

represent ROI-non-corrected and -corrected brain regions, respectively.

(B)-(D) Box plots showing comparisons of the relative fold-change in C/I ratio among brain

regions displaying significantly different C/I ratios ($P<0.05$) between WT and $Tbr1^{+/-}$ mouse

brains. Brain regions were further categorized into the three groups shown in (B)-(D), based on

the cause of differential C/I ratios: (B) brain regions exhibiting impaired contralateral BLA

axon projections; (C) brain regions showing significantly enhanced ipsilateral BLA axon

projections in $Tbr1^{+/-}$ mouse brains; (D) brain regions exhibiting a relative change in bilateral

projection. The regions shown in (B) and (C) are also present in **Figure 2C-2E**, whereas those

in (D) are not.

In the box plots, boxes show median and interquartile range and whiskers extend to show the

rest of the distribution except for outliers. Spots represent individual samples. Asterisks label

parental brain regions of the following offspring regions, bounded by horizontal lines. Regions

in red font were ROI-corrected. Colored bars under the charts represent the brain identities, as

shown in (A). Horizontal dotted lines represent 100% of the relative fold-change of the C/I ratio

compared to the average of the WT group. Sample sizes of mice are the same in (A)-(D) (WT,

n=13; $Tbr1^{+/-}$, n= 15).

1227

1228

Figure S4. Principal component analysis (PCA) of BLA-derived circuits in wild-type and

Tbr1[+/−] **mouse brains.**

(A)-(H) Data on oChIEF percentages in contralateral (A)-(D) or ipsilateral (E)-(H) brain regions

were subjected to PCA and reduced to the first two principal components (PCs).

(A), (E) Plots showing PC locations of individual mouse brain samples projected to the space

defined by PC1 (x-axis) and PC2 (y-axis).

(B), (F) Differentiation of PC locations between wild-type and *Tbr1*[+/−] mouse brains by KNN

classification along the PC1 or PC2 axis. For each KNN classification, the left panel shows a

parameter series of number of neighbors (k) used for KNN plotted against KNN classification

accuracies, and the right panel shows the most appropriate k parameters with relatively high

classification accuracy that were used to compare (wild-type versus *Tbr1*[+/−] data) the real

classification accuracy (indicated by the red line) with the classification accuracy distributions

of PC locations shuffled 1000-times.

(C), (G) Left panels show PC2 (C) or PC1 (G) loadings versus the significance level (U test *P*

value) of volume-based quantification of oChIEF percentage (data from **Figure 2C**) in each

contralateral (C) or ipsilateral (G) brain region. Right panels show correlations of top 10 PC2

(C) or PC1 (G) loadings with their significance level of volume-based quantification of oChIEF

percentage. (D), Contralateral brain regions with the top 10 PC2 loadings and their

corresponding PC1 loadings. (H), Ipsilateral brain regions with the top 10 PC1 loadings and

their corresponding PC2 loadings. Sample sizes of mice are identical in (A)-(H) (WT, n=13;

Tbr1$^{+/-}$, n= 15).

1250

1251

Figure S5. Comparison of C-FOS expression in the affected BLA projection regions between the WT ctrl, *Tbr1*^+/− ctrl, WT TBS and *Tbr1*^+/− TBS groups.

(A) Volume-based quantification of relative C-FOS density in affected BLA projection regions. Box plots show the comparisons of relative C-FOS density among affected regions for all BLA axonal projections in the contralateral (upper panel) and ipsilateral brain hemispheres (lower panel), except for the regions of white matter. Boxes show median and interquartile range, and whiskers extend to the rest of the distribution except for outliers. Spots represent individual samples. A Kruskal–Wallis test followed by a post-hoc Dunn's test was used for statistical analysis. Asterisks label parental brain regions of the following offspring regions, bounded by horizontal lines. Red font indicates ROI-corrected brain regions. Colored bars under the charts represent brain identities. The horizontal dotted line represents 100% of the relative C-FOS density compared to the average of the WT group.

(B) 3D maps of binned and averaged C-FOS+ cubes in affected BLA projection regions for the four experimental groups. The sizes of spots represent the normalized C-FOS densities within binned cubes (100 μm^3). Region color codes are as applied in (A)-(B). Sample sizes of mice are identical in (A), (B) (WT ctrl, n=7; *Tbr1*^+/− ctrl, n=8; WT TBS, n=7; *Tbr1*^+/−TBS, n=7).

1268

Figure S6. Slice-based quantification of oChIEF and C-FOS signals.

(A) Workflow of slice-based quantification. The pixel distributions of oChIEF and the distributions of C-FOS$^+$ cells along the AP axis of each brain region were calculated and compared between experimental groups.

(B) Slice-based quantification of C-FOS expression in affected BLA projection regions for the four pairwise-compared groups (left to right, *Tbr1*$^{+/-}$ ctrl vs. WT ctrl, WT TBS vs. WT ctrl, *Tbr1*$^{+/-}$ TBS vs. *Tbr1*$^{+/-}$ ctrl, *Tbr1*$^{+/-}$ TBS vs. WT TBS). The sample size of each group is the same as in **Figure 3**. Each scatterplot shows the summation of C-FOS distributional differences along the AP axis (sum of ΔC-FOS) versus the significance (Adjusted *P* value) for each brain

1278   region of the pairwise-compared groups. Statistical analysis is the same as described in **Figure**

1279   **3**. Vertical and horizontal dotted lines represent the sum of ΔC-FOS = 0 and Adjusted *P* value

1280   = 0.05, respectively. Brain region color codes are as described in **Figure 3**. Open and filled

1281   circles represent ROI-non-corrected and -corrected brain regions, respectively.

Figure S7. Comparison of volume-based and slice-based analysis of binarized oChIEF signals and summary of the region compositions in the communities of activity correlation networks.

(A)-(B) Binarized oChIEF percentages (volume-based analysis) versus oChIEF pixel distributions along the AP axis (slice-based analysis) used for structure-wise comparisons between the contralateral (A) and ipsilateral (B) hemispheres of wild-type and $Tbr1^{+/-}$ mouse brains. The dataset and plots in the volume-based analysis are similar to those in **Figure 2C** except the compared regions represent a subset from the **Figure 2C** dataset to facilitate comparison with the slice-based analysis. Thus, the brain regions compared in this figure are identical for both volume-based and slice-based analyses. Statistical analysis of volume-based quantification was performed as described in **Figure 2C**. Scatterplots of slice-based analysis showing the summation of oChIEF difference along the AP axis (sum of ΔoChIEF) versus significance (Friedman test P value, determined by Friedman one-way repeated measure analysis) for each brain region. Vertical and horizontal dotted lines represent the sum of ΔoChIEF = 0 and Friedman test P value = 0.05, respectively. Brain region color codes are as described in **Figure 2C**. Open and filled circles represent ROI-non-corrected and -corrected brain regions, respectively. Sample sizes of mice in (A) and (B) are identical to those described in **Figures 2** and **4**, respectively.

(C) Pie charts summarizing the differences in structure-wise comparisons between the volume-based and slice-based analyses shown in (A) and (B).

(D) The various plots represent different communities (rows) in different networks (columns). The plots show the numbers of brain regions involved in different brain areas. The data for the contralateral and ipsilateral brain hemispheres are presented separately, with numbers of bilaterally involved brain regions also indicated.

Supporting Tables and Videos

Table S1 Results for oChIEF-Citrine and C-FOS signals using volume-based analyses.

Table S2 Results for oChIEF-Citrine and C-FOS signals using slice-based analyses.

Table S3 Network analysis based on C-FOS expression.

Video S1 Coronal brain slice series of binarized oChIEF signals (green) superimposed on a CCFv3 template (gray). Wild-type (WT), upper; *Tbr1*$^{+/-}$, lower.

Video S2 Summarized 3D stacks with binarized and binned (10 μm^3) oChIEF signals from 13 WT (upper) and 15 *Tbr1*$^{+/-}$ (lower) mouse brains.