1

WHAT IF…? COMPUTATIONAL ANATOMY IN THE GRANULAR LAYER OF

THE CEREBELLAR CORTEX

Mike Gilbert[a] (corresponding author), Anders Rasmussen[b]

[a]School of Psychology, College of Life and Environmental Sciences, University of

Birmingham B15 2TT, UK

ORCID: 0000-0003-4415-6915

[b]Department of Experimental Medical Science, Lund University, BMC F10, 22184 Lund,

Sweden

ORCID: 0000-0003-4477-369X

Correspondence: m.gilbert.1@bham.ac.uk

ABSTRACT/SUMMARY

Neuronal firing rates code values which the brain computes. It is thought that, at a collective level, information is represented in high resolution which is replicable with computer-like high fidelity and allows computations to be powerful. We challenge that in the cerebellum. We find plausibly that there is a computational effect, unaided and unlearned, of the combination of anatomy and the idea that the cerebellum is the physical form of a computational network of linear units. A unit can be a cell, part of a cell or some other structure. Units randomly sample afferent data. Anatomy mimics random sampling of the whole afferent layer of firing rates, with replacement, notwithstanding that a single unit receives unchanging contact from a tiny fraction of cells in the afferent layer. Cell morphologies and network geometry are explicable as bespoke adaptations for this function. Input data are converted to a (very) low-resolution, coarsely-pixellated internal picture. This is functional. It provides an otherwise elusive mechanism to orchestrate firing of large cell groups – microzones – downstream, and it solves the problem of working blind with unreliable information.

Keywords: cerebellum; theory; hypothesis; model; computational anatomy; cerebellar cortex; granular layer; network; input layer; code

1. **Introduction**

There is now a substantial body of evidence that rate-coded information is linearly conserved in communication between cerebellar neurons [1-8], and that cells are adapted for faithful transmission [9-15]. Firing rates that linearly code behavioural and task metrics have been observed 'at all levels of the cerebellar circuit' [16 p.239, citing 30 sources]. Evidence of linear signalling in the granular layer is rich. To decongest the main text, we include a detailed review in the Methods section to which we cross refer from the main text.

The inner layer of the cerebellar cortex, the granular layer, receives the main excitatory input to the cerebellum, from mossy fibres. Mossy fibres contact both granule cells and large inhibitory interneurons, Golgi cells. Golgi cells in turn inhibit granule cells. The granule cell axon rises into the outer layer, the molecular layer, where it divides in two to form parallel fibres, which lie parallel to the cerebellar surface and other parallel fibres. Parallel fibres make contact in passing on Purkinje cells, which carry the sole output of the cerebellar cortex.

In the traditional learning model, the function of the granular layer is to generate patterns of active and inactive parallel fibres that are repeated if the cerebellum

receives the same mossy fibre input. Following training, the response of Purkinje cells

to parallel fibre firing rates is displaced by the compound effect of learned changes of

synaptic transmission strengths. Replicability of patterns is necessary (and assumed) so

that signals are received by the correct set of modified synapses. Learned changes of

synaptic weights are also a core feature of artificial neural network learning models.

However, it is not clear that input to the cerebellum is ever in the same pattern of

active cells firing at the same permutation of rates. Around 4,000 mossy fibres

innervate the region that supplies parallel fibre input even to a single Purkinje cell.

Moreover, it is difficult to reconcile evidence of linear signalling with network models

that use synaptic weight changes since the function of learning is to displace the naive

response to firing rates.

Neuron-to-neuron transmission involves the complex interaction of a large array of

often co-dependent biophysical and other variables which are specific to the cell types

making and receiving contact. The significance of evidence of linearity is that despite

the seeming clamour, linear relationships are in fact conserved. There are a number of

examples that the cerebellum is adapted to disengage or otherwise compensate for

intrinsic non-linear properties of neurons and synapses [13, 14, 17, 18]. This is –

plausibly, we submit – part of a general strategy of isolating the effect of parameters

that code information and controlling for other variables. In this view, the cerebellum

is a physical implementation of networked linear units. A unit can be a cell, part of a

cell or some other structure. Units sample afferent data. Signalling by a layer of units

at any given instant in time is the physiological form of a data set. Anatomy mimics

random sampling of the whole afferent layer of firing rates, with replacement,

notwithstanding that a single unit receives unchanging contact from a tiny fraction of

cells in the afferent layer. Cell morphologies and network geometry are explicable as

bespoke adaptations for this function. In this contention, neuronal sums are not

complex but have a statistical effect at collective level which is exploited to code

information. Sampling units provide model parameters. We field evidence in the main

text to explain the derivation of model parameters. Parameter values are supplied by

the literature. Our main findings are summarised in the Discussion section. We include

a statement of the relevant parts of the hypothesis at or near the beginning of each

section.

2. Mossy fibres

2.1 Anatomy, 'ranks' and 'files'

Mossy fibres give rise to sagittally-extending collaterals, the direction of the long axis

of microzones (Fig 1). These in turn give rise to multiple distal branches, each ending in

a cluster of terminals [19-22]. Terminal clusters are sagittally aligned and separated by

a randomly variable distance typically not less than 200 μm [19, 20] measured from

edge to edge of clusters (so they are separated by a greater distance measured from

centre to centre). Terminals themselves are intimately and randomly intermingled with

terminals of other mossy fibres.

Accordingly, mossy fibres do not terminate at a point but a region with dimensions:

the field occupied by a terminal cluster. A single mossy fibre gives rise to several

terminal branches and each of those to several terminals, such that a sagittal strip of

the granular layer receives multiple copies of every signal at each of multiple locations.

Cluster size is variable (because it depends on the number of terminals and how tightly

clustered they are), averaging 200 μm sagittally x 150 μm mediolaterally [20]. We term

the average dimensions of a cluster a 'field'. The granular layer is anatomically

seamless. A field is simply a nominal division that we use for modelling. A sagittal row

of 75–100 fields laid end to end has the area (viewed from the cerebellar surface) of a

mid-range microzone [23-26]. We term a row of fields a 'rank'. Ranks are functionally

but not anatomically discrete.

Each terminal is ensheathed by a semi-permeable membrane which restricts

neurotransmitter diffusion, a structure taken together termed a glomerulus. Excitation

of granule cells and Golgi cells by mossy fibres, and inhibition of granule cells by Golgi

cells, all take place here.

109   **Figure 1**

2.2 Hypothesis

Mossy fibre anatomy is an anatomical solution to a computational problem. The cerebellum exploits statistical effects of random sampling called the central limit theorem. For the effects to occur, samples must be independent, also termed sampling 'with replacement' (in the illustration sometimes used, coloured balls selected at random from a bag are put back in the bag between samplings). Clearly, physical replacement is impossible because neither a terminal cluster nor a terminal can be in more than one place at the same time. However, anatomy simulates replacement. The distributed probabilities of the number of terminal branches per mossy fibre are the same as the probabilities of the number of repeat samplings if rates received by a rank were in fact independently randomly sampled by fields. Similarly, the distributed probabilities of the number of terminals per cluster are the same as the distributed probabilities of the number of repeat samplings if rates received by a field were in fact randomly sampled with replacement by individual target neurons. Note that it is the population of *rates* that are hypothetically sampled with replacement, and not *signals*. The population of signals includes copies of rates because of branching and clustering. Copies are an anatomical device to mimic sampling rates with replacement.

2.3 Terminal branching simulates sampling with replacement

If each mossy fibre could terminate in only one field, a mossy fibre 'sampled' by one field clearly cannot also be sampled by any others. Termination by a single mossy fibre in a random number of terminal branches, at multiple locations, removes that restriction.

In theory, sampling with replacement should mean any number of fields can receive the same signal, up to all of them. Clearly this is not true – there is still an anatomical limit. But in practice even with real replacement the number of repeat samplings is constrained by a probability distribution. If, then, the number of terminal branches per mossy fibre has the same probability distribution, branching has the same result as if firing rates received by each field in a rank were in fact a random sample with replacement of firing rates received by the rank as a whole.

We can test that by calculating the probability distribution in the hypothetical case that mossy fibres did not branch terminally and rates were in fact sampled with replacement. The probability, p, that a given rate is selected k times in taking N samples, sample size x, assuming mossy fibres each give rise to an average of m terminal branches, is given by

$$\left[{Nx!}/{k!\,(Nx-k)!} \right] * \left[{1}/{(Nx/m)} \right]^k * \left[1 - \left({1}/{(Nx/m)} \right) \right]^{Nx-k}$$

148   The results are shown in Table 1.

149   **Table 1. Probability p of k repeat selections of a single mossy fibre**

k	0	1	2	3	4	5	6	7	8	9	10	11
p	0.007	0.034	0.084	0.140	0.175	0.175	0.146	0.104	0.065	0.036	0.018	0.008

150
151   Although data are scarce, the number of repeat selections is consistent with what we

152   know about the biological range of the number of terminal clusters per mossy fibre

153   [20, 27]. Estimated by this method, the mean is 4–5.

154 2.4 Terminal clusters simulate sampling with replacement

The principle is the same here but at local, field scale. Mossy fibre terminal clustering,

and random intermingling with terminals of other mossy fibres (Fig 1), means that

contact on any particular cell does not mean the same mossy fibre cannot also contact

cells somewhere else. Again, to approximate independence, it is only necessary for the

number of repeat selections of a signal to have the same probability distribution as if

there was actual replacement.

A practical problem with the strategy of clustering is that a target could receive the

same signal twice (or more). The risk is mitigated by taking small samples. A sample

size of 4 is small enough for a low risk. Granule cells average 4 dendrites, each

receiving contact from a single terminal, and we estimate that Golgi cells receive an

average of 4 inputs to each of their basal dendrites (see derivation of parameters in

the next section). Given m x 100 terminals and an independent and equal chance that

contact is by any terminal, the probability of an n^{th} contact by the same mossy fibre on

a single dendrite is:

$$P_n = \left[\prod_{x=2}^{n} \left(\frac{m - x + 1}{100m - x + 1} \right) \right] \times \frac{y!}{x!\,(y - x)!}$$

where y is the number of inputs to a single dendrite and m is the number of terminals

per cluster, provided by (for illustration) 100 mossy fibres. Calculated in this way,

around one in every twenty samples contains a duplicate, and less than two in a

thousand contain more than one. The actual fractions are smaller because the number

of mossy fibres that innervate a field is higher, around 180 (see Methods, Derivation of

mossy fibre rates).

Quasi-sampling by each field of mossy fibre firing rates received as input to the rank as

a whole, and by cells in each field of rates received by that field, is a biological facsimile

of simultaneous random sampling with replacement by single neurons of input rates to

a whole rank.

3. Golgi cells

3.1 Hypothesis

Despite forming an anatomically seamless dense mesh of axonal filaments that fills the

granular layer, Golgi cells work in groups: ensembles. An ensemble is the population of

a sagittal row of three fields. Groups are not anatomically bounded or even

functionally discrete, but function as a single unit for the anatomical reason that the

axonal range of all Golgi cells in a 3-field row includes all or most of the middle field.

Put another way, each field defines a functional population of Golgi cells – an

ensemble – being those cells with convergent input to that field. Ensembles perform a

computational conversion of mossy fibre rates into inhibition of granule cells. The

computation is (chiefly) a statistical result of random sampling, in two steps. At the

first step, Golgi cell basal dendrites receive contact from a random sample of mossy

fibres that innervate a field (strictly speaking, they randomly sample terminals); and in

the second step, middle-field glomeruli each receive innervation by a random sample

of Golgi cells. The effect of the computation is to synchronise inhibition of granule cells

in the middle field. The strength of inhibition is linearly related to the mean of mossy

fibre firing rates received as input to the entire rank.

Ensembles conflate sagittally anatomically and functionally. Ensembles in a sagittal

strip all return the same output (because they all perform the same computation on

input data with the same frequency distribution). Effectively: the Golgi cell population

of a whole rank functions as a single unit.

202   3.2 The Golgi cell computation

Say a population of values (with any frequency distribution) is randomly sampled.

There are n samples, sample size m. If we take the mean (or sum) of each sample, and

n is large, the sample means have a normal or near-normal distribution, by the central

limit theorem. (This works even if m is a range.) The new distribution is centred on the

same mean as the sampled distribution, but narrower. Modest values of n or m give an

untidy result. However, if the new distribution is then itself randomly sampled, and the

process is repeated, the results get tidier and narrower.

We envisage that in motor circuits during behaviour signalling is analogue, avoiding the

noise caused by discrete signals. So, for example, Golgi cell dendritic signalling is a

sustained state under modulation by fluctuating mossy fibre rates. Similarly, GABA

spillover into the intraglomerular space, almost exclusively the agent of granule cell

inhibition, becomes elevated by a build-up during behaviour, so that this too is at a

sustained level, controlled by afferent Golgi cell rates. Thus, at each step, data are

coded at any moment in a snapshot of a state of change.

We tested the hypothesis *in silico* by simulating a rank. Derivation of model

parameters is described in: Methods, Derivation of ensemble dimensions and model

parameters. Mossy fibre rates sampled by Golgi cells in each field were obtained by

sampling from a distribution representing signals received as input to the rank as a

whole, which were randomly generated subject to constraints on the shape of the

distribution. This was modified to take into account mossy fibre terminal branching

and clustering, random variation of the number of terminals per cluster and clusters

per cell, and the fact that most clusters fall partly inside and partly outside a field (see

Methods, Derivation of mossy fibre rates).

This gave the number of mossy fibre signals, firing rates and the number of copies of

each rate received by each field, and therefore input to an ensemble territory,

populated by 30 Golgi cells, 10 per field. There are ~700 mossy fibre terminals per

field. As far as we know (and we propose), contact by mossy fibres on Golgi cells is at

random at field scale, i.e., inputs to a single basal dendrite are provided by a random

sample of local terminals. The number of physiological samplings is equal to the

number of basal dendrites per field; sample size is the number of mossy fibres which

make contact on each dendrite. The effect is sustained dendritic depolarisation under

modulation by a linear function of input rates (see Methods, Linear transmission of

mossy fibres to Golgi cells), which we take as the mean.

Taking the sample means generates a new set of values (Fig 2 column 2) with a smaller

range and centred on the mean, which is linearly related to the mean of the sampled

distribution. Basal dendritic charge transfer is passive and little filtered [11, 28].

Integration of dendritic states at the Golgi soma is represented by taking the mean of

dendritic charge, for each simulated cell. Golgi cell firing rates have a linear

relationship with the amplitude of depolarising current [29], suggesting (and we

assume) a linear conversion of somatic membrane potential into the Golgi cell firing

rate.

The output is a third data set, Golgi cell firing rates (Fig 2 column 3), on the same

normalised scale. Each glomerulus in the middle field receives innervation from a

random subset of Golgi cells afferent to the field, and accordingly independently

randomly samples Golgi cell firing rates. Golgi cells inhibit granule cells in the

glomerulus. A granule cell extends a single dendrite into a glomerulus, typically

receiving synaptic contact from two or three Golgi cells [30]. However, inhibition of

granule cells by Golgi cells is almost entirely by GABA spillover into the intraglomerular

space [5], via extrasynaptic receptors [31, 32] (see Methods, Linear transmission of

Golgi cells to granule cells). This effectively increases the Golgi cell to granule

convergence ratio, since the number of Golgi cells that are functionally afferent to a

granule cell is not the number that make direct contact, but the number afferent to a

glomerulus. The membrane around a glomerulus restricts GABA diffusion so that

signalling is by modulation of the intraglomerular concentration of GABA, which in turn

reflects the rate it is released. The rate it is released by a single cell is proportional to

the firing rate of that cell. The sum of rates is proportional to the mean. The

convergence ratio is sample size; the number of samples is the number of glomeruli

per field ($\sim$700, the same as terminals). Intraglomerular GABA concentration is under

continuous modulation at a sustained level which we took to be a linear function of

the mean of afferent Golgi cell rates, given by the sample means. The number of Golgi

cells afferent to a glomerulus is unreported, as far as we know. We take it as a random

number in the range 8–12, generated for each glomerulus. We tested other ratios to

confirm that the ratio does not affect the ensemble computation (Supplementary

Materials).

This generates a final population of values (Fig 2 columns 4 and 5), the output of the

ensemble conversion. There are 700 outputs, representing glomeruli in the middle

field. All granule cells that extend a dendrite into the same glomerulus receive equal

inhibition (of that dendrite).

3.3 Results

3.3.1 An ensemble

There is a progressive narrowing of the data range at each step. Inputs to the system are converted to narrowly-grouped and normally-distributed outputs. The narrow output range indicates that inhibition is locally synchronised, i.e., granule cells in a field receive co-modulated inhibition. Inhibition has a linear relationship with mossy fibre rates, acting through the ensemble conversion. These results are independent of the shape of mossy fibre rates.

Figure 2

3.3.2 A rank

We next used the ensemble conversion to derive inhibition of granule cells in a sagittal row of 100 fields, a rank. The results are shown in Figure 3. As ensembles all perform the same computation and take as input (at any moment) the same frequency distribution of input rates, or nearly the same distribution, they all return the same output, or nearly the same output. There are 700 outputs per field. The mean of outputs, taken for each field, has a narrow range centred on the mean of mossy fibre input rates to the whole rank. Field-to-field variation of output appears to be inherent

in the mechanism – i.e., caused by uncertainty of the outcome of random sampling –

so it has a predictable range which the system can be built to tolerate.

Conclusion: Golgi cells in a rank act as a single functional unit. Ensembles conflate in

the sagittal direction functionally as they do anatomically. As a result, the whole

granule cell population *of a rank* receive synchronised inhibition which is linearly

related to the mean of mossy fibre rates.

Figure 3

299 4. Granule cells

300 4.1 The fraction of active granule cells is reliably regulated by

301 feed-forward inhibition at low density

302 4.1.1 Hypothesis

Anatomy simulates independent random sampling by a single granule cell of the

population of mossy fibre input rates to the whole rank (section 2). There is a

competition for influence between mossy fibre excitation and Golgi cell inhibition of

granule cells. The competition takes place in the glomerulus. During behaviour, this

takes place ceaselessly. The strength of inhibition is synchronised between glomeruli

that populate a rank (section 3), so that all mossy fibre signals compete, at any

moment, against the same inhibitory threshold. The outcome in each glomerulus is

independent of the outcome at other terminals, including other terminals that

innervate the same granule cell. To fire, a granule cell must receive a minimum

number of mossy fibre inputs that prevail in the glomerular competition. The

probability that enough prevail depends, at any moment, on the strength of inhibition.

Since inhibition is synchronised, the probability is the same for all granule cells at any

time. The number that fire is very near the expected proportion (the number predicted

by probability) because granule cells are numerous. Inhibition has a linear relationship

with mossy fibre firing rates (section 3). This is always conserved (by the Golgi cell

ensemble conversion) regardless of the mossy fibre distribution (section 3). So, the

probability is invariant, and therefore so is the proportion of active granule cells.

Note: the idea of an all-or-nothing result of the intraglomerular 'balance' of excitation

and inhibition has appeared previously [33].

4.1.2 Anatomy

Granule cells receive input to each dendrite from a single mossy fibre. Granule cells

have 3–5 dendrites, averaging 4 (the simulation assumes 4, but the result is the same

with 3–5). Each dendrite of a single granule cell extends into a different glomerulus.

Estimates vary of the number of granule cells that receive contact from a single mossy

fibre terminal. 50 is mid-range [34, 35], but estimates vary significantly [2]. 50 is

consistent with our estimates of the number of terminals and granule cells in a field.

The granule cell sampling unit is the whole cell since it receives only a single input rate

to each dendrite. Sample size is therefore 4, and the number of samples per field is the

number of granule cells, which we estimate at 8,750 (see Methods, Derivation of

mossy fibre rates).

4.1.3 Feed-forward inhibition regulates the probability that granule cells

fire

Excitation and inhibition compete for influence in a glomerulus. It is unclear if there is

always an emphatic winner, but a swing in the dominant direction may be amplified by

positive feedback, acting through presynaptic receptors. Activation of presynaptic

$GABA_B$ receptors on mossy fibres inhibits glutamate release [36], while activation of

presynaptic mGluR2 receptors on Golgi cells inhibits GABA release [37]. We postulate

that only mossy fibre signals which prevail in the glomerular competition contribute

significantly to granule cell depolarisation. To fire, a granule cell must receive a

minimum number of competitive mossy fibre signals – thought to be three [38, 39].

Certainly, one is too few [5, 40, 41].

The probability that a given granule cell receives three competitive signals and

therefore fires is a function of the percentage of mossy fibre signals that are

competitive. The expected number of granule cells that fire in a field is given by

$$E_{fld} = T \sum_{x=j}^{n} \left[\frac{n!}{(n-x)!x!} [P(e > i)]^x [1 - P(e > i)]^{n-x} \right]$$

where T is the total number of granule cells per field, n is the number of dendrites per

granule cell, and j is the minimum number of dendrites that must receive dominant

excitation, $e > i$. The proportion of granule cells that fire is near the expected number

by the law of large numbers. The percentage of mossy fibre signals that are

competitive is always the same because inhibition scales with the mean of mossy fibre

rates (because Golgi cells translate mossy fibre firing rates into proportional

inhibition). As the percentage is unvarying, so is the probability.

Probability is an abstraction with no direct physiological counterpart. The simulation

does not include a binomial function but derives output from input by comparing

excitatory to inhibitory input to each glomerulus, the way we claim it is derived

physiologically. The equation is simply shorthand – biology exploits chance and large

numbers to generate the same result (as if inputs were passed through the equation).

4.1.4 Does physiology have that outcome?

To test that, we incorporated the Golgi cell ensemble conversion into a model that

recodes mossy fibre signals into granule cell signals. The model compares mossy fibre

signal frequency with inhibition at each of 4 dendrites of each of 8,750 granule cells

per field, in each of 100 fields. Mossy fibre input rates to each field are derived per

section 3 and Methods, so taking into account, *inter alia*, that a field receives a

randomly variable number of copies of most signals, and most terminal clusters lie only

partly inside a field. Inhibition – outputs of the ensemble conversion, 700 per field – is

provided by the Golgi cell model. Excitation and inhibition received by each glomerulus

are randomly paired. Pairing is used to identify the subset of granule cells that fire

(cells that receive at least 3 mossy fibre signals which exceed the inhibitory threshold).

Output of the model is the number of granule cells that fire in each field, the mean of

mossy fibre rates they each receive (the sample means), and the mean of the sample

means in each field ('field means'). The results are shown in Figure 4.

4.1.5 Results

Feed-forward inhibition – mossy fibre $\rightarrow$ Golgi cell $\rightarrow$ granule cell – reliably and closely

regulates the fraction of granule cells that are active at low density. The Golgi cell

computation may be the primary regulator of the number of granule cells that fire.

Field-to-field random variability of the number is inversely proportional to the fraction

that are active, suggesting that, if primary regulation is in this form, the fraction is low.

We would nonetheless expect to see greater precision, suggesting the physiological

presence of a supplementary mechanism(s) that we have not represented in the

model.

A local feedback mechanism may be provided by granule cell ascending axon contact

on Golgi cells. The ascending portion of the granule cell axon makes synaptic contact

on local Golgi cells, on basal dendrites [11]. A Golgi cell receives ~400 functional

synaptic contacts from this source, around 25% of the granule cells that a Golgi cell

receives contact from in total. During sustained mossy fibre discharge, synapses from

local granule cells contribute equivalent charge to Golgi cells as mossy fibre synapses

and 'display similarly fast kinetics' [42]. Short-latency inhibitory feedback may add

homeostatic fine adjustment to regulation at local level, at field scale.

It is a long-standing idea [43] that granule cell activity may be regulated by feedback

via parallel fibres, which contact Golgi cell apical dendrites. We have previously

proposed that gap junctions between apical dendrites may provide the means for

Golgi cells to detect the density of parallel fibre activity [44]. Feedback by this pathway

is not sensitive to local fluctuation of granule cell active numbers, and may be the

weakest regulator.

The results are the same with any (tested) distribution of mossy fibre rates and also

whether mossy fibres are all active or some are active and others are not. Active cells

can be in any random permutation.

Figure 4

4.2 Granule cells encode mossy fibre rate information

collectively in rank-sized strips

4.2.1 Hypothesis

Anatomy simulates independent random sampling by a single granule cell of the

population of mossy fibre input rates to the whole rank (section 2). The granule cell

firing rate is a linear function of depolarising somatic charge, in turn a linear function

of the mean (or sum) of dominant excitatory input rates. The linear relationship is

undiluted by inhibition because dominant signals transiently displace a significant

effect of inhibition. Derived in this way, granule cell rates are collectively well

predicted by mossy fibre rates, by the central limit theorem. This provides the

substrate for granule cells to collectively encode mossy fibre rate information. Code

groups are functionally defined by the topography of mossy fibre input to the

cerebellum, as Golgi cell networks are. The output of a rank is the smallest unit of

granule cell encoded information. The permutation of active mossy fibres, and of

granule cells, and how firing rates are distributed among active cells, all code nothing.

4.2.2 Do granule cell firing rates – the output of a rank – have the

hypothesised features?

We ran the same simulation as section 4.1 to test whether, and how, mossy fibre rates

are related to granule cell rates, and if fields in a rank all have the same output. We

take granule cell firing rates to be a linear function of the mean of dominant input

rates to each active cell, which we take as the mean itself. Field means are therefore

the mean of granule cell firing rates in each field. The results are shown in Figures 5

and 6. We present evidence relating to mossy fibre → granule cell linear transmission

in the Methods section (Methods, Linear transmission of mossy fibres to granule cells).

The narrowing and predicted shape of the granule cell frequency distribution is evident

in all conditions. The field means approach a straight line, the test that a rank operates

as a single functional unit. When mossy fibre rates are normally distributed (SD 12.5

Hz), the granule cell distribution is very narrow (and normal if rates are counted in

narrower bins) and the field means are very close to a straight line. The field means

follow the mean of mossy fibre rates at a precisely fixed distance. When mossy fibre

rates have a random uniform distribution, the field means remain almost a straight

line, and the distance from the mossy fibre mean varies with the size of the mossy

fibre range.

A straight line indicates that fields all return the same output – not the same firing

rates, but the same narrowed range, clustered around the mean, and attracted to a

normal distribution (more evident with narrowly and/or normally distributed mossy

fibre rates). This holds even though a single granule cell receives a tiny fraction of

mossy fibre rates, and each field receives only a small random sample of mossy fibre

signals received by a rank, and regardless of the shape of the mossy fibre distribution.

This is not in itself necessarily in fatal conflict with the traditional cerebellar learning

model, but has the alternative interpretation that granule cell firing conserves mossy

fibre rate information, coded in the output of a whole rank. This is therefore the

minimum resolution of granule cell information, but not the minimum unit of code.

The same information is coded in any random sample of granule cell signals over a

modest minimum number. It is immaterial (i.e. codes nothing) which particular cells

are active or which cells fire at what rates, or even what rates those are, provided any

random sample of rates is as if drawn from the same frequency distribution, with

replacement. The output of a rank can be likened by analogy to a pixel: it is only

divisible into smaller pieces of the same information.

452   **Figures 5 and 6**

5. **Discussion**

There is a general expectation that the representation of information in the brain is in high resolution and replicable with high fidelity, and that those features make it precise and provide the inputs to sophisticated computations. We challenge that. We propose in this paper:

i) There is a computational effect, unaided and unlearned, of the triad of cell morphologies, network architecture and linear transmission, for which the granular layer is highly adapted.

ii) Recoding in the granular layer exploits the central limit theorem. Statistical properties of granule cell rates code mossy fibre rate information. The pattern of active granule cells, and which cells fire at what rates, code nothing.

iii) High-dimensional mossy fibre information recodes as a very low-resolution granule cell code. Large cell groups code information in their collective activity. At any instant, a whole group codes the same information. A code group functions as a single unit and spans the cerebellar network sagittally. So, information has spatial dimensions, the dimensions of a code group.

iv) Granule cell code groups mirror microzonal organisation of the molecular layer.

Accordingly, granule cell encoded information has a striped pattern viewed from the

cerebellar surface. This can be likened by analogy to pixellation.

v) While firing of a whole code group is the minimum unit of information, it is not the

minimum unit of code. Inside pixel boundaries the same information is coded in any

random sample of granule cell firing rates.

vi) There is a huge sacrifice of data resolution on entry to the cerebellum, counter to

the general expectation that more data should inform more powerful brain

computations. This is functional. Coded in this form, granule cell data can drive

coherent firing of large cell groups downstream. Microzones lie at right angles to

parallel fibres, presenting a large target. The form of the granule cell code means the

same information is received by all locations of sagittal plane, so that microzone-

grouped Purkinje cells receive the same information without receiving the same

signals. An estimated 350,000 parallel fibres pass through a Purkinje cell dendritic

territory [45-48]. If even only a small fraction of parallel fibres is active (0.5%), and

allowing for the fact that only one in two make contact, and that a majority of

synapses are severely depressed or silent [49, 50], a single Purkinje cell still receives

around 200 parallel fibre signals at working synapses at any time. A granule cell code in

this form provides a means to modulate firing of Purkinje cells in unison by a chain of

rate codes. There is evidence that Purkinje cells linearly code the strength of granule

cell inputs [6-8] and behavioural metrics [3, 51-54].

vii) A statistical code may be a physiological strategy to mitigate variable performance

of single neurons and synapses, a long-known impediment to precise and replicable

biological information capture. Learning models depend on pattern replication for

signals to match up with the correct synaptic weights. Yet it is known that single cells

are not metronomically consistent performers. The response of cells of the same type

can vary in the same conditions, and so can the response of a single cell to the same

stimulus at different times. Moreover, the granular layer is *designed* to amplify the

difference if the pattern of active mossy fibres should vary only by a modest number of

cells. Indeed, even that is not necessary – the same cells firing at different rates also

generates a different pattern of active granule cells, and therefore parallel fibres. Small

differences of input are converted by decorrelation to wholesale changes to output.

Possibly, and we submit, the cerebellum actively eschews reliance on pattern

replication in favour of group code which gives a more (because statistically) reliable

outcome.

A perennial issue for network theory is that biological networks do not seem to have

much information about the signals they receive. Input to the cerebellum is eclectic

and in multiple modalities. In some cases, a single granule cell can receive multi-modal

input. In the traditional cerebellar model, supervision solves this problem. It is

unnecessary for the cerebellum itself to discriminate because it is an assumption of the

model that externally sourced instruction signals train the correct response.

We propose instead the solution that resolution of high-dimensional input data is

sacrificed. The cerebellum disregards source and modality data below a physical scale

which mirrors topography. It is therefore unnecessary to discriminate below that scale.

Mapping studies show that mossy fibre signals (evoked by cutaneous stimulation)

share the topographical organisation of climbing fibres [55-57]. At sub-topographical

dimensions, signal source and type are simply discounted. Therefore, networks do not

need to be adapted to the source or type of input they receive – anatomically,

networks are interchangeable, so network design can be modular. This feature makes

it functionally flexible, allowing the almost unchanging basic cerebellar wiring to

survive dramatic upheavals of the vertebrate phenotype [58, 59].

521   **METHODS**

DERIVATION OF MOSSY FIBRE RATES

The number of mossy fibres that innervate a rank

As far as is known, fields receive a random sample mossy fibre input rates to a rank. To

simulate that, we need the size of the sampled population – the number of mossy fibre signals

received by a rank – and sample size, the number received by a field.

Despite the repeating 'crystalline' architecture of the cerebellar cortex, at local level there are

anatomical random variables. Variables are: i) the number of mossy fibres that innervate a

field, ii) the number of terminals they each end in, and iii) the number of those that are inside

field limits.

The number of mossy fibres afferent to a field has been estimated at 100 [20]. However, that

number is a field-sized fraction of the general population. The actual number is larger because

mossy fibres end in a dispersed group (cluster) of terminals, and most groups straddle field

limits.

We work from an estimate of the total number of terminals per field, which is, in turn, derived

from convergent estimates of the number of granule cells per field. (Estimate 1) There are an

estimated 1.92×10^6 granule cells per μl [45]. The number of fields which fit in that volume is

$(1,000/150)^2 \times (1,000/200) = $ ~222, assuming a field depth of 150 μm [20]. The number of

granule cells in a field is, therefore 1,920,000/222 = 8,649. (Estimate 2) An estimated 350,000

parallel fibres pass through a single Purkinje cell's territory [45]. Parallel fibres run about 3 mm

in each direction from the granule cell soma [60, 61]. The Purkinje cell dendritic arbour fills the

molecular layer vertically and averages ~200 μm wide. Therefore, 6,000/150 = 40 fields fit in a

row that provides parallel fibres that pass through a Purkinje cell dendritic territory, giving

350,000/40 = 8,750 granule cells per field. Estimates vary of the number of granule cells which

extend a dendrite into a single glomerulus. 50 is mid-range. (8,750/50) x 4 = 700 glomeruli per

field.

Granule cells have 3–5 dendrites, averaging 4, and receive input to each of them from a single

mossy fibre (so an average granule cell receives 4 inputs). Estimates of divergence of a single

mossy fibre terminal onto granule cells vary [2]. It is unclear how much this reflects biological

diversity versus differences in opinion and methods. Here we use a mid-range estimate of 1:50

[34, 35]. Assuming a mossy fibre terminal receives a single granule cell dendrite from each of d

granule cells, and given an average of 4 dendrites per granule cell and z granule cells per field,

the total number of mossy fibre terminals in a field is $N = (z \times 4)/d = 700$ given $z = 8,750$ and d

$= 50$. A convergent estimate is provided by multiplying the per field number of mossy fibres by

the average number of terminals per cluster.

Cluster size varies. We take the range as 4–12 terminals per cluster [20]. We assume the

number varies at random and cluster sizes occur with equal probability. A maximum of 7 fits in

a field (because a field shares the dimensions of the area that encloses a mean cluster). Fields

are not anatomically bounded, so many clusters straddle field limits. To reflect that, we

assume there is a 1/7 probability that a 7-terminal cluster (or larger) has 7 terminals inside a

field and also the same probability of any other number. A 6-terminal cluster has a 2/7 chance

that all 6 terminals are inside and a 1/7 chance of any other number, and so on.

Table 1 shows the derivation of the expected (i.e., predicted by probability) ratios of terminals

contributed to a field by each cluster size (R1), not counting terminals that lie outside when a

cluster straddles field boundaries; and the ratios of mossy fibres contributing $n = 1$–7 terminals

(R2), regardless of cluster size. R1 ratios are given by the sum of numbers in each row. So, for

example, the ratio of cells that end in 8 terminals to cells that end in 5 is 28:25. R2 ratios are

given by the number of appearances of each number in the grid – so the expected ratio of cells

which contribute 7 terminals to the number which contribute 4 is 6:12. The ratios effectively

give the distributed probability of possible states. For example, there is a probability of 10/(6 +

8 + 10 + 12 + (3 x 9)) = 0.159 that a mossy fibre contributes 5 terminals to a field (regardless of

cluster size). Similarly, there is a probability of ((6 x 28)/((6 x 28) + 27 + 25 + 22))/7 = 0.099 that

a given cell contributes 7 terminals to a field.

Table 1. Derivation of expected ratios

size	number of terminals inside							R1
12	7	6	5	4	3	2	1	**28**
11	7	6	5	4	3	2	1	**28**
10	7	6	5	4	3	2	1	**28**
9	7	6	5	4	3	2	1	**28**
8	7	6	5	4	3	2	1	**28**
7	7	6	5	4	3	2	1	**28**
6	6	6	5	4	3	2	1	**27**
5	5	5	5	4	3	2	1	**25**
4	4	4	4	4	3	2	1	**22**
R2	6	8	10	12	9	9	9	

If R1 ratios are denoted by a series $x_4, x_5 \ldots x_{12}$, and a given element in the series is x_j, the

fraction of terminals provided by a given cluster size is $x_j/(x_4 + x_5 + \cdots + x_{12})$, and this

fraction is multiplied by N to give the number of terminals. The number of terminals is divided

by the expected mean number of terminals that fall inside field limits (for that cluster size) to

give the expected number of mossy fibres that provide them. The number of j-terminal mossy

fibres that innervate a field is therefore given by

$$N \left[{x_j} \middle/ {\sum_{i=1}^{n} x_i} \right] \middle/ \left[{x_j}\middle/{7} \right]$$

The results are shown in Table 2. The equal number of mossy fibres (mfs) in each column is

what we would expect, because we made it an assumption that all sizes occur with equal

probability, and mf values are expected values – i.e., in proportions predicted by their

probability. The mf total is the expected number of mossy fibres afferent to a field (which

contribute at least one terminal).

Table 2. The expected number of mossy fibres which innervate a field

cluster size	12	11	10	9	8	7	6	5	4	total
terminals	81	81	81	81	81	81	78.1	72.3	63.6	700
mfs	20.25	20.25	20.25	20.25	20.25	20.25	20.25	20.25	20.25	182.25

590   R2 ratios – the relative incidence of afferent mossy fibres that contribute 7 terminals to a field,

591   6 terminals, 5 and so on, down to 1 – and the expected number of mossy fibres that contribute

592   them, are shown in Table 3.

593   **Table 3. The number of mossy fibres which contribute *n* = 1–7 terminals to a field**

n	7	6	5	4	3	2	1	total
R2	6	8	10	12	9	9	9	63
R3	42	48	50	48	27	18	9	
mfs	17.34	23.15	28.92	34.72	26.04	26.04	26.04	182.25
%age	9.52	12.69	15.87	19.05	14.29	14.29	14.29	

R3 is *n* x R2, giving the ratio of terminals provided by mossy fibres with *n* terminals inside a

field. %age is given by (R2/63) x 100. this is the percentage of mossy fibres with *n* terminals

inside a field.

From here, the number of mossy fibres that innervate a rank is given by 182.25 x 100 divided

by the mean number of clusters per mossy fibre per rank. If a rank receives, on average, five

clusters per mossy fibre, ~3,645 mossy fibres innervate a rank. This is the estimate used in the

simulation unless stated otherwise.

Assumptions. Cluster sizes occur with equal probability. The probabilities of each outcome

when a cluster straddles field boundaries (how many terminals are inside and how many

outside) are as stated. A single mossy fibre ends in an average of 5 terminal clusters per rank.

Mossy fibre input rates to a field

Terminal clustering means a field receives multiple copies of most signals. The proportions of

mossy fibres afferent to a simulated field that contribute each possible number of terminals

(1–7) are generated in simulated steps. In the first step, the simulation randomly generates,

for each field, the cluster size ratio. This means, the relative proportions of afferent mossy

fibres (which contribute at least 1 terminal to a field) that end in 4 terminals, 5, 6 and so on up

to 12, assuming they occur with equal probability, disregarding how many terminals fall inside

and outside field boundaries. In step two, the ratio of cluster sizes is weighted to reflect the

greater probability that large clusters contribute more terminals to a field. In step 3, ratios are

converted to the fraction of terminals contributed by each cluster size. When multiplied by the

total number of terminals, this gives the number of terminals contributed by each cluster size.

In step four, the values obtained in step three are divided up between cells that contribute a

single terminal, 2 terminals, 3 and so on, to reflect straddling of field limits. The proportions

give the number of mossy fibres afferent to a simulated field that contribute a single terminal,

two, three and so on up to seven, the maximum.

There is no field-level effect of mossy fibre terminal branching. Terminal clusters arising from

the same mossy fibre are separated by a minimum distance, so that a field cannot contain

terminals of more than one cluster per mossy fibre.

Terminal clustering means that fields contain a randomly variable number of copies of most

signals. In the simulation, copies are added after sampling of mossy fibre rates to obtain rates

received by a field. An alternative order would be to add copies to the sampled population and

then sample that. We follow anatomy: the number of *rates* received by a field is equal to the

number of afferent mossy fibres; the number of *signals* contains duplicate rates.

DERIVATION OF ENSEMBLE DIMENSIONS AND MODEL PARAMETERS

4 Purkinje cells fit in a field, based on field size, thickness of the Purkinje cell dendritic plate – 9

μm [62] – and the distance between them – ~40 μm [63]. 'In man, the ratio [of Purkinje cells

to Golgi cells] is 1꞉1.5, while in the monkey and cat it is almost 1꞉1.9 and in the rat 1꞉3.3' [64].

Possibly, estimates are low because smaller Golgi cells at deeper level were not counted [63

p.121]. We take it that there are 10 Golgi cells per field, a ratio of 1:2.5 with 4 Purkinje cells.

Golgi cell morphology creates ensemble dimensions. The Golgi cell axon ramifies profusely,

giving rise to a dense plexus. The axonal field is sagittally elongated – that is, in the direction of

the long axis of a microzone (mean range 650 +/- 179 μm by 180 +/- 40 μm [in mice: 65]) – and

is the depth of the granular layer, which it fills vertically. Fine, beaded axon fibres enter

glomeruli, where they inhibit granule cells [66]. Because of the dimensions and orientation of

the Golgi cell plexus, all Golgi cells in a sagittal row of three fields inhibit substantially the

whole of the middle field. Each field defines a functional population of Golgi cells – an

ensemble – being those cells with convergent input to that field. A single Golgi cell contacts

thousands of granule cells [66, 67].

Contact by mossy fibres on a Golgi cell is on basal dendrites. 4–6 basal dendrites arise from the

Golgi cell soma. We assume 5, so a population of ensemble-grouped Golgi cells has a

combined total of 30 x 5 = 150 basal dendrites. The number of mossy fibres which contact a

Golgi cell is unreported. Dendrites radiate from the cell body. A single dendrite has a range of

50–100 μm [63]. The longest dimension of a field – which contains ~700 mossy fibre terminals

– is 200 μm. We estimate the mean number of mossy fibres that contact a single dendrite is 4,

so 20 per cell, or ~3% of the population of terminals (or 0.6% per dendrite).

Glomeruli individually receive innervation from a random sample of Golgi cells afferent to a

field. It is unknown how many Golgi cells innervate a glomerulus.

LINEAR TRANSMISSION OF MOSSY FIBRES TO GOLGI CELLS

It is part of our proposal that Golgi cell basal dendritic signals linearly reflect mossy fibre rates,

so that mossy fibre rate information is conserved proportionally in Golgi cell firing rates. Golgi

cells are large interneurons whose cell bodies and basal dendrites lie in the granular layer.

Mossy fibres directly contact Golgi cell basal dendrites. Golgi cell firing frequency increases

linearly with the amplitude of depolarising current [29 p.845]. 'Golgi cells can follow peripheral

signals in a continuous fashion, modulating their frequency with the intensity of the stimulus

[citing [68, 69]]' [70]. As a result, during movement, the 'output of Golgi cells varies smoothly

as a function of input....They are capable of modulating their output to reflect their input with

little noise or signal distortion' [12]. 'Sensory-evoked Golgi-cell inhibition scales proportionally

with the level of mossy fiber excitatory synaptic input', such that inhibition reliably conveys

mossy fibre rate information [4]. On this evidence: mossy fibre rate information is conserved

proportionally in Golgi cell firing rates.

Golgi cells extend their basal dendrites into glomeruli [66, 71]. Contact on them by mossy

fibres is multi-synaptic [63], contributing to a reliable, rapid (submillisecond) response. The

fast rise time and weak distance dependence of the amplitude and timing of Golgi cell EPSCs

evoked by mossy fibre signals, suggests that basal dendrites introduce relatively little filtering

[11, 28], as expected for large-diameter dendrites. We note that the pause in Golgi cell firing

following a discrete stimulus under anaesthesia [4, 72, 73] disappears from Golgi cell

recordings during locomotion [69].

During behaviour in freely-moving animals, mossy fibre activity is dense: a high fraction of

mossy fibres are active [74]. Both mossy fibres [75-77] and Golgi cells [69] have been reported

to fire with a sustained, time-varying signature. In the behaving animal, Golgi cell basal

dendritic membrane potential is a continuous variable under modulation by sustained inputs,

we submit. As charge transfer to the soma is passive, polarisation of the soma is likewise

sustained, under modulation by dendritic states.

It is worth noting that very high mossy fibre rates can be generated in experimental conditions.

However, the typical physiological range is 50–300 Hz [76]. In the simulations, we use the

physiological rates, within stated constraints on the shape of the distribution.

Other control of Golgi cells? Do Golgi cells receive a significant influence of input from other

sources? The evidence is incomplete but currently suggests other influence is absent or weak.

Contrary to early reports, neither Purkinje cells nor climbing fibres contact Golgi cells [78, who

give references], and only a modest minority of inhibitory inputs to Golgi cells (if any) are from

molecular layer interneurons [79], which generate weak synaptic currents [80], consistent with

either extremely weak or wholly absent innervation [81]. There is conflicting evidence whether

Golgi cells make inhibitory synaptic contact on each other [81, 82]. Our simulation does not

include an effect on Golgi cell firing by other sources of input.

LINEAR TRANSMISSION OF GOLGI CELLS TO GRANULE CELLS

'In the adult rat cerebellum, each granule cell dendrite receives 2.6 ± 0.55 synaptic contacts

from Golgi axon terminals' [30] citing [35].[1] However, the large majority of inhibition (98%) is

by spillover [5], where neurotransmitter released into the synaptic cleft spills out into the

glomerular space. Golgi cells release GABA, an inhibitory neurotransmitter. This is detected by

high-affinity $GABA_A$ receptors located perisynaptically and extrasynaptically on granule cells

[31, 32]. Even synaptically-received signals have most of their effect (i.e., the large majority of

charge transfer is mediated) via spillover.[2]

Golgi cells fire spontaneously [83] at a time-varying rate in the behaving animal, so that

downstream, a glomerulus receives continuous convergent input. As a result, there is a

sustained build-up of glomerular GABA during behaviour [84] at an adjustable concentration

controlled by the mean of afferent Golgi cell firing rates [85]. A glomerulus randomly samples

Golgi cells afferent to a field. Golgi cells provide the only inhibitory input to granule cells.

Signalling by spillover is sometimes assumed to be ambient and slow. However, the action of

glomerular GABA spillover has a fast phasic component – not as fast as synaptic transmission

(~1 ms) but with a rise time of only a few milliseconds [85]. Unlike the spiky appearance of

synaptically-induced inhibitory postsynaptic currents, spillover [4] generates a sustained

---

[1] Data are sparse. The original data are an estimate derived from visual inspection of slices thirty years ago so far without later corroboration, except from Rossi and Hamann 1998, who derive a lower estimate but propose the disparity is because their estimate is too low.
[2] Because of the long duration of IPSCs generated by spillover (recorded in slices), the total charge carried is three times that of IPSCs generated by directly connected terminals. 30. Rossi, D.J. and M. Hamann, *Spillover-mediated transmission at inhibitory synapses promoted by high affinity alpha6 subunit GABA(A) receptors and glomerular geometry.* Neuron, 1998. **20**(4): p. 783-95.

outward current. The response time to multiple inputs is likely faster than the experimentally

isolated response to a discrete signal (because the combined rate of GABA release is higher).

Golgi cell oscillations. Golgi cells fire autonomously. Under anaesthesia, firing falls into a slow

oscillating pattern [86]. Under excitatory synaptic input, however, this disappears [82].

Anaesthesia 'has a strong influence on spontaneous activity of Golgi cells' [78]. Discharge at

low rates with irregular timing under anaesthesia [83] is replaced by higher rates with more

regular timing without anaesthesia [68, 69, 87]. (This paragraph is a cautionary note about

evidence obtained under anaesthesia, which has been used to argue that oscillations provide a

form of signalling.)

LINEAR TRANSMISSION OF MOSSY FIBRES TO GRANULE CELLS

There is significant evidence that granule cell firing rates are linearly related to input rates, but

also evidence thought to be unfavourable to that conclusion. We take them in turn.

The short and equal length of granule cell dendrites would suggest light and equal filtering.

The mossy fibre-granule cell connection has a range of adaptations which support high-fidelity

transmission of high frequency signals across a wide bandwidth [9]. Faithful transmission of

rate information is reported [1, 2]. Vesicle release and replenishment are fast [10, 13] and

postsynaptic AMPA receptors operate in their linear range [13], where they are resistant to

desensitisation [88]. Multiple contacts are made by a mossy fibre with each granule cell [35].

Conversion of depolarising somatic charge into granule cell firing is a simple function, such that

granule cells 'have a relatively linear and uncomplicated conversion of depolarisation level to

spike rate' [41 p.2393, citing Jorntell and Ekerot 2006 and D'Angelo et al 1998].

Glutamate spillover enhances precision and reliability of transmission [13, 89]. During

behaviour, at sustained physiological mossy fibre rates, there may be a build-up of

intraglomerular glutamate, increasing the relative influence of spillover in a balance with

synaptic transmission, assisted by short-term synaptic depression. In these conditions,

spillover may dominate, as it dominates inhibitory transmission [5, 85]. Spillover from multiple

release sites (200-400 at a single mossy fibre terminal) is plausibly sufficient to mitigate

variability of vesicular size, release probability, and the number of release points at synapses

made on any individual cell, increasing fidelity and equality of excitation of granule cells.

Against that, there are reported to be heterogeneous synaptic weights. Different strength and

short-term dynamics of the mossy fibre to granule cell connection have been reported in

vestibular slices, with $GABA_A$ receptors blocked. The amplitude of the response, measured at

the soma, is reported to be statistically correlated to the source of the mossy fibre that is

stimulated [90], suggesting that the strength of the connection depends on the source.

However, the presence of weights does not necessarily mean that the function is to modulate

firing rates. Rather, it is suggested that the function may temporally code the source of inputs

because different combinations have different response onset times. An alternative, which

they also mention, is that some inputs may be in a supporting role, reminiscent of 'driver' and

'modulatory' signals in the thalamus and cortex [91]. Feasibly, signals in a supporting role must

be present for the postsynaptic granule cell to fire, but do not affect the granule cell firing rate.

Grants

Funding: This work was supported by the Leverhulme Trust [grant number ECF-2022-079] (Mike Gilbert); the Swedish Research Council [grant number 2020-01468]; the Per-Eric och Ulla Schybergs Foundation [grant number 42630]; and the Crafoord Foundation [grant numbers 20180704, 20200729, 20220776, & 20230655] (Anders Rasmussen).

Acknowledgement

Dr Mike Evans, School of Mathematics, University of Leeds, LS2 9JT, UK, read an early draft of the paper and provided feedback.

Author contributions

M.G. Researched, conceived and developed the ideas, performed the analysis, wrote the code, prepared the figures and wrote the manuscript. A.R. provided expertise and feedback, and reviewed and provided editorial comments on the manuscript.

Declaration of interests

The authors declare no competing interests.

REFERENCES

1.   Rancz, E.A., et al., *High-fidelity transmission of sensory information by single cerebellar*
mossy fibre boutons. Nature, 2007. **450**(7173): p. 1245-8.
2.   Ritzau-Jost, A., et al., *Ultrafast action potentials mediate kilohertz signaling at a central*
synapse. Neuron, 2014. **84**(1): p. 152-163.
3.   Jelitai, M., et al., *Dendritic excitation-inhibition balance shapes cerebellar output*
during motor behaviour. Nat Commun, 2016. **7**: p. 13722.
4.   Duguid, I., et al., *Control of cerebellar granule cell output by sensory-evoked Golgi cell*
inhibition. Proc Natl Acad Sci U S A, 2015. **112**(42): p. 13099-104.
5.   Duguid, I., et al., *Tonic inhibition enhances fidelity of sensory information transmission*
in the cerebellar cortex. J Neurosci, 2012. **32**(32): p. 11132-43.
6.   Walter, J.T. and K. Khodakhah, *The linear computational algorithm of cerebellar*
Purkinje cells. J Neurosci, 2006. **26**(50): p. 12861-72.
7.   Walter, J.T. and K. Khodakhah, *The advantages of linear information processing for*
cerebellar computation. Proc Natl Acad Sci U S A, 2009. **106**(11): p. 4471-6.
8.   Park, S.M., E. Tara, and K. Khodakhah, *Efficient generation of reciprocal signals by*
inhibition. J Neurophysiol, 2012. **107**(9): p. 2453-62.
9.   Delvendahl, I. and S. Hallermann, *The Cerebellar Mossy Fiber Synapse as a Model for*
High-Frequency Transmission in the Mammalian CNS. Trends Neurosci, 2016. **39**(11): p.
722-737.
10.   Saviane, C. and R.A. Silver, *Fast vesicle reloading and a large pool sustain high*
bandwidth transmission at a central synapse. Nature, 2006. **439**(7079): p. 983-7.
11.   Cesana, E., et al. *Novel granule cell-Golgi cell excitatory input in the cerebellar granular*
layer. in *FENS Abstr.* 2010.
12.   Moore, J.W. and D.E.J. Blazis, *11 - Simulation of a Classically Conditioned Response: A*
Cerebellar Neural Network Implementation of the Sutton–Barto–Desmond Model, in
Neural Models of Plasticity, J.H. Byrne and W.O. Berry, Editors. 1989, Academic Press.
p. 187-207.
13.   Sargent, P.B., et al., *Rapid vesicular release, quantal variability, and spillover contribute*
to the precision and reliability of transmission at a glomerular synapse. J Neurosci,
2005. **25**(36): p. 8173-87.
14.   Turecek, J., S.L. Jackman, and W.G. Regehr, *Synaptic Specializations Support*
Frequency-Independent Purkinje Cell Output from the Cerebellar Cortex. Cell Reports,
2016. **17**(12): p. 12.
15.   Turecek, J., S.L. Jackman, and W.G. Regehr, *Synaptotagmin 7 confers frequency*
invariance onto specialized depressing synapses. Nature, 2017. **551**: p. 503.
16.   Raymond, J.L. and J.F. Medina, *Computational Principles of Supervised Learning in the*
Cerebellum. Annu Rev Neurosci, 2018. **41**: p. 233-253.
17.   Alvina, K., et al., *Questioning the role of rebound firing in the cerebellum.* Nat Neurosci,
2008. **11**(11): p. 1256-8.
18.   Schonewille, M., et al., *Purkinje cells in awake behaving animals operate at the upstate*
membrane potential. Nat Neurosci, 2006. **9**(4): p. 459-61; author reply 461.

19.   Wu, H.S., I. Sugihara, and Y. Shinoda, *Projection patterns of single mossy fibers*
originating from the lateral reticular nucleus in the rat cerebellar cortex and nuclei. J
Comp Neurol, 1999. **411**(1): p. 97-118.

20.   Sultan, F. and D. Heck, *Detection of sequences in the cerebellar cortex: numerical*
estimate of the possible number of tidal-wave inducing sequences represented. J
Physiol Paris, 2003. **97**(4-6): p. 591-600.

21.   Shinoda, Y. and I. Sugihara, *Axonal Trajectories of Single Climbing and Mossy Fiber*
Neurons in the Cerebellar Cortex and Nucleus, in *Handbook of the Cerebellum and*
Cerebellar Disorders, S.J.D. Manto M., Rossi F., Gruol D.L., Koibuchi N., Editor. 2013,
Springer, Dordrecht.

22.   Shinoda, Y., et al., *The entire trajectory of single climbing and mossy fibers in the*
cerebellar nuclei and cortex. Prog Brain Res, 2000. **124**: p. 173-86.

23.   Garwicz, M., C.F. Ekerot, and H. Jorntell, *Organizational principles of cerebellar*
neuronal circuitry. News in Physiological Sciences, 1998. **13**: p. 28-34.

24.   Oscarsson, O., *Functional units of the cerebellum - sagittal zones and microzones.*
Trends in Neurosciences, 1979. **2**: p. 143-145.

25.   Ozden, I., et al., *Reliable coding emerges from coactivation of climbing fibers in*
microbands of cerebellar Purkinje neurons. J Neurosci, 2009. **29**(34): p. 10463-73.

26.   Dean, P., et al., *The cerebellar microcircuit as an adaptive filter: experimental and*
computational evidence. Nat Rev Neurosci, 2010. **11**(1): p. 30-43.

27.   Eccles, J.C., M. Ito, and J. Szentágothai, *The cerebellum as a neuronal machine.* 1967,
Berlin, New York etc.: Springer-Verlag. 335 p.

28.   Kanichay, R.T. and R.A. Silver, *Synaptic and cellular properties of the feedforward*
inhibitory circuit within the input layer of the cerebellar cortex. J Neurosci, 2008.
28(36): p. 8955-67.

29.   Dieudonné, S., *Submillisecond kinetics and low efficacy of parallel fibre-Golgi cell*
synaptic currents in the rat cerebellum. J Physiol, 1998. **510 (Pt 3)**: p. 845-66.

30.   Rossi, D.J. and M. Hamann, *Spillover-mediated transmission at inhibitory synapses*
promoted by high affinity alpha6 subunit GABA(A) receptors and glomerular geometry.
Neuron, 1998. **20**(4): p. 783-95.

31.   Nusser, Z., W. Sieghart, and P. Somogyi, *Segregation of different GABAA receptors to*
synaptic and extrasynaptic membranes of cerebellar granule cells. J Neurosci, 1998.
18(5): p. 1693-703.

32.   Brickley, S.G., S.G. Cull-Candy, and M. Farrant, *Development of a tonic form of synaptic*
inhibition in rat cerebellar granule cells resulting from persistent activation of GABAA
receptors. J Physiol, 1996. **497 (Pt 3)**: p. 753-9.

33.   Gilbert, M. and R.C. Miall, *How and Why the Cerebellum Recodes Input Signals: An*
Alternative to Machine Learning. Neuroscientist, 2021a. **DOI:**
10.1177/1073858420986795.

34.   Gao, Z., et al., *Excitatory Cerebellar Nucleocortical Circuit Provides Internal*
Amplification during Associative Conditioning. Neuron, 2016. **89**(3): p. 645-57.

35.   Jakab, R.L. and J. Hámori, *Quantitative morphology and synaptology of cerebellar*
glomeruli in the rat. Anat Embryol (Berl), 1988. **179**(1): p. 81-8.

36. Mitchell, S.J. and R.A. Silver, *GABA spillover from single inhibitory axons suppresses*
low-frequency excitatory transmission at the cerebellar glomerulus. J Neurosci, 2000a.
20(23): p. 8651-8.
37. Mitchell, S.J. and R.A. Silver, *Glutamate spillover suppresses inhibition by activating*
presynaptic mGluRs. Nature, 2000b. **404**(6777): p. 498-502.
38. Billings, G., et al., *Network structure within the cerebellar input layer enables lossless*
sparse encoding. Neuron, 2014. **83**(4): p. 960-74.
39. Jörntell, H. and C.F. Ekerot, *Properties of somatosensory synaptic integration in*
cerebellar granule cells in vivo. J Neurosci, 2006. **26**(45): p. 11786-97.
40. Chadderton, P., T.W. Margrie, and M. Hausser, *Integration of quanta in cerebellar*
granule cells during sensory processing. Nature, 2004. **428**(6985): p. 856-60.
41. Bengtsson, F. and H. Jörntell, *Sensory transmission in cerebellar granule cells relies on*
similarly coded mossy fiber inputs. Proc Natl Acad Sci U S A, 2009. **106**(7): p. 2389-94.
42. Cesana, E., et al., *Granule cell ascending axon excitatory synapses onto Golgi cells*
implement a potent feedback circuit in the cerebellar granular layer. J Neurosci, 2013.
33(30): p. 12430-46.
43. Albus, J.S., *A Theory of Cerebellar Function.* Mathematical Biosciences, 1971. **10**(1-2):
p. 25-61.
44. Gilbert, M. and A. Rasmussen, *Gap Junctions May Have A Computational Function In*
The Cerebellum: A Hypothesis. The Cerebellum, 2024.
45. Harvey, R.J. and R.M. Napper, *Quantitative study of granule and Purkinje cells in the*
cerebellar cortex of the rat. J Comp Neurol, 1988. **274**(2): p. 151-7.
46. Harvey, R.J. and R.M. Napper, *Quantitative studies on the mammalian cerebellum.*
Prog Neurobiol, 1991. **36**(6): p. 437-63.
47. Napper, R.M. and R.J. Harvey, *Number of parallel fiber synapses on an individual*
Purkinje cell in the cerebellum of the rat. J Comp Neurol, 1988. **274**(2): p. 168-77.
48. Napper, R.M. and R.J. Harvey, *Quantitative study of the Purkinje cell dendritic spines in*
the rat cerebellum. J Comp Neurol, 1988. **274**(2): p. 158-67.
49. Isope, P. and B. Barbour, *Properties of unitary granule cell-->Purkinje cell synapses in*
adult rat cerebellar slices. J Neurosci, 2002. **22**(22): p. 9668-78.
50. Ekerot, C.F. and H. Jörntell, *Parallel fibre receptive fields of Purkinje cells and*
interneurons are climbing fibre-specific. Eur J Neurosci, 2001. **13**(7): p. 1303-10.
51. Dugué, G.P., et al., *Cerebellar re-encoding of self-generated head movements.* Elife,
2017. **6**.
52. Chen, S., G.J. Augustine, and P. Chadderton, *The cerebellum linearly encodes whisker*
position during voluntary movement. eLife, 2016. **5**: e10509-e10509.
53. Herzfeld, D.J., et al., *Encoding of action by the Purkinje cells of the cerebellum.* Nature,
2015. **526**(7573): p. 439-42.
54. Hong, S., et al., *Multiplexed coding by cerebellar Purkinje neurons.* Elife, 2016. **5**.
55. Apps, R. and M. Garwicz, *Anatomical and physiological foundations of cerebellar*
information processing. Nat Rev Neurosci, 2005. **6**(4): p. 297-311.
56. Apps, R. and R. Hawkes, *Cerebellar cortical organization: a one-map hypothesis.* Nat
Rev Neurosci, 2009. **10**(9): p. 670-81.

892 57. Ji, Z. and R. Hawkes, *Topography of purkinje cell compartments and mossy fiber*
893   *terminal fields in lobules ii and iii of the rat cerebellar cortex: Spinocerebellar and*
894   *cuneocerebellar projections.* Neuroscience, 1994. **61**(4): p. 935-954.
895 58. Larsell, O., *The Comparative Anatomy and Histology of the Cerebellum from Myxinoids*
896   *through Birds.* 1967: The University of Minnesota Press.
897 59. Larsell, O., *The Comparative Anatomy and Histology of the Cerebellum from*
898   *Monotremes through Apes.* 1970: The University of Minnesota Press, Minneapolis.
899 60. Mugnaini, E., *The length of cerebellar parallel fibers in chicken and rhesus monkey.* J
900   Comp Neurol, 1983. **220**(1): p. 7-15.
901 61. Brand, S., A.L. Dahl, and E. Mugnaini, *The length of parallel fibers in the cat cerebellar*
902   *cortex. An experimental light and electron microscopic study.* Exp Brain Res, 1976.
903   **26**(1): p. 39-58.
904 62. Hámori, J., *Anatomy and neurochemical anatomy of the cerebellum*, in *Cerebellar*
905   *degenerations : clinical neurobiology*, A. Plaitakis, Editor. 1992, Kluwer Academic.
906 63. Palay, S.L. and V. Chan-Palay, *Cerebellar cortex : cytology and organization.* 1974,
907   Berlin: Springer.
908 64. Lange, W., *Regional differences in the distribution of golgi cells in the cerebellar cortex*
909   *of man and some other mammals.* Cell Tissue Res, 1974. **153**(2): p. 219-26.
910 65. Barmack, N.H. and V. Yakhnitsa, *Functions of interneurons in mouse cerebellum.* J
911   Neurosci, 2008. **28**(5): p. 1140-52.
912 66. Hámori, J. and J. Szentágothai, *Participation of Golgi neuron processes in the cerebellar*
913   *glomeruli: an electron microscope study.* Exp Brain Res, 1966. **2**(1): p. 35-48.
914 67. D'Angelo, E., et al., *The cerebellar Golgi cell and spatiotemporal organization of*
915   *granular layer activity.* Front Neural Circuits, 2013. **7**: p. 93.
916 68. Miles, F.A., et al., *Long-term adaptive changes in primate vestibuloocular reflex. III.*
917   *Electrophysiological observations in flocculus of normal monkeys.* J Neurophysiol, 1980.
918   **43**(5): p. 1437-76.
919 69. Edgley, S.A. and M. Lidierth, *The discharges of cerebellar Golgi cells during locomotion*
920   *in the cat.* J Physiol, 1987. **392**: p. 315-32.
921 70. Galliano, E., P. Mazzarello, and E. D'Angelo, *Discovery and rediscoveries of Golgi cells.* J
922   Physiol, 2010. **588**(Pt 19): p. 3639-55.
923 71. Eccles, J.C., R. Llinás, and K. Sasaki, *Golgi cell inhibition in the cerebellar cortex.* Nature,
924   1964. **204**: p. 1265-6.
925 72. Holtzman, T., et al., *Different responses of rat cerebellar Purkinje cells and Golgi cells*
926   *evoked by widespread convergent sensory inputs.* J Physiol, 2006. **574**(Pt 2): p. 491-
927   507.
928 73. Vos, B.P., A. Volny-Luraghi, and E. De Schutter, *Cerebellar Golgi cells in the rat:*
929   *receptive fields and timing of responses to facial stimulation.* Eur J Neurosci, 1999.
930   **11**(8): p. 2621-34.
931 74. Hartmann, M.J. and J.M. Bower, *Tactile responses in the granule cell layer of cerebellar*
932   *folium crus IIa of freely behaving rats.* J Neurosci, 2001. **21**(10): p. 3549-63.
933 75. Arenz, A., et al., *The contribution of single synapses to sensory representation in vivo.*
934   Science, 2008. **321**(5891): p. 977-80.
935 76. van Kan, P.L., A.R. Gibson, and J.C. Houk, *Movement-related inputs to intermediate*
936   *cerebellum of the monkey.* J Neurophysiol, 1993. **69**(1): p. 74-94.

77.   Giovannucci, A., et al., *Cerebellar granule cells acquire a widespread predictive*
feedback signal during motor learning. Nat Neurosci, 2017.
78.   Pietrajtis, K. and S. Dieudonné, *Golgi Neurons*, in *Handbook of the Cerebellum and*
Cerebellar Disorders, M. Manto, et al., Editors. 2013, Springer Netherlands: Dordrecht.
p. 829-852.
79.   Eyre, M.D. and Z. Nusser, *Only a Minority of the Inhibitory Inputs to Cerebellar Golgi*
Cells Originates from Local GABAergic Cells. eNeuro, 2016. **3**(2).
80.   Dumoulin, A., A. Triller, and S. Dieudonné, *IPSC kinetics at identified GABAergic and*
mixed GABAergic and glycinergic synapses onto cerebellar Golgi cells. J Neurosci, 2001.
21(16): p. 6045-57.
81.   Hull, C. and W.G. Regehr, *Identification of an inhibitory circuit that regulates cerebellar*
Golgi cell activity. Neuron, 2012. **73**(1): p. 149-58.
82.   Vervaeke, K., et al., *Rapid desynchronization of an electrically coupled interneuron*
network with sparse excitatory synaptic input. Neuron, 2010. **67**(3): p. 435-51.
83.   Ruigrok, T.J., R.A. Hensbroek, and J.I. Simpson, *Spontaneous activity signatures of*
morphologically identified interneurons in the vestibulocerebellum. J Neurosci, 2011.
31(2): p. 712-24.
84.   Rossi, D.J., M. Hamann, and D. Attwell, *Multiple modes of GABAergic inhibition of rat*
cerebellar granule cells. J Physiol, 2003. **548**(Pt 1): p. 97-110.
85.   Mapelli, L., S. Solinas, and E. D'Angelo, *Integration and regulation of glomerular*
inhibition in the cerebellar granular layer circuit. Front Cell Neurosci, 2014. **8**: p. 55.
86.   Ros, H., et al., *Neocortical networks entrain neuronal circuits in cerebellar cortex.* J
Neurosci, 2009. **29**(33): p. 10309-20.
87.   Rasmussen, A., et al., *Golgi cell activity during eyeblink conditioning in decerebrate*
ferrets. Cerebellum, 2014. **13**(1): p. 42-5.
88.   DiGregorio, D.A., et al., *Desensitization properties of AMPA receptors at the cerebellar*
mossy fiber granule cell synapse. J Neurosci, 2007. **27**(31): p. 8344-57.
89.   DiGregorio, D.A., Z. Nusser, and R.A. Silver, *Spillover of glutamate onto synaptic AMPA*
receptors enhances fast transmission at a cerebellar synapse. Neuron, 2002. **35**(3): p.
521-33.
90.   Chabrol, F.P., et al., *Synaptic diversity enables temporal coding of coincident*
multisensory inputs in single neurons. Nat Neurosci, 2015. **18**(5): p. 718-27.
91.   Sherman, S.M. and R.W. Guillery, *Distinct functions for direct and transthalamic*
corticocortical connections. J Neurophysiol, 2011. **106**(3): p. 1068-77.

Fig 1. Mossy fibre termination. A Sagittal section of the rat cerebellum (obtained by

multi-photon microscopy, courtesy Tom Deerinck). The outer layer of the cerebellar

cortex, the molecular layer, is stained green and the inner layer, the granular layer, is

blue. **B** Output of a procedure used to group mossy fibre terminals into clusters [20 fig

6], reproduced with permission. All terminals belong to one mossy fibre, which

terminates in two strips. Clusters are shown in different shades of grey. **C** Schematic of

an area of the folded cerebellar cortex. The strip inside the red dotted line is the width

and depth of a microzone but only part of its length. Termination of mossy fibre signals

is not at a point but at several randomly spaced, sagittally-aligned locations. Yellow

zones are terminal locations of a single mossy fibre. Size of locations varies depending

partly on the number of terminals in each cluster. Pink zones are terminal locations of

a second mossy fibre. In reality, zones overlap extensively and terminals are

intermingled. A mossy fibre mediolateral axon may give rise to several sagittal

branches, of which here only one is shown. The zone enclosed by a blue line has the

dimensions of a 'field'. A field is a nominal division of the granular layer with the

average dimensions of a mossy fibre terminal cluster – 200 µm x 150 µm – which we

use to quantify granular layer computations. A strip of the granular layer with the

dimensions of a sagittal row of 75–100 fields is a 'rank'. Viewed from the cerebellar

surface, a rank has the dimensions of a mid-range estimate of a microzone, but its long

sides are less cleanly defined. **D** Section of a microzone viewed obliquely from the

cerebellar surface, to illustrate the size and random intermingling of mossy fibre

terminals. The molecular layer is represented by squares of about the dimensions of a

Purkinje cell territory. The wire frame boxes are an exploded view of the orange fields.

The right box shows randomly distributed terminals of 5 mossy fibres (each a different

colour). We estimate that ~180 mossy fibres contribute terminals to a field, and a field

contains a total of around 700 terminals. Proportions of terminal size to field size are

preserved. The left box shows terminals of 20 mossy fibres.

Fig 2. Golgi cell ensemble conversion. Column 1 We randomly generated (3,645)

values representing mossy fibre rates with a uniform, normal, twin-peaked or

discontinuous distribution, simulating input to a row of 100 fields. Rate data were

randomly sampled to derive input rates to each field, and copies added to reflect the

fact that each afferent cell contributes a randomly variable number of terminals. Red

data represent copies-included signals received by a row of three fields. See Methods

for derivation of rates. **Column 2** The first-column red data were randomly sampled

150 times, sample size 4, representing mossy fibre input to a 3-field population of 30

Golgi cells, each with 5 basal dendrites (the mean) which each receive contact from 4

mossy fibres. Data are the sample means, representing dendritic depolarisation.

Column 3 We next randomly sampled column 2 data *without* replacement 30 times,

sample size 5, and took the sample means, representing somatic integration of

dendritic signals. **Column 4** Column 3 data were randomly sampled 700 times, sample

size 8–12, reflecting convergent input from 3 fields of Golgi cells onto 700 glomeruli in

the middle field (which receives the output of a three-field ensemble), and took the

sample means, representing GABA spillover that is a linear function of the mean of

afferent rates. Inhibition of granule cells is almost exclusively mediated by spillover.

Column 5 The column 4 data are presented as a histogram. **All rows** Column 1 mossy

fibre data are converted to narrowly-focussed inhibition of granule cells. The mean of

the column 4/5 data is linearly related to the mean of the mossy fibre distribution. This

result is independent of the shape of the original distribution.

1021

1022

1023

Fig 3. The Golgi cell population of a rank functions as a single unit. A Simulation of

Golgi cells in a sagittal row of 100 fields, taking the mossy fibre distribution on the left

as input (see Fig 2 and Methods for derivation of mossy fibre rates). **Top right**: pink:

the mean of mossy fibre input rates to each field; red: the mean with duplicate signals

added; dashed grey line: the mean of input rates to the whole rank. **Bottom right**: red:

same as top right; blue: mean, for each field, of glomerulus data derived in the same

way as Figure 2 columns 4 and 5 (the 'inhibitory field mean'); pale blue: SD of

glomerulus data. **B** Same as (A) but sampling a different mossy fibre distribution.

A and B

Red data closely track the pink data in top right panels, indicating that duplicate signals

(because the mossy fibre axon ends in a cluster of terminals) do not skew the mean of

firing rates received by a field.

Blue data across all fields are in a narrowly circumscribed range. Focus varies with the

bandwidth of the sampled distribution (mossy fibre rates). A normal mossy fibre

distribution with a range of over 100 Hz is converted to a blue range of ~3 Hz.

Inhibitory field means are tightly grouped: inhibition of granule cells is synchronised in

a rank.

The range of the blue data is centred on the grey dashed line – the mean of mossy

fibre input rates received by the whole rank – and varies independently of the red

data. That is, unpredictability inherent in the mechanism (because the outcome of

random sampling is inherently uncertain) is not compounded; random variation of the

red data doesn't flow through to amplify random variation of the blue data (in fact the

blue data have a slightly smaller range).

C The same simulation. Left: the sampled distribution (mossy fibre input rates to a

rank); right: the inhibitory field means (we use the means to compare fields). The y axis

of the right panel has the same range as the x axis of the left panel. Panels are tall to

improve visual comparison to other rows, where the y axis has a smaller range. **D** and **E**

The same, moving the sampled distribution to the right. Inhibition of granule cells is

precisely and linearly coupled to the mean of mossy fibre input rates to the whole

rank.

1054

1055

1056

1057

1058

Fig 4. Regulation of granule cell activity by the feed-forward pathway (mossy

fibres→Golgi cells→granule cells) We incorporated the Golgi cell ensemble

conversion into a simulation of the conversion of mossy fibre signals into granule cell

signals (recoding). One of the outputs of recoding is regulation of the percentage of

active granule cells. We tested if the recoding computation causes the percentage to

be stable, i.e., in the same narrow range in all fields regardless of the distribution of

mossy fibre rates. **Column 1** Mossy fibre firing rates were randomly generated with a

uniform, discontinuous, three-peak or normal distribution, to represent input to a row

of 100 fields. Rates, and rates received by each field, were derived in the same way as

earlier Figures. Red data are rates received by a single randomly-selected field,

including duplicate signals. y axes are scaled to the data – all distributions contain the

same number of signals. **Column 2** The percentage of granule cells that fire in each

field. The Golgi cell computation includes no adjustment to sparsen granule cell

activity. The white line and shaded area are mean and SD. **Column 3** The same, with an

adjustment (a co-efficient) to sparsen granule cell activity. y axes have 1/10 the range

of column 2. The percentage is confined to a narrow range. This result is independent

of the sampled (i.e., mossy fibre) distribution, suggesting that regulation is

independent of fluctuation of mossy fibre rates. Nonetheless, field-to-field variation

relative to the mean is significant. Homeostatic local fine tuning may be provided

physiologically by feedback via granule cell ascending axon contact on Golgi cells, not

represented in the simulation.

1080

Fig 5. A rank is a single recoding unit. The same simulation as Figure 4 but data are granule cell firing rates or derived from granule cell firing rates. **Column 1** We tested the effect of changing the shape of the mossy fibre distribution (without changing the number of signals). **Column 2** Pale green: sample means – the mean of input rates to each granule cell in a single field selected at random; blue: sample means of the subset of granule cells which meet the conditions to fire. In rows A–C, ~30% fire. In rows D–F, ~1% fire, y axes are rescaled and green data are omitted. **Column 3** Red: mean of mossy fibre input rates to each field in a rank, including duplicate signals; pink: SD of mossy fibre input rates to each field; blue: field means (i.e., the mean of granule cell firing rates in each field); pale blue: SD of granule cell firing rates in each field.

Observations

Regardless of the shape and large range of the mossy fibre distribution, the field means always converge towards a straight line which is a fixed distance from the overall mean of mossy fibre rates.

The amount and direction of field-to-field random fluctuation of the blue data is independent of local fluctuation of the red data, and in a smaller range. Uncertainty inherent in random sampling – first by single fields of input rates to a rank, and then by Golgi cells and granule cells of input rates to a field – is not compounded.

Fig 6. Granule cells collectively code mossy fibre rate information. Column 1 Mossy

fibre rates were generated in the same way as earlier Figures, with either a normal (SD

12.5) or uniform distribution representing input to a row of 100 fields, with copies

added – the sampled distribution. **Column 2** Histogram of granule cell firing rates in a

randomly selected field, obtained in the same way as in the bottom 3 rows of Fig 5.

Column 3 Grey line: the mean of the sampled distribution; blue line: field means. **A–C**

The fields means are precisely and linearly related to the mean of mossy fibre rates.

Blue data approach a straight line which is parallel to the grey line at a mean distance

of 15 Hz (SD 1.21 Hz). **D–F** If the shape and mean of a uniform sampled distribution are

unchanging, the distance between the blue and grey lines is linearly related to the

range.

1113