1

THE CEREBELLUM CONVERTS INPUT DATA INTO HYPER LOW-RESOLUTION 3-D PIXELS. WHO KNEW?

Mike Gilbert[a] (corresponding author), Anders Rasmussen[b]

[a]School of Psychology, College of Life and Environmental Sciences, University of Birmingham B15 2TT, UK

ORCID: 0000-0003-4415-6915

[b]Department of Experimental Medical Science, Lund University, BMC F10, 22184 Lund, Sweden

ORCID: 0000-0003-4477-369X

Correspondence: m.gilbert.1@bham.ac.uk

ABSTRACT

Our scope is the inner layer of the cerebellar cortex, the granular layer, where input to the cerebellum is received. We ask how input signals are converted into an internal code, what form that has, and how it solves some fundamental problems that come with using neurons to represent information. While there is a computational element, and the ideas are quantified with a computer simulation, the approach is primarily evidence-led and aimed at experimenters rather than the computational community. Network models are often simplified to provide a noiseless medium for sophisticated computations. We propose, with evidence, the reverse: physiology is highly adapted to provide a noiseless medium for straightforward computations, which are understandable with general science numeracy. We find that input data are converted to a hyper low-resolution, pixel-like internal code which spans the cerebellar network side-to-side. A pixel is the smallest unit of information but not the smallest unit of code – the same information is coded in any random sample of signals. Code in this form mitigates inconsistent performance by single neurons and synapses, explains how the cerebellum can work 'blind' (below a topographically-set scale threshold), and converts untidy input rates into coordinated firing of large cell groups – microzones – downstream.

Keywords: cerebellum; theory; model; granular layer; network; code

1. Introduction

It is generally assumed that brain-coded information is rich and replicable with computer-like high fidelity, and that those features make it precise and allow computations to be powerful. At least, that is not contentious. In many neural network learning models, including cerebellar models, computations are implemented by learned synaptic changes that selectively adjust synaptic transmission strength. This has been a mainstay of much modelling for 50 years. We propose that in the cerebellum, plausibly, all of that may not be true.

The inner layer of the cerebellar cortex, the granular layer, receives the main excitatory input to the cerebellum, from mossy fibres. Mossy fibres contact both granule cells and large inhibitory interneurons, Golgi cells. Golgi cells in turn inhibit granule cells. The granule cell axon rises into the outer layer, the molecular layer, where it divides in two to form parallel fibres, which lie parallel to the cerebellar surface and other parallel fibres. Parallel fibres make contact in passing on Purkinje cells, which carry the sole output of the cerebellar cortex.

We attempt to explain anatomy in the light of evidence of linear signalling. By anatomy we mean cell morphologies and connectivity, and bespoke adaptations of synaptic

contact. Inevitably, this involves attention to anatomical detail. The principles,

however, address generic problems that come with using nerve cells to code

information.

Speaking generally, neuron-to-neuron transmission involves the complex interaction of

a large array of often co-dependent biophysical and other variables which are specific

to the cell types making and receiving contact. With all the noise, it is difficult to

imagine how information is represented or moved around without catastrophic

corruption of data. Yet despite the seeming clamour, linear relationships are in fact

conserved. There is now a substantial body of evidence that rate-coded information is

linearly conserved in communication between cerebellar neurons [1-8], and that cells

are adapted for faithful transmission [9-15]. Firing rates that linearly code behavioural

and task metrics have been observed 'at all levels of the cerebellar circuit' [16 p.239,

citing 30 sources]. There are a number of examples that the cerebellum is adapted to

disengage or otherwise compensate for intrinsic non-linear properties of neurons and

synapses [13, 14, 17, 18]. This is – we submit – part of a general strategy of isolating

the effect of parameters that code information while eliminating or mitigating

interference by other variables. Evidence of linear signalling in the granular layer is

particularly rich. We include a review in the Methods section to which we cross refer

from the main text.

We will argue that the granular layer is in effect the physiological form of a network of

linear units which randomly sample firing rates in the afferent cell layer, within

topographically-defined boundaries. As sampling units have identified anatomical

counterparts, anatomy provides model parameters. We field evidence in the main text

to explain the derivation of model parameters. Since cerebellar cell morphologies and

connections are well studied, parameter values (usually a range) are available from the

literature. We state in the next section the proposal that biology simulates a network

of linear units. We expand on that in statements at or near the beginning of each

section, headed 'Hypothesis', then describe (in each section) how it was tested, and

the results.

Classically, neural network models solve a problem. The nature of the problem – and

therefore the function of the network – is a proposal and becomes an assumption of

the model. The solution is a computational mechanism; the mechanism is the model.

The direction is function → mechanism (the first two steps of Marr's three levels of

analysis; the third step is to consult the evidence to see if the known physiology could

provide the physical hardware to implement the mechanism). We work in reverse. We

propose first a mechanism that can explain the evidence, then infer function from the

mechanism. High-level ideas come last. We use modelling in a different role, to

quantify and test the ideas by simulating the mechanism, to verify that it works with

physiological parameters. The direction is evidence → mechanism → function. This is

simply a different approach and not in competition with other methods. However, it is

important (and we raise it here) because while there is a computational element, and

the ideas are quantified with a computer simulation, it is not a computational paper.

Simply, as firing rates represent values, we describe – what we think are – the code

and computations, but as they are straightforward, they can be described in words and

require only general science numeracy.

To aid navigation, our key findings are: **1.** Anatomy (cell morphologies and wiring)

performs a passive computation unaided and unlearned. **2.** There is no algorithm. **3.**

Physiology is the physical form of a network of linear units (meaning, output of a unit is

a linear function of inputs). A unit can be a cell, part of a cell or some other structure,

but not a single synapse. **4.** External signals are converted into a low-resolution

internal code that occupies volume – 3-D pixels. Accordingly, internally-coded

information has spatial dimensions, the dimensions of cell groups that code together

(code groups). **5.** Code groups span the cerebellar network from side to side (sagittal

direction, mirroring division of the outer layer of the cerebellar cortex into

microzones). **6.** Information is coded in the combined activity of code-grouped cells,

and code groups communicate by a facsimile of independent random sampling, for

which physiology is highly adapted. **7.** The code is analogue, under modulation of

continuous input during behaviour. **8.** Coding and data processing exploit statistical

effects of random sampling. **9.** A pixel (the information represented by a code group) is

the smallest unit of information. Inside pixel dimensions, information is indivisible –

i.e., any smaller number of randomly selected cells (over a minimum number) still

represents the same information. **10.** By the same token, a pixel is not the smallest

unit of code – the same information is coded in any random sample of firing rates in a

code group. **11.** The form of the code is key to function. It has these results. (i) Neural

representation of information does not depend on replicability of neural activity (a

solution to the problem that input data are 'unreliable' in the sense that firing of single

neurons may not code information consistently or precisely, and synaptic transmission

involves random variables). (ii) The cerebellum can work 'blind' below a

topographically-set scale threshold (because it does not need to differentiate between

signals below pixel scale). (iii) It converts input rates – with any frequency distribution

– into coordinated firing of large cell groups – microzones – downstream (because

target cells can receive the same code without receiving the same signals).

127 2. **Linear units**

The cerebellum is the physical implementation of a computational network of linear

units. Linear refers to the conversion function – output is a linear function of inputs,

such as the weighted sum or mean of inputs. A unit can be a cell, part of a cell or some

other structure, but never a single synapse. Units form layers that data are passed

through. This happens continuously. Data are coded, at any instant, in a snapshot of

the collective output of a layer. Where – as in most cases – units receive as inputs

values represented by firing rates in the afferent layer, they randomly sample afferent

rates. Anatomy mimics random sampling of the whole cell layer afferent to a group

(we will explain how). The result is a facsimile of random sampling by individual units

of the entire afferent population of firing rates, with replacement, notwithstanding

that a single unit receives unchanging contact from a tiny fraction of cells in the

afferent layer. Cell morphologies and network geometry are explicable as bespoke

adaptations for this purpose.

Random sampling, we will argue, is inside spatial dimensions that are functionally

defined by anatomy, although they are not physically bounded. The result is that firing

rates of successive layers are linearly related by rate statistics at collective level, at that

scale. Signalling is analogue, so that signalling and computations are continuous during

behaviour.

147 3. **Mossy fibres**

148 **3.1 Anatomy, 'ranks' and 'files'**

The molecular layer of the cerebellar cortex is divided functionally into long, thin strips

called microzones [19]. Mossy fibres give rise to sagittally-extending collaterals, the

direction of the long axis of microzones (Fig 1). These in turn give rise to multiple distal

branches, each ending in a cluster of terminals [20-23]. Terminal clusters are sagittally

aligned and separated by a randomly variable distance typically not less than 200 µm

[20, 21] measured from edge to edge of clusters (so they are separated by a greater

distance measured from centre to centre). Terminals themselves are intimately and

randomly intermingled with terminals of other mossy fibres.

Accordingly, mossy fibres do not terminate at a point but a region with dimensions:

the field occupied by a terminal cluster. A single mossy fibre gives rise to several

terminal branches and each of those to several terminals, such that a sagittal strip of

the granular layer receives multiple copies of every signal at each of multiple locations.

Cluster size is variable (because it depends on the number of terminals and how tightly

clustered they are), averaging 200 µm sagittally x 150 µm mediolaterally [21]. We term

the average dimensions of a cluster a 'field'. The granular layer is anatomically

seamless. A field is simply a nominal division that we use for modelling. A sagittal row

of 75–100 fields laid end to end has the area (viewed from the cerebellar surface) of a

mid-range microzone [19, 24-26]. We term a row of fields a 'rank'. Ranks are

functionally but not anatomically discrete. Ranks are parallel to microzones but they

are divisions of the granular layer, while microzones are divisions of the molecular

layer.

Each terminal is ensheathed by a semi-permeable membrane which restricts

neurotransmitter diffusion, a structure taken together termed a glomerulus. Excitation

of granule cells and Golgi cells by mossy fibres, and inhibition of granule cells by Golgi

cells, all take place here.

Fig 1. Mossy fibre termination. A Sagittal section of the rat cerebellum (obtained by

multi-photon microscopy, courtesy Tom Deerinck). The outer layer of the cerebellar

cortex, the molecular layer, is stained green and the inner layer, the granular layer, is

blue. **B** Output of a procedure used to group mossy fibre terminals into clusters [21 fig

6], reproduced with permission. All terminals belong to one mossy fibre, which

terminates in two strips. Clusters are shown in different shades of grey. **C** Schematic of

an area of the folded cerebellar cortex. The strip inside the red dotted line is the width

and depth of a microzone but only part of its length. Yellow zones are terminal

locations of a single mossy fibre. Pink zones are terminal locations of a second mossy

fibre. In reality, zones overlap extensively and terminals are intermingled. The zone

enclosed by a blue line has the dimensions of a 'field', a nominal division of a rank with

the average dimensions of a mossy fibre terminal cluster. **D** Section of a microzone

viewed obliquely from the cerebellar surface, to illustrate the size and random

intermingling of mossy fibre terminals. The wire frame boxes are an exploded view of

the orange fields. The right box shows randomly distributed terminals of 5 mossy fibres

(each a different colour). Proportions of terminal size to field size are preserved. The

left box shows terminals of 20 mossy fibres (out of ~180 that contribute terminals to a

field).

195 3.2 Hypothesis

Mossy fibre anatomy is an anatomical solution to a computational problem. The

cerebellum exploits statistical effects of random sampling called the central limit

theorem. For the effects to occur, samples must be independent, also termed sampling

'with replacement' (in the illustration sometimes used, coloured balls selected at

random from a bag are put back in the bag between samplings). Clearly, physical

replacement is impossible because neither a terminal cluster nor a terminal can be in

more than one place at the same time. However, anatomy simulates replacement.

Terminal branching simulates sampling with replacement by fields in a rank of input

rates to the whole rank – the distributed probabilities of the number of terminal

branches per mossy fibre are the same as the probabilities of the number of repeat

samplings if rates received by a rank were in fact independently randomly sampled by

fields. Terminal clustering simulates sampling with replacement by cells in a field of

input rates to the field – the distributed probabilities of the number of terminals per

cluster are the same as the distributed probabilities of the number of repeat samplings

if rates received by a field were in fact randomly sampled with replacement by

individual target neurons. The combined effect is a biological facsimile of sampling

with replacement by individual neurons of firing rates received as input to a whole

rank, notwithstanding that they receive fixed contact from a tiny fraction of afferent

cells. Note that it is the population of *rates* that are hypothetically sampled with

replacement, and not *signals.* The population of signals includes copies of rates

because of branching and clustering.

3.3 Terminal branching

If each mossy fibre could terminate in only one field, a mossy fibre 'sampled' by one

field clearly cannot also be sampled by any others. Termination by a single mossy fibre

in a random number of terminal branches, at multiple locations, removes that

restriction.

In theory, sampling with replacement should mean any number of fields can receive

the same signal, up to all of them. Clearly this is not true – there is still an anatomical

limit. But in practice even with real replacement the number of repeat samplings is

constrained by a probability distribution. If, then, the number of terminal branches per

mossy fibre has the same probability distribution, branching has the same result as if

firing rates received by each field in a rank were in fact a random sample with

replacement of firing rates received by the rank as a whole.

We can test that by calculating the probability distribution in the hypothetical case

that mossy fibres did not branch terminally and rates were in fact sampled with

replacement. The probability, p, that a given rate is selected k times in taking N

samples, sample size x, assuming mossy fibres each give rise to an average of m

terminal branches, is given by

$$\left[Nx! \big/ k! \, (Nx - k)! \right] * \left[1 \big/ (Nx/m) \right]^{k} * \left[1 - \left(1 \big/ (Nx/m) \right) \right]^{Nx-k}$$

236   The results are shown in Table 1.

237   **Table 1. Probability p of k repeat selections of a single mossy fibre**

k	0	1	2	3	4	5	6	7	8	9	10	11
p	0.007	0.034	0.084	0.140	0.175	0.175	0.146	0.104	0.065	0.036	0.018	0.008

239   Although data are scarce, the number of repeat selections is consistent with what we

240   know about the biological range of the number of terminal clusters per mossy fibre

241   [21, 27]. Estimated by this method, the mean is 4–5.

3.4 Terminal clusters

The principle is the same here but at local, field scale. Mossy fibre terminal clustering,

and random intermingling with terminals of other mossy fibres (Fig 1), means that

contact on any particular cell does not mean the same mossy fibre cannot also contact

cells somewhere else. Again, to approximate independence, it is only necessary for the

number of repeat selections of a signal to have the same probability distribution as if

there was actual replacement.

A practical problem with the strategy of clustering is that a target could receive the

same signal twice (or more). The risk is mitigated by taking small samples. A sample

size of 4 is small enough for a low risk. Granule cells average 4 dendrites, each

receiving contact from a single terminal, and we estimate that Golgi cells receive an

average of 4 inputs to each of their basal dendrites (see derivation of parameters in

the next section). Given m x 100 terminals and an independent and equal chance that

contact is by any terminal, the probability of an n^{th} contact by the same mossy fibre on

a single dendrite is:

$$P_n = \left[\prod_{x=2}^{n} \left(\frac{m - x + 1}{100m - x + 1} \right) \right] \times \frac{y!}{x! \, (y - x)!}$$

where y is the number of inputs to a single dendrite and m is the number of terminals

per cluster, provided by (for illustration) 100 mossy fibres. Calculated in this way,

around one in every twenty samples contains a duplicate, and less than two in a

thousand contain more than one. The actual fractions are smaller because the number

of mossy fibres that innervate a field is higher, around 180 (see Methods, Derivation of

mossy fibre rates).

Quasi-sampling by each field of mossy fibre firing rates received as input to the rank as

a whole, and by cells in each field of rates received by that field, is a biological facsimile

of simultaneous random sampling with replacement by single neurons of input rates to

a whole rank.

4. Golgi cells

4.1 Hypothesis

Anatomical detail is critical to computational performance and the functional division of the granular layer into groups of cells that work together. Despite forming an anatomically seamless carpet of cells, Golgi cells work in groups: ensembles. An ensemble is the population of a sagittal row of three fields. Groups are not anatomically bounded or even functionally discrete, but function as a single unit for the anatomical reason that the axonal range of all Golgi cells in a 3-field row includes all or most of the middle field. Put another way, each field defines a functional population of Golgi cells – an ensemble – being those cells with convergent input to that field.

Ensembles perform a computational conversion of mossy fibre rates received by a three-field row into inhibition of granule cells. Say a population of values (with any frequency distribution) is randomly sampled. There are n samples, sample size m. If we take the mean (or sum) of each sample, and n is large, the sample means have a normal or near-normal distribution, by the central limit theorem. (This works even if m is a range.) The new distribution is centred on the same mean as the sampled distribution, but narrower. Modest values of n or m give an untidy result. However, if

the new distribution is then itself randomly sampled, and the process is repeated, the

results get tidier and narrower.

The ensemble computation is (chiefly) a statistical result of random sampling, in two

steps. At the first step, Golgi cell basal dendrites receive contact from a random

sample of mossy fibres that innervate their location (strictly speaking, they randomly

sample terminals); and in the second step, middle-field glomeruli each receive

innervation by a random sample of Golgi cells. The effect of the computation is to

synchronise inhibition of granule cells in the middle field. The strength of inhibition is

linearly related to the mean of mossy fibre firing rates received as input to the entire

rank. Ensembles conflate sagittally anatomically and functionally. Ensembles in a

sagittal strip all return the same output (because they all perform the same

computation on input data sampled from the same distribution). Effectively: the Golgi

cell population of a whole rank functions as a single unit.

4.2 The Golgi cell computation

We envisage that in motor circuits during behaviour signalling is analogue, avoiding the

noise caused by discrete signals. So, for example, Golgi cell dendritic signalling is a

sustained state under modulation by fluctuating mossy fibre rates. Similarly, GABA

spillover into the intraglomerular space, almost exclusively the agent of granule cell

inhibition, becomes elevated by a build-up during behaviour, so that this too is at a

sustained level, controlled by afferent Golgi cell rates. Thus, at each step, data are

coded at any moment in a snapshot of a state of change.

We tested the hypothesis *in silico* by simulating a rank. Derivation of model

parameters is described in: Methods, Derivation of ensemble dimensions and model

parameters. Mossy fibre rates sampled by Golgi cells in each field were obtained by

sampling from a distribution representing signals received as input to a rank, which

were randomly generated subject to constraints on the shape of the distribution. This

was modified to take into account mossy fibre terminal branching and clustering,

random variation of the number of terminals per cluster and clusters per cell, and the

fact that most clusters fall partly inside and partly outside a field (see Methods,

Derivation of mossy fibre rates).

This gave the number of mossy fibre signals, firing rates and the number of copies of

each rate received by each field, and therefore input to an ensemble territory,

populated by 30 Golgi cells, 10 per field. There are ~700 mossy fibre terminals per

field. As far as we know (and we propose), contact by mossy fibres on Golgi cells is at

random at field scale, i.e., signals received by a single basal dendrite are provided by a

random sample of local terminals. The number of physiological samplings is equal to

the number of basal dendrites per field; sample size is the number of mossy fibres

which make contact on each dendrite. The effect is sustained dendritic depolarisation

under modulation by a linear function of input rates (see Methods, Linear transmission

of mossy fibres to Golgi cells), which we take as the mean.

Taking the sample means generates a new set of values (Fig 2 column 2) with a smaller

range and centred on the mean, which is linearly related to the mean of the sampled

distribution. Basal dendritic charge transfer is passive and little filtered [11, 28].

Integration of dendritic states at the Golgi soma is represented by taking the mean of

dendritic charge, for each simulated cell. Golgi cell firing rates have a linear

relationship with the amplitude of depolarising current [29], suggesting (and we

assume) a linear conversion of somatic membrane potential into the Golgi cell firing

rate.

The output is a third data set, Golgi cell firing rates (Fig 2 column 3), on the same

normalised scale. Each glomerulus in the middle field receives innervation from a

random subset of Golgi cells afferent to the field, and accordingly a random sample of

Golgi cell firing rates. Golgi cells inhibit granule cells in the glomerulus. A granule cell

extends a single dendrite into a glomerulus, typically receiving synaptic contact from

two or three Golgi cells [30]. However, inhibition of granule cells by Golgi cells is

almost entirely by GABA spillover into the intraglomerular space [5], via extrasynaptic

receptors [31, 32] (see Methods, Linear transmission of Golgi cells to granule cells).

This effectively increases the Golgi cell to granule convergence ratio, since the number

of Golgi cells that are functionally afferent to a granule cell is not the number that

make direct contact, but the number afferent to a glomerulus. The membrane around

a glomerulus restricts GABA diffusion so that signalling is by modulation of the

intraglomerular concentration of GABA, which in turn reflects the rate it is released.

The rate of release by a single cell is proportional to the firing rate of that cell.

Intraglomerular GABA concentration is under continuous modulation at a sustained

level which we take as a linear function of the mean of afferent Golgi cell rates (the

sum also works). Sample size is provided by the Golgi cell to glomerulus convergence

ratio. The number of Golgi cells afferent to a glomerulus is unreported, as far as we

know. We take it as a random number in the range 8–12, generated for each

glomerulus. We tested other ratios to confirm that the ratio does not materially affect

the ensemble computation (Supplementary Materials). The number of samples is the

number of glomeruli per field (~700, the same as terminals)

This generates a final population of values (Fig 2 columns 4 and 5), the output of the

ensemble conversion. There are 700 outputs, representing glomeruli in the middle

field. All granule cells that extend a dendrite into the same glomerulus receive equal

inhibition (of that dendrite).

Fig 2. Golgi cell ensemble conversion. Column 1 We randomly generated (3,645) values representing mossy fibre rates with a uniform, normal, twin-peaked or discontinuous distribution, simulating input to a row of 100 fields. Rate data were randomly sampled to derive input rates to each field, and copies added to reflect the fact that each afferent cell contributes a randomly variable number of terminals. Red data represent copies-included signals received by a row of three fields. See Methods for derivation of rates. **Column 2** The first-column red data were randomly sampled 150 times, sample size 4, representing mossy fibre input to a 3-field population of 30 Golgi cells, each with 5 basal dendrites (the mean) which each receive contact from 4 mossy fibres. Data are the sample means, representing dendritic depolarisation. **Column 3** We next randomly sampled column 2 data *without* replacement 30 times, sample size 5, and took the sample means, representing somatic integration of dendritic signals. **Column 4** Column 3 data were randomly sampled 700 times, sample size 8–12, reflecting convergent input from 3 fields of Golgi cells onto 700 glomeruli in the middle field (which receives the output of a three-field ensemble), and took the sample means, representing GABA spillover that is a linear function of the mean of afferent rates. Inhibition of granule cells is almost exclusively mediated by spillover. **Column 5** The column 4 data are presented as a histogram. **All rows** Column 1 mossy fibre data are converted to narrowly-focussed inhibition of granule cells. The mean of the column 4/5 data is linearly related to the mean of the mossy fibre distribution. This result is independent of the shape of the original distribution.

4.3 Results

4.3.1 An ensemble

There is a progressive narrowing of the data range at each step. Inputs to an ensemble (mossy fibre signals received by a row of three fields) are converted to narrowly-grouped and normally-distributed outputs (representing GABA concentration in middle-field glomeruli). The narrow output range indicates that inhibition is locally synchronised, i.e., granule cells in the middle field receive co-modulated inhibition. Inhibition has a linear relationship with mossy fibre rates, acting through the ensemble conversion. These results are independent of the frequency distribution of mossy fibre rates – input to the system may be in any permutation of firing rates, with the same result.

4.3.2 A rank

We next used the ensemble conversion to derive inhibition of granule cells in a sagittal row of 100 fields, a rank. The results are shown in Figure 3. As ensembles all perform the same computation and effectively randomly sample (with replacement) the same data (the distribution of mossy fibre input rates to the whole rank), they all return the same output, or nearly the same output. There are 700 outputs per field, representing glomeruli, as before. The mean of outputs, taken for each field, has a narrow range centred on the mean of mossy fibre input rates to the parent rank. Focus varies with

the bandwidth of the sampled distribution (mossy fibre rates). A normal mossy fibre

distribution with a range of over 100 Hz is converted to a blue range of ~3 Hz.

In Figure 3 rows A and B, blue data in the bottom right panel represent the mean, for

each field, of glomerulus data. The range of the blue data is centred on the grey

dashed line – the mean of mossy fibre input rates received by the whole rank – and

varies independently of the red data. That is, field-to-field variation of output appears

to be inherent in the mechanism – i.e., caused by inherent uncertainty of the outcome

of random sampling. However, it is not compounded – random variation of the red

data doesn't flow through to amplify random variation of the blue data (in fact the

blue data have a slightly smaller range).

Conclusion: Golgi cells in a rank act as a single functional unit. Ensembles conflate in

the sagittal direction functionally as they do anatomically. As a result, the whole

granule cell population *of a rank* receive synchronised inhibition which is linearly

related to, and varies with, the mean of mossy fibre rates.

Fig 3. The Golgi cell population of a rank functions as a single unit. A Simulation of Golgi cells in a sagittal row of 100 fields, taking the mossy fibre distribution on the left as input (see Fig 2 and Methods for derivation of mossy fibre rates). **Top right**: pink: the mean of mossy fibre input rates to each field; red: the mean with duplicate signals added; dashed grey line: the mean of input rates to the whole rank. **Bottom right**: red: same as top right; blue: mean, for each field, of glomerulus data derived in the same way as Figure 2 columns 4 and 5 (the 'inhibitory field mean'); pale blue: SD of glomerulus data. **B** Same as (A) but sampling a different mossy fibre distribution. **A and B** Red data closely track the pink data in top right panels, indicating that duplicate signals (because the mossy fibre axon ends in a cluster of terminals) do not skew the mean of firing rates received by a field. Inhibitory field means are tightly grouped: inhibition of granule cells is synchronised in a rank. **C** The same simulation. Left: the sampled distribution (mossy fibre input rates to a rank); right: the inhibitory field means (we use the means to compare fields). The y axis of the right panel has the same range as the x axis of the left panel. Panels are tall to improve visual comparison to other rows. **D** and **E** The same, moving the sampled distribution to the right. Inhibition of granule cells is precisely and linearly coupled to the mean of mossy fibre input rates to the whole rank.

5. Granule cells

5.1 The fraction of active granule cells is reliably regulated by feed-forward inhibition at low density

5.1.1 Hypothesis

Anatomy simulates independent random sampling by a single granule cell of the population of mossy fibre input rates to the whole rank (section 3). There is a competition for influence between mossy fibre excitation and Golgi cell inhibition of granule cells. The competition takes place in the glomerulus. During behaviour, this takes place ceaselessly. The strength of inhibition is synchronised between glomeruli that populate a rank (section 4), so that all mossy fibre signals compete, at any moment, against the same inhibitory threshold. The outcome in each glomerulus is independent of the outcome at other terminals, including other terminals that innervate the same granule cell. To fire, a granule cell must receive a minimum number of mossy fibre inputs that prevail in the glomerular competition. The probability that enough prevail depends, at any moment, on the strength of inhibition. Since inhibition is synchronised, the probability is the same for all granule cells at any given instant in time. The number that fire is very near the expected proportion (the number predicted by probability) because granule cells are numerous. The probability

is also the same at all times because inhibition has a linear relationship with mossy

fibre firing rates (section 4). This is always conserved (by the Golgi cell ensemble

conversion) regardless of the mossy fibre distribution (section 4). So, the probability is

invariant, and therefore so is the proportion of active granule cells.

Note: the idea of an all-or-nothing result of the intraglomerular 'balance' of excitation

and inhibition has appeared previously [33].

5.1.2 Anatomy

Granule cells receive input to each dendrite from a single mossy fibre. Granule cells

have 3–5 dendrites, averaging 4 (the simulation assumes 4, but the result is the same

with 3–5). Each dendrite of a single granule cell extends into a different glomerulus.

Estimates vary of the number of granule cells that receive contact from a single mossy

fibre terminal. 50 is mid-range [34, 35], but estimates vary significantly [2]. 50 is

consistent with our estimates of the number of terminals and granule cells in a field.

The granule cell sampling unit is the whole cell since it receives only a single input rate

to each dendrite. Sample size is therefore 4, and the number of samples per field is the

number of granule cells, which we estimate at 8,750 (see Methods, Derivation of

mossy fibre rates).

Excitation and inhibition compete for influence in a glomerulus. It is unclear if there is

always an emphatic winner, but a swing in the dominant direction may be amplified by

positive feedback, acting through presynaptic receptors. Activation of presynaptic

GABA$_B$ receptors on mossy fibres inhibits glutamate release [36], while activation of

presynaptic mGluR2 receptors on Golgi cells inhibits GABA release [37]. We postulate

that only mossy fibre signals which prevail in the glomerular competition contribute

significantly to granule cell depolarisation.

5.1.3 Feed-forward inhibition regulates the probability that granule cells

fire

The probability that a given granule cell receives three competitive signals and

therefore fires is a function of the percentage of mossy fibre signals that are

competitive. The expected number of granule cells that fire in a field is given by

$$E_{fld} = T \sum_{x=j}^{n} \left[\frac{n!}{(n-x)!x!} [P(e>i)]^x [1-P(e>i)]^{n-x} \right]$$

where T is the total number of granule cells per field, n is the number of dendrites per

granule cell, and j is the minimum number of dendrites that must receive dominant

excitation, $e > i$. The proportion of granule cells that fire is near the expected number

by the law of large numbers. The percentage of mossy fibre signals that are

competitive is always the same because inhibition scales with the mean of mossy fibre

rates (because Golgi cells translate mossy fibre firing rates into proportional

inhibition). As the percentage is unvarying, so is the probability.

503 5.1.4 Does physiology have that outcome?

To test that, we incorporated the Golgi cell ensemble conversion into a model that

recodes mossy fibre signals into granule cell signals. Probability is an abstraction with

no direct physiological counterpart. The model does not include a binomial function

but derives output from input by comparing excitatory to inhibitory input to each

glomerulus, the way we claim it is derived physiologically. The equation in the previous

section is simply shorthand – biology exploits chance and large numbers to generate

the same result as if inputs were passed through the equation.

The model compares mossy fibre signal frequency with inhibition at each of 4

dendrites of each of 8,750 granule cells per field, in each of 100 fields. Mossy fibre

input rates to each field are derived per section 4 and Methods, so taking into account,

inter alia, that a field receives a randomly variable number of copies of most signals,

and most terminal clusters lie only partly inside a field. Inhibition – outputs of the

ensemble conversion, 700 per field – is provided by the Golgi cell model. Excitation and

inhibition received by each glomerulus are randomly paired. Pairing is used to identify

the subset of granule cells that fire. We assume that, to fire, a granule cell must

receive at least 3 mossy fibre signals which exceed the inhibitory threshold. In

experimental conditions, using stimulation that excites very high mossy fibre rates,

granule cells can be induced to fire in response to only one input [1]. However, other

accounts suggest one is too few [5, 38, 39], and three are needed [40, 41]. Output of

the model is the number of granule cells that fire in each field, the mean of mossy fibre

rates they each receive (the sample means), and the mean of the sample means in

each field ('field means'). The results are shown in Figure 4.

5.1.5 Results

Feed-forward inhibition – mossy fibre → Golgi cell → granule cell – reliably and closely

regulates the fraction of granule cells that are active at low density. Field-to-field

random variability of the number is inversely proportional to the fraction that are

active, suggesting that, if primary regulation is provided by feed-forward inhibition, the

fraction is low. We would nonetheless expect to see greater precision, suggesting the

physiological presence of a supplementary mechanism(s) that we have not

represented in the model.

A local feedback mechanism may be provided by granule cell ascending axon contact

on Golgi cells. The ascending portion of the granule cell axon makes synaptic contact

on local Golgi cells, on basal dendrites [11]. A Golgi cell receives ~400 functional

synaptic contacts from this source, around 25% of the granule cells that a Golgi cell

receives contact from in total. During sustained mossy fibre discharge, synapses from

local granule cells contribute equivalent charge to Golgi cells as mossy fibre synapses

and 'display similarly fast kinetics' [42]. Short-latency inhibitory feedback may add

homeostatic fine adjustment to feed-forward regulation at local level, we propose.

It is a long-standing idea [43] that granule cell activity may be regulated by feedback

via parallel fibres, which contact Golgi cell apical dendrites. We have previously

proposed that gap junctions between apical dendrites may provide the means for

Golgi cells to detect the density of parallel fibre activity [44]. Feedback by this pathway

is not sensitive to local fluctuation of granule cell active numbers, and may be the

weakest regulator.

The results are the same with any (tested) distribution of mossy fibre rates and also

whether mossy fibres are all active or some are active and others are not. Active cells

can be in any random permutation.

Fig 4. Regulation of granule cell activity by the feed-forward pathway (mossy fibres→Golgi cells→granule cells) We incorporated the Golgi cell ensemble conversion into a simulation of the conversion of mossy fibre signals into granule cell signals (recoding). One of the outputs of recoding is regulation of the percentage of active granule cells. We tested if the recoding computation causes the percentage to be stable, i.e., in the same narrow range in all fields regardless of mossy fibre rates. **Column 1** Mossy fibre firing rates were randomly generated with a uniform, discontinuous, three-peak or normal distribution, to represent input to a row of 100 fields. Rates, and rates received by each field, were derived in the same way as earlier Figures. Red data are rates received by a single randomly-selected field, including duplicate signals. y axes are scaled to the data – all distributions contain the same

number of signals. **Column 2** The percentage of granule cells that fire in each field. The
Golgi cell computation includes no adjustment to sparsen granule cell activity. The
white line and shaded area are mean and SD. **Column 3** The same, with an adjustment
(a co-efficient) to sparsen granule cell activity. y axes have 1/10 the range of column 2.
The percentage is confined to a narrow range. This result is independent of the
sampled (i.e., mossy fibre) distribution, suggesting that regulation is independent of
fluctuation of mossy fibre rates. Nonetheless, field-to-field variation relative to the
mean is significant. Homeostatic local fine tuning may be provided physiologically by
feedback via granule cell ascending axon contact on Golgi cells, not represented in the
simulation.

576 5.2 Granule cells encode mossy fibre rate information

577 collectively in rank-sized strips

578 5.2.1 Hypothesis

Anatomy simulates independent random sampling by a single granule cell of the

population of mossy fibre input rates to the whole rank (section 3). The granule cell

firing rate is a linear function of depolarising somatic charge, in turn a linear function

of the mean (or sum) of excitatory input rates that prevail against inhibition. We

postulate that the linear relationship is undiluted by inhibition because dominant

signals displace a significant effect of inhibition. Derived in this way, granule cell rates

are collectively well predicted by mossy fibre rates, by the central limit theorem. This

provides the substrate for granule cells to collectively encode mossy fibre rate

information. Granule cell code groups are functionally defined by the topography of

mossy fibre input to the cerebellum, as Golgi cell networks are. The output of a rank is

the smallest unit of granule cell encoded information. The permutation of active mossy

fibres, and of granule cells, and how firing rates are distributed among active cells, all

code nothing.

5.2.2 Do granule cell firing rates – the output of a rank – have the

hypothesised features?

We ran the same simulation as section 5.1 to test whether, and how, mossy fibre rates

are related to granule cell rates, and if fields in a rank all have the same output. We

take granule cell firing rates to be a linear function of the mean of dominant input

rates to each active cell, which we take as the mean itself. The results are shown in

Figures 5 and 6. We present evidence relating to mossy fibre → granule cell linear

transmission in the Methods section (Methods, Linear transmission of mossy fibres to

granule cells).

The narrowing and predicted shape of the granule cell frequency distribution is evident

in all conditions. The field means (representing the mean of granule cell rates in each

field) approach a straight line, the test that a rank operates as a single functional unit.

When mossy fibre rates are normally distributed (SD 12.5 Hz), the granule cell

distribution is very narrow (and normal if rates are counted in narrower bins) and the

field means are very close to a straight line. The field means follow the mean of mossy

fibre rates at a precisely fixed distance. In all tested conditions, granule cell rates have

a narrowed range, which is the same in all fields, clustered around the mean, and

attracted to a normal distribution (more evident with narrowly and/or normally

distributed mossy fibre rates). This holds even though a single granule cell receives a

tiny fraction of mossy fibre rates, and each field receives only a small fraction of mossy

fibre signals received by a rank, and regardless of the shape of the mossy fibre

distribution.

We conclude that, plausibly, granule cell firing conserves mossy fibre rate information,

coded in the output of a whole rank. This is therefore the minimum resolution of

granule cell information. However, it is not the minimum unit of code. The same

information is coded in any random sample of granule cell signals, over a minimum

number. It is immaterial (i.e. codes nothing) which particular cells are active or which

cells fire at what rates, or even what rates those are, provided any random sample of

rates is as if drawn from the same frequency distribution, with replacement. The

output of a rank can be likened by analogy to a pixel: it is only divisible into smaller

pieces of the same information.

Fig 5. A rank functions as a single unit. The same simulation as Figure 4 but data are granule cell firing rates or derived from granule cell firing rates. **Column 1** Mossy fibre rates were generated in the same way as earlier figures (the number of signals is the same in all rows). **Column 2** Pale green: sample means – the mean of input rates to each granule cell in a single field selected at random; blue: sample means of the subset of granule cells which meet the conditions to fire. In rows A–C, ~30% fire. In rows D–F, ~1% fire, y axes are rescaled and green data are omitted. **Column 3** Red: mean of mossy fibre input rates to each field in a rank, including duplicate signals; pink: SD of mossy fibre input rates to each field; blue: field means (i.e., the mean of granule cell firing rates in each field); pale blue: SD of granule cell firing rates in each field. Regardless of the shape and large range of the mossy fibre distribution, the field means always converge towards a straight line which is a fixed distance from the overall mean of mossy fibre rates. The amount and direction of field-to-field random fluctuation of the blue data is independent of local fluctuation of the red data, and in a smaller range. Uncertainty inherent in random sampling – first by single fields of input rates to a rank, and then by Golgi cells and granule cells of input rates to a field – is not compounded.

Fig 6. Granule cells collectively code mossy fibre rate information. Column 1 Mossy fibre rates were generated in the same way as earlier Figures, with either a normal (SD 12.5) or uniform distribution representing input to a row of 100 fields, with copies added – the sampled distribution. **Column 2** Histogram of granule cell firing rates in a randomly selected field, obtained in the same way as in the bottom 3 rows of Fig 5. **Column 3** Grey line: the mean of the sampled distribution; blue line: field means. **A–C** The fields means are precisely and linearly related to the mean of mossy fibre rates. Blue data approach a straight line which is parallel to the grey line at a mean distance of 15 Hz (SD 1.21 Hz). **D–F** If the shape and mean of a uniform sampled distribution are unchanging, the distance between the blue and grey lines is linearly related to the range.

6. **Discussion**

6.1 Conclusions

Network models are often a simplified representation of physiology to provide a substrate for sophisticated computations. We find the reverse: physiology is highly adapted to provide a substrate for straightforward computations. In summary we find:

1. There is a computational effect, unaided and unlearned, of the combination of cell morphologies, network architecture and linear units, for which the granular layer is highly adapted. There is no algorithm.

2. Physiology is the physical form of a network of linear units. A unit may be a cell, part

of a cell or some other structure. Units randomly sample afferent activity. Code groups

communicate by a facsimile of independent random sampling. Anatomy replicates the

same result as if single neurons independently randomly sample the entire population

of firing rates in the afferent cell layer, notwithstanding that single cells receive fixed

contact from a tiny fraction of afferent cells.

3. Data processing exploits statistical effects of random sampling, such that i) inhibition

of granule cells in a rank is synchronised and tracks mossy fibre rates; and ii) granule

cell rates are in a condensed range, reliably (because statistically) well-predicted at

collective level, and again linearly related to mossy fibre rates.

4. High-dimensional mossy fibre information recodes as a very low-resolution granule

cell code. Large granule cell groups function as a single unit, coding information in their

collective activity.

5. At any instant, a whole group codes the same information. Accordingly, information

has spatial dimensions, which can be likened to 3-dimensional pixels. Cell groups take

their dimensions from a combination of topography and cell type morphologies, rather

than having physical boundaries.

6. Pixels span the cerebellar network in the sagittal direction, from side to side,

mirroring microzonal organisation of the molecular layer. Accordingly, granule cell

encoded information has a striped pattern viewed from the cerebellar surface.

7. While firing of a whole code group is the minimum unit of information, it is not the

minimum unit of code. Inside code group boundaries the same information is coded in

any random sample of granule cell firing rates. We might say for convenience of

reference the code is homogenised.

8. The shape of data and homogenised code are key to function, having results

discussed below.

6.2 Control of Purkinje cells

There is a huge sacrifice of data resolution on entry to the cerebellum, counter to the

general expectation that data processing is sophisticated. This is functional. Coded in

this form, granule cell data can drive coherent firing of large cell groups downstream.

Microzones lie at right angles to parallel fibres and parallel to code groups. The

statistical form of the granule cell code means the same information is received by all

locations of sagittal plane, so that microzone-grouped Purkinje cells can receive the

same information without receiving the same signals. An estimated 350,000 parallel

fibres pass through a Purkinje cell dendritic territory [45-48]. If even only a small

fraction of parallel fibres is active (0.5%), and allowing for the fact that only one in two

make contact, and that a majority of synapses are severely depressed or silent [49, 50],

a single Purkinje cell still receives around 200 parallel fibre signals at working synapses

at any time. A granule cell code in this form provides a means to modulate firing of

Purkinje cells in unison by a chain of rate codes: mossy fibre → granule cell → Purkinje

cell. There is evidence that Purkinje cells linearly code the strength of granule cell

inputs [6-8] and linearly code behavioural metrics [3, 51-54].

6.3 Replicability

A statistical code may be a physiological strategy to mitigate variable performance of

single neurons and synapses, a long-known impediment to precise and replicable

biological information capture. The traditional cerebellar learning model [43, 55, 56]

relies on (and assumes) replicability of patterns of active and inactive parallel fibres,

because pattern replication is necessary for signals to match up with the correct

training-modified synaptic weights. Learned changes of synaptic weights are also a

core feature of artificial neural network learning models.

However, this may not be safe. First, it is not clear that input to the cerebellum is ever

replicated. Around 4,000 mossy fibres innervate the region that supplies parallel fibre

input to a single Purkinje cell. It may not be a safe assumption that conditions

upstream are routinely repeated with precision sufficient to consistently (or perhaps

ever) generate a repeat of the same inputs. Second, single neurons and synapses are

not reliably consistent performers. Neurons even of the same type are by no means

replicas, and synaptic transmission involves random variables. Third, the granular layer

is *designed* to amplify the effect on the pattern of active parallel fibres if the pattern of

active mossy fibres should vary, even only by a modest number of cells. Indeed, even

that is not necessary – the same pattern of active mossy fibres but firing in a different

permutation of rates also generates a different pattern of active granule cells, and

therefore parallel fibres. Small differences of input (the pattern of active mossy fibres

and/or permutation of rates they fire at) are converted by decorrelation to wholesale

changes to output (the pattern of active parallel fibres). Possibly, and we submit, the

cerebellum actively eschews reliance on pattern replication in favour of group code

which gives a more (because statistically) reliable outcome.

6.4 Working 'blind'

A perennial issue for network theory is that biological networks do not seem to have

much information about the signals they receive. Input to the cerebellum is eclectic

and in multiple modalities. In some cases, a single granule cell can receive multi-modal

input. In the traditional cerebellar model, supervision solves this problem. It is

unnecessary for the cerebellum itself to discriminate because it is an assumption of the

model that externally-sourced instruction signals train the correct response.

We propose instead the solution that resolution of high-dimensional input data is

sacrificed. The cerebellum disregards source and modality data below a physical scale

threshold. It is therefore unnecessary to discriminate below that scale. Mapping

studies show that mossy fibre signals (evoked by cutaneous stimulation) share the

topographical organisation of climbing fibres [57-59]. At sub-topographical

dimensions, signal source and type are simply discounted. Therefore, networks do not

need to be adapted to the source or type of input they receive – anatomically,

networks are interchangeable, so network design can be modular. This feature may

have been necessary to allow basic cerebellar wiring to survive almost unchanged

through dramatic upheavals of the vertebrate phenotype [60, 61].

 METHODS

DERIVATION OF MOSSY FIBRE RATES

The number of mossy fibres that innervate a rank

As far as is known, fields receive a random sample mossy fibre input rates to a rank. To
simulate that, we need the size of the sampled population – the number of mossy fibre signals
received by a rank – and sample size, the number received by a field.

Despite the repeating 'crystalline' architecture of the cerebellar cortex, at local level there are
anatomical random variables. Variables are: i) the number of mossy fibres that innervate a
field, ii) the number of terminals they each end in, and iii) the number of those that are inside
field limits.

The number of mossy fibres afferent to a field has been estimated at 100 [21]. However, that
number is a field-sized fraction of the general population. The actual number is larger because
mossy fibres end in a dispersed group (cluster) of terminals, and most groups straddle field
limits.

We work from an estimate of the total number of terminals per field, which is, in turn, derived
from convergent estimates of the number of granule cells per field. (Estimate 1) There are an
estimated 1.92×10^6 granule cells per µl [45]. The number of fields which fit in that volume is
$(1,000/150)^2 \times (1,000/200) = \sim222$, assuming a field depth of 150 µm [21]. The number of
granule cells in a field is, therefore 1,920,000/222 = 8,649. (Estimate 2) An estimated 350,000
parallel fibres pass through a single Purkinje cell's territory [45]. Parallel fibres run about 3 mm
in each direction from the granule cell soma [62, 63]. The Purkinje cell dendritic arbour fills the
molecular layer vertically and averages ~200 µm wide. Therefore, 6,000/150 = 40 fields fit in a
row that provides parallel fibres that pass through a Purkinje cell dendritic territory, giving
350,000/40 = 8,750 granule cells per field. Estimates vary of the number of granule cells which
extend a dendrite into a single glomerulus. 50 is mid-range. (8,750/50) x 4 = 700 glomeruli per
field.

Granule cells have 3–5 dendrites, averaging 4, and receive input to each of them from a single

mossy fibre (so an average granule cell receives 4 inputs). Estimates of divergence of a single

mossy fibre terminal onto granule cells vary [2]. It is unclear how much this reflects biological

diversity versus differences in opinion and methods. Here we use a mid-range estimate of 1:50

[34, 35]. Assuming a mossy fibre terminal receives a single granule cell dendrite from each of d

granule cells, and given an average of 4 dendrites per granule cell and z granule cells per field,

the total number of mossy fibre terminals in a field is $N = (z \times 4)/d = 700$ given $z = 8,750$ and d

$= 50$. A convergent estimate is provided by multiplying the per field number of mossy fibres by

the average number of terminals per cluster.

Cluster size varies. We take the range as 4–12 terminals per cluster [21]. We assume the

number varies at random and cluster sizes occur with equal probability. A maximum of 7 fits in

a field (because a field shares the dimensions of the area that encloses a mean cluster). Fields

are not anatomically bounded, so many clusters straddle field limits. To reflect that, we

assume there is a 1/7 probability that a 7-terminal cluster (or larger) has 7 terminals inside a

field and also the same probability of any other number. A 6-terminal cluster has a 2/7 chance

that all 6 terminals are inside and a 1/7 chance of any other number, and so on.

Table 1 shows the derivation of the expected (i.e., predicted by probability) ratios of terminals

contributed to a field by each cluster size (R1), not counting terminals that lie outside when a

cluster straddles field boundaries; and the ratios of mossy fibres contributing n = 1–7 terminals

(R2), regardless of cluster size. R1 ratios are given by the sum of numbers in each row. So, for

example, the ratio of cells that end in 8 terminals to cells that end in 5 is 28:25. R2 ratios are

given by the number of appearances of each number in the grid – so the expected ratio of cells

which contribute 7 terminals to the number which contribute 4 is 6:12. The ratios effectively

give the distributed probability of possible states. For example, there is a probability of 10/(6 +

8 + 10 + 12 + (3 x 9)) = 0.159 that a mossy fibre contributes 5 terminals to a field (regardless of

cluster size). Similarly, there is a probability of ((6 x 28)/((6 x 28) + 27 + 25 + 22))/7 = 0.099 that

a given cell contributes 7 terminals to a field.

Table 1. Derivation of expected ratios

size	number of terminals inside							R1
12	7	6	5	4	3	2	1	**28**
11	7	6	5	4	3	2	1	**28**
10	7	6	5	4	3	2	1	**28**
9	7	6	5	4	3	2	1	**28**
8	7	6	5	4	3	2	1	**28**
7	7	6	5	4	3	2	1	**28**
6	6	6	5	4	3	2	1	**27**
5	5	5	5	4	3	2	1	**25**
4	4	4	4	4	3	2	1	**22**
R2	6	8	10	12	9	9	9	

If R1 ratios are denoted by a series $x_4, x_5 \ldots x_{12}$, and a given element in the series is x_j, the

fraction of terminals provided by a given cluster size is $x_j/(x_4 + x_5 + \cdots + x_{12})$, and this

fraction is multiplied by N to give the number of terminals. The number of terminals is divided

by the expected mean number of terminals that fall inside field limits (for that cluster size) to

give the expected number of mossy fibres that provide them. The number of j-terminal mossy

fibres that innervate a field is therefore given by

$$N \left[x_j \Big/ \sum_{i=1}^{n} x_i \right] \Big/ \left[x_j / 7 \right]$$

The results are shown in Table 2. The equal number of mossy fibres (mfs) in each column is

what we would expect, because we made it an assumption that all sizes occur with equal

probability, and mf values are expected values – i.e., in proportions predicted by their

probability. The mf total is the expected number of mossy fibres afferent to a field (which

contribute at least one terminal).

Table 2. The expected number of mossy fibres which innervate a field

cluster size	12	11	10	9	8	7	6	5	4	total
terminals	81	81	81	81	81	81	78.1	72.3	63.6	700
mfs	20.25	20.25	20.25	20.25	20.25	20.25	20.25	20.25	20.25	182.25

821   R2 ratios – the relative incidence of afferent mossy fibres that contribute 7 terminals to a field,

822   6 terminals, 5 and so on, down to 1 – and the expected number of mossy fibres that contribute

823   them, are shown in Table 3.

824   **Table 3. The number of mossy fibres which contribute *n* = 1–7 terminals to a field**

n	7	6	5	4	3	2	1	total
R2	6	8	10	12	9	9	9	63
R3	42	48	50	48	27	18	9	
mfs	17.34	23.15	28.92	34.72	26.04	26.04	26.04	182.25
%age	9.52	12.69	15.87	19.05	14.29	14.29	14.29	

R3 is *n* x R2, giving the ratio of terminals provided by mossy fibres with *n* terminals inside a

field. %age is given by (R2/63) x 100. this is the percentage of mossy fibres with *n* terminals

inside a field.

From here, the number of mossy fibres that innervate a rank is given by 182.25 x 100 divided

by the mean number of clusters per mossy fibre per rank. If a rank receives, on average, five

clusters per mossy fibre, ~3,645 mossy fibres innervate a rank. This is the estimate used in the

simulation unless stated otherwise.

Assumptions. Cluster sizes occur with equal probability. The probabilities of each outcome

when a cluster straddles field boundaries (how many terminals are inside and how many

outside) are as stated. A single mossy fibre ends in an average of 5 terminal clusters per rank.

Mossy fibre input rates to a field

Terminal clustering means a field receives multiple copies of most signals. The proportions of

mossy fibres afferent to a simulated field that contribute each possible number of terminals

(1–7) are generated in simulated steps. In the first step, the simulation randomly generates,

for each field, the cluster size ratio. This means, the relative proportions of afferent mossy

fibres (which contribute at least 1 terminal to a field) that end in 4 terminals, 5, 6 and so on up

to 12, assuming they occur with equal probability, disregarding how many terminals fall inside

and outside field boundaries. In step two, the ratio of cluster sizes is weighted to reflect the

greater probability that large clusters contribute more terminals to a field. In step 3, ratios are

converted to the fraction of terminals contributed by each cluster size. When multiplied by the

total number of terminals, this gives the number of terminals contributed by each cluster size.

In step four, the values obtained in step three are divided up between cells that contribute a

single terminal, 2 terminals, 3 and so on, to reflect straddling of field limits. The proportions

give the number of mossy fibres afferent to a simulated field that contribute a single terminal,

two, three and so on up to seven, the maximum.

There is no field-level effect of mossy fibre terminal branching. Terminal clusters arising from

the same mossy fibre are separated by a minimum distance, so that a field cannot contain

terminals of more than one cluster per mossy fibre.

Terminal clustering means that fields contain a randomly variable number of copies of most

signals. In the simulation, copies are added after sampling of mossy fibre rates to obtain rates

received by a field. An alternative order would be to add copies to the sampled population and

then sample that. We follow anatomy: the number of *rates* received by a field is equal to the

number of afferent mossy fibres; the number of *signals* contains duplicate rates.

DERIVATION OF ENSEMBLE DIMENSIONS AND MODEL PARAMETERS

4 Purkinje cells fit in a field, based on field size, thickness of the Purkinje cell dendritic plate – 9

μm [64] – and the distance between them – ~40 μm [65]. 'In man, the ratio [of Purkinje cells

to Golgi cells] is 1⊡1.5, while in the monkey and cat it is almost 1⊡1.9 and in the rat 1⊡3.3' [66].

Possibly, estimates are low because smaller Golgi cells at deeper level were not counted [65

p.121]. We take it that there are 10 Golgi cells per field, a ratio of 1:2.5 with 4 Purkinje cells.

Golgi cell morphology creates ensemble dimensions. The Golgi cell axon ramifies profusely,

giving rise to a dense plexus. The axonal field is sagittally elongated – that is, in the direction of

the long axis of a microzone (mean range 650 +/- 179 µm by 180 +/- 40 µm [in mice: 67]) – and

is the depth of the granular layer, which it fills vertically. Fine, beaded axon fibres enter

glomeruli, where they inhibit granule cells [68]. Because of the dimensions and orientation of

the Golgi cell plexus, all Golgi cells in a sagittal row of three fields inhibit substantially the

whole of the middle field. Each field defines a functional population of Golgi cells – an

ensemble – being those cells with convergent input to that field. A single Golgi cell contacts

thousands of granule cells [68, 69].

Contact by mossy fibres on a Golgi cell is on basal dendrites. 4–6 basal dendrites arise from the

Golgi cell soma. We assume 5, so a population of ensemble-grouped Golgi cells has a

combined total of 30 x 5 = 150 basal dendrites. The number of mossy fibres which contact a

Golgi cell is unreported. Dendrites radiate from the cell body. A single dendrite has a range of

50–100 µm [65]. The longest dimension of a field – which contains ~700 mossy fibre terminals

– is 200 µm. We estimate the mean number of mossy fibres that contact a single dendrite is 4,

so 20 per cell, or ~3% of the population of terminals (or 0.6% per dendrite).

Glomeruli individually receive innervation from a random sample of Golgi cells afferent to a

field. It is unknown how many Golgi cells innervate a glomerulus.

LINEAR TRANSMISSION OF MOSSY FIBRES TO GOLGI CELLS

It is part of our proposal that Golgi cell basal dendritic signals linearly reflect mossy fibre rates,

so that mossy fibre rate information is conserved proportionally in Golgi cell firing rates. Golgi

cells are large interneurons whose cell bodies and basal dendrites lie in the granular layer.

Mossy fibres directly contact Golgi cell basal dendrites. Golgi cell firing frequency increases

linearly with the amplitude of depolarising current [29 p.845]. 'Golgi cells can follow peripheral

signals in a continuous fashion, modulating their frequency with the intensity of the stimulus

[citing [70, 71]]' [72]. As a result, during movement, the 'output of Golgi cells varies smoothly

as a function of input....They are capable of modulating their output to reflect their input with

little noise or signal distortion' [12]. 'Sensory-evoked Golgi-cell inhibition scales proportionally

with the level of mossy fiber excitatory synaptic input', such that inhibition reliably conveys

mossy fibre rate information [4]. On this evidence: mossy fibre rate information is conserved

proportionally in Golgi cell firing rates.

Golgi cells extend their basal dendrites into glomeruli [68, 73]. Contact on them by mossy

fibres is multi-synaptic [65], contributing to a reliable, rapid (submillisecond) response. The

fast rise time and weak distance dependence of the amplitude and timing of Golgi cell EPSCs

evoked by mossy fibre signals, suggests that basal dendrites introduce relatively little filtering

[11, 28], as expected for large-diameter dendrites. We note that the pause in Golgi cell firing

following a discrete stimulus under anaesthesia [4, 74, 75] disappears from Golgi cell

recordings during locomotion [71].

During behaviour in freely-moving animals, mossy fibre activity is dense: a high fraction of

mossy fibres are active [76]. Both mossy fibres [77-79] and Golgi cells [71] have been reported

to fire with a sustained, time-varying signature. In the behaving animal, Golgi cell basal

dendritic membrane potential is a continuous variable under modulation by sustained inputs,

we submit. As charge transfer to the soma is passive, polarisation of the soma is likewise

sustained, under modulation by dendritic states.

It is worth noting that very high mossy fibre rates can be generated in experimental conditions.

However, the typical physiological range is 50–300 Hz [78]. In the simulations, we use the

physiological rates, within stated constraints on the shape of the distribution.

Other control of Golgi cells? Do Golgi cells receive a significant influence of input from other

sources? The evidence is incomplete but currently suggests other influence is absent or weak.

Contrary to early reports, neither Purkinje cells nor climbing fibres contact Golgi cells [80, who

give references], and only a modest minority of inhibitory inputs to Golgi cells (if any) are from

molecular layer interneurons [81], which generate weak synaptic currents [82], consistent with

either extremely weak or wholly absent innervation [83]. There is conflicting evidence whether

Golgi cells make inhibitory synaptic contact on each other [83, 84]. Our simulation does not

include an effect on Golgi cell firing by other sources of input.

LINEAR TRANSMISSION OF GOLGI CELLS TO GRANULE CELLS

'In the adult rat cerebellum, each granule cell dendrite receives 2.6 ± 0.55 synaptic contacts

from Golgi axon terminals' [30] citing [35].[1] However, the large majority of inhibition (98%) is

by spillover [5], where neurotransmitter released into the synaptic cleft spills out into the

glomerular space. Golgi cells release GABA, an inhibitory neurotransmitter. This is detected by

high-affinity $GABA_A$ receptors located perisynaptically and extrasynaptically on granule cells

[31, 32]. Even synaptically-received signals have most of their effect (i.e., the large majority of

charge transfer is mediated) via spillover.[2]

Golgi cells fire spontaneously [85] at a time-varying rate in the behaving animal, so that

downstream, a glomerulus receives continuous convergent input. As a result, there is a

sustained build-up of glomerular GABA during behaviour [86] at an adjustable concentration

controlled by the mean of afferent Golgi cell firing rates [87]. A glomerulus randomly samples

Golgi cells afferent to a field. Golgi cells provide the only inhibitory input to granule cells.

Signalling by spillover is sometimes assumed to be ambient and slow. However, the action of

glomerular GABA spillover has a fast phasic component – not as fast as synaptic transmission

(~1 ms) but with a rise time of only a few milliseconds [87]. Unlike the spiky appearance of

synaptically-induced inhibitory postsynaptic currents, spillover [4] generates a sustained

---

[1] Data are sparse. The original data are an estimate derived from visual inspection of slices thirty years ago so far without later corroboration, except from Rossi and Hamann 1998, who derive a lower estimate but propose the disparity is because their estimate is too low.
[2] Because of the long duration of IPSCs generated by spillover (recorded in slices), the total charge carried is three times that of IPSCs generated by directly connected terminals. 30.   Rossi, D.J. and M. Hamann, *Spillover-mediated transmission at inhibitory synapses promoted by high affinity alpha6 subunit GABA(A) receptors and glomerular geometry*. Neuron, 1998. **20**(4): p. 783-95.

outward current. The response time to multiple inputs is likely faster than the experimentally

isolated response to a discrete signal (because the combined rate of GABA release is higher).

Golgi cell oscillations. Golgi cells fire autonomously. Under anaesthesia, firing falls into a slow

oscillating pattern [88]. Under excitatory synaptic input, however, this disappears [84].

Anaesthesia 'has a strong influence on spontaneous activity of Golgi cells' [80]. Discharge at

low rates with irregular timing under anaesthesia [85] is replaced by higher rates with more

regular timing without anaesthesia [70, 71, 89]. (This paragraph is a cautionary note about

evidence obtained under anaesthesia, which has been used to argue that oscillations provide a

form of signalling.)

LINEAR TRANSMISSION OF MOSSY FIBRES TO GRANULE CELLS

There is significant evidence that granule cell firing rates are linearly related to input rates, but

also evidence thought to be unfavourable to that conclusion. We take them in turn.

The short and equal length of granule cell dendrites would suggest light and equal filtering.

The mossy fibre-granule cell connection has a range of adaptations which support high-fidelity

transmission of high frequency signals across a wide bandwidth [9]. Faithful transmission of

rate information is reported [1, 2]. Vesicle release and replenishment are fast [10, 13] and

postsynaptic AMPA receptors operate in their linear range [13], where they are resistant to

desensitisation [90]. Multiple contacts are made by a mossy fibre with each granule cell [35].

Conversion of depolarising somatic charge into granule cell firing is a simple function, such that

granule cells 'have a relatively linear and uncomplicated conversion of depolarisation level to

spike rate' [39 p.2393, citing Jorntell and Ekerot 2006 and D'Angelo et al 1998].

Glutamate spillover enhances precision and reliability of transmission [13, 91]. During

behaviour, at sustained physiological mossy fibre rates, there may be a build-up of

intraglomerular glutamate, increasing the relative influence of spillover in a balance with

synaptic transmission, assisted by short-term synaptic depression. In these conditions,

spillover may dominate, as it dominates inhibitory transmission [5, 87]. Spillover from multiple

release sites (200-400 at a single mossy fibre terminal) is plausibly sufficient to mitigate

variability of vesicular size, release probability, and the number of release points at synapses

made on any individual cell, increasing fidelity and equality of excitation of granule cells.

Against that, there are reported to be heterogeneous synaptic weights. Different strength and

short-term dynamics of the mossy fibre to granule cell connection have been reported in

vestibular slices, with $GABA_A$ receptors blocked. The amplitude of the response, measured at

the soma, is reported to be statistically correlated to the source of the mossy fibre that is

stimulated [92], suggesting that the strength of the connection depends on the source.

However, the presence of weights does not necessarily mean that the function is to modulate

firing rates. Rather, it is suggested that the function may temporally code the source of inputs

because different combinations have different response onset times. An alternative, which

they also mention, is that some inputs may be in a supporting role, reminiscent of 'driver' and

'modulatory' signals in the thalamus and cortex [93]. Feasibly, signals in a supporting role must

be present for the postsynaptic granule cell to fire, but do not affect the granule cell firing rate.

Grants

Funding: This work was supported by the Leverhulme Trust [grant number ECF-2022-079] (Mike Gilbert); the Swedish Research Council [grant number 2020-01468]; the Per-Eric och Ulla Schybergs Foundation [grant number 42630]; and the Crafoord Foundation [grant numbers 20180704, 20200729, 20220776, & 20230655] (Anders Rasmussen).

Acknowledgement

Dr Mike Evans, School of Mathematics, University of Leeds, LS2 9JT, UK, read an early draft of the paper and provided feedback.

Author contributions

M.G. Researched, conceived and developed the ideas, performed the analysis, wrote the code, prepared the figures and wrote the manuscript. A.R. provided expertise and feedback, and reviewed and provided editorial comments on the manuscript.

Declaration of interests

The authors declare no competing interests.

995   **REFERENCES**

996   1.   Rancz, E.A., et al., *High-fidelity transmission of sensory information by single cerebellar*
997   *mossy fibre boutons.* Nature, 2007. **450**(7173): p. 1245-8.
998   2.   Ritzau-Jost, A., et al., *Ultrafast action potentials mediate kilohertz signaling at a central*
999   *synapse.* Neuron, 2014. **84**(1): p. 152-163.
1000   3.   Jelitai, M., et al., *Dendritic excitation-inhibition balance shapes cerebellar output*
1001   *during motor behaviour.* Nat Commun, 2016. **7**: p. 13722.
1002   4.   Duguid, I., et al., *Control of cerebellar granule cell output by sensory-evoked Golgi cell*
1003   *inhibition.* Proc Natl Acad Sci U S A, 2015. **112**(42): p. 13099-104.
1004   5.   Duguid, I., et al., *Tonic inhibition enhances fidelity of sensory information transmission*
1005   *in the cerebellar cortex.* J Neurosci, 2012. **32**(32): p. 11132-43.
1006   6.   Walter, J.T. and K. Khodakhah, *The linear computational algorithm of cerebellar*
1007   *Purkinje cells.* J Neurosci, 2006. **26**(50): p. 12861-72.
1008   7.   Walter, J.T. and K. Khodakhah, *The advantages of linear information processing for*
1009   *cerebellar computation.* Proc Natl Acad Sci U S A, 2009. **106**(11): p. 4471-6.
1010   8.   Park, S.M., E. Tara, and K. Khodakhah, *Efficient generation of reciprocal signals by*
1011   *inhibition.* J Neurophysiol, 2012. **107**(9): p. 2453-62.
1012   9.   Delvendahl, I. and S. Hallermann, *The Cerebellar Mossy Fiber Synapse as a Model for*
1013   *High-Frequency Transmission in the Mammalian CNS.* Trends Neurosci, 2016. **39**(11): p.
1014   722-737.
1015   10.   Saviane, C. and R.A. Silver, *Fast vesicle reloading and a large pool sustain high*
1016   *bandwidth transmission at a central synapse.* Nature, 2006. **439**(7079): p. 983-7.
1017   11.   Cesana, E., et al. *Novel granule cell-Golgi cell excitatory input in the cerebellar granular*
1018   *layer.* in *FENS Abstr.* 2010.
1019   12.   Moore, J.W. and D.E.J. Blazis, *11 - Simulation of a Classically Conditioned Response: A*
1020   *Cerebellar Neural Network Implementation of the Sutton–Barto–Desmond Model*, in
1021   *Neural Models of Plasticity*, J.H. Byrne and W.O. Berry, Editors. 1989, Academic Press.
1022   p. 187-207.
1023   13.   Sargent, P.B., et al., *Rapid vesicular release, quantal variability, and spillover contribute*
1024   *to the precision and reliability of transmission at a glomerular synapse.* J Neurosci,
1025   2005. **25**(36): p. 8173-87.
1026   14.   Turecek, J., S.L. Jackman, and W.G. Regehr, *Synaptic Specializations Support*
1027   *Frequency-Independent Purkinje Cell Output from the Cerebellar Cortex.* Cell Reports,
1028   2016. **17**(12): p. 12.
1029   15.   Turecek, J., S.L. Jackman, and W.G. Regehr, *Synaptotagmin 7 confers frequency*
1030   *invariance onto specialized depressing synapses.* Nature, 2017. **551**: p. 503.
1031   16.   Raymond, J.L. and J.F. Medina, *Computational Principles of Supervised Learning in the*
1032   *Cerebellum.* Annu Rev Neurosci, 2018. **41**: p. 233-253.
1033   17.   Alvina, K., et al., *Questioning the role of rebound firing in the cerebellum.* Nat Neurosci,
1034   2008. **11**(11): p. 1256-8.
1035   18.   Schonewille, M., et al., *Purkinje cells in awake behaving animals operate at the upstate*
1036   *membrane potential.* Nat Neurosci, 2006. **9**(4): p. 459-61; author reply 461.

19. Oscarsson, O., *Functional units of the cerebellum - sagittal zones and microzones.*
Trends in Neurosciences, 1979. **2**: p. 143-145.
20. Wu, H.S., I. Sugihara, and Y. Shinoda, *Projection patterns of single mossy fibers*
originating from the lateral reticular nucleus in the rat cerebellar cortex and nuclei. J
Comp Neurol, 1999. **411**(1): p. 97-118.
21. Sultan, F. and D. Heck, *Detection of sequences in the cerebellar cortex: numerical*
estimate of the possible number of tidal-wave inducing sequences represented. J
Physiol Paris, 2003. **97**(4-6): p. 591-600.
22. Shinoda, Y. and I. Sugihara, *Axonal Trajectories of Single Climbing and Mossy Fiber*
Neurons in the Cerebellar Cortex and Nucleus, in *Handbook of the Cerebellum and*
Cerebellar Disorders, S.J.D. Manto M., Rossi F., Gruol D.L., Koibuchi N., Editor. 2013,
Springer, Dordrecht.
23. Shinoda, Y., et al., *The entire trajectory of single climbing and mossy fibers in the*
cerebellar nuclei and cortex. Prog Brain Res, 2000. **124**: p. 173-86.
24. Garwicz, M., C.F. Ekerot, and H. Jorntell, *Organizational principles of cerebellar*
neuronal circuitry. News in Physiological Sciences, 1998. **13**: p. 28-34.
25. Ozden, I., et al., *Reliable coding emerges from coactivation of climbing fibers in*
microbands of cerebellar Purkinje neurons. J Neurosci, 2009. **29**(34): p. 10463-73.
26. Dean, P., et al., *The cerebellar microcircuit as an adaptive filter: experimental and*
computational evidence. Nat Rev Neurosci, 2010. **11**(1): p. 30-43.
27. Eccles, J.C., M. Ito, and J. Szentágothai, *The cerebellum as a neuronal machine.* 1967,
Berlin, New York etc.: Springer-Verlag. 335 p.
28. Kanichay, R.T. and R.A. Silver, *Synaptic and cellular properties of the feedforward*
inhibitory circuit within the input layer of the cerebellar cortex. J Neurosci, 2008.
28(36): p. 8955-67.
29. Dieudonné, S., *Submillisecond kinetics and low efficacy of parallel fibre-Golgi cell*
synaptic currents in the rat cerebellum. J Physiol, 1998. **510 (Pt 3)**: p. 845-66.
30. Rossi, D.J. and M. Hamann, *Spillover-mediated transmission at inhibitory synapses*
promoted by high affinity alpha6 subunit GABA(A) receptors and glomerular geometry.
Neuron, 1998. **20**(4): p. 783-95.
31. Nusser, Z., W. Sieghart, and P. Somogyi, *Segregation of different GABAA receptors to*
synaptic and extrasynaptic membranes of cerebellar granule cells. J Neurosci, 1998.
18(5): p. 1693-703.
32. Brickley, S.G., S.G. Cull-Candy, and M. Farrant, *Development of a tonic form of synaptic*
inhibition in rat cerebellar granule cells resulting from persistent activation of GABAA
receptors. J Physiol, 1996. **497 (Pt 3)**: p. 753-9.
33. Gilbert, M. and R.C. Miall, *How and Why the Cerebellum Recodes Input Signals: An*
Alternative to Machine Learning. Neuroscientist, 2021a. **DOI:**
10.1177/1073858420986795.
34. Gao, Z., et al., *Excitatory Cerebellar Nucleocortical Circuit Provides Internal*
Amplification during Associative Conditioning. Neuron, 2016. **89**(3): p. 645-57.
35. Jakab, R.L. and J. Hámori, *Quantitative morphology and synaptology of cerebellar*
glomeruli in the rat. Anat Embryol (Berl), 1988. **179**(1): p. 81-8.

36.   Mitchell, S.J. and R.A. Silver, *GABA spillover from single inhibitory axons suppresses*
1081   *low-frequency excitatory transmission at the cerebellar glomerulus.* J Neurosci, 2000a.
1082   **20**(23): p. 8651-8.
1083   37.   Mitchell, S.J. and R.A. Silver, *Glutamate spillover suppresses inhibition by activating*
1084   *presynaptic mGluRs.* Nature, 2000b. **404**(6777): p. 498-502.
1085   38.   Chadderton, P., T.W. Margrie, and M. Hausser, *Integration of quanta in cerebellar*
1086   *granule cells during sensory processing.* Nature, 2004. **428**(6985): p. 856-60.
1087   39.   Bengtsson, F. and H. Jörntell, *Sensory transmission in cerebellar granule cells relies on*
1088   *similarly coded mossy fiber inputs.* Proc Natl Acad Sci U S A, 2009. **106**(7): p. 2389-94.
1089   40.   Billings, G., et al., *Network structure within the cerebellar input layer enables lossless*
1090   *sparse encoding.* Neuron, 2014. **83**(4): p. 960-74.
1091   41.   Jörntell, H. and C.F. Ekerot, *Properties of somatosensory synaptic integration in*
1092   *cerebellar granule cells in vivo.* J Neurosci, 2006. **26**(45): p. 11786-97.
1093   42.   Cesana, E., et al., *Granule cell ascending axon excitatory synapses onto Golgi cells*
1094   *implement a potent feedback circuit in the cerebellar granular layer.* J Neurosci, 2013.
1095   **33**(30): p. 12430-46.
1096   43.   Albus, J.S., *A Theory of Cerebellar Function.* Mathematical Biosciences, 1971. **10**(1-2):
1097   p. 25-61.
1098   44.   Gilbert, M. and A. Rasmussen, *Gap Junctions May Have A Computational Function In*
1099   *The Cerebellum: A Hypothesis.* The Cerebellum, 2024.
1100   45.   Harvey, R.J. and R.M. Napper, *Quantitative study of granule and Purkinje cells in the*
1101   *cerebellar cortex of the rat.* J Comp Neurol, 1988. **274**(2): p. 151-7.
1102   46.   Harvey, R.J. and R.M. Napper, *Quantitative studies on the mammalian cerebellum.*
1103   Prog Neurobiol, 1991. **36**(6): p. 437-63.
1104   47.   Napper, R.M. and R.J. Harvey, *Number of parallel fiber synapses on an individual*
1105   *Purkinje cell in the cerebellum of the rat.* J Comp Neurol, 1988. **274**(2): p. 168-77.
1106   48.   Napper, R.M. and R.J. Harvey, *Quantitative study of the Purkinje cell dendritic spines in*
1107   *the rat cerebellum.* J Comp Neurol, 1988. **274**(2): p. 158-67.
1108   49.   Isope, P. and B. Barbour, *Properties of unitary granule cell-->Purkinje cell synapses in*
1109   *adult rat cerebellar slices.* J Neurosci, 2002. **22**(22): p. 9668-78.
1110   50.   Ekerot, C.F. and H. Jörntell, *Parallel fibre receptive fields of Purkinje cells and*
1111   *interneurons are climbing fibre-specific.* Eur J Neurosci, 2001. **13**(7): p. 1303-10.
1112   51.   Dugué, G.P., et al., *Cerebellar re-encoding of self-generated head movements.* Elife,
1113   2017. **6**.
1114   52.   Chen, S., G.J. Augustine, and P. Chadderton, *The cerebellum linearly encodes whisker*
1115   *position during voluntary movement.* eLife, 2016. **5**: p. e10509-e10509.
1116   53.   Herzfeld, D.J., et al., *Encoding of action by the Purkinje cells of the cerebellum.* Nature,
1117   2015. **526**(7573): p. 439-42.
1118   54.   Hong, S., et al., *Multiplexed coding by cerebellar Purkinje neurons.* Elife, 2016. **5**.
1119   55.   Fujita, M., *Adaptive filter model of the cerebellum.* Biol Cybern, 1982. **45**(3): p. 195-
1120   206.
1121   56.   Marr, D., *A theory of cerebellar cortex.* J Physiol, 1969. **202**(2): p. 437-70.
1122   57.   Apps, R. and M. Garwicz, *Anatomical and physiological foundations of cerebellar*
1123   *information processing.* Nat Rev Neurosci, 2005. **6**(4): p. 297-311.

1124   58.   Apps, R. and R. Hawkes, *Cerebellar cortical organization: a one-map hypothesis.* Nat
1125   Rev Neurosci, 2009. **10**(9): p. 670-81.
1126   59.   Ji, Z. and R. Hawkes, *Topography of purkinje cell compartments and mossy fiber*
1127   *terminal fields in lobules ii and iii of the rat cerebellar cortex: Spinocerebellar and*
1128   *cuneocerebellar projections.* Neuroscience, 1994. **61**(4): p. 935-954.
1129   60.   Larsell, O., *The Comparative Anatomy and Histology of the Cerebellum from Myxinoids*
1130   *through Birds.* 1967: The University of Minnesota Press.
1131   61.   Larsell, O., *The Comparative Anatomy and Histology of the Cerebellum from*
1132   *Monotremes through Apes.* 1970: The University of Minnesota Press, Minneapolis.
1133   62.   Mugnaini, E., *The length of cerebellar parallel fibers in chicken and rhesus monkey.* J
1134   Comp Neurol, 1983. **220**(1): p. 7-15.
1135   63.   Brand, S., A.L. Dahl, and E. Mugnaini, *The length of parallel fibers in the cat cerebellar*
1136   *cortex. An experimental light and electron microscopic study.* Exp Brain Res, 1976.
1137   **26**(1): p. 39-58.
1138   64.   Hámori, J., *Anatomy and neurochemical anatomy of the cerebellum*, in *Cerebellar*
1139   *degenerations : clinical neurobiology*, A. Plaitakis, Editor. 1992, Kluwer Academic.
1140   65.   Palay, S.L. and V. Chan-Palay, *Cerebellar cortex : cytology and organization.* 1974,
1141   Berlin: Springer.
1142   66.   Lange, W., *Regional differences in the distribution of golgi cells in the cerebellar cortex*
1143   *of man and some other mammals.* Cell Tissue Res, 1974. **153**(2): p. 219-26.
1144   67.   Barmack, N.H. and V. Yakhnitsa, *Functions of interneurons in mouse cerebellum.* J
1145   Neurosci, 2008. **28**(5): p. 1140-52.
1146   68.   Hámori, J. and J. Szentágothai, *Participation of Golgi neuron processes in the cerebellar*
1147   *glomeruli: an electron microscope study.* Exp Brain Res, 1966. **2**(1): p. 35-48.
1148   69.   D'Angelo, E., et al., *The cerebellar Golgi cell and spatiotemporal organization of*
1149   *granular layer activity.* Front Neural Circuits, 2013. **7**: p. 93.
1150   70.   Miles, F.A., et al., *Long-term adaptive changes in primate vestibuloocular reflex. III.*
1151   *Electrophysiological observations in flocculus of normal monkeys.* J Neurophysiol, 1980.
1152   **43**(5): p. 1437-76.
1153   71.   Edgley, S.A. and M. Lidierth, *The discharges of cerebellar Golgi cells during locomotion*
1154   *in the cat.* J Physiol, 1987. **392**: p. 315-32.
1155   72.   Galliano, E., P. Mazzarello, and E. D'Angelo, *Discovery and rediscoveries of Golgi cells.* J
1156   Physiol, 2010. **588**(Pt 19): p. 3639-55.
1157   73.   Eccles, J.C., R. Llinás, and K. Sasaki, *Golgi cell inhibition in the cerebellar cortex.* Nature,
1158   1964. **204**: p. 1265-6.
1159   74.   Holtzman, T., et al., *Different responses of rat cerebellar Purkinje cells and Golgi cells*
1160   *evoked by widespread convergent sensory inputs.* J Physiol, 2006. **574**(Pt 2): p. 491-
1161   507.
1162   75.   Vos, B.P., A. Volny-Luraghi, and E. De Schutter, *Cerebellar Golgi cells in the rat:*
1163   *receptive fields and timing of responses to facial stimulation.* Eur J Neurosci, 1999.
1164   **11**(8): p. 2621-34.
1165   76.   Hartmann, M.J. and J.M. Bower, *Tactile responses in the granule cell layer of cerebellar*
1166   *folium crus IIa of freely behaving rats.* J Neurosci, 2001. **21**(10): p. 3549-63.
1167   77.   Arenz, A., et al., *The contribution of single synapses to sensory representation in vivo.*
1168   Science, 2008. **321**(5891): p. 977-80.

78. van Kan, P.L., A.R. Gibson, and J.C. Houk, *Movement-related inputs to intermediate*
cerebellum of the monkey. J Neurophysiol, 1993. **69**(1): p. 74-94.
79. Giovannucci, A., et al., *Cerebellar granule cells acquire a widespread predictive*
feedback signal during motor learning. Nat Neurosci, 2017.
80. Pietrajtis, K. and S. Dieudonné, *Golgi Neurons*, in *Handbook of the Cerebellum and*
Cerebellar Disorders, M. Manto, et al., Editors. 2013, Springer Netherlands: Dordrecht.
p. 829-852.
81. Eyre, M.D. and Z. Nusser, *Only a Minority of the Inhibitory Inputs to Cerebellar Golgi*
Cells Originates from Local GABAergic Cells. eNeuro, 2016. **3**(2).
82. Dumoulin, A., A. Triller, and S. Dieudonné, *IPSC kinetics at identified GABAergic and*
mixed GABAergic and glycinergic synapses onto cerebellar Golgi cells. J Neurosci, 2001.
21(16): p. 6045-57.
83. Hull, C. and W.G. Regehr, *Identification of an inhibitory circuit that regulates cerebellar*
Golgi cell activity. Neuron, 2012. **73**(1): p. 149-58.
84. Vervaeke, K., et al., *Rapid desynchronization of an electrically coupled interneuron*
network with sparse excitatory synaptic input. Neuron, 2010. **67**(3): p. 435-51.
85. Ruigrok, T.J., R.A. Hensbroek, and J.I. Simpson, *Spontaneous activity signatures of*
morphologically identified interneurons in the vestibulocerebellum. J Neurosci, 2011.
31(2): p. 712-24.
86. Rossi, D.J., M. Hamann, and D. Attwell, *Multiple modes of GABAergic inhibition of rat*
cerebellar granule cells. J Physiol, 2003. **548**(Pt 1): p. 97-110.
87. Mapelli, L., S. Solinas, and E. D'Angelo, *Integration and regulation of glomerular*
inhibition in the cerebellar granular layer circuit. Front Cell Neurosci, 2014. **8**: p. 55.
88. Ros, H., et al., *Neocortical networks entrain neuronal circuits in cerebellar cortex.* J
Neurosci, 2009. **29**(33): p. 10309-20.
89. Rasmussen, A., et al., *Golgi cell activity during eyeblink conditioning in decerebrate*
ferrets. Cerebellum, 2014. **13**(1): p. 42-5.
90. DiGregorio, D.A., et al., *Desensitization properties of AMPA receptors at the cerebellar*
mossy fiber granule cell synapse. J Neurosci, 2007. **27**(31): p. 8344-57.
91. DiGregorio, D.A., Z. Nusser, and R.A. Silver, *Spillover of glutamate onto synaptic AMPA*
receptors enhances fast transmission at a cerebellar synapse. Neuron, 2002. **35**(3): p.
521-33.
92. Chabrol, F.P., et al., *Synaptic diversity enables temporal coding of coincident*
multisensory inputs in single neurons. Nat Neurosci, 2015. **18**(5): p. 718-27.
93. Sherman, S.M. and R.W. Guillery, *Distinct functions for direct and transthalamic*
corticocortical connections. J Neurophysiol, 2011. **106**(3): p. 1068-77.

1205