1

WHAT IF…? COMPUTATIONAL ANATOMY IN THE GRANULAR LAYER OF

THE CEREBELLAR CORTEX

Mike Gilbert[a] (corresponding author), Anders Rasmussen[b]

[a]School of Psychology, College of Life and Environmental Sciences, University of
Birmingham B15 2TT, UK

[b]Department of Experimental Medical Science, Lund University, BMC F10, 22184 Lund,
Sweden

Correspondence: m.gilbert.1@bham.ac.uk

Summary

Neural networks are believed to carry out sophisticated computations. In the influential traditional cerebellar learning model, and many artificial neural networks, computations are implemented by learned synaptic changes that selectively adjust synaptic transmission strength. These ideas have become widely influential. We propose an alternative in the form of a reinterpretation of the evidence relating to the granular layer of the cerebellar cortex, the entry point of the cerebellum. The main proposals are that there is a computational effect, unaided, of the anatomy and spatial organisation of granular layer cells and wiring, with the result that high resolution input data are converted on entry into a (very) low resolution internal code. Internally-coded information has spatial dimensions which mirror the functional organisation of the cerebellar cortex into long thin strips called microzones. Data are coded in the collective activity of granule cells at this scale. There is no finer internal resolution. This feels counter-intuitive because it is counter to the general expectation that high dimensionality increases the power of brain computations. However, it means the cerebellum is able to handle all input the same way, and it provides an otherwise elusive mechanism to orchestrate firing of large cell groups – microzones – downstream, which powerfully modulate firing of the output cells of the cerebellar network. There is no algorithm and the response is unlearned. The model is not abstract and has good evidence.

36 **Keywords:** cerebellum, theory, model, computational anatomy, cerebellar cortex,

37 granular layer, network, code.

1. Introduction

The cerebellum typically makes up 20–30 % of the vertebrate brain by mass and volume (about half that in humans). Most of that is made up of the cerebellar cortex, which is divided into two layers. The inner layer, the granular layer, receives the main excitatory input to the cerebellum, from mossy fibres. Mossy fibres contact both granule cells and large inhibitory interneurons, Golgi cells. Golgi cells in turn inhibit granule cells. Each mossy fibre terminal is surrounded by a semi-permeable membrane, a structure called a glomerulus. Golgi cells extend their axons into glomeruli, so granule cells receive both excitation and inhibition there. The granule cell axon rises into the outer layer of the cerebellar cortex, the molecular layer, where it divides in two. The two branches, termed parallel fibres, travel in opposite directions making synaptic contact in passing on Purkinje cells, which carry the sole output of the cerebellar cortex.

In the influential supervised learning model (Raymond and Medina, 2018, for an excellent review), control of Purkinje cells by granule cell rates is displaced by learned changes of parallel fibre synaptic weights (Albus, 1989; Brunel et al., 2004; Dean et al., 2010; Fujita, 1982). This obviates a requirement to explain how granule cell firing rates are derived from mossy fibre rates, or encode mossy fibre rate information, because it is implicit that a granule cell rate code would be redundant, because control of Purkinje cells is surrendered to learning.

There is accumulating evidence that rate-coded information is linearly conserved in

communication between cerebellar neurons (Duguid et al., 2015; Duguid et al., 2012;

Jelitai et al., 2016; Park et al., 2012; Rancz et al., 2007; Ritzau-Jost et al., 2014; Walter

and Khodakhah, 2006, 2009), and that cells are adapted for faithful transmission

(Cesana et al., 2010; Delvendahl and Hallermann, 2016; Moore and Blazis, 1989;

Sargent et al., 2005; Saviane and Silver, 2006; Turecek et al., 2016, 2017). Firing rates

that linearly code behavioural and task metrics have been observed 'at all levels of the

cerebellar circuit' (Raymond and Medina, 2018 p.239, citing 30 sources). Evidence of

linear signalling in the granular layer is rich. To decongest the main text, we include a

detailed review in the Methods section, to which we cross-refer from the text.

It is difficult to reconcile linear relationships with network models that use synaptic

weight changes because the theorised function of weight changes is to supplant a

purely rate-coded response of networked units. Modelling the behaviour of a large cell

network can feel like a blunt trauma anyway. For one thing, physiological units are not

generic. Cells and synapses even of the same type both vary. We work with 'sampling

units' – these can be cells, parts of cells, or some other structure. It forms part of our

proposal that the cerebellum exploits statistical effects of random sampling of afferent

rate data. Sampling units provide model parameters. We field evidence in the main

text to explain the derivation of model parameters. Parameter values are supplied by

the literature.

We use sampling units on the following basis in reasons. Cell-to-cell transmission

involves the complex interaction of a large array of often co-dependent biophysical

and other variables which are specific to the cell types making and receiving contact.

Models vary in how they address that. Broadly, models tend to be divided between

network models and biological neuron models. The difference is often partly a scale

preference – network models sacrifice biophysics and biological neuron models aren't

networks. On the face of it, the task is to identify (or select or abstract) and relate

parameters to give the expected outcome. We present the view that this is not a

modelling problem but a biological problem. We mean, the cerebellum must solve the

problem of disengaging or otherwise compensating for intrinsic non-linear properties

of neurons and synapses (Alvina et al., 2008; Sargent et al., 2005; Schonewille et al.,

2006; Turecek et al., 2016). The reason is to isolate the effect of variables that code

information and control for all other variables. Other variables are noise, including

biophysical variables. The significance of evidence of linearity is that despite the

seeming clamour, linear relationships are in fact isolated. In this view, the cerebellum

is made up of a repeating network of linear units because it is adapted to isolate

computational variables by an aggressive strategy of noise reduction. The units are

sampling units.

We take it, in general, that output of a sampling unit is a linear function of the mean of

input rates, $f(\langle r \rangle)$, which we take as the mean itself. We envisage that in motor

circuits during behaviour signalling is analogue, avoiding the noise caused by discrete

signals. So, for example, Golgi cell dendritic signalling is a sustained state under

modulation by fluctuating mossy fibre rates. Similarly, GABA spillover into the

intraglomerular space, almost exclusively the agent of granule cell inhibition, becomes

elevated by a build-up during behaviour, so that this too is at a sustained level,

controlled by afferent Golgi cell rates. Thus, at each step, data are coded at any

moment in a snapshot of a continuously variable state of change.

Historically, it has been challenging to imagine a useful function of linear units. We

propose it has a statistical outcome which emerges at scale. We submit that

(i) there are reliable statistical effects of concurrent unit computations, which relate

mossy fibre rates and granule cell rates at collective level; in

(ii) functionally-bounded cell groups, even though the granular layer is an anatomically

seamless mesh of neurons; and that this is

(iii) the unaided result of the triad of cell type morphologies, network geometry and

linear cerebellar signalling, and that it

(iv) provides a substrate for rate information coded in mossy fibre signals to be

converted to control and coordination of Purkinje cells without synaptic weight

changes in the historically expected role. This is because the form and spatial

dimensions of the granule cell code mean all locations of a microzone receive the same

code without receiving the same signals.

As we use a common function, we use a normalised scale. The output of the model is

the granule cell code, and a regulated fraction of active granule cells. We include a

statement of the relevant parts of the hypothesis at or near the beginning of each

section.

123   2. Mossy fibres

124   2.1 Anatomy, 'ranks' and 'files'

Mossy fibres give rise to sagittally-extending collaterals, the direction of the long axis

of microzones (Fig 1). These in turn give rise to multiple distal branches, each ending in

a cluster of terminals (Shinoda and Sugihara, 2013; Shinoda et al., 2000; Sultan and

Heck, 2003; Wu et al., 1999). Terminal clusters are sagittally aligned and separated by

a randomly variable distance typically not less than 200 µm (Sultan and Heck, 2003;

Wu et al., 1999) measured from edge to edge of clusters (so they are separated by a

greater distance measured from centre to centre). Terminals themselves are intimately

and randomly intermingled with terminals of other mossy fibres.

Accordingly, mossy fibres do not terminate at a point but a region with dimensions:

the field occupied by a terminal cluster. A single mossy fibre gives rise to several

terminal branches and each of those to several terminals, such that a sagittal strip of

the granular layer receives multiple copies of every signal at each of multiple locations.

Cluster size is variable (because it depends on the number of terminals and how tightly

clustered they are), averaging 200 µm sagittally x 150 µm mediolaterally (Sultan and

Heck, 2003). We term the average dimensions of a cluster a 'field'. The granular layer

is anatomically seamless. A field is simply a nominal division that we use for modelling.

A sagittal row of 75–100 fields laid end to end has the area (viewed from the cerebellar

surface) of a mid-range microzone (Dean et al., 2010; Garwicz et al., 1998; Oscarsson,

1979; Ozden et al., 2009). We term a row of fields a 'rank'. Fields in a rank are not

anatomically *or* functionally discrete – they are simply a device.

Figure 1

2.2 Hypothesis

Mossy fibre anatomy is an anatomical solution to a computational problem. The

cerebellum exploits statistical effects of random sampling called the central limit

theorem. For the effects to occur, samples must be independent, also termed sampling

'with replacement' (in the illustration sometimes used, coloured balls selected at

random from a bag are put back in the bag between samplings). Clearly, physical

replacement is impossible because neither a terminal cluster nor a terminal can be in

more than one place at the same time. However, anatomy simulates replacement. The

distributed probabilities of the number of terminal branches per mossy fibre are the

same as the probabilities of the number of repeat samplings if rates received by a rank

were in fact randomly sampled with replacement by fields. Similarly, the distributed

probabilities of the number of terminals per cluster are the same as the probabilities of

number of repeat samplings if rates received by a field were in fact randomly sampled

with replacement by individual neurons. Note that it is the population of *rates* that are

hypothetically sampled with replacement, and not *signals.* The population of signals

includes copies of rates because of branching and clustering. Copies are an anatomical

device to approximate the result of sampling rates with replacement.

2.3 Terminal branching simulates sampling with replacement

If each mossy fibre could terminate in only one field, a mossy fibre 'sampled' by one

field clearly cannot also be sampled by any others. Termination by a single mossy fibre

in a random number of terminal branches, at multiple locations, removes that

restriction.

In theory, sampling with replacement should mean any number of fields can receive

the same signal, up to all of them. Clearly this is not true – there is still an anatomical

limit. But in practice even with real replacement the number of repeat samplings is

constrained by a probability distribution. If, then, the number of terminal branches per

mossy fibre has the same probability distribution, branching has the same result as if

firing rates received by each field in a rank were in fact a random sample with

replacement of firing rates received by the rank as a whole.

We can test that by calculating the probability distribution in the hypothetical case

that mossy fibres did not branch terminally and rates were in fact sampled with

replacement. The probability, p, that a given rate is selected k times in taking N

samples, sample size x, assuming mossy fibres each give rise to an average of m

terminal branches, is given by

$$\left[{Nx!}/{k!\,(Nx-k)!}\right] * \left[1/{(Nx/m)}\right]^{k} * \left[1 - \left(1/{(Nx/m)}\right)\right]^{Nx-k}$$

184   The results are shown in Table 1.

185   **Table 1. Probability p of k repeat selections of a single mossy fibre**

k	0	1	2	3	4	5	6	7	8	9	10	11
p	.007	.034	.084	.140	.175	.175	.146	.104	.065	.036	.018	.008

Although data are scarce, the number of repeat selections is consistent with what we

know about the biological range of the number of terminal clusters per mossy fibre

(Eccles et al., 1967; Sultan and Heck, 2003). Estimated by this method, the mean is 4–

5. The anatomical cap on the number of fields that receive a signal (set by the number

of terminal branches per cell) has the same distributed probability as the probabilities

of the number of repeat samplings if the population of firing rates received by a rank

was randomly sampled with replacement 100 times, representing 100 fields. Anatomy

simulates replacement.

2.4 Terminal clusters simulate sampling with replacement

Terminal clustering of mossy fibres has the same function as terminal branching but at

field scale. As far as we know, granule cells and Golgi cells receive contact from a

random sample of mossy fibres that innervate a field, and more exactly a random

sample of terminals. If a mossy fibre terminal branch ended in a single terminal,

contact on any particular cell would place it beyond the range of most other cells.

Mossy fibre termination in a cluster of terminals, and random intermingling of

terminals (Fig 1), removes that restriction. Again, to approximate independence, it is

only necessary for the number of repeat selections of a signal to have the same

probability distribution as if there was actual replacement.

A practical problem with the strategy of clustering is that a target could receive the

same signal twice (or more). The risk is mitigated by taking small samples. A sample

size of 4 is small enough for a low risk. Granule cells average 4 dendrites, each

receiving contact from a single terminal, and we estimate that Golgi cells receive an

average of 4 inputs to each of their basal dendrites (see derivation of parameters in

the next section). Given m x 100 terminals and an independent and equal chance that

contact is by any terminal, the probability of an n^{th} contact by the same mossy fibre on

a single dendrite is:

$$P_n = \left[\prod_{x=2}^{n} \left(\frac{m - x + 1}{100m - x + 1} \right) \right] \times \frac{y!}{x!\,(y - x)!}$$

where y is the number of inputs to a single dendrite and m is the number of terminals

per cluster, provided by (for illustration) 100 mossy fibres. Calculated in this way,

around one in every twenty samples contains a duplicate, and less than two in a

thousand contain more than one. The actual fractions are smaller because the number

of mossy fibres that innervate a field is higher, around 180 (see Methods, Derivation of

mossy fibre rates).

So, each field receives, at any given instant in time, a quasi-independent random

sample of mossy fibre input rates to a rank, and single neurons individually receive a

quasi-independent random sample of input rates to a field. The combined effect is a

biological facsimile of simultaneous random sampling with replacement by single

neurons of input rates *to a whole rank*.

3. Golgi cells

3.1 Derivation of ensemble dimensions and model parameters

4 Purkinje cells fit in a field, based on field size, thickness of the Purkinje cell dendritic plate – 9 µm (Hámori, 1992) – and the distance between them – ~40 µm (Palay and Chan-Palay, 1974). 'In man, the ratio [of Purkinje cells to Golgi cells] is 1⬚1.5, while in the monkey and cat it is almost 1⬚1.9 and in the rat 1⬚3.3' (Lange, 1974). Possibly, estimates are low because smaller Golgi cells at deeper level were not counted (Palay and Chan-Palay, 1974 p.121). We take it that there are 10 Golgi cells per field, a ratio of 1:2.5 with 4 Purkinje cells.

Golgi cell morphology creates ensemble dimensions. The Golgi cell axon ramifies profusely, giving rise to a dense plexus. The axonal field is sagittally elongated – that is, in the direction of the long axis of a microzone (mean range 650 +/- 179 µm by 180 +/- 40 µm (in mice: Barmack and Yakhnitsa, 2008)) – and is the depth of the granular layer, which it fills vertically. Fine, beaded axon fibres enter glomeruli, where they inhibit granule cells (Hámori and Szentágothai, 1966). Because of the dimensions and 3-dimensional orientation of the Golgi cell plexus, all Golgi cells in a sagittal row of three fields inhibit substantially the whole of the middle field. Each field defines a functional population of Golgi cells – an ensemble – being those cells with convergent input to

that field. A single Golgi cell contacts thousands of granule cells (D'Angelo et al., 2013;

Hámori and Szentágothai, 1966).

Contact by mossy fibres on a Golgi cell is on basal dendrites. 4–6 basal dendrites arise

from the Golgi cell soma. We assume 5, so a population of ensemble-grouped Golgi

cells has a combined total of 30 x 5 = 150 basal dendrites. The number of mossy fibres

which contact a Golgi cell is unreported. Dendrites radiate from the cell body. A single

dendrite has a range of 50–100 μm (Palay and Chan-Palay, 1974). The longest

dimension of a field – which contains ~700 mossy fibre terminals – is 200 μm. We

estimate the mean number of mossy fibres that contact a single dendrite is 4, so 20

per cell, or ~3% of the population of terminals (or 0.6% per dendrite).

Glomeruli individually receive innervation from a random sample of Golgi cells afferent

to a field. It is unknown how many Golgi cells innervate a glomerulus.

3.2 Hypothesis

Despite forming an anatomically seamless dense mesh of axonal filaments that fills the

granular layer, Golgi cells work in groups: ensembles. An ensemble is the population of

a sagittal row of three fields. Groups are not anatomically bounded or even

functionally discrete, but simply function as a single unit for the anatomical reason that

the range of Golgi cells in 3 fields includes all or most of the middle field. Ensembles

perform a computational conversion of mossy fibre rates into inhibition of granule

cells. The computation is (chiefly) a statistical result of random sampling, in two steps.

At the first step, Golgi cell basal dendrites receive contact from a random sample of

mossy fibres that innervate a field (strictly speaking, they randomly sample terminals);

and in the second step, middle-field glomeruli each receive innervation by a random

sample of Golgi cells. The effect of the computation is to synchronise inhibition of

granule cells in the middle field. The strength of inhibition is linearly related to the

mean of mossy fibre firing rates received as input to the entire rank, even though only

a modest fraction of rates are received by three fields.

Ensembles conflate sagittally anatomically and functionally. Ensembles in a sagittal

strip all return the same output (because they all perform the same computation on a

large random sample of the same data). Effectively: the Golgi cell population of a

whole rank functions as a single unit.

3.3 The Golgi cell computation

Say a population of values (with any frequency distribution) is randomly sampled.

There are n samples, sample size m. If we take the mean (or sum) of each sample, and

n is large, the sample means have a normal or near-normal distribution, by the central

limit theorem. (This works even if m is a range.) The new distribution is centred on the

same mean as the sampled distribution, but narrower. Modest values of n or m give an

untidy result. However, if the new distribution is then itself randomly sampled, and the

process is repeated, the results get tidier and narrower.

We simulated random sampling by Golgi cells of mossy fibre rates, and downstream,

random sampling by glomeruli of Golgi cell rates, both with replacement. We randomly

generated mossy fibre rates received by each field in a rank, taking into account mossy

fibre terminal branching and clustering, random variation of the number of terminals

per cluster and clusters per cell, and the fact that most clusters fall partly inside and

partly outside a field (see Methods, Derivation of mossy fibre rates). Rates were

obtained by sampling from a distribution representing signals received as input to the

rank as a whole, which were randomly generated subject to constraints on the shape

of the distribution.

This gave the number of mossy fibre signals, firing rates and the number of copies of

each rate received by each field, and therefore input to an ensemble territory,

populated by 30 Golgi cells, 10 per field. There are ~700 mossy fibre terminals per

field. As far as we know (and we propose), contact by mossy fibres on Golgi cells is at

random at field scale, i.e., inputs to a single basal dendrite are provided by a random

sample of local terminals. The number of physiological samplings is equal to the

number of basal dendrites per field; sample size is the number of mossy fibres which

make contact on each dendrite. The effect is sustained dendritic depolarisation under

modulation by a linear function of input rates (see Methods, Linear transmission of

mossy fibres to Golgi cells), which we take as the mean.

Taking the sample means generates a new set of values (Fig 2 column 2) with a smaller

range and centred on the mean, which is linearly related to the mean of the sampled

distribution. Basal dendritic charge transfer is passive and little filtered (Cesana et al.,

2010; Kanichay and Silver, 2008). Integration of dendritic states at the Golgi soma is

represented by taking the mean of dendritic charge, for each simulated cell. Golgi cell

firing rates have a linear relationship with the amplitude of depolarising current

(Dieudonné, 1998), suggesting (and we assume) a linear conversion of somatic

membrane potential into the Golgi cell firing rate.

The output is a third data set, Golgi cell firing rates (Fig 2 column 3), on the same

normalised scale. Each glomerulus in the middle field receives innervation from a

random subset of Golgi cells afferent to the field, and accordingly independently

randomly samples Golgi cell firing rates. Golgi cells inhibit granule cells in the

glomerulus. A granule cell extends a single dendrite into a glomerulus, typically

receiving synaptic contact from two or three Golgi cells (Rossi and Hamann, 1998).

However, inhibition of granule cells by Golgi cells is almost entirely by GABA spillover

into the intraglomerular space (Duguid et al., 2012), via extrasynaptic receptors

(Brickley et al., 1996; Nusser et al., 1998) (Methods, Linear transmission of Golgi cells

to granule cells). This effectively increases the Golgi cell to granule convergence ratio,

since the number of Golgi cells that are functionally afferent to a granule cell is not the

number that make direct contact, but the number afferent to a glomerulus. The

membrane around a glomerulus restricts GABA diffusion so that signalling is by

modulation of the intraglomerular concentration of GABA, which in turn reflects the

rate it is released. The rate it is released by a single cell is proportional to the firing rate

of that cell. The sum of rates is proportional to the mean. The convergence ratio is

sample size; the number of samples is the number of glomeruli per field (~700, the

same as terminals). We took the sample means to represent intraglomerular GABA

concentration, which physiologically is under continuous modulation at a sustained

level which is a linear function of the mean of afferent Golgi cell rates. The number of

Golgi cells afferent to a glomerulus is unreported, as far as we know. We take it as a

random number in the range 8–12, generated for each glomerulus, to reflect likely

physiological variation. We tested other ratios to confirm that the ratio does not affect

the ensemble computation (Supplementary Materials).

This generates a final population of values (Fig 2 columns 4 and 5), the output of the

ensemble conversion. There are 700 outputs, the number of glomeruli in the middle

field, representing inhibition of granule cells. All granule cells that extend a dendrite

into the same glomerulus receive equal inhibition (of that dendrite).

3.4 Results

3.4.1 An ensemble

There is a progressive narrowing of the data range at each step. Inputs to the system

are converted to narrowly-grouped and normally-distributed outputs. Input values

represent mossy fibre signals received by a row of 3 fields; output values represent

inhibition of granule cells in the middle field by GABA spillover in glomeruli. The

narrow output range indicates that inhibition is locally synchronised, i.e., granule cells

in a field receive co-modulated inhibition. Inhibition is statistically constrained to a

linear relationship with mossy fibre rates, acting through the ensemble conversion.

These results are independent of the shape of mossy fibre rates.

Figure 2

3.4.2 A rank

We next used the ensemble conversion to derive inhibition of granule cells in a sagittal

row of 100 fields, a rank. The results are shown in Figure 3. As ensembles all perform

the same computation and take as input (at any moment) the same frequency

distribution of input rates, or nearly the same distribution, they all return the same

output, or nearly the same output. There are 700 outputs per field. The mean of

outputs, taken for each field, has a narrow range centred on the mean of mossy fibre

input rates to the whole rank. Field-to-field variation of output appears to be inherent

in the mechanism, i.e., caused by uncertainty of the outcome of random sampling, so it

has a predictable range which the system can be built to tolerate.

So: Golgi cells in a rank act as a single functional unit. Ensembles conflate in the sagittal

direction functionally as they do anatomically. As a result, the whole granule cell

population *of a rank* receive synchronised inhibition which is linearly related to the

mean of mossy fibre rates.

365   **Figure 3**

367 4. Granule cells

368 4.1 The fraction of active granule cells is reliably regulated by feed-

369 forward inhibition at low density

370 4.1.1 Hypothesis

Anatomy simulates independent random sampling by a single granule cell of the

population of mossy fibre input rates to the whole rank (section 2). There is a

competition for influence between mossy fibre excitation and Golgi cell inhibition of

granule cells. The competition takes place in the glomerulus. During behaviour, this

takes place ceaselessly. The strength of inhibition is synchronised between glomeruli

that populate a rank (section 3), so that all mossy fibre signals compete, at any

moment, against the same inhibitory threshold. The outcome in each glomerulus is

independent of the outcome at other terminals, including other terminals that

innervate the same granule cell. To fire, a granule cell must receive a minimum

number of mossy fibre inputs that prevail in the glomerular competition. The

probability that enough prevail depends, at any moment, on the strength of inhibition.

Since inhibition is synchronised, the probability is the same for all granule cells at any

time. The number that fire is very near the expected proportion (the number predicted

by probability) because granule cells are numerous. Because the linear relationship of

mossy fibre firing rates and inhibition is always conserved by the Golgi cell

computation, the probability is invariant, and therefore so is the proportion of active

granule cells.

Note: the idea of an all-or-nothing result of the intraglomerular 'balance' of excitation

and inhibition has appeared previously (Gilbert and Miall, 2021a).

4.1.2 Anatomy

Granule cells receive input to each dendrite from a single mossy fibre. Granule cells

have 3–5 dendrites, averaging 4 (the simulation assumes 4, but the result is the same

with 3–5). Each dendrite of a single granule cell extends into a different glomerulus.

Estimates vary of the number of granule cells that receive contact from a single mossy

fibre terminal. 50 is mid-range (Gao et al., 2016; Jakab and Hámori, 1988), but

estimates vary significantly (Ritzau-Jost et al., 2014). 50 is consistent with our

estimates of the number of terminals and granule cells in a field. The granule cell

sampling unit is the whole cell since it receives only a single input rate to each

dendrite. Sample size is therefore 4, and the number of samples per field is the

number of granule cells, which we estimate at 8,750 (see Methods).

4.1.3 Feed-forward inhibition regulates the probability that granule cells fire

Excitation and inhibition compete for influence in a glomerulus. It is unclear if there is

always an emphatic winner, but a swing in the dominant direction may be amplified by

positive feedback, acting through presynaptic receptors. Activation of presynaptic

$GABA_B$ receptors on mossy fibres inhibits glutamate release (Mitchell and Silver,

2000a), while activation of presynaptic mGluR2 receptors on Golgi cells inhibits GABA

release (Mitchell and Silver, 2000b). We postulate that only mossy fibre signals which

prevail in the glomerular competition contribute significantly to granule cell

depolarisation. To fire, a granule cell must receive a minimum number of competitive

mossy fibre signals – thought to be three (Billings et al., 2014; Jörntell and Ekerot,

2006). Certainly, one is too few (Bengtsson and Jörntell, 2009; Chadderton et al., 2004;

Duguid et al., 2012).

The probability that a given granule cell receives three competitive signals and

therefore fires is a function of the percentage of mossy fibre signals that are

competitive. The expected number of granule cells that fire in a field is given by

$$E_{fld} = T \sum_{x=j}^{n} \left(\binom{n}{x} [P(e > i)]^x [1 - P(e > i)]^{n-x} \right)$$

where T is the total number of granule cells per field, n is the number of dendrites per

granule cell, and j is the minimum number of dendrites that must receive dominant

excitation, $e > i$. The proportion of granule cells that fire is near the expected number

by the law of large numbers. The percentage of mossy fibre signals that are

competitive is always the same because inhibition is synchronised and scale with the

mean of mossy fibre rates (because Golgi cells translate mossy fibre firing rates into

proportional inhibition). As the percentage is unvarying, so is the probability.

4.1.4 Does physiology have that outcome?

Probability is an abstraction with no direct physiological counterpart. The simulation

does not include a binomial function but derives output from input by comparing

excitatory to inhibitory input to each glomerulus, the way we claim it is derived

physiologically. The equation is simply shorthand – biology exploits chance and large

numbers to generate the same result. To test that, we incorporated the Golgi cell

ensemble conversion into a model that recodes mossy fibre signals into granule cell

signals. The model compares mossy fibre signal frequency with inhibition at each of 4

dendrites of each of 8,750 granule cells per field, in each of 100 fields. Mossy fibre

input rates to each field are derived per section 3 and Methods, so taking into account,

inter alia, that a field receives a randomly variable number of copies of most signals,

and most terminal clusters lie only partly inside a field. Inhibition – outputs of the

ensemble conversion, 700 per field – is provided by the Golgi cell model. Excitation and

inhibition received by each glomerulus are randomly paired. Pairing is used to identify

the subset of granule cells that fire (cells that receive at least 3 mossy fibre signals

which exceed the inhibitory threshold). Output of the model is the number of granule

cells that fire in each field, the mean of mossy fibre rates they each receive (the sample

means), and the mean of the sample means in each field ('field means'). The results

are shown in Figure 4.

4.1.5 Results

Feed-forward inhibition – mossy fibre → Golgi cell → granule cell – may be the

primary regulator of the number of granule cells that fire. Field-to-field random

variability of the number is inversely proportional to the fraction that are active,

suggesting that, if primary regulation is in this form, the fraction is low. We anticipate

nonetheless a smaller range relative to the mean, indicating the physiological presence

of a supplementary mechanism that we have not represented in the model.

A local feedback mechanism may be provided by granule cell ascending axon contact

on Golgi cells. The ascending portion of the granule cell axon makes synaptic contact

on local Golgi cells, on basal dendrites (Cesana et al., 2010). A Golgi cell receives ~400

functional synaptic contacts from this source, around 25% of the granule cells that a

Golgi cell receives contact from in total. During sustained mossy fibre discharge,

synapses from local granule cells contribute equivalent charge to Golgi cells as mossy

fibre synapses and 'display similarly fast kinetics' (Cesana et al., 2013). Short-latency

inhibitory feedback may add homeostatic fine adjustment to regulation at local level,

at field scale.

It is a long-standing idea (Albus, 1971) that granule cell activity may be regulated by

feedback via parallel fibres, which contact Golgi cell apical dendrites. We have

previously proposed that gap junctions between apical dendrites may provide the

means for Golgi cells to detect the density of parallel fibre activity (Gilbert and

Rasmussen, 2024). However, feedback by this pathway is not sensitive to local

fluctuation of granule cell active numbers.

The results are the same with any (tested) distribution of mossy fibre rates and also

whether mossy fibres are all active or some are active and others are not. Active cells

can be in any random permutation, subject to there being a minimum number that are

active at any time.

470   **Figure 4**

4.2 Granule cells encode mossy fibre rate information at a collective level that mirrors organisation of the molecular layer into microzones

4.2.1 Hypothesis

Anatomy simulates independent random sampling by a single granule cell of the population of mossy fibre input rates to the whole rank (section 2). The granule cell firing rate is a linear function of depolarising somatic charge, in turn a linear function of the mean (or sum) of dendritic depolarisations. Cells that fire receive rate-proportional dendritic depolarisation that is undiluted by inhibition because they receive dominant mossy fibre signals. Derived in this way, granule cell rates are collectively well predicted by mossy fibre rates, by the central limit theorem. This provides the substrate for granule cells to collectively encode mossy fibre rate information. Code groups are functionally defined by the topography of mossy fibre input to the cerebellum, as Golgi cell networks are. The output of a rank is the smallest unit of granule cell encoded information. The permutation of active mossy fibres, and of granule cells, and how firing rates are distributed among active cells, all code nothing.

4.2.2 Do granule cell firing rates – the output of a rank – have the hypothesised features?

We ran the same simulation as section 4.1 to test whether and how mossy fibre rates

are related to granule cell rates, and if fields in a rank all have the same output. We

take granule cell firing rates to be a linear function of the mean of input rates to each

active cell, which we take as the mean itself. Field means are therefore the mean of

granule cell firing rates in each field. The results are shown in Figures 5 and 6. We

present evidence relating to mossy fibre $\rightarrow$ granule cell linear transmission in the

Methods section (Methods, Linear transmission of mossy fibres to granule cells).

The narrowing and predicted shape of the granule cell frequency distribution is evident

in all conditions. The field means approach a straight line, the test that a rank operates

as a single functional unit. When mossy fibre rates are normally distributed (SD 12.5

Hz), the granule cell distribution is very narrow (and normal if rates are counted in

narrower bins) and the field means are very close to a straight line. The field means

follow the mean of mossy fibre rates at a precisely fixed distance. When mossy fibre

rates have a random uniform distribution, the field means remain almost a straight

line, and the distance from the mossy fibre mean varies with the size of the mossy

fibre range.

A straight line indicates that fields all return the same output – not the same firing

rates, but the same narrowed range, clustered around the mean, and attracted to a

normal distribution (more evident with narrowly and/or normally distributed mossy

fibre rates). This holds even though a single granule cell receives a tiny fraction of

mossy fibre rates, and each field receives only a small random sample of mossy fibre

signals received as input to the rank as a whole, and it is independent of the shape of

the distribution of mossy fibre rates.

This is not in itself necessarily in fatal conflict with the traditional cerebellar learning

model, but has the alternative interpretation that granule cell firing conserves mossy

fibre rate information, coded in the output of a whole rank. This is therefore the

minimum resolution of granule cell information but with the great flexibility that the

same information is coded in any random sample of granule cell signals, at any

location. It does not matter which cells they are, so there is no need for patterns to be

repeated, or received at any particular set of synapses downstream. It is immaterial

which cells fire at what rates, or indeed what rates they are, provided any random

sample of rates is as if drawn from the same frequency distribution, with replacement.

The output of a rank can be likened by analogy to a pixel: it is only divisible into smaller

pieces of the same information.

525   **Figures 5 and 6**

5. **Discussion**

Neural networks are believed to carry out sophisticated computations. In the

influential traditional cerebellar learning model, and many artificial neural networks,

computations are implemented by learned synaptic changes that selectively adjust

synaptic transmission strength. We propose a rethink, taking the granular layer of the

cerebellar cortex, the input layer of the cerebellar network, as a physiologically-

detailed example. The proposals are quantified and tested in silico and empirically

testable. Some features are:

i) Information has geometry. Large cell groups talk to each other in a statistical rate

code. At any instant, a whole group codes the same information. Accordingly,

information has spatial dimensions, the dimensions of a code group.

ii) Granule cell code groups form long, thin strips, mirroring microzonal organisation of

the molecular layer, so internal code has a striped pattern. A strip/code group

functions as a single unit and spans the cerebellar network sagittally.

iii) Recoding exploits the central limit theorem. The mean of any random sample of

granule cell firing rates has a linear relationship with the mean of mossy fibre input

rates to the whole strip (and granule cell rates have a reduced range). The pattern of

active cells, and which cells fire at what rates, codes nothing.

iv) Accordingly, while firing of a whole code group is the minimum unit of information,

it is not the minimum unit of code, because inside those dimensions, the same

information is coded in any random sample of firing rates.

v) There is a computational effect, unaided, of the triad of cell morphologies, network

architecture and linear transmission. The cerebellum is highly adapted for this result.

No learning algorithm or learning is needed.

vi) There is a huge sacrifice of data resolution on entry to the cerebellum, counter to

the general expectation that more data should inform more powerful brain

computations.

vii) This is functional. Coded in this form, granule cell data can drive coherent firing of

large cell groups downstream.

viii) A statistical code may be a physiological strategy to mitigate variable performance

of single neurons and synapses, a long-known impediment to precise and replicable

biological information capture. It is a long-standing sometimes implicit notion that a

repeat of conditions that provide information to the brain generates a repeat of the

same brain activity. In fact it is at least not clear there is a faithful correlation, or

indeed that conditions are ever exactly repeated. Code that does not depend on high

fidelity may be a solution to working with unreliable information.

Microzones lie at right angles to parallel fibres, presenting a large target. As a result, a

Purkinje cell receives none of the same signals as almost all other Purkinje cells in the

same microzone. However, the form of the granule cell code means the same

information is received by all locations of sagittal plane, so that microzone-grouped

Purkinje cells receive the same information without receiving the same signals. An

estimated 350,000 parallel fibres pass through a Purkinje cell dendritic territory

(Harvey and Napper, 1988, 1991; Napper and Harvey, 1988a, b). If even only a small

fraction of parallel fibres is active (0.5%), and allowing for the fact that only one in two

make contact, and that a majority of synapses are severely depressed or silent (Ekerot

and Jörntell, 2001; Isope and Barbour, 2002), a single Purkinje cell still receives around

200 signals in any snapshot of parallel fibre activity. This provides a substrate to

control firing of Purkinje cells. Control is by a chain of rate codes. Purkinje cells linearly

code the strength of granule cell inputs (Park et al., 2012; Walter and Khodakhah,

2006, 2009) and behavioural metrics (Chen et al., 2016; Dugué et al., 2017; Herzfeld et

al., 2015; Hong et al., 2016; Jelitai et al., 2016). Learned changes of synaptic weights in

the form traditionally envisaged may not be needed. Also, it can explain how Purkinje

cell firing rates in a microzone are modulated in union.

A note on methodology. With our approach, the function of the computation is not

inherent in the mechanism. This breaks with a long tradition network modelling.

Generally, neural network models solve a problem it is proposed the network has. The

proposal is an assumption of the model; the solution is a computational mechanism;

the mechanism is the model. We don't propose a function, only a mechanism. We

work towards a function.

Computational anatomy provides a tool to habilitate linear signalling, historically not

popular with modellers because it makes unit computations less flexible.

METHODS

DERIVATION OF MOSSY FIBRE RATES

The number of mossy fibres that innervate a rank

As far as is known, fields receive a random sample mossy fibre input rates to a rank. To

simulate that, we need the size of the sampled population – the number of mossy fibre signals

received by a rank – and sample size, the number received by a field.

Despite the repeating 'crystalline' architecture of the cerebellar cortex, at local level there are

anatomical random variables. Variables are: i) the number of mossy fibres that innervate a

field, ii) the number of terminals they each end in, and iii) the number of those that are inside

field limits.

The number of mossy fibres afferent to a field has been estimated at 100 (Sultan and Heck,

2003). However, that number is a field-sized fraction of the general population. The actual

number is larger because mossy fibres end in a dispersed group (cluster) of terminals, and

most groups straddle field limits.

We work from an estimate of the total number of terminals per field, which is, in turn, derived

from convergent estimates of the number of granule cells per field. (Estimate 1) There are an

estimated 1.92×10^6 granule cells per µl (Harvey and Napper, 1988). The number of fields

which fit in that volume is $(1{,}000/150)^2 \times (1{,}000/200) = {\sim}222$, assuming a field depth of 150 µm

(Sultan and Heck, 2003). The number of granule cells in a field is, therefore $1{,}920{,}000/222 =$

8,649. (Estimate 2) An estimated 350,000 parallel fibres pass through a single Purkinje cell's

territory (Harvey and Napper, 1988). Parallel fibres run about 3 mm in each direction from the

granule cell soma (Brand et al., 1976; Mugnaini, 1983). The Purkinje cell dendritic arbour fills

the molecular layer vertically and averages ~200 µm wide. Therefore, $6{,}000/150 = 40$ fields fit

in a row that provides parallel fibres that pass through a Purkinje cell dendritic territory, giving

$350{,}000/40 = 8{,}750$ granule cells per field. Estimates vary of the number of granule cells which

extend a dendrite into a single glomerulus. 50 is mid-range. $(8{,}750/50) \times 4 = 700$ glomeruli per

field.

Granule cells have 3–5 dendrites, averaging 4, and receive input to each of them from a single

mossy fibre (so an average granule cell receives 4 inputs). Estimates of divergence of a single

mossy fibre terminal onto granule cells vary (Ritzau-Jost et al., 2014). It is unclear how much

this reflects biological diversity versus differences in opinion and methods. Here we use a mid-

range estimate of 1:50 (Gao et al., 2016; Jakab and Hámori, 1988). Assuming a mossy fibre

terminal receives a single granule cell dendrite from each of d granule cells, and given an

average of 4 dendrites per granule cell and z granule cells per field, the total number of mossy

fibre terminals in a field is $N = (z \times 4)/d = 700$ given $z = 8,750$ and $d = 50$. A convergent

estimate is provided by multiplying the per field number of mossy fibres by the average

number of terminals per cluster.

Cluster size varies. We take the range as 4–12 terminals per cluster (Sultan and Heck, 2003).

We assume the number varies at random and cluster sizes occur with equal probability. A

maximum of 7 fits in a field (because a field shares the dimensions of the area that encloses a

mean cluster). Fields are not anatomically bounded, so many clusters straddle field limits. To

reflect that, we assume there is a 1/7 probability that a 7-terminal cluster (or larger) has 7

terminals inside a field and also the same probability of any other number. A 6-terminal cluster

has a 2/7 chance that all 6 terminals are inside and a 1/7 chance of any other number, and so

on.

Table 1 shows the derivation of the expected (i.e., predicted by probability) ratios of terminals

contributed to a field by each cluster size (R1), not counting terminals that lie outside when a

cluster straddles field boundaries; and the ratios of mossy fibres contributing $n = 1$–7 terminals

(R2), regardless of cluster size. R1 ratios are given by the sum of numbers in each row. So, for

example, the ratio of cells that end in 8 terminals to cells that end in 5 is 28:25. R2 ratios are

given by the number of appearances of each number in the grid – so the expected ratio of cells

which contribute 7 terminals to the number which contribute 4 is 6:12. The ratios effectively

give the distributed probability of possible states. For example, there is a probability of 10/(6 +

8 + 10 + 12 + (3 x 9)) = 0.159 that a mossy fibre contributes 5 terminals to a field (regardless of

cluster size). Similarly, there is a probability of ((6 x 28)/((6 x 28) + 27 + 25 + 22))/7 = 0.099 that

a given cell contributes 7 terminals to a field.

Table 1. Derivation of expected ratios

size	number of terminals inside							R1
12	7	6	5	4	3	2	1	**28**
11	7	6	5	4	3	2	1	**28**
10	7	6	5	4	3	2	1	**28**
9	7	6	5	4	3	2	1	**28**
8	7	6	5	4	3	2	1	**28**
7	7	6	5	4	3	2	1	**28**
6	6	6	5	4	3	2	1	**27**
5	5	5	5	4	3	2	1	**25**
4	4	4	4	4	3	2	1	**22**
R2	6	8	10	12	9	9	9	

If R1 ratios are denoted by a series $x_4, x_5 \dots x_{12}$, and a given element in the series is x_j, the

fraction of terminals provided by a given cluster size is $x_j/(x_4 + x_5 + \cdots + x_{12})$, and this

fraction is multiplied by N to give the number of terminals. The number of terminals is divided

by the expected mean number of terminals that fall inside field limits (for that cluster size) to

give the expected number of mossy fibres that provide them. The number of j-terminal mossy

fibres that innervate a field is therefore given by

$$N\left[\left.x_j\middle/\textstyle\sum_{i=1}^{n} x_i\right.\right]\middle/\left[x_j/7\right]$$

654   The results are shown in Table 2. The equal number of mossy fibres (mfs) in each column is

655   what we would expect, because we made it an assumption that all sizes occur with equal

656   probability, and mf values are expected values – i.e., in proportions predicted by their

657 probability. The mf total is the expected number of mossy fibres afferent to a field (which

658 contribute at least one terminal).

Table 2. The expected number of mossy fibres which innervate a field

cluster size	12	11	10	9	8	7	6	5	4	total
terminals	81	81	81	81	81	81	78.1	72.3	63.6	700
mfs	20.25	20.25	20.25	20.25	20.25	20.25	20.25	20.25	20.25	182.25

659 (table label line 659)

660 R2 ratios – the relative incidence of afferent mossy fibres that contribute 7 terminals to a field,

661 6 terminals, 5 and so on, down to 1 – and the expected number of mossy fibres that contribute

662 them, are shown in Table 3.

663 **Table 3. The number of mossy fibres which contribute n = 1–7 terminals to a field**

n	7	6	5	4	3	2	1	total
R2	6	8	10	12	9	9	9	63
R3	42	48	50	48	27	18	9	
mfs	17.34	23.15	28.92	34.72	26.04	26.04	26.04	182.25
%age	9.52	12.69	15.87	19.05	14.29	14.29	14.29	

R3 is n x R2, giving the ratio of terminals provided by mossy fibres with n terminals inside a

field. %age is given by (R2/63) x 100. this is the percentage of mossy fibres with n terminals

inside a field.

From here, the number of mossy fibres that innervate a rank is given by 182.25 x 100 divided

by the mean number of clusters per mossy fibre per rank. If a rank receives, on average, five

clusters per mossy fibre, ~3,645 mossy fibres innervate a rank. This is the estimate used in the

simulation unless stated otherwise.

Assumptions. Cluster sizes occur with equal probability. The probabilities of each outcome

when a cluster straddles field boundaries (how many terminals are inside and how many

outside) are as stated. A single mossy fibre ends in an average of 5 terminal clusters per rank.

Mossy fibre input rates to a field

Terminal clustering means a field receives multiple copies of most signals. The proportions of

mossy fibres afferent to a simulated field that contribute each possible number of terminals

(1–7) are generated in simulated steps. In the first step, the simulation randomly generates,

for each field, the cluster size ratio. This means, the relative proportions of afferent mossy

fibres (which contribute at least 1 terminal to a field) that end in 4 terminals, 5, 6 and so on up

to 12, assuming they occur with equal probability, disregarding how many terminals fall inside

and outside field boundaries. In step two, the ratio of cluster sizes is weighted to reflect the

greater probability that large clusters contribute more terminals to a field. In step 3, ratios are

converted to the fraction of terminals contributed by each cluster size. When multiplied by the

total number of terminals, this gives the number of terminals contributed by each cluster size.

In step four, the values obtained in step three are divided up between cells that contribute a

single terminal, 2 terminals, 3 and so on, to reflect straddling of field limits. The proportions

give the number of mossy fibres afferent to a simulated field that contribute a single terminal,

two, three and so on up to seven, the maximum.

There is no field-level effect of mossy fibre terminal branching. Terminal clusters arising from

the same mossy fibre are separated by a minimum distance, so that a field cannot contain

terminals of more than one cluster per mossy fibre.

Terminal clustering means that fields contain a randomly variable number of copies of most

signals. In the simulation, copies are added after sampling of mossy fibre rates to obtain rates

received by a field. An alternative order would be to add copies to the sampled population and

then sample that. We follow anatomy: the number of *rates* received by a field is equal to the

number of afferent mossy fibres; the number of *signals* contains duplicate rates.

LINEAR TRANSMISSION OF MOSSY FIBRES TO GOLGI CELLS

It is part of our proposal that Golgi cell basal dendritic signals linearly reflect mossy fibre rates,

so that mossy fibre rate information is conserved proportionally in Golgi cell firing rates. Golgi

cells are large interneurons whose cell bodies and basal dendrites lie in the granular layer.

Mossy fibres directly contact Golgi cell basal dendrites. Golgi cell firing frequency increases

linearly with the amplitude of depolarising current (Dieudonné, 1998 p.845). 'Golgi cells can

follow peripheral signals in a continuous fashion, modulating their frequency with the intensity

of the stimulus [citing (Edgley and Lidierth, 1987; Miles et al., 1980)]' (Galliano et al., 2010). As

a result, during movement, the 'output of Golgi cells varies smoothly as a function of

input....They are capable of modulating their output to reflect their input with little noise or

signal distortion' (Moore and Blazis, 1989). 'Sensory-evoked Golgi-cell inhibition scales

proportionally with the level of mossy fiber excitatory synaptic input', such that inhibition

reliably conveys mossy fibre rate information (Duguid et al., 2015). On this evidence: mossy

fibre rate information is conserved proportionally in Golgi cell firing rates.

Golgi cells extend their basal dendrites into glomeruli (Eccles et al., 1964; Hámori and

Szentágothai, 1966). Contact on them by mossy fibres is multi-synaptic (Palay and Chan-Palay,

1974), contributing to a reliable, rapid (submillisecond) response. The fast rise time and weak

distance dependence of the amplitude and timing of Golgi cell EPSCs evoked by mossy fibre

signals, suggests that basal dendrites introduce relatively little filtering (Cesana et al., 2010;

Kanichay and Silver, 2008), as expected for large-diameter dendrites. We note that the pause

in Golgi cell firing following a discrete stimulus under anaesthesia (Duguid et al., 2015;

Holtzman et al., 2006; Vos et al., 1999) disappears from Golgi cell recordings during

locomotion (Edgley and Lidierth, 1987).

During behaviour in freely-moving animals, mossy fibre activity is dense: a high fraction of

mossy fibres are active (Hartmann and Bower, 2001). Both mossy fibres (Arenz et al., 2008;

Giovannucci et al., 2017; van Kan et al., 1993) and Golgi cells (Edgley and Lidierth, 1987) have

been reported to fire with a sustained, time-varying signature. In the behaving animal, Golgi

cell basal dendritic membrane potential is a continuous variable under modulation by

sustained inputs, we submit. As charge transfer to the soma is passive, polarisation of the

soma is likewise sustained, under modulation by dendritic states.

It is worth noting that very high mossy fibre rates can be generated in experimental conditions.

However, the typical physiological range is 50–300 Hz (van Kan et al., 1993). In the simulations,

we use the physiological rates, within stated constraints on the shape of the distribution.

Other control of Golgi cells? Do Golgi cells receive a significant influence of input from other sources? The evidence is incomplete but currently suggests other influence is absent or weak. Contrary to early reports, neither Purkinje cells nor climbing fibres contact Golgi cells (Pietrajtis and Dieudonné, 2013, who give references), and only a modest minority of inhibitory inputs to Golgi cells (if any) are from molecular layer interneurons (Eyre and Nusser, 2016), which generate weak synaptic currents (Dumoulin et al., 2001), consistent with either extremely weak or wholly absent innervation (Hull and Regehr, 2012). There is conflicting evidence whether Golgi cells make inhibitory synaptic contact on each other (Hull and Regehr, 2012; Vervaeke et al., 2010). Our simulation does not include an effect on Golgi cell firing by other sources of input.

LINEAR TRANSMISSION OF GOLGI CELLS TO GRANULE CELLS

'In the adult rat cerebellum, each granule cell dendrite receives 2.6 ± 0.55 synaptic contacts from Golgi axon terminals' (Rossi and Hamann, 1998) citing (Jakab and Hámori, 1988).[1] However, the large majority of inhibition (98%) is by spillover (Duguid et al., 2012), where neurotransmitter released into the synaptic cleft spills out into the glomerular space. Golgi cells release GABA, an inhibitory neurotransmitter. This is detected by high-affinity GABA$_A$ receptors located perisynaptically and extrasynaptically on granule cells (Brickley et al., 1996; Nusser et al., 1998). Even synaptically-received signals have most of their effect (i.e., the large majority of charge transfer is mediated) via spillover.[2]

---

[1] Data are sparse. The original data are an estimate derived from visual inspection of slices thirty years ago so far without later corroboration, except from Rossi and Hamann 1998, who derive a lower estimate but propose the disparity is because their estimate is too low.
[2] Because of the long duration of IPSCs generated by spillover (recorded in slices), the total charge carried is three times that of IPSCs generated by directly connected terminals. Rossi, D.J., and Hamann, M. (1998). Spillover-mediated transmission at inhibitory synapses promoted by high affinity alpha6 subunit GABA(A) receptors and glomerular geometry. Neuron 20, 783-795.

Golgi cells fire spontaneously (Ruigrok et al., 2011) at a time-varying rate in the behaving

animal, so that downstream, a glomerulus receives continuous convergent input. As a result,

there is a sustained build-up of glomerular GABA during behaviour (Rossi et al., 2003) at an

adjustable concentration controlled by the mean of afferent Golgi cell firing rates (Mapelli et

al., 2014). A glomerulus randomly samples Golgi cells afferent to a field. Golgi cells provide the

only inhibitory input to granule cells.

Signalling by spillover is sometimes assumed to be ambient and slow. However, the action of

glomerular GABA spillover has a fast phasic component – not as fast as synaptic transmission

(~1 ms) but with a rise time of only a few milliseconds (Mapelli et al., 2014). Unlike the spiky

appearance of synaptically-induced inhibitory postsynaptic currents, spillover (Duguid et al.,

2015) generates a sustained outward current. The response time to multiple inputs is likely

faster than the experimentally isolated response to a discrete signal (because the combined

rate of GABA release is higher).

Golgi cell oscillations. Golgi cells fire autonomously. Under anaesthesia, firing falls into a slow

oscillating pattern (Ros et al., 2009). Under excitatory synaptic input, however, this disappears

(Vervaeke et al., 2010). Anaesthesia 'has a strong influence on spontaneous activity of Golgi

cells' (Pietrajtis and Dieudonné, 2013). Discharge at low rates with irregular timing under

anaesthesia (Ruigrok et al., 2011) is replaced by higher rates with more regular timing without

anaesthesia (Edgley and Lidierth, 1987; Miles et al., 1980; Rasmussen et al., 2014). (This

paragraph is a cautionary note about evidence obtained under anaesthesia, which has been

used to argue that oscillations provide a form of signalling.)

LINEAR TRANSMISSION OF MOSSY FIBRES TO GRANULE CELLS

There is significant evidence that granule cell firing rates are linearly related to input rates, but

also evidence thought to be unfavourable to that conclusion. We take them in turn.

The short and equal length of granule cell dendrites would suggest light and equal filtering.

The mossy fibre-granule cell connection has a range of adaptations which support high-fidelity

transmission of high frequency signals across a wide bandwidth (Delvendahl and Hallermann,

2016). Faithful transmission of rate information is reported (Rancz et al., 2007; Ritzau-Jost et

al., 2014). Vesicle release and replenishment are fast (Sargent et al., 2005; Saviane and Silver,

2006) and postsynaptic AMPA receptors operate in their linear range (Sargent et al., 2005),

where they are resistant to desensitisation (DiGregorio et al., 2007). Multiple contacts are

made by a mossy fibre with each granule cell (Jakab and Hámori, 1988). Conversion of

depolarising somatic charge into granule cell firing is a simple function, such that granule cells

'have a relatively linear and uncomplicated conversion of depolarisation level to spike rate'

(Bengtsson and Jörntell, 2009 p.2393, citing Jorntell and Ekerot 2006 and D'Angelo et al 1998).

Glutamate spillover enhances precision and reliability of transmission (DiGregorio et al., 2002;

Sargent et al., 2005). During behaviour, at sustained physiological mossy fibre rates, there may

be a build-up of intraglomerular glutamate, increasing the relative influence of spillover in a

balance with synaptic transmission, assisted by short-term synaptic depression. In these

conditions, spillover may dominate, as it dominates inhibitory transmission (Duguid et al.,

2012; Mapelli et al., 2014). Spillover from multiple release sites (200-400 at a single mossy

fibre terminal) is plausibly sufficient to mitigate variability of vesicular size, release probability,

and the number of release points at synapses made on any individual cell, increasing fidelity

and equality of excitation of granule cells.

Against that, there are reported to be heterogeneous synaptic weights. Different strength and

short-term dynamics of the mossy fibre to granule cell connection have been reported in

vestibular slices, with GABA$_A$ receptors blocked. The amplitude of the response, measured at

the soma, is reported to be statistically correlated to the source of the mossy fibre that is

stimulated (Chabrol et al., 2015), suggesting that the strength of the connection depends on

the source.

However, the presence of weights does not necessarily mean that the function is to modulate

firing rates. Rather, it is suggested that the function may temporally code the source of inputs

because different combinations have different response onset times. An alternative, which

they also mention, is that some inputs may be in a supporting role, reminiscent of 'driver' and

'modulatory' signals in the thalamus and cortex (Sherman and Guillery, 2011). Feasibly, signals

in a supporting role must be present for the postsynaptic granule cell to fire, but do not affect
the granule cell firing rate.

Grants

811   Funding: This work was supported by the Leverhulme Trust [grant number ECF-2022-

812   079] (Mike Gilbert); the Swedish Research Council [grant number 2020-01468]; the

813   Per-Eric och Ulla Schybergs Foundation [grant number 42630]; and the Crafoord

814   Foundation [grant numbers 20180704, 20200729, 20220776, & 20230655] (Anders

815   Rasmussen).

Acknowledgement

Author contributions

820   M.G. Researched, conceived and developed the ideas, performed the analysis, wrote

821   the code, prepared the figures and wrote the manuscript. A.R. provided expertise and

822   feedback, and reviewed and provided editorial comments on the manuscript.

Declaration of interests

The authors declare no competing interests.

REFERENCES

Albus, J.S. (1971). A Theory of Cerebellar Function. Mathematical Biosciences 10, 25-61.
Albus, J.S. (1989). The Marr and Albus theories of the cerebellum-two early models of
associative memory. In Digest of Papers COMPCON Spring 89 Thirty-Fourth IEEE Computer
Society International Conference: Intellectual Leverage, pp. 577-582.
Alvina, K., Walter, J.T., Kohn, A., Ellis-Davies, G., and Khodakhah, K. (2008). Questioning the
role of rebound firing in the cerebellum. Nature neuroscience 11, 1256-1258.
Arenz, A., Silver, R.A., Schaefer, A.T., and Margrie, T.W. (2008). The contribution of single
synapses to sensory representation in vivo. Science (New York, NY) 321, 977-980.
Barmack, N.H., and Yakhnitsa, V. (2008). Functions of interneurons in mouse cerebellum. The
Journal of neuroscience : the official journal of the Society for Neuroscience 28, 1140-1152.
Bengtsson, F., and Jörntell, H. (2009). Sensory transmission in cerebellar granule cells relies on
similarly coded mossy fiber inputs. Proceedings of the National Academy of Sciences of the
United States of America 106, 2389-2394.
Billings, G., Piasini, E., Lorincz, A., Nusser, Z., and Silver, R.A. (2014). Network structure within
the cerebellar input layer enables lossless sparse encoding. Neuron 83, 960-974.
Brand, S., Dahl, A.L., and Mugnaini, E. (1976). The length of parallel fibers in the cat cerebellar
cortex. An experimental light and electron microscopic study. Experimental brain research 26,
39-58.
Brickley, S.G., Cull-Candy, S.G., and Farrant, M. (1996). Development of a tonic form of synaptic
inhibition in rat cerebellar granule cells resulting from persistent activation of GABAA
receptors. The Journal of physiology 497 (Pt 3), 753-759.
Brunel, N., Hakim, V., Isope, P., Nadal, J.P., and Barbour, B. (2004). Optimal information
storage and the distribution of synaptic weights: perceptron versus Purkinje cell. Neuron 43,
745-757.
Cesana, E., Dieudonné, S., Isope, P., Bidoret, C., D'Angelo, E., and Forti, L. (2010). Novel granule
cell-Golgi cell excitatory input in the cerebellar granular layer. In FENS Abstr.
Cesana, E., Pietrajtis, K., Bidoret, C., Isope, P., D'Angelo, E., Dieudonne, S., and Forti, L. (2013).
Granule cell ascending axon excitatory synapses onto Golgi cells implement a potent feedback
circuit in the cerebellar granular layer. The Journal of neuroscience : the official journal of the
Society for Neuroscience 33, 12430-12446.
Chabrol, F.P., Arenz, A., Wiechert, M.T., Margrie, T.W., and DiGregorio, D.A. (2015). Synaptic
diversity enables temporal coding of coincident multisensory inputs in single neurons. Nature
neuroscience 18, 718-727.
Chadderton, P., Margrie, T.W., and Hausser, M. (2004). Integration of quanta in cerebellar
granule cells during sensory processing. Nature 428, 856-860.
Chen, S., Augustine, G.J., and Chadderton, P. (2016). The cerebellum linearly encodes whisker
position during voluntary movement. eLife 5, e10509-e10509.
D'Angelo, E., Solinas, S., Mapelli, J., Gandolfi, D., Mapelli, L., and Prestori, F. (2013). The
cerebellar Golgi cell and spatiotemporal organization of granular layer activity. Frontiers in
neural circuits 7, 93.
Dean, P., Porrill, J., Ekerot, C.F., and Jörntell, H. (2010). The cerebellar microcircuit as an
adaptive filter: experimental and computational evidence. Nature reviews Neuroscience 11,
30-43.

Delvendahl, I., and Hallermann, S. (2016). The Cerebellar Mossy Fiber Synapse as a Model for
High-Frequency Transmission in the Mammalian CNS. Trends Neurosci 39, 722-737.
Dieudonné, S. (1998). Submillisecond kinetics and low efficacy of parallel fibre-Golgi cell
synaptic currents in the rat cerebellum. The Journal of physiology 510 (Pt 3), 845-866.
DiGregorio, D.A., Nusser, Z., and Silver, R.A. (2002). Spillover of glutamate onto synaptic AMPA
receptors enhances fast transmission at a cerebellar synapse. Neuron 35, 521-533.
DiGregorio, D.A., Rothman, J.S., Nielsen, T.A., and Silver, R.A. (2007). Desensitization properties
of AMPA receptors at the cerebellar mossy fiber granule cell synapse. The Journal of
neuroscience : the official journal of the Society for Neuroscience 27, 8344-8357.
Dugué, G.P., Tihy, M., Gourévitch, B., and Léna, C. (2017). Cerebellar re-encoding of self-
generated head movements. eLife 6.
Duguid, I., Branco, T., Chadderton, P., Arlt, C., Powell, K., and Hausser, M. (2015). Control of
cerebellar granule cell output by sensory-evoked Golgi cell inhibition. Proceedings of the
National Academy of Sciences of the United States of America 112, 13099-13104.
Duguid, I., Branco, T., London, M., Chadderton, P., and Hausser, M. (2012). Tonic inhibition
enhances fidelity of sensory information transmission in the cerebellar cortex. The Journal of
neuroscience : the official journal of the Society for Neuroscience 32, 11132-11143.
Dumoulin, A., Triller, A., and Dieudonné, S. (2001). IPSC kinetics at identified GABAergic and
mixed GABAergic and glycinergic synapses onto cerebellar Golgi cells. The Journal of
neuroscience : the official journal of the Society for Neuroscience 21, 6045-6057.
Eccles, J.C., Ito, M., and Szentágothai, J. (1967). The cerebellum as a neuronal machine (Berlin,
New York etc.: Springer-Verlag).
Eccles, J.C., Llinás, R., and Sasaki, K. (1964). Golgi cell inhibition in the cerebellar cortex. Nature
204, 1265-1266.
Edgley, S.A., and Lidierth, M. (1987). The discharges of cerebellar Golgi cells during locomotion
in the cat. The Journal of physiology 392, 315-332.
Ekerot, C.F., and Jörntell, H. (2001). Parallel fibre receptive fields of Purkinje cells and
interneurons are climbing fibre-specific. The European journal of neuroscience 13, 1303-1310.
Eyre, M.D., and Nusser, Z. (2016). Only a Minority of the Inhibitory Inputs to Cerebellar Golgi
Cells Originates from Local GABAergic Cells. eNeuro 3.
Fujita, M. (1982). Adaptive filter model of the cerebellum. Biological cybernetics 45, 195-206.
Galliano, E., Mazzarello, P., and D'Angelo, E. (2010). Discovery and rediscoveries of Golgi cells.
The Journal of physiology 588, 3639-3655.
Gao, Z., Proietti-Onori, M., Lin, Z., Ten Brinke, M.M., Boele, H.J., Potters, J.W., Ruigrok, T.J.,
Hoebeek, F.E., and De Zeeuw, C.I. (2016). Excitatory Cerebellar Nucleocortical Circuit Provides
Internal Amplification during Associative Conditioning. Neuron 89, 645-657.
Garwicz, M., Ekerot, C.F., and Jorntell, H. (1998). Organizational principles of cerebellar
neuronal circuitry. News in Physiological Sciences 13, 28-34.
Gilbert, M., and Miall, R.C. (2021a). How and Why the Cerebellum Recodes Input Signals: An
Alternative to Machine Learning. Neuroscientist DOI: 10.1177/1073858420986795.
Gilbert, M., and Rasmussen, A. (2024). Gap Junctions May Have A Computational Function In
The Cerebellum: A Hypothesis. The Cerebellum.
Giovannucci, A., Badura, A., Deverett, B., Najafi, F., Pereira, T.D., Gao, Z., Ozden, I., Kloth, A.D.,
Pnevmatikakis, E., Paninski, L., *et al.* (2017). Cerebellar granule cells acquire a widespread
predictive feedback signal during motor learning. Nature neuroscience.

Hámori, J. (1992). Anatomy and neurochemical anatomy of the cerebellum. In Cerebellar
degenerations : clinical neurobiology, A. Plaitakis, ed. (Kluwer Academic).
Hámori, J., and Szentágothai, J. (1966). Participation of Golgi neuron processes in the
cerebellar glomeruli: an electron microscope study. Experimental brain research 2, 35-48.
Hartmann, M.J., and Bower, J.M. (2001). Tactile responses in the granule cell layer of
cerebellar folium crus IIa of freely behaving rats. The Journal of neuroscience : the official
journal of the Society for Neuroscience 21, 3549-3563.
Harvey, R.J., and Napper, R.M. (1988). Quantitative study of granule and Purkinje cells in the
cerebellar cortex of the rat. The Journal of comparative neurology 274, 151-157.
Harvey, R.J., and Napper, R.M. (1991). Quantitative studies on the mammalian cerebellum.
Progress in neurobiology 36, 437-463.
Herzfeld, D.J., Kojima, Y., Soetedjo, R., and Shadmehr, R. (2015). Encoding of action by the
Purkinje cells of the cerebellum. Nature 526, 439-442.
Holtzman, T., Rajapaksa, T., Mostofi, A., and Edgley, S.A. (2006). Different responses of rat
cerebellar Purkinje cells and Golgi cells evoked by widespread convergent sensory inputs. The
Journal of physiology 574, 491-507.
Hong, S., Negrello, M., Junker, M., Smilgin, A., Thier, P., and De Schutter, E. (2016). Multiplexed
coding by cerebellar Purkinje neurons. eLife 5.
Hull, C., and Regehr, W.G. (2012). Identification of an inhibitory circuit that regulates cerebellar
Golgi cell activity. Neuron 73, 149-158.
Isope, P., and Barbour, B. (2002). Properties of unitary granule cell-->Purkinje cell synapses in
adult rat cerebellar slices. The Journal of neuroscience : the official journal of the Society for
Neuroscience 22, 9668-9678.
Jakab, R.L., and Hámori, J. (1988). Quantitative morphology and synaptology of cerebellar
glomeruli in the rat. Anatomy and embryology 179, 81-88.
Jelitai, M., Puggioni, P., Ishikawa, T., Rinaldi, A., and Duguid, I. (2016). Dendritic excitation-
inhibition balance shapes cerebellar output during motor behaviour. Nature communications
7, 13722.
Jörntell, H., and Ekerot, C.F. (2006). Properties of somatosensory synaptic integration in
cerebellar granule cells in vivo. The Journal of neuroscience : the official journal of the Society
for Neuroscience 26, 11786-11797.
Kanichay, R.T., and Silver, R.A. (2008). Synaptic and cellular properties of the feedforward
inhibitory circuit within the input layer of the cerebellar cortex. The Journal of neuroscience :
the official journal of the Society for Neuroscience 28, 8955-8967.
Lange, W. (1974). Regional differences in the distribution of golgi cells in the cerebellar cortex
of man and some other mammals. Cell and tissue research 153, 219-226.
Mapelli, L., Solinas, S., and D'Angelo, E. (2014). Integration and regulation of glomerular
inhibition in the cerebellar granular layer circuit. Frontiers in cellular neuroscience 8, 55.
Miles, F.A., Fuller, J.H., Braitman, D.J., and Dow, B.M. (1980). Long-term adaptive changes in
primate vestibuloocular reflex. III. Electrophysiological observations in flocculus of normal
monkeys. Journal of neurophysiology 43, 1437-1476.
Mitchell, S.J., and Silver, R.A. (2000a). GABA spillover from single inhibitory axons suppresses
low-frequency excitatory transmission at the cerebellar glomerulus. The Journal of
neuroscience : the official journal of the Society for Neuroscience 20, 8651-8658.

Mitchell, S.J., and Silver, R.A. (2000b). Glutamate spillover suppresses inhibition by activating
presynaptic mGluRs. Nature 404, 498-502.
Moore, J.W., and Blazis, D.E.J. (1989). 11 - Simulation of a Classically Conditioned Response: A
Cerebellar Neural Network Implementation of the Sutton–Barto–Desmond Model. In Neural
Models of Plasticity, J.H. Byrne, and W.O. Berry, eds. (Academic Press), pp. 187-207.
Mugnaini, E. (1983). The length of cerebellar parallel fibers in chicken and rhesus monkey. The
Journal of comparative neurology 220, 7-15.
Napper, R.M., and Harvey, R.J. (1988a). Number of parallel fiber synapses on an individual
Purkinje cell in the cerebellum of the rat. The Journal of comparative neurology 274, 168-177.
Napper, R.M., and Harvey, R.J. (1988b). Quantitative study of the Purkinje cell dendritic spines
in the rat cerebellum. The Journal of comparative neurology 274, 158-167.
Nusser, Z., Sieghart, W., and Somogyi, P. (1998). Segregation of different GABAA receptors to
synaptic and extrasynaptic membranes of cerebellar granule cells. The Journal of neuroscience
: the official journal of the Society for Neuroscience 18, 1693-1703.
Oscarsson, O. (1979). Functional units of the cerebellum - sagittal zones and microzones.
Trends in Neurosciences 2, 143-145.
Ozden, I., Sullivan, M.R., Lee, H.M., and Wang, S.S. (2009). Reliable coding emerges from
coactivation of climbing fibers in microbands of cerebellar Purkinje neurons. The Journal of
neuroscience : the official journal of the Society for Neuroscience 29, 10463-10473.
Palay, S.L., and Chan-Palay, V. (1974). Cerebellar cortex : cytology and organization (Berlin:
Springer).
Park, S.M., Tara, E., and Khodakhah, K. (2012). Efficient generation of reciprocal signals by
inhibition. Journal of neurophysiology 107, 2453-2462.
Pietrajtis, K., and Dieudonné, S. (2013). Golgi Neurons. In Handbook of the Cerebellum and
Cerebellar Disorders, M. Manto, J.D. Schmahmann, F. Rossi, D.L. Gruol, and N. Koibuchi, eds.
(Dordrecht: Springer Netherlands), pp. 829-852.
Rancz, E.A., Ishikawa, T., Duguid, I., Chadderton, P., Mahon, S., and Hausser, M. (2007). High-
fidelity transmission of sensory information by single cerebellar mossy fibre boutons. Nature
450, 1245-1248.
Rasmussen, A., Zucca, R., Jirenhed, D.A., Johansson, F., Ortenblad, C., Svensson, P., and
Hesslow, G. (2014). Golgi cell activity during eyeblink conditioning in decerebrate ferrets.
Cerebellum (London, England) 13, 42-45.
Raymond, J.L., and Medina, J.F. (2018). Computational Principles of Supervised Learning in the
Cerebellum. Annual review of neuroscience 41, 233-253.
Ritzau-Jost, A., Delvendahl, I., Rings, A., Byczkowicz, N., Harada, H., Shigemoto, R., Hirrlinger, J.,
Eilers, J., and Hallermann, S. (2014). Ultrafast action potentials mediate kilohertz signaling at a
central synapse. Neuron 84, 152-163.
Ros, H., Sachdev, R.N., Yu, Y., Sestan, N., and McCormick, D.A. (2009). Neocortical networks
entrain neuronal circuits in cerebellar cortex. The Journal of neuroscience : the official journal
of the Society for Neuroscience 29, 10309-10320.
Rossi, D.J., and Hamann, M. (1998). Spillover-mediated transmission at inhibitory synapses
promoted by high affinity alpha6 subunit GABA(A) receptors and glomerular geometry. Neuron
20, 783-795.
Rossi, D.J., Hamann, M., and Attwell, D. (2003). Multiple modes of GABAergic inhibition of rat
cerebellar granule cells. The Journal of physiology 548, 97-110.

Ruigrok, T.J., Hensbroek, R.A., and Simpson, J.I. (2011). Spontaneous activity signatures of
morphologically identified interneurons in the vestibulocerebellum. The Journal of
neuroscience : the official journal of the Society for Neuroscience 31, 712-724.
Sargent, P.B., Saviane, C., Nielsen, T.A., DiGregorio, D.A., and Silver, R.A. (2005). Rapid vesicular
release, quantal variability, and spillover contribute to the precision and reliability of
transmission at a glomerular synapse. The Journal of neuroscience : the official journal of the
Society for Neuroscience 25, 8173-8187.
Saviane, C., and Silver, R.A. (2006). Fast vesicle reloading and a large pool sustain high
bandwidth transmission at a central synapse. Nature 439, 983-987.
Schonewille, M., Khosrovani, S., Winkelman, B.H., Hoebeek, F.E., De Jeu, M.T., Larsen, I.M.,
Van der Burg, J., Schmolesky, M.T., Frens, M.A., and De Zeeuw, C.I. (2006). Purkinje cells in
awake behaving animals operate at the upstate membrane potential. Nature neuroscience 9,
459-461; author reply 461.
Sherman, S.M., and Guillery, R.W. (2011). Distinct functions for direct and transthalamic
corticocortical connections. Journal of neurophysiology 106, 1068-1077.
Shinoda, Y., and Sugihara, I. (2013). Axonal Trajectories of Single Climbing and Mossy Fiber
Neurons in the Cerebellar Cortex and Nucleus. In Handbook of the Cerebellum and Cerebellar
Disorders, S.J.D. Manto M., Rossi F., Gruol D.L., Koibuchi N., ed. (Springer, Dordrecht).
Shinoda, Y., Sugihara, I., Wu, H.S., and Sugiuchi, Y. (2000). The entire trajectory of single
climbing and mossy fibers in the cerebellar nuclei and cortex. Progress in brain research 124,
173-186.
Sultan, F., and Heck, D. (2003). Detection of sequences in the cerebellar cortex: numerical
estimate of the possible number of tidal-wave inducing sequences represented. Journal of
physiology, Paris 97, 591-600.
Turecek, J., Jackman, S.L., and Regehr, W.G. (2016). Synaptic Specializations Support
Frequency-Independent Purkinje Cell Output from the Cerebellar Cortex. Cell reports 17, 12.
Turecek, J., Jackman, S.L., and Regehr, W.G. (2017). Synaptotagmin 7 confers frequency
invariance onto specialized depressing synapses. Nature 551, 503.
van Kan, P.L., Gibson, A.R., and Houk, J.C. (1993). Movement-related inputs to intermediate
cerebellum of the monkey. Journal of neurophysiology 69, 74-94.
Vervaeke, K., Lorincz, A., Gleeson, P., Farinella, M., Nusser, Z., and Silver, R.A. (2010). Rapid
desynchronization of an electrically coupled interneuron network with sparse excitatory
synaptic input. Neuron 67, 435-451.
Vos, B.P., Volny-Luraghi, A., and De Schutter, E. (1999). Cerebellar Golgi cells in the rat:
receptive fields and timing of responses to facial stimulation. The European journal of
neuroscience 11, 2621-2634.
Walter, J.T., and Khodakhah, K. (2006). The linear computational algorithm of cerebellar
Purkinje cells. The Journal of neuroscience : the official journal of the Society for Neuroscience
26, 12861-12872.
Walter, J.T., and Khodakhah, K. (2009). The advantages of linear information processing for
cerebellar computation. Proceedings of the National Academy of Sciences of the United States
of America 106, 4471-4476.
Wu, H.S., Sugihara, I., and Shinoda, Y. (1999). Projection patterns of single mossy fibers
originating from the lateral reticular nucleus in the rat cerebellar cortex and nuclei. The Journal
of comparative neurology 411, 97-118.

A

B

C

Molecular
layer

Granular
layer

Mossy
fibre

Cerebellar y-axis [μm]

Cerebellar x-axis [μm]

D

Fig 1. Mossy fibre termination. A Sagittal section of the rat cerebellum (obtained by
multi-photon microscopy, courtesy Tom Deerinck). The outer layer of the cerebellar
cortex, the molecular layer, is stained green and the inner layer, the granular layer, is
blue. **B** Output of a procedure used to group mossy fibre terminals into clusters (Sultan
and Heck, 2003 fig 6), reproduced with permission. All terminals belong to one mossy
fibre, which terminates in two strips. Clusters are shown in different shades of grey. **C**
Schematic of an area of the folded cerebellar cortex. The strip inside the red dotted
line is the width and depth of a microzone but only part of its length. Termination of
mossy fibre signals is not at a point but at several randomly spaced, sagittally-aligned
locations. Yellow zones are terminal locations of a single mossy fibre. Size of locations
varies depending partly on the number of terminals in each cluster. Pink zones are
terminal locations of a second mossy fibre. In reality, zones overlap extensively and
terminals are intermingled. A mossy fibre mediolateral axon may give rise to several
sagittal branches, of which here only one is shown. The zone enclosed by a blue line
has the dimensions of a 'field'. A field is a nominal division of the granular layer with
the average dimensions of a mossy fibre terminal cluster – 200 μm x 150 μm – which
we use to quantify granular layer computations. A strip of the granular layer with the
dimensions of a sagittal row of 75–100 fields is a 'rank'. Viewed from the cerebellar
surface, a rank has the dimensions of a mid-range estimate of a microzone, but its long
sides are less cleanly defined. **D** Section of a microzone viewed obliquely from the
cerebellar surface, to illustrate the size and random intermingling of mossy fibre
terminals. The molecular layer is represented by squares of about the dimensions of a
Purkinje cell territory. The wire frame boxes are an exploded view of the orange fields.
The right box shows randomly distributed terminals of 5 mossy fibres (each a different
colour). We estimate that ~180 mossy fibres contribute terminals to a field, and a field
contains a total of around 700 terminals. Proportions of terminal size to field size are
preserved. The left box shows terminals of 20 mossy fibres.

1076

1077

Fig 2. Golgi cell ensemble conversion. Column 1 We randomly generated (3,645) values representing mossy fibre rates with a uniform, normal, twin-peaked or discontinuous distribution, simulating input to a row of 100 fields. Rate data were randomly sampled to derive input rates to each field, and copies added to reflect the fact that each afferent cell contributes a randomly variable number of terminals. Red data represent copies-included signals received by a row of three fields. See Methods for derivation of rates. **Column 2** The first-column red data were randomly sampled 150 times, sample size 4, representing mossy fibre input to a 3-field population of 30 Golgi cells, each with 5 basal dendrites (the mean) which each receive contact from 4 mossy fibres. Data are the sample means, representing dendritic depolarisation. **Column 3** We next randomly sampled column 2 data *without* replacement 30 times, sample size 5, and took the sample means, representing somatic integration of dendritic signals. **Column 4** Column 3 data were randomly sampled 700 times, sample size 8–12, reflecting convergent input from 3 fields of Golgi cells onto 700 glomeruli in the middle field (which receives the output of a three-field ensemble), and took the sample means, representing GABA spillover that is a linear function of the mean of afferent rates. Inhibition of granule cells is almost exclusively mediated by spillover. **Column 5** The column 4 data are presented as a histogram. **All rows** Column 1 mossy fibre data are converted to narrowly-focussed inhibition of granule cells. The mean of the column 4/5 data is linearly related to the mean of the mossy fibre distribution. This result is independent of the shape of the original distribution.

1100

Fig 3. The Golgi cell population of a rank functions as a single unit. A Simulation of
Golgi cells in a sagittal row of 100 fields, taking the mossy fibre distribution on the left
as input (see Fig 2 and Methods for derivation of mossy fibre rates). **Top right**: pink:

the mean of mossy fibre input rates to each field; red: the mean with duplicate signals
added; dashed grey line: the mean of input rates to the whole rank. **Bottom right**: red:
same as top right; blue: mean, for each field, of glomerulus data derived in the same
way as Figure 2 columns 4 and 5 (the 'inhibitory field mean'); pale blue: SD of
glomerulus data. **B** Same as (A) but sampling a different mossy fibre distribution.

A and B

Red data closely track the pink data in top right panels, indicating that duplicate signals
(because the mossy fibre axon ends in a cluster of terminals) do not skew the mean of
firing rates received by a field.

Blue data across all fields are in a narrowly circumscribed range. Focus varies with the
bandwidth of the sampled distribution (mossy fibre rates). A normal mossy fibre
distribution with a range of over 100 Hz is converted to a blue range of ~3 Hz.

Inhibitory field means are tightly grouped: inhibition of granule cells is synchronised in
a rank.

The range of the blue data is centred on the grey dashed line – the mean of mossy
fibre input rates received by the whole rank – and varies independently of the red
data. That is, unpredictability inherent in the mechanism (because the outcome of
random sampling is inherently uncertain) is not compounded; random variation of the
red data doesn't flow through to amplify random variation of the blue data (in fact the
blue data have a slightly smaller range).

C The same simulation. Left: the sampled distribution (mossy fibre input rates to a
rank); right: the inhibitory field means (we use the means to compare fields). The y axis
of the right panel has the same range as the x axis of the left panel. Panels are tall to
improve visual comparison to other rows, where the y axis has a smaller range. **D** and **E**
The same, moving the sampled distribution to the right. Inhibition of granule cells is
precisely and linearly coupled to the mean of mossy fibre input rates to the whole
rank.

1131

1132

1133

Fig 4. Regulation of granule cell activity by the feed-forward pathway (mossy fibres→Golgi cells→granule cells) We incorporated the Golgi cell ensemble conversion into a simulation of the conversion of mossy fibre signals into granule cell signals (recoding). One of the outputs of recoding is regulation of the percentage of active granule cells. We tested if the recoding computation causes the percentage to be stable, i.e., in the same narrow range in all fields regardless of the distribution of mossy fibre rates. **Column 1** Mossy fibre firing rates were randomly generated with a uniform, discontinuous, three-peak or normal distribution, to represent input to a row of 100 fields. Rates, and rates received by each field, were derived in the same way as earlier Figures. Red data are rates received by a single randomly-selected field, including duplicate signals. y axes are scaled to the data – all distributions contain the same number of signals. **Column 2** The percentage of granule cells that fire in each field. The Golgi cell computation includes no adjustment to sparsen granule cell activity. The white line and shaded area are mean and SD. **Column 3** The same, with an adjustment (a co-efficient) to sparsen granule cell activity. y axes have 1/10 the range of column 2. The percentage is confined to a narrow range. This result is independent of the sampled (i.e., mossy fibre) distribution, suggesting that regulation is

1151   independent of fluctuation of mossy fibre rates. Nonetheless, field-to-field variation
1152   relative to the mean is significant. Homeostatic local fine tuning may be provided
1153   physiologically by feedback via granule cell ascending axon contact on Golgi cells, not
1154   represented in the simulation.

1155

Fig 5. A rank is a single recoding unit. The same simulation as Figure 4 but data are
granule cell firing rates or derived from granule cell firing rates. **Column 1** We tested
the effect of changing the shape of the mossy fibre distribution (without changing the
number of signals). **Column 2** Pale green: sample means – the mean of input rates to
each granule cell in a single field selected at random; blue: sample means of the subset
of granule cells which meet the conditions to fire. In rows A–C, ~30% fire. In rows D–F,
~1% fire, y axes are rescaled and green data are omitted. **Column 3** Red: mean of
mossy fibre input rates to each field in a rank, including duplicate signals; pink: SD of
mossy fibre input rates to each field; blue: field means (i.e., the mean of granule cell
firing rates in each field); pale blue: SD of granule cell firing rates in each field.

Observations

Regardless of the shape and large range of the mossy fibre distribution, the field
means always converge towards a straight line which is a fixed distance from the
overall mean of mossy fibre rates.

The amount and direction of field-to-field random fluctuation of the blue data is
independent of local fluctuation of the red data, and in a smaller range. Uncertainty
inherent in random sampling – first by single fields of input rates to a rank, and then by
Golgi cells and granule cells of input rates to a field – is not compounded.

1175

1176

Fig 6. Granule cells collectively code mossy fibre rate information. Column 1 Mossy
fibre rates were generated in the same way as earlier Figures, with either a normal (SD
12.5) or uniform distribution representing input to a row of 100 fields, with copies
added – the sampled distribution. **Column 2** Histogram of granule cell firing rates in a
randomly selected field, obtained in the same way as in the bottom 3 rows of Fig 5.
Column 3 Grey line: the mean of the sampled distribution; blue line: field means. **A–C**
The fields means are precisely and linearly related to the mean of mossy fibre rates.
Blue data approach a straight line which is parallel to the grey line at a mean distance
of 15 Hz (SD 1.21 Hz). **D–F** If the shape and mean of a uniform sampled distribution are
unchanging, the distance between the blue and grey lines is linearly related to the
range.

1188