1

Breaking the code – pixels, a surprising solution to neural

network problems: the cerebellar input layer

Running head: Breaking the code: the cerebellar input layer

Mike Gilbert[a]* and Anders Rasmussen[b]

[a]School of Psychology, College of Life and Environmental Sciences, University of

Birmingham B15 2TT, UK

[b]Department of Experimental Medical Science, Lund University, BMC F10, 22184 Lund,

Sweden

* Corresponding author; email: m.gilbert.1@bham.ac.uk

ABSTRACT

The cerebellum makes up 20–30 % of the brain by mass and volume in most vertebrates.

It is thought that neuronal signal frequencies – firing rates – provide the physiological

form of values that are used in neural network computations. The cerebellum is made up

of repeating, anatomically overlapping networks. Neural network modelling has a long

tradition of learning algorithms that are implemented physiologically by learned changes

of synaptic transmission strength. In the cerebellum, the evidence has ultimately failed to

corroborate the predicted role of learning.

We ask, with a focus on the input layer of the cerebellar network:

1) If the agent of change is not synaptic memory, what is it? What is the physiological

mechanism of the cerebellar network computation?

2) What is the computation, and how is information represented by internal signals

activity – what is the code?

3) How is that related to the matrix-like internal architecture of the cerebellar cortex, and

the well-known functional organisation of the cortex into long, thin strips called

microzones?

Our approach is to propose a reinterpretation of the rich anatomical and biophysical

literature and test it with computer modelling. Modelling provides a test because

information about computationally-relevant parameters are detailed, so we can replicate

networks in high resolution, including anatomical noise. Using this method, tested *in*

silico, we found the evidence plausibly supports controversial conclusions. Using the

same numbering:

1) The input layer computation is the functional result of cell morphologies and network

architecture. Passing signals through a network provides the biological equivalent of

mathematical operations, which run in series and in parallel.

2) Information is coded in statistics of internal firing rates. At this collective level, there

are reliable statistical effects that conserve rate information between network layers.

3) Still more unexpectedly, multi-dimensional detail contained in input signals is

converted to an extremely low resolution – pixelated – internal code. However, as a

result, the internal code is intelligible by randomly sampling signals. That means

downstream targets can receive different signals but the same code, providing an

otherwise elusive mechanism to coordinate firing in microzones.

The proposals are bespoke because cell morphologies and the connections between

them are unique to the cerebellum. However, the idea that anatomy and 3-dimensional

network architecture implement mathematical operations unaided, if it holds water, is

likely to have implications for coding in other brain regions.

Keywords: theory, computational neuroscience, cerebellum, network, code

1. INTRODUCTION

Most of the cerebellum by mass and volume is made up of the cerebellar cortex, which is divided into two layers, an inner layer, the granular layer, and outer layer, the molecular layer. The main source of glutamatergic input to the cerebellum is from mossy fibres. Mossy fibres terminate in the granular layer, on granule cells and on Golgi cells, which inhibit granule cells. The granule cell axon rises into the outer layer where it divides in two to form parallel fibres. Parallel fibres lie parallel to the cerebellar surface and each other. Parallel fibres contact Purkinje cells, which carry the only output of the cerebellar cortex. Our focus is the granular layer.

In the traditional cerebellar model, the supervised learning model (1, an excellent review), control of Purkinje cells by granule cell rates is displaced by learning coded as parallel fibre synaptic weight changes (2-5). After training, when a Purkinje cell receives a remembered pattern, it fires at a learned rate. It is implicit that a granule cell code is redundant because control is surrendered to learning. Supervision is provided by instruction signals which would, in principle, permit Purkinje cells to be trained and controlled individually in this way.

We challenge that. Modelling physiological networks presents generic problems. It is thought to be a safe starting point that the variable rate at which neurons discharge action potentials – firing rates, sometimes called signal frequency – provides the physiological equivalent of numbers that are used to code information, and that neural

networks provide the physiological implementation of computations which take firing

rates as inputs. Relating that to physiology is not straightforward. Networks can be large

and typically contain multiple cell types. Cell ratios – the ratio of cells of each type to

other types – all differ. So do convergence and divergence ratios for each type onto other

types they contact. Physiological networks are usually not anatomically separate, often

sharing cells, sometimes most of them. Insentient networks don't 'know' what data

values are a measure of, or what scale they are measured on, especially problematic if

signals are mixed up. Biology is messy: most network parameter values are randomly

variable.

One modelling approach is to simplify the parameters – for example, replacing cells with

generic 'units'. Another is to model in abstract – how might an artificial system perform

the computation a network may have? There is a theoretical reliance on synaptic weights

to provide the mechanism of the network conversion. This has become unsafe. In the

cerebellum, experimental evidence has ultimately failed to provide support for the

expected role of long-term depression in motor learning (6-8) or for finely-graded

modification of synaptic weights predicted by learning models (9, 10). There is

accumulating evidence of linear transmission of rate-coded data, in conflict with models

that use experience to train synaptic weight adjustments. For example, cerebellar

neurons have been repeatedly found to fire at a rate that is a linear function of

behavioural and task metrics, and this 'has been found at all levels of the cerebellar

circuit' (1 p.239, citing 30 sources).

We asked, focussing on the input layer of the cerebellar network:

1) If the agent of the network conversion is not synaptic weights, what is the mechanism

and what is the computation? If transmission is linear, what is the computational effect

of cell and network anatomy?

2) How does the cerebellum encode information internally, in firing of granule cells? The

traditional model does not need a code because learning transfers control of Purkinje

cells from firing rates to synaptic weights. If control is instead by rates, as we propose,

what is the code?

3) How does that help to explain functional anatomy? Also, how is it functionally related

to the well-established organisation of Purkinje cells into microzones (11-13)?

Thinking about it in this way mitigates issues inherent in generic networks because the

model is not generic. Anatomical data are rich. If anatomy implements computations,

then knowing model parameters, we can model it. We are able to include detail that

makes cells and networks unique, including random variability of parameter values

(noise).

We find, using the same numbering:

1) Anatomy – cell morphologies and network input layer architecture – feasibly provides

the biological equivalent of mathematical operations. Computational performance relies

on linear transmission.

2) Granule cell rate information is collectively coded in statistical properties (mean and

frequency distribution) of the firing rates of co-active cells. Code in this form means

granule cells can read mossy fibre rates by taking random samples, and the granule cell

code can in turn be read by randomly sampling granule cell rates. It also means mossy

fibre rate information is conserved by a statistical relationship of external and internal

signals. The relationship is independent of the permutation of active inputs and the

permutation of the rates they each fire at. Anatomical noise does not impair

computational performance.

3) Still more surprisingly, high-dimensional input data are converted to a low-resolution

internal code. Information has spatial dimensions. The smallest unit of internal

information is a strip with microzone dimensions (viewed from the cerebellar surface)

but fuzzier boundaries. In this sense, the internal code is pixelated. The same information

is contained in any random sample of signals inside those boundaries, however. This

provides a substrate for mixed-up, multi-dimensional input signals to drive coherent

firing of microzone-grouped cells downstream.

Since anatomy is the key instrument of information processing in this contention, and

provides model parameters, each main section of the paper begins with a summary of

the literature. Evidence of linear transmission is also discussed in context.

The structure of the paper follows the steps of the computation. It is accordingly in three

parts headed 'Mossy fibres', 'Golgi cells' and 'Granule cells'. The second and third parts

include a results section and our interpretation of the results. The theoretical nature of

the work means there is no methods section. Our method is: (1) Propose the steps of the

network input layer computation. (2) Present evidence that neurophysiology implements

the steps. (3) Test it with a computer program that replicates the computationally

relevant metrics of anatomy, and includes noise.

'Model' can be ambiguous. In artificial neural networks, for example, it is a mathematical

statement of the proposal – the computation is synonymous with the model. *That is not*

our method. In our approach, modelling provides a way to quantify the proposals so that

we can test the computational effect of physiological parameters, if they are related the

way we claim they are. We can do that in high-resolution detail because there is detailed

anatomical information available. The steps of the computation are straightforward. The

main function of computer modelling is a test.

The proposition of linear transmission of rate information comes with an advantage.

Information transitions through several biophysical forms in transit even through a single

cell. This has tended to divide modelling approaches between internal steps of signalling,

to account for biophysical data, and network level signalling. Bringing them together is a

knotty problem partly because different states are measured on different scales. Linear

transmission makes the problem tractable. If firing rates are linearly related it is a

plausible proposition that biophysical transitions are a chain of functionally linear steps

regardless of the values they would take on a conventional scale. Indeed, non-linear

interim steps are improbable. Linear rate relationships accordingly provide a basis to

represent all steps on a common scale. Linear intracellular transmission adds steps to the

network computation. It forms part of our proposal that linear biophysical transitions

exist for this computational effect.

158 2. MOSSY FIBRES

159 2.1 Contents

A single mossy fibre has several, sagittally-aligned branch endings, each ending in a

cluster of terminals. A sagittal strip of the granular layer therefore receives multiple

copies of each mossy fibre signal at multiple locations. As far as we know, cerebellar

theory has not previously addressed this.

We derive in this section the number mossy fibres that innervate a strip, the number that

innervate defined local sectors, and the ratios of the number of those that contribute a

single terminal, two and so on. That is, the number of input rates received be each

sector, and the number of copies of each rate, taking into account random variation of

the number of terminals per cluster, and the fact that most clusters fall partly inside and

partly outside a sector. This information is used to generate mossy fibre signals that are

received as inputs to the simulations of Golgi cell computations and recoding of mossy

fibre into granule cell signals in sections 3 and 4. It is essential to very precisely simulate

anatomy to test that real networks can execute the computational function we say they

have (and that the computation is noise-tolerant).

This reveals that the important function of terminal branching is to simulate sampling

with replacement by sectors of mossy fibre input rates to a strip as a whole. Simulated

replacement has the computational role in coding that sectors each receive the same

mean of rates as the sampled distribution, notwithstanding relatively modest per sector

sample size.

2.2 Anatomy

The molecular layer of the cerebellar cortex is organised functionally into long thin strips

called microzones, around 100 times long as they are wide, populated by Purkinje cells

and interneurons. Some 35 million parallel fibres pass through a microzone, at right

angles, approaching from both sides, from a distance of up to 3 mm (14-17).

Mossy fibres give rise to sagittally-extending collaterals, the direction of the long axis of

microzones (Fig 1). These in turn give rise to multiple distal branches, each ending in a

cluster of terminals (18-21). Terminal clusters are sagittally aligned and separated by a

randomly variable distance typically not less than 200 μm (18, 19) measured from edge

to edge of clusters (so they are separated by a greater distance measured from centre to

centre). Terminals are intimately and randomly intermingled with terminals of other

mossy fibres.

So, mossy fibres do not terminate at a point but a region with dimensions: the field

occupied by a terminal cluster. Cluster size is variable (because it depends on the number

of terminals and how tightly clustered they are), averaging 200 µm sagittally x 150 µm

mediolaterally (19), a 'field' (a sector in section 2.1). A sagittal row of 75–100 fields (a

'rank') has the area of a microzone viewed from the cerebellar surface. Microzone

dimensions vary, but a mediolateral span of 150 µm is mid-range (5, 11-13). Where

microzones are functionally defined by the topography of termination of climbing fibres

on Purkinje cells, a rank is a division of the granular layer and has less sharply defined

boundaries. Mossy fibre input to the cerebellum evoked by cutaneous stimulation largely

shares the topographical organisation of climbing fibres (22-24).

Figure 1 about here please

2.3 Derivation of the size of the sampled population and sample size

Fields randomly sample mossy fibre input rates to a rank, we propose. To simulate that,

we need the size of the sampled population – the number of mossy fibre signals received

by a rank – and sample size, the number received by a field.

Despite the repeating 'crystalline' architecture of the cerebellar cortex, at local level

there are anatomical random variables. Variables are: i) the number of mossy fibres that

innervate a field, ii) the number of terminals they each end in, and iii) the number of

those that are inside field limits.

The number of mossy fibres afferent to a field has been estimated at 100 (19). However,

that number is a field-sized fraction of the general population. The actual number is

larger because mossy fibres end in a dispersed group (cluster) of terminals, and most

groups straddle field limits.

We work from an estimate of the total number of terminals per field, which is, in turn,

derived from convergent estimates of the number of granule cells per field. (Estimate 1)

There are an estimated 1.92×10^6 granule cells per µl (14). The number of fields which fit

in that volume is $(1,000/150)^2 \times (1,000/200) = \sim222$, assuming a field depth of 150 µm

(19). The number of granule cells in a field is, therefore 1,920,000/222 = 8,649. (Estimate

2) An estimated 350,000 parallel fibres pass through a single Purkinje cell's territory (14).

Parallel fibres run about 3 mm in each direction from the granule cell soma (16, 17). The

Purkinje cell dendritic arbour fills the molecular layer vertically and averages ~200 µm

wide. Therefore, 6,000/150 = 40 fields fit in a row that provides parallel fibres that pass

through a Purkinje cell dendritic territory, giving 350,000/40 = 8,750 granule cells per

field.

Granule cells have 3–5 dendrites, averaging 4, and receive input to each of them from a

single mossy fibre. Estimates of divergence of a single mossy fibre terminal onto granule

cells vary (25). It is unclear how much this reflects biological diversity versus differences

in opinion and methods. Here we use a mid-range estimate of 1:50 (26, 27). Assuming a

mossy fibre terminal receives a single granule cell dendrite from each of d granule cells,

and given an average of 4 dendrites per granule cell and z granule cells per field, the total

number of mossy fibre terminals in a field is $N = (z \times 4)/d = 700$ given $z = 8,750$ and $d =$

50. A convergent estimate is provided by multiplying the per field number of mossy

fibres by the average number of terminals per cluster.

Cluster size varies. We take the range as 4–12 terminals per cluster (19). We assume the

number varies at random and cluster sizes occur with equal probability. A maximum of 7

fits in a field (because a field shares the dimensions of the area that encloses a mean

cluster). Fields are not anatomically bounded, so many clusters straddle field limits. To

reflect that, we assume there is a 1/7 probability that a 7-terminal cluster (or larger) has

7 terminals inside a field and also the same probability of any other number. A 6-terminal

cluster has a 2/7 chance that all 6 terminals are inside and a 1/7 chance of any other

number, and so on.

Table 1 shows the derivation of the expected (i.e., predicted by probability) ratios of

terminals contributed to a field by each cluster size (R1), not counting terminals that lie

outside when a cluster straddles field boundaries; and the ratios of mossy fibres

contributing $n = 1$–7 terminals (R2), regardless of cluster size. R1 ratios are given by the

sum of numbers in each row. So, for example, the ratio of cells that end in 8 terminals to

cells that end in 5 is 28:25. R2 ratios are given by the number of appearances of each

number in the grid – so the expected ratio of cells which contribute 7 terminals to the

number which contribute 4 is 6:12. The ratios effectively give the distributed probability

of possible states. For example, there is a probability of 10/(6 + 8 + 10 + 12 + (3 x 9)) =

0.159 that a mossy fibre contributes 5 terminals to a field (regardless of cluster size).

Similarly, there is a probability of ((6 x 28)/((6 x 28) + 27 + 25 + 22))/7 = 0.099 that a given

cell contributes 7 terminals to a field.

Table 1. Derivation of expected ratios

size	number of terminals inside							R1
12	7	6	5	4	3	2	1	28
11	7	6	5	4	3	2	1	28
10	7	6	5	4	3	2	1	28
9	7	6	5	4	3	2	1	28
8	7	6	5	4	3	2	1	28
7	7	6	5	4	3	2	1	28
6	6	6	5	4	3	2	1	27
5	5	5	5	4	3	2	1	25
4	4	4	4	4	3	2	1	22
R2	6	8	10	12	9	9	9	

If R1 ratios are denoted by a series $x_4, x_5 \ldots x_{12}$, and a given element in the series is x_j,

the fraction of terminals provided by a given cluster size is $x_j/(x_4 + x_5 + \cdots + x_{12})$, and

this fraction is multiplied by N to give the number of terminals. The number of terminals

is divided by the expected mean number of terminals that fall inside field limits (for that

cluster size) to give the expected number of mossy fibres that provide them. The number

of j-terminal mossy fibres that innervate a field is therefore given by

$$N\left[\frac{x_j}{\sum_{i=1}^{n} x_i}\right]/\left[\frac{x_j}{7}\right]$$

The results are shown in Table 2. The equal number of mossy fibres (mfs) in each column

is what we would expect, because we made it an assumption that all sizes occur with

equal probability, and mf values are expected values – i.e., in proportions predicted by

their probability. The mf total is the expected number of mossy fibres afferent to a field

(which contribute at least one terminal).

Table 2. The expected number of mossy fibres which innervate a field

cluster size	12	11	10	9	8	7	6	5	4	total
terminals	81	81	81	81	81	81	78.1	72.3	63.6	700
mfs	20.25	20.25	20.25	20.25	20.25	20.25	20.25	20.25	20.25	182.25

271 R2 ratios – the relative incidence of afferent mossy fibres that contribute 7 terminals to a

272 field, 6 terminals, 5 and so on, down to 1 – and the expected number of mossy fibres that

273 contribute them, are shown in Table 3.

274 **Table 3. The number of mossy fibres which contribute n = 1–7 terminals to a field**

n	7	6	5	4	3	2	1	total
R2	6	8	10	12	9	9	9	63
R3	42	48	50	48	27	18	9	
mfs	17.34	23.15	28.92	34.72	26.04	26.04	26.04	182.25
%age	9.52	12.69	15.87	19.05	14.29	14.29	14.29	

R3 is n x R2, giving the ratio of terminals provided by mossy fibres with n terminals inside

a field. %age is given by (R2/63) x 100. this is the percentage of mossy fibres with n

terminals inside a field.

From here, the number of mossy fibres that innervate a rank is given by 182.25 x 100

divided by the mean number of clusters per mossy fibre per rank. If a rank receives, on

average, five clusters per mossy fibre, ~3,645 mossy fibres innervate a rank. This is the

estimate used in the simulation unless stated otherwise.

Assumptions. Cluster sizes occur with equal probability. The probabilities of each

outcome when a cluster straddles field boundaries (how many terminals are inside and

how many outside) are as stated. A single mossy fibre ends in an average of 5 terminal

clusters per rank.

287   2.4 Derivation of mossy fibre input rates to a field

Terminal clustering means a field receives multiple copies of most signals. This section

lists the steps in the simulation that generate mossy fibre firing rates received by a field.

As noted, at local level, anatomical parameters are random variables. Random variables

are randomly generated (in steps 1, 2 and 4 below) with the stated probabilities.

In the first step, the simulation randomly generates, for each field, the cluster size ratio.

This is the ratio of afferent mossy fibres (which contribute at least 1 terminal to a field)

that end in 4 terminals, 5, 6 and so on up to 12, assuming they occur with equal

probability. In step two, the ratio of cluster sizes is weighted to reflect the greater

probability that large clusters contribute more terminals to a field (with probabilities

reflecting the expected ratio of terminals contributed by each cluster size). In step 3,

ratios are converted to the fraction of terminals contributed by each cluster size. When

multiplied by the total number of terminals, this gives the number of terminals

contributed by each cluster size. In step four, the values obtained in step three are

divided up between cells that contribute a single terminal, 2 terminals, 3 and so on, to

reflect straddling of field limits. The proportions give the number of mossy fibres afferent

to a simulated field that contribute a single terminal, two, three and so on up to seven,

the maximum.

There is no field-level effect of mossy fibre terminal branching. Terminal clusters arising

from the same mossy fibre are separated by a minimum distance, so that a field cannot

contain terminals of more than one cluster per mossy fibre.

Terminal clustering means that fields contain a randomly variable number of copies of

most signals. In the simulation, copies are added after sampling of mossy fibre rates to

obtain rates received by a field. An alternative order would be to add copies to the

sampled population and then sample that. We follow anatomy: the number of *rates*

received by a field is equal to the number of afferent mossy fibres; the number of *signals*

contains duplicate rates.

2.5 Mossy fibre terminal branching simulates sampling with replacement

Termination of a single mossy fibre at multiple locations in a rank is an anatomical

solution to a computational problem. The problem is that the granular layer computation

exploits statistical effects which require that a Golgi cell at any location of a rank must

randomly sample the entire distribution of mossy fibre input rates to the whole rank. The

same statement applies to granule cells. There is an additional requirement. This is that

samples are independent, meaning, all samples have an equal probability that they

contain any given rate picked from the sampled distribution. If samples were being

drawn from a bag of coloured balls, balls would need to be replaced between samplings,

so that, as samples were withdrawn, it did not change what was left in the bag. Hence

sampling 'with replacement'. The physiological equivalent of random sampling is contact

from a random sample of the afferent cell population. All samples are 'selected'

concurrently. There is no replacement. Physiology must therefore simulate replacement.

It is worth stating clearly that this is not a simplification but a hypothesis.

If each mossy fibre could terminate in only one field, a mossy fibre 'sampled' by one field

cannot also be sampled by any others. However, termination by a single mossy fibre in a

random number of terminal branches, at multiple locations, removes that restriction.

The result is an approximation of sampling with replacement. In theory, sampling with

replacement should mean any number of fields can receive the same signal, up to all of

them. Clearly this is not true – there is an anatomical limit. But in practice even with real

replacement the number of repeat samplings is constrained by a probability distribution.

That is, at large numbers, the number of fields that can receive a signal would be limited

just as surely if there was actual replacement as it is in reality by anatomy.

That means we can test if terminal branching simulates replacement by calculating the

probability distribution in the hypothetical case that mossy fibres did not branch

terminally and rates were in fact sampled with replacement. The probability, p, that a

given rate is selected k times in taking N samples, sample size x, assuming mossy fibres

each give rise to an average of m terminal branches, is given by

344
$$\left[{Nx!}/{k!\,(Nx-k)!} \right] * \left[{1}/{(Nx/m)} \right]^{k} * \left[1 - \left({1}/{(Nx/m)} \right) \right]^{Nx-k}$$

346   The results are shown in Table 4.

347   **Table 4. Probability p of k repeat selections of a single mossy fibre**

k	0	1	2	3	4	5	6	7	8	9	10	11
p	.007	.034	.084	.140	.175	.175	.146	.104	.065	.036	.018	.008

The number of repeat selections is consistent with what we know about the biological

range of the number of terminal clusters per mossy fibre (19, 28), noting that data are

limited. Estimated by this method, the mean is 4–5. Put another way, the anatomical cap

on the number of fields that can receive a signal (set by the number of terminal branches

of that cell) is reflected in the simulation by the probability distribution. The simulation

reflects anatomy and anatomy simulates replacement.

2.6 The range of mossy fibre firing rates

Some very high mossy fibre rates can be generated in experimental conditions. However,

it is thought the typical physiological range is 50–300 Hz (29) citing (30). In the

simulations, rates are generated at random in that range, within stated constraints on

the shape of the distribution. Figures display the sampled distribution.

3. GOLGI CELLS

3.1 Contents

We propose a functional organisation of Golgi cells into 3-field groups. Ensembles

implement a computational conversion of mossy fibre firing rates received by an

ensemble territory into synchronised inhibition of granule cells in the middle field.

Ensembles are not anatomically bounded and merge functionally into sagittal strips so

that the unit of conversion is a rank. Ensemble-grouped Golgi cells all innervate

substantially the whole of the middle field: convergence onto the middle field defines the

group.

'Input' is to an ensemble territory – i.e., three fields – which is sampled by Golgi cells.

Output is inhibition of middle-field granule cells. More exactly, there are 700 outputs, the

number of glomeruli, that mediate inhibition. Some of the steps of the computation are

intracellular. It forms part of the proposal that in physiological practice the only variables

with significant influence are ensemble architecture and firing rates. A linear relationship

of input and output is evidenced. Anatomy provides parameters and the values they

take, next.

3.2 Anatomy

Anatomy implements ensemble computations. We test that by replicating an ensemble

in silico using computationally relevant metrics of anatomy. Anatomical evidence

provides parameter values, normally a range.

Golgi cells are large interneurons whose cell bodies lie in the granular layer. They are

most concentrated just below the Purkinje cell layer (31). The Golgi cell axon branches

profusely, giving rise to a dense plexus. The axonal field is sagittally elongated, that is, in

the direction of the long axis of a microzone – mean range 650 +/- 179 μm by 180 +/- 40

μm (in mice: 32) – and is the depth of the granular layer, which it fills vertically. Their

density (33) and large, profusely ramified axonal field (32) means that Golgi cell axonal

territories overlap extensively in the sagittal direction. The Golgi cell population of a row

of three fields all inhibit approximately the whole of the middle field. Convergence of

their axonal fields defines the size and territory of the smallest functional Golgi cell unit –

an ensemble – we suggest. Their combined range is longer – five fields. But convergence

on the middle field is functionally significant because it means all granule cells in the

middle field receive inhibition from a random sample of a three-field unit.

There is a significant population of Golgi cells at deeper level. Deeper cells tend to have

smaller cell bodies and to lack dendrites that extend into the molecular layer. However,

the axons of smaller Golgi cells are indistinguishable from large Golgi cells (31).

4 Purkinje cells fit in a field, based on an estimate of the number that span a microzone

(5). Estimates of the ratio of Golgi cells to Purkinje cells vary. 'In man, the ratio [of

Purkinje cells to Golgi cells] is 1:1.5, while in the monkey and cat it is almost 1:1.9 and in

the rat 1:3.3' (34). Estimates may be low where Golgi cells were identified by their large

size, so that smaller cells at deeper level were not counted (31 p.121). We take it that

there are 10 Golgi cells per field, a ratio of 1:2.5 with 4 Purkinje cells, so 30 in an

ensemble.

Contact by mossy fibres on a Golgi cell is on basal dendrites. 4–6 basal dendrites arise

from the Golgi cell soma. If we take it as 5, a population of ensemble grouped Golgi cells

has a combined total of 30 x 5 = 150 dendrites. The number of mossy fibres which

contact a Golgi cell is unreported. It has been estimated that a Golgi cell 'cannot receive

more than 10–12 MF [mossy fibre] inputs, given the total basolateral dendrites length

and the density of glomeruli' (35). Without more data it is challenging to estimate how

many make contact. Two per dendrite (10 between an average of 5 dendrites) is low.

Dendrites radiate from the cell body. A single dendrite has a range of 50–100 µm (31).

The longest dimension of a field – which contains ∼700 mossy fibre terminals – is 200

µm; on the other hand, there is little dendritic branching, so a dendrite occupies a path

not a volume. We take the number as 20, 4 per dendrite.

Golgi cells inhibit granule cells in a structure termed a glomerulus (31, 36, 37). Mossy

fibre terminals are ensheathed in a semi-permeable membrane which restricts

neurotransmitter diffusion. Excitation of granule cells by mossy fibres and inhibition by

Golgi cells takes place here. Most of the Golgi cell axonal plexus – which is confined to

the granular layer – is made up of very fine beaded fibres. These enter glomeruli, where

they contact granule cell dendrites (37). 'In the adult rat cerebellum, each granule cell

dendrite receives 2.6 ± 0.55 synaptic contacts from Golgi axon terminals' (38) citing (27).[1]

However, the large majority of inhibition (98%) (39) is by spillover, where

neurotransmitter released into the synaptic cleft spills out into the glomerular space.

Golgi cells release GABA, an inhibitory neurotransmitter. This is detected by high-affinity

$GABA_A$ receptors located perisynaptically and extrasynaptically on granule cells (40, 41).

Even synaptically-received signals have most of their effect (i.e., the large majority of

charge transfer is mediated) via spillover.[2]

---

[1] Data are sparse. The original data are an estimate derived from visual inspection of slices thirty years ago so far without later corroboration, except from Rossi and Hamann 1998, who derive a lower estimate but propose the disparity is because their estimate is too low.

[2] Because of the long duration of IPSCs generated by spillover (recorded in slices), the total charge carried is three times that of IPSCs generated by directly connected terminals. 38. Rossi DJ, Hamann M. Spillover-mediated transmission at inhibitory synapses promoted by high affinity alpha6 subunit GABA(A) receptors and glomerular geometry. Neuron. 1998;20(4):783-95.

As far as is known, inhibition received by a glomerulus is from a random sample of Golgi

cells afferent to a field. The number of Golgi cells with convergent input to a single

glomerulus is unknown. There is high divergence. Each Golgi cell inhibits a large number

of granule cells (32, 42). A single Golgi cell contacts thousands of granule cells (37). Given

a fixed cell ratio, the convergence ratio scales proportionally with divergence. We take

the ratio as a random variable in the range 8–12:1 (we test the effect of other ratios in

Fig 3).

3.3 The Golgi cell ensemble computation

It is thought that discharge rates of neurons provide the physiological form of values that

are used in network computations. That is, firing rates provide the biological equivalent

of numbers, and transitions that change rates can be represented mathematically.

We argue in this section that transmission through a Golgi cell ensemble is in biophysical

steps that can be represented as mathematical operations. Steps are a sequence; a single

step is a compound of multiple operations reflecting events that occur in parallel.

Information is represented at ensemble-wide level. Each step takes a population of

values and converts it to another population of values. Ensembles exploit statistical

effects of random sampling. There are two stages of random sampling. Ensembles

perform better than single cells because there are more layers of sampling and more

samples per layer.

Say a population of values (with any frequency distribution) is randomly sampled. There

are n samples, sample size m. If we take the mean (or sum) of each sample, and n is large,

the sample means have a normal or near-normal distribution, by the central limit

theorem. (This works even if m is a range.) The new distribution is centred on the same

mean as the sampled distribution, but narrower. Modest values of n or m give an untidy

result. However, if the new distribution is then itself randomly sampled, and the process

is repeated, the results get tidier and narrower.

To apply that to Golgi cell ensembles, the first-sampled population is of mossy fibre rates

received by three fields. Three fields contain a total of 30 Golgi cells. Rates are sampled

by Golgi cell basal dendrites. At five dendrites per cell, the number of samples is 30 x 5 =

150. Sample size is the number of mossy fibres which make contact per dendrite, which

we take to be 4.

The postsynaptic result – charge influx to dendrites – is a proportional function of input

rates. Otherwise, it would not convey rate information (43). As far as we know, there are

no data for transformation of multiple mossy fibre input rates into Golgi cell basal

dendritic depolarisation. We take it as the mean. (We discuss evidence for physiological

conservation of rate information in the next section 3.5.2.) Taking the sample means

creates a new set of values (Fig 2 column 2). The frequency distribution of the new values

is in a more condensed range and centred on the mean of the sampled distribution. Basal

dendritic charge transfer is passive and little filtered (44, 45). Integration of dendritic

states at the Golgi soma is represented by taking the mean of dendritic charge. This

generates a third population of values (Fig 2 column 3). Golgi cell firing rates have a linear

relationship with the amplitude of depolarising current (46). Thus, rate-proportional

dendritic charge influx is converted to Golgi cell firing rates.

The second stage of random sampling is of Golgi cell rates by glomeruli. Inhibition almost

exclusively by spillover (39-41) effectively increases sample size, as sample size is not the

number of Golgi cells that make direct contact on any particular granule cell, but the

number afferent to a glomerulus. Sample size is accordingly the number of Golgi cells

with convergent input to a glomerulus (taken as 8 to 12), and the number of samples is

the number of glomeruli per field (700). Rate-proportional spillover provides the

physiological equivalent of averaging afferent rates, represented by taking the sample

means. This generates a final population of values (Fig 2 columns 4 and 5), the output of

the ensemble conversion.

3.4 Results

3.4.1 A single ensemble

The results for a single ensemble are shown in Figure 2. Neural information has a shifting

form. The ensemble conversion has intracellular as well as between-cell steps and

therefore transitions through forms that are normally measured on different scales. We

use a common scale because our interest is confined to the computational effect of

architecture through the steps of the conversion, assuming a linear relationship of forms.

Figure 2 about here please

3.4.1.1 An ensemble converts randomly sampled mossy fibre rates into synchronised

inhibition of granule cells

Outputs of the simulation of a Golgi cell ensemble are graphed in Figure 2. Mossy fibre

firing rates received by a rank were generated as described in section 2 within

constraints on the frequency distribution (column 1). The distribution of firing rates

received by a row of three fields selected at random is shown (in red) in the same panel.

In column 2, the red data are themselves randomly sampled – 150 samples, sample size

4, representing 30 Golgi cells x 5 basal dendrites, and contact on each dendrite by 4

mossy fibres – and the sample means plotted, representing dendritic charge states. The

range is narrower than the sampled population (in all rows), but data are not tightly

grouped.

In column 3, the sample means are themselves sampled (without replacement) – 30

samples, sample size 5, representing 30 Golgi cells with 5 dendrites each – and the

samples averaged (and plotted), representing somatic charge states. Somatic data are in

a narrower range than dendritic data but still not tightly grouped.

In the final step (column 4) somatic data are sampled 700 times, reflecting convergent

input from 3 fields of Golgi cells onto 700 glomeruli in the middle field. Sample size is 8–

12, randomly generated for each sample; physiological sample size is unknown. Column 5

shows the column 4 data as a histogram, for comparison with column 1.

The result is that mossy fibre signal frequencies received as input to a rank are reliably

converted to a narrow distribution of values representing inhibition of granule cells, in a

randomly selected field. The same result is obtained regardless of the shape of the

sampled distribution. Ensembles convert input values to synchronised inhibition of

granule cells in the middle field.

3.4.1.2 Unknown parameters

We next explored the effect of varying model parameters for which physiological data

are unreported. We did that to verify that the conversion is not critically dependent on

our estimate of unknowns.

The results are shown in Figure 3. In 3a, the varied parameter is the number of mossy

fibres with convergent input to a single Golgi cell dendrite, and in 3b, it is the number of

Golgi cells with convergent input to a glomerulus. Both display the output of the

conversion, representing the mean of Golgi cell rates received by each glomerulus in the

field that receives ensemble output (so 700 values, the same as Figure 2, column 4). The

ratios are shown above each panel. In other figures, ratios are 4:1 (mossy fibres that

contact a Golgi cell basal dendrite) and 8–12:1 (Golgi cells that converge on a

glomerulus).

We found that increasing the convergence ratio condenses the plotted data range

(representing tighter alignment of inhibition within a field), but there is a diminishing

return. There is a stronger effect of increasing a low ratio and a progressively weakening

effect at higher ratios. The only condition that returned a large range was a very low

Golgi cell to glomerulus ratio. So: varying unknown parameters did not reveal a reason to

discard or revise the model, provided that the physiological Golgi cell to glomerulus

convergence ratio is not unexpectedly low.

Figure 3 about here please

3.4.2 A rank

We simulated a sagittal row of 100 fields, a rank. The sampled distribution, representing

mossy fibre rates received by the whole row, was generated as described in section 2, as

before. This time we took output as the mean of output values for each field (the

'glomerular field mean') – that is, for each field, the mean of values obtained the same

way as Figure 2, column 4. Within-field glomerular means are synchronised by statistical

constraint to a narrowly restricted range, so that taking the mean is a reasonable

measure to compare fields.

The results are shown in Figure 4. In panels 4A and B, the top long panel shows the mean

of mossy fibre rates received by each field in a row of 100, without copies (pink) and with

copies added ('mf field mean', red). Red data closely track the pink data, indicating that

adding copies does not skew the mean of firing rates received by a field.

The bottom long panel shows mf field mean (red) and glomerular field mean (blue), again

for 100 fields. As in Figure 2, the blue data are restricted to a focused range (the y axis

range is shorter than Figure 2). In Figure 4B, for example, the SD of output is ~3 Hz,

derived from mossy fibre rates with a range of ~190 Hz, SD 25 Hz. Within that range,

however, input and output vary randomly and independently from field to field, in a

bandwidth centred on the mean of mossy fibre rates received by the rank as a whole.

Panels 4C–E show the effect on the output of the rank conversion of changing the mean

of the sampled distribution without changing the shape or bandwidth.

Our findings are:

i) Field-to-field variability of the mf field mean is predominantly the result of the fact that

a field receives a random sample of mossy fibre rates, and not the effect of adding copies

to reflect terminal clustering;

ii) A rank of Golgi cells behaves like a single unit. Inhibition of the granule cell population

of a whole rank is synchronised. Glomerulus field means are statistically constrained to a

narrow range.

iii) The mf field mean and glomerular field mean were only related by having a narrowly

constrained range with the same mean, which was also the mean of the sampled

distribution (i.e., of mossy fibre rates received by the rank). This was unexpected. We

expected noise to be compounded by the chain of random samplings, such that a shift

(deviation from the mean) of the mf field mean would tend to cause a shift in the same

direction of the glomerular field mean. Instead, they varied independently. Indeed, the

output range was modestly narrower than the input range.

iv) The glomerular field mean is precisely and linearly coupled to the mean of the

sampled distribution.

Our interpretation is that Golgi cells work together in functionally unitary sagittal strips.

Inhibition of a rank is synchronised at an adjustable rate which is a linear function of the

mean of mossy fibre rates received by the whole rank. Synchrony is an automated

computational consequence of statistical effects of ensemble architecture.

Figure 4 about here please

3.5 Functional anatomy

The ensemble computation generates central-limit-theorem-predicted output. For that,

physiology must create conditions that simulate sampling with replacement, and which

control for other variables. Control for other variables means, the output of the

conversion must be exclusively the effect of ensemble architecture (which includes cell

morphology), and not other variables, because other variables are noise. Other variables

come in three types: mechanics of transmission that may mean it is not linear, time, and

other sources of control of Golgi cells. This section considers sampling with replacement,

linear transmission, and time.

3.5.1 Mossy fibre terminal clustering simulates sampling with replacement

Recoding exploits statistical effects (known as the central limit theorem) that depend on

replacement. Samples taken with replacement are 'independent' – there is no effect of

sampling on other samples. The physiological form of sampling is anatomical. A sample is

made up of the firing rates of cells that contact a sampling unit – in this case a Golgi cell

dendrite (not the whole cell; for granule cells, a unit is the whole cell). For independence,

each sample must be equally likely to contain any rate in the sampled distribution. This is

not anatomically achievable if mossy fibres each ended in a single terminal since clearly a

single terminal cannot be in more than one place at once. By terminating anywhere, a

cell has removed itself from the population that might contact other cells. Mossy fibre

termination in terminal clusters, and random intermingling of terminals (Fig 1), means

that a signal can be received by more than one target. Moreover, at local scale there is a

near equal probability that this happens, because a sampling unit falls within or passes

through a large majority of local cluster fields. It is a biological make-do, but the outcome

approximates sampling with replacement.

Clustering itself creates a potential problem, however. Physiological contact must

simulate sampling of mossy fires afferent to a field, not terminals, i.e., the population of

rates, not the population of signals, which includes duplicates. To approximate the result

of sampling rates only, a single sampling unit must not receive contact from the same

mossy fibre more than once. A result close to that is obtained with small sample size. It is

an imperfect solution because it does not fully eliminate duplication, but duplication is

unlikely, so rare. It is intuitive that small samples would contain duplication less often,

and we can show mathematically with granule cells that sample size of 4 is small enough.

If we assume that each dendrite of a granule cell has an equal chance of contact with

each of m x 100 mossy fibre terminals, the probability of an n^{th} contact with the same

mossy fibre is:

$$P_n = \left[\prod_{x=2}^{n} \left(\frac{m - x + 1}{100m - x + 1} \right) \right] \times \frac{y!}{x!\,(y - x)!}$$

where y is the number of granule cell dendrites and m is the number of terminals per

cluster. Calculated in this way, around 95% of granule cells receive input to each dendrite

from a different mossy fibre, and less than two in a thousand receive more than one

duplicate of the same signal.

So: the conversion of mossy fibre rates into inhibition of granule cells in functionally

unitary strips depends on anatomical mimicry of sampling with replacement. Sampling

units are single Golgi cell dendrites, not the whole cell (so the number of samples per

ensemble territory is 150 not 30). The sampled population is mossy fibres that innervate

a whole strip. Mossy fibre terminal branching and terminal clusters together have this

result. The same statements apply to granule cells with the difference that the sampling

unit is a whole cell.

3.5.2 Linear transmission

Direct evidence for or against linear conservation of rate information at each step of Golgi cell internal transmission is not available at the biophysical scale. However, evidence is available at cell and anatomical level.

Golgi cells fire autonomously. Under anaesthesia, firing falls into a slow oscillating pattern (47). Under excitatory synaptic input, however, this disappears (48). Anaesthesia 'has a strong influence on spontaneous activity of Golgi cells' (35). Discharge at low rates with irregular timing under anaesthesia (49) is replaced by higher rates with more regular timing without anaesthesia (50-52). (This paragraph is a cautionary note about evidence obtained under anaesthesia, which has been used to argue that oscillations provide a form of signalling.)

Contact on Golgi cells by mossy fibres is multi-synaptic (31), contributing to a reliable, rapid (submillisecond) response (low quantal release per synapse but collectively low uncertainty). The fast rise time and weak distance dependence of the amplitude and timing of Golgi cell EPSCs evoked by mossy fibre signals, suggests that basal dendrites introduce relatively little filtering (44, 45), as expected for large-diameter dendrites.

Contact is in glomeruli. A slow decay component associated with transmission by spillover is weak or absent, suggesting a modest spillover component of transmission. Possible reasons are individually modest release probability (53) and lower-affinity AMPA receptors (54).

'Sensory-evoked Golgi-cell inhibition [of a granule cell] scales proportionally with the

level of mossy fiber excitatory synaptic input' to the same cell, such that inhibition

reliably conveys mossy fibre rate information (43). Golgi cell 'firing frequency increases

linearly with the amplitude of depolarising current' (46 p.845). 'Golgi cells can follow

peripheral signals in a continuous fashion, modulating their frequency with the intensity

of the stimulus [citing (50, 51)]' (55). As a result, during movement, the 'output of Golgi

cells varies smoothly as a function of input.…They are capable of modulating their output

to reflect their input with little noise or signal distortion' (56).

So: mossy fibre rate information is conserved proportionally in Golgi cell firing rates. We

extrapolate that the steps in between are also linear, because it is more likely than a

linear outcome of non-linear steps.

There is a further stage in the pathway to inhibition of granule cells. Golgi cells provide

the only inhibitory input to granule cells. As noted, inhibition is almost exclusively by

spillover (39-41). Each glomerulus receives innervation by an independent random

sample of Golgi cells afferent to the field. Should spillover provide the physiological

equivalent of averaging afferent rates by diffusion in the intraglomerular space, as we

claim, linearity is preserved.

3.5.3 Time

We simulate a moment in time, a snapshot of concurrent activity. Our parameters are

linearly conserved rate information and anatomy. That is not a simplification – we argue

that the physiological computation is controlled for other variables, which are noise. It is

intuitive to think of signals as discrete episodes with a duration. However, that would

mean the combined effect of a set of signals would still include time-variable elements,

even in a snapshot, because integration would partly depend on their relative timing and

duration. We propose that the cerebellum side-steps that problem by primarily using

continuous variables for internal signalling, and that Golgi cells and glomeruli are in a

sustained state during periods of activity.

During behaviour in freely-moving animals, mossy fibre activity is dense: a high fraction

of mossy fibres are active (57). Both mossy fibres (30, 58, 59) and Golgi cells (50) have

been reported to fire with a sustained, time-varying signature, though it can depend on

modality and the manner of stimulation (60). This would suggest – and we propose – that

in the behaving animal Golgi cells receive time-varying but sustained input signals. The

large Golgi cell soma, where dendritic signals are integrated, is in this contention in a

state of continuously variable change. We note that the pause in Golgi cell firing

following a discrete stimulus under anaesthesia (43, 60, 61) disappears from Golgi cell

recordings during locomotion (50). Golgi cells fire spontaneously (49), at a rate that is

modulated by somatic charge, so that downstream, a glomerulus receives convergent

input at continuously variable rates. As a result, there is a sustained build-up of

glomerular GABA during behaviour (62) at an adjustable concentration controlled by the

mean of afferent Golgi cell firing rates (63).

On the face of it, glomeruli could cause a timing issue. Signalling by spillover is

sometimes assumed to be ambient and slow. However, the action of glomerular GABA

spillover has a fast phasic component – not as fast as synaptic transmission (∼1 ms) but

with a rise time of only a few milliseconds (63). The response time to a changing rate of

multiple active inputs should be faster than the experimentally isolated response to a

discrete signal. Unlike the spiky appearance of synaptically-induced inhibitory

postsynaptic currents, spillover (43) generates a sustained outward current.

Our interpretation is that glomeruli do not cause a special timing issue. A sustained state

of biophysical variables may provide a physiological solution to problems (extra variables)

that would be caused by coding data in discrete signals.

694   4. GRANULE CELLS

695   4.1 Contents

We next incorporated the Golgi cell ensemble conversion into a model that recodes

mossy fibre signals granule cell signals. It was first proposed over fifty years ago that

inhibition might be used to select (and change) which permutation of mossy fibre inputs

to a granule cell makes it fire (64) – so the same cell can code information combined in

different ways – or to adjust the number of inputs needed for a granule cell to fire (65),

as part of a mechanism that uses parallel fibre feedback to regulate the percentage that

are active.

We propose something functionally and mechanistically different. The function is to

maintain at a fixed number the minimum number of mossy fibre signals needed for a

single granule cell to fire, by auto-adjusting inhibition in step with the sampled

distribution. The sampled distribution is the population of mossy fibre rates received by a

whole rank. The function is to pixelate information encoded in granule cell signals. Pixels

are rank sized, long thin strips a hundred times longer than they are wide (or more). A

pixel is the smallest unit of granule cell-coded information, but not the smallest unit of

code. It is not necessary to receive the whole output of a rank to receive the information.

The same information is coded in any random sample of signals. Moreover, it can be

coded in a small fraction of the granule cell population, allowing the code to be 'sparse',

although we think not for the traditionally expected reasons. Primary auto-regulation

does not use feedback.

We describe the physiological mechanism and test it *in silico*. We take the same

approach as section 3, describing the physiology to explain the derivation of

computationally relevant variables and the values they take. We generate mossy fibre

rates as before and again start with a single field, and then simulate recoding by 100

fields to test whether they all return the same output, i.e., ranks recode as a unit.

Note: the idea that there may be an all-or-nothing result of the intraglomerular 'balance'

of excitation and inhibition (section 4.3) has appeared previously (66). The idea that

active granule cells may accordingly receive input rates with a narrowed bandwidth at

the higher end of the range also appears there.

4.2 Anatomy

The Golgi cell ensemble conversion forms part of the mechanism of granular layer

recoding. This section explains the derivation of additional parameters.

Granule cells receive input to each dendrite from a single mossy fibre. Granule cells have

3–5 dendrites, averaging 4 (the simulation assumes 4, but the result is the same with 3–

5). Contact is made in a glomerulus, as noted. Each dendrite of a single granule cell

extends into a different glomerulus. Estimates vary of the number of granule cells that

receive contact from a single terminal. 50 is mid-range (26, 27), although estimates vary

(25). 50 is consistent with our earlier estimates of the number of terminals and granule

cells in a field. A granule cell receives contact from a random sample of mossy fibres

afferent to a field, and therefore receives a random sample of mossy fibre rates. The

sampling unit is the whole cell since it receives only a single input rate to each dendrite.

Sample size is therefore 4, and the number of samples per field is the number of granule

cells, which we estimate above at 8,750.

4.3 The granule cell computation

A granule cell needs a minimum number of active mossy fibre inputs to fire – thought to be three (29, 67). Certainly, one is too few (39, 68, 69). We take it as three. Three signals is not in itself sufficient, as each of those signals must also prevail over spillover-mediated Golgi cell inhibition. Inhibition is provided by the Golgi cell ensemble conversion, so it is synchronised in a rank. It is not exactly synchronous. A single granule cell receives inhibition of each dendrite from an independent random sample of Golgi cells afferent to the field. Unitary recordings confirm that a granule cell receives inhibition from multiple Golgi cells (43). Nonetheless, it is in a focussed range so that all mossy fibre signals compete against inhibition in a narrow range.

In each glomerulus, excitation and inhibition compete for influence. Activation of presynaptic $GABA_B$ receptors on mossy fibres inhibits glutamate release (70), while activation of presynaptic mGluR2 receptors on Golgi cells inhibits GABA release (71). The outcome of the competition is independent of other mossy fibre terminals, even other terminals that provide input to the same granule cell. It is unclear if there is always an emphatic winner, but it is possible, because a swing in the dominant direction may be amplified by positive feedback. So even if all granule cells receive four active mossy fibre inputs, which we assume they do, only a subset fire, those that receive at least three mossy fibre winners.

There is a single round of sampling, by granule cells. We term the mean of Golgi cell rates

received by a single glomerulus the 'inhibitory threshold' and take it that a mossy fibre

signal must exceed the inhibitory threshold to count towards the minimum number

needed for a granule cell to fire. The simulation compares mossy fibre signal frequency

with the inhibitory threshold at each of 4 dendrites of each of 8,750 granule cells in each

of 100 fields. As discussed, there are 700 outputs of the ensemble conversion per field,

one for each glomerulus. Excitation and inhibition are paired at random. The comparison

is used to identify the subset of granule cells that fire (cells that receive at least 3 mossy

fibre signals which exceed the inhibitory threshold). The simulation takes the mean of

input rates received by each of those cells. Output of the model is accordingly the

number of granule cells that fire in each field, the mean of input rates they each receive,

and the average for each field ('GC field mean'). The results are shown in Figures 5 and 6.

4.4 Results

We find:

i) The frequency distribution of the means of mossy fibre rates received by each granule

cell is narrower than the sampled distribution, normal, and has the same mean. The

distribution of the means of active granule cells is still narrower because they receive

mossy fibre rates at the higher end of the range (Fig 5 column 2). This typically holds

even in a single field when only a very small percentage of granule cells fire ($\sim$1% active)

(Fig 5 column 2 row 6, Fig 6 column 2).[3]

ii) A sagittal row of fields which all sample the same population of mossy fibre rates, all

return the same output (Fig 5 column 3, Fig 6 column 3). The same output means, the

granule cell frequency distribution has the same mean, spread and shape. Accordingly,

any random sample of the entire output of a rank has the same statistics.

iii) (i) and (ii) are independent of the sampled distribution (mossy fibre input rates to a

rank) although a very wide mossy fibre range weakens the granule cell focus.

iv) The mean of the granule cell distribution follows the mean of the sampled

distribution. Given a fixed shape and bandwidth of the mossy fibre distribution, it follows

at a fixed distance (Fig 6 rows 1–3). Given a fixed shape and mean, increasing the mossy

fibre range causes the same relative increase in the granule cell mean (Fig 6 rows 4–6).

Our interpretation is that recoding pixelates multidimensional mossy fibre input,

converting it into a low-resolution internal code. Pixels are rank sized – there is no finer

grade of resolution. Any random sample of signals inside pixel boundaries (over a modest

minimum) codes the same information. There is a statistically conserved linear and

precise relationship between the mean of the sampled distribution and the mean of

firing rates of randomly sampled granule cells. Auto-modulation of Golgi cell inhibition of

---

[3] The focus of the granule cell range depends on the range of the sampled distribution (Fig 6 column 2 rows 4–6).

granule cells means the number of granule cells that fire is independent of mossy fibre

rates. It also regulates the number of co-active granule cells without feedback.

Figures 5 and 6 about here

4.5 Functional anatomy

The same comments apply here as the commentary on Golgi cells in section 3.5. The

treatment here is briefer because some of it is already covered.

If granule cells collectively code mossy fibre rate information in the form we propose,

weighted mossy fibre to granule cell transmission would interfere, so we would expect

transmission to be adapted to conserve rate information. The short and equal length of

granule cell dendrites would suggest light and equal filtering. The mossy fibre-granule

cell connection has multiple adaptations to transmit high frequency signals (72) across a

wide bandwidth. Faithful transmission of rate information is reported (25, 73). Vesicle

release and replenishment are fast (74) and postsynaptic AMPA receptors operate in

their linear range (75), where they are resistant to desensitisation (54). Conversion of

depolarising somatic charge into granule cell firing is a simple function such that granule

cells 'have a relatively linear and uncomplicated conversion of depolarisation level to

spike rate' (69 p.2393, citing Jorntell and Ekerot 2006 and D'Angelo et al 1998).

However, linear transmission is contentious because it has been reported that

transmission strength may be variable. There is variation between granule cells (and

trials) of EPSCs following mossy fibre stimulation *in vitro* (75), leading to the suggestion

that fast vesicular release may allow precise signalling with heterogeneous synaptic

weights. However, in this study stimulation was typically not above 2 Hz to avoid short-

term depression observed at higher frequencies, and GABA was blocked.

An alternative interpretation is that during behaviour, at behavioural mossy fibre rates,

there is glomerular build-up of intraglomerular glutamate, and signalling by spillover

becomes important, as it is for inhibitory signals (39, 63). Spillover from multiple release

sites (200-400 at a single mossy fibre terminal) is plausibly sufficient to mitigate

variability of vesicular size, release probability, and the number of release points at

synapses made on any individual cell. Indeed, short-term synaptic plasticity may have the

function of increasing the relative influence of spillover.

Still, there *are* weights. Different strength and short-term dynamics of the mossy fibre to

granule cell connection have been reported in vestibular slices, with $GABA_A$ receptors

blocked. The amplitude of the response, measured at the soma, is reported to be

statistically correlated to the source of the mossy fibre that is stimulated (76), suggesting

that the strength of the connection depends on the source.

However, the presence of weights does not necessarily mean that the function is to

modulate firing rates. Indeed, the authors of the study don't suggest that, but propose

that it may temporally code the source of inputs because different combinations have a

different response onset time. An alternative, which they mention, is that some inputs

may be in a supporting role, reminiscent of 'driver' and 'modulatory' signals in the

thalamus and cortex (77). Feasibly, signals in a supporting role must be present for the

postsynaptic granule cell to fire, but do not affect the granule cell firing rate.

The evidence is inconclusive. There is plausible evidence for linear transmission.

4.6 Physiology implements a binomial function

While mossy fibres are densely active, as noted, they may not all be active, and the

permutation of active and inactive cells may change with time. Indeed, doubtless during

behaviour it is in a state of flux. This does not impact function because the effect of silent

cells on the outcome of recoding is simply the same as the effect of weak signals.

Regulation by inhibition of the number of granule cells that fire has the same

computational result as silencing most mossy fibres, leaving a minority active. The 'silent'

cells are the ones that fire at rates below the inhibitory threshold – the losers in the

glomerular competition. The probability that a given cell fires depends on the percentage

of winners. Inhibition controls the probability by controlling the percentage. Because the

number of granule cells in a field is large, the number of active and inactive granule cells

is very near the expected proportions.

The proportions are given by a binomial function. In that sense, physiology implements a

binomial distribution. The expected number of granule cells that fire in a field is given by

$$E_{fld} = T \sum_{x=j}^{n} \left(\binom{n}{x} [P(e > i)]^x [1 - P(e > i)]^{n-x} \right)$$

where T is the total number of granule cells per field, n is the number of dendrites per

granule cell, and j is the minimum number that must receive excitation that prevails over

inhibition, given by $e > i$.

The simulation does not include a binomial function, it derives output from input by

comparing excitatory to inhibitory input to each glomerulus, glomerulus by glomerulus,

the way it is derived physiologically. The reason we mention the binomial formula is that

it is part of the functional context of synchronised inhibition that tracks mossy fibre rates.

As noted, there is a long-standing proposal that Golgi cell control of the number of

effective inputs needed to make a granule cell fire might be used to regulate the density

of parallel fibre signals activity (65). We counter propose that the opposite is necessary –

the minimum number of effective inputs must be unvarying for control of the number

that fire by probability. The primary controller is feedforward inhibition of granule cells

(and not parallel fibre feedback) for that reason: so that inhibition tracks mossy fibre

rates.

It is worth stating that a minimum number of effective inputs is also minimum sample

size, which is low anyway. For that reason, it is also safeguards the statistically reliable

encoding of mossy fibre rate information in granule cell firing.

5. DISCUSSION

5.1 Reprise

Network models often use acquired synaptic modification to implement computations.

For example, learning algorithms use synaptic change to store experience. In the

cerebellum, that has meant theories did not need to explain how granule cells code

information, because control of Purkinje cells by rates was displaced by a learned

response. We challenge that by proposing another way that input data are computed,

and how physiology is adapted for that function, focussing on the granular layer of the

cerebellar cortex.

We propose that Golgi cells work together as functional units – ensembles. The function

of ensembles is to convert mossy fibre input to a 3-field strip into synchronous inhibition

of granule cells in the middle field. The function of inhibition is to participate in recoding

which converts mossy fibre signals into granule cell signals. Ensembles are not discrete

but represent a minimum functional scale. Inhibition is synchronised between fields in a

rank, so that granule cells in a rank receive synchronised inhibition at an adjustable rate

which is a linear function of the mean of the sampled population (i.e., of mossy fibre

input rates to the whole rank).

The ensemble conversion forms part of a model of recoding that converts mossy fibre

rate information into a well-predicted granule cell code. Since at any moment all fields

randomly sample the same population of mossy fibre rates, and receive synchronised

inhibition, they all return the same output code. That is, recoding is unitised. The output

of a rank is the smallest unit of granule cell encoded information. Resolution is low: a

rank is a long thin strip of the granular layer. The mean firing rate of granule cells in each

field is a linear function of the mean of the sampled distribution. As all fields return the

same code, and information is collective coded in statistics, the same information is

coded in firing of *any* random sample of granule cells in a rank.

5.2 Pixels offer a substrate of control of Purkinje cells

We model a moment in time, a snapshot of a state of flux, in a single rank. The

cerebellum receives a torrent of data during movement. The region that provides parallel

fibre input to a microzone contains around 40 ranks, 20 each side, giving rise to parallel

fibres that intermingle in the molecular layer. We do not include other ranks.

With that caveat, granule cell code in this form provides a plausible substrate to drive

coherent firing of microzone-grouped cells downstream which populate a parallel strip. A

microzone is around 100 Purkinje cells long and 4 wide, and lies at right angles across the

path of parallel fibres, as noted. An estimated 350,000 parallel fibres pass through a

single Purkinje cell territory of which around half make contact (14, 15). So, all Purkinje

cells in a microzone receive contact from a different random sample of parallel fibres.

Most of them receive none of the same signals. Even two Purkinje cells lying directly on

beam receive contact from only (a number near to) half the same parallel fibres.

However, even if only a small fraction of parallel fibres are active, many make contact. If

0.5% are active, a single Purkinje cell receives ~875 signals, enough to code information

as statistics. Since the output of recoding is pixelated, and the same information is

contained in any random sample of granule cell signals inside (nominal) pixel boundaries,

and Purkinje cells randomly sample granule cell signals, all Purkinje cells receive the same

code, with the same timing.

5.3 A sparse granule cell code is economic and quiet

The recoding mechanism also provides regulation of the fraction of active granule cells.

Auto-modulation of inhibition decouples the fraction from mossy fibre rates. Without

adjustable inhibition of granule cells by Golgi cells, stronger mossy fibre rates would

mean that fewer mossy fibre signals would be needed to make a granule cell fire. So, if

rates went up, more would fire. In learning models, low density of active granule cells

increases storage capacity for pattern memory. In our model, Purkinje cells are

controlled by rates, not memory, so we have no use for the traditional reason. There are

other reasons for a low and stable number, however.

A low fraction saves energy. Granule cells account for more than half the cells in the

brain. The axons of granule cells are unmyelinated. The energy cost of transmission is

therefore higher than myelinated axons. A sparse code is accordingly potentially a large

energy saving. Granule cells fire in short bursts. A single burst is at a single firing rate.

Modulation of firing of parallel fibre targets is achieved by the turnover of parallel fibre

rates. A low fraction of granule cells firing in short bursts is a large saving.

Stability mitigates noise. Collective neural activity contains multiple variables all

potentially with a competing influence on downstream targets. Non-coding variables are

noise, each potentially capable of corrupting transmission. If nothing is coded in the

fraction of active granule cells, fluctuation is noise. A stable regulated percentage of

active granule cells therefore importantly eliminates a redundant variable.

Stability may be enhanced by granule cell ascending axon contact on Golgi cell basal

dendrites (45). A Golgi cell receives ~400 functional synaptic contacts from this source,

around 25% of the granule cells that a Golgi cell receives contact from in total. 'During

sustained mf [mossy fibre] discharge, synapses from local GrCs [granule cells] contribute

equivalent charge to GoCs [Golgi cells] as mf synapses' and 'display similarly fast kinetics'

(78), providing the substrate for potent local feedback that should further narrow the

range of the glomerular means both within and between fields.

5.4 Do Golgi cells receive a significant influence of excitatory or inhibitory

input from other sources?

Further regulation may be provided by feedback via parallel fibres received by dendrites

which some Golgi cells extend into the molecular layer. It is a long-standing idea that

feedback may contribute to self-regulation of the fraction of co-active parallel fibres (64,

65). Our expectation is that the strength of an effect by this pathway is modest (because

fine adjustments are sufficient for its function), consistent with a leading role in

regulation of feed-forward inhibition. Modulation of Golgi cell rates by single parallel

fibre signals is weak (46), although sub-linear integration is mitigated by gap junctions

(79). Also, only a modest number of parallel fibre signals are received at any time. A small

fraction of parallel fibres which pass through a Golgi cell territory make contact (~1,200

out of a number in the hundreds of thousands (78)). Of those, only some are active at

any time, and if the parallel fibre code is sparse, it is a small fraction. For example, if 1%

of parallel fibres are active, it is 12. This would suggest the effect of parallel fibre inputs

on Golgi cell firing is relatively weak, and control by direct input from mossy fibres is the

main form of control. Possibly, small modulations of Golgi cell somatic charge are

sufficient to make small adjustments that regulate a sparse code – that is, the number of

granule cells that need to be affected is relatively modest, so fine margins are sufficient.

As noted, not all Golgi cells have apical dendrites, so the collective effect of parallel fibre

feedback is presumably diluted because a significant population of Golgi cells don't

receive it.

Do Golgi cells receive a significant influence of input from other sources? The evidence is

incomplete but currently suggests there may not be any, or any that are significant.

Contrary to early reports, neither Purkinje cells nor climbing fibres contact Golgi cells (35,

who give references), and only a modest minority of inhibitory inputs to Golgi cells (if

any) are from molecular layer interneurons (80), which generate weak synaptic currents

(81), consistent with either extremely weak or wholly absent innervation (82). There is

conflicting evidence whether Golgi cells make inhibitory synaptic contact on each other

(48, 82). The simulation does not include an effect on Golgi cell firing by other sources of

input.

974 5.5 High-level problem solving: a comparison of the present proposals and

the supervised learning model

Physiological neural networks face some high-level problems which therefore neural

network models share. The concluding part of the Discussion compares our proposed

solution of those problems with the supervised learning model solution, the name of the

traditionally dominant class of cerebellar models.

1) Networks must distil output from input values. The traditional cerebellar model, and

many generic artificial neural networks, hold that computational derivation of output

from input is by modification of synaptic weights under climbing fibre tuition. However,

in the cerebellum, it has ultimately emerged that the evidence is unconvincing for that

and good for linear transmission. We propose, in conflict with the traditional model, that

the computation is mediated by cell type morphologies and network architecture, a

dynamic result of passive anatomy. As anatomy is known in detail, there is rich evidence

to provide parameters for a simulation, down to the level of biological messiness at a

local scale. Anatomy is underrated in theory. It tends to be thought of as merely a vehicle

to move data around, and therefore not evidence of anything. We propose that anatomy

has a computational function and therefore that evidence of anatomy is evidence of the

mechanism.

2) Signals must be correctly routed – not straightforward because in each layer of cells

they pass through, each signal is received by multiple targets and each of those targets

receives multiple signals, without an obvious key to the organisation of individual

connections, if any. The traditional theoretical solution is external supervision (as in

'supervised learning'). This is provided by climbing fibres which signal errors and teach

change at synapses which are active in a learning window. The model assumes that the

result of training is corrected motor output. It therefore does not matter which particular

pattern of parallel fibres is active. It also assumes that Purkinje cells individually receive

correctly routed signals, without explaining why they are organised into microzones.

We propose instead that internally encoded information has large spatial dimensions.

Unitised coding in topographically spatially-defined blocks, and in statistics, means there

is no need to route signals at a smaller scale. This is consistent with the organisation

reported in mapping studies and is a plausible substrate to drive coherent firing of

functionally-grouped Purkinje cells. It does not need supervision. In fact, we believe that

error signals are a good candidate to explain the vestibulo-ocular reflex but probably not

function of non-floccular circuits.

3) Insentient physiological networks must compute input firing rates without needing to

discriminate between signals on the basis of information they don't have – this can be

likened to receiving input values measured on different scales without knowing what the

scales are. Input to the cerebellum is eclectic and in multiple modalities. In some cases, a

single granule cell can receive multi-modal input. In the traditional model, supervision

solves this problem. It is unnecessary for the cerebellum itself to discriminate because

input patterns are selected for training by the timing of instruction signals.

Our alternative solution is that resolution of high-dimensional input data is sacrificed at a

surprisingly profligate level. The cerebellum disregards source and modality data below a

physical scale which mirrors topography. It is therefore unnecessary to discriminate

below that scale. Mapping studies show that mossy fibre signals (evoked by cutaneous

stimulation) share the topographical organisation of climbing fibres (22-24). Input data

are not normalised, as that would mean by definition that it was meaningfully converted

to a common internal scale. Rather, at sub-topographical dimensions, signal source and

type are simply discounted – they have no effect on the cerebellar computation or

therefore on output. This has the flexible advantage that networks do not need to be

adapted to the source or type of input they receive – anatomically, networks are

interchangeable, so network design can be modular.

1026

Glossary

1027

field A nominal division of the granular layer measuring 200 μm (sagittal) x 150 μm

(mediolateral), the average dimensions of a mossy fibre terminal cluster.

GC field mean The mean of input rates to active granule cells, calculated for each cell,

averaged for each field.

glomerular field mean The mean of inhibitory thresholds in a field.

mf field mean The mean of mossy fibre signal frequencies received as input to a field,

including copies.

rank A sagittal row of 100 fields.

inhibitory threshold The mean of Golgi cell rates received as convergent input to a

glomerulus. A mossy fibre signal must exceed this value to count towards the minimum

number of active inputs needed for a granule cell to fire.

1039

1040 **Grants**

1041 Funding: This work was supported by the Leverhulme Trust [grant number ECF-2022-079]

1042 (Mike Gilbert); the Swedish Research Council [grant number 2020-01468]; the Per-Eric

1043 och Ulla Schybergs Foundation [grant number 42630]; and the Crafoord Foundation

1044 [grant numbers 20180704, 20200729, 20220776, & 20230655] (Anders Rasmussen).

1045

1046 Funding sources had no involvement in the study.

1047

1048 **Acknowledgement**

1049 Dr Mike Evans, School of Mathematics, University of Leeds, LS2 9JT, UK, read an early

1050 draft of the paper and provided feedback.

REFERENCES

1.   Raymond JL, Medina JF. Computational Principles of Supervised Learning in the
Cerebellum. Annual review of neuroscience. 2018;41:233-53.

2.   Albus JS, editor The Marr and Albus theories of the cerebellum-two early models of
associative memory. Digest of Papers COMPCON Spring 89 Thirty-Fourth IEEE Computer Society
International Conference: Intellectual Leverage; 1989 Feb. 27 1989-March 3 1989.

3.   Brunel N, Hakim V, Isope P, Nadal JP, Barbour B. Optimal information storage and the
distribution of synaptic weights: perceptron versus Purkinje cell. Neuron. 2004;43(5):745-57.

4.   Fujita M. Adaptive filter model of the cerebellum. Biological cybernetics. 1982;45(3):195-
206.

5.   Dean P, Porrill J, Ekerot CF, Jörntell H. The cerebellar microcircuit as an adaptive filter:
experimental and computational evidence. Nature reviews Neuroscience. 2010;11(1):30-43.

6.   Welsh JP, Yamaguchi H, Zeng XH, Kojo M, Nakada Y, Takagi A, et al. Normal motor
learning during pharmacological prevention of Purkinje cell long-term depression. Proceedings of
the National Academy of Sciences of the United States of America. 2005;102(47):17166-71.

7.   Schonewille M, Gao Z, Boele HJ, Veloz MF, Amerika WE, Simek AA, et al. Reevaluating the
role of LTD in cerebellar motor learning. Neuron. 2011;70(1):43-50.

8.   ten Brinke MM, Boele HJ, Spanke JK, Potters JW, Kornysheva K, Wulff P, et al. Evolving
Models of Pavlovian Conditioning: Cerebellar Cortical Dynamics in Awake Behaving Mice. Cell
reports. 2015;13(9):1977-88.

9.   Isope P, Barbour B. Properties of unitary granule cell-->Purkinje cell synapses in adult rat
cerebellar slices. The Journal of neuroscience : the official journal of the Society for Neuroscience.
2002;22(22):9668-78.

10.   Ekerot CF, Jörntell H. Parallel fibre receptive fields of Purkinje cells and interneurons are
climbing fibre-specific. The European journal of neuroscience. 2001;13(7):1303-10.

11.   Oscarsson O. Functional units of the cerebellum - sagittal zones and microzones. Trends
in Neurosciences. 1979;2:143-5.

12.   Garwicz M, Ekerot CF, Jorntell H. Organizational principles of cerebellar neuronal
circuitry. News in Physiological Sciences. 1998;13:28-34.

13.   Ozden I, Sullivan MR, Lee HM, Wang SS. Reliable coding emerges from coactivation of
climbing fibers in microbands of cerebellar Purkinje neurons. The Journal of neuroscience : the
official journal of the Society for Neuroscience. 2009;29(34):10463-73.

14.   Harvey RJ, Napper RM. Quantitative study of granule and Purkinje cells in the cerebellar
cortex of the rat. The Journal of comparative neurology. 1988;274(2):151-7.

15.   Harvey RJ, Napper RM. Quantitative studies on the mammalian cerebellum. Progress in
neurobiology. 1991;36(6):437-63.

16.   Brand S, Dahl AL, Mugnaini E. The length of parallel fibers in the cat cerebellar cortex. An
experimental light and electron microscopic study. Experimental brain research. 1976;26(1):39-
58.

17.   Mugnaini E. The length of cerebellar parallel fibers in chicken and rhesus monkey. The
Journal of comparative neurology. 1983;220(1):7-15.

18.   Wu HS, Sugihara I, Shinoda Y. Projection patterns of single mossy fibers originating from
the lateral reticular nucleus in the rat cerebellar cortex and nuclei. The Journal of comparative
neurology. 1999;411(1):97-118.
19.   Sultan F, Heck D. Detection of sequences in the cerebellar cortex: numerical estimate of
the possible number of tidal-wave inducing sequences represented. Journal of physiology, Paris.
2003;97(4-6):591-600.
20.   Shinoda Y, Sugihara I. Axonal Trajectories of Single Climbing and Mossy Fiber Neurons in
the Cerebellar Cortex and Nucleus. In: Manto M. SJD, Rossi F., Gruol D.L., Koibuchi N., editor.
Handbook of the Cerebellum and Cerebellar Disorders: Springer, Dordrecht; 2013.
21.   Shinoda Y, Sugihara I, Wu HS, Sugiuchi Y. The entire trajectory of single climbing and
mossy fibers in the cerebellar nuclei and cortex. Progress in brain research. 2000;124:173-86.
22.   Apps R, Garwicz M. Anatomical and physiological foundations of cerebellar information
processing. Nature reviews Neuroscience. 2005;6(4):297-311.
23.   Apps R, Hawkes R. Cerebellar cortical organization: a one-map hypothesis. Nature
reviews Neuroscience. 2009;10(9):670-81.
24.   Ji Z, Hawkes R. Topography of purkinje cell compartments and mossy fiber terminal fields
in lobules ii and iii of the rat cerebellar cortex: Spinocerebellar and cuneocerebellar projections.
Neuroscience. 1994;61(4):935-54.
25.   Ritzau-Jost A, Delvendahl I, Rings A, Byczkowicz N, Harada H, Shigemoto R, et al. Ultrafast
action potentials mediate kilohertz signaling at a central synapse. Neuron. 2014;84(1):152-63.
26.   Gao Z, Proietti-Onori M, Lin Z, Ten Brinke MM, Boele HJ, Potters JW, et al. Excitatory
Cerebellar Nucleocortical Circuit Provides Internal Amplification during Associative Conditioning.
Neuron. 2016;89(3):645-57.
27.   Jakab RL, Hámori J. Quantitative morphology and synaptology of cerebellar glomeruli in
the rat. Anatomy and embryology. 1988;179(1):81-8.
28.   Eccles JC, Ito M, Szentágothai J. The cerebellum as a neuronal machine. Berlin, New York
etc.: Springer-Verlag; 1967. 335 p. p.
29.   Jörntell H, Ekerot CF. Properties of somatosensory synaptic integration in cerebellar
granule cells in vivo. The Journal of neuroscience : the official journal of the Society for
Neuroscience. 2006;26(45):11786-97.
30.   van Kan PL, Gibson AR, Houk JC. Movement-related inputs to intermediate cerebellum of
the monkey. Journal of neurophysiology. 1993;69(1):74-94.
31.   Palay SL, Chan-Palay V. Cerebellar cortex : cytology and organization. Berlin: Springer;
1974.
32.   Barmack NH, Yakhnitsa V. Functions of interneurons in mouse cerebellum. The Journal of
neuroscience : the official journal of the Society for Neuroscience. 2008;28(5):1140-52.
33.   Llinás RR, Negrello MN. Cerebellum 2015 [Available from:
http://www.scholarpedia.org/article/Cerebellum.
34.   Lange W. Regional differences in the distribution of golgi cells in the cerebellar cortex of
man and some other mammals. Cell and tissue research. 1974;153(2):219-26.
35.   Pietrajtis K, Dieudonné S. Golgi Neurons. In: Manto M, Schmahmann JD, Rossi F, Gruol
DL, Koibuchi N, editors. Handbook of the Cerebellum and Cerebellar Disorders. Dordrecht:
Springer Netherlands; 2013. p. 829-52.
36.   Eccles JC, Llinás R, Sasaki K. Golgi cell inhibition in the cerebellar cortex. Nature.
1964;204:1265-6.

37.   Hámori J, Szentágothai J. Participation of Golgi neuron processes in the cerebellar
glomeruli: an electron microscope study. Experimental brain research. 1966;2(1):35-48.
38.   Rossi DJ, Hamann M. Spillover-mediated transmission at inhibitory synapses promoted by
high affinity alpha6 subunit GABA(A) receptors and glomerular geometry. Neuron.
1998;20(4):783-95.
39.   Duguid I, Branco T, London M, Chadderton P, Hausser M. Tonic inhibition enhances
fidelity of sensory information transmission in the cerebellar cortex. The Journal of neuroscience
: the official journal of the Society for Neuroscience. 2012;32(32):11132-43.
40.   Nusser Z, Sieghart W, Somogyi P. Segregation of different GABAA receptors to synaptic
and extrasynaptic membranes of cerebellar granule cells. The Journal of neuroscience : the
official journal of the Society for Neuroscience. 1998;18(5):1693-703.
41.   Brickley SG, Cull-Candy SG, Farrant M. Development of a tonic form of synaptic inhibition
in rat cerebellar granule cells resulting from persistent activation of GABAA receptors. The
Journal of physiology. 1996;497 (Pt 3):753-9.
42.   D'Angelo E, Solinas S, Mapelli J, Gandolfi D, Mapelli L, Prestori F. The cerebellar Golgi cell
and spatiotemporal organization of granular layer activity. Frontiers in neural circuits. 2013;7:93.
43.   Duguid I, Branco T, Chadderton P, Arlt C, Powell K, Hausser M. Control of cerebellar
granule cell output by sensory-evoked Golgi cell inhibition. Proceedings of the National Academy
of Sciences of the United States of America. 2015;112(42):13099-104.
44.   Kanichay RT, Silver RA. Synaptic and cellular properties of the feedforward inhibitory
circuit within the input layer of the cerebellar cortex. The Journal of neuroscience : the official
journal of the Society for Neuroscience. 2008;28(36):8955-67.
45.   Cesana E, Dieudonné S, Isope P, Bidoret C, D'Angelo E, Forti L, editors. Novel granule cell-
Golgi cell excitatory input in the cerebellar granular layer. FENS Abstr; 2010.
46.   Dieudonné S. Submillisecond kinetics and low efficacy of parallel fibre-Golgi cell synaptic
currents in the rat cerebellum. The Journal of physiology. 1998;510 (Pt 3):845-66.
47.   Ros H, Sachdev RN, Yu Y, Sestan N, McCormick DA. Neocortical networks entrain
neuronal circuits in cerebellar cortex. The Journal of neuroscience : the official journal of the
Society for Neuroscience. 2009;29(33):10309-20.
48.   Vervaeke K, Lorincz A, Gleeson P, Farinella M, Nusser Z, Silver RA. Rapid
desynchronization of an electrically coupled interneuron network with sparse excitatory synaptic
input. Neuron. 2010;67(3):435-51.
49.   Ruigrok TJ, Hensbroek RA, Simpson JI. Spontaneous activity signatures of morphologically
identified interneurons in the vestibulocerebellum. The Journal of neuroscience : the official
journal of the Society for Neuroscience. 2011;31(2):712-24.
50.   Edgley SA, Lidierth M. The discharges of cerebellar Golgi cells during locomotion in the
cat. The Journal of physiology. 1987;392:315-32.
51.   Miles FA, Fuller JH, Braitman DJ, Dow BM. Long-term adaptive changes in primate
vestibuloocular reflex. III. Electrophysiological observations in flocculus of normal monkeys.
Journal of neurophysiology. 1980;43(5):1437-76.
52.   Rasmussen A, Zucca R, Jirenhed DA, Johansson F, Ortenblad C, Svensson P, et al. Golgi
cell activity during eyeblink conditioning in decerebrate ferrets. Cerebellum (London, England).
2014;13(1):42-5.
53.   DiGregorio DA, Nusser Z, Silver RA. Spillover of glutamate onto synaptic AMPA receptors
enhances fast transmission at a cerebellar synapse. Neuron. 2002;35(3):521-33.

54.   DiGregorio DA, Rothman JS, Nielsen TA, Silver RA. Desensitization properties of AMPA
receptors at the cerebellar mossy fiber granule cell synapse. The Journal of neuroscience : the
official journal of the Society for Neuroscience. 2007;27(31):8344-57.
55.   Galliano E, Mazzarello P, D'Angelo E. Discovery and rediscoveries of Golgi cells. The
Journal of physiology. 2010;588(Pt 19):3639-55.
56.   Moore JW, Blazis DEJ. 11 - Simulation of a Classically Conditioned Response: A Cerebellar
Neural Network Implementation of the Sutton–Barto–Desmond Model. In: Byrne JH, Berry WO,
editors. Neural Models of Plasticity: Academic Press; 1989. p. 187-207.
57.   Hartmann MJ, Bower JM. Tactile responses in the granule cell layer of cerebellar folium
crus IIa of freely behaving rats. The Journal of neuroscience : the official journal of the Society for
Neuroscience. 2001;21(10):3549-63.
58.   Arenz A, Silver RA, Schaefer AT, Margrie TW. The contribution of single synapses to
sensory representation in vivo. Science (New York, NY). 2008;321(5891):977-80.
59.   Giovannucci A, Badura A, Deverett B, Najafi F, Pereira TD, Gao Z, et al. Cerebellar granule
cells acquire a widespread predictive feedback signal during motor learning. Nature
neuroscience. 2017.
60.   Holtzman T, Rajapaksa T, Mostofi A, Edgley SA. Different responses of rat cerebellar
Purkinje cells and Golgi cells evoked by widespread convergent sensory inputs. The Journal of
physiology. 2006;574(Pt 2):491-507.
61.   Vos BP, Volny-Luraghi A, De Schutter E. Cerebellar Golgi cells in the rat: receptive fields
and timing of responses to facial stimulation. The European journal of neuroscience.
1999;11(8):2621-34.
62.   Rossi DJ, Hamann M, Attwell D. Multiple modes of GABAergic inhibition of rat cerebellar
granule cells. The Journal of physiology. 2003;548(Pt 1):97-110.
63.   Mapelli L, Solinas S, D'Angelo E. Integration and regulation of glomerular inhibition in the
cerebellar granular layer circuit. Frontiers in cellular neuroscience. 2014;8:55.
64.   Marr D. A theory of cerebellar cortex. The Journal of physiology. 1969;202(2):437-70.
65.   Albus JS. A Theory of Cerebellar Function. Mathematical Biosciences. 1971;10(1-2):25-61.
66.   Gilbert M, Miall RC. How and Why the Cerebellum Recodes Input Signals: An Alternative
to Machine Learning. Neuroscientist. 2021a;DOI: 10.1177/1073858420986795.
67.   Billings G, Piasini E, Lorincz A, Nusser Z, Silver RA. Network structure within the cerebellar
input layer enables lossless sparse encoding. Neuron. 2014;83(4):960-74.
68.   Chadderton P, Margrie TW, Hausser M. Integration of quanta in cerebellar granule cells
during sensory processing. Nature. 2004;428(6985):856-60.
69.   Bengtsson F, Jörntell H. Sensory transmission in cerebellar granule cells relies on similarly
coded mossy fiber inputs. Proceedings of the National Academy of Sciences of the United States
of America. 2009;106(7):2389-94.
70.   Mitchell SJ, Silver RA. GABA spillover from single inhibitory axons suppresses low-
frequency excitatory transmission at the cerebellar glomerulus. The Journal of neuroscience : the
official journal of the Society for Neuroscience. 2000a;20(23):8651-8.
71.   Mitchell SJ, Silver RA. Glutamate spillover suppresses inhibition by activating presynaptic
mGluRs. Nature. 2000b;404(6777):498-502.
72.   Delvendahl I, Hallermann S. The Cerebellar Mossy Fiber Synapse as a Model for High-
Frequency Transmission in the Mammalian CNS. Trends Neurosci. 2016;39(11):722-37.

73.   Rancz EA, Ishikawa T, Duguid I, Chadderton P, Mahon S, Hausser M. High-fidelity
transmission of sensory information by single cerebellar mossy fibre boutons. Nature.
2007;450(7173):1245-8.
74.   Saviane C, Silver RA. Fast vesicle reloading and a large pool sustain high bandwidth
transmission at a central synapse. Nature. 2006;439(7079):983-7.
75.   Sargent PB, Saviane C, Nielsen TA, DiGregorio DA, Silver RA. Rapid vesicular release,
quantal variability, and spillover contribute to the precision and reliability of transmission at a
glomerular synapse. The Journal of neuroscience : the official journal of the Society for
Neuroscience. 2005;25(36):8173-87.
76.   Chabrol FP, Arenz A, Wiechert MT, Margrie TW, DiGregorio DA. Synaptic diversity enables
temporal coding of coincident multisensory inputs in single neurons. Nature neuroscience.
2015;18(5):718-27.
77.   Sherman SM, Guillery RW. Distinct functions for direct and transthalamic corticocortical
connections. Journal of neurophysiology. 2011;106(3):1068-77.
78.   Cesana E, Pietrajtis K, Bidoret C, Isope P, D'Angelo E, Dieudonne S, et al. Granule cell
ascending axon excitatory synapses onto Golgi cells implement a potent feedback circuit in the
cerebellar granular layer. The Journal of neuroscience : the official journal of the Society for
Neuroscience. 2013;33(30):12430-46.
79.   Vervaeke K, Lorincz A, Nusser Z, Silver RA. Gap junctions compensate for sublinear
dendritic integration in an inhibitory network. Science (New York, NY). 2012;335(6076):1624-8.
80.   Eyre MD, Nusser Z. Only a Minority of the Inhibitory Inputs to Cerebellar Golgi Cells
Originates from Local GABAergic Cells. eNeuro. 2016;3(2).
81.   Dumoulin A, Triller A, Dieudonné S. IPSC kinetics at identified GABAergic and mixed
GABAergic and glycinergic synapses onto cerebellar Golgi cells. The Journal of neuroscience : the
official journal of the Society for Neuroscience. 2001;21(16):6045-57.
82.   Hull C, Regehr WG. Identification of an inhibitory circuit that regulates cerebellar Golgi
cell activity. Neuron. 2012;73(1):149-58.

1253

1254

A

B

Cerebellar y-axis [µm]

Cerebellar x-axis [µm]

C

Molecular layer

Granular layer

Mossy fibre

D

1255

Fig 1. Mossy fibre termination in a rank. There is random interdigitation of branch

endings of mossy fibres co-afferent to a rank, and at local level random intermingling of

terminals of mossy fibres co-afferent to a field. **(A)** Sagittal section of the rat cerebellum

(obtained by multi-photon microscopy, courtesy Tom Deerinck). The outer layer, the

molecular layer, of the cerebellar cortex is stained green and the inner layer, the granular

layer, is blue. **(B)** Output of a procedure used to group mossy fibre terminal into clusters

(19 fig 6), reproduced with permission. All terminals belong to one mossy fibre, which

terminates in two strips. Clusters are shown in different shades of grey. **(C)** Schematic of

an area of the folded cerebellar cortex. The strip inside the red dotted line is the width

and depth of a microzone but only part of its length. Mossy fibre signals are not received

at a point but by a volume of the granular layer, at several randomly spaced, sagittally-

aligned locations. Yellow zones are terminal locations of a single mossy fibre. A mossy

fibre mediolateral axon may give rise to several sagittal branches, of which here only one

is shown. Size of locations varies depending partly on the number of terminals in each

cluster. Pink zones are terminal locations of a second mossy fibre. In reality, zones

overlap extensively and terminals are intermingled. The zone enclosed by a blue line has

the dimensions of a 'field'. A field is a nominal division of the granular layer with the

average dimensions of a mossy fibre terminal cluster, which we use to quantify granular

layer computations. A sagittal row of 100 fields is a 'rank'. **(D)** Section of a microzone

viewed obliquely from the cerebellar surface, to illustrate the size and random

intermingling of mossy fibre terminals. The molecular layer is represented by squares of

about the dimensions of a Purkinje cell territory. The wire frame boxes are an exploded

view of the orange fields. The right box shows randomly distributed terminals of 5 mossy

fibres (each a different colour). We estimate that ∼180 mossy fibres contribute terminals

to a field, and a field contains a total of around 700 terminals. Proportions of terminal

size to field size are preserved. The left box shows terminals of 20 mossy fibres.

1282

1283

Fig 2. Golgi cell ensemble conversion. Column 1 We randomly generated (3,645) mossy fibre rates with a uniform, normal, twin-peaked or discontinuous distribution representing input to a row of 100 fields. Rates were randomly sampled to represent input to individual fields and copies added to reflect variable size of terminal clusters and that most clusters straddle field boundaries. Red data represent copies-included signals received by a row of three fields. **Column 2** The first-column red data were randomly sampled 150 times, sample size 4, representing mossy fibre input to a 3-field population of 30 Golgi cells, each with 5 basal dendrites (the mean),and each of those receiving contact from 4 mossy fibres. Data are the sample means, representing dendritic depolarisation. **Column 3** We next randomly sampled column 2 data without replacement 30 times, sample size 5, and took the sample means, representing somatic integration of dendritic signals. **Column 4** We then took 700 random samples of column 3 data, sample size 8–12, reflecting convergent input from 3 fields of Golgi cells onto 700 glomeruli in the middle field (to which a three-field ensemble has output). Data are the sample means. **Column 5** The column 4 data are presented as a histogram.

1300

1301

Fig 3. Testing unreported parameters shows that reliability and precision are chiefly

impacted at different steps in the computation. All data were obtained in the same way

as Fig 2 column 4, but varying ratios for which physiological parameters have not been

reported. Data points are accordingly the means of randomly sampled Golgi cell rates

received as convergent input to each of 700 glomeruli, the population of the middle field

of a Golgi cell ensemble territory. Broken dark blue line: the mean of the sample means.

The sampled distribution was generated in the same way as Fig 2 row 4. (The sampled

distribution is the population of mossy fibre input rates to a row of 100 fields which

includes the ensemble territory.) Yellow line: the mean of the sampled distribution.

Varied ratios are the convergence ratio of mossy fibres onto a single Golgi cell basal

dendrite (4:1 in Fig 2) and the convergence ratio of Golgi cells onto a single glomerulus is

(a random number in the range 8–12:1 in Fig 2). In **(A)** we varied the first and in **(B)** the

second. The ratios are shown above each panel. y axes cover a shorter range than Fig 2.

Blue data – representing inhibition of granule cells – are the output of the ensemble

transformation. A narrower range represents tighter within-field synchrony of inhibition

received by granule cells. Findings were: (1) In both cases, there is a diminishing return of

increasing the ratio. There is a modest return of higher ratios than Fig 2, selected to be

physiologically plausible. (2) The blue mean varies in a small range around the yellow

mean, but within that, varies at random each time the simulation is run. In (A), it varies

from panel to panel because mossy fibre input rates to the simulated ensemble territory

are resampled each time. In (B), that step in the computation is performed only once and

the output used for all panels, so that the effect of changing the second ratio is not

compounded with changing the first. Here, the blue mean is the same in all panels, but

moves randomly when the simulation is rerun, in the same range as (A). So, reliability

(reproducibility of the glomerular field mean) is impacted at the step of sampling of

mossy fibre rates by Golgi cells, but not sampling of Golgi cell rates by glomeruli;

reliability was not obviously affected by varying the first-step ratio (not shown). Precision

(narrowness of the glomerular range) is impacted by both ratios, but chiefly at the

second step.

1331

1332

Fig 4. A whole rank of Golgi cells work as a functional unit. Synchronised inhibition of granule cells is precisely and linearly related to the mean of mossy fibre rates. (A) and **(B)** Simulation of conversion by Golgi cells of mossy fire signals into inhibition of a sagittal row of 100 fields, taking the mossy fibre distribution on the left as input. The top panel compares the mean of mossy fibre input rates to each field (pink) with the mean with copy signals added to reflect mossy fibre terminal clustering (red). The dashed grey line is the mean of the sampled distribution (i.e., the mean of mossy fibre input rates to the whole rank). The bottom panel compares the copies added data (red) with the glomerular field mean (blue line). SD of inhibitory thresholds is shown in pale blue. The y scale of the long panels is Hz. Results are: (1) In all top panels, red data closely track the pink data, indicating that adding copies does not skew the mean of firing rates received by a field. (2) Blue data across all fields are in a narrowly circumscribed range. The degree of precision depends on the mossy fibre range although the conversion works with any input distribution. A normal mossy fibre distribution with a 2SD range of over 100 Hz is converted to a blue range of ∼3 Hz. (3) Fields that sample the same input data (that data being: mossy fibre input rates to a rank) return the same output. This has the physiological counterpart that inhibition of granule cells is synchronised in a rank. (4) The blue range is centred on the grey dashed line – the mean of mossy fibre rates received by the rank as a whole – and varies independently of the red data. That is, field-to-field variation of mean mossy fibre input rates doesn't flow through to an effect on blue data, the output of the system. (5) We also tested setting sample size 'optimally' (i.e., higher,

at 24:1) for mossy fibres that contact a single Golgi cell dendrite, and Golgi cells that

inhibit a single glomerulus. This only very modestly improved performance: higher ratios

do not return a notably narrower blue range (than the range when parameters are set at

known or expected physiological values). The results are not shown. **(C)–(E)** The same

simulation, sampling the mossy fibre distribution on the left. The blue data are the

glomerular field means, i.e., mean inhibition of granule cells in each field (inhibition of

granule cells is condensed into a small range by the ensemble conversion, so we use the

means to compare fields). y axes of panels on the right of each pair have the same range

as x axes of panels on the left. Results: inhibition of granule cells is precisely and linearly

coupled to the mean of the sampled distribution.

1364

1365

1366

1367

Fig 5. Incorporating the Golgi cell conversion into a model that recodes input to the cerebellum into internal signals. We incorporated the Golgi cell ensemble conversion into a simulation of recoding in the cerebellar granular layer. Recoding converts input to the cerebellum from mossy fibres into firing of granule cells. **Column 1.** We randomly generated (3,645) mossy fibre rates with a uniform, discontinuous or normal distribution representing input to a row of 100 fields. Red data are rates received by a single field (field 2, selected at random), with copies added, as an example. y axes are scaled to the data. **Column 2.** Red data for each field were randomly sampled 8,750 times, sample size 4, representing a per field population of 8,750 granule cells, each with 4 dendrites, which each receive contact from a single mossy fibre terminal. The field 2 sample means are displayed as a histogram (pale green, top three rows only; there are the same number of green data in each panel). Blue data represent the subgroup of granule cells that fire (all rows). In the bottom three rows the subgroup is small (y axes are rescaled) – recoding is set (by adding a constant to inhibition) where around 1% of granule cells are active. **Column 3.** The y scale is Hz. The mean (line) and SD (shading) of the red and blue data for each field in a row of 100 fields. Blue line data are GC field means. **Column 4.** The percentage of granule cells that fire in each field. The white line and shaded area in the top three panels are mean and SD respectively. Results are: (1) Blue line data across all fields are in a narrowly circumscribed range. The degree of precision depends on the mossy fibre range although the conversion works with any input distribution. (2) Fields that sample the same input data (that data being: mossy fibre input rates to a whole

rank) return the same granule cell code. That is, the frequency distribution of the mean

of input rates to each active granule cell is centred on the narrowly-varying GC field

mean and attracted to normal. (3) A whole rank recodes as a single unit. Any random

sample of granule cells has the same mean and distribution, if, as we claim, information

is collectively encoded in rate statistics of co-active cells. It is not material that fields all

return a different permutation of output rates.

1395

1396

Fig 6. Granule cell rates collectively code mossy fibre rate information. Column 1 We

randomly generated (3,645) mossy fibre rates per section 2 with either a normal (SD

12.5) or uniform distribution representing input to a row of 100 fields, with copies added

– the sampled distribution. **Column 2** Frequency distribution of the mean of mossy fibre

input rates to each of the granule cells that fire in a randomly selected field, obtained

with the recoding model in the same way as in the bottom 3 rows of Fig 5. **Column 3**

Grey line: the mean of the sampled distribution; blue line: GC field means. **Column 4** The

percentage of active granule cells in each field. **(A)–(C)** The blue data are precisely and

linearly related to the mean of mossy fibre rates. Blue data track the mean of the

sampled distribution at a mean distance of 15 Hz (SD 1.21 Hz) in all rows, i.e., the

relationship is independent of mossy fibre rates if the shape and bandwidth of the

distribution of mossy fibre rates are approximately fixed. There is also no effect of

changing mossy fibre rates on the percentage of granule cells that fire. **(D)–(F)** If instead

the shape and mean of the sampled distribution are unchanging, there is a linear

relationship with the bandwidth. Increasing the bandwidth causes the same relative

increase in the mean distance between the mossy fibre mean and GC field means. There

is no effect on the percentage of granule cells that fire.