Research Article

Genetic dissection of BDNF and TrkB expression in glial cells

Changran Niu [1,2], Xinpei Yue [1,2], Juan Ji An [1], Haifei Xu [1], Baoji Xu [1,2,*]

[1] Department of Neuroscience, The Herbert Wertheim UF Scripps Institute for Biomedical

Innovation & Technology, University of Florida, Jupiter, 33458, FL

[2] Skaggs Graduate School of Chemical and Biological Sciences, The Scripps Research Institute,

La Jolla, 92037, CA

[*] Correspondence: baoji.xu@ufl.edu; Tel: (561) 228-2340

Abstract

The brain-derived neurotrophic factor (BDNF) and its high-affinity receptor tropomyosin-

related kinase receptor B (TrkB) are widely expressed in the central nervous system. It is well

documented that neurons express BDNF and full-length TrkB (TrkB.FL), and a lower level of

truncated TrkB (TrkB.T). With conflicting results, glial cells also have been reported to express

BDNF and TrkB. In the current study, we employed a more sensitive and reliable genetic method

to characterize the expression of BDNF and TrkB in glial cells in the mouse brain. We utilized

three Cre mouse strains in which Cre recombinase is expressed in the same cells as BDNF,

TrkB.FL, or all TrkB isoforms, and crossed them to Cre-dependent EGFP reporter mice to label

BDNF- or TrkB- expressing cells. We performed immunohistochemistry with glial cell markers to

examine the expression of BDNF and TrkB in microglia, astrocytes, and oligodendrocytes.

Surprisingly, we found no BDNF- or TrkB- expressing microglia in the brain and spinal cord.

Consistent with previous studies, most astrocytes only express TrkB.T in the adult brain.

Moreover, there are a small number of astrocytes and oligodendrocytes that express BDNF, the

function of which is to be determined. We also found that oligodendrocyte precursor cells, but not

mature oligodendrocytes, express both TrkB.FL and TrkB.T in the adult brain. These results not

only clarify the expression of BDNF and TrkB in glial cells, but also open opportunities to

investigate previously unidentified roles of BDNF and TrkB in glial cells.

Keywords: BDNF, TrkB, Cre, glia, microglia, astrocyte, oligodendrocyte

1. Introduction

Brain-derived neurotrophic factor (BDNF) is a neurotrophin that is widely expressed in the

central nervous system. BDNF signals via the tropomyosin-related kinase B (TrkB) receptor to

regulate neuronal proliferation, differentiation and survival, and synaptic function [1]. Binding of

BDNF to the full-length TrkB (TrkB.FL) receptor triggers dimerization of TrkB.FL and tyrosine

autophosphorylation, leading to activation of three main downstream signaling cascades

mediated by phospholipase C γ1 (PLCγ1), phosphoinositide 3-kinase, and RAS/RAF/mitogen-

activated protein kinase, respectively [2]. Alternative splicing of *Ntrk2* exons produces the

truncated TrkB (TrkB.T) isoforms that lack nearly all the intracellular domain including the tyrosine

kinase motif [3]. TrkB.T can negatively regulate TrkB.FL signaling by forming a TrkB.FL-TrkB.T

heterodimer and inhibiting the activation of TrkB.FL [4]. TrkB.T also functions to sequester and

translocate BDNF, induce neurite outgrowth, and activate downstream protein kinase C and PLCγ

via G-protein and inhibit Rho GTPase via dissociating Rho GDP dissociation inhibitor [4]. Because

of the diverse roles of the BDNF-TrkB pathway, dysregulation of the BDNF-TrkB signaling has

been implicated in various disease conditions, including cognitive impairment, obesity,

neurodegenerative diseases, and cancer [5].

BDNF-TrkB signaling pathway has been studied extensively in neurons. It is well-

documented that neurons express BDNF, TrkB.FL and a lower level of TrkB.T. Expression of

BDNF and TrkB has also been reported in glial cells. BDNF expressed in microglia was suggested

to be involved in neuropathic pain and motor-learning dependent synapse formation [6, 7].

However, other studies showed very low or no expression of BDNF in microglia [8]. Therefore, it

is debatable whether microglia express BDNF or not. It has been shown that astrocytes

predominantly express TrkB.T [1, 9]. Whether astrocytes also express TrkB.FL is not clear yet.

Some studies have shown TrkB.FL expression in astrocytes isolated from mouse brain [10, 11],

whereas other studies show no TrkB.FL expression in astrocytes [12, 13]. Astrocytes can

internalize proBDNF secreted by neurons, convert proBDNF to mature BDNF (mBDNF) and

release mBDNF for synaptic reuse [14, 15]. Astrocytes was shown to express BDNF *in vitro* which

regulates neuronal dendrite maturation [16], but whether astrocytes express BDNF *in vivo* under

physiological conditions is unknown. Oligodendrocytes have also been shown to express BDNF,

the reduction of which impairs presynaptic vesicular exocytosis [17]. Oligodendroglial expression

of TrkB is important for oligodendrocyte myelination and oligodendrocyte precursor cell (OPC)

proliferation [18-20]. But it has not been characterized which TrkB isoform oligodendrocytes

express and what percentage of oligodendrocytes express BDNF or TrkB.

To fill these knowledge gaps, in the current study, we employed a genetic method to label

and characterize the expression of BDNF and TrkB in glial cells, including microglia, astrocytes

and oligodendrocytes. We crossed the *Bdnf*[2A-Cre/+] [21], *Ntrk2*[2A-Cre/+], and *Ntrk2*[CreER/+] [22] mouse

strains to the Cre-dependent B6;129S4-Gt(*ROSA*)*26Sor*[tm9(EGFP/Rpl10a)Amc]/J (EGFP-L10a) reporter

mice [23] to generate double heterozygous mice in which cells expressing BDNF or TrkB were

labeled by EGFP. We then performed immunofluorescence staining of cell type markers with the

brain sections from these mice. Colocalization of EGFP and cell type markers was analyzed to

determine the expression of BDNF and TrkB in glial cells. There are several advantages of this

method: (1) high sensitivity because a very small amount of Cre expression can induce EGFP

expression in the reporter mice; (2) no involvement of BDNF and TrkB antibodies which have

non-specific immunoreactivity; (3) no need to isolate glia from the mouse brain, which involves

potential contamination of other cell types; (4) being able to observe expression of BDNF and

TrkB *in vivo* at the level of single cells.

Our results show that homeostatic and activated microglia do not express either BDNF or

TrkB. Astrocytes predominantly express TrkB.T, and a small astrocyte population express

TrkB.FL and BDNF in the adult mice. A subset of oligodendrocytes express BDNF in the

hippocampal CA1, as well as TrkB.T and TrkB.FL.

2. Materials and Methods

2.1 Animals

Mice of the EGFP-L10a/+ (B6;129S4-Gt(*ROSA*)*26Sor*$^{tm9(EGFP/Rpl10a)Amc}$/J; stock #024750)

and the *Bdnf*$^{2A-Cre/+}$ (B6.FVB-*Bdnf*$^{em1)cre\_Zak}$/J; stock #030189) strains were obtained from the

Jackson Laboratory. The *Ntrk2*$^{CreER/+}$ mouse strain was generously provided by Dr. David Ginty

at Harvard Medical School [22]. *Ntrk2*$^{2A-Cre/+}$, *Ntrk2*$^{CreER/+}$, and *Bdnf*$^{2A-Cre/+}$ mice were crossed to

EGFP-L10a/+ mice to generate *Ntrk2*$^{2A-Cre/+}$;EGFP-L10a/+, *Ntrk2*$^{CreER/+}$;EGFP-L10a/+ and *Bdnf*$^{2A-}$

$^{Cre/+}$;EGFPL10a/+ mice. All experiments were performed in accordance with relevant guidelines

and regulations regarding the use of experimental animals. The Animal Care and Use Committees

at the UF Scripps approved all animal procedures used in this study.

2.2 Generation of *Ntrk2*$^{2A-Cre/+}$ mice

We generated a *Ntrk2*$^{2A-Cre}$ mouse allele in which the DNA sequence encoding P2A-Cre

recombinase was inserted immediately before the stop codon for TrkB.FL at the *Ntrk2* locus using

the CRISPR/Cas9 technique. sgRNA (5'-AGTCAAGAGGTTCGTCGTGT-3') was designed using

the CRISPR tool (http://crispr.mit.edu) to minimize potential off-target effects. The DNA donor

plasmid contains two homology arms with 840-bp flanking the P2A-Cre sequence, which were

cloned into BamHI digested pBluescript II KS (-) vector using the Gibson assembly method (NEB,

E5510). To block further Cas9 targeting and recutting after undergoing homology-direct repair,

we introduced a silent mutation into the PAM motif of the sgRNA located within the 3' homology

arm in the donor plasmid by using Q5 Site-Directed Mutagenesis kit (NEB, E0054). The donor

plasmid was confirmed with DNA sequencing. Microinjection of a mixture of sgRNA, donor DNA,

and Cas9 protein was performed by the Genomic Modification Facility at Scripps Research.

Zygotes were cultured to the blastocyst stage *in vitro*. Genomic DNA from blastocysts was

extracted and PCR screen was performed to select blastocysts with successful homologous

recombination. Positive blastocysts were transferred into the oviduct of pseudo-pregnant females

to produce founder mice. Two out of twenty-eight pups were positive for the knockin, confirmed

by the genomic DNA PCR using the following primers: CCTCCTGGTGAGCAAACGAT (forward)

and GACATAGGGCCGGGATTCTC (reverse) for PCR across the 5' homology arm, and

CACCTCCATGCCTGTGTTTT (forward) and GGTTCTTGCGAACCTCATCA (reverse) for PCR

across the 3' homology arm. The founder mice were crossed to C57BL/6J to produce mice with

germline transmission. Cre expression in these lines were confirmed by *in situ* hybridization.

2.3 In situ hybridization

Freshly dissected mouse brains were quickly frozen in a 2-methylbutane (Fisher Scientific

#O3551-4) dry ice bath. Subsequently, 14-μm thick cryostat sections were collected onto

Superfrost Plus slides (Fisherbrand #12-550-15) and used for in situ hybridization. Fluorescence

in situ hybridization was performed using the RNAscope Multiplex Fluorescent Reagent Kit v2

(#323100, Advanced Cell diagnostics). Brain sections were fixed with pre-chilled 10% formalin at

4°C for 15 min, and dehydrated with 50%, 75% and 100% ethanol. Sections were air-dried for 20

min and followed by v2Hydrogen peroxide for 10 min and pretreated with protease IV for 30 min,

washed 2 times in PBS and incubated with the target probes (Ntrk2-C3, #423611-C3; Cre-C2,

#31228-C2) for 2 hours at 40°C. Signals were amplified by subsequent incubation with v2Amp1

for 30 min, v2Amp2 for 30 min, v2Amp3 for 15 min, v2C2/C3HPR for 15 min and followed by TSA

Plus Fluorescein/Cyanine5 (NEL754001KT, Akoya Biosciences) for 30 min at 40°C. Slides were

counterstained with DAPI, and images were acquired using a Nikon C2+ confocal microscope.

2.4 Immunohistochemistry

Mice were perfused with phosphate-buffered saline (PBS) and then 4% paraformaldehyde

(PFA) in PBS. The brain was dissected and postfixed overnight in 4% PFA-PBS. Mouse brains

were soaked in 30% sucrose-PBS for cryoprotection. Brains were sectioned into 40-μm-thick

slices using a sliding microtome (Leica). Brain sections were blocked in 0.3% Triton X-100 and

10% normal horse serum before overnight incubation in primary antibody at 4 °C. The following

primary antibodies were used: guinea pig anti-Iba1(1:500; Synaptic System #234308), goat anti-

SOX9 (1:2000; R&D Systems #AF3075), mouse anti-GFAP (1:500; Cell Signaling Technology

#3670S), mouse anti-Olig2 (1:250; Millipore #MABN50), rabbit anti-ASPA (1:1500; GeneTex

#GTX113389), chicken anti-GFP (1:2000; abcam #13970), rabbit anti-DsRed (1:1000; TaKaRa

#632496). The brain sections were washed three times with PBS before incubation in secondary

antibody for 1 hour at room temperature. The brain sections were then counterstained with DAPI,

mounted onto microscope slides, and covered with mounting media.

2.5 LPS injection

LPS (Millipore #L6529) was dissolved in sterile saline to a final concentration of 0.1 μg/μL.

The mice were given daily saline or LPS injection at a dose of 500 μg/kg body weight for seven

consecutive days. The mice were harvested the next day after the last treatment.

2.6 Tamoxifen injection

Tamoxifen (Sigma #T5648-5G) was dissolved in 100% ethanol to a concentration of 20

mg/ml. An equal volume of corn oil was added to the tamoxifen solution. The corn oil/ethanol

mixture was vortexed, centrifuged for 10 mins at 13,000 rpm, and then placed in a vacuum

centrifuge for 30 mins to evaporate ethanol. Tamoxifen was given to $Ntrk2^{CreER/+}$;EGFP-L10a/+

mice at the age of 6 weeks through daily i.p. injection at a dose of 150 mg/kg body weight for 10

consecutive days to induce translocation of the Cre-ER fusion protein to the nucleus. The mice

were given three weeks to recover before they were harvested.

2.7 Plasmids and viruses

pAAV8-EF1a-Nuc-flox(mCherry)-EGFP was obtained from Addgene (plasmid #112677, a

gift from Brandon Harvey [24]). For higher expression in glial cells, the EF1a promoter was

replaced by the CAG promoter from pAAV-CAG-mNeonGreen (Addgene plasmid #99134, a gift

from Viviana Gradinaru [25]). pAAV8-EF1a-Nuc-flox(mCherry)-EGFP was digested with AgeI and

BamHI, and the CAG promoter fragment was amplified by PCR (forward primer: GAAT

ACCGGTCGTTACATAACTTACGGTAAATG;   reverse   primer:

GAATGGATCCCGCCCGCCGCGCGCTT). PCR product was digested with AgeI and BamHI and

ligated into the backbone of pAAV8-EF1a-Nuc-flox(mCherry)-EGFP. The resulting pAAV8-CAG-

Nuc-flox(mCherry)-EGFP plasmid was confirmed by sequencing and used to package AAV

viruses. AAV viruses were purified with TaKaRa AAVpro Purification Kit Maxi (#6666).

2.8 Stereotaxic injection of adeno-associated virus

Mice were anesthetized using isoflurane and securely positioned on a stereotaxic holder

(David Kopf, Model 940). A small incision was made to expose the skull, and subsequently a

small hole was drilled on the skull above an injection site. For injection, a Nanofil 33-gauge needle

(World Precision Instruments, #NF33BV-2) was slowly inserted into the CA1 region (AP: -1.60

mm; ML: ± 0.80 mm; DV: −1.80 mm). AAV8-CAG-Nuc-flox(mCherry)-EGFP (150 nl at 3.12E+12

vg/mL) was then injected at a rate of 30 nl/min using a micro syringe pump (World Precision

Instruments, SYS-MICRO4). The needle was then withdrawn after staying in the same position

for 5 min. Following the injection, mice received Loxicom (5 mg/kg) for analgesia and were

returned to their home cages.

2.9 Imaging and Cell number quantification

Fluorescent images were captured using a Nikon C2+ confocal microscope. Cell numbers

were quantified using the cell counter plugin in FIJI software.

3. Results

3.1 Homeostatic and activated microglia in the cortex, hippocampus and spinal cord do

not express BDNF

$Bdnf^{2A\text{-}Cre/+}$ mice that express Cre recombinase in BDNF-expressing cells were crossed to

Cre-dependent EGFPL10a/+ reporter mice to generate $Bdnf^{2A\text{-}Cre/+}$;EGFPL10a/+ mice. Cre-

mediated excision of *loxP*-flanked STOP sequence results in constitutive expression of EGFP

localized in the nucleus and cytoplasm of Cre-expressing cells. EGFP is thus expressed in cells

that express BDNF at any time during the lifespan of a $Bdnf^{2A\text{-}Cre/+}$;EGFP-L10a/+ mouse. We first

used NeuN as a marker to label neurons. As expected, most neurons in the cortex and

hippocampal CA1 express BDNF (Fig. 1A-F).

Next, we wanted to know whether microglia express BDNF. For this, we used Iba1

antibody to label microglia in $Bdnf^{2A\text{-}Cre/+}$;EGFP-L10a/+ mice. Surprisingly, we did not observe

EGFP expression in nuclei of Iba1$^+$ microglia in the somatomotor cortex (MO), hippocampal CA1

stratum radiatum (SR), or the spinal cord of 4-month-old $Bdnf^{2A\text{-}Cre/+}$;EGFPL10a/+ mice, indicating

that homeostatic microglia do not express BDNF under physiological conditions (Fig. 1G-O).

However, we did observe EGFP puncta in some microglial cell bodies and processes, outside

microglial nuclei (Fig. 1P-S). Immunostaining of CD68, a microglial lysosomal marker, showed

that the EGFP puncta are always colocalized with CD68 (Fig. 1P-S). This indicates that these

EGFP signals in microglia come from BDNF-expressing cells engulfed by microglia and are not

from BDNF expression in microglia.

Figure 1. Homeostatic microglia do not express BDNF. (**A-F**) BDNF expression in NeuN+ neurons in the MO (A-C) and CA1 (D-F). (**G-O**) Homeostatic microglia do not express BDNF in the MO (G-I), CA1 (J-L), and spinal cord (M-O). (**P-S**) Colocalization of EGFP puncta signals with CD68+ microglia lysosomes. Arrows denote EGFP signal inside microglial cytoplasm. Arrowheads denote EGFP signal inside microglial processes. Scale bar = 50 μm. n = 2 mice.

Microglia can exist in homeostatic or activated states. To investigate whether activated

microglia express BDNF under pathological conditions, we injected lipopolysaccharide (LPS) to

induce microglial activation in adult $Bdnf^{2A\text{-}Cre/+}$;EGFPL10a/+ mice. CD68 is used as a marker for

microglia activation, as activated microglia express a higher level of CD68. LPS treatment resulted

in a significant increase in CD68 intensity per microglia (Fig. 2). We did not observe EGFP signal

in Iba1$^+$ microglial nuclei in LPS-injected mice, suggesting that activated microglia also do not

express BDNF (Fig. 2E-H). Therefore, our results suggest that both homeostatic and activated

microglia do not express BDNF.

Figure 2. Activated microglia do not express BDNF. (**A-H**) Immunostaining of Iba1 and CD68 in the hippocampus

of $Bdnf^{2A\text{-}Cre/+}$;EGFPL10a/+ mice injected with saline (A-D) or LPS (E-H). Arrows indicate colocalization of Iba1 and

CD68, and no EGFP inside microglia. Scale bar = 50 μm. (**I**) Quantification of CD68 intensity per microglia. n = 1 35

microglia from one saline mouse and 50 microglia from 2 LPS mice . *p = 0.0204 by two-sided *t*-test. Data are shown

as mean ± s.e.m.

3.2 Some astrocytes express BDNF in the cortex and hippocampus

Whether astrocytes express BDNF *in vivo* under physiological conditions is unknown. To

answer this question, we labeled astrocytes in $Bdnf^{2A\text{-}Cre/+}$;EGFPL10a/+ mice using SOX9

antibody for the somatosensory cortex (SS) and GFAP antibody for the hippocampus. We

observed that a small portion of astrocytes (2.6% in the SS and 18.3% in the CA1 SR) are EGFP$^+$

in the *Bdnf*[2A-Cre/+];EGFPL10a/+ mice, indicating that these cells express BDNF at some point

during the lifetime of the mice (Fig. 3A-H). The EGFP signals in these mice are cumulative,

because once the EGFP expression in a cell is turned on by Cre, it is never turned off even if the

cell stops to express Cre. Therefore, we cannot know when BDNF is expressed in these mice.

Figure 3. A small subset of astrocytes express BDNF in the hippocampus. (**A-H**) Immunostaining of SOX9 (A-C)

and GFAP (E-G) in the SS (A-C) and CA1 SR (E-G) of *Bdnf*[2A-Cre/+];EGFPL10a/+ mice, and (D, H) quantification of

colocalization. (**I-M**) Immunostaining of GFAP in the CA1 SR of adult *Bdnf*[2A-Cre/+] mice injected with the mCherry-to-

EGFP color switch AAV, and (M) quantification of colocalization. Scale bar = 50 μm. Arrowheads denote EGFP-

expressing astrocytes. Arrows denote astrocytes that do not express EGFP.

To test whether astrocytes in adult mice still express BDNF, we injected a Cre-dependent

mCherry-to-EGFP color switch adeno-associated virus (AAV), AAV8-CAG-Nuc-flox(mCherry)-

EGFP, into the hippocampal CA1 of 2-month-old *Bdnf*[2A-Cre/+] mice. In these mice, transduced Cre-

negative cells will express mCherry, while transduced Cre-positive cells will express EGFP.

Around 87% astrocytes were transduced by the AAV, and 10.6% of the transduced astrocytes

express BDNF in adult mice (Fig. 3I-M). This suggests that a small portion of astrocytes in the

CA1 SR still express BDNF in adult mice. Because this number is smaller than the cumulative

18.3% in $Bdnf^{2A-Cre/+}$;EGFPL10a/+ mice, there could be two possibilities: (1) some astrocytes only

express BDNF early during development and stop BDNF expression in adults; (2) at each different

time point, there is a different small subset of astrocytes that express BDNF, and that accumulates

to the 18.3% seen in $Bdnf^{2A-Cre/+}$;EGFPL10a/+ mice. To figure out which possibility is true, future

studies can investigate BDNF expression at developmental stages.

3.3 A small portion of oligodendrocytes express BDNF in the hippocampal CA1 SR

Next, we examined BDNF expression in Olig2$^+$ oligodendrocytes. We found that most

Olig2$^+$ oligodendrocytes in the SS do not express BDNF, and 4.4% of Olig2$^+$ oligodendrocytes in

the CA1 SR express BDNF at some point during the lifespan of $Bdnf^{2A-Cre/+}$;EGFPL10a/+ mice

(Fig. 4A-H). To find out whether these oligodendrocytes express BDNF in adult mice, we injected

the mCherry-to-EGFP color switch virus to the hippocampus of $Bdnf^{2A-Cre/+}$ mice and examined

colocalization of Olig2 with mCherry and EGFP. Out of the transduced Olig2$^+$ cells, 14.5% express

BDNF in adult mice (Fig. 4I-M). It is unexpected that the percentage of BDNF-expressing

oligodendrocytes in adult is higher than that of cumulative expression in $Bdnf^{2A-Cre/+}$;EGFPL10a/+

mice. One possible explanation is that AAV injection might cause tissue damage or myelin

damage, which induced the increase in BDNF expression in oligodendrocytes.

Figure 4. A small subset of oligodendrocytes express BDNF in the hippocampus. (**A-H**) Immunostaining of Olig2

in the SS (A-C) and CA1 SR (E-G) of adult *Bdnf*[2A-Cre/+];EGFPL10a/+ mice, and (D,H) quantification of colocalization. (**I-**

M) Immunostaining of Olig2 in the CA1 SR of adult *Bdnf*[2A-Cre/+] mice injected with the mCherry-to-EGFP color switch

AAV, and (M) quantification of colocalization. Scale bar = 50 μm. Arrowheads denote EGFP-expressing

oligodendrocytes. Arrows denote oligodendrocytes that do not express EGFP.

274 3.4 Generation of *Ntrk2*[2A-Cre] allele.

To label cells that express TrkB.FL, we generated the *Ntrk2*[2A-Cre/+] mouse strain which

express Cre recombinase in TrkB.FL-expressing cells. We employed the CRISPR/Cas9

technology to insert the P2A-Cre sequence into the *Ntrk2* locus immediately before the stop codon

for the TrkB.FL (Fig. 5A). Mice with the insertion were identified using genomic DNA PCR (Fig.

5B). There was a high colocalization between *Cre* mRNA and *Ntrk2* mRNA for TrkB-FL in the

cortex of *Ntrk2*[2A-Cre/+] mice (Fig. 5C-D), validating the allele.

Figure 5. Generation of *Ntrk2²ᴬ⁻ᶜʳᵉ* mice. (A) Strategy for insertion of the P2A-Cre sequence into the *Ntrk2* locus. **(B)** Genomic DNA PCR to identify *Ntrk2²ᴬ⁻ᶜʳᵉ/⁺* mice. One primer is located outside each of the two homology arms in the targeting vector. **(C, D)** In situ hybridization of the cortex.

3.5 Microglia in the cortex, hippocampus or spinal cord do not express TrkB.FL or TrkB.T

To study TrkB.FL expression in glial cells, we crossed *Ntrk2*²ᴬ⁻ᶜʳᵉ/⁺ mice that express Cre in TrkB.FL-expressing cells to the Cre-dependent EGFP-L10a/+ reporter mice to generate *Ntrk2*²ᴬ⁻ᶜʳᵉ/⁺;EGFP-L10a/+ mice in which TrkB.FL expression is genetically labeled by EGFP. As expected, all neurons in the MO and hippocampal CA1 express TrkB.FL (Fig. 6A-F). Next, we examined whether microglia express TrkB.FL. No colocalization of EGFP and Iba1 was found in the MO or hippocampal CA1 SR of 4-month-old *Ntrk2*²ᴬ⁻ᶜʳᵉ/⁺;EGFP-L10a/+ mice, indicating that microglia in these regions do not express TrkB.FL any time up to the time when the mice were perfused (Fig. 6G-L).

Figure 6. Microglia do not express TrkB.FL. (**A-F**) Colocalization of NeuN⁺ neurons and EGFP in the MO and CA1 of *Ntrk2*[2A-Cre/+];EGFP-L10a/+ mice. (**G-L**) Homeostatic microglia do not express EGFP in the MO and CA1 SR of *Ntrk2*[2A-Cre/+];EGFP-L10a/+ mice. Scale bar = 50 μm. n = 2 mice.

Figure 7. Microglia do not express either TrkB.FL or TrkB.T in adult mice. (**A-F**) Colocalization of NeuN[+] neurons and EGFP in the MO and CA1 of *Ntrk2*[CreER/+];EGFP-L10a/+ mice. (**G-L**) Homeostatic microglia are not labeled by EGFP in the MO and CA1 SR of *Ntrk2*[CreER/+];EGFP-L10a/+ mice. Scale bar = 50 μm. n = 2 mice.

To see whether microglia express TrkB.T, we utilized *Ntrk2*[CreER/+] mice that express Cre-ER under the *Ntrk2* promoter. We crossed *Ntrk2*[CreER/+] mice to Cre-dependent EGFP-L10a/+ reporter mice to generate *Ntrk2*[CreER/+];EGFP-L10a/+ mice, in which cells expressing either TrkB.FL or TrkB.T are genetically labeled by EGFP upon tamoxifen exposure. As expected, neurons in the cortex and hippocampal CA1 are labeled by EGFP due to their expression of TrkB.FL and TrkB.T. However, only a subset of neurons were labeled by EGFP (Fig. 7A-F). Because almost all neurons should express TrkB.FL in adult mice (Fig. 6A-F), this suggests that

the Cre induction by tamoxifen was lower than 100%. Our data showed no colocalization of EGFP

and Iba1 in either MO or CA1 SR region (Fig. 7G-L). This means that Iba1[+] microglia do not

express either TrkB.FL or TrkB.T in adult mice.

3.5 Astrocytes predominantly express TrkB.T and a small subset of astrocytes express

TrkB.FL

Astrocytes have been shown to predominantly express TrkB.T; however, whether

astrocytes also express TrkB.FL is not as clear. To investigate the expression of TrkB.FL in

astrocytes, we quantified colocalization of EGFP and astrocyte markers, SOX9 and GFAP, in

Ntrk2[2A-Cre/+];EGFP-L10a/+ mice. Our analyses showed that a small portion of astrocytes (16.8%

in SS and 18.0% in hippocampal CA1 SR) express TrkB.FL at some point during the lifespan of

the mice (Fig. 8A-H). To test whether astrocytes in adult mice express TrkB.FL, we injected the

Cre-dependent mCherry-to-EGFP color switch AAV to the hippocampus of 2-month-old *Ntrk2*[2A-

Cre/+] mice [24]. 92.7% of astrocytes were transduced by the AAV virus, and only 3.6% of the

transduced cells express Cre (Fig. 8I-M). These results suggest that only a very small subset of

astrocytes express TrkB.FL in adult mice. Because this number is smaller than the 18.0% seen

in *Ntrk2*[2A-Cre/+];EGFP-L10a/+ mice. The two possibilities discussed before can also apply here: (1)

A small subset of astrocytes express TrkB.FL only early during development and stop the

expression in adult mice; (2) at each different time point, there is a different small subset of

astrocytes that express TrkB.FL, and that accumulates to the 18% seen in *Ntrk2*[2A-Cre/+];EGFP-

L10a/+ mice.

Figure 8. Astrocytes predominantly express TrkB.T in adult mice. (**A-G**) Immunostaining of SOX9 (A-C) or GFAP (D-G) in the SS (A-C) and CA1 SR (E-G) of *Ntrk2*[2A-Cre/+];EGFP-L10a/+ mice, and (D,H) quantification of colocalization. (**I-M**) Immunostaining of GFAP in the CA1 SR of adult *Ntrk2*[2A-Cre/+] mice injected with the mCherry-to-EGFP color switch AAV, and the quantification of the colocalization of GFAP and EGFP. (**N-U**) Immunostaining of SOX9 (A-C) or GFAP (D-G) in the SS (A-C) and CA1 SR (E-G) of *Ntrk2*[CreER/+];EGFP-L10a/+ mice, and the quantification of colocalization. Scale bar = 50 µm. Arrowheads denote EGFP-expressing astrocytes. Arrows denote astrocytes that do not express EGFP.

Next, we examined expression of TrkB.T using *Ntrk2*[CreER/+];EGFP-L10a/+ mice where cells expressing either TRKB.FL or TrkB.T are labeled by EGFP. Tamoxifen was given at the age of 6 weeks to induce Cre recombination. We found that 43.4% astrocytes express either TrkB.FL or

345 TrkB.T in the SS, and 66.2% in the CA1 SR (Fig. 8N-U). Because we showed that most astrocytes

346 do not express TrkB.FL in the CA1 SR of adult mice (Fig. 8I-M), this means most of the EGFP$^+$

347 astrocytes in the $Ntrk2^{CreER/+}$;EGFP-L10a/+ mice express TrkB.T only. Due to the incomplete Cre

348 induction described above, the actual percentage of astrocytes that express TrkB in

349 $Ntrk2^{CreER/+}$;EGFP-L10a/+ mice could be higher than what we show.

3.6 Oligodendrocytes express TrkB.FL and TrkB.T in adult brains

Lastly, we investigated TrkB expression in oligodendrocytes. Colocalization of

oligodendrocyte marker Olig2 and EGFP in $Ntrk2^{2A-Cre/+}$;EGFP-L10a/+ mice suggests that a

subset of oligodendrocytes (15.8% in SS and 43.4% in CA1 SR) express TrkB.FL at some point

during their lifespan (Fig. 9A-H). To examine whether oligodendrocytes express TrkB.FL in adult

mice, the Cre-dependent mCherry-to-EGFP color switch AAV was injected to the hippocampus

of adult $Ntrk2^{2A-Cre/+}$ mice. The transduction rate of oligodendrocytes is lower than that of

astrocytes. Around 27.8% of oligodendrocytes were transduced by the AAV. We observed that

21.1% of transduced Olig2$^+$ oligodendrocytes express TrkB.FL in the adult mouse brains (Fig. 9I-

M).

To find out whether oligodendrocytes also express TrkB.T, we quantified colocalization of

Olig2 and EGFP in tamoxifen-treated $Ntrk2^{CreER/+}$;GFPL10/+ mice. We found that 8.8% of Olig2$^+$

oligodendrocytes are colocalized with EGFP in the SS, indicating that these cells express either

TrkB.FL or TrkB.T in the adult mice. In the CA1 SR, 38.4% of Olig2$^+$ oligodendrocytes express

either TrkB.FL or TrkB.T in the adult mice (Fig. 9R-U). Because we showed 21.1% of Olig2$^+$

oligodendrocytes express TrkB.FL in the CA1 SR of adult mice, this means that there are at least

another 17% of Olig2$^+$ oligodendrocytes that express TrkB.T in the adult mouse brains. Olig2

labels both oligodendrocyte precursor cells (OPCs) and mature oligodendrocytes. To differentiate

between these two populations, we used aspartoacylase (ASPA) as a marker to mark mature

oligodendrocytes only. Our results show that nearly all mature oligodendrocytes do not express

either TrkB.FL or TrkB.T in the CA1 SR of adult mice, which suggests that only OPCs express

TrkB.FL and TrkB.T in adult mice (Fig. 9V-Y).

Figure 9. OPCs express TrkB.FL and TrkB.T in adult mice. (**A-G**) Immunostaining of Olig2 in the SS (A-C) and CA1 SR (E-G) of *Ntrk2*[2A-Cre/+];EGFP-L10a/+ mice, and (D,H) quantification of colocalization. (**I-M**) Immunostaining of Olig2 in the CA1 SR of adult *Ntrk2*[2A-Cre/+] mice injected with the mCherry-to-EGFP color switch AAV, and quantification of the colocalization of GFAP and EGFP. (**N-U**) Immunostaining of Olig2 in the SS (A-C) and CA1 SR (E-G) of *Ntrk2*[CreER/+];EGFP-L10a/+ mice, and (Q,U) quantification of colocalization. (**V-Y**) Immunostaining of ASPA in the CA1 SR (of *Ntrk2*[CreER/+];EGFP-L10a/+ mice, and (Y) quantification of colocalization. Scale bar = 50 µm. Arrowheads denote EGFP-expressing oligodendrocytes. Arrows denote oligodendrocytes that do not express EGFP.

4. Discussion

Whether microglia express BDNF and TrkB or not has been debated. Microglia were discovered to express BDNF *in vitro* at the level of mRNA [26, 27] and protein [28] decades ago. LPS stimulation was suggested to increase BDNF secretion from cultured microglia [26, 28]. *In vivo* experiments showed BDNF expression in activated microglia [29]. BDNF from spinal microglia were suggested to be responsible for pain hypersensitivity [7]. Microglial BDNF was also reported to promote motor learning-dependent synapse formation [6]. Following these studies, many researchers showed the knockdown of microglial expression of BDNF disrupts biological processes including self-renew and proliferation of hippocampal neurons [30], nerve injury-induced pyramidal neuron hypersensitivity in the SS cortex [31], and mechanical allodynia induced by high-frequency stimulation [32]. Microglial BDNF was also suggested to be critical for (*R*)-ketamine-mediated microglial activation in the medial prefrontal cortex of chronic social defeat stress mice [33]. However, other studies using $Bdnf^{Laz/+}$ reporter mice and transcriptomic analysis show that microglia do not express significant level of BDNF in the spinal cord [8, 34]. This inconsistency likely comes from the lack of binding specificity of BDNF antibodies, as many previous efforts to characterize BDNF expression relied on immunohistochemistry. To avoid the use of BDNF antibodies and increase sensitivity, we took advantage of the $Bdnf^{2A-cre/+}$ knockin mice that express Cre recombinase under the *Bdnf* promoter and the EGFPL10a/+ reporter mice that express EGFP localized in cell bodies but not dendrites and axons for better characterization of colocalization. Our data suggest that both resting and activated microglia do not express BDNF in the MO cortex, hippocampal CA1, and spinal cord. These results are consistent with findings from transcriptomic analysis that showed microglia do not express BDNF in the spinal cord [8], and from another group that recently demonstrated resting microglia and microglia activated by ATP do not express BDNF transcriptionally and translationally in the mouse motor cortex [35]. Therefore, the phenotypes seen after deleting the microglial *Bdnf* gene likely come from artifacts.

However, our data do not exclude the possibility that microglia may have the potential to

internalize and recycle BDNF expressed by neurons and other cell types in the CNS.

Using similar method, we also show that microglia do not express the TrkB receptor in the

brain and spinal cord. Our data show microglia do not express TrkB.FL from embryo to adult, and

do not express TrkB.T in adult mice. This means that at least in adult mice, microglia do not

respond to BDNF through the TrkB receptor. However, it is still likely that BDNF can signal through

the low affinity receptor p75^{NTR} in microglia, as studies have suggested a subset of microglia

express p75^{NTR} and its expression is upregulated upon infection in mice [36]. Another study

suggested no significant p75^{NTR} expression in wild-type microglia, but the expression is increased

in 5xFAD mice and Alzheimer's patients [37], suggesting a potential role of BDNF-p75^{NTR} signaling

in neuroinflammation.

Cultured astrocytes have been shown to express BDNF and the expression is increased

by KCl, carbachol, and glutamate [38, 39]. Astrocytic BDNF regulates neuronal dendrite

complexity and spine number *in vitro* [16]. Studies have suggested that cortical layer II/III

astrocytes can incorporate extracellular proBDNF through p75^{NTR}-mediated endocytosis and

cleave proBDNF to mBDNF [15]. But whether and what percentage of astrocytes produce BDNF

in vivo under physiological condition remain elusive. Our results show that a small subset of

astrocytes express BDNF in the SS cortex (14.6%) and hippocampal CA1 SR (18.3%) under

physiological condition in mice. In adult mice, 10% of astrocytes in CA1 SR express BDNF. This

is consistent with other studies that showed a relatively low baseline level of mBDNF in astrocytes

[40], but an upregulation of astrocytic BDNF expression under pathological conditions induced

by MK-801, an N-methyl-D-aspartic acid (NMDA) receptor antagonist, *in vitro* [41] and treatment

of glatiramer acetate in R6/2 HD mouse model *in vivo* [40]. Although only 10% of astrocytes

express BDNF in the hippocampus under physiological condition, astrocytes seem to be an

important source of BDNF for normal CNS functions. Studies using BDNF knock-down mouse

models found that astrocytic BDNF is required for oligodendrogenesis after white matter damage

[42]. In the experimental autoimmune encephalomyelitis (EAE) model of multiple sclerosis,

astrocyte specific BDNF knockout led to more severe clinical course with increased axonal injury

and loss [43].

It is known that astrocytes predominantly express TrkB.T, which plays important roles in

mediating calcium release from intracellular stores in astrocytes [9] and controlling astrocyte

morphological maturation [11]. In the adult brain, only less than 5% astrocytes express TrkB.FL

in the hippocampal CA1 SR. These data are consistent with the current belief that astrocytes

dominantly express TrkB.T in the adult mice. We also show that cumulatively through embryo to

adult, 18% astrocytes in the hippocampal CA1 SR have expressed TrkB.FL, suggesting it is likely

that some astrocytes only express TrkB.FL early during embryonic and postnatal developmental

stages. This corroborates a previous *in vitro* study that showed radial glia (the main astrocytic

precursor cells) and proliferating cortical astrocytes in culture express mRNAs for both TrkB.FL

and TrkB.T, whereas differentiating astrocytes predominantly express TrkB.T and downregulate

TrkB.FL expression [44]. These data suggest that BDNF-TrkB.FL signaling may play a role in

astrocyte development and maturation.

Our data show that a very small subset (less than 5%) of oligodendrocytes in the

hippocampal CA1 SR express BDNF. Contradictory to the studies that showed BDNF expression

in cortical oligodendrocytes [45, 46], our data showed that most Olig2[+] cells in the cortex do not

express BDNF. Oligodendroglial BDNF expression has been reported by a few studies. A study

has showed that oligodendrocytes in the spinal cord and optic nerve express BDNF mRNA and

the expression is increased in Long-Evans shaker rats that lack CNS myelin due to the MBP gene

mutation [47]. Another study found that fingolimod treatment resulted in increased BDNF mRNA

concomitant with increased OPC differentiation in rat OPC culture [48]. Our results confirm the

expression of BDNF in oligodendrocytes in certain brain regions. And our observation of

increased BDNF-expressing oligodendrocytes in AAV injected mice may suggest that BDNF

expression is increased when myelin is disrupted, as a previous study has reported increased

BDNF level in oligodendrocytes after injury [49].

Our results also showed that only OPCs, but not mature oligodendrocytes, express

TrkB.FL and TrkB.T in the hippocampal CA1 SR of adult mice. BDNF-TrkB signaling in OPCs has

been suggested to have important functions. OPC-specific TrkB deletion resulted in impaired

activity-dependent myelination [18]. BDNF and its structural mimetic TDP6 have been shown to

promote myelination and myelin repair via oligodendroglial TrkB receptor [20, 50, 51]. Future

studies are necessary to determine which TrkB isoform(s) are responsible for mediating these

effects.

In summary, we characterized the expression of BDNF and TrkB in microglia, astrocytes,

and oligodendrocytes in the current study. Our results clarify that microglia in the examined

regions do not express BDNF or TrkB. Our findings also open opportunities to investigate

previously unidentified roles of BDNF and TrkB in astrocytes and oligodendrocytes.

Author Contributions

C.N. and B.X. designed research; C.N., X.Y., J.J.A., and H.X. performed experiments; C.N.

analyzed data; C.N. wrote and B.X. edited the paper.

Funding

This work was supported by the grants from the National Institutes of Health to BX (R01

DK103335, R01 DK105954, and R01 MH125187).

Data Availability Statement

Original data generated and analyzed during this study are included in this published article.

Acknowledgments

We thank David Ginty for the *Ntrk2^{CreER}* mouse strain and Zhonghou Wang for injecting viruses into the hippocampus in a few mice.

Conflicts of Interest

The authors declared that they have no competing financial interests.

References

1. Tejeda, G.S. and M. Díaz-Guerra, *Integral Characterization of Defective BDNF/TrkB Signalling in Neurological and Psychiatric Disorders Leads the Way to New Therapies.* Int J Mol Sci, 2017. **18**(2).

2. Xu, B. and X. Xie, *Neurotrophic factor control of satiety and body weight.* Nat Rev Neurosci, 2016. **17**(5): p. 282-92.

3. Baxter, G.T., et al., *Signal transduction mediated by the truncated trkB receptor isoforms, trkB.T1 and trkB.T2.* J Neurosci, 1997. **17**(8): p. 2683-90.

4. Fenner, B.M., *Truncated TrkB: beyond a dominant negative receptor.* Cytokine Growth Factor Rev, 2012. **23**(1-2): p. 15-24.

5. Gupta, V.K., et al., *TrkB receptor signalling: implications in neurodegenerative, psychiatric and proliferative disorders.* Int J Mol Sci, 2013. **14**(5): p. 10122-42.

6. Parkhurst, C.N., et al., *Microglia promote learning-dependent synapse formation through brain-derived neurotrophic factor.* Cell, 2013. **155**(7): p. 1596-609.

7. Coull, J.A., et al., *BDNF from microglia causes the shift in neuronal anion gradient underlying neuropathic pain.* Nature, 2005. **438**(7070): p. 1017-21.

8. Denk, F., et al., *Persistent Alterations in Microglial Enhancers in a Model of Chronic Pain.* Cell Rep, 2016. **15**(8): p. 1771-81.

9.   Rose, C.R., et al., *Truncated TrkB-T1 mediates neurotrophin-evoked calcium signalling in glia*

cells. Nature, 2003. **426**(6962): p. 74-8.

10. Fernández-García, S., et al., *Astrocytic BDNF and TrkB regulate severity and neuronal activity*

in mouse models of temporal lobe epilepsy. Cell Death Dis, 2020. **11**(6): p. 411.

11. Holt, L.M., et al., *Astrocyte morphogenesis is dependent on BDNF signaling via astrocytic*

TrkB.T1. Elife, 2019. **8**.

12. Aroeira, R.I., A.M. Sebastião, and C.A. Valente, *BDNF, via truncated TrkB receptor,*

modulates GlyT1 and GlyT2 in astrocytes. Glia, 2015. **63**(12): p. 2181-97.

13. Ameroso, D., et al., *Astrocytic BDNF signaling within the ventromedial hypothalamus*

regulates energy homeostasis. Nat Metab, 2022. **4**(5): p. 627-643.

14. Vignoli, B., et al., *Peri-Synaptic Glia Recycles Brain-Derived Neurotrophic Factor for LTP*

Stabilization and Memory Retention. Neuron, 2016. **92**(4): p. 873-887.

15. Vignoli, B., et al., *Astrocytic microdomains from mouse cortex gain molecular control over*

long-term information storage and memory retention. Commun Biol, 2021. **4**(1): p. 1152.

16. de Pins, B., et al., *Conditional BDNF Delivery from Astrocytes Rescues Memory Deficits,*

Spine Density, and Synaptic Properties in the 5xFAD Mouse Model of Alzheimer Disease. J

Neurosci, 2019. **39**(13): p. 2441-2458.

17. Jang, M., et al., *Oligodendrocytes regulate presynaptic properties and neurotransmission*

through BDNF signaling in the mouse brainstem. Elife, 2019. **8**.

18. Geraghty, A.C., et al., *Loss of Adaptive Myelination Contributes to Methotrexate*

Chemotherapy-Related Cognitive Impairment. Neuron, 2019. **103**(2): p. 250-265.e8.

19. Wong, A.W., et al., *Oligodendroglial expression of TrkB independently regulates myelination*

and progenitor cell proliferation. J Neurosci, 2013. **33**(11): p. 4947-57.

20. Xiao, J., et al., *Brain-derived neurotrophic factor promotes central nervous system myelination*

via a direct effect upon oligodendrocytes. Neurosignals, 2010. **18**(3): p. 186-202.

21. Tan, C.L., et al., *Warm-Sensitive Neurons that Control Body Temperature.* Cell, 2016. **167**(1):
p. 47-59.e15.

22. Rutlin, M., et al., *The cellular and molecular basis of direction selectivity of Aδ-LTMRs.* Cell,
2014. **159**(7): p. 1640-51.

23. Liu, J., et al., *Cell-specific translational profiling in acute kidney injury.* J Clin Invest, 2014.
124(3): p. 1242-54.

24. Bäck, S., et al., *Neuron-Specific Genome Modification in the Adult Rat Brain Using CRISPR-*
Cas9 Transgenic Rats. Neuron, 2019. **102**(1): p. 105-119.e8.

25. Chan, K.Y., et al., *Engineered AAVs for efficient noninvasive gene delivery to the central and*
peripheral nervous systems. Nat Neurosci, 2017. **20**(8): p. 1172-1179.

26. Miwa, T., et al., *Lipopolysaccharide enhances synthesis of brain-derived neurotrophic factor*
in cultured rat microglia. J Neurosci Res, 1997. **50**(6): p. 1023-9.

27. Elkabes, S., E.M. DiCicco-Bloom, and I.B. Black, *Brain microglia/macrophages express*
neurotrophins that selectively regulate microglial proliferation and function. J Neurosci, 1996.
16(8): p. 2508-21.

28. Nakajima, K., et al., *Neurotrophin secretion from cultured microglia.* J Neurosci Res, 2001.
65(4): p. 322-31.

29. Soontornniyomkij, V., et al., *Expression of brain-derived neurotrophic factor protein in*
activated microglia of human immunodeficiency virus type 1 encephalitis. Neuropathol Appl
Neurobiol, 1998. **24**(6): p. 453-60.

30. Harley, S.B.R., et al., *Selective Ablation of BDNF from Microglia Reveals Novel Roles in Self-*
Renewal and Hippocampal Neurogenesis. J Neurosci, 2021. **41**(19): p. 4172-4186.

31. Huang, L., et al., *BDNF produced by cerebral microglia promotes cortical plasticity and pain*
hypersensitivity after peripheral nerve injury. PLoS Biol, 2021. **19**(7): p. e3001337.

32. Zhou, L.J., et al., *Microglia Are Indispensable for Synaptic Plasticity in the Spinal Dorsal Horn*
and Chronic Pain. Cell Rep, 2019. **27**(13): p. 3844-3859.e6.

33. Yao, W., et al., *Microglial ERK-NRBP1-CREB-BDNF signaling in sustained antidepressant*

actions of (R)-ketamine. Mol Psychiatry, 2022. **27**(3): p. 1618-1629.

34. Dembo, T., et al., *Primary Afferent-Derived BDNF Contributes Minimally to the Processing of*

Pain and Itch. eNeuro, 2018. **5**(6).

35. Honey, D., et al., *Analysis of microglial BDNF function and expression in the motor cortex.*

Front Cell Neurosci, 2022. **16**: p. 961276.

36. Düsedau, H.P., et al., *p75(NTR) regulates brain mononuclear cell function and neuronal*

structure in Toxoplasma infection-induced neuroinflammation. Glia, 2019. **67**(1): p. 193-211.

37. Capsoni, S., et al., *The chemokine CXCL12 mediates the anti-amyloidogenic action of*

painless human nerve growth factor. Brain, 2017. **140**(1): p. 201-217.

38. Wu, H., W.J. Friedman, and C.F. Dreyfus, *Differential regulation of neurotrophin expression*

in basal forebrain astrocytes by neuronal signals. J Neurosci Res, 2004. **76**(1): p. 76-85.

39. Jean, Y.Y., L.D. Lercher, and C.F. Dreyfus, *Glutamate elicits release of BDNF from basal*

forebrain astrocytes in a process dependent on metabotropic receptors and the PLC pathway.

Neuron Glia Biol, 2008. **4**(1): p. 35-42.

40. Reick, C., et al., *Expression of brain-derived neurotrophic factor in astrocytes - Beneficial*

effects of glatiramer acetate in the R6/2 and YAC128 mouse models of Huntington's disease.

Exp Neurol, 2016. **285**(Pt A): p. 12-23.

41. Yu, W., et al., *Reactive Transformation and Increased BDNF Signaling by Hippocampal*

Astrocytes in Response to MK-801. PLoS One, 2015. **10**(12): p. e0145651.

42. Miyamoto, N., et al., *Astrocytes Promote Oligodendrogenesis after White Matter Damage via*

Brain-Derived Neurotrophic Factor. J Neurosci, 2015. **35**(41): p. 14002-8.

43. Linker, R.A., et al., *Functional role of brain-derived neurotrophic factor in neuroprotective*

autoimmunity: therapeutic implications in a model of multiple sclerosis. Brain, 2010. **133**(Pt 8):

p. 2248-63.

44. Climent, E., et al., *Astrocytes in culture express the full-length Trk-B receptor and respond to*

brain derived neurotrophic factor by changing intracellular calcium levels: effect of ethanol

exposure in rats. Neurosci Lett, 2000. **288**(1): p. 53-6.

45. Dai, X., et al., *The trophic role of oligodendrocytes in the basal forebrain.* J Neurosci, 2003.

23(13): p. 5846-53.

46. Bagayogo, I.P. and C.F. Dreyfus, *Regulated release of BDNF by cortical oligodendrocytes is*

mediated through metabotropic glutamate receptors and the PLC pathway. ASN Neuro, 2009.

1(1).

47. Smith, C.M., E. Cooksey, and I.D. Duncan, *Myelin loss does not lead to axonal degeneration*

in a long-lived model of chronic demyelination. J Neurosci, 2013. **33**(6): p. 2718-27.

48. Khani-Habibabadi, F., et al., *Hotair and Malat1 Long Noncoding RNAs Regulate Bdnf*

Expression and Oligodendrocyte Precursor Cell Differentiation. Mol Neurobiol, 2022. **59**(7): p.

4209-4222.

49. Dougherty, K.D., C.F. Dreyfus, and I.B. Black, *Brain-derived neurotrophic factor in astrocytes,*

oligodendrocytes, and microglia/macrophages after spinal cord injury. Neurobiol Dis, 2000.

7(6 Pt B): p. 574-85.

50. Wong, A.W., et al., *TDP6, a brain-derived neurotrophic factor-based trkB peptide mimetic,*

promotes oligodendrocyte myelination. Mol Cell Neurosci, 2014. **63**: p. 132-40.

51. Fletcher, J.L., et al., *Targeting TrkB with a Brain-Derived Neurotrophic Factor Mimetic*

Promotes Myelin Repair in the Brain. J Neurosci, 2018. **38**(32): p. 7088-7099.