Assimilation efficiencies and elimination rates of silver, cadmium and zinc accumulated

by trophic pathway in *Gammarus fossarum*

Ophélia Gestin[a,b,c,*] (gestin.ophelia@gmail.com), Christelle Lopes[a] (christelle.lopes@univ-lyon1.fr),

Nicolas Delorme[c] (nicolas.delorme@inrae.fr), Laura Garnero[c] (laura.garnero@inrae.fr), Olivier

Geffard[c] (olivier.geffard@inrae.fr) and Thomas Lacoue-Labarthe[b] (tlacouel@univ-lr.fr)

[a] Univ Lyon, *Université Lyon 1, CNRS, Laboratoire de Biométrie et Biologie Évolutive UMR 5558,*

69622 Villeurbanne, France

[b] *Littoral Environnement et Sociétés (LIENSs), UMR 7266 CNRS - Université de la Rochelle, 2 rue*

Olympe de Gouges, 17000 La Rochelle, France

[c] *INRAE, RiverLy, Ecotoxicology Laboratory, 5 Avenue de la Doua, CS20244, 69625 Villeurbanne*

Cedex, France

*Corresponding author

Abstract

To improve the assessment of metal toxicity in aquatic organisms, it is important to consider the different uptake pathways (i.e. trophic or aqueous). The bioaccumulation of dissolved metals such as Cd and Zn in gammarids is beginning to be well described. However, there are very few data on the contribution of the dietary pathway, and its associated toxicokinetic parameters. Among these, the assimilation efficiency (AE) is an essential parameter for the implementation of models that take the trophic pathway into account. This study aims to estimate the assimilation efficiencies and elimination rates of two types of food, i.e. alder leaves and chironomid larvae, contaminated with three metals (Ag, Cd and Zn) of major concern for the Water Framework Directive (WFD). The pulse-chase-feeding method was used. Gammarids were fed with alder leaves or chironomid larvae previously contaminated with ^{110m}Ag, ^{109}Cd or ^{65}Zn, for a short period of time (1 to 5 hours), followed by an elimination phase of 14 days. At different time points, the gammarids were placed alive on the gamma detector to individually quantify whole body concentrations of ^{110m}Ag, ^{109}Cd or ^{65}Zn. Our results indicate that: i) Cd has the highest assimilation efficiency (39% for leaves and 19% for larvae), followed by Zn (15% for leaves and 9% for larvae) and Ag (5% for leaves); ii) for Cd and Zn, the AE were higher when gammarids were fed with leaves than with larvae; iii) the elimination rates of metals seem to depend more on the food matrix than on the metal assimilated; and thus iv) the biological half-life calculated from the k_{es} is 5.1 days for Ag, between 4.9 and 13 days for Cd and between 3.8 and 13 days for Zn.

Keywords: Trophic transfer, Amphipods, Metals, Alder leaves, Chironomids larvae, Dietary pathway

1. Introduction

In the past decades, a number of works have demonstrated the strong capacity of some freshwater organisms to accumulate contaminants from the aquatic environment (Timmermans et al., 1992; Dutton & Fisher, 2011). Consequently, some freshwater invertebrates have been used as bioindicators of good ecological status for French freshwater systems, based on the chemical concentrations recorded in the whole body (Besse et al., 2012; Pellet et al., 2014; Lebrun et al., 2020). The ubiquitous gammarid *Gammarus fossarum* is widely used for this property, as it accumulates metals as a grazer of leaf litter and is an opportunistic predator of invertebrate prey (Kunz et al., 2010; Pellet et al., 2014; Filipović Marijić et al., 2016). Although the bioaccumulation processes of dissolved metals have been extensively documented (Vellinger et al., 2012; Urien et al., 2017; Gestin et al., 2022, 2023), less is known about the dietary assimilation and elimination of metals in this species. As exposure conditions are more difficult to standardise than for aqueous exposures, the dietary route in aquatic invertebrates is often poorly understood. However, the dietary pathway could account for a significant metal entry in invertebrates (Borgmann et al., 1989; Pellet et al., 2014), which highlights the need to consider this contribution in order to avoid misinterpretation of recorded metal concentrations in this species (Franklin et al., 2005; Besse et al., 2013; Lebrun et al., 2015; Conti et al., 2016; Hadji et al., 2016).

The few studies focusing on freshwater invertebrates have shown that the uptake from the trophic pathway predominates for some metals and metalloids, such as for Cd, Cu and Se in *Hyallela azteca* (Borgmann et al., 2007) or Cd in *Gammarus pulex* (Pellet et al., 2014). In *Daphnia magna*, however, the uptake from the aqueous pathway predominates for Zn (Memmert, 1987). Previous results suggest that metal accumulation, detoxification and elimination, and thus potential toxicity, are strongly dependent on the uptake pathway. In this

sense, Croteau & Luoma (2009) called for the need to delineate the bioaccumulation

mechanisms and toxicokinetics in different organisms, as these processes are metal and species

specific.

Assimilation efficiency (AE) is one of the most important parameters used to quantify trophic

transfer. This parameter directly reflects the amount of metals available from food and is a key

physiological parameter for understanding the trophic transfer mechanisms (Croteau & Luoma,

2009). The AE could be partly defined by the fraction of metal in food that crosses the intestinal

epithelial barrier to accumulate in the body of the consumer (Wang & Fisher, 1999a; Nunez-

Nogueira et al., 2006; Croteau & Luoma, 2008). AE values can be used to estimate the influence

of consumer species, contaminants, temperature, etc. (Wang & Fisher, 1999a; Wang & Wong,

2003). Among these, the type of food can greatly influence the AE of metals and thus their

bioaccumulation efficiency (Wang & Fisher, 1999a; Nunez-Nogueira et al., 2006; Dubois &

Hare, 2009; Pouil et al., 2016). It has already been established that metals from plant matrices

are less assimilated by organisms than elements of animal origin, because the digestion of

compounds such as fibres or hemicelluloses is slower and more complex (Xu & Pascoe, 1994;

Nunez-Nogueira et al., 2006). It is therefore important to consider dietary diversity in

omnivorous organisms, such as gammarids, in trophic pathway studies.

The pulse-chase-feeding approach is commonly used to determine the AE values (Calow &

Fletcher, 1972; Warnau et al., 1996; Metian et al., 2007; Pouil et al., 2017) and consists of

exposing the organism once to a labelled food source and tracking the elimination of the tracer

over time (Wang & Fisher, 1999a; Pellet et al., 2014). A kinetic modelling approach is applied

to the depuration data to determine the metal-specific AE and the depuration rates (k_e). The

pulse-chase-feeding technique is widely used and adapted to work with different tools, such as

isotopic tracers (stable and radioactive ones) (Wang & Fisher, 1999a; Baines & Fisher, 2002;

Pouil et al., 2017). In the case of metals, gamma-emitting radiotracers are among the most

powerful tracers for accurately quantifying small amounts of metal. They also allow individual

tracking of metal uptake and loss over time, as whole-body gamma detection is non-destructive

(Wang & Fisher, 1999a). Furthermore, it is conventionally accepted that radioactive isotopes

of metals have identical properties to their stable counterparts (Warnau & Bustamante, 2007).

This confers the ability to assess metal accumulation and elimination rates by measuring only

the concentrations of radioisotopes, thus avoiding the measurement of the background of stable

metals (Cresswell et al., 2017).

A toxicokinetic model for the aqueous pathway has already been developed for zinc and

cadmium (Gestin et al., 2022). However, to be able to set up this kind of model considering the

trophic pathway, it is important to have preliminary data concerning the AE of metals. In this

framework, the aim of this study was to fill the knowledge gap on metal assimilation efficiency

in gammarids, by determining the AE values and the elimination rates of Ag, Cd and Zn in the

amphipod *G. fossarum*. A pulse-chase-feeding method was used, taking into account two

contrasting types of food in the gammarid diet: alder leaves and chironomid larvae. After a

short period of exposure to radiolabelled food, gammarids were transferred to depuration

conditions and metal loss in tagged individuals was monitored for 14 days. Nonlinear least

squares (NLS) models were fitted to the experimental data to determine the AE, k_e and

biological half-life of metals in gammarids. The values were then compared to discuss the

influence of metal and food type on dietary bioaccumulation.

2. Material and methods

2.1 Organisms and alder leaves: origins, collection and maintenance

Adult male gammarids (*Gammarus fossarum*: 27.3 ± 5.7 mg wet weight) were sampled from an uncontaminated watercress pound (Saint-Maurice-de-Rémens, France) (Gestin et al., 2021). They were then transported to the LIENSs laboratory premises (La Rochelle), where they were kept under constant bubbling in Evian® water (see characteristics in Table S1) with a dark/light cycle of 8:16h, at a temperature of 12.0 ± 0.5°C. They were fed with alder (*Alnus glutinosa*) leaves *ad libitum* and kept for a 7-day acclimation period before the experiments.

Alder leaves (*A. glutinosa*) used for trophic exposure were collected from a clean site (Les Ardillats, France). They were leached (i.e. soaked in well water for several days, with daily renewal of the water) in the ecotoxicology laboratory of INRAE Lyon-Villeurbanne in order to depurate the toxic tannins for gammarid. The wet leaves were then cut into discs of 6 mm diameter, avoiding the large veins. Finally, the leaf discs were placed in a container filled with well water (see characteristics in Table S1) and transferred to the LIENSs laboratory.

Chironomid (*Chironomus riparius*) egg masses from the INRAE laboratory were transferred to the LIENSs laboratory. The eggs were placed in beakers containing powdered silica and filled with Evian® water and kept at room temperature (about 20°C) for few days until hatching. The larvae were then fed with fish pellets until the third larval instar.

2.2 Radiotracers

Throughout the experiment, the rearing and laboratory materials were decontaminated with 1/10 diluted detergent (Decon® 90 solution), 1/10 diluted HCl solution (hydrochloric acid S.G. 32%, certified AR for analysis; Fischer Scientific®) and distilled water. The radioisotope carrier-free solutions were purchased from Eckert & Ziegler Isotope Products Inc., Valencia, USA. The ^{109}Cd and ^{65}Zn solutions were in their chloride forms (i.e. $CdCl_2$ and $ZnCl_2$), 0.1M

HCl and the ^{110m}Ag solution was in its nitrate form (i.e. AgNO$_3$), 0.1M HNO$_3$. These three

stock solutions were diluted to intermediate solutions to achieve appropriate radiotracer

concentrations in the experimental setup. These radiotracer additions to the media did not

change the pH of the water. For each radioisotope, the detectors were calibrated with two in-

house liquid standards of the appropriate geometry, *i.e.* a 500 µL "gammarid" geometry and a

10 mL "water" geometry. Two NaI detectors coupled to InterWinner 7.0 software (ITECH

Instruments$^{®}$) were used to count the radioisotope activities (in Bq, Bq.g^{-1} or Bq.mL^{-1}) present

in all samples.

2.3 Contamination of food

The leaf discs (~ 108 in total, i.e. 36 for each metal) were placed in 10 mL of Evian$^{®}$ water

contaminated with 2010, 2030 or 2000 Bq.mL^{-1} of ^{109}Cd, ^{65}Zn and ^{110m}Ag, respectively, for

7 days. They were then placed in 0.1 L beakers of clean Evian$^{®}$ water for 5 days to remove the

weakly adsorbed radioisotopes on the leaf walls. At the end of the procedures, 12 discs for each

metal were randomly counted by gamma spectrometry at the end of these procedures, to follow

the radiolabeling of the leaf.

At the end of the third instar (i.e. L3, when they start to turn red), nine pools of 10 chironomid

larvae each (i.e. 90 larvae in total) were collected, placed in 100 mL beakers filled with Evian$^{®}$

water (i.e. n=10 per beaker) and exposed for 4 days to 201, 203 or 200 Bq.mL^{-1} for ^{109}Cd, ^{65}Zn

and ^{110m}Ag (3 pools per radiotracer). This short exposure time was chosen to avoid the

emergence of radiolabelled larvae in mosquitoes in the laboratory. At the end of the exposure

period, the larvae were gently rinsed (i.e. rapidly soaked in 3 successive baths of clean water),

then individually frozen and all counted by gamma-ray spectrometry. No radiotracer activity

was detected in chironomid larvae exposed to ^{110m}Ag.

 2.4 Experimental procedure to obtain assimilation efficiencies of gammarids

The Ag AE value and elimination rate were not determined with chironomids as food source.

Prior to the pulse-chase feeding experiment, gammarids (n = 22 for each alder leaf/metal

condition and n = 14 for each chironomid/metal condition) were starved for 2 days and then

individually placed in 250 mL beakers filled with Evian® water (Fig. 1). For each metal, they

were then either exposed to either two radiolabelled leaf discs for 3-5 h, or to one radiolabelled

thawed chironomid larva for 1 h (Fig. 1). For the leaf-feeding condition, the 14 gammarids that

ate the most were whole-body gamma-counted alive immediately after the pulse-chase-feeding

period. For the larval feeding conditions, all the 14 exposed gammarids were gamma-counted.

Selected gammarids, as described below in section 3.1, (i.e. 13 ^{109}Cd-leaf-fed gammarids, 11

^{109}Cd-chironomid-fed gammarids, 13 ^{65}Zn-leaf-fed gammarids, 13 ^{65}Zn-chironomid-fed

gammarids and 14 ^{110m}Ag-leaf-fed gammarids) were then transferred under depuration

conditions, *i.e.* 200 mL beakers filled with non-radiolabelled with Evian® water (*i.e.* closed

circuit, constantly aerated, T = 12.0 ± 0.5°C). Each beaker contained 7 radiolabelled gammarids

individually separated by handmade baskets (i.e., plastic mesh 11 cm high and 8.6 cm in

diameter, with a mesh size of 0.5 cm; Fig. 1). A non-radiolabelled individual was added to each

beaker to control for potential recycling of radioisotopes through to leaching from gammarid

depuration. Gammarids were fed *ad libitum* with uncontaminated alder leaves throughout the

depuration period (Fig. 1).

Whole-body radiotracer activities in gammarids were tracked individually during two weeks of

depuration (Tables S2 and S3). Individuals were sampled and counted at days 0.5, 1, 1.5, 2, 4,

7, 9, 11 and 14 (Fig. 1). Gammarids were removed from the beakers using a 10 mL pipette,

rinsed with clean water and placed in a plastic box (Caubère®, ref 1210) containing 500 µL of

clean Evian® water (Fig. 1). Counting times of live gammarids varied between 10 and 20 min

in order to minimise stress. The counting uncertainties did not exceed 5% for Zn for all samples,

whereas the uncertainties increased with decreasing activities in the gammarid samples, without

exceeding 20% and 15% for Cd and Ag, respectively. The radiotracer activities in the water

were monitored daily and the water was changed at least every two days to minimize tracer

contamination in the water. Mortality was also monitored daily.

Figure 1. The different steps of the pulse-chase-feeding.

2.5 Estimation of assimilation efficiencies and depuration parameters of gammarids

The depuration of radiotracers by gammarids was expressed as the percentage of remaining

activity (i.e. radioactivity at time t divided by the initial radioactivity measured in the organism

at the beginning of the depuration period; Tables S2 and S3; see Warnau et al., 1996). The

kinetic parameters and the assimilation efficiencies with respect to the food type and metal were

estimated using a nonlinear least squares (NLS) approach modelling (see the supplementary

material file for the script), following the two-component exponential equation:

$$A(t)_{f,r} = \left(100 - AE_{f,r}\right) \times e^{-k_{es,f,r} \times t} + AE_{f,r} \times e^{-k_{el,f,r} \times t} \qquad (1)$$

where $A(t)_{f,r}$ is the remaining activity (%) at time t, with respect to the type of food f and the

radioisotope r, $AE_{f,r}$ is the assimilation efficiency, $k_{es,f,r}$ and $k_{el,f,r}$ the elimination rates of

the first short-term phase (i.e. elimination of the non-assimilated metal fraction, which is rapidly

depurated) and of the second long-term phase (i.e. elimination of the assimilated metal fraction,

which is slowly depurated), respectively. The indexes f correspond to the type of food ingested

by the gammarids, with f =1 for alder leaves and f =2 for chironomid larvae; and the indexes

r correspond to the radioisotope tested, with r =1 for [109]Cd, r =2 for [65]Zn and r =3 for [110m]Ag.

The biological half-life $T_{b\frac{1}{2},f,r}$ for each metal with respect to the food source was calculated

from the elimination rate constant (k_{es} or k_{el}), based on the equation:

$$Tb_{\frac{1}{2},f,r} = \frac{ln2}{k_e} \qquad (2)$$

The elimination kinetic parameters and the assimilation efficiencies were estimated using R

software.

3. Results and discussion

Estimation of metal accumulation efficiencies is an essential first step in understanding bioaccumulation mechanisms. Furthermore, AE values are key inputs for the implementation of toxicokinetic models to study these mechanisms (Van Campenhout et al., 2009). For the gammarid *G. fossarum*, this study provides estimates of the assimilation efficiency and of the two elimination rates of biphasic digestion after the ingestion of food (i.e. of animal or plant origin) contaminated with one of the three metals studied here (Ag, Cd and Zn).

3.1 Data quality evaluation and selection

The ^{109}Cd is a low energy photon emitter with a gamma ray peak at 88 keV (3.79%), which means that a sample must have enough activity to stand out from the background compared to ^{65}Zn (1115 keV) or ^{110m}Ag (658 keV for its main gamma ray). Therefore, in order to obtain organisms that have eaten enough and to be able to determine the curve correctly, only individuals that reached a whole-body activity of 150 Bq and above immediately after the single feeding (t0) were kept to follow the metal loss kinetics (Fig. 1 and S1). This means that individuals that did not graze enough during the exposure period due to repletion or stress were dismissed. For the gammarids fed with Cd-enriched chironomids (n=11), only 4 gammarids in a first set of 14 individuals showed a body activity above 150 Bq at t0 (Fig. S1). In a second batch of 14 gammarids, 7 individuals ingested more than 150 Bq of ^{109}Cd activities (Fig. S1, Tables S2 and S3). For the Cd leaf-feeding condition (n=13), only 1 of 14 gammarids ingested less than 150 Bq of ^{109}Cd at t0, and so was therefore removed from the experiment. In each condition (n=13) of dietary Zn exposure (i.e. both leaves and larvae as sources), one gammarid died during the first day of the experiment. In the leaf-feeding Ag condition (n=14), all 14 gammarids remained alive throughout the depuration phase.

 3.2. Metal elimination kinetics patterns

253   Kinetic models were fitted to whole body concentrations over time following a pulse-chase-

254   feeding of metal-contaminated alder leaves. The metal elimination kinetics showed a biphasic

255   pattern (Fig. 2), expressing the presence of two metal pools: a first poorly retained pool

256   characterised by a high rate of elimination (k_{es}), and a second slowly eliminated pool (k_{el}).

257
258   **Figure 2.** Influence of the metal and the type of matrix used for feeding on the elimination of metals ingested
259   by the gammarids during the depuration phase, expressed as the percentage of activity remaining in the
260   gammarids compared to the end of the pulse-chase phase (t0) as a function of the depuration time (hours).
261   Gammarids were contaminated by cadmium for a) and b); zinc for c) and d); and silver for e). Food
262   corresponded to leaf discs for a), c) and e); and to chironomid larvae for b) and d).

Regardless of the metal, gammarids eliminated the highest proportions of ingested trace elements during the first 48 h after exposure to contaminated leaves. Elimination rates were similar among metals with k_{es} values of 0.18 ± 0.01 h^{-1}, 0.15 ± 0.03 h^{-1} and 0.14 ± 0.01 h^{-1} for Zn, Cd and Ag respectively (Table 1). Thus, these values imply that, after a 5 h period of grazing, these metal pools had a $Tb_{s1/2}$ of 4-5 hours (Table 1). These biological half-lives, which are similar among metals, are consistent with the intestinal transit time reported for *Gammarus pulex* (16 hours; Pellet et al., 2014), suggesting that these fractions were not assimilated and were excreted in feces. This hypothesis, stimulated by the ingestion of non-radiolabelled alder leaves since the gammarids were placed in depuration conditions, is supported by this significant release of radiotracers during the first 48 hours of the elimination phase.

Table 1. Parameters estimated by fitting an NLS model from pulse-chase feeding data, with EA (%), k_{el} (d^{-1}) and k_{es} (d^{-1}) with their respective standard deviations, t-value, Pr(>ItI) and percentiles (95%). Calculated biological half-lives times of metals as a function of diet type with their respective standard deviations and percentiles (95%).

| Metal | Food type | Parameters | Assimilation efficiencies | | | | | Biological half-life | | |
| | | | Estimate values | t value | Pr(>ItI) | Percentiles | | Calculated values | Percentiles | |
						2.5%	97.5%		2.5%	97.5%
Ag	Alder leaves	k_{es}	0.14 ± 0.01	14	<2E-16	0.12	0.16	5.1 ± 0.4	4.4	5.9
		AE	5.2 ± 1.1	4.8	4.2E-6	3.0	7.3	/	/	/
		k_{el}	0	/	/	/	/	∞	/	/
Cd	Chironomid larvae	k_{es}	0.053 ± 0.005	11	<2E-16	0.044	0.063	13 ± 1	11	16
		AE	19 ± 2	11	<2E-16	15	22	/	/	/
		k_{el}	0	/	/	/	/	∞	/	/
	Alder leaves	k_{es}	0.15 ± 0.03	5.5	3.0e-7	0.09	0.20	4.9 ± 1.0	3.6	7.2
		AE	39 ± 3	12	<2E-16	33	45	/	/	/
		k_{el}	0.0058 ± 0.0010	5.6	1.8e-7	0.0037	0.0078	124 ± 25	89	185
Zn	Chironomid larvae	k_{es}	0.055 ± 0.004	15	<2E-16	0.047	0.062	13 ± 1	11	15
		AE	9.1 ± 1.2	7.3	2.6E-11	6.7	12	/	/	/
		k_{el}	0	/	/	/	/	∞	/	/
	Alder leaves	k_{es}	0.18 ± 0.01	13	<2E-16	0.16	0.21	3.8 ± 0.3	3.3	4.4
		AE	15 ± 2	8.4	8.0E-14	11	18	/	/	/
		k_{el}	0.0061 ± 0.0014	4.4	2.3E-5	0.0034	0.0089	120 ± 35	78	203

The second kinetic phase was characterised by a lower elimination rate, with k_{el} values of

0.0061 ± 0.0014 h^{-1}, 0.0058 ± 0.0010 h^{-1} and zero for Zn, Cd and Ag, corresponding to k_{el} of

$Tb_{1/2} = 120 \pm 35$ h, 124 ± 25 h and ∞, respectively (Table 1). This longer elimination rate is

considered to be due to the physiological turnover of the assimilated metal fraction. It is

assumed that this pool crosses the intestinal barrier, is distributed to different organs and is

taken over by detoxification mechanisms, as already described for Cd and Zn in invertebrates

(Nunez-Nogueira et al., 2006). It is noteworthy that the $Tb_{1/2}$ of Cd and Zn are equivalent,

suggesting that metabolism of both trace elements requires similar physiological mechanisms.

In contrast, the assimilated pool of Ag was not eliminated by the gammarid, which contrasts

with a shorter Ag retention time than other metals, such as Cd or Zn, reported in daphnia (Lam

& Wang, 2006), in cuttlefish and in turbot (Bustamante et al., 2004; Pouil et al., 2015). This

phenomenon has already been observed in *Lymnea stagnalis* (k_{el} not different from 0)

suggesting that Ag is strongly retained in internal tissues (Croteau et al., 2011). This could be

explained by the fact that Ag is an element that reacts very strongly with sulphur compounds

(Lam & Wang, 2006). In bivalves, Ag has indeed been shown to be stored in the epithelial cells

of the intestinal cells, as Ag_2S (Berthet et al., 1992), which are already known to be a site of

accumulation and detoxification of other metals such as Cd and Cu in gammarids (Pellet et al.,

2014). In this highly stable Ag_2S form, silver is found in the insoluble fraction (Berthet et al.,

1992), and remains indefinitely sequestered in tissues in a non-metabolically available form

(Fowler et al., 2004).

3.3. The AE values of metals after feeding by contaminated alder leaves

According to the second depuration kinetic components, the assimilation efficiencies of metals

in alder leaves fed to gammarids were ranked as follows: Cd ($39 \pm 3\%$) > Zn ($15 \pm 2\%$) > Ag

($5.2 \pm 1.1\%$) (Fig. 2; Table 1). These contrasting AE values among elements were also found

in *Daphnia magna*, with AE of 30-70%, 7-66% and 1-2%, for Cd, Zn and Ag, respectively (Yu

& Wang, 2002; Lam & Wang, 2006). Consistent with our results, a higher AE for Cd (5-47%)

than for Zn (1.4%) was also found in the leaf-fed gammarid *Gammarus pulex* (Xu & Pascoe,

1994; Pellet et al., 2014). To our knowledge, there are no data on Ag AE in amphipods, although

similar AE values ranging from 3 to 23% have been reported in the marine copepods

Acartia torua, *Acartia hudsonica* and *Temova longicornis*. Compared to the Ag AE determined

here in *G. fossarum*, higher AE values were reported for this metal in the cephalopod

Sepia officinalis (19 ± 3%) (Bustamante et al., 2004) and in the gastropod *Lymnaea stagnalis*

(73 ± 5%) (Croteau et al., 2011), whereas it did not exceed 1% in the fish *Scophthalmus*

maximus (0.32%) (Reinfelder & Fisher, 1991; Lam & Wang, 2006; Pouil et al., 2015). Thus, all

these data show large disparities confirming that there are no general rules for predicting AE

among metals and biological species (Wang & Fisher, 1999a; Pouil et al., 2018).

3.4. The effect of food type on metal trophic transfer

The AE values are intrinsically linked to both biotic and abiotic factors that modulate the

digestive physiology, metal bioavailability and bioaccessibility (Pouil et al., 2018). Among

these factors, the type of food and its amount significantly influence the AE values, as reported

in fish (Pouil et al., 2017), while it is less evident in copepods, as reviewed by Wang & Fisher

(1999b). More specifically, the AE of Cd in *Dreissena polymorpha* has been estimated to range

from 19 to 72% for 8 types of food (Roditi & Fisher, 1999). The AE of Zn has also been reported

to be highly variable in fish, ranging from 2 to 97% (with a median of 20%) as a function of

the predator species and the food sources (Pouil et al., 2018). Mechanistically, these AE

variations are thought to be driven by the subcellular distribution of metals in food (i.e. the

storage form). Theoretically, metals are considered to be trophically available for transfer to

consumers when they are associated with organelles, enzymes and metal-binding proteins (i.e.

Trophically Available Metals, abbreviated TAM), i.e. metals found in the soluble fraction.

Conversely, metals associated with the insoluble fraction (i.e. metal-rich granules or cellular

debris) may not be assimilated (Wallace & Lopez, 1996; Wallace & Luoma, 2003).

Nevertheless, in practice, the relationship between TAM and AE is less clear due to the complex

organ and subcellular distribution of metals in metazoan prey (see Pouil et al., 2016).

In the present study, the AE of Cd and Zn were 2- and 1.5-fold higher when the ingested food

was alder leaves ($39 \pm 3\%$ and $15 \pm 2\%$, respectively; Fig. 2 and Table 1) compared to

chironomid larvae ($19 \pm 2\%$ and $9.1 \pm 1.2\%$, respectively; Fig. 2 and Table 1). This is in

contrast to what is usually reported for omnivorous organisms, where the trophic transfer of

metals from plant sources is lower than that from animal prey because the digestion process of

plant materials is slower and more complex (Xu & Pascoe, 1994; Nunez-Nogueira et al., 2006).

For example, the shrimp *Penaeus indicus* fed with Cd-contaminated algae showed an AE of

$42.4 \pm 5.1\%$, a value 1.8 time lower than that reported for shrimp fed with Cd-contaminated

squid (*i.e.* $74.6 \pm 8.5\%$). In contrast, the AE of Zn was not dependent on the type of diet (i.e.

AE of *P. indicus* fed with squid $= 57.9 \pm 7.3\%$; AE of *P. indicus* fed with algae $= 59.1 \pm 8.4\%$)

(Nunez-Nogueira et al., 2006).

Regarding our results, it is noteworthy that live chironomid larvae were contaminated with the

radiotracers whereas only dead leaves were radiolabelled. More than the plant or animal

character, we suspect a large passive adsorption of metals on the walls of inert dead leaves

compared to a poor absorption within the leaf structures. Once ingested, the metal fraction

should be easily desorbed under a free form ion and thus more bioavailable, increasing the

assimilation efficiency for the gammarid. On the contrary, metals actively ingested by live

chironomid larvae were distributed among tissues, incorporated into cells and taken up by

detoxification mechanisms, thus binding to various subcellular components. Therefore, the

higher AE obtained for leaf-fed gammarids than for chironomid-fed gammarids could be related

to the sorption processes and the TAM fractions in live vs. inert food sources rather than to the

contrasting efficiencies of digestion of plant and animal materials.

As found for AE, both k_{es} and k_{el} of Cd and Zn varied more with the food matrix than with the

metal itself. With $Tb_{s1/2}$ of 13.0 ± 1.2 and 13.0 ± 0.9 h, for Cd and Zn, respectively, the short-

lived metal pool was 2 to 3 times longer in chironomid-fed gammarids than in leaf-fed

gammarids (Table 1). In addition, the long-term biological half-life $Tb_{l1/2}$ tended to infinity for

both elements when gammarids were fed with chironomids (Table 1). This strong retention of

metals is consistent with the previously stated hypothesis that the two food sources have

different TAM (Rainbow et al., 2011) which, once assimilated, should also determine the fate

of the metal in the gammarid. Thus, the leaf-adsorbed metal assimilated in a free form or in a

weak ionic complex is likely to be taken up more rapidly by detoxification and/or elimination

processes than the assimilated metal strongly bound to components and concretions of

chironomid cells. Nunez-Nogeira et al (2006) also found a higher k_e of Cd in the shrimp

P. indicus contaminated with squid meat $(0.009 \pm 0.003 \text{ d}^{-1})$ than when contaminated with

algae $(0.004 \pm 0.001 \text{ d}^{-1})$, confirming that Cd assimilated from a vegetable matrix is eliminated

more rapidly than from a complex multicellular organism. However, Zn was excreted more

rapidly after ingestion of animal than plant matrices (Nunez-Nogueira et al., 2006), contrary to

what we found in *G. fossarum*. These observations suggest that Zn and Cd undergo different

physiological bioaccumulation processes depending on the type of food ingested, hence the

importance of better understanding the toxicokinetics of ingested metals and what influences

it.

5. Conclusion

To better improve and predict the fate of metals in organisms and their potential transfer within food webs, it is important to understand the key factor controlling metal bioaccumulation in organisms, such as diet as a contamination pathway (Xu & Pascoe, 1994; Hadji et al., 2016). As a first step, the results presented here fill in the gaps regarding the critical physiological parameters governing bioaccumulation and trophic transfer mechanisms: assimilation efficiency (AE) and elimination rates. Furthermore, AE is a key input for the implementation of toxicokinetic models to study bioaccumulation mechanisms and the fate of metals in organisms (Van Campenhout et al., 2009). By exposing the gammarid, *G. fossarum*, to two types of food (i.e. of animal or plant origin) contaminated with Ag, Cd and Zn, this study reported that, regardless of the type of food, Cd had the highest AE values (39 and 19%, respectively), followed by Zn (15 and 9%, respectively) and Ag (5%, determined for plant only). In turn, the food matrix modulated the metal elimination rates, as k_{es} and k_{el} for Cd and Zn were higher when metals were ingested *via* leaves than *via* chironomids. This implies a shorter retention time for metals presumably adsorbed on an inert matrix compared to metals metabolically assimilated by live chironomids.

Thus, by providing AE estimates, we have shown that AE in gammarids depends on metals and food sources, as has already been demonstrated in fish (Pouil et al., 2018) or marine invertebrates (Bustamante et al., 2004). It is well known that the metal subcellular partitioning and its storage form in the food source is an important parameter affecting the AE of predators. Therefore, it would be interesting to demonstrate the relationship between bioavailable metal fractions in food (e.g. TAM, Rainbow et al., 2011) and AE in order to better predict trophic transfer based on food type.

Finally, the use of AE data could greatly improve toxicokinetic models to better understand the ADME (i.e. Absorption, Distribution, Metabolisation and Elimination) mechanisms of metals

in small invertebrates. In particular, the data obtained here has enabled us to improve our

knowledge and set up a toxicokinetic model of Cd accumulation *via* the trophic pathway (Gestin

et al., *submitted*). The study of organotropism over time in relation to the metal accumulation

pathway in gammarids would be an important direction for future research to further unravel

the bioaccumulation mechanisms (Franklin et al., 2005; Urien et al., 2016) and to implement

toxicokinetic models.

Acknowledgments

This work has been supported by the APPROve project funded by the ANR (ANR-18-CE34-

0013-01). We thank the « Radioecology lab » of the LIENSs and Competent Radiological

Protection Persons (Christine Dupuy and Thomas Lacoue-Labarthe) for their technical support.

This work benefitted from the French GDR "Aquatic Ecotoxicology" framework which aims

at fostering stimulating scientific discussions and collaborations for more integrative

approaches.

Funding

This work has been supported by the APPROve project funded by the ANR (ANR-18-CE34-

0013-01).

420
Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of

interest in relation to the content of the article.

References

Baines, S.B., & Fisher, N.S. (2002). Assimilation and retention of selenium and other trace
elements from crustacean food by juvenile striped bass (*Morone saxatilis*). *Limnology*
and Oceanography, *47*, 646–655. https://doi.org/10.4319/lo.2002.47.3.0646

Berthet, B., Amiard, J.C., Amiard-Triquet, C., Martoja, M., & Jeantet, A.Y. (1992). Bioaccumulation, toxicity and physico-chemical speciation of silver in bivalve molluscs: ecotoxicological and health consequences. *Science of the Total Environment*, *125*, 97–122. https://doi.org/10.1016/0048-9697(92)90385-6

Besse, J.P., Coquery, M., Lopes, C., Chaumot, A., Budzinski, H., Labadie, P., & Geffard, O. (2013). Caged *Gammarus fossarum* (Crustacea) as a robust tool for the characterization of bioavailable contamination levels in continental waters: Towards the determination of threshold values. *Water Research*, *47*, 650–660. https://doi.org/10.1016/j.watres.2012.10.024

Besse, J.P., Geffard, O., & Coquery, M. (2012). Relevance and applicability of active biomonitoring in continental waters under the Water Framework Directive. *TrAC - Trends in Analytical Chemistry*, *36*, 113–127. https://doi.org/10.1016/j.trac.2012.04.004

Borgmann, U., Couillard, Y., & Grapentine, L.C. (2007). Relative contribution of food and water to 27 metals and metalloids accumulated by caged *Hyalella azteca* in two rivers affected by metal mining. *Environmental Pollution*, *145*, 753–765. https://doi.org/10.1016/j.envpol.2006.05.020

Borgmann, U., Ralph, K.M., & Norwood, W.P. (1989). Toxicity test procedures for *Hyalella azteca*, and chronic toxicity of cadmium and pentachlorophenol to *H. azteca, Gammarus fasciatus*, and *Daphnia magna*. *Archives of Environmental Contamination and Toxicology*, *18*, 756–764. https://doi.org/10.1007/BF01225013

Bustamante, P., Teyssié, J.L., Danis, B., Fowler, S.W., Miramand, P., Cotret, O., & Warnau, M. (2004). Uptake, transfer and distribution of silver and cobalt in tissues of the common cuttlefish *Sepia officinalis* at different stages of its life cycle. *Marine Ecology Progress Series*, *269*, 185–195. https://doi.org/10.3354/meps269185

Calow, P., & Fletcher, C.R. (1972). A new radiotracer technique involving 14C and 51Cr, for estimating the assimilation efficiencies of aquatic, primary consumers. *Oecologia*, *9*, 155–170. https://doi.org/10.1007/BF00345880

Van Campenhout, K., Bervoets, L., Redeker, E.S., & Blust, R. (2009). A kinetic model for the relative contribution of waterborne and dietary cadmium and zinc in the common carp (*Cyprinus carpio*). *Environmental Toxicology and Chemistry*, *28*, 209–219. https://doi.org/10.1897/08-136.1

Conti, E., Dattilo, S., Costa, G., & Puglisi, C. (2016). Bioaccumulation of trace elements in the sandhopper *Talitrus saltator* (Montagu) from the Ionian sandy coasts of Sicily. *Ecotoxicology and Environmental Safety*, *129*, 57–65. https://doi.org/10.1016/j.ecoenv.2016.03.008

Cresswell, T., Metian, M., Golding, L.A., & Wood, M.D. (2017). Aquatic live animal radiotracing studies for ecotoxicological applications: Addressing fundamental methodological deficiencies. *Journal of Environmental Radioactivity*, *178–179*, 453–460. https://doi.org/10.1016/j.jenvrad.2017.05.017

Croteau, M.N., & Luoma, S.N. (2008). A biodynamic understanding of dietborne metal uptake by a freshwater invertebrate. *Environmental Science and Technology*, *42*, 1801–1806. https://doi.org/10.1021/es7022913

Croteau, M.N., & Luoma, S.N. (2009). Predicting dietborne metal toxicity from metal
influxes. *Environmental Science and Technology*, *43*, 4915–4921.
https://doi.org/10.1021/es9007454

Croteau, M.N., Misra, S.K., Luoma, S.N., & Valsami-Jones, E. (2011). Silver
bioaccumulation dynamics in a freshwater invertebrate after aqueous and dietary
exposures to nanosized and ionic Ag. *Environmental Science and Technology*, *45*,
6600–6607. https://doi.org/10.1021/es200880c

Dubois, M., & Hare, L. (2009). Subcellular distribution of cadmium in two aquatic
invertebrates: Change over time and relationship to Cd assimilation and loss by a
predatory insect. *Environmental Science and Technology*, *43*, 356–361.
https://doi.org/10.1021/es801406r

Dutton, J., & Fisher, N.S. (2011). Bioaccumulation of As, Cd, Cr, Hg(II), and MeHg in
killifish (*Fundulus heteroclitus*) from amphipod and worm prey. *Science of the Total*
Environment, *409*, 3438–3447. https://doi.org/10.1016/j.scitotenv.2011.05.022

Filipović Marijić, V., Dragun, Z., Sertić Perić, M., Matoničkin Kepčija, R., Gulin, V., Velki,
M., Ečimović, S., Hackenberger, B.K., & Erk, M. (2016). Investigation of the soluble
metals in tissue as biological response pattern to environmental pollutants (*Gammarus*
fossarum example). *Chemosphere*, *154*, 300–309.
https://doi.org/10.1016/j.chemosphere.2016.03.058

Fowler, S.W., Teyssié, J.L., Cotret, O., Danis, B., Rouleau, C., & Warnau, M. (2004).
Applied radiotracer techniques for studying pollutant bioaccumulation in selected marine
organisms (jellyfish, crabs and sea stars). *Nukleonika*, *49*, 97–100

Franklin, N.M., Glover, C.N., Nicol, J.A., & Wood, C.M. (2005). Calcium/cadmium
interactions at uptake surfaces in rainbow trout: Waterborne versus dietary routes of
exposure. *Environmental Toxicology and Chemistry*, *24*, 2954–2964.
https://doi.org/10.1897/05-007R.1

Gestin, O., Geffard, O., Delorme, N., Garnero, L., Lacoue-Labarthe, T., & Lopes, C. (2024).
Bioaccumulation, organotropism and fate of cadmium in Gammarus fossarum exposed
through dietary pathway. Manuscript submitted for publication.

Gestin, O., Lacoue-Labarthe, T., Coquery, M., Delorme, N., Garnero, L., Dherret, L., Ciccia,
T., Geffard, O., & Lopes, C. (2021). One and multi-compartments toxico-kinetic
modeling to understand metals' organotropism and fate in *Gammarus fossarum*.
Environment International, *156*.
https://doi.org/https://doi.org/10.1016/j.envint.2021.106625

Gestin, O., Lacoue-Labarthe, T., Delorme, N., Garnero, L., Geffard, O., & Lopes, C. (2023).
Influence of the exposure concentration of dissolved cadmium on its organotropism,
toxicokinetic and fate in *Gammarus fossarum*. *Environment International*, *171*.
https://doi.org/10.1016/j.envint.2022.107673

Gestin, O., Lopes, C., Delorme, N., Garnero, L., Geffard, O., & Lacoue-Labarthe, T. (2022).
Organ-specific accumulation of cadmium and zinc in *Gammarus fossarum* exposed to
environmentally relevant metal concentrations. *Environmental Pollution*, *308*.
https://doi.org/10.1016/j.envpol.2022.119625

Hadji, R., Urien, N., Uher, E., Fechner, L.C., & Lebrun, J.D. (2016). Contribution of aqueous
and dietary uptakes to lead (Pb) bioaccumulation in *Gammarus pulex*: From
multipathway modeling to in situ validation. *Ecotoxicology and Environmental Safety*,
129, 257–263. https://doi.org/10.1016/j.ecoenv.2016.03.033

Kunz, P.Y., Kienle, C., & Gerhardt, A. (2010). Gammarus *Spp. in Aquatic Ecotoxicology and*
Water Quality Assessment: Toward Integrated Multilevel Tests. In D. Whitacre (Ed.),
Reviews of environmental contamination and toxicology Springer, New York.

Lam, I.K.S., & Wang, W.-X. (2006). Accumulation and elimination of aqueous and dietary
silver in *Daphnia magna*. *Chemosphere*, *64*, 26–35.
https://doi.org/10.1016/j.chemosphere.2005.12.023

Lebrun, J.D., Geffard, O., Urien, N., François, A., Uher, E., & Fechner, L.C. (2015). Seasonal
variability and inter-species comparison of metal bioaccumulation in caged gammarids
under urban diffuse contamination gradient: Implications for biomonitoring
investigations. *Science of the Total Environment, The*, *511*, 501–508.
https://doi.org/10.1016/j.scitotenv.2014.12.078

Lebrun, J.D., Uher, E., Urien, N., & Tales, E. (2020). Ecological factors governing
distribution of gammarid species and their metal bioaccumulation abilities at the Seine
basin scale. *Ecological Indicators*, *118*, 106726.
https://doi.org/10.1016/j.ecolind.2020.106726

Memmert, U. (1987). Bioaccumulation of zinc in two freshwater organisms (Daphnia magna,
crustacea and Brachydanio rerio, pisces). *Water Research*, *21*, 99–106.
https://doi.org/10.1016/0043-1354(87)90104-7

Metian, M., Warnau, M., Oberhänsli, F., Teyssié, J.L., & Bustamante, P. (2007). Interspecific
comparison of Cd bioaccumulation in European Pectinidae (*Chlamys varia* and *Pecten*
maximus). *Journal of Experimental Marine Biology and Ecology*, *353*, 58–67.
https://doi.org/10.1016/j.jembe.2007.09.001

Nunez-Nogueira, G., Rainbow, P.S., & Smith, B.D. (2006). Assimilation efficiency of zinc
and cadmium in the decapod crustacean *Penaeus indicus*. *Journal of Experimental*
Marine Biology and Ecology, *332*, 75–83. https://doi.org/10.1016/j.jembe.2005.11.004

Pellet, B., Ayrault, S., Tusseau-Vuillemin, M.-H., & Gourlay-Francé, C. (2014). Quantifying
diet-borne metal uptake in *Gammarus pulex* using stable isotope tracers. *Ecotoxicology*
and Environmental Safety, *110*, 182–189. https://doi.org/10.1016/j.ecoenv.2014.09.001

Pouil, S., Bustamante, P., Warnau, M., & Metian, M. (2018). Overview of trace element
trophic transfer in fish through the concept of assimilation efficiency. *Marine Ecology*
Progress Series, *588*, 243–254. https://doi.org/10.3354/meps12452

Pouil, S., Warnau, M., Oberhänsli, F., Teyssié, J.L., Bustamante, P., & Metian, M. (2016).
Influence of food on the assimilation of essential elements (Co, Mn, and Zn) by turbot
Scophthalmus maximus. *Marine Ecology Progress Series*, *550*, 207–218.
https://doi.org/10.3354/meps11716

Pouil, S., Warnau, M., Oberhänsli, F., Teyssié, J.L., Bustamante, P., & Metian, M. (2017).
Comparing single-feeding and multi-feeding approaches for experimentally assessing
trophic transfer of metals in fish. *Environmental Toxicology and Chemistry*, *36*, 1227–

1234. https://doi.org/10.1002/etc.3646

Pouil, S., Warnau, M., Oberhänsli, F., Teyssié, J.L., & Metian, M. (2015). Trophic transfer of
110mAg in the turbot *Scophthalmus maximus* through natural prey and compounded
feed. *Journal of Environmental Radioactivity*, *150*, 189–194.
https://doi.org/10.1016/j.jenvrad.2015.08.016

Rainbow, P.S., Luoma, S.N., & Wang, W.X. (2011). Trophically available metal - A variable
feast. *Environmental Pollution*, *159*, 2347–2349.
https://doi.org/10.1016/j.envpol.2011.06.040

Reinfelder, J.R., & Fisher, N.S. (1991). The assimilation of elements ingested by marine
copepods. *Science*, *251*, 794–796. https://doi.org/10.1126/science.251.4995.794

Roditi, H.A., & Fisher, N.S. (1999). Rates and routes of trace element uptake in zebra
mussels. *Limnology and Oceanography*, *44*, 1730–1749.
https://doi.org/10.4319/lo.1999.44.7.1730

Timmermans, K.R., Spijkerman, E., Tonkes, M., & Govers, H.A.J. (1992). Cadmium and zinc
uptake by two species of aquatic invertebrate predators from dietary and aqueous
sources. *Canadian Journal of Fisheries and Aquatic Sciences*, *49*, 655–662.
https://doi.org/10.1139/f92-074

Urien, N., Farfarana, A., Uher, E., Fechner, L.C., Chaumot, A., Geffard, O., & Lebrun, J.D.
(2017). Comparison in waterborne Cu, Ni and Pb bioaccumulation kinetics between
different gammarid species and populations: Natural variability and influence of metal
exposure history. *Aquatic Toxicology*, *193*, 245–255.
https://doi.org/10.1016/j.aquatox.2017.10.016

Vellinger, C., Parant, M., Rousselle, P., Immel, F., Wagner, P., & Usseglio-Polatera, P.
(2012). Comparison of arsenate and cadmium toxicity in a freshwater amphipod
(*Gammarus pulex*). *Environmental Pollution*, *160*, 66–73.
https://doi.org/10.1016/j.envpol.2011.09.002

Wallace, W.G., & Lopez, G.R. (1996). Relationship between subcellular cadmium
distribution in prey and cadmium trophic transfer to a predator. *Estuaries*, *19*, 923–930.
https://doi.org/10.2307/1352308

Wallace, W.G., & Luoma, S.N. (2003). Subcellular compartmentalization of Cd and Zn in
two bivalves. II. Significance of trophically available metal (TAM). *Marine Ecology*
Progress Series, *257*, 125–137

Wang, W.-X., & Fisher, N.S. (1999)(a). Assimilation efficiencies of chemical contaminants in
aquatic invertebrates: A synthesis. *Environmental Toxicology and Chemistry*, *18*, 2034–
2045. https://doi.org/10.1897/1551-5028(1999)018<2034:AEOCCI>2.3.CO;2

Wang, W.-X., & Fisher, N.S. (1999)(b). Delineating metal accumulation pathways for marine
invertebrates. *The Science of the Total Environment*, *237/238*, 459–472.
https://doi.org/10.1016/s0048-9697(99)00158-8

Wang, W.-X., & Wong, R.S.K. (2003). Bioaccumulation kinetics and exposure pathways of
inorganic mercury and methylmercury in a marine fish, the sweetlips *Plectorhinchus*
gibbosus. *Marine Ecology Progress Series*, *261*, 257–268.
https://doi.org/10.3354/meps261257

Warnau, M., & Bustamante, P. (2007). Radiotracer techniques: A unique tool in marine
ecotoxicological studies. *Environmental Bioindicators*, *2*, 217–218.
https://doi.org/10.1080/15555270701714822

Warnau, M., Teyssié, J.L., & Fowler, S.W. (1996). Biokinetics of selected heavy metals and
radionuclides in the common Mediterranean echinoid *Paracentrotus lividus*: Sea water
and food exposures. *Marine Ecology Progress Series*, *141*, 83–94.
https://doi.org/10.3354/meps141083

Xu, Q., & Pascoe, D. (1994). The importance of food and water as sources of zinc during
exposure of *Gammarus pulex* (Amphipoda). *Archives of Environmental Contamination*
and Toxicology, *26*, 459–465. https://doi.org/10.1007/BF00214147

Yu, R., & Wang, W. (2002). Trace metal assimilation and release budget in *Daphnia magna*.
Limnology and Oceanography, *47*, 495–504