1 **Postsynaptic mitochondria are positioned to support functional diversity of dendritic spines**
2
3

4 Connon I. Thomas[1], Melissa A. Ryan[1,3], Naomi Kamasawa[1], Benjamin Scholl[2]
5
6
7 [1] Electron Microscopy Core Facility, Max Planck Florida Institute for Neuroscience, 1 Max Planck Way,
8 Jupiter, FL 33458, USA
9 [2] Department of Neuroscience, Perelman School of Medicine at the University of Pennsylvania, 415 Curie
10 Blvd, Philadelphia, PA, 19104, USA
11 [3] Present Address: Department of Neuroscience, Baylor College of Medicine, Houston, TX, 77030, USA
12
13 Corresponding author: Benjamin Scholl, benjamin.scholl@pennmedicine.upenn.edu

14
Abstract: 165
Introduction: 867
Methods: 1810
Results: 2548
Discussion: 1682
Legends: 722
Figures: 6
22

23 Acknowledgements: The authors thank Nicole Shultz and Rachel Satterfield for help with perfusions and
24 fixative preparation, Tal Laviv for comments, the Fitzpatrick lab for useful discussions, and the MPFI ARC
25 for animal care. The authors thank the GENIE project for access to GCaMP6.

26
27 Author contributions: B.S. and C.I.T. conceived experiments. B.S. performed biological experiments.
28 C.I.T. and M.A.R. performed electron microscopy and image processing with guidance from N.K.. C.I.T.,
29 M.A.R., and B.S. performed volumetric reconstruction. C.I.T. and B.S. analyzed data. C.I.T. and B.S.
30 wrote the paper with help from N.K. and M.A.R.
31
32 Competing financial interests: The authors declare no competing financial interests.
33

34 Funding Sources: NIH R00 EY031137 (B.S.), NIH R21 EY034992 (B.S.), Whitehall Award (B.S.), Max
35 Planck Florida Institute for Neuroscience (C.T., N.K.)

Abstract

Postsynaptic mitochondria are critical to the development, plasticity, and maintenance of synaptic inputs. However, their relationship to synaptic structure and functional activity is unknown. We examined a correlative dataset from ferret visual cortex with *in vivo* two-photon calcium imaging of dendritic spines during visual stimulation and electron microscopy (EM) reconstructions of spine ultrastructure, investigating mitochondrial abundance near functionally- and structurally-characterized spines. Surprisingly, we found no correlation to structural measures of synaptic strength. Instead, we found that mitochondria are positioned near spines with orientation preferences that are dissimilar to the somatic preference. Additionally, we found that mitochondria are positioned near groups of spines with heterogeneous orientation preferences. For a subset of spines with mitochondrion in the head or neck, synapses were larger and exhibited greater selectivity to visual stimuli than those without a mitochondrion. Our data suggest mitochondria are not necessarily positioned to support the energy needs of strong spines, but rather support the structurally and functionally diverse inputs innervating the basal dendrites of cortical neurons.

Introduction

By volume, mitochondria are the most prolific organelle in neuronal compartments. Mitochondria are observed within cell bodies, dendritic processes, axons, and occasionally dendritic protrusions (Cameron et al., 1991; Li et al., 2004; Kasthuri et al., 2015; Faitg et al., 2021; Pekkurnaz and Wang, 2022). Their abundance is thought to reflect a critical role: supporting the energy demands of neuronal communication through a supply of adenosine triphosphate (ATP) (Datta and Jaiswal, 2021). Neuronal communication occurs through synapses, and the process of transforming action potentials into graded postsynaptic depolarization is one of the most energy-expensive processes in the brain (Rossi and Pekkurnaz, 2019). ATP supplied by mitochondria is needed for postsynaptic glutamate receptor activation, presynaptic vesicle endo/exocytosis pathways, glutamate recycling, and Ca^{2+} channel activity (Attwell and Laughlin, 2001; Harris et al., 2012). Mitochondria are also involved in local protein synthesis, protein transport, and Ca^{2+} buffering critical for synapse maintenance and structural dynamics (Lipton and Whittingham, 1982; Li et al., 2004; Cheng et al., 2010; Harris et al., 2012; Rangaraju et al., 2014; Datta and Jaiswal, 2021). While there is growing evidence that mitochondria support synaptic activity, how this support is achieved is largely unknown. Some research suggests that ATP might diffuse from distant sources (Pathak et al., 2015), but the limited rate of diffusion of ATP in the cytosol implies the importance of spatial positioning close to synaptic compartments (Belles et al., 1987; Hubley et al., 1996; Devine and Kittler, 2018). Thus, it is unclear to what extent mitochondrial positioning is orchestrated within neuronal compartments to meet the needs of local synaptic inputs.

Adding further complexity, mitochondria are heterogeneous in morphology and density, in a manner
dependent on subcellular compartment (i.e. presynapse vs. postsynapse). Presynaptic mitochondria are
generally smaller, shorter, and less complex than those in dendrites of the same brain area (Delgado et
al., 2019; Faitg et al., 2021; Turner et al., 2022). Ultrastructural correlations reveal a strong link between
synaptic strength and mitochondrial abundance: larger boutons, which typically contain a greater number
of presynaptic vesicles and are correlated with synapse strength and efficacy (Schikorski and Stevens,
1997), contain larger mitochondria and, occasionally, a greater number of mitochondria (Wilke et al.,
2013; Cserép et al., 2018; Rodriguez-Moreno et al., 2018). Scaling to meet presynaptic energy demands
has also been shown in the calyx of Held nerve terminal where synapse functional maturation correlates
with a dramatic increase in mitochondrial volume and organization (Thomas et al., 2019). However, while
evidence grows for the specific recruitment of mitochondria to the presynapse, the nature of *postsynaptic*
mitochondria is largely unknown (Faitg et al., 2021).

Postsynaptic mitochondria are primarily located within the dendritic shafts, rarely found extending into
dendritic spines or inside the spine head (Cameron et al., 1991; Li et al., 2004; Kasthuri et al., 2015).
Dendritic mitochondria are largely thought to provide local energetic support, based on studies relating
mitochondrial and synapse abundance. Mitochondrial density along the dendrite co-varies with synapse
density in the neocortex (Turner et al., 2022). Additionally, dendritic mitochondria are found closer to
excitatory synapses than expected by chance in retinal ganglion cells, and a similar relationship has been
shown in cultured neurons (Li et al., 2004; Chang et al., 2006; Faits et al., 2016). *In vivo* measurements of
mitochondria Ca^{2+} influx correlate with cytosolic calcium influx (Datta and Jaiswal, 2021), further
supporting a view of supporting local synaptic activity via calcium buffering. However, a missing piece of
this framework is whether dendritic mitochondria are organized to support the structure and function of
individual dendritic spines and their synaptic activity.

A spatial relationship between mitochondria and dendritic spines may be critically important. Dendritic
spines exhibit a wide diversity of morphologies and functional response properties compared to the
somatic output (Scholl et al., 2021). Given the wide collection of synaptic inputs, conveying distinct
information (Scholl and Fitzpatrick, 2020) at different input rates (Goris et al., 2015), it is reasonable to
suggest that mitochondria are precisely positioned to support highly-active synapses onto dendritic spines
or those with greater influence on somatic action potentials. In addition, dendrites can exhibit
homogeneous or heterogeneous clustering of synaptic activity (Wilson et al., 2016), and functional
organization motifs may require differential energetic support. However, it is an open question whether
mitochondria are preferentially positioned to support certain populations of dendritic spines over others
based on their structure or sensory-driven activity.

We address this question by combining *in vivo* two-photon Ca^{2+} imaging of dendritic spines driven by

visual stimuli and serial block-face scanning electron microscopy (SBF-SEM) reconstructions of L2/3

pyramidal neurons of ferret visual cortex (Scholl et al., 2021). Using a deep learning method for

unsupervised 3D identification of mitochondria in the EM volumes, we characterized the 3D volume of

dendritic mitochondria near individual dendritic spines and compared it to structural and functional

measurements. While mitochondrial volume was unrelated to dendritic spine morphology, we discovered

a surprising functional correlation: local mitochondria were positioned near spines differentially-tuned to

the somatic output and those in regions with functional heterogeneity. Additionally, spines that contain a

mitochondrion in the head or neck were larger and more selective to visual stimuli. Our novel dataset and

analysis suggest that dendritic mitochondria are precisely positioned to support functional synaptic

diversity of cortical neurons.

Results

Characterizing postsynaptic mitochondria in 3D EM volumes from ferret visual cortex

To investigate subcellular organization of dendritic mitochondria, we returned to a previously published

CLEM dataset (Scholl et al., 2021). This dataset contains individual layer 2/3 neurons from the ferret

visual cortex for which basal dendrites were examined *in vivo* with two-photon calcium imaging of

GCaMP6s during visual stimulation, and then subsequently recovered for SBF-SEM and volumetric 3D

reconstruction to characterize ultrastructural anatomy. In this study, we continued to extract ultrastructural

information from the dendrites imaged *in vivo*, focusing on postsynaptic mitochondria. To identify

postsynaptic mitochondria, we used an unsupervised method for 3D identification of mitochondria in the

EM volumes (see Methods). This method quickly and accurately identified mitochondria within the

dendritic shaft (**Figure 1A**). From these 3D annotations, we first characterized the basic structure and

organization of postsynaptic mitochondria in our dataset (n = 324 mitochondria identified within 63

dendritic segments from 4 cells collected from 2 animals).

Mitochondria were distributed throughout the dendrites, with infrequent large (> 10 micron) gaps between

individual mitochondria (median inter-mitochondrion distance = 3.025 µm, IQR = 5.281 µm, **Figure 1A,**

B). Mitochondria in the basal dendrites of ferret layer 2/3 cortical cells were typically small (median

volume = 0.248 µm^3, IQR = 0.443 µm^3, surface area = 2.537 µm^2, IQR = 3.144 µm^2, **Figures 1D-E**), with

a short length through the dendrite (maximum length median = 1.137 µm, IQR = 0.881 µm, **Figures 1F)**,

in comparison to *in vitro* and *in vivo* reports of mouse cortical neurons (Lewis et al., 2018). While most

mitochondria were positioned within the dendritic shaft, we did observe a small fraction of mitochondrion

positioned either inside the spine neck or head (4.9%, 10/203 spines, **Figure 1C**). Of these spines,

mitochondria were found entirely within the spine head (2/10), at the junction between the head and neck

(5/10), or within the neck (3/10). Mitochondria in spines were significantly smaller and more spherical in

shape than dendritic mitochondria (median volume = 0.050 μm^3, IQR = 0.047 μm^3, surface area = 0.658

μm^2, IQR = 0.540 μm^2, maximum length = 0.590 μm, IQR = 0.249 μm; all p < 0.0001, Mann-Whitney *U*

tests; **Figures 1D-F**). These mitochondria were never continuations of dendritic mitochondria.

Dendritic mitochondria organization is independent of dendritic spine morphology

We hypothesized that dendritic mitochondria are strategically positioned to support synapses on dendritic

spines in a structural- or activity-dependent manner, as found for axons (Smith et al., 2016). To test this

hypothesis, we measured the total mitochondria volume within a given distance from the base of the

spine neck (**Figure 2A**). We chose this metric over measuring distance to the nearest mitochondrion for

three reasons: (1) distance to a mitochondrion is ill defined and could be measured from either the center

or nearest membrane, (2) larger mitochondria produce more ATP and provide a greater capacity for

calcium buffering to support nearby synapses, and (3) multiple mitochondria were occasionally clustered

nearby individual spines. In presynaptic boutons, a critical inflection-point distance has been identified,

where, the presence of a mitochondrion less than 3 μm from the synapse results in an increase in active

zone size and docked vesicle number (Smith et al., 2016). Based on this observation, we measured

mitochondria volume using two radii: 1 μm and 5 μm (**Figure 2A**). In our dataset, the median spine neck

length is roughly 1.8 μm; therefore, 1 and 5 μm distances represent short and extended ranges relative to

the synapse. In our population, 56.8% of spines had no mitochondria volume within 1 μm and 12.1% of

spines had none within 5 μm. Expectedly, there was greater mitochondria volume for a 5 μm radius

compared to 1 μm nearby individual spines (1 μm radius median volume = 0.221 μm^3, IQR = 0.278 μm^3; 5

μm radius median volume = 0.771 μm^3, IQR = 0.561 μm^3; n = 69 spines, p < 0.001, Mann-Whitney *U*

test), although these two measures were significantly correlated (Spearman's r = 0.239, p = 0.0238;

Figure 2B). For all subsequent analyses unless otherwise stated, we focused *only* on spines with

measurable mitochondrial volume at each radius.

Dendritic spines differ in shape and size (e.g. Figure 1B). As spine structure is highly correlated with

synaptic strength (Bartol et al., 2015; Holler et al., 2021), we first examined whether mitochondria were

preferentially located nearby stronger or weaker synapses, focusing on spines without a mitochondrion in

the head or neck. Spine head volume and postsynaptic density (PSD) area, features that are positively

correlated with synapse strength, were uncorrelated with mitochondria volume in a 1 μm radius (Spine

head volume: r = -0.032, p = 0.403, n = 60; PSD area: r = 0.066, p = 0.309, n = 60 spines; **Figure 2C, G**)

or a 5 μm radius (Spine head volume: r = -0.034, p = 0.357, n = 121; PSD area: r = -0.072, p = 0.216, n =

121 spines; **Figure 2D, H**). Spine neck length was also uncorrelated with mitochondria volume at both

radii (1 μm: r = 0.185, p = 0.079, n = 60 spines; 5 μm: r = 0.040, p = 0.333, n = 121 spines; **Figure 2E, F**).

We did observe a positive correlation between local 1 μm radius mitochondria volume and the distance of

a dendritic spine from the cell body (r = 0.325, p = 0.006, n = 60 spines; **Figure 2I**). While this relationship

was not observed for mitochondria within a 5 μm radius (5 μm: r = 0.104, p = 0.129, n = 121 spines;

Figure 2J), there is a simple explanation: the volume fraction of dendrite occupied by mitochondria

increases with distance from the soma (Turner et al., 2022). Dendrites are thicker near the soma and

possess few dendritic spines.

Dendritic mitochondria are positioned near dendritic spines with diverse visual response properties

As dendritic mitochondria position was unrelated to dendritic spine anatomy, we next considered the

visually-driven calcium activity of individual dendritic spines. In ferret V1, neurons and dendritic spines

have exquisite response specificity to the orientation of drifting gratings (Wilson et al., 2016). Somatic

orientation tuning is thought to arise from the co-activation of inputs sharing similar tuning (Scholl et al.,

2021). Given the importance of co-tuned inputs supporting somatic selectivity, we hypothesized that

dendritic mitochondria would be strategically positioned for energetic support and maintenance of co-

tuned inputs.

To test this hypothesis, we examined spines in our EM volumes which were also examined with *in vivo*

two-photon calcium imaging during the presentation of oriented drifting gratings (**Figure 3A**). As

described previously (Scholl et al., 2021), evoked calcium signals were measured to characterize the

orientation tuning of individual dendritic spines and the corresponding somatic output (**Figure 3B**). Each

of the 4 cells in our dataset was similarly tuned for orientation (selectivity range = 0.40 – 0.57, mean =

0.46 $\pm$ 0.08 s.d.). For each tuned spine (selectivity > 0.1, n = 148/164 spines; see Methods), we

calculated orientation preference and compared this value to the orientation tuning of each parent soma

($\Delta\theta_{pref}$; see Methods). As shown in Figure 3B, spines can be co-tuned with the soma or differentially-

tuned. We found that mitochondrial volume within a 1 µm radius was correlated with difference in

orientation preference relative to the soma (bootstrapped spearman's r = 0.315 $\pm$ 0.11 s.e.m., p = 0.010, n

= 54 spines; **Figure 3C**). This relationship could not be attributed to different properties of co-tuned ($\Delta\theta_{pref}$

< 22.5˚, n = 68) and differentially-tuned spines ($\Delta\theta_{pref}$ > 67.5˚, n = 30) (selectivity: p = 0.12, max response

amplitude: p = 0.58). When increasing the radius to 5 µm, however, no correlation was observed

(bootstrapped Spearman's r = 0.017$\pm$ 0.10 s.e.m., p = 0.432, n = 107 spines; **Figure 3D**). This trend was

found across all 4 cells examined (1 µm: mean Spearman's r = 0.49 ± 0.22 s.d.; 5 µm: mean Spearman's

r = 0.09 ± 0.13 s.d.). We also considered other spine response properties related to tuning preference;

specifically, orientation selectivity and response amplitude at the preferred orientation. For either metric,

we observed no relationship to mitochondria within 1 µm radius (selectivity: 1 µm: Spearman's r = -0.081,

p = 0.269, n = 60; max response amplitude: 1 µm: Spearman's r = -0.179, p = 0.078, n = 64) but did see a

weak, significant relationship to both at a 5 µm radius (selectivity: Spearman's r = 0.175, p = 0.027, n =

121; max response amplitude: Spearman's r =-0.166, p = 0.030, n = 129). This weak relationship

suggests that within some dendritic segments, a greater volume of mitochondria leads to rapid removal of

calcium out of the spine and cytoplasm, in agreement with a recent modeling study (Leung et al., 2021).

We next wondered whether mitochondria might be positioned near clusters of spines with diverse

orientation preferences. Within the same dendritic segment, local populations of spines can either share

the same orientation (homogeneous; (Scholl et al., 2017) or exhibit diversity (heterogeneous; (Wilson et

al., 2016). We measured the average orientation difference (i.e. local heterogeneity) of all spines within a

5 µm neighborhood of a given spine (**Figure 4A**; see Methods). We choose 5 µm because this value is

strongly associated with the spatial extent of synaptic clustering and provides adequate sample sizes for

computing the average orientation differences (n > 2 spines). Similar to spine-soma difference in

orientation preference, we found that local heterogeneity was strongly correlated with mitochondria

volume within a 1 µm radius (r = 0.563, p = 0.0005, n = 31 spines; **Figure 4B**), and not within 5 µm (r =

0.017, p = 0.432, n = 107 spines; **Figure 4C**). Importantly, while local heterogeneity was correlated with

spine co-tuning (circular r = 0.33, p = 0.0024, n = 100 spines), this correlation was entirely due to inputs

that closely matched the somatic preference. Restricting to only differentially-tuned inputs ($\Delta\theta_{pref} > 11.25$

deg) yielded no correlation between spine-soma similarity and local heterogeneity (circular r = 0.03, p =

0.82, n = 71 spines). Focusing on these spines, local heterogeneity was still strongly correlated with

mitochondria volume within 1 µm (Spearman's r = 0.52, p = 0.0052, n = 50).

Our data show that dendritic mitochondria are not observed near co-tuned spines or homogeneous

clusters, but these measurements are based on orientation tuning preference, a metric that does not

necessarily reflect local synaptic activity. Thus, we wondered whether a similar relationship exists for

correlated activity between spines or the total calcium activity from groups of spines. As functional

similarity coincides with spine co-activity (Scholl et al., 2017), dendritic mitochondria should not be

positioned near spines with greater co-activity or total activity. We examined both the average local

temporal correlation of calcium activity (**Figure 4D**) and trial-by-trial average calcium activity for all spines

within a 5 µm radius of a given spine (**Figure 4G**; see Methods). Local temporal correlation was

calculated using the average Pearson correlation coefficient of time-varying ΔF/F traces between a target

spine and all neighboring spines within 5 µm. We found no relationship for either measurement within a 1

µm radius (local temporal correlation: r = 0.015, p = 0.469, n = 28 spines; local average calcium activity: r

= 0.160, p = 0.131, n = 51 spines; **Figure 4E, H**), nor for local temporal correlation at a 5 µm radius (r = -

0.152, p = 0.141, n = 52 spines; **Figure 4F**). We did, however, observe a considerable negative

correlation between mitochondria volume within a 5 µm radius and local average calcium activity (r = -

0.320, p = 0.0004, n = 105 spines; **Figure 4I**). As this correlation was only observed at an extended

radius, it suggests that dendritic regions with a particularly large volume of mitochondria, calcium is

quickly buffered and calcium activity is attenuated within dendritic spines.

Dendritic spines with mitochondria are large and have high tuning selectivity

Our analyses have revealed that dendritic mitochondria are proximal to differentially-tuned spines and

dendritic regions of local heterogeneity. These analyses have excluded dendritic spines containing

mitochondria in the spine neck or head (**Figure 1C**). We then examined whether these unique spines (n =

10) displayed distinct structural or functional properties.

Spines with a mitochondrion in the head or neck had a significantly larger spine head volume compared

to the majority of spines that do not (mito+: median = 0.608 μm^3, n = 10 spines; mito–: median = 0.311

μm^3, n = 157 spines, p = 0.0014, Mann-Whitney *U*-test, p = 0.01, permutation test; **Figure 5A**). A similar

trend was observed for PSD area, but it was not significant (mito +: median = 0.396 μm^2, n = 10 spines;

mito–: median = 0.244 μm^2, n = 157 spines, p = 0.087, Mann-Whitney *U*-test, p = 0.12, permutation test;

Figure 5B). Other anatomical properties such as spine neck length (mito +: median = 1.850 μm, n = 10

spines; mito–: median = 1.65 μm, n = 157 spines, p = 0.382, Mann-Whitney *U*-test, p = 0.14, permutation

test; **Figure 5C**) and distance of the spine from the soma (mito +: median = 52.32 μm, n = 10 spines;

mito–: median = 50.56 μm, n = 157 spines, p = 0.898, Mann-Whitney *U*-test, p = 0.42, permutation test;

Figure 5D) showed no difference between spine groups. Given the difference in size between spines with

and without a mitochondrion, we wondered about spines nearby a dendritic mitochondrion (within 1 μm, n

= 65) compared to those with no measurable mitochondria within 5 μm (n = 19). For these groups, we

found no difference in spine volume (within 1 μm: median = 0.29 μm^3, IQR = 0.41 μm^3; no mitochondria

within 5 μm: median = 0.40 μm^3, IQR = 0.37 μm^3; Mann-Whitney ranksum test, p = 0.67) or PSD area

(within 1 μm: median = 0.26 μm^2, IQR = 0.33 μm^2; no mitochondria within 5 μm: median = 0.31 μm^2, IQR

= 0.36 μm^2; Mann-Whitney ranksum test, p = 0.49).

Comparing other functional response properties, we found only a single metric showing a significant

difference: orientation tuning selectivity. Spines with a mitochondrion were more selective for oriented

gratings (mito +: median = 0.396, n = 10 spines; mito–: median = 0.268, n = 161 spines, p = 0.016, Mann-

Whitney U-test, p = 0.03, permutation test; **Figure 6A**). Unlike for dendritic mitochondria, we saw no

difference in spine-soma difference of orientation preference (mito +: median = 10.62, n = 10 spines;

mito–: median = 30.17, p = 0.451, n = 146 spines, Mann-Whitney U-test, p = 0.23, permutation test;

Figure 6B). Again, we made a similar examination of spines near dendritic mitochondria and those

without mitochondria within 5 μm. We found no difference orientation selectivity (1 μm: median = 0.29,

IQR = 0.28; no mitochondria within 5 μm: median = 0.28, IQR = 0.15; Mann-Whitney ranksum test, p =

0.74) or mismatch to soma orientation (1 μm: median = 0.54 deg, IQR = 0.86 deg; no mitochondria within

5 μm: median = 0.46 deg, IQR = 0.47 deg; Mann-Whitney ranksum test, p = 0.75). Taken together, these

data suggest mitochondria positioned within dendritic spines may serve a different purpose than those

within the dendrites; supporting stronger synaptic inputs, which have been shown to be more selective for

features of visual stimuli (Scholl et al., 2021).

Discussion

We used a correlative light and electron microscopy dataset (Scholl et al., 2021) to characterize
mitochondria within basal dendrites of L2/3 pyramidal neurons of the ferret V1 and relate their abundance
near spines to an array of ultrastructural and functional properties of those spines. Based on 3D
reconstructions using deep learning segmentation, mitochondria were found throughout the dendrites and
had a fragmented morphology similar to those observed recently in CA1 pyramidal neuron basal
dendrites (Virga et al., 2023). We report no relationship to measures of synaptic strength such as spine
head volume or PSD area to dendritic mitochondrial volume within either a short or extended (1 or 5 μm,
respectively) distance from the base of the spine neck. We did observe significant correlations between
the volume of nearby mitochondrial and functional diversity–for spine orientation preference similarity to
the soma, and for local dendritic orientation preference heterogeneity. We also found that spines having
mitochondria within the head or neck have significantly larger heads than those that do not and are more
selective for visual stimuli, but are no different in other measures of structure and function. Taken
together, our results point toward a system where dendritic mitochondria are positioned to support
functional diversity, but not synaptic strength as has been previously shown in the presynaptic terminal.
Furthermore, though rare, mitochondria within spines do support synaptic strength and selectivity,
suggesting that mitochondria are flexible to support different facets of postsynaptic signaling depending
on the compartment in which they are present.
300
301   *Mitochondria and dendritic spine anatomy*
302
In this study, we investigated whether dendritic mitochondria are preferentially positioned to support
certain populations of dendritic spines over others, based on their structural or sensory-driven activity. We
were surprised to find that dendritic mitochondria were not positioned near strong synapses, as measured
by spine volume or PSD area. Considering that glutamate receptors and ion channels might take up 42%
of the O2 consumed by oxidative phosphorylation, and that larger spines have greater quantities of
receptors (Holtmaat and Svoboda, 2009), we assumed that mitochondria would need to be positioned
nearby for oxidative support. This is in contrast to studies of the presynapse: presynaptic terminal active
zone area and mitochondria volume are strongly correlated, and mitochondria proximity is correlated with
the number of nearby docked vesicles (Smith et al., 2016; Thomas et al., 2019). Our results suggest
dendritic mitochondria are not necessarily positioned to support the energy needs of large spines (i.e.
strong synapses) through oxidative phosphorylation, at least in the pyramidal neuron basal dendrites we
observed, and opens new questions into the nature of mitochondria.
315
316   *Mitochondria and spine functional properties*
317
We hypothesized that dendritic mitochondria would be positioned near synaptic inputs co-tuned with the
somatic output (i.e. similar orientation preference). Our rationale was that co-activation of co-tuned inputs

are critical for shaping somatic selectivity, and these inputs can often be found to be clustered together to

trigger nonlinear dendritic events (Wilson et al., 2016; Scholl et al., 2017, 2021). Surprisingly, we found

the opposite: mitochondria are present in greater volumes near differentially-tuned spines. This includes

spines that had an orientation preference mismatching the soma and spines within dendritic regions of

heterogeneity in orientation preference. What might be the cause of such a striking deviation from our

original hypothesis?

One possibility is that regions of high spine functional diversity are undergoing plasticity, such as spine

pruning, *de novo* formation, or strength modifications. Recent *in vivo* studies have demonstrated that

mitochondria are recruited to dendritic regions undergoing structural dynamics (Faits et al., 2016;

Dromard et al., 2021); however, it is unclear whether this is the case for our own EM dataset as we were

only able to study a single developmental time point (adolescent, visually-mature ferret). Another

possibility is that differentially-tuned spines and heterogeneous clusters reside nearby a greater number

of GABAergic inputs, supported energetically or buffered by local mitochondria, which could thereby act

to sculpt away excitatory input. It is also possible that co-tuned spines will be more co-activated with and

infiltrated by back-propagating somatic action potentials (Palmer and Stuart, 2009; Popovic et al., 2015),

as compared to differentially-tuned spines. Back-propagating action potentials may alter spine head

depolarization, PSD composition, or calcium dynamics; differences in spine activation may lead to

different levels of support needed from mitochondria. Finally, it is possible our observations are specific

to the ferret or visual cortex. Unfortunately, there are not enough reported measurements of postsynaptic

mitochondria to compare between animal models, let alone brain areas, and it remains to be shown if our

results will hold for layer 2/3 pyramidal neurons in mouse V1. Nevertheless, as ferrets are a highly visual

animal with a primate-like organization of the early visual system, our observations may reflect specific

visual processing demands (e.g. extracting motion information from incoming signals).

Of course, there are limitations precluding our ability to assess dendritic spine structure-function

relationships. Visual cortical neurons exhibit a high degree of functional heterogeneity and we probed

only a single stimulus dimension (orientation). *In vivo* spine calcium imaging is biased towards larger

spines and can only resolve synaptic events involving NMDA receptors or voltage-gated calcium

channels (Scholl et al., 2021). Spine calcium signals are likely not to directly reflect underlying electrical

signals (Koester and Johnston, 2005; Sobczyk et al., 2005). In addition, the kinetics of GCaMP6s make it

difficult to characterize other functional properties of synapses such activity frequency during visual

stimulation. In summary, although we cannot provide a clear reason for our findings, our data highlight the

need for greater investigation into dendritic mitochondria, their subcellular organization within cells, and

their relationship to patterns of synaptic activity.

Mitochondrial structure and distribution

Several factors influence the size and abundance of neuronal mitochondria including brain region,

neuronal compartment, cell type, animal age, and health (Delgado et al., 2019; Glancy et al., 2020).

Despite this variability, dendritic mitochondria are generally longer and take up a greater fraction of

neurite volume compared to axonal mitochondria, which tend to be more fragmented and, in some cases,

more mobile. Previous reports in mouse hippocampal cell culture and rat hippocampus report dendrites

with long, overlapping mitochondria that form stable compartments of ~35 µm length or more (Popov et

al., 2005; Rangaraju et al., 2019). Studies of mouse cortex using EM or light microscopy report

mitochondria lengths of roughly 1 to 10 µm (Lewis et al., 2018; Turner et al., 2022). Our results are closer

to those of mouse cortex, but are smaller (median length just over 1 µm, 10th percentile = 0.55 µm, 90th

percentile = 2.49 µm). The mitochondrial morphology we observe is most similar to a recent report from

mouse CA1 pyramidal neuron basal dendrites, where mitochondria are ~1 µm long and have a similar

size distribution (Virga et al., 2023). This similarity could be due to the fact that dendritic mitochondrial

morphology is cell type and neuronal-compartment specific (e.g. apical vs. basal dendrites). Another

possibility is mitochondrial fragmentation, as we observe in layer 2/3 ferret basal dendrites, is linked to

synaptic excitability. In support of this hypothesis, Virga et al. (2023) demonstrated that low levels of

excitability and activity is linked to longer mitochondria. Thus, presynaptic neurons driving basal synapses

in carnivore V1 may operate at relatively high firing rates (Mante et al., 2008), leading to more fragmented

mitochondria than compared to mouse V1.

Another variable feature across brain regions is the presence of mitochondria within spines. In

hippocampal cell cultures, mitochondria enter about 8-10% of spines, typically as extensions of a

dendritic mitochondrion (Li et al., 2004). EM reconstructions of mouse neocortex showed only 0.21% of

spines (n = 3/1,425) contained mitochondria (Kasthuri et al., 2015). In contrast, for olfactory bulb granule

cells, ~87% of spines contain mitochondria (Cameron et al., 1991). We found that 4.9% of spines contain

mitochondria, however, since we specifically examined layer 2/3 neuron basal dendrites proximal to the

soma, it is difficult to ascertain where this value stands within the literature and future studies will need to

investigate variations due to brain area, cell type, and compartment. Another interesting distinction in our

dataset was that spine mitochondria were roughly spherical or oblong, never continuations of a dendritic

mitochondrion. It is likely that these mitochondria are small fragmentations of nearby dendritic

mitochondria trafficked into the spine, possibly through interactions with the cytoskeleton (MacAskill and

Kittler, 2010), supporting the idea that spine mitochondria serve different role than dendritic mitochondria.

Previous observations in cultured neurons suggest that dendritic mitochondria are stable, and similar

evidence has been shown in retinal ganglion cell dendrites from the intact retina (Chang et al., 2006; Faits

et al., 2016; Rangaraju et al., 2019). Experiments have also shown oxidative phosphorylation by

mitochondria is needed for dendritic protein translation, not glycolysis (Rangaraju et al., 2019), and that it

contributes a majority of the energy consumed at the postsynaptic site by receptor and ion channel

activation (Harris et al., 2012). However, taking our observations into account, it's difficult to conceive

such a system could exist where mitochondria are simultaneously immobile and provide significant

quantities of energy for synaptic activity through oxidative phosphorylation, yet have no positional bias

towards large, strong synapses. We expect there exists a balance between oxidative phosphorylation and

glycolysis supplying energy at the synapse. This balance could shift depending on brain region, cell type,

compartment, activity level, and mitochondrial morphology and mobility. In this way, dendritic areas

receiving synaptic inputs with high functional diversity require greater local mitochondrial ATP production,

while baseline activity is supported through local glycolysis or ATP diffusion from more distant sources.

Additionally, the role that mitochondria serve at the synapse may shift toward one of calcium buffering

and plasticity support. Local depletion of mitochondria has been shown to reduce plasticity (Rangaraju et

al., 2019), and by blocking fission, spines have impaired structural and functional LTP. Induction of LTP

causes increased dendritic mitochondrial Ca^{2+} and a burst of fission (Divakaruni et al., 2018). Thus, once

spines become established, mitochondria may localize to input-diverse dendritic regions to support

synaptic plasticity. More investigation into the support functions of postsynaptic mitochondria is needed to

elucidate their different contributions to synaptic function, plasticity, and development.

Figure 1: Volumetric characterization of postsynaptic mitochondria in L2/3 pyramidal neurons of ferret
visual cortex. ***A.*** Example EM reconstruction of a dendrite with mitochondria (purple). ***B.*** Cumulative
frequency distribution of inter-mitochondria distance. ***C.*** EM reconstructions of all functionally-
characterized spines that contain mitochondria in the head or neck of the spine. Mitochondria were
observed either fully within the head, at the base of the head near the opening of the neck, or in the neck.
Electron micrographs show cross sections through three example spines having mitochondria
(arrowheads) at the base of the head (purple), fully within the head (yellow), or in the neck (pink). ***D.***
Cumulative frequency distribution of mitochondria volume, comparing mitochondria found in dendrites
(gray) versus those found in spines (black). ***E.*** Mitochondria surface area ***F.*** Maximum mitochondria
length. Mann-Whitney *U* tests; ****p < 0.0001.

Figure 2: Dendritic mitochondria are positioned to support a diversity of anatomical strengths. ***A.***
Schematic illustration of the method used to measure dendritic mitochondria volume near structurally and
functionally characterized spines in EM reconstructions. A sphere of a short (1 µm, blue) or extended
radius (5 µm, orange) from the base of the spine neck (black dot) was used to sample the nearby volume
of mitochondria for each spine. ***B.*** Correlation plot of the volume of mitochondria present within short and
extended radii for individual spines. Only spines that had dendritic mitochondria within 5 µm were
considered. ***C, D.*** Correlation of spine volume and mitochondria volume within a short or extended
distance. ***E, F.*** Correlation of spine neck length and mitochondria volume. ***G, H.*** Correlation of PSD area
and mitochondria volume. ***I, J.*** Correlation of the distance between a spine/soma and mitochondria
volume. r = Spearman's correlation coefficient. Correlation plots exclude spines that have zero
mitochondria volume within the corresponding measurement radius.

Figure 3: Dendritic mitochondria are positioned to support spines with a diversity of orientation tuning. ***A.***
2-photon average projection image of a short segment of GCaMP6s-labeled dendrite and the
corresponding EM reconstruction. Functionally characterized spines are indicated with rainbow
colors/arrows, mitochondria in the EM reconstruction are purple. ***B.*** Peak ΔF/F responses in response to
visual stimulus for two example spines shown in ***A*** (teal and indigo), showing two different orientation
preferences. The teal spine has a preference that is aligned with the average preference of the soma
(vertical gray bar), while the indigo spine has a preference that is dissimilar to the soma. Spine data are
mean ± s.e.m. (n = 8 stimulus trials). ***C, D.*** Correlation of spine-soma difference of orientation preference
and mitochondria volume for short (1 μm, blue) and extended (5 μm radius, orange) distances from the
base of a spine. r = Spearman's correlation coefficient. Correlation plots exclude spines that have zero
mitochondria volume within the corresponding measurement radius.

Figure 4: Dendritic mitochondria are positioned locally in regions having diverse synaptic orientation

preferences and broadly in areas of low local Ca^{2+} activity. **A.** Illustration of average difference in

orientation preference between a target spine (arrowhead) and neighboring spines within 5 µm. **B, C.**

Correlation of local orientation difference and mitochondria volume for short (1 µm, blue) and extended (5

µm radius, orange) distances form the base of a spine. **D.** Illustration of average local temporal correlation

of spines within 5 µm of the target spine. Scale bar is 1 sec and 100% ΔF/F. **E, F.** Correlation of local

temporal correlation and mitochondria volume. **G.** Illustration of local Ca^{2+} activity, shown as mean trial-to-

trial ΔF/F, within 5 µm of the target spine. **H, I.** Correlation of local average Ca^{2+} activity and mitochondria

volume. r = Spearman's correlation coefficient. Correlation plots exclude spines that have zero

mitochondria volume within the corresponding measurement radius.

Figure 5: Spines having a mitochondrion in the head or neck are larger. **A.** Cumulative frequency

distribution of spine head volume for spines that have a mitochondrion in the head or neck (black, n = 10

spines, shown in Figure 1C) versus spines that do not (gray). **B.** PSD area. **C.** Neck length. **D.** Distance

between the spine and soma. Mann-Whitney *U* tests, **p < 0.01.

Figure 6: Spines having a mitochondrion in the head or neck are more selective for visual features. **A.**

Cumulative frequency distribution of orientation selectivity for spines that have a mitochondrion in the

head or neck (black, n = 10 spines, shown in Figure 1C) versus spines that do not (gray). **B.** Spine-soma

difference of orientation preference. Mann-Whitney *U* tests, *p < 0.05.

Methods

All procedures were performed according to NIH guidelines and approved by the Institutional Animal Care
and Use Committee at Max Planck Florida Institute for Neuroscience.

Animals, Viral Injections, and Cranial Windows

Information on animals, survival viral injections, cranial window implantation are described in depth in our
previous studies (Scholl et al., 2017, 2019, 2021). Briefly, layer 2/3 neurons of the primary visual cortex of
female ferrets (n = 3, Marshall Farms) were driven to express the calcium indicator GCaMP6s via a Cre-
dependent AAV expression system. On the day of imaging, a cranial window was surgically implanted to
image calcium activity of cortical pyramidal neurons during presentation of visual stimuli.

Two-Photon Imaging

Two-photon imaging was performed using a Bergamo II microscope (Thorlabs) running Scanimage
(Vidrio Technologies) with 940 nm dispersion-compensated excitation provided by an Insight DS+
(Spectraphysics). For spine and axon imaging, power after the objective was limited to <50 mW. Images
were collected at 30 Hz using bidirectional scanning with 512 × 512 pixel resolution or with custom ROIs
(region of interest; framerate range: 22 - 50 Hz). Somatic imaging was performed with a resolution of
0.488 - 0.098 µm/pixel. Dendritic spine imaging was performed with a resolution of 0.164 - 0.065
µm/pixel.

Visual Stimuli

Visual stimuli were generated using Psychopy (Peirce, 2007). The monitor was placed 25 cm from the
animal. Receptive field locations for each cell were hand mapped and the spatial frequency optimized
(range: 0.04 to 0.20 cpd). For each soma and dendritic segment, square-wave or sine-wave drifting
gratings were presented at 22.5 degree increments to each eye independently (2 second duration, 1
second ISI, 8-10 trials for each field of view). Drifting gratings of different directions (0 – 315°) were
presented independently to both eyes with a temporal frequency of 4 Hz.

Two-Photon Imaging Analysis

Imaging data were excluded from analysis if motion along the z-axis was detected. Dendrite images were
corrected for in-plane motion via a 2D cross-correlation based approach in MATLAB or using a piecewise

non-rigid motion correction algorithm (Pnevmatikakis and Giovannucci, 2017). ROIs were drawn in

ImageJ; dendritic ROIs spanned contiguous dendritic segments and spine ROIs were fit with custom

software. Mean pixel values for ROIs were computed over the imaging time series and imported into

MATLAB. ΔF/F was computed with a time-averaged median or percentile filter (10th percentile). We

subtracted a scaled version of the dendritic signal to remove back-propagating action potentials as

performed previously (Wilson et al., 2016). ΔF/F traces were synchronized to stimulus triggers sent from

Psychopy and collected by Spike2.

Peak responses to bars and gratings were computed using the Fourier analysis to calculate mean

and modulation amplitudes for each stimulus presentation, which were summed together. Spines were

included for analysis if the mean peak for the preferred stimulus was >10%, the SNR at the preferred

stimulus was > 1, and spines were weakly correlated with the dendritic signal (Spearman's correlation, r <

0.4). Some spine traces contained negative events after subtraction, so correlations were computed

ignoring negative values. Preferred orientation for each spine was calculated by fitting responses with a

double Gaussian tuning curve (Wilson et al., 2016) using lsqcurvefit (Matlab). Orientation selectivity was

computed by calculating the vector strength of mean responses (Wilson et al., 2016).

For each tuned spine (orientation selectivity > 0.1), we calculated orientation preference

difference between the spine and parent soma as in Wilson et al. (2016). To measure the average

orientation difference (i.e. local heterogeneity), we calculated the orientation preference difference

between a target spine and all surrounding spines within 5 microns on the same dendrite. For targets with

n > 2 neighbors, we then computed the circular mean. Average spatiotemporal correlation for a given

spine was calculated similarly: the Pearson correlation between time-varying ΔF/F traces was computed

between a target spine and all neighbors within 5 µm, and then we calculated the average of this

population. To calculate trial-by-trial average calcium activity, we averaged the amplitudes of evoked

calcium transients between a target spine and neighbors within 5 µm for all visual stimulus trials. A grand

mean was then computed across all trials.

Perfusion, fixation, and slice preparation for fluorescence imaging

Anesthetized animals were immediately perfused with 2% paraformaldehyde and 2 – 2.5%

glutaraldehyde in a 0.1 M sodium cacodylate buffer (pH 7.4). Following removal, brains were sliced at 80

µm parallel to the area flattened by the cranial window. Slices were quickly imaged at low magnification

(20x, 0.848 × 0.848 µm/pixel) using a Leica CLSM TCS SP5 II running LAS AF (ver. 3.0, Leica) with 488

nm laser excitation. Fluorescence of GCaMP6s was used to locate the target cell in this view.

Autofluorescence resulting from glutaraldehyde fixation also produced signal within the tissue slices. A

field of view large enough to cover roughly one-fourth of the slice, with the target cell included, was

captured using image tiling. This process was performed to identify the fluorescent cell of interest and for

slice-level correlation in later steps of the workflow. The slice containing the target cell was then imaged

with the same 20x objective at higher pixel resolution (0.360 × 0.360 × 0.976 µm/voxel) to obtain a z-

stack of the full depth of the slice and immediate region surrounding the target cell. CLSM imaging

required approximately 1-3 hours to find the cell of interest within a slice and capture a tiled slice overview

and higher resolution z-stack.

Sample preparation for SBF-SEM imaging

Samples were prepared as previously described (Scholl et al., 2021; Thomas et al., 2021). Briefly, tissue

pieces were incubated in an aqueous solution of 2% osmium tetroxide buffered in 0.1 M sodium

cacodylate for 45 minutes at room temperature (RT). Tissue was not rinsed and the osmium solution was

replaced with cacodylate buffered 2.5% potassium ferrocyanide for 45 minutes at RT in the dark. Tissue

was rinsed with water 2 × 10 minutes, which was repeated between consecutive steps. Tissue was

incubated at room temperature for 20 minutes in aqueous 1% thiocarbohydrizide dissolved at 60°C,

aqueous 1% osmium tetroxide for 45 minutes at RT, and then 1% uranyl acetate in 25% ethanol for 20

minutes at RT in the dark. Tissue was rinsed then left in water overnight at 4°C. The following day, tissue

was stained with Walton's lead aspartate for 30 minutes at 60°C. Tissue was then dehydrated in a graded

ethanol series (30, 50, 70, 90, 100%), 1:1 ethanol to acetone, then 100% dry acetone. Tissue was

infiltrated using 3:1 acetone to Durcupan resin (Sigma Aldrich) for 2 hours, 1:1 acetone to resin for 2

hours, and 1:3 acetone to resin overnight, then flat embedded in 100% resin on a glass slide and covered

with an Aclar sheet at 60°C for 2 days. The tissue was trimmed to less than 1 × 1 mm in size, the empty

resin at the surface was shaved to expose tissue surface using an ultramicrotome (UC7, Leica), then

turned downwards to be remounted to a metal pin with conductive silver epoxy (CircuitWorks,

Chemtronics).

SBF-SEM image acquisition and volume data handling

Samples were imaged using a correlative approach as previously described (Thomas et al., 2021).

Briefly, tissue was sectioned and imaged using 3View2XP and Digital Micrograph (ver. 3.30.1909.0,

Gatan Microscopy Suite) installed on a Gemini SEM300 (Carl Zeiss Microscopy LLC.) equipped with an

OnPoint BSE detector (Gatan, Inc.). The detector magnification was calibrated within SmartSEM imaging

software (ver. 6.0, Carl Zeiss Microscopy LLC.) and Digital Micrograph with a 500 nm cross line grating

standard. A low magnification image of each block face was manually matched to its corresponding depth

in the CLSM Z-stack using blood vessels and cell bodies as fiducials. Final imaging was performed at 2.0

- 2.2 kV accelerating voltage, 20 or 30 µm aperture, working distance of ~5 mm, 0.5 - 1.2 µs pixel dwell

time, 5.7 - 7 nm per pixel, knife speed of 0.1 mm/sec with oscillation, and 56 - 84 nm section thickness.

Acquisition was automated and ranged from several days to several weeks depending on the size of the

ROI and imaging conditions. The true section thickness was measured using mitochondria diameter

calibrations (Fiala and Harris, 2001). Calibration for each block was required, since variation in thickness

can occur due to heating, charging from the electron beam, resin polymerization, and tissue fixation and

staining quality (Starborg et al., 2013; Hughes et al., 2014).

Serial tiled images were exported as TIFFs to TrakEM2 (Schindelin et al., 2012) within ImageJ

(ver. 1.52p) to montage tiles, then aligned using Scale-Invariant Feature Transform image alignment with

linear feature correspondences and rigid transformation (Lowe, 2004). Once aligned, images were

inverted and contrast normalized.

SBF-SEM image analysis

Aligned images were exported to Microscopy Image Browser (ver. 2.51, 2.6, Belevich et al.,

2016) for segmentation of dendrites, spines, postsynaptic densities (PSDs), and boutons. Three

annotators preformed segmentation and the segmentations of each annotator were proof-read by an

experienced annotator (~1000 hours of segmentation experience) prior to quantification. Binary labels

files were imported to Amira (ver. 6.7, 2019.1, Thermo Fisher Scientific) which was used to create 3D

surface models of each dendrite, spine, and PSD. Once reconstructed, the model of each dendrite was

manually overlaid onto its corresponding two-photon image using Adobe Photoshop for re-identification of

individual spines. Amira was used to measure the volume of spine heads, surface area of PSDs, and

spine neck length.

For mitochondrial analyses, segmentations were moved into ORS Dragonfly (ver. 2021.3,

2022.1). Within Dragonfly, a pre-built U-net++ architecture was trained on a small set of manual

mitochondria segmentations, then applied for segmentation of all mitochondria within target dendrites.

Dendrite segmentations were used as a mask to apply the trained mitochondria network, and

mitochondria segmentations were proofread for accuracy. Mitochondria morphology was measured using

tools within Dragonfly, while inter-mitochondria distances were measured in Microscopy Image Browser

using the measure tool. For quantification of mitochondria volume near spines, a sphere of either 1 µm or

5 µm radius was applied with the center at the base of each spine neck, and the volume of mitochondria

intersected by each sphere was measured. Mitochondria within spines were segmented manually and

these spines (n = 10) were not included in correlation analyses. In addition, analyses excluded spines

with 0 μm^3 mitochondrial volume within the relevant radii unless otherwise stated.

Statistics

Statistical analyses are described in the main text and in figure legends. We used non-parametric

statistical analyses (Wilcoxon sign-rank test, Mann-Whitney rank-sum test) or permutation tests to avoid

assumptions about the distributions of the data. Permutation tests used equivalent sample sizes for each

distribution and compared data to the null hypothesis. Statistical analysis was performed in MATLAB

(2021b) or GraphPad Prism (ver. 9). Circular correlation coefficients were computed for tests with circular

variables. For all other tests, Spearman's correlation coefficient was computed. All correlation significance

tests were one-sided. Quantitative approaches were not used to determine if the data met the

assumptions of the parametric tests.

References

Attwell D, Laughlin SB (2001) An Energy Budget for Signaling in the Grey Matter of the Brain. J Cereb Blood Flow Metab 21:1133–1145.

Bartol TM, Bromer C, Kinney J, Chirillo MA, Bourne JN, Harris KM, Sejnowski TJ (2015) Nanoconnectomic upper bound on the variability of synaptic plasticity. eLife 4:e10778.

Belevich I, Joensuu M, Kumar D, Vihinen H, Jokitalo E (2016) Microscopy Image Browser: A Platform for Segmentation and Analysis of Multidimensional Datasets. PLOS Biol 14:e1002340.

Belles B, Hescheler J, Trube G (1987) Changes of membrane currents in cardiac cells induced by long whole-cell recordings and tolbutamide. Pflüg Arch - Eur J Physiol 409:582–588.

Cameron HA, Kaliszewski CK, Greer CA (1991) Organization of mitochondria in olfactory bulb granule cell dendritic spines. Synapse 8:107–118.

Chang DTW, Honick AS, Reynolds IJ (2006) Mitochondrial Trafficking to Synapses in Cultured Primary Cortical Neurons. J Neurosci 26:7035–7045.

Cheng A, Hou Y, Mattson MP (2010) Mitochondria and Neuroplasticity. ASN Neuro 2:AN20100019.

Cserép C, Pósfai B, Schwarcz AD, Dénes Á (2018) Mitochondrial Ultrastructure Is Coupled to Synaptic Performance at Axonal Release Sites. eNeuro 5:ENEURO.0390-17.2018.

Datta S, Jaiswal M (2021) Mitochondrial calcium at the synapse. Mitochondrion 59:135–153.

Delgado T, Petralia RS, Freeman DW, Sedlacek M, Wang Y-X, Brenowitz SD, Sheu S-H, Gu JW, Kapogiannis D, Mattson MP, Yao PJ (2019) Comparing 3D ultrastructure of presynaptic and postsynaptic mitochondria. Biol Open 8:bio044834.

Devine MJ, Kittler JT (2018) Mitochondria at the neuronal presynapse in health and disease. Nat Rev Neurosci 19:63–80.

Divakaruni SS, Van Dyke AM, Chandra R, LeGates TA, Contreras M, Dharmasri PA, Higgs HN, Lobo MK, Thompson SM, Blanpied TA (2018) Long-Term Potentiation Requires a Rapid Burst of Dendritic Mitochondrial Fission during Induction. Neuron 100:860-875.e7.

Dromard Y, Arango-Lievano M, Fontanaud P, Tricaud N, Jeanneteau F (2021) Dual imaging of dendritic spines and mitochondria in vivo reveals hotspots of plasticity and metabolic adaptation to stress. Neurobiol Stress 15:100402.

Faitg J, Lacefield C, Davey T, White K, Laws R, Kosmidis S, Reeve AK, Kandel ER, Vincent AE, Picard M (2021) 3D neuronal mitochondrial morphology in axons, dendrites, and somata of the aging mouse hippocampus. Cell Rep 36:109509.

Faits MC, Zhang C, Soto F, Kerschensteiner D (2016) Dendritic mitochondria reach stable positions during circuit development Youle RJ, ed. eLife 5:e11583.

Fiala JC, Harris KM (2001) Cylindrical diameters method for calibrating section thickness in serial electron microscopy. J Microsc 202:468–472.

Glancy B, Kim Y, Katti P, Willingham TB (2020) The Functional Impact of Mitochondrial Structure Across Subcellular Scales. Front Physiol 11:541040.

Goris RLT, Simoncelli EP, Movshon JA (2015) Origin and Function of Tuning Diversity in Macaque Visual Cortex. Neuron 88:819–831.

Harris JJ, Jolivet R, Attwell D (2012) Synaptic Energy Use and Supply. Neuron 75:762–777.

Holler S, Köstinger G, Martin KAC, Schuhknecht GFP, Stratford KJ (2021) Structure and function of a neocortical synapse. Nature 591:111–116.

Holtmaat A, Svoboda K (2009) Experience-dependent structural synaptic plasticity in the mammalian brain. Nat Rev Neurosci 10:647–658.

Hubley MJ, Locke BR, Moerland TS (1996) The effects of temperature, pH, and magnesium on the diffusion coefficient of ATP in solutions of physiological ionic strength. Biochim Biophys Acta BBA - Gen Subj 1291:115–121.

Hughes L, Hawes C, Monteith S, Vaughan S (2014) Serial block face scanning electron

microscopy—the future of cell ultrastructure imaging. Protoplasma 251:395–401.

Kasthuri N et al. (2015) Saturated Reconstruction of a Volume of Neocortex. Cell 162:648–661.

Koester, Helmut J., and Daniel Johnston. "Target Cell-Dependent Normalization of Transmitter Release at Neocortical Synapses." *Science* 308, no. 5723 (May 6, 2005): 863–66. https://doi.org/10.1126/science.1100815.

Leung A, Ohadi D, Pekkurnaz G, Rangamani P (2021) Systems modeling predicts that mitochondria ER contact sites regulate the postsynaptic energy landscape. Npj Syst Biol Appl 7:1–14.

Lewis TL, Kwon S-K, Lee A, Shaw R, Polleux F (2018) MFF-dependent mitochondrial fission regulates presynaptic release and axon branching by limiting axonal mitochondria size. Nat Commun 9:5008.

Li Z, Okamoto K-I, Hayashi Y, Sheng M (2004) The Importance of Dendritic Mitochondria in the Morphogenesis and Plasticity of Spines and Synapses. Cell 119:873–887.

Lipton P, Whittingham TS (1982) Reduced ATP concentration as a basis for synaptic transmission failure during hypoxia in the in vitro guinea-pig hippocampus. J Physiol 325:51–65.

Lowe DG (2004) Distinctive Image Features from Scale-Invariant Keypoints. Int J Comput Vis 60:91–110.

MacAskill AF, Kittler JT (2010) Control of mitochondrial transport and localization in neurons. Trends Cell Biol 20:102–112.

Mante V, Bonin V, Carandini M (2008) Functional Mechanisms Shaping Lateral Geniculate Responses to Artificial and Natural Stimuli. Neuron 58:625–638.

Palmer LM, Stuart GJ (2009) Membrane potential changes in dendritic spines during action potentials and synaptic input. J Neurosci 29:6897–6903.

Pathak D, Shields LY, Mendelsohn BA, Haddad D, Lin W, Gerencser AA, Kim H, Brand MD, Edwards RH, Nakamura K (2015) The Role of Mitochondrially Derived ATP in Synaptic Vesicle Recycling. J Biol Chem 290:22325–22336.

Peirce JW (2007) PsychoPy—Psychophysics software in Python. J Neurosci Methods 162:8–13.

Pekkurnaz G, Wang X (2022) Mitochondrial heterogeneity and homeostasis through the lens of a neuron. Nat Metab 4:802–812.

Pnevmatikakis EA, Giovannucci A (2017) NoRMCorre: An online algorithm for piecewise rigid motion correction of calcium imaging data. J Neurosci Methods 291:83–94.

Popov V, Medvedev NI, Davies HA, Stewart MG (2005) Mitochondria form a filamentous reticular network in hippocampal dendrites but are present as discrete bodies in axons: A three-dimensional ultrastructural study. J Comp Neurol 492:50–65.

Popovic MA, Carnevale N, Rozsa B, Zecevic D (2015) Electrical behaviour of dendritic spines as revealed by voltage imaging. Nat Commun 6:8436.

Rangaraju V, Calloway N, Ryan TA (2014) Activity-Driven Local ATP Synthesis Is Required for Synaptic Function. Cell 156:825–835.

Rangaraju V, Lauterbach M, Schuman EM (2019) Spatially Stable Mitochondrial Compartments Fuel Local Translation during Plasticity. Cell 176:73-84.e15.

Rodriguez-Moreno J, Rollenhagen A, Arlandis J, Santuy A, Merchan-Pérez A, DeFelipe J, Lübke JHR, Clasca F (2018) Quantitative 3D Ultrastructure of Thalamocortical Synapses from the "Lemniscal" Ventral Posteromedial Nucleus in Mouse Barrel Cortex. Cereb Cortex 28:3159–3175.

Rossi MJ, Pekkurnaz G (2019) Powerhouse of the mind: mitochondrial plasticity at the synapse. Curr Opin Neurobiol 57:149–155.

Schikorski T, Stevens CF (1997) Quantitative Ultrastructural Analysis of Hippocampal Excitatory Synapses. J Neurosci 17:5858–5867.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,

Rueden C, Saalfeld S, Schmid B (2012) Fiji: an open-source platform for biological-
image analysis. Nat Methods 9:676–682.
Sobczyk, Aleksander, Volker Scheuss, and Karel Svoboda. "NMDA Receptor Subunit-
Dependent [Ca ^{2} +] Signaling in Individual Hippocampal Dendritic Spines." *The Journal*
of Neuroscience 25, no. 26 (June 29, 2005): 6037–46.
https://doi.org/10.1523/JNEUROSCI.1221-05.2005.
Scholl B, Fitzpatrick D (2020) Cortical synaptic architecture supports flexible sensory
computations. Curr Opin Neurobiol 64:41–45.
Scholl B, Thomas CI, Ryan MA, Kamasawa N, Fitzpatrick D (2021) Cortical response selectivity
derives from strength in numbers of synapses. Nature 590:111–114.
Scholl B, Wilson DE, Fitzpatrick D (2017) Local Order within Global Disorder: Synaptic
Architecture of Visual Space. Neuron 96:1127-1138.e4.
Smith HL, Bourne JN, Cao G, Chirillo MA, Ostroff LE, Watson DJ, Harris KM (2016)
Mitochondrial support of persistent presynaptic vesicle mobilization with age-dependent
synaptic growth after LTP Häusser M, ed. eLife 5:e15275.
Starborg T, Kalson NS, Lu Y, Mironov A, Cootes TF, Holmes DF, Kadler KE (2013) Using
transmission electron microscopy and 3View® to determine collagen fibril size and
three-dimensional organization. Nat Protoc 8:1433–1448.
Thomas CI, Keine C, Okayama S, Satterfield R, Musgrove M, Guerrero-Given D, Kamasawa N,
Young SM (2019) Presynaptic Mitochondria Volume and Abundance Increase during
Development of a High-Fidelity Synapse. J Neurosci 39:7994–8012.
Thomas CI, Ryan MA, Scholl B, Guerrero-Given D, Fitzpatrick D, Kamasawa N (2021)
Targeting Functionally Characterized Synaptic Architecture Using Inherent Fiducials and
3D Correlative Microscopy. Microsc Microanal 27:156–169.
Turner NL et al. (2022) Reconstruction of neocortex: Organelles, compartments, cells, circuits,
and activity. Cell 185:1082-1100.e24.
Virga DM, Hamilton S, Osei B, Morgan A, Zamponi E, Park NJ, Hewitt VL, Zhang D, Gonzalez
KC, Bloss E, Polleux F, Lewis TL (2023) Activity-dependent subcellular
compartmentalization of dendritic mitochondria structure in CA1 pyramidal neurons.
:2023.03.25.534233 Available at:
https://www.biorxiv.org/content/10.1101/2023.03.25.534233v1 [Accessed April 15,
2023].
Wilke SA, Antonios JK, Bushong EA, Badkoobehi A, Malek E, Hwang M, Terada M, Ellisman
MH, Ghosh A (2013) Deconstructing Complexity: Serial Block-Face Electron Microscopic
Analysis of the Hippocampal Mossy Fiber Synapse. J Neurosci 33:507–522.
Wilson DE, Whitney DE, Scholl B, Fitzpatrick D (2016) Orientation selectivity and the functional
clustering of synaptic inputs in primary visual cortex. Nat Neurosci 19:1003–1009.