Neuroendocrine Coupling of Interoceptive Bacteria-Derived Cues to Foraging Behavior in

C. elegans

Sonia A. Boor[1,2], Joshua D. Meisel[2,3], Dennis H. Kim[1,*]

1. Division of Infectious Diseases, Department of Pediatrics, Boston Children's Hospital

and Harvard Medical School, Boston, MA 02115

2. Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139

3. Department of Molecular Biology, Massachusetts General Hospital, Boston, MA 02114

* Correspondence to dennis.kim@childrens.harvard.edu

The authors declare no competing interests.

Author contributions:

SAB: Conceptualization, Formal analysis, Validation, Investigation, Visualization,

Methodology, Writing—original draft, Writing—review and editing

JDM: Investigation, Visualization, Writing – review and editing

DHK: Conceptualization, Supervision, Funding acquisition, Visualization, Methodology,

Writing—original draft, Writing—review and editing

Summary

Animal internal state is modulated by nutrient intake, resulting in behavioral responses to changing food conditions. DAF-7 is a neuroendocrine TGF-beta ligand that regulates diverse food-related behaviors of *Caenorhabditis elegans,* including foraging behavior. Here, we show that in *C. elegans,* interoceptive cues from the ingestion of bacterial food inhibit the expression of DAF-7, a neuroendocrine TGF-beta ligand, from the ASJ pair of sensory neurons, whereas food deprivation in the presence of external chemosensory cues from bacteria promotes the expression of DAF-7 from the ASJ neurons. We show that SCD-2, the *C. elegans* ortholog of mammalian Anaplastic Lymphoma Kinase (ALK), which has been implicated in the central control of metabolism of mammals, functions in the AIA interneurons to regulate foraging behavior and cell-non-autonomously control the expression of DAF-7 from the ASJ neurons. Our data establish an SCD-2-dependent neuroendocrine DAF-7 gene expression feedback loop that couples the ingestion of bacterial food to foraging behavior.

Introduction

Internal states, such as fear, arousal, and hunger, are shaped by the integration of information about internal and external conditions and can result in the modulation of various physiological and behavioral outputs (Flavell et al., 2022). As animals encounter different food environments, information about internal nutritional status and external food quality can elicit transitions between internal states that either favor the exploration of new places or exploitation of the current environment. Many animals, including mammals, zebrafish, *Drosophila melanogaster*, and the roundworm *Caenorhabditis elegans,* increase exploration in response to food deprivation in order to increase chances of a food encounter (Ben Arous et al., 2009;

Connolly, 1966; Gutman et al., 2007; Herbers, 1981; Johnson et al., 2020; Overton & Williams,

2004; Russell et al., 1987). However, the mechanisms behind how animals couple changes in

food availability to internal states that influence foraging behavior are not well understood.

Insight into the cellular and organismal mechanisms governing internal states may enhance

understanding of the dysregulation of internal states that is thought to contribute to many human

psychiatric and neurological diseases (Flavell et al., 2022; Yap & Greenberg, 2018).

C. elegans forage for microbes that grow on decaying organic matter, where they

encounter fluctuations in not only the quantity of nutritious bacterial food available but also in

the quality and pathogenicity of this bacterial food (Kim & Flavell, 2020). As they navigate their

food environments, *C. elegans* exhibit two-state foraging and feeding behavior known as

roaming and dwelling (Ben Arous et al., 2009; Flavell et al., 2013; Flavell et al., 2020; Fujiwara

et al., 2002). On abundant nutritious food, animals spend about 80% of their time dwelling and

20% of their time roaming; as food becomes scarcer or lower in quality, animals increase the

proportion of their time roaming (Ben Arous et al., 2009). The internal states that underlie *C.*

elegans roaming and dwelling responses to changing food conditions present an experimentally

tractable paradigm in which to study how internal states are regulated. Furthermore, the shared

modulators of internal states and high degree of genetic conservation between *C. elegans* and

humans suggest that understanding *C. elegans* feeding and foraging behavior could have relevant

implications for human health and disease.

Aided by a complete connectome (White et al., 1986), a growing body of work has

examined neural circuits and their effects on foraging behavior in *C. elegans* (Ji et al., 2021;

Pradhan et al., 2019). The excitation or inhibition of neurons in these circuits elicits rapid

behavioral modulation as well as persistent behavioral states through recurrent neuronal firing.

Neuronal dynamics are additionally influenced by neuromodulators (Bargmann, 2012). For

instance, the activation of a small group of neurons can induce transitions between roaming and

dwelling states through conserved serotonin and pigment dispersing factor signaling (Flavell et

al., 2013; Ji et al., 2021). Other neuromodulators such as dopamine and octopamine can further

influence foraging behavior and food-dependent locomotion (Churgin et al., 2017; Oranth et al.,

2018; Sawin et al., 2000). Neuromodulators play a key role in mediating internal states by

functioning at titratable levels over longer timescales and farther distances than direct synaptic

signaling (Flavell et al., 2022; Sengupta, 2013). Activity-dependent gene expression in neurons

has been established to have key roles in the development and plasticity of neuronal circuits, but

less is known about how changes in neuronal gene expression may shape internal states driving

behavior (Yap & Greenberg, 2018). As transcription occurs on a slower timescale than neuronal

firing or neuromodulator release, dynamic gene expression could be important in regulating the

persistence of internal states.

We have been studying the dynamic temporal and neuron-specific expression of *daf-7*,

which encodes a TGF-beta ligand that is involved in the neuroendocrine regulation of a diverse

range of behaviors in *C. elegans*, including the dauer developmental decision, longevity,

metabolism, and feeding and foraging behavior (Ben Arous et al., 2009; Greer et al., 2008; Ren

et al., 1996; Schackwitz et al., 1996; Shaw et al., 2007). Expression of *daf-7* is restricted to a

limited set of sensory neurons, including the ASI neurons (Meisel et al., 2014; Ren et al., 1996;

Schackwitz et al., 1996). DAF-7 expression from the ASI neurons was shown to respond to

changing environmental conditions, such as crowding and food levels (Entchev et al., 2015; Ren

et al., 1996; Schackwitz et al., 1996). Previously, we observed that highly dynamic *daf-7*

expression can be observed in the ASJ neurons, with induction of expression in response to

secondary metabolites produced by pathogenic *Pseudomonas aeruginosa* PA14, which is

necessary for pathogen avoidance behavior (Meisel et al., 2014). In addition, we have shown that

upon the onset of reproductive maturity, male *C. elegans* upregulate *daf-7* expression in their

ASJ neurons to promote male mate-searching behavior (Hilbert & Kim, 2017).

Here, we have identified how the ingestion of bacterial food regulates the expression of a

single gene in two neurons to shape internal state dynamics driving foraging behavior in *C.*

elegans. We observed that *daf-7* transcription levels in the ASJ neurons couples foraging

behavior to changes in bacterial food ingestion, under the control of the highly conserved

receptor tyrosine kinase SCD-2/ALK. The relationship between gene transcription and

behavioral states across organisms is largely uncharacterized (Yap & Greenberg, 2018), and our

results establish a role for SCD-2/ALK-regulated dynamic neuronal gene expression in shaping

internal states underlying foraging behavior in response to changing nutritional conditions.

Results

Ingested food inhibits *daf-7* expression in the ASJ neurons

Our lab has previously reported that when adult hermaphrodites are fed the normal *E. coli*

OP50 food source, *pdaf-7::GFP* expression is restricted to the ASI chemosensory neurons

(Hilbert & Kim, 2017; Meisel et al., 2014). In contrast, we observed that when animals were fed

OP50 treated with aztreonam, an antibiotic that causes the bacteria to form inedible long strands

(Gruninger et al., 2008), *pdaf-7::GFP* was expressed in both the ASI and ASJ neurons (Figures

1A, 1C, 1D). These data suggested that ingestion of food inhibits *daf-7* expression in the ASJ

neurons. However, we did not observe *daf-7* expression in the ASJ neurons in the complete

absence of food ("Empty"), which indicated that some bacterial component of non-ingestible

aztreonam-treated *E. coli* OP50 was required for the induction of *daf-7* in the ASJ neurons in the

absence of ingested food (Figures 1A, 1C, 1D). The non-ingestible food signal that results in the

upregulation of *daf-7* expression in the ASJ neurons on aztreonam-treated food could be

diffusible, volatile, or mechanosensory. To identify the nature of the non-ingestible food cue, we

exposed animals to food that had been seeded on the lid of the plate (exposing animals to only

the volatile food cues) ("Lid") or underneath the agar (exposing animals only to diffusible food

cues) ("Bottom") in the absence of ingestible food and probed *daf-7* expression. We observed

induction of *daf-7* in the ASJ neurons when OP50 is seeded under the agar of the plates, but not

when present on the lid of the plate, suggesting that a bacteria-derived food signal that diffuses

through the agar is necessary for the induction of *daf-7* expression in ASJ in the absence of

ingested food. (Figures 1A-F). Thus, these data suggest that in hermaphrodites feeding on *E. coli*

OP50, there is an external water-soluble diffusible cue from the bacteria that stimulates *daf-7*

expression from the ASJ neurons, but this expression is inhibited by a second bacteria-derived,

interoceptive cue generated from the ingestion of bacteria (Figure 1B).

We next asked whether the inhibition of *daf-7* expression in the ASJ neurons by ingestion

of bacterial food was mediated by the early detection of ingested food or the delayed nutritional

effects of bacterial food consumption on organismal physiology. The kinetics of *pdaf-7::GFP*

induction in the ASJ neurons when animals were moved from ingestible to non-ingestible food

showed a rapid upregulation within 2-3 h for the GFP fluorescence in the ASJ neurons to become

visible (Figure 1G). Given the delay for GFP folding and accumulation to visible levels, these

data suggested that the absence of ingested food rapidly led to the induction of *daf-7* expression

in the ASJ neurons. Recent work has described the interoceptive sensing of ingested food by the

acid-sensing ion channels (ASICs) encoded by *del-3* and *del-7*, which are expressed in the minor

neurites of the NSM neurons where they detect food in the pharynx and mediate behavioral

slowing upon encountering food (Rhoades et al., 2019). We observed that *del-3;del-7* animals

expressed *daf-7* in the ASJ neurons even on ingestible food (Figure 1H). These observations are

consistent with the hypothesis that the detection of an interoceptive ingested food signal in the

pharynx, mediated by DEL-3 and DEL-7, inhibits *daf-7* expression in the ASJ neurons.

***daf-7* expression in the ASJ neurons promotes roaming behavior**

Next, we sought to understand how changes in *daf-7* expression in the ASJ neurons in

response to changing food conditions might modulate food-dependent behavior. Feeding *C.*

elegans alternate between two distinct behavioral states known as roaming and dwelling (Ben

Arous et al., 2009; Fujiwara et al., 2002). On nutritious food, wild-type animals spend most of

their time dwelling, a feeding state marked by a low movement speed and high body bending

angle. As food conditions worsen, animals will decrease the fraction of time dwelling and

increase the fraction of time roaming, a foraging state marked by high speed and low curvature

(Ben Arous et al., 2009). Prior genetic analysis of roaming and dwelling has revealed a role for

DAF-7 in promoting roaming behavior, with *daf-7* animals spending a decreased fraction of time

roaming compared to wild-type animals (Ben Arous et al., 2009). Using both worm-tracking

software to quantify roaming and dwelling (*WormLab*, 2020) and an exploration assay to

measure the general activity levels of animals ((Flavell et al., 2013), see Materials and Methods),

we observed that *daf-7(e1372)* and *daf-7(ok3125)* animals spent a markedly decreased

proportion of time in the roaming state than wild-type animals, consistent with these prior

findings (Figure 2A). To determine whether we could detect modulation of roaming and

dwelling behaviors caused by changes in *daf-7* expression in the ASJ neurons, we adopted two

complementary approaches. First, we found that in a *daf-7(ok3125)* background, rescue of *daf-7*

cDNA under the ASJ-specific *trx-1* promoter partially restored the fraction of time animals spent

in the roaming state (Figure 2B). Second, we examined exploratory behavior in animals that had

daf-7 deleted in the ASJ neurons, using a floxed allele of *daf-7* and Cre expressed under the *trx-1*

promoter. We found that animals with an ASJ-specific *daf-7* deletion explored less of the lawn

than animals with wild-type expression of *daf-7* (Figure 2C).

The relationships between ingested food and *daf-7* expression in the ASJ neurons, *daf-7*

expression in the ASJ neurons and roaming, and roaming and amount of food ingested suggest a

neuroendocrine gene expression positive-feedback loop that couples the ingestion of bacterial

food to the modulation of foraging behavior through *daf-7* expression in the ASJ neurons (Figure

2D). When animals are on abundant edible food, the ingestion of this food inhibits *daf-7*

expression in the ASJ neurons and attenuates the proportion of time in the roaming state. When

animals are removed from edible food, upregulation of *daf-7* expression in the ASJ neurons

promotes an increased proportion of time in the roaming state. We hypothesized that this

transcriptional feedback loop might function in the persistence of internal states underlying

roaming and dwelling behaviors, so we measured the duration of roaming and dwelling states in

animals lacking DAF-7 signaling.

Representative trace files of wild-type, *daf-7(e1372),* and *daf-7(ok3125)* animals show

how DAF-7 can influence roaming and dwelling state duration (Figure 2E). Quantification of

state duration across multiple animals revealed that *daf-7* animals exhibited shortened roaming

states relative to wild-type animals as predicted by the positive feedback loop (Figure 2F). We

did not observe a difference in dwelling state duration among the genotypes examined, although

we suspect the wide variation dwelling state durations exhibited by animals precluded the

detection of a statistically significant difference with our analysis.

Known modulators of foraging behavior also regulate *daf-7* expression

The positive-feedback loop supported by our data (Figure 2D) implies that *daf-7*

expression in the ASJ neurons could be increased in genetic backgrounds favoring roaming

behavior and hence a diminished ingestion of bacterial food. Serotonin has been shown to

promote long dwelling states, and thus animals with mutations in *tph-1,* the gene encoding the

enzyme that catalyzes the rate-limiting step of serotonin biosynthesis, spend a greater proportion

of their time in the roaming state (Figure 3A) (Flavell et al., 2013; Ji et al., 2021). Consistent

with the correlation between *daf-7* expression in the ASJ neurons and roaming, we see that *tph-

1(n4622)* animals constitutively express *daf-7* in the ASJ neurons on both ingestible and non-

ingestible food, with this *daf-7* expression retaining sensitivity to changes in food environments

(Figure 3B).

Pigment dispersing factor (PDF) signaling has also been previously shown to modulate

roaming and dwelling behavior, with animals carrying loss-of-function mutations in the PDF

receptor gene *pdfr-1* showing a decreased fraction of their time in the roaming state than wild-

type animals (Figure 3C) (Flavell et al., 2013; Ji et al., 2021). Consistent with the correlation

between roaming and *daf-7* expression in ASJ, we observed that animals with loss-of-function

mutations in *pdfr-1* do not upregulate *daf-7* transcription in the ASJ neurons on non-ingestible

food (Figure 3D). From a screen done in parallel to this work, we recovered a putative gain-of-

function allele of *pdfr-1*, *syb3826,* carrying a S325F substitution (Boor, 2022). We observed that

pdfr-1(syb3826) animals exhibited increased roaming behavior compared with wild-type animals

(Figure 3C) and constitutive expression of *daf-7* in the ASJ neurons when fed ingestible food,

which is further upregulated on non-ingestible food (Figure 3D).

SCD-2 controls *daf-7* expression in the ASJ neurons and roaming

207   To identify additional genetic factors involved in coupling the ingestion of food to *daf-7*

expression in the ASJ neurons and its subsequent effect on roaming, we were guided by the

isolation of an allele of *hen-1, qd259,* in a screen for genes regulating the expression of *daf-7*

from the ASJ neurons in response to *P. aeruginosa*. Through genomic rescue experiments, we

confirmed that the *hen-1* mutation was the causative lesion in this strain, and animals with

mutations in *scd-2,* encoding the receptor of HEN-1, share this phenotype (Figure 4A). SCD-2

has characterized for its role in the food-dependent developmental arrest known as dauer

diapause (Reiner et al., 2008), and Anaplastic Lymphoma Kinase (ALK), the human ortholog of

SCD-2, has been implicated in metabolic phenotypes in humans, mice, and *Drosophila* (Orthofer

et al., 2020). Taken together, these observations led us to question if SCD-2 might be involved in

the regulation of *daf-7* expression in the ASJ neurons in response to changes in food ingestion.

We found that animals with loss-of-function alleles of either *scd-2* or *hen-1* showed reduced

upregulation of *daf-7* in the ASJ neurons in response to non-ingestible food compared to wild-

type animals, implicating SCD-2 in this *daf-7* expression response (Figures 4B and 4D).

221   To further examine the role of SCD-2 mediating the *daf-7* expression response to

ingested food, we generated a gain-of-function allele. Gain-of-function point mutations in ALK

are associated with neuroblastoma and most often occur in the kinase domain, with 85% of all

ALK point mutations seen at either F1174 or R1275 (Franco et al., 2013). F1174L, the most

common point mutation, results in ALK autophosphorylation and cytokine-independent growth

(Chen et al., 2008; George et al., 2008; Hallberg & Palmer, 2013; Holla et al., 2017; Janoueix-

Lerosey et al., 2008). Analysis of a sequence alignment of SCD-2 and ALK revealed that F1174

in ALK was conserved as F1029 in SCD-2, and we were able to create a gain-of-function allele

of *scd-2* by engineering a F1029L substitution (*scd-2(syb2455)*) (Figure 4C) (Boor, 2022). To

infer whether SCD-2 activity was sufficient to induce expression of *daf*-7 in the ASJ neurons in

the absence of ingestible bacterial food, we examined the gain-of-function *scd-2(syb2455)*

mutant and observed that these animals constitutively expressed *daf-7* in the ASJ neurons even in

the presence of ingestible food (Figure 4D). We further observed that the magnitude of

upregulation of *daf-7* expression in the ASJ neurons when animals were moved from ingestible

food to non-ingestible food was reduced in *scd-2(syb2455)* to levels only about one-fourth of

those seen in wild-type animals (Figure 4D). In contrast, these *scd-2(syb2455)* gain-of-function

animals still show robust upregulation of *daf-7* expression in the ASJ neurons when exposed to

P. aeruginosa, supporting a direct role for SCD-2 in the response to ingested food rather than a

global control of *daf-7* expression in the ASJ neurons (Figure 4A).

Expression of *scd-2* cDNA under the AIA-specific *gcy-28.d* promoter has been shown to

be sufficient to rescue behavioral phenotypes of *scd-2* mutants that have been attributed to a role

for AIA-expressed SCD-2 in sensory integration (Shinkai et al., 2011). To ask whether SCD-2

could be functioning in the AIA neurons to regulate *daf-7* expression in the ASJ neurons, we

generated a strain with the coding sequence of SCD-2 floxed and introduced a transgene

expressing Cre under the *gcy-28.d* promoter. Compared to animals containing a floxed *scd-2*

allele without Cre, animals with AIA-specific Cre expression resulting in AIA-specific deletion

of *scd-2* showed reduced upregulation of *daf-7* expression in the ASJ neurons in response to non-

ingestible food, suggesting that SCD-2 activity in the AIA neurons acts cell-non-autonomously

to control *daf-7* expression in the ASJ neurons (Figure 4E).

Our finding that SCD-2 regulates the *daf-7* expression in response to ingested food

predicts that SCD-2 would also promote roaming behavior. Consistent with this hypothesis, we

observed that animals with loss-of-function mutations in *scd-2* spent a lower proportion of time

roaming and explored less of the lawn in an exploration assay than wild-type animals (Figure

4F). Animals with a loss-of-function allele of *hen-1* also roamed less than wild type (Figure 4F).

Furthermore, as predicted by their regulation of *daf-7* expression in the ASJ neurons, *scd-2* and

hen-1 animals exhibit shorter roaming states and unchanged dwelling state durations compared

to wild-type animals (Figure 4G). When we examined the roaming and dwelling behavior of *scd-*

2(syb2455) gain-of-function animals, we found that these mutants showed increased roaming

behavior and lawn exploration as compared to wild-type animals and exhibited longer roaming

states (Figure 4H and 4I). As with the regulation of *daf-7* expression in the ASJ neurons, we

observed that Cre-lox-mediated deletion of *scd-2* specifically in the AIA neurons reduced both

roaming and exploration compared to floxed *scd-2* animals without Cre, suggesting that SCD-2

functions in the AIA interneurons to promote roaming behavior (Figure 4J).

A neuronal gene expression corelate of internal state dynamics driving foraging behavior

Considering our positive feedback model, we reasoned that conditions under which we

have observed *daf-7* expression in the ASJ neurons would correlate with increased roaming

behavior. Previous work from our lab has shown that upon reaching reproductive maturity, male

C. elegans induce *daf-7* expression in their ASJ neurons, which contributes to male mate-

searching behavior (Figure 5A) (Hilbert & Kim, 2017). Consistent with the behaviors that have

been reported for males and hermaphrodites in bacterial food-leaving assays, we observed that

wild-type males spend a greater proportion of their time in the roaming state than hermaphrodites

(Figure 5B). We also observed that the *daf-7* expression in the ASJ neurons in males was

sensitive to the inhibitory effects of ingested food, although the magnitude of the fold increase in

daf-7 expression from the ASJ neurons was about half that observed for hermaphrodites (Figure

5A).

We have also previously reported an upregulation in *daf-7* expression in the ASJ neurons

when animals are on pathogenic *P. aeruginosa* PA14 in response to specific secondary

metabolites phenazine-1-carboxamide and pyochelin (Figure 5C) (Meisel et al., 2014). Analysis

of roaming and dwelling on PA14 revealed that animals significantly increase the fraction of

their time in the roaming state on PA14 compared to nonpathogenic *E. coli* OP50 (Figure 5D).

We observed that upon exposure to the nonpathogenic PA14 mutant ΔgacA, which results in an

intermediate level of *daf-7* expression in the ASJ neurons (Figure 5C) (Meisel et al., 2014),

animals roamed an intermediate amount between animals on OP50 and pathogenic PA14 (Figure

5D).

Discussion

A neuroendocrine gene expression feedback loop couples the ingestion of bacterial food to

foraging behavior

The data we present here support a model in which bacterial food ingestion regulates

roaming and dwelling behavior in part via SCD-2-dependent *daf-7* expression in the ASJ

neurons. Our identification of the AIA neurons as the site of SCD-2 action is consistent with a

role for the AIA neurons in relaying food signals in other contexts. AIA has been characterized

as a downstream synaptic target of multiple food sensory neurons to integrate information about

the food environment and influence chemoreceptor expression in other neurons (Dobosiewicz et

al., 2019; McLachlan et al., 2022).

We have observed that the expression of *daf-7* in the ASJ neurons responds to two

opposing food cues. An interoceptive cue derived from the sensation of ingested food in the

pharynx inhibits the expression of *daf-7* in the ASJ neurons, while the presence of a diffusible

non-ingested food cue induces the expression of *daf-7* in the ASJ neurons (Figure 1B). In the

presence of both food cues, as on abundant ingestible food, the interoceptive food signal is

epistatic to the external food signal, promoting exploitation of the current food environment with

low levels of *daf-7* expression in the ASJ neurons. Our identification of two opposing food cues

is consistent with findings from Wexler et al., who reported that aztreonam-treated bacteria could

restore *daf-7* expression in the ASJ neurons of starved males, indicative of an external food cue

that is upregulating *daf-7* expression in the ASJ neurons (Wexler et al., 2020).

As discussed above, the relationship between foraging behavior, food ingestion, and

SCD-2-dependent *daf-7* expression has led us to posit a positive-feedback loop that couples food

ingestion with foraging behavior (Figure 2D). As animals in a patchy food environment

encounter a region with less edible food, our model predicts that a reduction of ingested food

would result in *daf-7* expression induction in the ASJ neurons via SCD-2. This increase in *daf-7*

expression is correlated with an internal state that favors roaming. As animals roam, they ingest

less bacteria, further promoting SCD-2-dependent *daf-7* expression in the ASJ neurons,

stabilizing a roaming-prone state. Once a roaming animal encounters edible food and begins to

eat, the interoceptive sensation of food in the pharynx results in a downregulation of *daf-7*

expression in the ASJ neurons, promoting a dwelling state in which the animal can continue to

eat. As additional food is ingested, this further reduces *daf-7* expression in the ASJ neurons,

stabilizing this internal state that favors dwelling.

A number of recent studies in *C. elegans* point to a key role for neuronal transcriptional

responses in the regulation of behavioral plasticity in response to changing environmental stimuli

such as food and temperature (Harris et al., 2023; Kyani-Rogers et al., 2022; Ryan et al., 2014;

Wexler et al., 2020). For example, differential expression of the diacetyl-sensing chemoreceptor

ODR-10 in the AWA neurons in response to changing food conditions, which functions

downstream of *daf-7* expression in the ASJ neurons of males, has been shown to modulate food-

leaving mate-searching behavior (Ryan et al., 2014; Wexler et al., 2020). We anticipate that the

characterization of neuron-specific transcriptional changes that are correlated with *daf-7*

expression in the ASJ neurons may yield additional mechanistic insights into how changing

environmental and endogenous ingested bacterial food levels modulate internal states driving

foraging behavior.

SCD-2/ALK controls a neuroendocrine response to the ingestion of bacterial food

Here, we show that *daf-7* expression in the ASJ neurons is regulated by the receptor

tyrosine kinase SCD-2. SCD-2 and its ligand, HEN-1, have been implicated sensory integration,

associative learning, and memory (Ishihara et al., 2002; Shinkai et al., 2011; Wolfe et al., 2019),

and SCD-2 was initially characterized for its role in regulating dauer formation, a

developmentally arrested state induced by stressful growth conditions, including lack of

nutritious food (Golden & Riddle, 1982, 1984a, 1984b, 1984c; Inoue & Thomas, 2000; Reiner et

al., 2008). SCD-2 is the *C. elegans* ortholog of Anaplastic Lymphoma Kinase (ALK). In humans,

ALK is expressed primarily in the nervous system and influences cell proliferation,

differentiation, and survival in response to external stimuli (Iwahara et al., 1997). Genetic

translocations resulting in gene fusions are common in *ALK*-dependent human cancers, including

non-small cell lung cancer (NSSLC), diffuse large B cell lymphoma (DLBCL), squamous cell

carcinoma (SCC), and renal cell carcinoma (RCC) (Hallberg & Palmer, 2013; Holla et al., 2017;

Morris et al., 1994). Recent studies have suggested a physiological role for ALK in the neuronal

control of metabolism. A genome-wide association study (GWAS) looking for genetic variants

associated with thinness identified a variant in the first intron of *ALK* (Orthofer et al., 2020).

Additional studies in *Drosophila* and mice have further implicated ALK in triglyceride

accumulation, starvation survival, and metabolic response to high-fat diets (Cheng et al., 2011;

Orthofer et al., 2020; Woodling et al., 2020).

We have observed a role for SCD-2 in regulating foraging behavior and neuroendocrine

gene expression in response to changing food conditions. We and others have observed that

animals with loss-of-function mutations in *scd-2* behave as if they are in the constant presence of

ingested "good" food; *scd-2* animals dwell more, reduce upregulation of *daf-7* expression in the

ASJ neurons in the absence of ingested food, and fail to enter dauer under poor food conditions

(Reiner et al., 2008). In contrast, animals with the gain-of-function *scd-2(syb2455)* allele act like

animals exposed to "bad" food conditions, as evidenced by increased roaming, constitutive

expression of *daf-7* in their ASJ neurons, and food-independent constitutive dauer entry (Boor,

2022).

The role for SCD-2 in regulating the physiological response to ingested food in *C.*

elegans is consistent with the growing body of work tying ALK and its orthologs to thinness and

metabolic phenotypes in humans, mice, and *Drosophila* (Cheng et al., 2011; Orthofer et al.,

2020; Woodling et al., 2020). Our observations of *scd-2 C. elegans* are consistent with

phenotypes seen in *ALK^{-/-}* mice and *Alk* RNAi *Drosophila*, including increased energy

expenditure and reduced triglyceride accumulation (Orthofer et al., 2020). Since the phenotypes

of *scd-2* animals are associated with abundant food in *C. elegans,* we might expect these *ALK^{-/-}*

mice and *Alk* RNAi *Drosophila* to mimic animals that are constantly eating and engage in

compensatory mechanisms of increased energy expenditure and triglyceride metabolism. Further

characterization of the mechanisms by with SCD-2 and ALK regulate food-dependent behavior,

gene expression, and metabolism could reveal new factors governing body weight and have

implications in the treatment and prevention of obesity.

daf-7 expression in the ASJ neurons is correlated with internal state favoring exploration

The response of *daf-7* expression in the ASJ neurons to changes in food conditions and

its effect on foraging behavior provides an example of the influence of gene expression on

internal states. We have identified a transcriptional switch of one gene in two cells that can

influence foraging behavior in a manner as equally simple as but on a longer time scale than

neuronal firing (Figure 5E). This *daf-7*-expressing, pro-roaming internal state might drive other

exploratory behaviors that our lab and others have observed under conditions were *daf-7* is

expressed in the ASJ neurons. Male *C. elegans* have been observed to upregulate the expression

of *daf-7* in their ASJ neurons upon reaching reproductive maturity, and this *daf-7* expression has

been implicated in driving mate-searching behavior (Hilbert & Kim, 2017). Males must find a

hermaphrodite mate to reproduce – a requirement not shared by hermaphrodites – and they often

must leave food to do so. The constitutive expression of *daf-7* in the ASJ neurons of adult males

could be due to a reduced sensitivity to ingested food allowing for prioritization of searching for

a mate over feeding. This is consistent with our observation that *daf-7* expression in the ASJ

neurons is upregulated less in response to non-ingestible food in males than in hermaphrodites.

Similarly, our lab has previously reported that two secondary metabolites of pathogenic

P. aeruginosa can result in the upregulation of *daf-7* expression in the ASJ neurons, which

contributes to pathogen avoidance behavior (Meisel et al., 2014). The results presented here

suggest that perhaps these secondary metabolites are tapping into this foraging circuit and

overriding other inputs that regulate *daf-7* expression in the ASJ neurons to promote the animals

moving away from the pathogen in a manner like roaming animals seeking nutritious food. In the

natural environment of *C. elegans*, pathogenic and nutritious bacteria coexist, and animals must

employ mechanisms to avoid infection while still obtaining adequate nutrition. Low levels of

nutritious food, the presence of pathogen, or an absence of mates each represents a suboptimal

environment and may induce a common internal state that promotes exploration and enhances

the chances of encountering more favorable conditions.

Materials and Methods

C. elegans **strains:** *C. elegans* was maintained on *E. coli* OP50 as previously described

(Brenner, 1974). Daf-c strains were grown at 16°C. See Table 1 for a complete list of strains used

in this study.

Design of potential *scd-2* gain-of-function alleles: We performed a protein alignment of *C.*

elegans SCD-2 and human ALK in NCBI BLAST (Figure 4C). Using a list of known oncogenic

ALK mutations (Holla et al., 2017), we screened these residues for conservation or similarity

between the *C. elegans* and human protein sequences. Genome editing was done by SunyBiotech

using CRISPR technology. Alleles were evaluated in a dauer assay for a gain-of-function Daf-c

phenotype, and validated in trans-heterozygote analysis with *scd-2(sa249)* for a dominant

phenotype (Boor, 2022).

Preparation of food condition plates: Unless otherwise indicated, all assays were performed on

NGM plates with no peptone. Aztreonam-treated bacteria was prepared as previously reported

(Gruninger et al., 2008), and 40 µL aztreonam-treated food was added to plates containing 10

µg/mL aztreonam. "Fed" plates for *daf-7* quantification were seeded with 40 µL OP50 grown

overnight in a shaking LB culture at 37˚C. "Bottom" plates were seeded with 250 µL OP50, and

the agar was inverted with a spatula immediately prior to adding animals. "Lid" plates were

prepared by pouring a spot of NGM agar on the inside of the lid of the plate and seeding 100 µL

of OP50 on to the spot. PA14 was grown as previously described and seeded onto SKA plates, as

were *E. coli* OP50 controls for these experiments (Meisel et al., 2014).

pdaf-7::gfp **quantification assays:** Plates of gravid animals were bleached and eggs were

dropped onto NGM plates seeded with OP50 and grown at 20˚C for 67 hours, unless otherwise

noted. On the day of imaging, 15-30 day one adult animals were transferred by picking to assay

plates, where they were incubated for 5 hours at 20˚C. Animals to be imaged were mounted on

glass slides with agarose pads and 50 mM sodium azide or 5 mM Levamisole. All imaging for

pictures were conducted on the Zeiss Axioimager Z1. Quantification of GFP brightness was

derived from maximum fluorescence values within the ASJ neurons in FIJI.

Exploration assay: The exploration assay was performed loosely as previously described with

several modifications (Flavell et al., 2013). Thirty-five millimeter NGM plates with no peptone

were seeded with 500 µL OP50 grown overnight in LB so that the lawn covered the entire

surface of the plate. On the day of the assay, one day-one adult animal was placed on the plate

and allowed to explore for two hours before being removed. The plate was then superimposed on

a grid of 3.5 mm squares and the number of squares crossed by the *C. elegans* tracks was

manually counted.

Roaming/dwelling assay: Animals were egg-laid and grown to day-one adults. 10 cm NGM

plates without peptone were seeded with 2 mL stationary phase *E. coli* OP50 grown overnight in

LB. Assays were performed with ~20 animals inside a 6 cm copper ring placed in the center of

the seeded plate. After one hour for the animals to adjust to their new environment, videos were

recorded at 3.75 frames per second for 1.5-3 hours. Videos were analyzed using MBF

Biosciences WormLab software (*WormLab*, 2020).

Measurements of speed and bending angle (midpoint) were averaged over 10 second

intervals, and values for each 10-second interval were plotted on a scatter plot of speed (μm/sec)

vs. bending angle (degrees). Quantification of fraction of time spent roaming or dwelling was

done by segregating the points of the scatter plots by a horizontal line whose placement was

based on the distribution of points in the control condition for each experiment. Values for this

speed cutoff ranged from y=7 to y=12. Points falling above the line were classified as roaming,

and those below the line were classified as dwelling. The fraction of time spent roaming was

calculated for each animal based off the speed cutoff. Outliers were removes using a ROUT test

with Q = 1%. The durations of roaming and dwelling states were determined manually by

analyzing average speed values for successive ten-second intervals. Only states lasting longer

than five intervals (fifty seconds) were considered.

Statistics: All statistical analysis was performed using the GraphPad Prism software (Graphpad

Prism, RRID: SCR_002798). Statistical tests used are indicated in each figure legend.

Acknowledgements: We thank Bob Horvitz and the Caenorhabditis Genetics Center, which is

funded by the NIH Office of Research Infrastructure Programs (P40 OD010440), for strains, and

Cori Bargmann for Cre plasmids. We thank current and past members of the Kim lab for

discussions. Finally, we acknowledge financial support from NIH grant R35GM141794.

Strain Name	Genotype	Source
N2	Wild type	Caenorhabditis Genetics Center (CGC)
JT249	*scd-2(sa249)*	CGC
RB783	*scd-2(ok565)*	CGC
JC2154	*hen-1(tm501)*	CGC
PHX2455	*scd-2(syb2455)*	This study/SunyBiotech
FK181	*ksIs2[pdaf-7::gfp; rol-6(su1006)]*	CGC
ZD2540	*ksIs2; scd-2(sa249)*	This study
ZD930	*ksIs2; scd-2(ok565)*	This study
ZD918	*ksIs2; hen-1(tm501)*	This study
ZD2605	*ksIs2; scd-2(syb2455)*	This study
CB1372	*daf-7(e1372)*	CGC
ZD715	*daf-7(ok3125)*	(Meisel et al., 2014)
ZD695	*daf-7(ok3125);qdEx34[ptrx-1::daf-7;pges-1::GFP]*	(Meisel et al., 2014)
ZD696	*daf-7(ok3125);qdEx35[ptrx-1::daf-7;pges-1::GFP]*	(Meisel et al., 2014)
ZD2632	*ksIs2; del-3(ok2613);del-7(ok1187)*	This study
MT14984	*tph-1(n4622)*	Horvitz Lab
ZD667	*ksIs2; tph-1(n4622)*	This study
ZD2079	*pdfr-1(ok3425)*	(Hilbert & Kim, 2018)
PHX3826	*pdfr-1(syb3826)*	This study/SunyBiotech
ZD1987	*ksIs2; pdfr-1(ok3425); him-5(e1490)*	(Hilbert & Kim, 2018)
ZD2633	*ksIs2; pdfr-1(syb3826); him-5(e1490)*	This study
ZD2721	*scd-2(syb5845 syb6052) Backcrossed x2*	This study
ZD2722	*daf-7(syb5855 syb5965) qdEx[ptrx-1::Cre; pofm-1::gfp] Backcrossed x2*	This study
ZD2752	*scd-2(syb5845 syb6052); ex[gcy-28.dp::cre; ofm-1p::gfp]*	This study
ZD2766	*scd-2(syb5845 syb6052); ex[gcy-28.dp::cre; ofm-1p::gfp]; ksIs2*	This study
ZD1005	*him-5(e1390); ksIs2*	(Hilbert & Kim, 2017)

461
462   **Table 1.** Complete list of *C. elegans* strains used in this study.

464 **Figure 1. Ingestion of bacterial food inhibits *daf-7* expression in the ASJ neurons.**

465   A. Schematic of experimental exposure to varied food conditions. Animals were grown to

466   adulthood on edible *E. coli* OP50, then transferred to various experimental food

467   conditions: "Fed": *E. coli* OP50; "Aztreonam": OP50 treated with aztreonam; "Bottom":

OP50 was seeded underneath the agar of the plate; "Lid": Animals placed on agar with no food but where food was seeded on a spot of agar on the inside of the lid of the plate; "Empty": no food.

B. Model for the convergence of external and ingested food signals on *daf-7* expression in the ASJ neurons.

C. *pdaf-7::gfp* expression pattern in animals under different food conditions, from left to right: "Fed," "Aztreonam," "Bottom," "Lid," "Empty." Filled triangles indicate the ASI neurons; open triangles indicate the ASJ neurons. Scale bar indicates 50 μm.

D. E. F. Maximum fluorescence values of *pdaf-7::gfp* in the ASJ neurons of adult animals under various food conditions. Each point represents an individual animal, and error bars indicate standard deviation. ****$p<0.0001$, ns not significant as determined by an unpaired two-tailed t-test.

G. Maximum fluorescence values of *pdaf-7::gfp* in the ASJ neurons of adult animals at various time points after being moved from "Fed" to "Bottom" conditions. Each point represents an individual animal, and error bars indicate standard deviation. ****$p<0.0001$, * $p<0.05$, ns not significant as determined by an unpaired two-tailed t-test.

H. Maximum fluorescence values of *pdaf-7::gfp* in the ASJ neurons of adult wild type and *del-3(ok2613); del-7(ok1187)* under "Fed" conditions. Each point represents an individual animal, and error bars indicate standard deviation. ***$p<0.001$, as determined by an unpaired two-tailed t-test.

489
Figure 2. *daf-7* expression in the ASJ neurons promotes roaming.

A. Fraction of time roaming (left), and number of squares entered in exploration assay

(right) of wild type, *daf-7(e1372)* and *daf-7(ok3125)* animals. Each point represents an

individual animal, and error bars indicate standard deviation ****p<0.0001, **p<0.01,

*p<0.05 as determined by an unpaired two-tailed t-test.

B. Fraction of time roaming of wild type, *daf-7(ok3125),* and two independent lines where

daf-7 cDNA was expressed under the *trx-1* promoter in a *daf-7(ok3125)* background.

Each point represents an individual animal, and error bars indicate standard deviation.

**p<0.01, *p<0.05 as determined by an unpaired two-tailed t-test.

C. Fraction of time roaming (left), and number of squares entered in exploration assay

(right) of wild type, *daf-7(e1372), daf-7(ok3125),* and a floxed *daf-7* strain with and

without Cre expressed under the *trx-1* promoter. Each point represents an individual

animal, and error bars indicate standard deviation. ** p<0.01, *p<0.05 as determined by

an unpaired two-tailed t-test.

D. Model of positive-feedback relationship between food ingestion, *daf-7* expression in the

ASJ neurons, and roaming.

E. Sample trace files of representative individual wild-type, *daf-7(e1372),* and *daf-*

7(ok3125) animals. R = roaming, D= dwelling.

F. Duration of dwelling (open circles) and roaming (closed circles) states for wild type, *daf-*

7(e1372) and *daf-7(ok3125)* animals. Each point represents a discrete roaming or

dwelling period. Error bars indicate standard deviation. ***p<0.001, *p<0.05, ns not

significant as determined by an unpaired two-tailed t-test.

Figure 3. Known modulators of roaming and dwelling affect *daf-7* expression in the ASJ neurons

A. Fraction of time roaming of wild type and *tph-1(n4622)* animals. Each point represents an individual animal, and error bars indicate standard deviation. ****p<0.0001 as determined by an unpaired two-tailed t-test.

B. Maximum fluorescence values of *pdaf-7::gfp* in the ASJ neurons of adult wild type and *tph-1(n4622)* animals under "Fed" and "Bottom" conditions. Each point represents an individual animal, and error bars indicate standard deviation. ****p<0.0001, ***p<0.001 as determined by an unpaired two-tailed t-test.

C. Fraction of time roaming of wild type, *pdfr-1(ok3425),* and *pdfr-1(syb3826)* animals. Each point represents an individual animal, and error bars indicate standard deviation. ****p<0.0001, **p<0.01 as determined by an unpaired two-tailed t-test.

526   D. Maximum fluorescence values of *pdaf-7::gfp* in the ASJ neurons of adult wild type, *pdfr-*

527   *1(ok3425),* and *pdfr-1(syb3826)* animals under "Fed" and "Bottom" conditions. Each

528   point represents an individual animal, and error bars indicate standard deviation.

529   ***p<0.001, **p<0.01, ns not significant as determined by an unpaired two-tailed t-test.

Figure 4. SCD-2 regulates *daf-7* expression in the ASJ neurons and roaming behavior

A. Maximum fluorescence values of *pdaf-7::gfp* in the ASJ neurons of adult wild type, *scd-*

2(sa249), and *scd-2(syb2455)* animals exposed to *E. coli* OP50 and *P. aeruginosa* PA14.

Each point represents an individual animal, and error bars indicate standard deviation.

****$p < 0.0001$, as determined by an unpaired two-tailed t-test.

B. Maximum fluorescence values of *pdaf-7::gfp* in the ASJ neurons of adult wild type, *hen-*

1(tm501), scd-2(ok565) and *scd-2(sa249)* animals under "Fed" and "Bottom" conditions.

Each point represents an individual animal, and error bars indicate standard deviation.

****$p < 0.0001$, *$p < 0.05$ as determined by an unpaired two-tailed t-test.

C. Sequence alignment of the kinase domains of human ALK (top) and *C. elegans* SCD-2

(bottom). Amino acids highlighted in black are identical and those in gray are similar.

F1174/F1029 is outlined in green.

D. Maximum fluorescence values of *pdaf-7::gfp* in the ASJ neurons of adult wild type, *scd-*

2(ok565), scd-2(sa249), and *scd-2(syb2455)* animals under "Fed" and "Bottom"

conditions. Each point represents an individual animal, and error bars indicate standard

deviation. ****$p < 0.0001$, *$p < 0.05$, ns not significant as determined by an unpaired two-

tailed t-test.

E. Maximum fluorescence values of *pdaf-7::gfp* in the ASJ neurons of adult wild type,

floxed *scd-2, scd-2(sa249), scd-2(ok565),* and floxed *scd-2* animals with a transgene

expressing Cre under the AIA-specific *gcy-28.d* promoter under "Fed" and "Bottom"

conditions. Each point represents an individual animal, and error bars indicate standard

deviation. **$p < 0.01$ as determined by an unpaired two-tailed t-test.

F. Fraction of time spent roaming (left) and number of squares entered in exploration assay

(right) for wild type, *scd-2(sa249), scd-2(ok565)* and *hen-1(tm501)* animals. Each point

represents an individual animal. Error bars indicate standard deviation. ****$p < 0.0001$,

***$p < 0.001$, *$p < 0.05$ as determined by an unpaired two-tailed t-test.

G. Duration of roaming (closed circles) and dwelling (open circles) states for wild type, *scd-*

2(ok565) and *hen-1(tm501*. Each point represents a discrete roaming or dwelling period.

Error bars indicate standard deviation. **$p < 0.01$, *$p < 0.05$, ns not significant as

determined by an unpaired two-tailed t-test.

H. Fraction of time spent roaming (left) and number of squares entered in exploration assay

(right) for wild type and *scd-2(syb2455)* animals. Each point represents an individual

animal. Error bars indicate standard deviation. ****$p < 0.0001$, **$p < 0.01$ as determined by

an unpaired two-tailed t-test.

I. Duration of roaming (closed circles) and dwelling (open circles) states for wild type and

scd-2(syb2455) animals. Each point represents a discrete roaming or dwelling period.

Error bars indicate standard deviation. ***$p < 0.001$, ns not significant as determined by

an unpaired two-tailed t-test.

J. Left: Fraction of time spent roaming in worm tracker assay for *scd-2(sa249), scd-*

2(ok565) and floxed *scd-2* with or without a transgene expressing Cre under the AIA-

specific *gcy-28.d* promoter. Right: number of squares entered in exploration assay for

wild type, *scd-2(sa249), scd-2(ok565),* and floxed *scd-2* with or without a transgene

expressing Cre under the AIA-specific *gcy-28.d* promoter. Each point represents an

individual animal. Error bars indicate standard deviation. **$p < 0.01$, *$p < 0.05$, ns not

significant as determined by an unpaired two-tailed t-test.

577
Figure 5. DAF-7 expression in the ASJ neurons correlates with an increase in roaming

behavior under various conditions

A. Maximum fluorescence values of *pdaf-7::gfp* in the ASJ neurons of adult hermaphrodites

and males under "Fed" and "Aztreonam" conditions. Each point represents an individual

animal, and error bars indicate standard deviation. **p<0.01 as determined by an

unpaired t-test.

B. Fraction of time roaming of wild-type hermaphrodites and males. Each point represents

an individual animal, and error bars indicate standard deviation. ****$p<0.0001$ as

determined by an unpaired t-test.

C. Maximum fluorescence values of *pdaf-7::gfp* in the ASJ neurons of adult hermaphrodites

fed *E. coli* OP50, *P. aeruginosa* PA14 or *P. aeruginosa* PA14ΔgacA. Each point

represents an individual animal, and error bars indicate standard deviation.

****$p<0.0001$, **$p<0.01$ as determined by an unpaired t-test.

D. Fraction of time roaming of wild-type hermaphrodites on *E. coli* OP50, *P. aeruginosa*

PA14, or PA14 ΔgacA. Each point represents an individual animal, and error bars

indicate standard deviation. ****$p<0.0001$, ***$p<0.001$, *$p<0.05$ as determined by an

unpaired t-test.

E. *daf-7* expression in the ASJ neurons responds to environmental conditions and is

correlated with internal state. Under favorable conditions, *daf-7* is not expressed in the

ASJ neurons, and this is correlated with an internal state that favors exploitation of the

animal's current environment. Under unfavorable conditions, *daf-7* expression is induced

in the ASJ neurons, consistent with an internal state that favors exploration.

References

Bargmann, C. I. (2012). Beyond the connectome: how neuromodulators shape neural circuits. *Bioessays*, *34*(6), 458-465. https://doi.org/10.1002/bies.201100185

Ben Arous, J., Laffont, S., & Chatenay, D. (2009). Molecular and sensory basis of a food related two-state behavior in C. elegans. *PLoS One*, *4*(10), e7584. https://doi.org/10.1371/journal.pone.0007584

Boor, S. (2022). *Genetic Analysis of Bacterial Food Perception and its Influence on Foraging Behavior in C. elegans* Massachusetts Institute of Technology]. Cambridge, MA.

Brenner, S. (1974). The genetics of Caenorhabditis elegans. *Genetics*, *77*(1), 71-94. https://doi.org/10.1093/genetics/77.1.71

Chen, Y., Takita, J., Choi, Y. L., Kato, M., Ohira, M., Sanada, M., Wang, L., Soda, M., Kikuchi, A., Igarashi, T., Nakagawara, A., Hayashi, Y., Mano, H., & Ogawa, S. (2008). Oncogenic mutations of ALK kinase in neuroblastoma. *Nature*, *455*(7215), 971-974. https://doi.org/10.1038/nature07399

Cheng, L. Y., Bailey, A. P., Leevers, S. J., Ragan, T. J., Driscoll, P. C., & Gould, A. P. (2011). Anaplastic lymphoma kinase spares organ growth during nutrient restriction in Drosophila. *Cell*, *146*(3), 435-447. https://doi.org/10.1016/j.cell.2011.06.040

Churgin, M. A., McCloskey, R. J., Peters, E., & Fang-Yen, C. (2017). Antagonistic Serotonergic and Octopaminergic Neural Circuits Mediate Food-Dependent Locomotory Behavior in Caenorhabditis elegans. *J Neurosci*, *37*(33), 7811-7823. https://doi.org/10.1523/JNEUROSCI.2636-16.2017

Connolly, K. J. (1966). Locomotor activity in Drosophila as a function of food deprivation. *Nature*, *209*(5019), 224. https://doi.org/10.1038/209224a0

Dobosiewicz, M., Liu, Q., & Bargmann, C. I. (2019). Reliability of an interneuron response depends on an integrated sensory state. *Elife*, *8*. https://doi.org/10.7554/eLife.50566

Entchev, E. V., Patel, D. S., Zhan, M., Steele, A. J., Lu, H., & Ch'ng, Q. (2015). A gene-expression-based neural code for food abundance that modulates lifespan. *Elife*, *4*, e06259. https://doi.org/10.7554/eLife.06259

Flavell, S. W., Gogolla, N., Lovett-Barron, M., & Zelikowsky, M. (2022). The emergence and influence of internal states. *Neuron*, *110*(16), 2545-2570. https://doi.org/10.1016/j.neuron.2022.04.030

Flavell, S. W., Pokala, N., Macosko, E. Z., Albrecht, D. R., Larsch, J., & Bargmann, C. I. (2013). Serotonin and the neuropeptide PDF initiate and extend opposing behavioral states in C. elegans. *Cell*, *154*(5), 1023-1035. https://doi.org/10.1016/j.cell.2013.08.001

Flavell, S. W., Raizen, D. M., & You, Y. J. (2020). Behavioral States. *Genetics*, *216*(2), 315-332. https://doi.org/10.1534/genetics.120.303539

Franco, R., Rocco, G., Marino, F. Z., Pirozzi, G., Normanno, N., Morabito, A., Sperlongano, P., Stiuso, P., Luce, A., Botti, G., & Caraglia, M. (2013). Anaplastic lymphoma kinase: a glimmer of hope in lung cancer treatment? *Expert Rev Anticancer Ther*, *13*(4), 407-420. https://doi.org/10.1586/era.13.18

Fujiwara, M., Sengupta, P., & McIntire, S. L. (2002). Regulation of body size and behavioral state of C. elegans by sensory perception and the EGL-4 cGMP-dependent protein kinase. *Neuron*, *36*(6), 1091-1102. https://doi.org/10.1016/s0896-6273(02)01093-0

George, R. E., Sanda, T., Hanna, M., Frohling, S., Luther, W., 2nd, Zhang, J., Ahn, Y., Zhou, W., London, W. B., McGrady, P., Xue, L., Zozulya, S., Gregor, V. E., Webb, T. R., Gray, N. S., Gilliland, D. G., Diller, L., Greulich, H., Morris, S. W., . . . Look, A. T. (2008).

Activating mutations in ALK provide a therapeutic target in neuroblastoma. *Nature*, *455*(7215), 975-978. https://doi.org/10.1038/nature07397

Golden, J. W., & Riddle, D. L. (1982). A pheromone influences larval development in the nematode Caenorhabditis elegans. *Science*, *218*(4572), 578-580. https://doi.org/10.1126/science.6896933

Golden, J. W., & Riddle, D. L. (1984a). ACaenorhabditis elegans dauer-inducing pheromone and an antagonistic component of the food supply. *J Chem Ecol*, *10*(8), 1265-1280. https://doi.org/10.1007/BF00988553

Golden, J. W., & Riddle, D. L. (1984b). The Caenorhabditis elegans dauer larva: developmental effects of pheromone, food, and temperature. *Dev Biol*, *102*(2), 368-378. https://doi.org/10.1016/0012-1606(84)90201-x

Golden, J. W., & Riddle, D. L. (1984c). A pheromone-induced developmental switch in Caenorhabditis elegans: Temperature-sensitive mutants reveal a wild-type temperature-dependent process. *Proc Natl Acad Sci U S A*, *81*(3), 819-823. https://doi.org/10.1073/pnas.81.3.819

Greer, E. R., Perez, C. L., Van Gilst, M. R., Lee, B. H., & Ashrafi, K. (2008). Neural and molecular dissection of a C. elegans sensory circuit that regulates fat and feeding. *Cell Metab*, *8*(2), 118-131. https://doi.org/10.1016/j.cmet.2008.06.005

Gruninger, T. R., Gualberto, D. G., & Garcia, L. R. (2008). Sensory perception of food and insulin-like signals influence seizure susceptibility. *PLoS Genet*, *4*(7), e1000117. https://doi.org/10.1371/journal.pgen.1000117

Gutman, R., Yosha, D., Choshniak, I., & Kronfeld-Schor, N. (2007). Two strategies for coping with food shortage in desert golden spiny mice. *Physiol Behav*, *90*(1), 95-102. https://doi.org/10.1016/j.physbeh.2006.08.033

Hallberg, B., & Palmer, R. H. (2013). Mechanistic insight into ALK receptor tyrosine kinase in human cancer biology. *Nat Rev Cancer*, *13*(10), 685-700. https://doi.org/10.1038/nrc3580

Harris, N., Bates, S. G., Zhuang, Z., Bernstein, M., Stonemetz, J. M., Hill, T. J., Yu, Y. V., Calarco, J. A., & Sengupta, P. (2023). Molecular encoding of stimulus features in a single sensory neuron type enables neuronal and behavioral plasticity. *Curr Biol*, *33*(8), 1487-1501 e1487. https://doi.org/10.1016/j.cub.2023.02.073

Herbers, J. M. (1981). Time resources and laziness in animals. *Oecologia*, *49*(2), 252-262. https://doi.org/10.1007/BF00349198

Hilbert, Z. A., & Kim, D. H. (2017). Sexually dimorphic control of gene expression in sensory neurons regulates decision-making behavior in C. elegans. *Elife*, *6*. https://doi.org/10.7554/eLife.21166

Hilbert, Z. A., & Kim, D. H. (2018). PDF-1 neuropeptide signaling regulates sexually dimorphic gene expression in shared sensory neurons of C. elegans. *Elife*, *7*. https://doi.org/10.7554/eLife.36547

Holla, V. R., Elamin, Y. Y., Bailey, A. M., Johnson, A. M., Litzenburger, B. C., Khotskaya, Y. B., Sanchez, N. S., Zeng, J., Shufean, M. A., Shaw, K. R., Mendelsohn, J., Mills, G. B., Meric-Bernstam, F., & Simon, G. R. (2017). ALK: a tyrosine kinase target for cancer therapy. *Cold Spring Harb Mol Case Stud*, *3*(1), a001115. https://doi.org/10.1101/mcs.a001115

Inoue, T., & Thomas, J. H. (2000). Suppressors of transforming growth factor-beta pathway mutants in the Caenorhabditis elegans dauer formation pathway. *Genetics*, *156*(3), 1035-1046. https://doi.org/10.1093/genetics/156.3.1035

Ishihara, T., Iino, Y., Mohri, A., Mori, I., Gengyo-Ando, K., Mitani, S., & Katsura, I. (2002). HEN-1, a secretory protein with an LDL receptor motif, regulates sensory integration and learning in Caenorhabditis elegans. *Cell*, *109*(5), 639-649. https://doi.org/10.1016/s0092-8674(02)00748-1

Iwahara, T., Fujimoto, J., Wen, D., Cupples, R., Bucay, N., Arakawa, T., Mori, S., Ratzkin, B., & Yamamoto, T. (1997). Molecular characterization of ALK, a receptor tyrosine kinase expressed specifically in the nervous system. *Oncogene*, *14*(4), 439-449. https://doi.org/10.1038/sj.onc.1200849

Janoueix-Lerosey, I., Lequin, D., Brugieres, L., Ribeiro, A., de Pontual, L., Combaret, V., Raynal, V., Puisieux, A., Schleiermacher, G., Pierron, G., Valteau-Couanet, D., Frebourg, T., Michon, J., Lyonnet, S., Amiel, J., & Delattre, O. (2008). Somatic and germline activating mutations of the ALK kinase receptor in neuroblastoma. *Nature*, *455*(7215), 967-970. https://doi.org/10.1038/nature07398

Ji, N., Madan, G. K., Fabre, G. I., Dayan, A., Baker, C. M., Kramer, T. S., Nwabudike, I., & Flavell, S. W. (2021). A neural circuit for flexible control of persistent behavioral states. *Elife*, *10*. https://doi.org/10.7554/eLife.62889

Johnson, R. E., Linderman, S., Panier, T., Wee, C. L., Song, E., Herrera, K. J., Miller, A., & Engert, F. (2020). Probabilistic Models of Larval Zebrafish Behavior Reveal Structure on Many Scales. *Curr Biol*, *30*(1), 70-82 e74. https://doi.org/10.1016/j.cub.2019.11.026

Kim, D. H., & Flavell, S. W. (2020). Host-microbe interactions and the behavior of Caenorhabditis elegans. *J Neurogenet*, *34*(3-4), 500-509. https://doi.org/10.1080/01677063.2020.1802724

Kyani-Rogers, T., Philbrook, A., McLachlan, I. G., Flavell, S. W., O'Donnell, M. P., & Sengupta, P. (2022). Developmental history modulates adult olfactory behavioral preferences via regulation of chemoreceptor expression in Caenorhabditiselegans. *Genetics*, *222*(3). https://doi.org/10.1093/genetics/iyac143

McLachlan, I. G., Kramer, T. S., Dua, M., DiLoreto, E. M., Gomes, M. A., Dag, U., Srinivasan, J., & Flavell, S. W. (2022). Diverse states and stimuli tune olfactory receptor expression levels to modulate food-seeking behavior. *Elife*, *11*. https://doi.org/10.7554/eLife.79557

Meisel, J. D., Panda, O., Mahanti, P., Schroeder, F. C., & Kim, D. H. (2014). Chemosensation of bacterial secondary metabolites modulates neuroendocrine signaling and behavior of C. elegans. *Cell*, *159*(2), 267-280. https://doi.org/10.1016/j.cell.2014.09.011

Morris, S. W., Kirstein, M. N., Valentine, M. B., Dittmer, K. G., Shapiro, D. N., Saltman, D. L., & Look, A. T. (1994). Fusion of a kinase gene, ALK, to a nucleolar protein gene, NPM, in non-Hodgkin's lymphoma. *Science*, *263*(5151), 1281-1284. https://doi.org/10.1126/science.8122112

Oranth, A., Schultheis, C., Tolstenkov, O., Erbguth, K., Nagpal, J., Hain, D., Brauner, M., Wabnig, S., Steuer Costa, W., McWhirter, R. D., Zels, S., Palumbos, S., Miller Iii, D. M., Beets, I., & Gottschalk, A. (2018). Food Sensation Modulates Locomotion by Dopamine and Neuropeptide Signaling in a Distributed Neuronal Network. *Neuron*, *100*(6), 1414-1428 e1410. https://doi.org/10.1016/j.neuron.2018.10.024

Orthofer, M., Valsesia, A., Magi, R., Wang, Q. P., Kaczanowska, J., Kozieradzki, I., Leopoldi, A., Cikes, D., Zopf, L. M., Tretiakov, E. O., Demetz, E., Hilbe, R., Boehm, A., Ticevic, M., Noukas, M., Jais, A., Spirk, K., Clark, T., Amann, S., . . . Penninger, J. M. (2020). Identification of ALK in Thinness. *Cell*, *181*(6), 1246-1262 e1222. https://doi.org/10.1016/j.cell.2020.04.034

738   Overton, J. M., & Williams, T. D. (2004). Behavioral and physiologic responses to caloric
739   restriction in mice. *Physiol Behav*, *81*(5), 749-754.
740   https://doi.org/10.1016/j.physbeh.2004.04.025

741   Pradhan, S., Quilez, S., Homer, K., & Hendricks, M. (2019). Environmental Programming of
742   Adult Foraging Behavior in C. elegans. *Curr Biol*, *29*(17), 2867-2879 e2864.
743   https://doi.org/10.1016/j.cub.2019.07.045

744   Reiner, D. J., Ailion, M., Thomas, J. H., & Meyer, B. J. (2008). C. elegans anaplastic lymphoma
745   kinase ortholog SCD-2 controls dauer formation by modulating TGF-beta signaling. *Curr*
746   *Biol*, *18*(15), 1101-1109. https://doi.org/10.1016/j.cub.2008.06.060

747   Ren, P., Lim, C. S., Johnsen, R., Albert, P. S., Pilgrim, D., & Riddle, D. L. (1996). Control of C.
748   elegans larval development by neuronal expression of a TGF-beta homolog. *Science*,
749   *274*(5291), 1389-1391. https://doi.org/10.1126/science.274.5291.1389

750   Rhoades, J. L., Nelson, J. C., Nwabudike, I., Yu, S. K., McLachlan, I. G., Madan, G. K., Abebe,
751   E., Powers, J. R., Colon-Ramos, D. A., & Flavell, S. W. (2019). ASICs Mediate Food
752   Responses in an Enteric Serotonergic Neuron that Controls Foraging Behaviors. *Cell*,
753   *176*(1-2), 85-97 e14. https://doi.org/10.1016/j.cell.2018.11.023

754   Russell, J. C., Epling, W. F., Pierce, D., Amy, R. M., & Boer, D. P. (1987). Induction of
755   voluntary prolonged running by rats. *J Appl Physiol (1985)*, *63*(6), 2549-2553.
756   https://doi.org/10.1152/jappl.1987.63.6.2549

757   Ryan, D. A., Miller, R. M., Lee, K., Neal, S. J., Fagan, K. A., Sengupta, P., & Portman, D. S.
758   (2014). Sex, age, and hunger regulate behavioral prioritization through dynamic
759   modulation of chemoreceptor expression. *Curr Biol*, *24*(21), 2509-2517.
760   https://doi.org/10.1016/j.cub.2014.09.032

761   Sawin, E. R., Ranganathan, R., & Horvitz, H. R. (2000). C. elegans locomotory rate is modulated
762   by the environment through a dopaminergic pathway and by experience through a
763   serotonergic pathway. *Neuron*, *26*(3), 619-631. https://doi.org/10.1016/s0896-
764   6273(00)81199-x

765   Schackwitz, W. S., Inoue, T., & Thomas, J. H. (1996). Chemosensory neurons function in
766   parallel to mediate a pheromone response in C. elegans. *Neuron*, *17*(4), 719-728.
767   https://doi.org/10.1016/s0896-6273(00)80203-2

768   Sengupta, P. (2013). The belly rules the nose: feeding state-dependent modulation of peripheral
769   chemosensory responses. *Curr Opin Neurobiol*, *23*(1), 68-75.
770   https://doi.org/10.1016/j.conb.2012.08.001

771   Shaw, W. M., Luo, S., Landis, J., Ashraf, J., & Murphy, C. T. (2007). The C. elegans TGF-beta
772   Dauer pathway regulates longevity via insulin signaling. *Curr Biol*, *17*(19), 1635-1645.
773   https://doi.org/10.1016/j.cub.2007.08.058

774   Shinkai, Y., Yamamoto, Y., Fujiwara, M., Tabata, T., Murayama, T., Hirotsu, T., Ikeda, D. D.,
775   Tsunozaki, M., Iino, Y., Bargmann, C. I., Katsura, I., & Ishihara, T. (2011). Behavioral
776   choice between conflicting alternatives is regulated by a receptor guanylyl cyclase, GCY-
777   28, and a receptor tyrosine kinase, SCD-2, in AIA interneurons of Caenorhabditis
778   elegans. *J Neurosci*, *31*(8), 3007-3015. https://doi.org/10.1523/JNEUROSCI.4691-
779   10.2011

780   Wexler, L. R., Miller, R. M., & Portman, D. S. (2020). C. elegans Males Integrate Food Signals
781   and Biological Sex to Modulate State-Dependent Chemosensation and Behavioral
782   Prioritization. *Curr Biol*, *30*(14), 2695-2706 e2694.
783   https://doi.org/10.1016/j.cub.2020.05.006

White, J. G., Southgate, E., Thomson, J. N., & Brenner, S. (1986). The structure of the nervous
system of the nematode Caenorhabditis elegans. *Philos Trans R Soc Lond B Biol Sci*,
314(1165), 1-340. https://doi.org/10.1098/rstb.1986.0056
Wolfe, G. S., Tong, V. W., Povse, E., Merritt, D. M., Stegeman, G. W., Flibotte, S., & van der
Kooy, D. (2019). A Receptor Tyrosine Kinase Plays Separate Roles in Sensory
Integration and Associative Learning in C. elegans. *eNeuro*, *6*(4).
https://doi.org/10.1523/ENEURO.0244-18.2019
Woodling, N. S., Aleyakpo, B., Dyson, M. C., Minkley, L. J., Rajasingam, A., Dobson, A. J.,
Leung, K. H. C., Pomposova, S., Fuentealba, M., Alic, N., & Partridge, L. (2020). The
neuronal receptor tyrosine kinase Alk is a target for longevity. *Aging Cell*, *19*(5), e13137.
https://doi.org/10.1111/acel.13137
WormLab. In. (2020). MBF Bioscience.
Yap, E. L., & Greenberg, M. E. (2018). Activity-Regulated Transcription: Bridging the Gap
between Neural Activity and Behavior. *Neuron*, *100*(2), 330-348.
https://doi.org/10.1016/j.neuron.2018.10.013
799