Disentangling the effects of hearing loss and age on amplitude modulation frequency selectivity

Jonathan Regev[1,a], Helia Relaño-Iborra[1], Johannes Zaar[1,2], and Torsten Dau[1,3]

[1] *Hearing Systems Section, Department of Health Technology, Technical University of Denmark, Kongens Lyngby,*

2800, Denmark

[2] *Eriksholm Research Centre, Snekkersten, 3070, Denmark*

[3] *Copenhagen Hearing and Balance Center, Copenhagen University Hospital, Rigshospitalet, Copenhagen, 2100,*

Denmark

Abstract

The processing and perception of amplitude modulation (AM) in the auditory system reflect a frequency-selective process, often described as a modulation filterbank. Previous studies on perceptual AM masking reported similar results for older listeners with hearing impairment (HI) and young listeners with normal hearing (NH), suggesting no effects of age or hearing loss on AM frequency selectivity. However, recent evidence has shown that age, independently of hearing loss, adversely affects AM frequency selectivity. Hence, this study aimed to disentangle the effects of hearing loss and age. A simultaneous AM masking paradigm was employed, utilizing a sinusoidal carrier at 2.8 kHz, narrow-band noise modulation maskers, and target modulation frequencies of 4, 16, 64, and 128 Hz. The results obtained from young (n=3, 24-30 years) and older (n=10, 63-77 years) HI listeners were compared to previously obtained data from young and older NH listeners. Notably, the HI listeners generally exhibited lower (unmasked) AM detection thresholds and greater AM frequency selectivity

---

[a] Corresponding author: joreg@dtu.dk

than their NH counterparts in both age groups. Overall, the results suggest that age negatively affects AM frequency selectivity for both NH and HI listeners, while hearing loss improves AM detection and AM selectivity, likely due to the loss of peripheral compression.

I. INTRODUCTION

Most sounds encountered in everyday environments are complex signals that exhibit dynamic fluctuations over time (Singh and Theunissen, 2003; Elliott and Theunissen, 2009). These fluctuations include variations in the temporal envelope, commonly referred to as amplitude modulation (AM). The auditory system demonstrates an acute sensitivity to AM, as evidenced by perceptual studies (e.g., Viemeister, 1979) and physiological investigations (see Joris *et al.*, 2004 for a comprehensive review). Similar to spectral masking effects in the audio-frequency domain, masking effects also occur in the AM frequency domain, resulting in reduced sensitivity to a target AM in the presence of a masking AM (Bacon and Grantham, 1989; Houtgast, 1989; Dau *et al.*, 1997a, 1997b; Ewert and Dau, 2000; Sek and Moore, 2003). Specifically, AM masking patterns provide evidence for a frequency-selective process, where the amount of AM masking decreases as the spectral distance between the masker and the target increases (Bacon and Grantham, 1989; Houtgast, 1989; Dau *et al.*, 1997a; Ewert and Dau, 2000; Ewert *et al.*, 2002; Sek and Moore, 2003; Moore *et al.*, 2009; Sek *et al.*, 2015; Füllgrabe *et al.*, 2021a, 2021b; Conroy *et al.*, 2023).

This frequency-selective characteristic of AM processing has been modelled using the concept of a modulation filterbank, based on the idea that AM fluctuations are decomposed through an array of relatively broad bandpass modulation filters with a constant quality (Q) factor of approximately 1-2 (e.g., Dau *et al.*, 1997a, 1999; Ewert and Dau, 2000). Computational modelling studies have successfully applied the modulation filterbank concept to simulate data from various experimental paradigms, including simultaneous and non-simultaneous spectral and temporal signal detection and masking conditions (Dau *et al.*, 1997a, 1997b, 1999; Verhey *et al.*, 1999; Ewert and Dau, 2000; Ewert *et al.*, 2002; Piechowiak *et al.*, 2007; Jepsen *et al.*, 2008; Jepsen and Dau, 2011; King *et al.*, 2019), sound texture perception (McDermott and Simoncelli, 2011; McDermott *et al.*, 2013; McWalter and Dau, 2015, 2017), auditory stream segregation (Elhilali *et al.*, 2009; Christiansen *et al.*, 2014), and speech

intelligibility (Jørgensen and Dau, 2011; Jørgensen et al., 2013; Relaño-Iborra et al., 2016, 2019; Zaar

and Dau, 2017; Zaar et al., 2017; Steinmetzger et al., 2019; Zaar and Carney, 2022; for a review, see

Relaño-Iborra and Dau, 2022). Furthermore, the modulation filterbank is conceptually consistent with

the temporal dimension of a 'two-dimensional' spectro-temporal modulation filterbank, inspired by

neural responses to spectro-temporally varying stimuli in the auditory cortex of ferrets (Kowalski *et*

al., 1996; Depireux *et al.*, 2001) and supported by data from perceptual learning and masking conditions

(Sabin *et al.*, 2012; Oetjen and Verhey, 2015, 2017; Conroy *et al.*, 2022), as well as models of speech

intelligibility (Elhilali *et al.*, 2003; Chi *et al.*, 2005; Zilany and Bruce, 2007; Chabot-Leclerc *et al.*, 2014).

Overall, the versatility of the modulation filterbank model suggests that AM frequency selectivity is

an essential auditory processing feature for quantitatively predicting perceptual data obtained with

dynamically varying sounds. However, likely due to a lack of behavioral data, no modelling study was

conducted investigating the effects of reducing the selectivity of the modulation filterbank, either due

to age or hearing loss.

Several studies have investigated the effects of age and hearing loss on AM perception, yielding

mixed results. In terms of age effects, some studies showed a significant age-related deterioration in

AM detection (He *et al.*, 2008; Füllgrabe *et al.*, 2015; Wallaert *et al.*, 2016), while other investigations

did not find any effects on AM detection or AM depth discrimination (Schoof and Rosen, 2014;

Paraouty *et al.*, 2016; Schlittenlacher and Moore, 2016; Paraouty and Lorenzi, 2017). Studies comparing

listeners with normal hearing to listeners with hearing impairment (referred to hereafter as NH and

HI listeners, respectively) have found either similar or lower (i.e., better) AM detection thresholds for

the HI listeners (e.g., Moore *et al.*, 1992, 1996; Moore and Glasberg, 2001; Füllgrabe *et al.*, 2003; Sek *et*

al., 2015; Schlittenlacher and Moore, 2016; Wallaert *et al.*, 2017; Wiinberg *et al.*, 2019). The observed

improvement in AM detection with hearing loss has been suggested to result from the loss of cochlear

compression typically associated with sensorineural hearing loss (SNHL; Moore and Oxenham, 1998),

which may lead to an increased internal representation of AM depth for HI listeners (Moore *et al.*,

1996; Jennings *et al.*, 2018). Additionally, the amplified neural coding of the envelope observed in

mammals and humans with SNHL (Kale and Heinz, 2010, 2012; Henry *et al.*, 2014; Zhong *et al.*, 2014;

Millman *et al.*, 2017; Goossens *et al.*, 2018; Decruy *et al.*, 2020) may contribute, at least in part, to the

effective increase in perceived AM depth with hearing loss. However, hearing loss has been shown to

have a detrimental effect on supra-threshold AM depth discrimination (Schlittenlacher and Moore,

2016; Wiinberg *et al.*, 2019), suggesting that the increase in perceived AM depth is not necessarily

beneficial for AM depth discrimination.

Behavioral studies on AM frequency selectivity, where age and hearing loss were often

confounded, generally did not yield conclusive evidence of an effect of hearing loss (Takahashi and

Bacon, 1992; Lorenzi *et al.*, 1997; Sek *et al.*, 2015). These studies focused on masked-threshold patterns

(MTPs), which assess the detection threshold of an AM target in relation to the (center) frequency of

a masker of fixed modulation depth. Takahashi and Bacon (1992) compared MTPs (using noise

carriers and sinusoidal modulation targets and maskers) from ten young NH listeners and three groups

of ten older listeners with mild hearing loss (classified as 'age-appropriate' normal hearing). While a

larger proportion of the older listeners could not complete some conditions of the AM masking task,

no combined effect of age and hearing loss on AM frequency selectivity was found. Lorenzi *et al.*

(1997) employed a similar approach with four young NH and three young HI listeners, finding

comparable MTPs across the two groups, despite observations suggesting that HI listeners may

experience a greater masking effect from modulation maskers located below the target modulation

frequency. Additionally, Sek *et al.* (2015), using sinusoidal carriers and sinusoidal modulation maskers,

found no differences in AM masking patterns between six young NH and nine older HI listeners.

These results suggested that neither age nor hearing loss affect AM frequency selectivity (Sek *et al.*,

2015).

However, recent results by Regev *et al.* (2023) showed a reduction in AM frequency selectivity

associated with age when comparing young and older listeners with clinically normal hearing. These

results are qualitatively consistent with the observations of reduced temporal selectivity inferred from

auditory cortical functional magnetic resonance imaging (fMRI) responses by Erb *et al.* (2020). Regev

et al. (2023), using sinusoidal carriers and noise modulation maskers, compared MTPs for eleven young

NH listeners and ten older NH listeners (showing only a small mismatch in absolute hearing threshold

at the test frequency of 10 dB on average; see Section IV in Regev *et al.*, 2023 for an in-depth

discussion). They found an age-related increase (i.e., deterioration) in AM detection thresholds and a

decrease in AM frequency selectivity, with estimated modulation-filter Q factors that were

approximately halved for the older NH listeners compared to the young NH listeners. The contrast

between these recent findings and previous studies that did not identify a combined effect of age and

hearing loss on AM frequency selectivity (Takahashi and Bacon, 1992; Sek *et al.*, 2015) may have

occurred because hearing loss and age have opposite effects with regard to AM frequency selectivity.

In other words, hearing loss may 'counteract' the detrimental effects of ageing. However, a direct

comparison of MTPs between young NH, older NH, and similarly aged older HI listeners, tested

using the same experimental approach, has not yet been conducted.

The present study therefore aimed to independently evaluate the effects of hearing loss and age

on AM frequency selectivity by collecting data from HI listeners who were approximately age-matched

to the older NH listeners included in Regev *et al.* (2023), using the same experimental approach. Young

HI listeners were also included to explore whether a perceptual 'benefit' from hearing loss could be

observed in the absence of ageing effects. The study involved ten older and three young HI listeners,

ranging in age from 63 to 77 and 24 to 30 years, respectively. MTPs were collected using an AM

masking paradigm with sinusoidal carriers and bandpass Gaussian-noise modulation maskers (Ewert

et al., 2002). Additionally, unmasked AM detection thresholds were obtained using sinusoidal carriers.

The data obtained from the older HI listeners were compared to those obtained from the older NH

listeners reported in Regev *et al.* (2023) to assess the effects of hearing loss on AM frequency selectivity.

The data from the older HI listeners were also compared to those for the young NH listeners of Regev

et al. (2023) to determine whether the lack of a combined effect of age and hearing loss observed in

previous studies could be replicated (Takahashi and Bacon, 1992; Sek *et al.*, 2015). The data reported

in Regev *et al.* (2023) are publicly available from Regev *et al.* (2024). Despite the limited number of

young HI listeners, their data were included to provide insight into the effects of hearing loss in the

absence of ageing.

II.   METHODS

A. Listeners

Ten older HI listeners, aged 63-77 years (mean age = 70.4 years, 2 female) and three young HI

listeners, aged 24-30 years (mean age = 26 years, all female) participated in the study. On average, the

older HI listeners were 4.5 years older than the older NH listeners tested in Regev *et al.* (2023), while

the young HI listeners were less than one year older than the young NH listeners. All listening

experiments were conducted monaurally. The individual audiograms for the test ears are shown in

Figure 1. The older HI listeners had a high-frequency sloping hearing loss, classified as an N2 or N3

standard audiogram (Bisgaard *et al.*, 2010).[1] The young HI listeners had more heterogeneous

audiograms. For each listener, the test ear was selected as the one where the audiometric threshold at

3 kHz was in the range 50 to 60 dB HL. These limits were chosen to ensure that the listeners had a

significant hearing loss near the AM carrier frequency (2.8 kHz) and, at the same time, that the sound

pressure level (SPL) would not exceed 85 dB, to avoid discomfort. All listeners had a SNHL, as

---

[1] The classification of the collected audiogram as one of the standard audiograms was performed using the lowest mean squared error between the standard audiograms and the audiometric thresholds at frequencies from 1 to 8 kHz.

indicated by air-bone gaps ≤ 15 dB.[2] Working memory capacity was assessed with a reverse digit span

test (RDS), implemented binaurally, as described in Fuglsang *et al.* (2020), with an added linear gain

according to the Cambridge formula (CamEq; Moore and Glasberg, 1998) using a maximum-gain

limit of 30 dB at any frequency. The young and older HI listeners were found to have similar average

RDS scores (0.56 and 0.53 on a normalized scale, respectively), which are also comparable to the

average RDS scores reported for the young NH listeners in Regev *et al.* (2023) and slightly, but not

significantly, higher than for Regev *et al.*'s older NH cohort (0.59 and 0.42, respectively).

All participants gave written informed consent. Ethical approval for the study was provided by

the Science Ethics Committee of the Capital Region in Denmark (reference H-16036391). All

participants were paid for their time.

Figure 1: Individual and average audiograms for the test ear, for all participants. The left and right

panels show the audiograms for the young and older HI listeners, respectively. Individual audiograms

are shown in light gray, and group averages and standard deviations are shown in black.

---

[2] Except for one young HI listener who showed a gap of 20 dB at 4 kHz and one older HI listener who showed a gap of 20 dB at 1 kHz.

B. Procedure and apparatus

The experimental conditions and parameters were the same as described in Regev *et al.* (2023), with a few modifications to compensate for audibility loss. The stimulus presentation level in all experiments was adjusted to be 30 dB sensation level (SL) for all HI listeners, based on the hearing threshold at the carrier frequency, in an approach similar to that of Sek *et al.* (2015). This resulted in presentation levels of 78-80 dB SPL for the young HI listeners and 73-85 dB SPL for the older HI listeners (79 dB SPL on average for both groups).

All listening tests were conducted in double-walled soundproof booths. All thresholds were measured using a 3-interval 3-alternative forced-choice (3I-3AFC) paradigm with a 1-up 2-down tracking procedure, measuring the 70.7%-correct point on the psychometric function (Levitt, 1971). The experiments were implemented using the AFC toolbox in MATLAB (Ewert, 2013). The stimuli were digitally generated in MATLAB (The Mathworks Inc., 2015; Natick, Massachusetts) with a sampling frequency of 48 kHz, converted to analog using an RME Fireface soundcard (Haimhausen, Germany), and played through Sennheiser HDA200 headphones (Wedemark, Germany). The output level was calibrated using a GRAS RA0039 ear simulator (Holte, Denmark) and the frequency response of the headphones was compensated using an inverse digital filter to achieve a flat response at the tympanic membrane. The listeners received visual feedback during the tests ("correct"/"incorrect"). The order of testing was the same for both groups, as summarized in Appendix A.

The hearing threshold for the unmodulated pure-tone carrier at 2.8 kHz was measured adaptively. The signal had a duration of 300 ms, including 50-ms raised-cosine ramps, and the intervals were separated by 600 ms of silence. The step size was initially 8 dB, was then reduced to 4 dB after the first lower reversal, and was finally reduced to 2 dB after the second lower reversal. For any run, six reversals were obtained using the smallest step size and the threshold was defined as the average SPL

of the target tone across them. The threshold measurement was repeated at least three times, and the

final threshold was computed as the average across all repetitions.

C. Experimental conditions

1. Stimuli

The AM detection task and masked-threshold patterns (MTPs; Ewert and Dau, 2000) shared

common aspects of the stimuli. For both tasks, the thresholds were measured for sinusoidal AM at

target modulation frequencies of 4, 16, 64, and 128 Hz, imposed on a 2.8-kHz sinusoidal carrier. The

carrier had a duration of 600 ms, including 50-ms raised-cosine onset and offset ramps. The target

modulation (and masker modulation in the MTP task) had a duration of 500 ms, including 50-ms

raised-cosine onset and offset ramps, and was temporally centered in the carrier. The target

modulation was imposed with a 0-degree starting phase. The stimulus intervals were separated by 500

ms of silence. The stimuli for all three intervals were calibrated to the desired presentation level to

ensure that no level cues were available. The tracking variable was the target's modulation depth,

expressed on a dB scale as $M = 20log_{10}(m)$, where m is the modulation depth on a linear scale. The

step size was initially 4 dB, was reduced to 2 dB after the first lower reversal, and was finally reduced

to 1 dB after the second lower reversal.

a. AM detection thresholds

For the AM detection task, the starting modulation depth of the target was -5 dB. For any run, six

reversals were obtained using the smallest step size and the threshold was defined as the average target

modulation depth across them. The measurement at each target modulation frequency was repeated

three times, and the final threshold was defined as the average across repetitions. Additional

measurements were included if the standard error across repetitions exceeded 2 dB, until that limit

was met.[3] All listeners completed two training runs (at the target modulation frequencies of 4 and 128

Hz). Figure 2 shows an example of the stimulus envelope for the reference interval (i.e., the

unmodulated carrier, in panel A) and the target interval (i.e., the carrier modulated with the target

modulation, in panel B) for a target modulation frequency of 128 Hz.

b. Masked-threshold patterns

For the MTP task, the target and masker modulations were applied simultaneously to the carrier

using a multiplicative approach, as described in Regev *et al.* (2023). The masker modulation was a

narrow-band Gaussian noise with a fixed bandwidth corresponding to ½-octave when centered on

the target modulation frequency. For target modulation frequencies of 4 and 16 Hz, the masker

modulation was centered at frequencies ranging from -2 to +2 octaves relative to the target modulation

frequency, in steps of 2/3 octaves. In order to avoid cues from spectrally resolved modulation

sidebands, the highest masker-modulation center frequency was limited to 128 Hz, such that the

highest masker-modulation center frequencies corresponded to +2/3 and 0 octaves for the target

modulation frequencies of 64 and 128 Hz, respectively. For these target modulation frequencies, a

masked threshold was also measured for masker modulations centered around 4 Hz, corresponding

to -4 and -5 octaves relative to the target modulation frequencies, respectively. For these additional

conditions, the lower cutoff frequency of the masker modulation was set to 1 Hz, thus effectively

reducing the masker's bandwidth. The resulting masker-modulation bandwidths and center

frequencies for all conditions are summarized in Appendix B. For each measurement run, a 2-s long

segment of the masker modulation was created. For each interval, a 500-ms segment, randomly cut

out of the long masker modulation, was applied to the carrier. The root-mean square (rms) modulation

depth of the masker modulation was fixed at -10 dB.

---

[3] Overall, 8 measurements were added for 2 of the young HI listeners and 1 measurement was added for a single older HI listener.

The starting modulation depth of the target was 0 dB. For any run, eight reversals were obtained

using the smallest step size and the threshold was defined as the average target modulation depth

across them. The measurement at each masker-modulation center frequency was repeated three times,

and the final threshold was defined as the average across repetitions. To avoid overmodulation, the

maximum allowed target modulation depth was 0 dB, and a safety check was introduced during the

stimulus generation to minimize any potential residual overmodulation caused by the masker

modulation. If the procedure required a target modulation depth above 0 dB more than three times

over the course of a run, the run was stopped and an extra run was carried out to replace it. If this

happened more than 5 times for any condition, the threshold was considered as unmeasurable. This

was the case for a single older HI listener, whose thresholds could not be estimated for masker-

modulation center frequencies of +2/3 and +4/3 octaves relative to the 4 Hz target modulation

frequency. This listener's masked thresholds at 4 Hz were therefore not included in the statistical

analyses, or in the figures below, such that the MTP analyses at 4 Hz included only nine older HI

listeners. Additional measurements were included if the standard error across repetitions exceeded 2

dB, until that limit was met.[4] For each MTP, the listeners completed three training runs (at the masker-

modulation center frequencies of -2, 0, and +2 octaves relative to the target modulation frequency).

Figure 2 shows an example of the stimulus envelope for the reference interval (i.e., the carrier

modulated with the masker modulation, in panel C) and the target interval (i.e., the carrier modulated

with both the target and masker modulations, in panel D), for a target modulation frequency of 128

Hz and a masker-modulation center frequency of 64 Hz.

---

[4] Overall, a total of 2 measurements were added for a single young HI listener, and 7 measurements for 3 of the older HI listeners, due to aborted measurement runs. Five measurements were added for 2 of the young HI listeners and 10 measurements for 4 of the older HI listeners, due to excessive standard errors. Masked thresholds could not be estimated for a single older HI listener, at masker modulation center frequencies of +2/3 and +4/3 octaves relative to the 4 Hz target modulation frequency.

Figure 2: Examples of the temporal envelopes of the stimuli used in the AM detection and MTP experiments. Panel A: Reference interval in the AM detection task, represented by an unmodulated sinusoidal carrier (2.8 kHz). Panel B: Target interval in the AM detection experiment, containing the same carrier as in Panel A modulated by sinusoidal target modulation (128 Hz). Panel C: Reference interval in the MTP experiment, containing the same carrier as in Panel A modulated by narrowband-noise masker modulation (between 43.6 and 88.2 Hz, centered at 64 Hz). Panel D: Target interval in the MTP experiment, with the same carrier as in Panel A modulated by the sinusoidal target modulation and the narrowband-noise masker modulation.

2. Estimates of AM frequency selectivity

Ewert *et al.* (2002) and Regev *et al.* (2023) used the envelope power spectrum model of masking (EPSM; Ewert and Dau, 2000) to quantify the AM frequency selectivity reflected in their MTPs, by finding the Q-factor of the best-fitting bandpass modulation filter to the experimental data. However, high levels of asymmetry were observed in the MTPs in the present study (as described in detail below),

which are not consistent with the EPSM's assumption of symmetry (see section IV.D for a discussion

of this aspect). Hence, a simplified approach was used here to quantify AM frequency selectivity. The

best-fitting linear approximation (in terms of the least-squared error) was obtained separately for each

skirt using the group-average masked thresholds for the masker locations ranging from -2 to 0 and 0

to +2 octaves. Each linear fit was constrained to cross the on-target masked threshold. The bandwidth

of the asymmetrical modulation filter was then estimated by measuring the spectral distance between

the 3-dB-down points on the fits obtained for each skirt when interpolating across masker locations

ranging from -4 to +4 octaves. For the 128-Hz pattern, the high-frequency cutoff was estimated

assuming symmetry around the target modulation frequency. Finally, the metric quantifying AM

frequency selectivity was obtained as an overall 'Q-factor' by computing the ratio of the filter's

characteristic (modulation) frequency to the fitted bandwidth. The overall Q-factors describing the

young NH and older NH average MTPs, obtained by Regev *et al.* (2023), were also re-evaluated using

this linear approximation.

D. Statistical analysis

Analyses of variances (ANOVAs) of linear mixed-effects model fittings were used to assess the

differences between the results for the older HI listeners obtained in the present study and the data

for the NH listeners obtained by Regev *et al.* (2023). Each model and corresponding ANOVA were

applied twice. First, to estimate the effects of hearing loss in isolation from age effects, the results for

the older HI listeners were compared with the results for the older NH listeners. Second, using an

analysis comparable to that employed in previous studies, the results for the older HI listeners were

compared with the results obtained for the young NH listeners. For the analysis of the AM detection

thresholds, the listeners were considered as a random effect and group classification and modulation

frequency were considered as fixed effects. The MTPs were analyzed separately for each target

modulation frequency. The listeners were considered as a random effect and group classification and masker-modulation center frequency were considered as fixed effects. Levene's test was used to assess the homoskedasticity of the residuals and Shapiro-Wilk's test was used to assess the normality of the residual's distribution.[5] Post-hoc analyses were conducted using estimated marginal means and a Holm-Bonferroni correction criterion was applied for multiple comparisons. The level of statistical significance was set to 5% in all analyses. Due to the low number of young HI listeners (3), no statistical analysis was performed to compare their data to those of the other groups. This limitation is further discussed in section IV.D.

---

[5] Levene's test was not passed for one of the models applied, while Shapiro-Wilk's test was not passed for 3 of the other models. However, the deviations were minor, and the model predictions matched the average group data accurately. Hence, no transformations of the data were applied in order to keep all applied analyses consistent.

III.   RESULTS

A. AM detection thresholds

Figure 3 (Color online): Average AM detection thresholds for each listener group as a function of modulation frequency. The data for the young HI listeners are indicated by diamonds connected by solid lines. The data for the older HI listeners are shown as circles connected by solid lines. The results for the young and the older NH listeners (dashed lines; data from Regev et al., 2023) are shown as triangles and squares, respectively. Means and standard errors are shown. The stars indicate the level of statistical significance of the differences in thresholds between the older HI and older NH listeners (dark stars), and the older HI and young NH listeners (light stars; *$p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

Figure 3 shows the mean AM detection thresholds as a function of modulation frequency for the young HI (diamonds), older HI listeners (circles), young NH (squares), and older NH listeners (triangles), the latter two datasets from Regev *et al.* (2023). Substantial levels of individual variability were observed (see supplementary Figure 1). The statistical analysis comparing the thresholds for the

older HI and the older NH listeners revealed a significant main effect of group [$F_{1,18} = 21.21, p <$

0.001] and modulation frequency [$F_{3,54} = 22.84, p < 0.001$]. The post-hoc analysis revealed that

the older HI listeners had significantly lower (i.e., better) AM detection thresholds than the older NH

listeners at all modulation frequencies [effect size (expressed as difference in M; ΔM) $= -6.0$ dB,

standard error (SE) $= 2.0$ dB; $\Delta M = -8.5$ dB, $SE = 2.0$ dB; $\Delta M = -7.4$ dB, $SE = 2.0$ dB; $\Delta M =$

-7.0 dB, $SE = 2.0$ dB, for 4, 16, 64, and 128 Hz, respectively].

The statistical analysis comparing the thresholds for the older HI and the young NH listeners

revealed a significant main effect of group [$F_{1,19} = 8.13, p = 0.01$] and modulation frequency

[$F_{3,57} = 14.66, p < 0.001$], and a significant interaction [$F_{3,57} = 5.26, p = 0.003$]. The post-hoc

analysis revealed that the older HI listeners had significantly lower AM detection thresholds than the

young NH listeners for modulation frequencies of 16 Hz [$\Delta M = -7.6$ dB, $SE = 1.8$ dB] and 64 Hz

[$\Delta M = -4.9$ dB, $SE = 1.8$ dB].

297   **B. Masked-threshold patterns (MTPs)**

Figure 4 (Color online): Average MTPs (filled symbols) and unmasked AM detection thresholds (open symbols, replotted from Figure 3) for the older HI and the older NH listeners. The data for the HI listeners are indicated by the circles connected by solid lines and the data for the NH listeners (from Regev *et al.*, 2023) are shown as triangles connected by dashed lines. Each panel represents the results for one target modulation frequency: 4 Hz (top left), 16 Hz (top right), 64 Hz (bottom left), and 128 Hz (bottom right). The data from one of the older HI listeners were excluded from the 4-Hz panel for both the MTP and AM detection threshold. Means and standard errors are shown. Stars indicate the level of statistical significance of the differences in masked thresholds between the two groups (* *p < 0.05, ** p < 0.01, *** p < 0.001*).

Figure 4 shows the mean MTPs for the older HI (circles) and the older NH listeners (triangles;

data from Regev *et al.*, 2023). Each panel shows the MTPs for a specific target modulation frequency.

The mean AM detection thresholds are indicated by the open symbols (replotted from Figure 3). The

statistical analysis comparing the older HI and the older NH listeners revealed a significant main effect

of group for all target modulation frequencies [$F_{1,17} = 7.31, p = 0.015$; $F_{1,18} = 20.18, p < 0.001$;

$F_{1,18} = 8.70, p = 0.009$; $F_{1,18} = 14.43, p = 0.001$, for 4, 16, 64, and 128 Hz, respectively]. A

significant interaction between listener group and masker-modulation center frequency was found for

the target modulation frequencies of 4 Hz [$F_{6,102} = 3.36, p = 0.005$] and 16 Hz [$F_{6,108} = 4.64, p <$

0.001], whereas the interaction approached, but did not reach, significance for the target modulation

frequency of 64 Hz [$F_{5,90} = 2.01, p = 0.085$]. For the target modulation frequency of 4 Hz, the post-

hoc analysis revealed that the older HI listeners had significantly lower (better) masked thresholds

than the older NH listeners for the masker modulations centered at -2, -4/3, and -2/3 octaves relative

to the target frequency. At 16 Hz, the post-hoc analysis revealed that the older HI listeners had

significantly lower masked thresholds than the older NH listeners for the masker modulations

centered at -2, -4/3, and +2 octaves relative to the target frequency. At 64 Hz, the post-hoc analysis

revealed that the older HI listeners had significantly lower masked thresholds than the older NH

listeners for the masker modulations centered at -4, -2, and +2/3 octaves relative to the target

frequency. Finally, at 128 Hz, the post-hoc analysis revealed that the older HI listeners had significantly

lower masked thresholds than the older NH listeners for the masker modulations centered at -5, -2, -

4/3, and 0 octaves relative to the target frequency. The details of the post-hoc analyses can be found

in Appendix B. Overall, the MTPs were sharper for the older HI listeners than for the older NH

listeners, for the target modulation frequencies of 4 Hz (on the low-frequency skirt), 16 Hz, and 64

Hz. Conversely, the high-frequency skirts of the 4-Hz MTPs suggests similar selectivity for the older

HI and NH listeners. Finally, the MTP at 128 Hz reflects a general improvement (i.e., decrease) of the

masked thresholds for the older HI listeners, but an overall broader pattern, compared to that for the

older NH listeners, as shown by the fitted Q-factors displayed in Table 1 and discussed further below.

High levels of asymmetry were also observed for the older HI listener's MTPs, as quantified by the

slopes of the obtained linear fits (see supplementary Figure 2). At the target modulation frequency of

4 Hz, this was reflected by a steeper slope of the low-frequency skirt (3 dB/octave) than the high-

frequency skirt (1 dB/octave) and an off-target peak threshold (for the masker centered at +2/3

octaves relative to the target frequency). A similar effect was found at 16 Hz, with a steeper slope of

the low-frequency skirt (5.6 dB/octave) than of the high-frequency skirt (4.6 dB/octave). At 64 Hz,

the older HI listeners gave a much shallower slope of the low-frequency skirt (3.6 dB/octave) than of

the high-frequency skirt (9.3 dB/octave), although the latter was computed based only on the average

masked threshold at +2/3 octaves.

Figure 5 (Color online): Average MTPs (filled symbols) and unmasked AM detection thresholds (open symbols, replotted from Figure 3). The data for the young and older HI listeners (solid lines) are indicated by the diamonds and circles, respectively. The data for the young NH listeners are shown as dashed lines and squares (data from Regev *et al.*, 2023). Each panel represents the results for one target modulation frequency: 4 Hz (top left), 16 Hz (top right), 64 Hz (bottom left), and 128 Hz (bottom right). The data from one of the older HI listeners were excluded from the 4-Hz panel for both the MTP and AM detection thresholds. Means and standard errors are shown. Stars indicate the level of statistical significance of the differences in masked thresholds between the young NH and older HI listeners (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Figure 5 shows the mean MTPs for the young HI (diamonds) and older HI listeners (circles) and the data from the young NH listeners (squares) obtained by Regev *et al.* (2023). Each panel shows the

MTPs for a specific target modulation frequency. The mean AM detection thresholds are also shown

and indicated by the open symbols (replotted from Figure 3). The statistical analysis comparing the

older HI and the young NH listeners showed no main effect of group for any target modulation

frequency. However, there was a significant interaction between listener group and masker-

modulation center frequency for all target modulation frequencies [$F_{6,108} = 3.95, p = 0.001$;

$F_{6,114} = 8.1, p < 0.001$; $F_{5,95} = 4.12, p = 0.002$; $F_{4,76} = 6.3, p < 0.001$, for 4, 16, 64, and 128

Hz, respectively]. For the target modulation frequency of 4 Hz, the post-hoc analysis showed that the

older HI listeners had significantly higher (worse) masked thresholds than the young NH listeners for

the masker centered at +2 octaves relative to the target. At 16 Hz, the older HI listeners had

significantly lower (i.e., better) masked thresholds for the masker centered at -2 octaves relative to the

target frequency and significantly higher masked thresholds than the young NH listeners for the

masker centered at +2/3 octaves above the target frequency. At 64 Hz, the older HI listeners had

significantly lower masked thresholds than the young NH listeners for the masker centered at +2/3

octaves above the target frequency. Finally, at 128 Hz, the older HI listeners had significantly lower

masked thresholds than the young NH listeners for the masker centered at the target modulation

frequency. Details of the post-hoc analyses can be found in Appendix B.

Although no statistical analysis including the data for the young HI listeners was performed, due to

the low number of participants, their results provide insights into the effects of hearing loss in the

absence of ageing. Similar to the MTPs for the older HI listeners, high levels of asymmetry were

observed in the MTPs of the young HI listeners. At the target modulation frequency of 4 Hz, this was

reflected by a steeper slope of the low-frequency skirt (4.8 dB/octave) than the high-frequency skirt

(3.3 dB/octave). At 16 Hz, the low-frequency skirt (6 dB/octave) was again steeper than the high-

frequency skirt (5 dB/octave). At 64 Hz, the young HI listeners displayed a much shallower low-

frequency skirt (5.6 dB/octave) than high-frequency skirt (10.6 dB/octave), although the latter was

computed based only on the average masked threshold at +2/3 octaves.

Table 1 summarizes the overall Q-factors fitted to the average data for each listener group and

shows the mean square errors (MSEs, in dB) of the fit in parentheses. The table also shows the ratio

of the Q-factors for each group to the Q-factor for the young NH listeners. A ratio higher than 1

indicates sharper tuning for the listener group in question relative to the young NH listeners. For the

older NH listeners, the reduced Q-factor ratio reflects the age-related reduction of AM frequency

selectivity shown by Regev *et al.* (2023). In contrast, the results for the older HI listeners suggest

sharper tuning for this group than for the older NH listeners for the modulation frequencies of 4, 16,

and 64 Hz, and reduced AM frequency selectivity at 128 Hz. Compared to the young NH listeners,

the older HI listeners show similar, or slightly greater, AM frequency selectivity for the modulation

frequencies of 16 and 64 Hz, and broader tuning at 4 and 128 Hz. The young HI listeners' results

suggest generally greater AM frequency selectivity than for the older HI listeners, and even higher Q-

factors (i.e., sharper tuning) than the young NH listeners for the modulation frequencies of 4, 16, and

64 Hz.

Table 1: Overall fitted Q-factors, defined as the ratio of the filter's characteristic frequency to its

bandwidth estimated at the 3-dB-down point, and MSEs for the modulation filters fitted to the average

group data using an asymmetric linear fitting approach. The table also shows the ratio of Q-factors

for each group to the Q-factor for the young NH listeners.

Target modulation frequency [Hz]	Estimated Q-factor *(Overall MSE [dB])*				Q/Q_{YNH}		
	YNH	**ONH**	**YHI**	**OHI**	**ONH**	**YHI**	**OHI**
4	0.77 *(0.35)*	0.10 *(0.16)*	0.81 *(0.78)*	0.14 *(0.30)*	0.13	1.05	0.18
16	0.96 *(0.13)*	0.75 *(0.44)*	1.24 *(0.38)*	1.13 *(0.30)*	0.78	1.29	1.18
64	1.38 *(0.15)*	1.14 *(0.09)*	1.90 *(1.0)*	1.45 *(0.14)*	0.83	1.38	1.05
128	0.96 *(0.05)*	0.40 *(0.10)*	0.82 *(0.98)*	0.27 *(0.96)*	0.42	0.85	0.28

IV. DISCUSSION

A. Summary of main findings

This study aimed to investigate the potentially opposite effects of hearing loss and age on AM frequency selectivity. These opposing effects were suggested based on recent findings of an age-related reduction in AM frequency selectivity (Regev et al., 2023), which appeared to contradict previous conclusions that neither age nor hearing loss affect AM frequency selectivity (Takahashi and Bacon, 1992; Sek et al., 2015). Behavioral MTPs and unmasked AM detection thresholds were collected from three young and ten older HI listeners (24-30 and 63-77 years). The results were compared to the data obtained from young and older NH listeners by Regev et al. (2023), publicly available from Regev et al. (2024). Both HI groups showed lower AM detection thresholds than their similarly aged NH counterparts, and the older HI listeners also showed lower detection thresholds than the young NH listeners for target modulation frequencies above 4 Hz (Figure 3). Furthermore, for target modulation frequencies below 128 Hz, the older HI listeners generally showed greater AM frequency selectivity than the older NH listeners (Figure 4). When compared to the young NH listeners, the older HI listeners displayed an asymmetric broadening of the MTP at the target modulation frequency of 4 Hz (with a shallower high-frequency skirt), but similar or even slightly sharper tuning at 16 and 64 Hz (Figure 5). At the target modulation frequency of 128 Hz, the older HI listeners showed lower masked thresholds overall than the older NH listeners (with an average offset of 3.4 dB) and similar masked thresholds to the young NH listeners for the off-target maskers, effectively indicating reduced AM frequency selectivity compared to both NH groups. The data from the young HI listeners suggested greater AM frequency selectivity than the older HI listeners, and generally sharper tuning than the young NH listeners (as can be seen in Figure 5). Notably, both the young and older HI listeners showed high levels of asymmetry in their MTPs, with generally steeper low-frequency than high-frequency skirts. Finally, the best-fitting Q-factor supported previous observations suggesting that AM

frequency selectivity for HI listeners is similar to, or greater than, that for similarly aged NH listeners,

except for the highest target modulation rate of 128 Hz (Table 1).

B. Opposite effects of hearing loss and age

The results for both the AM detection and MTP tasks suggest a general perceptual benefit from

hearing loss, in terms of increased detectability of the target modulation and a sharpening of the

masked patterns. This benefit effectively counteracts the detrimental effects of age on AM detection

and AM frequency selectivity. The improvement of AM detection thresholds for HI listeners

compared to NH listeners is consistent with previous studies (e.g., Bacon and Gleitman, 1992; Moore

and Glasberg, 2001; Sek *et al.*, 2015; Schlittenlacher and Moore, 2016; Wallaert *et al.*, 2017; Wiinberg

et al., 2019). It has been suggested that this improvement is due to an effectively increased internal

representation of the signal's AM depth resulting from the loss of compression commonly associated

with SNHL (Moore *et al.*, 1996; Jennings *et al.*, 2018). The trends observed for the young HI listeners

support the notion that greater age is detrimental to AM detection whereas SNHL provides a benefit.

Previous AM masking studies comparing young NH and older HI listeners using low target

modulation frequencies ($\leq$ 16 Hz) found no differences in MTPs (Takahashi and Bacon, 1992; Sek *et*

al., 2015) and concluded that neither SNHL nor age affect AM frequency selectivity. The present

results are broadly consistent with the observation that MTPs are similar for young NH and older HI

listeners for low modulation frequencies, although an asymmetric broadening of the 4-Hz MTP was

found for the older HI listeners, with a shallower high-frequency skirt. However, the data also showed

that SNHL generally improves AM frequency selectivity for listeners of similar age for target

modulation frequencies below 128 Hz. In contrast, at 128 Hz, the masked thresholds were lower for

the older HI listeners than for the older NH listeners and effectively indicated a reduction in AM

frequency selectivity for the older HI listeners. Additionally, a slight broadening of the pattern at 128

Hz was observed for young HI listeners compared to the young NH listeners.

The results of the present study show that young HI listeners generally perform better than older

HI listeners, as shown by Regev *et al.* (2023) for NH listeners. Furthermore, although a sharpening of

the MTPs for HI listeners was not observed on both skirts of the patterns, HI listeners generally

outperformed similarly aged NH listeners in terms of AM detection thresholds and tuning of the

MTPs. Factors that may have influenced these results, such as the difference in presentation levels

across the NH and HI listeners, are discussed in section IV.D.

C. Potential mechanism underlying the perceptual benefit of hearing loss

The underlying auditory processes that account for the sharper MTPs for the older HI listeners

than for the older NH listeners may not be straightforward. It seems unlikely that SNHL would

improve AM frequency selectivity by 'restoring' the bandwidth of the hypothetical modulation filters.

Instead, the greater AM frequency selectivity observed here for the HI listeners may result from the

reduced peripheral compression that is typically associated with SNHL. One notable characteristic of

the data for the HI listeners is the strong asymmetry of the MTPs, with a steeper slope on the low-

frequency than the high-frequency skirt. This was also found by Regev *et al.* (2023) for their NH

listeners, although the effect was less prominent than in the present study. The asymmetry suggests

that an additional cue for the detection of the target may be available when the modulation maskers

are centered below - but not above - the target modulation frequency. The existence of such a cue was

discussed in previous studies by Bacon and Grantham (1989) and Strickland and Viemeister (1996),

who found negative masking when an AM masker was centered below the target modulation

frequency. They argued that listeners may be able to detect the target modulation in the temporal dips

of the masker modulation, or 'listen in the dips', which is a detection cue analogous to the release from

masking provided by envelope fluctuations in spectral masking tasks (e.g., Buus, 1985). In the present

study, the random dips occurring in the modulation masker would be longer than (or close to) one

cycle of the target for maskers centered below the target, while this would not be the case for

modulation maskers centered above the target. It is possible that these local temporal features aid the

detection of the target in the presence of low-frequency maskers. The data suggest that such a cue, if

present, may be more salient for HI listeners than for NH listeners. The loss of cochlear compression

for HI listeners might increase the target's AM depth glimpsed in the temporal valleys of the masker,

further enhancing the detectability of the target. This notion is consistent with the findings of Wallaert

et al. (2017), which showed a greater improvement in AM detection thresholds for HI listeners than

for NH listeners when increasing the numbers of cycles of the target modulation, suggesting enhanced

temporal integration for the HI listeners. This may imply that HI listeners require fewer cycles of the

target modulation to detect its presence, potentially contributing to the improved performance

measured on the lower skirt of the MTPs.

If this detection mechanism indeed mediated the sharper MTPs for HI listeners, investigating AM

frequency selectivity using an AM forward-masking paradigm should eliminate the perceptual benefit

of SNHL. AM forward masking has been used as an alternative approach for investigating AM

frequency selectivity (Wojtczak and Viemeister, 2005; Moore *et al.*, 2009; Füllgrabe *et al.*, 2021a, 2021b)

because it avoids extraneous cues resulting from the interaction of the target and masker modulation.

While it remains unclear whether AM simultaneous and AM forward-masking paradigms reveal the

same underlying AM frequency-selective process, a study comparing age-matched NH and HI

listeners using such a paradigm could provide valuable insights into the origins of the observed

asymmetry. However, such a study has not been conducted thus far. Additionally, computational

auditory models could be employed to assess the plausibility of 'listening in the dips' as a mechanism

underlying the present results.

The hypothesis that increased sensitivity to the temporal envelope, resulting from the loss of

cochlear compression, offers a perceptual advantage for HI listeners is supported by the findings of

Bianchi *et al.* (2016). They demonstrated that fundamental-frequency discrimination of unresolved

complex tones was comparable for older HI and young NH listeners when the reduced cochlear

compression resulted in an effective amplification of the modulation power of the stimulus (i.e., when

the envelope was maximally peaky), while performance was worse for the older HI listeners when the

envelope was much flatter and hence largely unaffected. Partially connected with the loss of cochlear

compression, the better-than-normal AM detection thresholds and AM frequency selectivity observed

here for the HI listeners may be related to the increased neural coding of the envelope which has been

shown both in animal models and humans (Kale and Heinz, 2010, 2012; Henry *et al.*, 2014; Zhong *et*

al., 2014; Millman *et al.*, 2017; Goossens *et al.*, 2018; Decruy *et al.*, 2020). This increased internal

representation of the envelope, while it may reflect a listening advantage for simple AM detection and

masking paradigms such as the ones considered in the present study, has been suggested ultimately to

be detrimental to the intelligibility of speech in noise (Millman *et al.*, 2017; Goossens *et al.*, 2018;

Decruy *et al.*, 2020). Hence, it could potentially contribute to the difficulties experienced by HI listeners

in complex listening scenarios. Similar conclusions were drawn by studies reporting a connection

between poorer speech intelligibility and increased neural coding of the envelope in older NH listeners

(e.g., Goossens *et al.*, 2018; Decruy *et al.*, 2019).

D. Limitations of the study

The presentation levels used in the present study were not directly comparable to those employed

by Regev *et al.* (2023) with NH listeners. Regev *et al.* used a fixed level of 65 dB SPL (corresponding

to average sensation levels of 67 and 57 dB SL for the young and older NH listeners, respectively),

while the present study adjusted the level for the HI listeners to be 30 dB SL (corresponding to 79 dB

SPL on average), following a similar approach to Sek *et al.* (2015). This methodological difference

implies that the presentation level was both lower in terms of SL (by 27-37 dB) and higher in terms

of SPL (by 14 dB) for the HI listeners than for the NH listeners. Previous studies have highlighted

the effect of the interaction between presentation level and hearing status on AM detection thresholds,

often finding lower AM detection thresholds for HI listeners than for NH listeners at equal SL, but

similar thresholds at equal SPL (Bacon and Gleitman, 1992; Moore and Glasberg, 2001; Füllgrabe *et*

al., 2003; Schlittenlacher and Moore, 2016; Wallaert *et al.*, 2017; Moore *et al.*, 2019), although some

studies still found differences at equal SPL (Sek *et al.*, 2015; Jennings *et al.*, 2018). Hence, it is possible

that the present results were influenced by the difference in presentation levels between the NH and

HI listeners. However, the present finding that the HI listeners showed lower AM detection thresholds

than the NH listeners at both lower SLs and higher SPLs seems consistent with previous studies that

reported similar results (Füllgrabe *et al.*, 2003; Jennings *et al.*, 2018). Füllgrabe *et al.* (2003) observed an

increased difference in AM detection thresholds between NH and HI listeners when the experiment

was repeated with a fixed SL for all participants, relative to the condition where the HI listeners were

tested with a higher SPL and lower SL than the NH listeners. Additionally, Wojtczak (2011)

investigated the effects of carrier level on AM frequency selectivity for NH listeners and found that a

reduction in SL led to an increase in AM detection thresholds and a broadening of the MTPs.

Therefore, it is likely that testing NH and HI listeners at equal SL would have further increased the

observed differences between the two groups in the present study. However, it remains unclear

whether lower AM detection thresholds and sharper MTPs for the HI listeners than for the NH

listeners would still be found if they were tested at equal SPL.

It is plausible that the underlying pathology of the hearing loss was different for the young and

older HI listeners, potentially affecting the effects on sound perception. Although all HI listeners had

a SNHL, the origin of the hearing loss in the young HI listeners was unfortunately unknown.

Additionally, the small number of young HI listeners included in this study, along with the significant

variability in their audiograms (except for the frequency of interest), limited the conclusions that could

be drawn from their data. In spite of these aspects, the results for the young HI listeners in this study

may offer some valuable insights. Indeed, the data obtained across the three young HI listeners were

consistent with the observations made based on the other listener groups. Specifically, young listeners

exhibited sharper tuning than older listeners with the same hearing status, and HI listeners generally

demonstrated sharper tuning than similarly aged NH listeners. Conducting further investigations with

a larger sample of young HI listeners may help validate the trends identified in the present study.

Finally, the method used to quantify AM frequency selectivity in the present study was limited by

the pronounced asymmetry of the MTPs. Previous studies quantified the tuning of the hypothetical

modulation filters using the EPSM (Ewert and Dau, 2000; Ewert *et al.*, 2002; Regev *et al.*, 2023), which

assumes symmetrical filters centered around the characteristic modulation frequency. However, the

present results do not support this assumption, indicating a need to revise the EPSM to incorporate

more flexible filter shapes. Similar conclusions were drawn by Regev *et al.* (2023), who observed large

variability in individual MTPs. Consequently, the Q-factor values reported in this study for the young

and older NH listeners differ from those reported by Regev *et al.* (2023) due to the different

approaches used to quantify AM frequency selectivity. However, the overall finding of lower Q factors

for the older NH listeners than for the young NH listeners is consistent between the two

investigations.

V.   CONCLUSIONS

This study investigated the effects of hearing loss on AM frequency selectivity by employing a

simultaneous AM masking task. Ten older and three young HI listeners were tested, ranging in age

from 63-77 and 24-30 years, respectively. These age ranges were approximately matched to those of

the young and older NH listeners tested by Regev *et al.* (2023). Although no statistical analysis

including the young HI data could be conducted, due to the limited number of young HI listeners,

their results provide valuable insight into the effects of hearing loss in the absence of ageing. The HI

listeners showed lower AM detection thresholds and generally sharper MTPs than the NH listeners,

suggesting that hearing loss may counteract the detrimental effects of age on AM frequency selectivity.

The MTPs showed strong asymmetry, with generally steeper low-frequency skirts. A potential

mechanism for explaining these results is the detection of the target modulation in the dips of low-

frequency masker modulations, coupled with the effective increase in AM depth resulting from the

loss of cochlear compression. This suggests that the observed sharpening of the MTPs for HI listeners

may be linked to the loss of cochlear compression rather than a 'restoration' of the bandwidth of the

underlying hypothetical modulation filters. Differences in presentation level between the HI and NH

listeners may have influenced the threshold differences, so the results of this study motivate further

investigations into the effects of hearing loss on AM masking and AM frequency selectivity, as well as

their interaction with age-related effects. In addition to the loss of cochlear compression, the lower

AM detection thresholds and greater AM frequency selectivity observed in HI listeners may be

associated with increased neural coding of the envelope, which has been suggested to be detrimental

to speech intelligibility in complex listening scenarios. Hence, future studies examining the effects of

hearing loss on AM masking can offer valuable insights into the underlying mechanisms that

contribute to the perceptual benefits observed for HI listeners and their implications for complex

auditory perception and speech intelligibility.

ACKNOWLEDGMENTS

The authors thank Borgný Súsonnudóttir Hansen for her contribution to the data collection, as well as Erik Kjærbøl and Jesper Borchorst Yde for their precious help in recruiting young HI participants. We also thank Stephan Ewert for helpful suggestions and for kindly sharing experimental details and data from earlier publications, Jens Hjortkjær and Jonatan Märcher-Rørsted for kindly providing their implementations of the reverse digit span test and of the Cambridge formula (CamEQ; Moore and Glasberg, 1998), as well as Raul Sanchez-Lopez and Beverly Wright for helpful insights and discussions. We are thankful to Brian C. J. Moore and two other anonymous reviewers for their insightful comments during the review process. This study was supported by the Oticon Centre of Excellence for Hearing and Speech Sciences (CHeSS).

AUTHOR DECLARATIONS

Conflict of Interest

The authors have no conflict of interest to report.

Ethics Approval

All participants gave written informed consent and were financially compensated for their time. Ethical approval for the study was provided by the Science Ethics Committee of the Capital Region in Denmark (reference H-16036391).

DATA AVAILABILITY

The data that support the findings of this study are openly available in the repository "Dataset for: 'Disentangling the effects of hearing loss and age on amplitude modulation frequency selectivity'" in DTU Data, at http://doi.org/10.11583/DTU.25134611.

APPENDIX

A. Experimental protocol

Table 2 summarizes the experimental protocol used for each HI listener (both young and older).

Modulation rate discrimination thresholds were also collected but were not considered in the present

study.

Table 2: Experimental protocol implemented for all HI listeners, including the tests of hearing

threshold at the carrier frequency (HT), modulation rate discrimination (MRD), masked threshold

pattern (MTP), and reverse digit span (RDS).

Session	*Test*	*Order*
1	HT followed by MRD	Fixed within session
2-5	MTP 4 Hz	Random across sessions
	MTP 16 Hz	
	MTP 64 Hz	
	MTP 128 Hz followed by AM detection	
Any (2-5)	RDS	Randomly added to any session

B. Parameters and post-hoc analyses for masked-threshold patterns

Table 3 summarizes the post-hoc analyses of the ANOVA comparing the MTPs for the older HI and NH listeners (Figure 4).

Table 3: Summary of post-hoc analyses of the MTPs comparing older HI and older NH listeners, for each target modulation frequency. The effect sizes (ΔM) and standard errors (*SEs*) are reported for each pairwise comparison. Significant p-values are highlighted in bold.

Target frequency f_m [Hz]	*Effective masker-modulation bandwidth [Hz]*	*Masker-modulation center frequencies [Hz]*	*Masker-modulation position [octaves re target f_m]*	*ΔM [dB]*	*SE [dB]*	*Corrected p-value*
4	1.4	1	-2	-4.6	1.1	**< 0.001**
		1.6	-4/3	-3.7		**0.001**
		2.5	-2/3	-2.2		**0.047**
		4	0	-1.3		0.237
		6.3	2/3	-0.8		0.472
		10.1	4/3	-1.6		0.147
		16	2	-1.7		0.134
16	5.6	4	-2	-4.5	0.8	**< 0.001**
		6.3	-4/3	-2.3		**0.003**
		10.1	-2/3	-1		0.196

		16	0	0.1		0.935
		25.4	2/3	-1		0.2
		40.3	4/3	-1		0.179
		64	2	-2.9		**< 0.001**
64	15.1	4	-4	-4.6	1.2	**< 0.001**
		16	-2	-3		**0.014**
	22.3	25.4	-4/3	-2.2		0.061
		40.3	-2/3	-1.7		0.152
		64	0	-1.4		0.249
		101.6	2/3	-3		**0.013**
128	27.2	4	-5	-4.9	1.2	**< 0.001**
		32	-2	-3.5		**0.005**
	44.6	50.8	-4/3	-2.7		**0.027**
		80.6	-2/3	-2.2		0.074
		128	0	-3.8		**0.003**

635   Table 4 summarizes the post-hoc analyses of the ANOVA comparing the MTPs for the older HI

636   and young NH listeners (Figure 5).

638   Table 4: Summary of post-hoc analyses of the MTPs comparing older HI and young NH listeners,

639   for each target modulation frequency. The effect sizes (ΔM) and standard errors (SEs) are reported

640   for each pairwise comparison. The significant p-values are highlighted in bold.

Target frequency f_m [Hz]	Effective masker-modulation bandwidth [Hz]	Masker-modulation center frequencies [Hz]	Masker-modulation position [octaves re target f_m]	ΔM [dB]	SE [dB]	Corrected p-value
4	1.4	1	-2	-0.2	1.1	0.873
		1.6	-4/3	-0.5		0.662
		2.5	-2/3	-0.4		0.747
		4	0	-1.4		0.24
		6.3	2/3	0.4		0.736
		10.1	4/3	1.7		0.15
		16	2	3.2		**0.008**
16	5.6	4	-2	-3.3	0.8	**< 0.001**
		6.3	-4/3	-1.4		0.081
		10.1	-2/3	-0.6		0.405
		16	0	0.7		0.371
		25.4	2/3	2		**0.013**
		40.3	4/3	1.1		0.16
		64	2	-0.8		0.304
64	15.1	4	-4	-1.7	1.1	0.126

		16	-2	-0.5		0.637
		25.4	-4/3	0.6		0.597
	22.3	40.3	-2/3	0.5		0.648
		64	0	-1.5		0.191
		101.6	2/3	-3.1		**0.008**
	27.2	4	-5	-0.8		0.434
		32	-2	0.7		0.542
128		50.8	-4/3	0.6	1.1	0.594
	44.6	80.6	-2/3	-1.1		0.302
		128	0	-3.9		**< 0.001**

REFERENCES

Bacon, S. P., and Gleitman, R. M. (**1992**). "Modulation detection in subjects with relatively flat hearing losses," J Speech Lang Hear Res, **35**, 642–653. doi:10.1044/jshr.3503.642

Bacon, S. P., and Grantham, D. W. (**1989**). "Modulation masking: Effects of modulation frequency, depth, and phase," J Acoust Soc Am, **85**, 2575–2580. doi:10.1121/1.397751

Bianchi, F., Fereczkowski, M., Zaar, J., Santurette, S., and Dau, T. (**2016**). "Complex-tone pitch discrimination in listeners with sensorineural hearing loss," Trends Hear, **20**, 1–15. doi:10.1177/2331216516655793

Bisgaard, N., Vlaming, M. S. M. G., and Dahlquist, M. (**2010**). "Standard audiograms for the IEC 60118-15 measurement procedure," Trends Amplif, **14**, 113–120. doi:10.1177/1084713810379609

654   Buus, S. (**1985**). "Release from masking caused by envelope fluctuations," J Acoust Soc Am, **78**,

655   1958–1965. doi:10.1121/1.392652

656   Chabot-Leclerc, A., Jørgensen, S., and Dau, T. (**2014**). "The role of auditory spectro-temporal

657   modulation filtering and the decision metric for speech intelligibility prediction," J

658   Acoust Soc Am, **135**, 3502–3512. doi:10.1121/1.4873517

659   Chi, T., Ru, P., and Shamma, S. A. (**2005**). "Multiresolution spectrotemporal analysis of complex

660   sounds," J Acoust Soc Am, **118**, 887–906. doi:10.1121/1.1945807

661   Christiansen, S. K., Jepsen, M. L., and Dau, T. (**2014**). "Effects of tonotopicity, adaptation,

662   modulation tuning, and temporal coherence in 'primitive' auditory stream segregation,"

663   J Acoust Soc Am, **135**, 323–333. doi:10.1121/1.4845675

664   Conroy, C., Buss, E., and Kidd, G. (**2023**). "Cues to reduce modulation informational masking,"

665   J Acoust Soc Am, **153**, 274–285. doi:10.1121/10.0016867

666   Conroy, C., Byrne, A. J., and Kidd, G. (**2022**). "Forward masking of spectrotemporal modulation

667   detection," J Acoust Soc Am, **151**, 1181–1190. doi:10.1121/10.0009404

668   Dau, T., Kollmeier, B., and Kohlrausch, A. (**1997a**). "Modeling auditory processing of amplitude

669   modulation. I. Detection and masking with narrow-band carriers," J Acoust Soc Am, **102**,

670   2892–2905. doi:10.1121/1.420344

671   Dau, T., Kollmeier, B., and Kohlrausch, A. (**1997b**). "Modeling auditory processing of amplitude

672   modulation. II. Spectral and temporal integration," J Acoust Soc Am, **102**, 2906–2919.

673   doi:10.1121/1.420345

674   Dau, T., Verhey, J., and Kohlrausch, A. (**1999**). "Intrinsic envelope fluctuations and modulation-

675   detection thresholds for narrow-band noise carriers," J Acoust Soc Am, **106**, 2752–2760.

676   doi:10.1121/1.428103

Decruy, L., Vanthornhout, J., and Francart, T. (**2019**). "Evidence for enhanced neural tracking of the speech envelope underlying age-related speech-in-noise difficulties," J Neurophysiol, **122**, 601–615. doi:10.1152/jn.00687.2018

Decruy, L., Vanthornhout, J., and Francart, T. (**2020**). "Hearing impairment is associated with enhanced neural tracking of the speech envelope," Hear Res, **393**, 107961. doi:10.1016/j.heares.2020.107961

Depireux, D. A., Simon, J. Z., Klein, D. J., and Shamma, S. A. (**2001**). "Spectro-temporal response field characterization with dynamic ripples in ferret primary auditory cortex," J Neurophysiol, **85**, 1220–1234. doi:10.1152/jn.2001.85.3.1220

Elhilali, M., Chi, T., and Shamma, S. A. (**2003**). "A spectro-temporal modulation index (STMI) for assessment of speech intelligibility," Speech Commun, **41**, 331–348. doi:10.1016/S0167-6393(02)00134-6

Elhilali, M., Ma, L., Micheyl, C., Oxenham, A. J., and Shamma, S. A. (**2009**). "Temporal coherence in the perceptual organization and cortical representation of auditory scenes," Neuron, **61**, 317–329. doi:10.1016/j.neuron.2008.12.005

Elliott, T. M., and Theunissen, F. E. (**2009**). "The modulation transfer function for speech intelligibility," PLoS Comput Biol, **5**, e1000302. doi:10.1371/journal.pcbi.1000302

Erb, J., Schmitt, L.-M., and Obleser, J. (**2020**). "Temporal selectivity declines in the aging human auditory cortex," Elife, **9**, 1–21. doi:10.7554/eLife.55300

Ewert, S. D. (**2013**). "AFC - A modular framework for running psychoacoustic experiments and computational perception models," Proceedings of the International Conference on Acoustics AIA-DAGA 2013, Merano, Italy, 1326–1329.

Ewert, S. D., and Dau, T. (**2000**). "Characterizing frequency selectivity for envelope fluctuations," J Acoust Soc Am, **108**, 1181–1196. doi:10.1121/1.1288665

Ewert, S. D., Verhey, J. L., and Dau, T. (**2002**). "Spectro-temporal processing in the envelope-

frequency domain," J Acoust Soc Am, **112**, 2921–2931. doi:10.1121/1.1515735

Fuglsang, S. A., Märcher-Rørsted, J., Dau, T., and Hjortkjær, J. (**2020**). "Effects of sensorineural

hearing loss on cortical synchronization to competing speech during selective attention,"

J Neurosci, **40**, 2562–2572. doi:10.1523/JNEUROSCI.1936-19.2020

Füllgrabe, C., Meyer, B., and Lorenzi, C. (**2003**). "Effect of cochlear damage on the detection of

complex temporal envelopes," Hear Res, **178**, 35–43. doi:10.1016/S0378-5955(03)00027-

3

Füllgrabe, C., Moore, B. C. J., and Stone, M. A. (**2015**). "Age-group differences in speech

identification despite matched audiometrically normal hearing: contributions from

auditory temporal processing and cognition," Front Aging Neurosci, **6**, 1–25.

doi:10.3389/fnagi.2014.00347

Füllgrabe, C., Sęk, A., and Moore, B. C. J. (**2021a**). "Frequency selectivity in the modulation

domain estimated using forward masking: Effects of masker modulation depth and

masker-signal delay," Hear Res, **405**, 108244. doi:10.1016/j.heares.2021.108244

Füllgrabe, C., Sęk, A., and Moore, B. C. J. (**2021b**). "Forward masking of amplitude modulation

across ears and its tuning in the modulation domain," J Acoust Soc Am, **149**, 1764–1771.

doi:10.1121/10.0003598

Goossens, T., Vercammen, C., Wouters, J., and van Wieringen, A. (**2018**). "Neural envelope

encoding predicts speech perception performance for normal-hearing and hearing-

impaired adults," Hear Res, **370**, 189–200. doi:10.1016/j.heares.2018.07.012

He, N., Mills, J. H., Ahlstrom, J. B., and Dubno, J. R. (**2008**). "Age-related differences in the

temporal modulation transfer function with pure-tone carriers," J Acoust Soc Am, **124**,

3841–3849. doi:10.1121/1.2998779

Henry, K. S., Kale, S., and Heinz, M. G. (**2014**). "Noise-induced hearing loss increases the

temporal precision of complex envelope coding by auditory-nerve fibers," Front Syst

Neurosci, **8**, 1–10. doi:10.3389/fnsys.2014.00020

Houtgast, T. (**1989**). "Frequency selectivity in amplitude-modulation detection," J Acoust Soc

Am, **85**, 1676–1680. doi:10.1121/1.397956

Jennings, S. G., Chen, J., Fultz, S. E., Ahlstrom, J. B., and Dubno, J. R. (**2018**). "Amplitude

modulation detection with a short-duration carrier: Effects of a precursor and hearing

loss," J Acoust Soc Am, **143**, 2232–2243. doi:10.1121/1.5031122

Jepsen, M. L., and Dau, T. (**2011**). "Characterizing auditory processing and perception in

individual listeners with sensorineural hearing loss," J Acoust Soc Am, **129**, 262–281.

doi:10.1121/1.3518768

Jepsen, M. L., Ewert, S. D., and Dau, T. (**2008**). "A computational model of human auditory

signal processing and perception," J Acoust Soc Am, **124**, 422–438.

Jørgensen, S., and Dau, T. (**2011**). "Predicting speech intelligibility based on the signal-to-noise

envelope power ratio after modulation-frequency selective processing," J Acoust Soc

Am, **130**, 1475–1487. doi:10.1121/1.3621502

Jørgensen, S., Ewert, S. D., and Dau, T. (**2013**). "A multi-resolution envelope-power based model

for speech intelligibility," J Acoust Soc Am, **134**, 436–446. doi:10.1121/1.4807563

Joris, P. X., Schreiner, C. E., and Rees, A. (**2004**). "Neural processing of amplitude-modulated

sounds," Physiol Rev, **84**, 541–577. doi:10.1152/physrev.00029.2003

Kale, S., and Heinz, M. G. (**2010**). "Envelope coding in auditory nerve fibers following noise-

induced hearing loss," J Assoc Res Otolaryngol, **11**, 657–673. doi:10.1007/s10162-010-

0223-6

Kale, S., and Heinz, M. G. (**2012**). "Temporal modulation transfer functions measured from
auditory-nerve responses following sensorineural hearing loss," Hear Res, **286**, 64–75.
doi:10.1016/j.heares.2012.02.004

King, A., Varnet, L., and Lorenzi, C. (**2019**). "Accounting for masking of frequency modulation
by amplitude modulation with the modulation filter-bank concept," J Acoust Soc Am,
145, 2277–2293. doi:10.1121/1.5094344

Kowalski, N., Depireux, D. A., and Shamma, S. A. (**1996**). "Analysis of dynamic spectra in ferret
primary auditory cortex. I. Characteristics of single-unit responses to moving ripple
spectra," J Neurophysiol, **76**, 3503–3523. doi:10.1152/jn.1996.76.5.3503

Levitt, H. (**1971**). "Transformed up-down methods in psychoacoustics," J Acoust Soc Am, **49**,
467–477. doi:10.1121/1.1912375

Lorenzi, C., Micheyl, C., Berthommier, F., and Portalier, S. (**1997**). "Modulation masking in
listeners with sensorineural hearing loss," J Speech Lang Hear Res, **40**, 200–207.
doi:10.1044/jslhr.4001.200

McDermott, J. H., Schemitsch, M., and Simoncelli, E. P. (**2013**). "Summary statistics in auditory
perception," Nat Neurosci, **16**, 493–498. doi:10.1038/nn.3347

McDermott, J. H., and Simoncelli, E. P. (**2011**). "Sound texture perception via statistics of the
auditory periphery: Evidence from sound synthesis," Neuron, **71**, 926–940.
doi:10.1016/j.neuron.2011.06.032

McWalter, R., and Dau, T. (**2015**). "Statistical representation of sound textures in the impaired
auditory system," Proceedings of the International Symposium on Auditory and
Audiological Research, 5, Nyborg, Denmark, 186–196.

McWalter, R., and Dau, T. (**2017**). "Cascaded amplitude modulations in sound texture
perception," Front Neurosci, **11**, 1–12. doi:10.3389/fnins.2017.00485

Millman, R. E., Mattys, S. L., Gouws, A. D., and Prendergast, G. (**2017**). "Magnified neural

envelope coding predicts deficits in speech perception in noise," J Neurosci, **37**, 7727–

7736. doi:10.1523/JNEUROSCI.2722-16.2017

Moore, B. C. J., Füllgrabe, C., and Sek, A. (**2009**). "Estimation of the center frequency of the

highest modulation filter," J Acoust Soc Am, **125**, 1075–1081. doi:10.1121/1.3056562

Moore, B. C. J., and Glasberg, B. R. (**1998**). "Use of a loudness model for hearing-aid fitting. I.

Linear hearing aids," Br J Audiol, **32**, 317–335. doi:10.3109/03005364000000083

Moore, B. C. J., and Glasberg, B. R. (**2001**). "Temporal modulation transfer functions obtained

using sinusoidal carriers with normally hearing and hearing-impaired listeners," J Acoust

Soc Am, **110**, 1067–1073. doi:10.1121/1.1385177

Moore, B. C. J., Mariathasan, S., and Sęk, A. P. (**2019**). "Effects of age and hearing loss on the

discrimination of amplitude and frequency modulation for 2- and 10-Hz rates," Trends

Hear, **23**, 233121651985396. doi:10.1177/2331216519853963

Moore, B. C. J., and Oxenham, A. J. (**1998**). "Psychoacoustic consequences of compression in

the peripheral auditory system.," Psychol Rev, **105**, 108–124. doi:10.1037/0033-

295X.105.1.108

Moore, B. C. J., Shailer, M. J., and Schooneveldt, G. P. (**1992**). "Temporal modulation transfer

functions for band-limited noise in subjects with cochlear hearing loss," Br J Audiol, **26**,

229–237. doi:10.3109/03005369209076641

Moore, B. C. J., Wojtczak, M., and Vickers, D. A. (**1996**). "Effect of loudness recruitment on the

perception of amplitude modulation," J Acoust Soc Am, **100**, 481–489.

doi:10.1121/1.415861

Oetjen, A., and Verhey, J. L. (**2015**). "Spectro-temporal modulation masking patterns reveal

frequency selectivity," J Acoust Soc Am, **137**, 714–723. doi:10.1121/1.4906171

Oetjen, A., and Verhey, J. L. (**2017**). "Characteristics of spectro-temporal modulation frequency

selectivity in humans," J Acoust Soc Am, **141**, 1887–1895. doi:10.1121/1.4976537

Paraouty, N., Ewert, S. D., Wallaert, N., and Lorenzi, C. (**2016**). "Interactions between amplitude

modulation and frequency modulation processing: Effects of age and hearing loss," J

Acoust Soc Am, **140**, 121–131. doi:10.1121/1.4955078

Paraouty, N., and Lorenzi, C. (**2017**). "Using individual differences to assess modulation-

processing mechanisms and age effects," Hear Res, **344**, 38–49.

doi:10.1016/j.heares.2016.10.024

Piechowiak, T., Ewert, S. D., and Dau, T. (**2007**). "Modeling comodulation masking release using

an equalization-cancellation mechanism," J Acoust Soc Am, **121**, 2111–2126.

doi:10.1121/1.2534227

Regev, J., Zaar, J., Relaño-Iborra, H., and Dau, T. (**2023**). "Age-related reduction of amplitude

modulation frequency selectivity," J Acoust Soc Am, **153**, 2298. doi:10.1121/10.0017835

Regev, J., Zaar, J., Relaño-Iborra, H., and Dau, T. (**2024**). "Dataset for: 'Age-related reduction of

amplitude modulation frequency selectivity,'" Technical University of Denmark, Dataset.

doi:10.11583/DTU.25134527.v1

Relaño-Iborra, H., and Dau, T. (**2022**). "Speech intelligibility prediction based on modulation

frequency-selective processing," Hear Res, **426**, 108610, doi:

10.1016/j.heares.2022.108610. doi:10.1016/j.heares.2022.108610

Relaño-Iborra, H., May, T., Zaar, J., Scheidiger, C., and Dau, T. (**2016**). "Predicting speech

intelligibility based on a correlation metric in the envelope power spectrum domain," J

Acoust Soc Am, **140**, 2670–2679. doi:10.1121/1.4964505

Relaño-Iborra, H., Zaar, J., and Dau, T. (**2019**). "A speech-based computational auditory signal

819   processing and perception model," J Acoust Soc Am, **146**, 3306–3317.

820   doi:10.1121/1.5129114

821   Sabin, A. T., Eddins, D. A., and Wright, B. A. (**2012**). "Perceptual learning evidence for tuning

822   to spectrotemporal modulation in the human auditory system," J Neurosci, **32**, 6542–

823   6549. doi:10.1523/JNEUROSCI.5732-11.2012

824   Schlittenlacher, J., and Moore, B. C. J. (**2016**). "Discrimination of amplitude-modulation depth

825   by subjects with normal and impaired hearing," J Acoust Soc Am, **140**, 3487–3495.

826   doi:10.1121/1.4966117

827   Schoof, T., and Rosen, S. (**2014**). "The role of auditory and cognitive factors in understanding

828   speech in noise by normal-hearing older listeners," Front Aging Neurosci, **6**, 1–14.

829   doi:10.3389/fnagi.2014.00307

830   Sek, A., Baer, T., Crinnion, W., Springgay, A., and Moore, B. C. J. (**2015**). "Modulation masking

831   within and across carriers for subjects with normal and impaired hearing," J Acoust Soc

832   Am, **138**, 1143–1153. doi:10.1121/1.4928135

833   Sek, A., and Moore, B. C. J. (**2003**). "Testing the concept of a modulation filter bank: The

834   audibility of component modulation and detection of phase change in three-component

835   modulators," J Acoust Soc Am, **113**, 2801–2811. doi:10.1121/1.1564020

836   Singh, N. C., and Theunissen, F. E. (**2003**). "Modulation spectra of natural sounds and

837   ethological theories of auditory processing," J Acoust Soc Am, **114**, 3394–3411.

838   doi:10.1121/1.1624067

839   Steinmetzger, K., Zaar, J., Relaño-Iborra, H., Rosen, S., and Dau, T. (**2019**). "Predicting the

840   effects of periodicity on the intelligibility of masked speech: An evaluation of different

modelling approaches and their limitations," J Acoust Soc Am, **146**, 2562–2576. doi:10.1121/1.5129050

Strickland, E. A., and Viemeister, N. F. (**1996**). "Cues for discrimination of envelopes," J Acoust Soc Am, **99**, 3638–3646. doi:10.1121/1.414962

Takahashi, G. A., and Bacon, S. P. (**1992**). "Modulation detection, modulation masking, and speech understanding in noise in the elderly," J Speech Lang Hear Res, **35**, 1410–1421. doi:10.1044/jshr.3506.1410

The Mathworks Inc. (**2015**). "MATLAB 2015b - MathWorks," www.mathworks.com/products/matlab,.

Verhey, J. L., Dau, T., and Kollmeier, B. (**1999**). "Within-channel cues in comodulation masking release (CMR): Experiments and model predictions using a modulation-filterbank model," J Acoust Soc Am, **106**, 2733–2745. doi:10.1121/1.428101

Viemeister, N. F. (**1979**). "Temporal modulation transfer functions based upon modulation thresholds," J Acoust Soc Am, **66**, 1364–1380. doi:10.1121/1.383531

Wallaert, N., Moore, B. C. J., Ewert, S. D., and Lorenzi, C. (**2017**). "Sensorineural hearing loss enhances auditory sensitivity and temporal integration for amplitude modulation," J Acoust Soc Am, **141**, 971–980. doi:10.1121/1.4976080

Wallaert, N., Moore, B. C. J., and Lorenzi, C. (**2016**). "Comparing the effects of age on amplitude modulation and frequency modulation detection," J Acoust Soc Am, **139**, 3088–3096. doi:10.1121/1.4953019

Wiinberg, A., Jepsen, M. L., Epp, B., and Dau, T. (**2019**). "Effects of hearing loss and fast-acting compression on amplitude modulation perception and speech intelligibility," Ear Hear, **40**, 45–54. doi:10.1097/AUD.0000000000000589

Wojtczak, M. (**2011**). "The effect of carrier level on tuning in amplitude-modulation masking," J Acoust Soc Am, **130**, 3916–3925. doi:10.1121/1.3658475

Wojtczak, M., and Viemeister, N. F. (**2005**). "Forward masking of amplitude modulation: Basic characteristics," J Acoust Soc Am, **118**, 3198–3210. doi:10.1121/1.2042970

Zaar, J., and Carney, L. H. (**2022**). "Predicting speech intelligibility in hearing-impaired listeners using a physiologically inspired auditory model," Hear Res, **426**, 108553. doi:10.1016/j.heares.2022.108553

Zaar, J., and Dau, T. (**2017**). "Predicting consonant recognition and confusions in normal-hearing listeners," J Acoust Soc Am, **141**, 1051–1064. doi:10.1121/1.4976054

Zaar, J., Schmitt, N., Derleth, R.-P., DiNino, M., Arenberg, J. G., and Dau, T. (**2017**). "Predicting effects of hearing-instrument signal processing on consonant perception," J Acoust Soc Am, **142**, 3216–3226. doi:10.1121/1.5011737

Zhong, Z., Henry, K. S., and Heinz, M. G. (**2014**). "Sensorineural hearing loss amplifies neural coding of envelope information in the central auditory system of chinchillas," Hear Res, **309**, 55–62. doi:10.1016/j.heares.2013.11.006

Zilany, M. S. A., and Bruce, I. C. (**2007**). "Predictions of speech intelligibility with a model of the normal and impaired auditory-periphery," 2007 3rd International IEEE/EMBS Conference on Neural Engineering, IEEE, 481–485. doi:10.1109/CNE.2007.369714