Disentangling the effects of hearing loss and age on amplitude modulation frequency selectivity

Jonathan Regev[1,a], Helia Relaño-Iborra[1], Johannes Zaar[1,2], and Torsten Dau[1,3]

[1] *Hearing Systems Section, Department of Health Technology, Technical University of Denmark, Kongens Lyngby, 2800, Denmark*

[2] *Eriksholm Research Centre, Snekkersten, 3070, Denmark*

[3] *Copenhagen Hearing and Balance Center, Copenhagen University Hospital, Rigshospitalet, Copenhagen, 2100, Denmark*

Abstract

The processing and perception of amplitude modulations (AMs) in the auditory system reflect a frequency-selective process, often described as a modulation filterbank. Previous studies on perceptual AM masking reported similar results for older listeners with hearing impairment (HI) and young listeners with normal hearing (NH), suggesting no effects of age nor hearing loss on AM frequency selectivity. However, recent evidence has shown that age, independently of hearing loss, is detrimental to AM frequency selectivity. Hence, the present study aimed to disentangle the effects of hearing loss and age. A simultaneous AM masking paradigm was employed, utilizing a sinusoidal carrier at 2.8 kHz, narrow-band noise modulation maskers, and target modulation frequencies of 4, 16, 64, and 128 Hz. The results obtained from older (n=10, 63-77 years) and young HI listeners (n=3, 24-30 years) were

---

[a] Corresponding author: joreg@dtu.dk

compared to data from young and older NH listeners. Notably, the HI listeners generally exhibited lower (unmasked) AM detection thresholds and greater AM frequency selectivity compared to their NH counterparts of similar age. These findings suggest that age negatively affects AM frequency selectivity in both NH and HI listeners, while hearing loss improves AM detection and AM selectivity, likely due to the loss of peripheral compression.

I.   INTRODUCTION

Most sounds encountered in everyday environments are complex signals that exhibit dynamic fluctuatations over time (Singh and Theunissen, 2003; Elliott and Theunissen, 2009). These fluctuations include variations in the temporal envelope, commonly referred to as amplitude modulations (AMs). The auditory system demonstrates a remarkable sensitivity to AMs, as evidenced by perceptual studies (e.g., Viemeister, 1979) and physiological investigations (see Joris *et al.*, 2004 for a comprehensive review). Similar to spectral masking effects in the audio-frequency domain, masking effects also occur in the AM frequency domain, resulting in reduced sensitivity to a target AM in the presence of a masking AM (Bacon and Grantham, 1989; Houtgast, 1989; Dau *et al.*, 1997a, 1997b; Ewert and Dau, 2000; Sek and Moore, 2003). Specifically, AM masking patterns provide evidence for a frequency-selective process, where the amount of AM masking decreases as the spectral distance between the masker and the target increases (Bacon and Grantham, 1989; Houtgast, 1989; Dau *et al.*, 1997a; Ewert and Dau, 2000; Ewert *et al.*, 2002; Sek and Moore, 2003; Moore *et al.*, 2009; Sek *et al.*, 2015; Füllgrabe *et al.*, 2021a, 2021b; Conroy *et al.*, 2023).

This frequency-selective characteristic of AM processing has been modelled using the concept of a modulation filterbank, which proposes that AM fluctuations are decomposed through an array of relatively broad bandpass modulation filters with a constant quality (Q) factor of approximately 1-2 (e.g., Dau *et al.*, 1997a, 1999; Ewert and Dau, 2000). Computational modelling studies have successfully applied the modulation filterbank concept to simulate data from various experimental paradigms, including simultaneous and non-simultaneous spectral and temporal signal detection and masking conditions (Dau *et al.*, 1997a, 1997b; Dau and Verhey, 1999; Verhey *et al.*, 1999; Ewert and Dau, 2000; Ewert *et al.*, 2002; Piechowiak *et al.*, 2007; Jepsen *et al.*, 2008; Jepsen and Dau, 2011; King *et al.*, 2019), sound texture perception (McDermott and Simoncelli, 2011; McDermott *et al.*, 2013; McWalter and Dau, 2015, 2017), auditory stream segregation (Elhilali *et al.*, 2009; Christiansen *et al.*, 2014), and speech

intelligibility (Jørgensen and Dau, 2011; Jørgensen et al., 2013; Relaño-Iborra et al., 2016, 2019; Zaar

and Dau, 2017; Zaar et al., 2017; Steinmetzger et al., 2019; Zaar and Carney, 2022; for a review, see

Relaño-Iborra and Dau, 2022). Furthermore, the modulation filterbank is conceptually consistent with

the temporal dimension of a 'two-dimensional' spectro-temporal modulation filterbank, inspired by

neural responses to spectro-temporally varying stimuli in the auditory cortex of ferrets (Kowalski *et*

al., 1996; Depireux *et al.*, 2001) and supported by data from perceptual learning and masking conditions

(Sabin *et al.*, 2012; Oetjen and Verhey, 2015, 2017; Conroy *et al.*, 2022), as well as models of speech

intelligibility (Elhilali *et al.*, 2003; Chi *et al.*, 2005; Zilany and Bruce, 2007; Chabot-Leclerc *et al.*, 2014).

Overall, the versatility of the modulation filterbank model suggests that AM frequency selectivity is

an essential auditory processing feature for quantitatively predicting perceptual data obtained with

dynamically varying sounds.

Several studies have investigated the effects of age and hearing loss on AM perception, yielding

mixed results. In terms of age effects, some studies showed a significant age-related deterioration in

AM detection (He *et al.*, 2008; Füllgrabe *et al.*, 2015; Wallaert *et al.*, 2016), while other investigations

did not find any effects on AM detection or AM depth discrimination (Schoof and Rosen, 2014;

Paraouty *et al.*, 2016; Schlittenlacher and Moore, 2016; Paraouty and Lorenzi, 2017). Studies comparing

listeners with normal hearing to listeners with hearing impairment (referred to hereafter as NH and

HI listeners, respectively) have found either similar or lower (i.e., better) AM detection thresholds for

the HI listeners (e.g., Moore *et al.*, 1992, 1996; Moore and Glasberg, 2001; Füllgrabe *et al.*, 2003; Sek *et*

al., 2015; Schlittenlacher and Moore, 2016; Wallaert *et al.*, 2017; Wiinberg *et al.*, 2019). The observed

improvement in AM detection with hearing loss has been suggested to result from the loss of cochlear

compression typically associated with sensorineural hearing loss (SNHL; Moore and Oxenham, 1998),

which may lead to an increased internal representation of AM depth in HI listeners (Moore *et al.*, 1996;

Jennings *et al.*, 2018). Additionally, the amplified neural coding of the envelope observed in mammals

and humans with SNHL (Kale and Heinz, 2010, 2012; Henry *et al.*, 2014; Zhong *et al.*, 2014; Millman

et al., 2017; Goossens *et al.*, 2018; Decruy *et al.*, 2020) may contribute, at least in part, to the effective

increase in perceived AM depth with hearing loss. However, hearing loss has been shown to have a

detrimental effect on supra-threshold AM depth discrimination (Schlittenlacher and Moore, 2016;

Wiinberg *et al.*, 2019), suggesting that the increase in perceived AM depth may not necessarily be

beneficial to AM depth discrimination.

Behavioral studies on AM frequency selectivity generally did not yield conclusive evidence of an

effect of hearing loss (Takahashi and Bacon, 1992; Lorenzi *et al.*, 1997; Sek *et al.*, 2015). These studies

focused on masked-threshold patterns (MTPs), which assess the detection threshold of an AM target

in relation to the (center) frequency of a masker of fixed modulation depth. Takahashi and Bacon

(1992) compared MTPs (using noise carriers and sinusoidal modulation targets and maskers) from ten

young NH listeners and three groups of ten older listeners with mild hearing loss (classified as 'age-

appropriate' normal hearing). While the older listeners faced more difficulties overall in completing

the AM masking task, no simultaneous effect of age and hearing loss on AM frequency selectivity was

found. Lorenzi *et al.* (1997) employed a similar approach with four young NH and three young HI

listeners, observing comparable MTPs across both groups, despite a potential higher masking effect

of modulation maskers located below the target modulation frequency in the HI listener group.

Additionally, Sek *et al.* (2015), using sinusoidal carriers and sinusoidal modulation maskers, found no

differences in AM masking patterns between six young NH and nine older HI listeners. These results

suggested that neither age nor hearing loss affect AM frequency selectivity (Sek *et al.*, 2015).

However, recent results by Regev *et al.* (2023) showed a reduction in AM frequency selectivity

associated with age when comparing young and older listeners with clinically normal hearing. These

results are qualitatively consistent with the observations of reduced temporal selectivity inferred from

auditory cortical functional magnetic resonance imaging (fMRI) responses by Erb *et al.* (2020). Regev

et al. (2023), using sinusoidal carriers and noise modulation maskers, compared MTPs from eleven

young NH listeners and ten older NH listeners. They found an age-related increase (i.e., deterioration)

in AM detection thresholds and a decrease in AM frequency selectivity, with estimated modulation

filter Q factors that were approximately halved in the older NH listeners compared to the young NH

listeners. Thus, the contrast between these recent findings and previous studies that did not identify a

simultaneous effect of age and hearing loss on AM frequency selectivity (Takahashi and Bacon, 1992;

Sek *et al.*, 2015) may suggest that hearing loss and age have antagonistic effects with regards to AM

frequency selectivity. In other words, hearing loss may 'counteract' the detrimental effects of ageing.

However, a direct comparison of MTPs between young NH listeners and similarly aged groups of

older NH and HI listeners tested using the same experimental approach has not yet been conducted.

The present study therefore aimed to independently evaluate the effects of hearing loss and age

on AM frequency selectivity by collecting data from HI listeners who were approximately age-matched

to the older NH listeners included in Regev *et al.* (2023), using the same experimental approach. Young

HI listeners were also included to explore whether a perceptual 'benefit' from hearing loss could be

observed in the absence of ageing effects. The study involved ten older and three young HI listeners,

ranging in age from 63 to 77 and 24 to 30 years, respectively. MTPs were collected using an AM

masking paradigm with sinusoidal carriers and bandpass Gaussian-noise modulation maskers (Ewert

et al., 2002). Additionally, unmasked AM detection thresholds were obtained using sinusoidal carriers.

The data obtained from the older HI listeners were compared to those obtained from the older NH

listeners reported in Regev *et al.* (2023) to assess the effects of hearing loss on AM frequency selectivity.

Furthermore, the data from the older HI listeners were also compared to the young NH listener results

of Regev *et al.* (2023) to determine whether the lack of a simultaneous effect of age and hearing loss

observed in previous studies could be replicated (Takahashi and Bacon, 1992; Sek *et al.*, 2015). Despite

the limited number of young HI listeners , their data were included to provide insights into the effects

of hearing loss in the absence of ageing. To quantify the AM frequency selectivity reflected in the data

at the listener group level, a simple approach applying linear fits separately to each skirt of the MTPs

was employed.

II.   METHODS

A. Listeners

Ten older HI listeners, aged 63-77 years (mean age = 70.4 years, 2 female) and three young HI

listeners, aged 24-30 years (mean age = 26 years, all female) participated in the study. On average, the

older HI listeners were 4.5 years older than the older NH listeners tested in Regev *et al.* (2023), while

the young HI listeners were less than one year older than the young NH listeners. All listening

experiments were conducted monaurally. The individual audiograms in the test ear are shown in Figure

1. The older HI listeners had a high-frequency sloping hearing loss, classified as an N2 or N3 standard

audiogram (Bisgaard *et al.*, 2010).[1] The young HI listeners had more heterogeneous audiograms. For

each listener, the test ear was selected as the one where the audiometric threshold at 3 kHz was within

50 to 60 dB HL. These limits were chosen to ensure that the listeners had a significant hearing loss

around the AM carrier frequency (2.8 kHz) and, at the same time, that the stimuli's sound pressure

level (SPL) would not exceed 85 dB to avoid discomfort. All listeners had a SNHL, as indicated by

air-bone gaps ≤ 15 dB.[2] Working memory capacity was assessed with a reverse digit span test (RDS),

implemented binaurally as described in Fuglsang *et al.* (2020) with an added linear gain according to

the Cambridge formula (CamEq; Moore and Glasberg, 1998) using a maximum-gain limit of 30 dB at

any frequency. The young and older HI listeners were found to have similar average RDS scores (0.56

and 0.53 on a normalized scale, respectively), which are also comparable to the average RDS scores

---

[1] The classification of the collected audiogram as one of the standard audiograms was performed using the lowest mean squared error between the standard audiograms and the audiometric thresholds at frequencies from 1 to 8 kHz.
[2] Except for one young HI listener who showed a gap of 20 dB at 4 kHz and one older HI listener who showed a gap of 20 dB 1 kHz.

reported for the young NH listeners in Regev *et al.* (2023) and slightly, but not significantly, higher

than for Regev *et al.*'s older NH cohort (0.59 and 0.42, respectively).

All participants gave written informed consent. Ethical approval for the study was provided by

the Science Ethics Committee of the Capital Region in Denmark (reference H-16036391). All

participants were financially compensated for their time.

Figure 1: Individual and average audiograms in the test ear, for all participants. The left and right

panels show the audiograms for the young and older HI listeners, respectively. Individual audiograms

are shown in light gray, and group averages and standard deviations are shown in black.

B. Procedure and apparatus

The experimental conditions and parameters were the same as described in Regev *et al.* (2023),

with few modifications introduced to compensate for audibility loss. The stimulus presentation level

in all experiments was adjusted to be 30 dB sensation level (SL) for all HI listeners, based on their

hearing threshold at the carrier frequency, in an approach similar to that of Sek *et al.* (2015). This

resulted in presentation levels of 78-80 dB SPL for the young HI listeners and 73-85 dB SPL for the

older HI listeners (79 dB SPL on average for both groups).

The hearing threshold for the unmodulated pure-tone carrier at 2.8 kHz was measured adaptively.

The signal had a duration of 300 ms, including 50-ms raised-cosine ramps, and the intervals were

separated by 600 ms of silence. The step size was initially 8 dB, was then reduced to 4 dB after the

first lower reversal, and was finally reduced to 2 dB after the second lower reversal. The threshold was

defined as the average SPL of the target tone across the last six reversals. The threshold measurement

was repeated at least three times, and the final threshold was computed as the average across all

repetitions.

All listening tests were conducted in double-walled soundproof booths. All thresholds were

measured using a 3-interval 3-alternative forced-choice (3I-3AFC) paradigm with a 1-up 2-down

tracking procedure, measuring the 70.7%-correct point on the psychometric function (Levitt, 1971).

The experiments were implemented using the AFC toolbox in MATLAB (Ewert, 2013). The stimuli

were digitally generated in MATLAB (The Mathworks Inc., 2015; Natick, Massachusetts) with a

sampling frequency of 48 kHz, converted to analog using an RME Fireface soundcard (Haimhausen,

Germany), and played through Sennheiser HDA200 headphones (Wedemark, Germany). The output

level was calibrated using a GRAS RA0039 ear simulator (Holte, Denmark) and the frequency

response of the headphones was compensated using an inverse digital filter. The listeners received

visual feedback during the tests ("correct"/"incorrect"). The order of testing was the same for both

groups, as summarized in Appendix A.

C. Experimental conditions

1. Stimuli

The AM detection task and masked-threshold patterns (MTPs; Ewert and Dau, 2000) shared

common aspects of the stimuli. In both conditions, the thresholds were measured for sinusoidal AM

at target modulation frequencies of 4, 16, 64, and 128 Hz, imposed on a 2.8-kHz sinusoidal carrier.

The carrier had a duration of 600 ms, including 50-ms raised-cosine onset and offset ramps. The target

modulation (and masker modulation in the MTP task) had a duration of 500 ms, including 50-ms

raised-cosine onset and offset ramps, and was temporally centered in the carrier. The target

modulation was imposed with a 0-degree starting phase. The stimulus intervals were separated by 500

ms of silence. The stimuli of all three intervals were calibrated to the desired presentation level to

ensure that no level cues were available. The tracking variable was the target's modulation depth,

expressed on a dB scale as $M = 20log_{10}(m)$, where m is the modulation depth on a linear scale. The

step size was initially 4 dB, was reduced to 2 dB after the first lower reversal, and was finally reduced

to 1 dB after the second lower reversal.

a. AM detection thresholds

For the AM detection task, the starting modulation depth of the target was -5 dB. The threshold

for any measurement run was defined as the average target modulation depth across the last six

reversals. The measurement at each target modulation frequency was repeated three times, and the

final threshold was defined as the average across repetitions. Additional measurements were included

if the standard error across repetitions exceeded 2 dB, until that limit was respected.[3] All listeners

completed two training runs (at the target modulation frequencies of 4 and 128 Hz). Figure 2 shows

an example of the stimulus envelope for the reference interval (i.e., the unmodulated carrier, in panel

A) and the target interval (i.e., the carrier modulated with the target modulation, in panel B) for a target

modulation frequency of 128 Hz.

b. Masked-threshold patterns

For the MTP task, the target and masker modulations were applied simultaneously on the carrier

using a multiplicative approach, as described in Regev *et al.* (2023). The masker modulation was a

narrow-band Gaussian noise with a fixed bandwidth corresponding to ½-octave when centered on

the target modulation frequency. For target modulation frequencies of 4 and 16 Hz, the masker

modulation was centered at frequencies ranging from -2 to +2 octaves relative to the target modulation

---

[3] Overall, 8 measurements were added across 2 of the young HI listeners and 1 measurement was added for a single older HI listener.

frequency, in steps of 2/3 octaves. In order to avoid cues from spectrally resolved modulation

sidebands, the highest masker-modulation center frequency was limited to 128 Hz, such that the

masker-modulation center frequencies only ranged from -2 to +2/3 and 0 octaves for the target

modulation frequencies of 64 and 128 Hz, respectively. Furthermore, for these target modulation

frequencies, a masked threshold was also measured for a masker modulation centered around 4 Hz,

corresponding to -4 and -5 octaves relative to the target modulation frequencies, respectively. For

these additional conditions, the lower cutoff frequency of the masker modulation was set to 1 Hz,

thus effectively reducing the masker's bandwidth. The resulting masker-modulation bandwidths and

center frequencies for all conditions are summarized in Appendix B. For each measurement run, a 2

s-long generation of the masker modulation was created. For each interval, a 500-ms segment,

randomly cut out of the long masker modulation, was then applied on the carrier. The root-mean

square (rms) modulation depth of the masker modulation was fixed to -10 dB.

The starting modulation depth of the target was 0 dB. The threshold for any measurement run

was defined as the average target modulation depth across the last eight reversals. The measurement

at each masker-modulation center frequency was repeated three times, and the final threshold was

defined as the average across repetitions. To avoid overmodulation, the maximum allowed modulation

depth was 0 dB. If the procedure required a modulation depth higher than 0 dB more than three times

over the course of a measurement run, the run was stopped and an extra measurement run was carried

out to replace it. If this happened more than 5 times for any condition, the threshold was considered

as unmeasurable. This was the case for a single older HI listener, where thresholds could not be

estimated at masker-modulation center frequencies of +2/3 and +4/3 octaves relative to the 4 Hz

target modulation frequency. This listener's masked thresholds at 4 Hz were therefore not included in

the statistical analyses, or in the figures below, such that the MTP analyses at 4 Hz included only nine

older HI listeners. Additional measurements were also included if the standard error across repetitions

exceeded 2 dB, until that limit was respected.[4] For each MTP, the listeners completed three training

run (at the masker-modulation center frequencies of -2, 0, and +2 octaves relative to the target

modulation frequency). Figure 2 shows an example of the stimulus envelope for the reference interval

(i.e., the carrier modulated with the masker modulation, in panel C) and the target interval (i.e., the

carrier modulated with both the target and masker modulations, in panel D), for a target modulation

frequency of 128 Hz and a masker-modulation center frequency of 64 Hz.

Figure 2: Examples of the temporal envelopes of the stimuli used in the AM detection and MTP

experiments. Panel A: Reference interval in the AM detection task, represented by an unmodulated

sinusoidal carrier (2.8 kHz). Panel B: Target interval in the AM detection experiment, containing the

---

[4] Overall, a total of 2 measurements were added for a single young HI listener, and 7 measurements for 3 of the older HI listeners, due to aborted measurement runs. Furthermore, 5 measurements were added for 2 of the young HI listeners and 10 measurements for 4 of the older HI listeners, due to excessive standard errors. Masked thresholds could not be estimated for a single older HI listener, at masker modulation center frequencies of +2/3 and +4/3 octaves relative to the 4-Hz target modulation frequency.

same carrier as in Panel A modulated by a sinusoidal target modulation (128 Hz). Panel C: Reference

interval in the MTP experiment, containing the same carrier as in Panel A modulated by a narrowband-

noise masker modulation (between 43.6 and 88.2 Hz, centered at 64 Hz). Panel D: Target interval in

the MTP experiment, with the same carrier as in Panel A modulated by the sinusoidal target

modulation and the narrowband-noise masker modulation.

2. Estimates of AM frequency selectivity

Ewert *et al.* (2002) and Regev *et al.* (2023) used the envelope power spectrum model of masking

(EPSM; Ewert and Dau, 2000) to quantify the AM frequency selectivity reflected in their MTPs, by

finding the Q-factor of the best-fitting bandpass modulation filter representing the experimental data.

However, high levels of asymmetry were observed in the MTPs in the present study (as described in

detail below), which are not consistent with the EPSM's assumption of symmetry. Hence, a simplified

approach was used here to quantify AM frequency selectivity. The best-fitting linear approximation

(in terms of the least-squared error) was obtained separately for each skirt using the masked thresholds

at the masker locations ranging from -2 to +2 octaves. Each linear fit was constrained to crossing the

on-target masked threshold. The bandwidth of the asymmetrical modulation filter was then estimated

by measuring the spectral distance between the 3-dB-down points on the fits obtained for each skirt

when interpolating across masker locations ranging from -4 to +4 octaves. For the 128-Hz pattern,

the high-frequency cutoff was estimated by assuming symmetry around the target modulation

frequency. Finally, the metric quantifying AM frequency selectivity was obtained as an overall 'Q-

factor' by computing the ratio of the filter's characteristic (modulation) frequency to the fitted

bandwidth.

D. Statistical analysis

Analyses of variances (ANOVAs) of linear mixed-effects model fittings were used to assess the differences between the results for the older HI listeners obtained in the present study and the data for the NH listeners obtained in Regev *et al.* (2023). Each model and corresponding ANOVA were applied twice. First, to estimate the effects of hearing loss in isolation from age effects, the results for the older HI listeners were compared with the results for the older NH listeners. Second, using an analysis comparable to that employed in previous studies, the results for the older HI listeners were compared with the results obtained for the young NH listeners. For the analysis of the AM detection thresholds, the listeners were considered as a random effect and group classification and modulation frequency were considered as fixed effects. The MTPs were analyzed separately for each target modulation frequency. The listeners were considered as a random effect and group classification and masker-modulation center frequency were considered as fixed effects. Levene's test was used to assess the homoskedasticity of the residuals and Shapiro-Wilk's test was used to assess the normality of residual's distribution.[5] Post-hoc analyses were conducted using estimated marginal means and a Holm-Bonferroni correction criterion was applied for multiple comparisons. The level of statistical significance was set to 5% in all analyses. Due to the low number of young HI listeners (3), no statistical analysis was performed to compare their data to those of the other groups.

---

[5] Levene's test was not passed for one of the models applied, while Shapiro-Wilk's test was not passed for 3 of the other models. However, the deviations were minor, and the model predictions matched the average group data accurately. Hence, no transformations of the data were applied in order to keep all applied analyses consistent.

III. RESULTS

A. AM detection thresholds

Figure 3 (Color online): Average AM detection thresholds for each listener group as a function of modulation frequency. The data for the young HI listeners are indicated by diamonds connected by solid lines. The data for the older HI listeners are shown as circles connected by solid lines. The results for the young and the older NH listeners (dashed lines; reproduced from Regev *et al.*, 2023) are shown as triangles and squares, respectively. Means and standard errors are shown. The stars indicate the level of statistical significance of the differences in thresholds between the older HI and older NH listeners (dark stars), and the older HI and young NH listeners (light stars; *$p < 0.05$, **$p < 0.01$, *** $p < 0.001$).

Figure 3 shows the mean AM detection thresholds as a function of modulation frequency, for both the young HI (diamonds) and the older HI listeners (circles), as well the young NH (squares) and the older NH listeners (triangles) reproduced from Regev *et al.* (2023). Substantial levels of individual variability were observed in the data (see supplementary Figure 1). The statistical analysis comparing the thresholds for the older HI and the older NH listeners revealed a significant main effect of group

$[F_{1,18} = 21.21, p < 0.001]$ and modulation frequency $[F_{3,54} = 22.84, p < 0.001]$. The post-hoc

analysis revealed that the older HI listeners had significantly lower (i.e., better) AM detection

thresholds than the older NH listeners at all modulation frequencies [effect size $(M) = -6.04$ dB,

standard error $(SE) = 1.995$ dB; $M = -8.47$ dB, $SE = 1.995$ dB; $M = -7.43$ dB, $SE = 1.995$

dB; $M = -7.05$ dB, $SE = 1.995$ dB, for 4, 16, 64, and 128 Hz, respectively].

The statistical analysis comparing the thresholds for the older HI and the young NH listeners

revealed a significant main effect of group $[F_{1,19} = 8.13, p = 0.01]$ and modulation frequency

$[F_{3,57} = 14.66, p < 0.001]$, as well as a significant interaction between both factors $[F_{3,57} = $

$5.26, p = 0.003]$. The post-hoc analysis revealed that the older HI listeners had significantly lower

AM detection thresholds than the young NH listeners at modulation frequencies of 16 Hz $[M = $

-7.6 dB, $SE = 1.82$ dB] and 64 Hz $[M = -4.86$ dB, $SE = 1.82$ dB].

B. Masked-threshold patterns (MTPs)

Figure 4 (Color online): Average MTPs (filled symbols) and AM detection thresholds (open symbols, replotted from Figure 3[6]) for the older HI and the older NH listeners. The data for the HI listeners are indicated by the circles connected by solid lines and the data for the NH listeners (reproduced from Regev *et al.*, 2023) are shown as triangles connected by dashed lines. Each panel represents the results for a specific target modulation frequency: 4 Hz (top left), 16 Hz (top right), 64 Hz (bottom left), and 128 Hz (bottom right). Note that the data from one of the older HI listeners was excluded from the 4-Hz panel for both the MTP and AM detection threshold. Means and standard errors are shown. Stars indicate the level of statistical significance of the differences between in masked thresholds between the two groups (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

---

[6] The AM detection threshold for a target modulation frequency of 4 Hz reflects the data for only 9 of the 10 older HI listeners.

Figure 4 shows the mean MTPs for the older HI (circles) and the older NH listeners (triangles;

reproduced from Regev *et al.*, 2023). Each panel shows the MTPs for a specific target modulation

frequency. The mean AM detection thresholds are also shown, indicated by the open symbols

(replotted from Figure 3[6]). The statistical analysis comparing the older HI and the older NH listeners

revealed a significant main effect of group at all target modulation frequencies $[F_{1,17} = 7.31, p =$

$0.015; F_{1,18} = 20.18, p < 0.001; F_{1,18} = 8.70, p = 0.009; F_{1,18} = 14.43, p = 0.001,$ for 4, 16,

64, and 128 Hz, respectively]. A significant interaction between listener group and masker-modulation

center frequency was found at the target modulation frequencies of 4 Hz $[F_{6,102} = 3.36, p = 0.005]$

and 16 Hz $[F_{6,108} = 4.64, p < 0.001]$, whereas the interaction approached, but did not reach,

significance at the target modulation frequency of 64 Hz $[F_{5,90} = 2.01, p = 0.085]$. At the target

modulation frequency of 4 Hz, the post-hoc analysis revealed that the older HI listeners had

significantly lower (better) masked thresholds than the older NH listeners for the masker modulations

centered at -2, -4/3, and -2/3 octaves relative to the target frequency. At 16 Hz, the post-hoc analysis

revealed that the older HI listeners had significantly lower masked thresholds than the older NH

listeners for the masker modulations centered at -2, -4/3, and +2 octaves relative to the target

frequency. At 64 Hz, the post-hoc analysis revealed that the older HI listeners had significantly lower

masked thresholds than the older NH listeners for the masker modulations centered at -4, -2, and

+2/3 octaves relative to the target frequency. Finally, at 128 Hz, the post-hoc analysis revealed that

the older HI listeners had significantly lower masked thresholds than the older NH listeners for the

masker modulations centered at -5, -2, -4/3, and 0 octaves relative to the target frequency. The details

of the post-hoc analyses can be found in Appendix B. Overall, a sharpening of the MTPs was found

for the older HI listeners, as compared to the older NH listeners, at the target modulation frequencies

of 4 Hz (on the low-frequency skirt), 16 Hz, and 64 Hz. Conversely, the high-frequency skirt of the

4-Hz MTP suggests a similar selectivity for the older HI and NH listeners. Finally, the MTP at 128

Hz reflects a general improvement (i.e., decrease) of the masked thresholds for the older HI listeners,

but an overall broader pattern, compared to the older NH listeners. High levels of asymmetry were

also observed in the older HI listener's MTPs, as quantified by the slopes of the obtained linear fits

(see supplementary Figure 2). At the target modulation frequency of 4 Hz, this was reflected by a

steeper slope of the low-frequency skirt (3 dB/octave) than the high-frequency skirt (1 dB/octave)

and an off-target peak threshold (for the masker centered at +2/3 octaves relative to the target

frequency). A similar observation was made at 16 Hz, with a steeper slope of the low-frequency skirt

(5.6 dB/octave) than that that of the high-frequency skirt (4.6 dB/octave). At 64 Hz, the older HI

listeners displayed a much shallower slope of the low-frequency skirt (3.6 dB/octave) compared to

the high-frequency side (9.3 dB/octave), although the latter was computed based only on the average

masked threshold at +2/3 octaves.

Figure 5 (Color online): Average MTPs (filled symbols) and AM detection thresholds (open symbols, replotted from Figure 3[6]) for the older HI and young NH listeners. The data for the older HI listeners are shown as circles connected by solid lines and the data for the young NH listeners (reproduced from Regev et al., 2023) are indicated by the squares connected by the dashed lines. Each panel represents the results for a specific target modulation frequency: 4 Hz (top left), 16 Hz (top right), 64 Hz (bottom left), and 128 Hz (bottom right). Note that the data from one of the older HI listeners was excluded from the 4-Hz panel for both the MTP and AM detection threshold. Means and standard errors are shown. Stars indicate the level of statistical significance of the differences in masked thresholds btween the two groups (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

Figure 5 shows the mean MTPs obtained from the older HI (circles) and the younger NH listeners (squares; reproduced from Regev *et al.*, 2023). Each panel shows the MTPs for a specific target modulation frequency. The mean AM detection thresholds are also shown and indicated by the open

symbols (replotted from Figure 3[6]). The statistical analysis comparing the older HI and the young NH

listeners revealed no main effect of group for any of the target modulation frequencies. However, a

significant interaction between listener group and masker-modulation center frequency was found at

all target modulation frequencies [$F_{6,108} = 3.95, p = 0.001$; $F_{6,114} = 8.1, p < 0.001$; $F_{5,95} =$

$4.12, p = 0.002$; $F_{4,76} = 6.3, p < 0.001$, for 4, 16, 64, and 128 Hz, respectively]. At the target

modulation frequency of 4 Hz, the post-hoc analysis revealed that the older HI listeners had

significantly higher (worse) masked thresholds than the young NH listeners for the masker

modulations centered at +2 octaves relative to the target. At 16 Hz, the post-hoc analysis revealed that

the older HI listeners had significantly lower (i.e., better) masked thresholds for the masker

modulations centered at -2 octaves relative to the target frequency, and significantly higher masked

thresholds than the young NH listeners for masker modulations centered at +2/3 octaves above

relative to the target frequency. At 64 Hz, the post-hoc analysis revealed that the older HI listeners

had significantly lower masked thresholds than the young NH listeners for the masker modulations

centered at +2/3 octaves relative to the target frequency. Finally, at 128 Hz, the post-hoc analysis

revealed that the older HI listeners had significantly lower masked thresholds than the young NH

listeners for the masker modulations centered at the target modulation frequency. The details of the

post-hoc analyses can be found in Appendix B.

Figure 6 (Color online): Average MTPs (filled symbols) and AM detection thresholds (open symbols,

replotted from Figure 3[6]). The data for the young and older HI listeners (solid lines) are indicated by

the diamonds and circles, respectively. The data for the young NH listeners (reproduced from Regev

et al., 2023) are shown as dashed lines and squares. Each panel represents the results for a specific

target modulation frequency: 4 Hz (top left), 16 Hz (top right), 64 Hz (bottom left), and 128 Hz

(bottom right). Note that the data from one of the older HI listeners was excluded from the 4-Hz

panel for both the MTP and AM detection threshold. Means and standard errors are shown.

Figure 6 shows the mean MTPs for both the young HI (diamonds) and the older HI listeners

(circles) as well as the data from the young NH listeners (squares) as obtained in Regev *et al.* (2023).

Each panel shows the MTPs for a specific target modulation frequency. The mean AM detection

thresholds are also shown and indicated by the open symbols (replotted from Figure 3[6]).

Similar to the MTPs obtained from the older HI listeners, high levels of asymmetry were also observed in the MTPs of the young HI listeners. At the target modulation frequency of 4 Hz, this was reflected by a steeper slope of the low-frequency skirt (4.8 dB/octave) than the high-frequency skirt (3.3 dB/octave). A similar observation was made at 16 Hz, with a steeper slope of the low-frequency skirt (6 dB/octave) than that of the high-frequency skirt (5 dB/octave). At 64 Hz, the young HI listeners displayed a much shallower low-frequency skirt (5.6 dB/octave) compared to the high-frequency side (10.6 dB/octave), although the latter was computed based only on the average masked threshold at +2/3 octaves.

Table 1 summarizes the overall Q-factors (representing the ratio of the filter's characteristic frequency to its bandwidth estimated at the 3-dB-down point) fitted to the average data of each listener group and indicates the fits' mean square errors (MSEs, in dB) in parentheses. The table also shows the ratio of the fitted Q-factors for each group to the Q-factor fitted for the young NH listeners, thus providing a comparison to the baseline represented by the young NH listeners. A ratio higher than 1 indicates sharper tuning for the listener group in question as compared to the young NH listeners, and vice-versa. For the older NH listeners, the reduced Q-factor ratio reflects the age-related reduction of AM frequency selectivity already shown by Regev *et al.* (2023). In contrast, the results for the older HI listeners suggest a sharper tuning for this group as compared to the older NH listeners at the modulation frequencies of 4, 16, and 64 Hz, and reduced AM frequency selectivity at 128 Hz. Compared to the young NH listeners, the results for the older HI listeners show a similar, or slightly greater, AM frequency selectivity at the modulation frequencies of 16 and 64 Hz, and a broader tuning at 4 and 128 Hz. The young HI listeners' results suggest generally greater AM frequency selectivity compared to the older HI listeners, and even higher Q-factors (i.e., sharper tuning) than the young NH listeners at the modulation frequencies of 4, 16, and 64 Hz.

398   Table 1: Overall fitted Q-factors, defined as the ratio of the filter's characteristic frequency to its

399   bandwidth estimated at the 3-dB-down point, and MSEs for the modulation filters fitted to the average

400   group data using an asymmetric linear fitting approach. The table also shows the ratio of fitted Q-

401   factors for each group to the Q-factor fitted for the young NH listeners (representing the baseline).

Target modulation frequency [Hz]	Estimated "Q-factor" *(Overall MSE [dB])*				Q/Q_{YNH}		
	YNH	**ONH**	**YHI**	**OHI**	**ONH**	**YHI**	**OHI**
4	0.77 *(0.35)*	0.10 *(0.16)*	0.81 *(0.78)*	0.14 *(0.30)*	0.13	1.05	0.18
16	0.96 *(0.14)*	0.75 *(0.44)*	1.24 *(0.38)*	1.13 *(0.30)*	0.78	1.29	1.18
64	1.38 *(0.15)*	1.14 *(0.09)*	1.90 *(1.0)*	1.45 *(0.14)*	0.83	1.38	1.05
128	0.96 *(0.05)*	0.40 *(0.10)*	0.82 *(0.98)*	0.27 *(0.96)*	0.42	0.85	0.28

IV. DISCUSSION

A. Summary of main findings

This study aimed to investigate the potentially 'antagonistic' effects of hearing loss and age on AM frequency selectivity. These opposing effects were suggested based on recent findings of an age-related reduction in AM frequency selectivity (Regev et al., 2023), which appeared to contradict previous conclusions that neither age nor hearing loss affect AM frequency selectivity (Takahashi and Bacon, 1992; Sek *et al.*, 2015). Behavioral MTPs and unmasked AM detection thresholds were collected from three young and ten older HI listeners (24-30 and 63-77 years), and the results were compared to the data obtained from young and older NH listeners in Regev *et al.* (2023). Both HI groups demonstrated lower (i.e., better) AM detection thresholds compard to their similarly aged NH counterparts, and the older HI listeners also exhibited lower detection thresholds than the young NH listeners for target modulation frequencies above 4 Hz (Figure 3). Furthermore, for target modulation frequencies below 128 Hz, the older HI listeners generally showed greater AM frequency selectivity than the older NH listeners (Figure 4). When compared to the young NH listeners, the older HI listeners displayed an asymmetric broadening of the MTP at the target modulation frequency of 4 Hz (with a shallower high-frequency skirt), but similar or even slightly sharper tuning at 16 and 64 Hz (Figure 5). At the target modulation frequency of 128 Hz, the older HI listeners showed lower masked thresholds overall compared to the older NH listeners (with an average offset of approximately 3.4 dB) and similar masked thresholds to the young NH listeners for the off-target maskers, effectively indicating reduced AM frequency selectivity compared to both NH groups. The data from the young HI listeners suggested greater AM frequency selectivity compared to the older HI listeners, as well as generally sharper tuning compared to the young NH listeners (Figure 6). Notably, both the young and older HI listeners exhibited high levels of asymmetry in their MTPs, with generally steeper low-frequency skirts compared to the high-frequency side. Finally, the best-fitting Q-factor obtained using a simple linear

fitting approach supported the previous observations and suggested that AM frequency selectivity in

HI listeners is similar to, or greater than, that in similarly aged NH listeners, except at the highest

target modulation rate of 128 Hz (Table 1).

B. Antagonistic effects of hearing loss and age

The results obtained from both the AM detection and MTP tasks suggest a general perceptual

benefit from hearing loss, in terms of an increased detectability of the target modulation and a

sharpening of the masked patterns. This benefit effectively counteracts the detrimental effects of age

on AM detection and AM frequency selectivity. The improvement of AM detection thresholds in HI

listeners compared to NH listeners is consistent with previous studies (e.g., Bacon and Gleitman, 1992;

Moore and Glasberg, 2001; Sek *et al.*, 2015; Schlittenlacher and Moore, 2016; Wallaert *et al.*, 2017;

Wiinberg *et al.*, 2019). It has been suggested that this improvement is due to an effectively increased

internal representation of the signal's AM depth resulting from the loss of compression commonly

associated with SNHL (Moore *et al.*, 1996; Jennings *et al.*, 2018). The trends observed in the data from

the young HI listeners support the notion that age is detrimental to AM detection whereas SNHL

provides a perceptual benefit. Conversely, previous AM masking studies comparing young NH and

older HI listeners at low target modulation frequencies ($\leq$ 16 Hz) found no differences in MTPs

(Takahashi and Bacon, 1992; Sek *et al.*, 2015) and concluded that neither SNHL nor age affect AM

frequency selectivity. The present results are broadly consistent with the observation that MTPs are

similar for young NH and older HI listeners at low modulation frequencies, although an asymmetric

broadening of the 4-Hz MTP was found for the older HI listeners, with a shallower high-frequency

skirt. However, the collected data also showed that SNHL generally improves AM frequency

selectivity for listeners of similar age at target modulation frequencies below 128 Hz. In contrast, the

data obtained at 128 Hz consistently showed lower masked thresholds for the older HI listeners

compared to the older NH listeners, indicating a reduction in AM frequency selectivity. Additionally,

a slight broadening of the pattern at 128 Hz was observed for young HI listeners compared to the

young NH group.

The results of the present study reveal that young listeners generally perform better than older

listeners with the same hearing status, as already shown by Regev *et al.* (2023). Furthermore, although

a sharpening of the MTPs in HI listeners was not observed on each skirt of the patterns, HI listeners

generally outperformed similarly aged NH listeners in terms of AM detection thresholds and tuning

of the MTPs.

C. Potential mechanism underlying the perceptual benefit of hearing loss

The underlying auditory processes that account for the sharper MTPs observed in the older HI

listeners compared to older NH listeners may not be straightforward. It seems unlikely that SNHL

would improve AM frequency selectivity by 'restoring' the bandwidth of the hypothetical modulation

filters. Instead, the greater AM frequency selectivity observed here in the HI listeners may be an

'apparent' effect resulting from the reduced peripheral compression which is typically associated with

SNHL. One notable characteristic of the data obtained from the HI listeners is the strong asymmetry

of the MTPs, with a steeper slope on the low-frequency skirt compared to the high-frequency side.

This observation was also made by Regev *et al.* (2023) regarding their NH listener data, although the

effect was less prominent than in the present study. The asymmetry suggests that an additional cue

for the detection of the target may be available when the modulation maskers are centered below -

but not above - the target modulation frequency. The existence of such a cue was discussed in previous

studies by Bacon and Grantham (1989) and Strickland and Viemeister (1996), who observed negative

masking when an AM target was masked by low-frequency AM maskers relative to the target

modulation frequency. They argued that listeners may be able to detect the target modulation in the

temporal dips of the masker modulation, or 'listen in the dips', which is a detection cue analogous to

the release from masking provided by envelope fluctuations in spectral masking tasks (e.g., Buus,

1985). In the present study, the random dips occurring in the modulation masker would be longer

than (or close to) one cycle of the target for masker modulations with center frequencies below the

target, while this would not be the case for modulation maskers centered above the target.

Consequently, it is possible that these local temporal features in the stimulus aid the detection of the

target in the presence of low-frequency maskers. Furthermore, the data suggest that such a cue, if

present, may be more salient for HI listeners than for NH listeners. The loss of cochlear compression

in HI listeners might increase the target's AM depth glimpsed in the temporal valleys of the masker,

further enhancing the detectability of the target. This notion is also consistent with the findings of

Wallaert *et al.* (2017), which showed a greater improvement in AM detection thresholds for HI listeners

compared to NH listeners when increasing the numbers of cycles of the target modulation, suggesting

an enhanced temporal integration in the HI listeners. This may imply that HI listeners require fewer

cycles of the target modulation to detect its presence, potentially contributing to the improved

performance measured in the lower skirt of the MTPS.

If this detection mechanism indeed mediated the sharper MTPs in HI listeners, investigating AM

frequency selectivity using an AM forward-masking paradigm should eliminate the perceptual benefit

of SNHL. AM forward masking has been used as an alternative approach to investigate AM frequency

selectivity (Wojtczak and Viemeister, 2005; Moore *et al.*, 2009; Füllgrabe *et al.*, 2021a, 2021b) because

it avoids extraneous cues resulting from the interaction between target and masker modulations. While

it remains unclear whether AM simultaneous and AM forward-masking paradigms reveal the same

underlying AM frequency-selective process, a study comparing age-matched NH and HI listeners

using such a paradigm could provide valuable insights into the origins of the observed asymmetry.

However, such a study has not been conducted thus far. Additionally, computational auditory models

could be employed to assess the plausibility of 'listening in the dips' as a mechanism underlying the

present results.

The hypothesis that increased sensitivity to the temporal envelope, resulting from the loss of

cochlear compression, offers a perceptual advantage for HI listeners is also supported by the findings

of Bianchi *et al.* (2016). Their study demonstrated that pitch-discrimination performance for

unresolved complex tones was comparable between older HI and young NH listeners when the

stimulus envelope could be effectively amplified through reduced compression (i.e., when it was

maximally peaky). However, performance was worse for the older HI listeners in other cases. In

addition to, or partially connected with, the loss of cochlear compression, the better-than-normal AM

detection thresholds and AM frequency selectivity observed in the HI listeners may also be related to

the increased neural coding of the envelope which has been shown both in animal models and humans

(Kale and Heinz, 2010, 2012; Henry *et al.*, 2014; Zhong *et al.*, 2014; Millman *et al.*, 2017; Goossens *et*

al., 2018; Decruy *et al.*, 2020). This increased internal representation of the envelope, while it may

reflect a listening advantage in simple AM detection and masking paradigms such as the ones

considered in the present study, has been suggested to ultimately be detrimental to speech intelligibility

in noise (Millman *et al.*, 2017; Goossens *et al.*, 2018; Decruy *et al.*, 2020). Hence, it could potentially

contribute to the difficulties often experienced by HI listeners in complex listening scenarios.

D. Limitations of the study

The presentation levels used in the present study were not directly comparable to those employed

by Regev *et al.* (2023) with NH listeners. Regev *et al.* used a fixed 65 dB SPL (corresponding to average

levels of 67 and 57 dB SL for the young and older NH listeners, respectively), while the present study

adjusted the level for the HI listeners to be at 30 dB SL (corresponding to 79 dB SPL on average),

following a similar approach to Sek *et al.* (2015). This methodological difference implies that the

presentation level was both lower in terms of SL (by 27-37 dB) and higher in terms of SPL (by 14 dB)

for the HI listeners compared to the NH listeners. Previous studies have highlighted the effect of the

interaction between presentation level and hearing status on AM detection thresholds, often finding

lower AM detection thresholds for HI listeners than for NH listeners at equal SL, but similar

thresholds at equal SPL (Bacon and Gleitman, 1992; Moore and Glasberg, 2001; Füllgrabe et al., 2003;

Schlittenlacher and Moore, 2016; Wallaert et al., 2017; Moore et al., 2019), although some studies still

found differences at equal SPL (Sek et al., 2015; Jennings et al., 2018). Hence, it is possible that the

present results were influenced by the difference in presentation levels across the two listener groups.

However, the present finding that the HI listeners showed lower AM detection thresholds than the

NH listeners at both lower SLs and higher SPLs seems consistent with previous studies that reported

similar results (Füllgrabe et al., 2003; Jennings et al., 2018). Füllgrabe et al. (2003) observed an increased

difference in AM detection thresholds between NH and HI listeners when the experiment was

repeated with a fixed SL for all participants. Additionally, Wojtczak (2011) investigated the effects of

carrier level on AM frequency selectivity with NH listeners and found that a reduction in SL led to an

increase in AM detection thresholds and a broadening of the MTPs. Therefore, it is likely that

investigating NH and HI listeners at equal SL would have further increased the observed differences

between the two groups in the present study. However, it remains unclear whether lower AM

detection thresholds and sharper MTPs would still be found in the HI listeners compared to the NH

listeners if they were tested at equal SPL.

The small number of young HI listeners included in this study, along with the significant variability

in their audiograms (except for the frequency of interest), limited the conclusions that could be drawn

from their data. Despite the small group size, the results from the young HI listeners in this study may

offer some valuable insights. Indeed, the data obtained across the three young HI listeners were

consistent with the observations made based on the other listener groups. Specifically, young listeners

exhibited sharper tuning than older listeners with the same hearing status, and HI listeners generally

demonstrated sharper tuning compared to similarly aged NH listeners. Conducting further

investigations with a larger sample of young HI listeners may help validate the trends identified in the

present study.

Finally, the method used to quantify AM frequency selectivity in the present study was limited by

the pronounced asymmetry observed in the MTPs. Previous studies quantified the tuning of the

hypothetical modulation filters using the EPSM (Ewert and Dau, 2000; Ewert *et al.*, 2002; Regev *et al.*,

2023), which assumes symmetrical filters centered around the characteristic modulation frequency.

However, the present results do not support this assumption, indicating the need for revising the

EPSM to incoporate more flexible filter shapes. Similar conclusions were drawn by Regev *et al.* (2023),

who observed large variability in individual MTPs. Consequently, the Q-factor values reported in this

study for the young and older NH listeners differ from those reported by Regev *et al.* (2023) due to

the different approaches used to quantify AM frequency selectivity. However, the overall finding of

lower Q factors for the older NH listeners compared to the young NH listeners remains consistent

between the two investigations.

V.   CONCLUSION

The present study investigated the effects of hearing loss on AM frequency selectivity by

employing a simultaneous AM masking task. Ten older and three young HI listeners were included in

the study, ranging in age from 63-77 and 24-30 years, respectively. These age ranges were

approximately matched to the young and older NH listeners tested in the study conducted by Regev

et al. (2023). The findings revealed that HI listeners exhibited lower AM detection thresholds and

generally sharper MTPs compared to the NH listeners, suggesting that hearing loss may counteract

the detrimental effects of age on AM frequency selectivity. The MTPs showed strong levels of

asymmetry, with generally steeper low-frequency skirts. A potential mechanism proposed to explain

these results was the detection of the target modulation in the dips of low-frequency masker

modulations, coupled with the effective increase in AM depth resulting from the loss of cochlear

compression. This suggests that the observed sharpening of the MTPs in HI listeners may be linked

to the loss of cochlear compression rather than a 'restoration' of the bandwidth of underlying

hypothetical modulation filters. Although differences in presentation levels between the HI and NH

listeners may have influenced the threshold differences, the results of this study motivate further

investigations into the effects of hearing loss on AM masking and AM frequency selectivity, as well as

their interaction with age-related effects. Furthermore, in addition to the loss of cochlear compression,

the lower AM detection thresholds and greater AM frequency selectivity observed in HI listeners may

be associated with an increased neural coding of the envelope, which has been suggested to be

detrimental to speech intelligibility in complex listening scenarios. Hence, future studies examining

the effects of hearing loss on AM masking can offer valuabe insights into the underlying mechanisms

that contribute to the perceptual benefits observed in HI listeners and their implications for complex

auditory perception and speech intelligibility.

ACKNOWLEDGMENTS

The authors thank Borgný Súsonnudóttir Hansen for her contribution to the data collection, as

well as Erik Kjærbøl and Jesper Borchorst Yde for their precious help in recruiting young HI

participants. The authors also thank Stephan Ewert for helpful suggestions and for kindly sharing

experimental details and data from earlier publications, Jens Hjortkjær and Jonatan Märcher-Rørsted

for kindly providing their implementations of the reverse digit span test and of the Cambridge formula

(CamEQ; Moore and Glasberg, 1998), as well as Raul Sanchez-Lopez and Beverly Wright for helpful

insights and discussions. This study was supported by the Oticon Centre of Excellence for Hearing

and Speech Sciences (CHeSS).

AUTHOR DECLARATIONS

Conflict of Interest

The authors have no conflict of interest to report.

Ethics Approval

All participants gave written informed consent and were financially compensated for their time.

Ethical approval for the study was provided by the Science Ethics Committee of the Capital Region

in Denmark (reference H-16036391).

DATA AVAILABILITY

The data that support the findings of this study are available from the corresponding author upon

request.

APPENDIX

A. Experimental protocol

Table 2 summarizes the experimental protocol used for each HI listener (both young and older).

Modulation rate discrimination thresholds were also collected, in addition to experiments described

above, but the results were not considered in the present study.

Table 2: Experimental protocol implemented for all HI listeners, including the tests of modulation

rate discrimination (MRD), masked threshold pattern (MTP), hearing threshold at the carrier

frequency (HT), and reverse digit span (RDS).

Session	*Test*	*Order*
1	HT followed by MRD	Fixed within session
2-5	MTP 4 Hz	Random across sessions
	MTP 16 Hz	
	MTP 64 Hz	
	MTP 128 Hz followed by AM detection	
Any (2-5)	RDS	Randomly added to any session

B. Parameters and post-hoc analyses for masked-threshold patterns

Table 3 summarizes the post-hoc analysis results on the ANOVA comparing the MTPs for the

older HI and NH listeners (Figure 4).

Table 3: Summary of post-hoc analysis results on the MTPs comparing older HI and older NH

listeners, for each target modulation frequency. The effect sizes (M) and standard errors (SEs) are

reported for each pairwise comparison. The significant p-values are highlighted in bold.

Target frequency f_m [Hz]	Effective masker-modulation bandwidth [Hz]	Masker-modulation center frequencies [Hz]	Masker-modulation position [octaves re target f_m]	M [dB]	SE [dB]	Corrected p-value
4	1.4	1	-2	-4.6	1.1	**< 0.001**
		1.6	-4/3	-3.7		**0.001**
		2.5	-2/3	-2.2		**0.047**
		4	0	-1.3		0.237
		6.3	2/3	-0.8		0.472
		10.1	4/3	-1.6		0.147
		16	2	-1.7		0.134
16	5.6	4	-2	-4.5	0.8	**< 0.001**
		6.3	-4/3	-2.3		**0.003**
		10.1	-2/3	-1		0.196
		16	0	0.1		0.935

		25.4	2/3	-1		0.2
		40.3	4/3	-1		0.179
		64	2	-2.9		**< 0.001**
64	15.1	4	-4	-4.6	1.2	**< 0.001**
	22.3	16	-2	-3		**0.014**
		25.4	-4/3	-2.2		0.061
		40.3	-2/3	-1.7		0.152
		64	0	-1.4		0.249
		101.6	2/3	-3		**0.013**
128	27.2	4	-5	-4.9	1.2	**< 0.001**
	44.6	32	-2	-3.5		**0.005**
		50.8	-4/3	-2.7		**0.027**
		80.6	-2/3	-2.2		0.074
		128	0	-3.8		**0.003**

623   Table 4 summarizes the post-hoc analysis results on the ANOVA comparing the MTPs for the

624   older HI and young NH listeners (Figure 5).

626   Table 4: Summary of post-hoc analysis results on the MTPs comparing older HI and young NH

627   listeners, for each target modulation frequency. The effect sizes (M) and standard errors (SEs) are

628   reported for each pairwise comparison. The significant p-values are highlighted in bold.

Target frequency f_m [Hz]	Effective masker-modulation bandwidth [Hz]	Masker-modulation center frequencies [Hz]	Masker-modulation position [octaves re target f_m]	M [dB]	SE [dB]	Corrected p-value
4	1.4	1	-2	-0.2	1.1	0.873
		1.6	-4/3	-0.5		0.662
		2.5	-2/3	-0.4		0.747
		4	0	-1.4		0.24
		6.3	2/3	0.4		0.736
		10.1	4/3	1.7		0.15
		16	2	3.2		**0.008**
16	5.6	4	-2	-3.3	0.8	**< 0.001**
		6.3	-4/3	-1.4		0.081
		10.1	-2/3	-0.6		0.405
		16	0	0.7		0.371
		25.4	2/3	2		**0.013**
		40.3	4/3	1.1		0.16
		64	2	-0.8		0.304
64	15.1	4	-4	-1.7	1.1	0.126

		16	-2	-0.5		0.637
		25.4	-4/3	0.6		0.597
	22.3	40.3	-2/3	0.5		0.648
		64	0	-1.5		0.191
		101.6	2/3	-3.1		**0.008**
	27.2	4	-5	-0.8		0.434
128		32	-2	0.7		0.542
	44.6	50.8	-4/3	0.6	1.1	0.594
		80.6	-2/3	-1.1		0.302
		128	0	-3.9		**< 0.001**

REFERENCES

Bacon, S. P., and Gleitman, R. M. (**1992**). "Modulation Detection in Subjects With Relatively Flat Hearing Losses," Journal of Speech, Language, and Hearing Research, **35**, 642–653. doi:10.1044/jshr.3503.642

Bacon, S. P., and Grantham, D. W. (**1989**). "Modulation masking: Effects of modulation frequency, depth, and phase," J Acoust Soc Am, **85**, 2575–2580. doi:10.1121/1.397751

Bianchi, F., Fereczkowski, M., Zaar, J., Santurette, S., and Dau, T. (**2016**). "Complex-Tone Pitch Discrimination in Listeners with Sensorineural Hearing Loss," Trends Hear, **20**, 1–15. doi:10.1177/2331216516655793

Bisgaard, N., Vlaming, M. S. M. G., and Dahlquist, M. (**2010**). "Standard Audiograms for the IEC 60118-15 Measurement Procedure," Trends Amplif, **14**, 113–120. doi:10.1177/1084713810379609

643 Buus, S. (**1985**). "Release from masking caused by envelope fluctuations," Journal of the

644   Acoustical Society of America, **78**, 1958–1965. doi:10.1121/1.392652

645 Chabot-Leclerc, A., Jørgensen, S., and Dau, T. (**2014**). "The role of auditory spectro-temporal

646   modulation filtering and the decision metric for speech intelligibility prediction," J

647   Acoust Soc Am, **135**, 3502–3512. doi:10.1121/1.4873517

648 Chi, T., Ru, P., and Shamma, S. A. (**2005**). "Multiresolution spectrotemporal analysis of complex

649   sounds," J Acoust Soc Am, **118**, 887–906. doi:10.1121/1.1945807

650 Christiansen, S. K., Jepsen, M. L., and Dau, T. (**2014**). "Effects of tonotopicity, adaptation,

651   modulation tuning, and temporal coherence in 'primitive' auditory stream segregation,"

652   J Acoust Soc Am, **135**, 323–333. doi:10.1121/1.4845675

653 Conroy, C., Buss, E., and Kidd, G. (**2023**). "Cues to reduce modulation informational masking,"

654   J Acoust Soc Am, **153**, 274–285. doi:10.1121/10.0016867

655 Conroy, C., Byrne, A. J., and Kidd, G. (**2022**). "Forward masking of spectrotemporal modulation

656   detection," J Acoust Soc Am, **151**, 1181–1190. doi:10.1121/10.0009404

657 Dau, T., Kollmeier, B., and Kohlrausch, A. (**1997a**). "Modeling auditory processing of amplitude

658   modulation. I. Detection and masking with narrow-band carriers," J Acoust Soc Am, **102**,

659   2892–2905. doi:10.1121/1.420344

660 Dau, T., Kollmeier, B., and Kohlrausch, A. (**1997b**). "Modeling auditory processing of amplitude

661   modulation. II. Spectral and temporal integration," J Acoust Soc Am, **102**, 2906–2919.

662   doi:10.1121/1.420345

663 Dau, T., Verhey, J., and Kohlrausch, A. (**1999**). "Intrinsic envelope fluctuations and modulation-

664   detection thresholds for narrow-band noise carriers," J Acoust Soc Am, **106**, 2752–2760.

665   doi:10.1121/1.428103

Dau, T., and Verhey, J. L. (**1999**). "Modeling Across-Frequency Processing Of Amplitude

Modulation," In T. Dau, V. Hohmann, and B. Kollmeier (Eds.), Psychophysics,

Physiology and Models of Hearing, World Scientific, Singapore, pp. 229–234.

doi:10.1142/9789812818140_0038

Decruy, L., Vanthornhout, J., and Francart, T. (**2020**). "Hearing impairment is associated with

enhanced neural tracking of the speech envelope," Hear Res, **393**, 107961.

doi:10.1016/j.heares.2020.107961

Depireux, D. A., Simon, J. Z., Klein, D. J., and Shamma, S. A. (**2001**). "Spectro-Temporal

Response Field Characterization With Dynamic Ripples in Ferret Primary Auditory

Cortex," J Neurophysiol, **85**, 1220–1234. doi:10.1152/jn.2001.85.3.1220

Elhilali, M., Chi, T., and Shamma, S. A. (**2003**). "A spectro-temporal modulation index (STMI)

for assessment of speech intelligibility," Speech Commun, **41**, 331–348.

doi:10.1016/S0167-6393(02)00134-6

Elhilali, M., Ma, L., Micheyl, C., Oxenham, A. J., and Shamma, S. A. (**2009**). "Temporal

Coherence in the Perceptual Organization and Cortical Representation of Auditory

Scenes," Neuron, **61**, 317–329. doi:10.1016/j.neuron.2008.12.005

Elliott, T. M., and Theunissen, F. E. (**2009**). "The Modulation Transfer Function for Speech

Intelligibility," (K. J. Friston, Ed.) PLoS Comput Biol, **5**, e1000302.

doi:10.1371/journal.pcbi.1000302

Erb, J., Schmitt, L.-M., and Obleser, J. (**2020**). "Temporal selectivity declines in the aging human

auditory cortex," Elife, **9**, 1–21. doi:10.7554/eLife.55300

Ewert, S. D. (**2013**). "AFC—A modular framework for running psychoacoustic experiments and

computational perception models," Proceedings of the International Conference on

Acoustics AIA-DAGA 2013, Merano, Italy, pp. 1326–1329.

Ewert, S. D., and Dau, T. (**2000**). "Characterizing frequency selectivity for envelope

fluctuations," J Acoust Soc Am, **108**, 1181–1196. doi:10.1121/1.1288665

Ewert, S. D., Verhey, J. L., and Dau, T. (**2002**). "Spectro-temporal processing in the envelope-

frequency domain," J Acoust Soc Am, **112**, 2921–2931. doi:10.1121/1.1515735

Fuglsang, S. A., Märcher-Rørsted, J., Dau, T., and Hjortkjær, J. (**2020**). "Effects of sensorineural

hearing loss on cortical synchronization to competing speech during selective attention,"

Journal of Neuroscience, **40**, 2562–2572. doi:10.1523/JNEUROSCI.1936-19.2020

Füllgrabe, C., Meyer, B., and Lorenzi, C. (**2003**). "Effect of cochlear damage on the detection of

complex temporal envelopes," Hear Res, **178**, 35–43. doi:10.1016/S0378-5955(03)00027-

3

Füllgrabe, C., Moore, B. C. J., and Stone, M. A. (**2015**). "Age-group differences in speech

identification despite matched audiometrically normal hearing: contributions from

auditory temporal processing and cognition," Front Aging Neurosci, **6**, 1–25.

doi:10.3389/fnagi.2014.00347

Füllgrabe, C., Sęk, A., and Moore, B. C. J. (**2021a**). "Frequency selectivity in the modulation

domain estimated using forward masking: Effects of masker modulation depth and

masker-signal delay," Hear Res, **405**, 108244. doi:10.1016/j.heares.2021.108244

Füllgrabe, C., Sęk, A., and Moore, B. C. J. (**2021b**). "Forward masking of amplitude modulation

across ears and its tuning in the modulation domain," J Acoust Soc Am, **149**, 1764–1771.

doi:10.1121/10.0003598

Goossens, T., Vercammen, C., Wouters, J., and van Wieringen, A. (**2018**). "Neural envelope

encoding predicts speech perception performance for normal-hearing and hearing-

impaired adults," Hear Res, **370**, 189–200. doi:10.1016/j.heares.2018.07.012

He, N., Mills, J. H., Ahlstrom, J. B., and Dubno, J. R. (**2008**). "Age-related differences in the

temporal modulation transfer function with pure-tone carriers," J Acoust Soc Am, **124**,

3841–3849. doi:10.1121/1.2998779

Henry, K. S., Kale, S., and Heinz, M. G. (**2014**). "Noise-induced hearing loss increases the

temporal precision of complex envelope coding by auditory-nerve fibers," Front Syst

Neurosci, **8**, 1–10. doi:10.3389/fnsys.2014.00020

Houtgast, T. (**1989**). "Frequency Selectivity In Amplitude-Modulation Detection," Journal of the

Acoustical Society of America, **85**, 1676–1680. doi:10.1121/1.397956

Jennings, S. G., Chen, J., Fultz, S. E., Ahlstrom, J. B., and Dubno, J. R. (**2018**). "Amplitude

modulation detection with a short-duration carrier: Effects of a precursor and hearing

loss," J Acoust Soc Am, **143**, 2232–2243. doi:10.1121/1.5031122

Jepsen, M. L., and Dau, T. (**2011**). "Characterizing auditory processing and perception in

individual listeners with sensorineural hearing loss," J Acoust Soc Am, **129**, 262–281.

doi:10.1121/1.3518768

Jepsen, M. L., Ewert, S. D., and Dau, T. (**2008**). "A computational model of human auditory

signal processing and perception," J Acoust Soc Am, **124**, 422–438.

Jørgensen, S., and Dau, T. (**2011**). "Predicting speech intelligibility based on the signal-to-noise

envelope power ratio after modulation-frequency selective processing," J Acoust Soc

Am, **130**, 1475–1487. doi:10.1121/1.3621502

Jørgensen, S., Ewert, S. D., and Dau, T. (**2013**). "A multi-resolution envelope-power based model

for speech intelligibility," J Acoust Soc Am, **134**, 436–446. doi:10.1121/1.4807563

Joris, P. X., Schreiner, C. E., and Rees, A. (**2004**). "Neural Processing of Amplitude-Modulated

Sounds," Physiol Rev, **84**, 541–577. doi:10.1152/physrev.00029.2003

Kale, S., and Heinz, M. G. (**2010**). "Envelope Coding in Auditory Nerve Fibers Following Noise-
Induced Hearing Loss," Journal of the Association for Research in Otolaryngology, **11**,
657–673. doi:10.1007/s10162-010-0223-6

Kale, S., and Heinz, M. G. (**2012**). "Temporal modulation transfer functions measured from
auditory-nerve responses following sensorineural hearing loss," Hear Res, **286**, 64–75.
doi:10.1016/j.heares.2012.02.004

King, A., Varnet, L., and Lorenzi, C. (**2019**). "Accounting for masking of frequency modulation
by amplitude modulation with the modulation filter-bank concept," J Acoust Soc Am,
145, 2277–2293. doi:10.1121/1.5094344

Kowalski, N., Depireux, D. A., and Shamma, S. A. (**1996**). "Analysis of dynamic spectra in ferret
primary auditory cortex. I. Characteristics of single-unit responses to moving ripple
spectra," https://doi-org.proxy.findit.cvt.dk/10.1152/jn.1996.76.5.3503, **76**, 3503–3523.
doi:10.1152/JN.1996.76.5.3503

Levitt, H. (**1971**). "Transformed Up-Down Methods in Psychoacoustics," J Acoust Soc Am, **49**,
467–477. doi:10.1121/1.1912375

Lorenzi, C., Micheyl, C., Berthommier, F., and Portalier, S. (**1997**). "Modulation Masking in
Listeners With Sensorineural Hearing Loss," Journal of Speech, Language, and Hearing
Research, **40**, 200–207. doi:10.1044/jslhr.4001.200

McDermott, J. H., Schemitsch, M., and Simoncelli, E. P. (**2013**). "Summary statistics in auditory
perception," Nat Neurosci, **16**, 493–498. doi:10.1038/nn.3347

McDermott, J. H., and Simoncelli, E. P. (**2011**). "Sound texture perception via statistics of the
auditory periphery: Evidence from sound synthesis," Neuron, **71**, 926–940.
doi:10.1016/j.neuron.2011.06.032

McWalter, R., and Dau, T. (**2015**). "Statistical representation of sound textures in the impaired

auditory system," Proceedings of the International Symposium on Auditory and

Audiological Research, 5, Nyborg, Denmark, pp. 186–196.

McWalter, R., and Dau, T. (**2017**). "Cascaded amplitude modulations in sound texture

perception," Front Neurosci, **11**, 1–12. doi:10.3389/fnins.2017.00485

Millman, R. E., Mattys, S. L., Gouws, A. D., and Prendergast, G. (**2017**). "Magnified neural

envelope coding predicts deficits in speech perception in noise," Journal of

Neuroscience, **37**, 7727–7736. doi:10.1523/JNEUROSCI.2722-16.2017

Moore, B. C. J., Füllgrabe, C., and Sek, A. (**2009**). "Estimation of the center frequency of the

highest modulation filter," J Acoust Soc Am, **125**, 1075–1081. doi:10.1121/1.3056562

Moore, B. C. J., and Glasberg, B. R. (**1998**). "Use of a loudness model for hearing-aid fitting. I.

Linear hearing aids," Br J Audiol, **32**, 317–335. doi:10.3109/03005364000000083

Moore, B. C. J., and Glasberg, B. R. (**2001**). "Temporal modulation transfer functions obtained

using sinusoidal carriers with normally hearing and hearing-impaired listeners," J Acoust

Soc Am, **110**, 1067–1073. doi:10.1121/1.1385177

Moore, B. C. J., and Oxenham, A. J. (**1998**). "Psychoacoustic consequences of compression in

the peripheral auditory system.," Psychol Rev, **105**, 108–124. doi:10.1037/0033-

295X.105.1.108

Moore, B. C. J., Shailer, M. J., and Schooneveldt, G. P. (**1992**). "Temporal modulation transfer

functions for band-limited noise in subjects with cochlear hearing loss," Br J Audiol, **26**,

229–237. doi:10.3109/03005369209076641

Moore, B. C. J., Wojtczak, M., and Vickers, D. A. (**1996**). "Effect of loudness recruitment on the

perception of amplitude modulation," J Acoust Soc Am, **100**, 481–489.

doi:10.1121/1.415861

Oetjen, A., and Verhey, J. L. (**2015**). "Spectro-temporal modulation masking patterns reveal

frequency selectivity," J Acoust Soc Am, **137**, 714–723. doi:10.1121/1.4906171

Oetjen, A., and Verhey, J. L. (**2017**). "Characteristics of spectro-temporal modulation frequency

selectivity in humans," J Acoust Soc Am, **141**, 1887–1895. doi:10.1121/1.4976537

Paraouty, N., Ewert, S. D., Wallaert, N., and Lorenzi, C. (**2016**). "Interactions between amplitude

modulation and frequency modulation processing: Effects of age and hearing loss," J

Acoust Soc Am, **140**, 121–131. doi:10.1121/1.4955078

Paraouty, N., and Lorenzi, C. (**2017**). "Using individual differences to assess modulation-

processing mechanisms and age effects," Hear Res, **344**, 38–49.

doi:10.1016/j.heares.2016.10.024

Piechowiak, T., Ewert, S. D., and Dau, T. (**2007**). "Modeling comodulation masking release using

an equalization-cancellation mechanism," J Acoust Soc Am, **121**, 2111–2126.

doi:10.1121/1.2534227

Regev, J., Zaar, J., Relaño-Iborra, H., and Dau, T. (**2023**). "Age-related reduction of amplitude

modulation frequency selectivity," J Acoust Soc Am, **153**, 2298–2311.

doi:10.1121/10.0017835

Relaño-Iborra, H., and Dau, T. (**2022**). "Speech intelligibility prediction based on modulation

frequency-selective processing," Hear Res, , doi: 10.1016/j.heares.2022.108610.

doi:10.1016/j.heares.2022.108610

Relaño-Iborra, H., May, T., Zaar, J., Scheidiger, C., and Dau, T. (**2016**). "Predicting speech

intelligibility based on a correlation metric in the envelope power spectrum domain," J

Acoust Soc Am, **140**, 2670–2679. doi:10.1121/1.4964505

Relaño-Iborra, H., Zaar, J., and Dau, T. (**2019**). "A speech-based computational auditory signal

processing and perception model," J Acoust Soc Am, **146**, 3306–3317.

doi:10.1121/1.5129114

Sabin, A. T., Eddins, D. A., and Wright, B. A. (**2012**). "Perceptual learning evidence for tuning

to spectrotemporal modulation in the human auditory system," Journal of Neuroscience,

32, 6542–6549. doi:10.1523/JNEUROSCI.5732-11.2012

Schlittenlacher, J., and Moore, B. C. J. (**2016**). "Discrimination of amplitude-modulation depth

by subjects with normal and impaired hearing," J Acoust Soc Am, **140**, 3487–3495.

doi:10.1121/1.4966117

Schoof, T., and Rosen, S. (**2014**). "The role of auditory and cognitive factors in understanding

speech in noise by normal-hearing older listeners," Front Aging Neurosci, **6**, 1–14.

doi:10.3389/fnagi.2014.00307

Sek, A., Baer, T., Crinnion, W., Springgay, A., and Moore, B. C. J. (**2015**). "Modulation masking

within and across carriers for subjects with normal and impaired hearing," J Acoust Soc

Am, **138**, 1143–1153. doi:10.1121/1.4928135

Sek, A., and Moore, B. C. J. (**2003**). "Testing the concept of a modulation filter bank: The

audibility of component modulation and detection of phase change in three-component

modulators," J Acoust Soc Am, **113**, 2801–2811. doi:10.1121/1.1564020

Singh, N. C., and Theunissen, F. E. (**2003**). "Modulation spectra of natural sounds and

ethological theories of auditory processing," J Acoust Soc Am, **114**, 3394–3411.

doi:10.1121/1.1624067

Steinmetzger, K., Zaar, J., Relaño-Iborra, H., Rosen, S., and Dau, T. (**2019**). "Predicting the

effects of periodicity on the intelligibility of masked speech: An evaluation of different

modelling approaches and their limitations," J Acoust Soc Am, **146**, 2562–2576.

doi:10.1121/1.5129050

Strickland, E. A., and Viemeister, N. F. (**1996**). "Cues for discrimination of envelopes," J Acoust

Soc Am, **99**, 3638–3646. doi:10.1121/1.414962

Takahashi, G. A., and Bacon, S. P. (**1992**). "Modulation Detection, Modulation Masking, and

Speech Understanding in Noise in the Elderly," Journal of Speech, Language, and

Hearing Research, **35**, 1410–1421. doi:10.1044/jshr.3506.1410

The   Mathworks   Inc.   (**2015**).   "MATLAB   2015b   -   MathWorks,"

www.mathworks.com/products/matlab. Last viewed January 1, 2016

Verhey, J. L., Dau, T., and Kollmeier, B. (**1999**). "Within-channel cues in comodulation masking

release (CMR): Experiments and model predictions using a modulation-filterbank

model," J Acoust Soc Am, **106**, 2733–2745. doi:10.1121/1.428101

Viemeister, N. F. (**1979**). "Temporal modulation transfer functions based upon modulation

thresholds," J Acoust Soc Am, **66**, 1364–1380. doi:10.1121/1.383531

Wallaert, N., Moore, B. C. J., Ewert, S. D., and Lorenzi, C. (**2017**). "Sensorineural hearing loss

enhances auditory sensitivity and temporal integration for amplitude modulation," J

Acoust Soc Am, **141**, 971–980. doi:10.1121/1.4976080

Wallaert, N., Moore, B. C. J., and Lorenzi, C. (**2016**). "Comparing the effects of age on amplitude

modulation and frequency modulation detection," J Acoust Soc Am, **139**, 3088–3096.

doi:10.1121/1.4953019

Wiinberg, A., Jepsen, M. L., Epp, B., and Dau, T. (**2019**). "Effects of Hearing Loss and Fast-

Acting Compression on Amplitude Modulation Perception and Speech Intelligibility,"

Ear Hear, **40**, 45–54. doi:10.1097/AUD.0000000000000589

Wojtczak, M. (**2011**). "The effect of carrier level on tuning in amplitude-modulation masking," J

Acoust Soc Am, **130**, 3916–3925. doi:10.1121/1.3658475

Wojtczak, M., and Viemeister, N. F. (**2005**). "Forward masking of amplitude modulation: Basic

characteristics," J Acoust Soc Am, **118**, 3198–3210. doi:10.1121/1.2042970

Zaar, J., and Carney, L. H. (**2022**). "Predicting speech intelligibility in hearing-impaired listeners

using a physiologically inspired auditory model," Hear Res, , doi:

10.1016/J.HEARES.2022.108553. doi:10.1016/J.HEARES.2022.108553

Zaar, J., and Dau, T. (**2017**). "Predicting consonant recognition and confusions in normal-hearing

listeners," J Acoust Soc Am, **141**, 1051–1064. doi:10.1121/1.4976054

Zaar, J., Schmitt, N., Derleth, R.-P., DiNino, M., Arenberg, J. G., and Dau, T. (**2017**). "Predicting

effects of hearing-instrument signal processing on consonant perception," J Acoust Soc

Am, **142**, 3216–3226. doi:10.1121/1.5011737

Zhong, Z., Henry, K. S., and Heinz, M. G. (**2014**). "Sensorineural hearing loss amplifies neural

coding of envelope information in the central auditory system of chinchillas," Hear Res,

309, 55–62. doi:10.1016/j.heares.2013.11.006

Zilany, M. S. A., and Bruce, I. C. (**2007**). "Predictions of Speech Intelligibility with a Model of

the Normal and Impaired Auditory-periphery," 2007 3rd International IEEE/EMBS

Conference on Neural Engineering, IEEE, pp. 481–485. doi:10.1109/CNE.2007.369714