EpiGePT: a Pretrained Transformer model for epigenomics

Zijing Gao[1,#], Qiao Liu[2,#,*], Wanwen Zeng[2], Wing Hung Wong[2,3,*] and Rui Jiang[1,*]

[1] Ministry of Education Key Laboratory of Bioinformatics, Research Department of Bioinformatics at the Beijing National Research Center for Information Science and Technology, Center for Synthetic and Systems Biology, Department of Automation, Tsinghua University, Beijing 100084, China;

[2] Department of Statistics, Stanford University, Stanford, CA 94305, USA;

[3] Department of Biomedical Data Science, Bio-X Program, Center for Personal Dynamic Regulomes, Stanford University, Stanford, CA 94305, USA;

[*] To whom correspondence should be addressed.

[#] The first two authors contributed equally.

E-mail: liuqiao@stanford.edu, whwong@stanford.edu, ruijiang@tsinghua.edu.cn

Abstract

The transformer-based models, such as GPT-3[1] and DALL-E[2], have achieved unprecedented breakthroughs in the field of natural language processing and computer vision. The inherent similarities between natural language and biological sequences have prompted a new wave of inferring the grammatical rules underneath the biological sequences. In genomic study, it is worth noting that DNA sequences alone cannot explain all the gene activities due to epigenetic mechanism. To investigate this problem, we propose EpiGePT, a new transformer-based language pretrained model in epigenomics, for predicting genome-wide epigenomic signals by considering the mechanistic modeling of transcriptional regulation. Specifically, EpiGePT takes the context-specific activities of transcription factors (TFs) into consideration, which could offer deeper biological insights comparing to models trained on DNA sequence only. In a series of experiments, EpiGePT demonstrates state-of-the-art performance in a diverse epigenomic signals prediction tasks as well as new prediction tasks by fine-tuning. Furthermore, EpiGePT is capable of learning the cell-type-specific long-range interactions through the self-attention mechanism and interpreting the genetic variants that associated with human diseases. We expect that the advances of EpiGePT can shed light on understanding the complex regulatory mechanisms in gene regulation. We provide free online prediction service of EpiGePT through https://health.tsinghua.edu.cn/epigept/.

Introduction

One of the fundamental problems in genomic study is how to decode and interpret the human genome sequences in a complex manner. Progress toward this goal is largely hindered by the vast majority of non-coding regions[3]. For example, it remains unclear how the genomic variants in the noncoding regions lead to malfunctions of regulatory elements by disrupting the underlying regulatory syntax of DNA[4]. Inspired from the field of natural language processing, there exists a natural analogy between human language and DNA sequence where texts are made of words and DNA sequence can be characterized by nucleotides or k-mers. The inherent similarities between natural language and biological sequences provide new perspectives towards better understanding the complex DNA language.

Recently, generative pre-trained transformer (GPT) models have achieved unprecedented success in various domains, including computer vision and natural language processing (NLP)[1, 5]. Such pre-trained models can be readily tailored or adapted to various downstream tasks. To date, the application of generative pre-trained models in genomic study remains largely unexplored. It is noticeable that a number of machine learning-based approaches have been proposed for predicting various genomic and epigenomic signals, such as chromatin accessibility[6, 7], histone modification[8] or chromatin interactions[9, 10]. However, these methods are rather scattered, with specific models designed for specific prediction tasks. It is in an urgent need to develop a foundation model to facilitate multiple genomic and epigenomic prediction tasks and unveil the universal gene regulation rules.

To design a genomic foundation model, it is worth noting that the existing large language

models (LLMs) purely rely on the language context consisting of words and sentences while the DNA sequences cannot explain all the heritable and stable changes in gene activity due to epigenetic mechanisms. In other words, the genomic foundation model based on pure DNA sequence may largely ignore the context-specific information, thus lacking mechanistic interpretation of context-specific gene regulation. For example, using transformer-based language model to decode genome sequence has been attempted by a recent work Enformer[11]. However, Enformer is not capable of predicting the function of sequences in new cellular contexts, which largely limits its generalization power.

To overcome the above limitation, we proposed EpiGePT, a new transformer-based deep learning framework, to predict genome-wide epigenomic signals by taking the mechanistic modeling of transcriptional regulation into consideration. With EpiGePT, we are able to investigate how to utilize the power of transformer-based language model to help researchers uncover how trans-regulatory factors (e.g., TFs) regulate target genes by interacting with cis-regulatory elements and further lead to changes in different chromatin states. After pretraining on a diverse panel of cell line and tissue level data from the Encode database[12], EpiGePT is able to directly predict the genome-wide chromatin states in any new cellular context given the expression profile of a few hundreds of TFs or facilitate new prediction tasks (e.g., 3D genome interaction) with finetuning.

To the best of our knowledge, EpiGePT is the first pretrained Transformer model for epigenomics with mechanistic modeling of transcriptional regulation. EpiGePT differs from existing methods in the following three aspects. First, unlike the methods that take pure DNA sequence as input, EpiGePT additionally takes the context-specific information (e.g., TF

activities) as input, thus enabling genome-wide prediction power in any new cellular context.

Second, instead of using task-specific model to predict a single genomic and epigenomic signal,

EpiGePT is designed for simultaneously predicting multiple epigenomic signals of the same

genomic region through multi-task learning, thus improving learning efficiency and prediction

accuracy compared to the task-specific models. Third, many methods typically take short DNA

sequence (e.g., a few hundred or a thousand base pair) as input, which may not be adequate to

capture the complex syntax of DNA due to truncation. The long input DNA sequence (e.g.,

128kb) for EpiGePT greatly enhances the ability for the model to capture the long-range

interaction in the genome, which are crucial for understanding the gene regulation mechanism.

In a series of experiments, we illustrate that our method is superior to existing methods in a

various tasks of chromatin states prediction, as well as the variant effect prediction. We also

show that the self-attention mechanism greatly helps unveil the complex code in the

conformation of long-range chromatin interactions, such as promoter-enhancer interactions and

promoter-silencer interactions. EpiGePT is an example of how transformer-based language

model and large-scale pretrain can be used in genomics research to provide biological insights.

With the help of EpiGePT model, it is expected that researchers can dissect the comprehensive

genomic regulatory code given the cellular context information and accelerate research findings

in genomic study.

Results

Overview of EpiGePT model

We developed a novel Transformer[13]-based language model named EpiGePT to predict multiple chromatin states across different cell types. EpiGePT is a language model for cross-cell-type prediction of chromatin states by multi-task learning based on genome-wide pre-training on epigenomic data (**Fig. 1 and** Fig. S6). EpiGePT is composed of four modules, including a sequence module, a TF module, a transformer module, and a prediction module. The sequence module is responsible for processing the long DNA sequence of interest (e.g., 128 kb) by employing a series of convolutional and pooling blocks (e.g., 5) to extract a comprehensive set of sequence features. By reducing the input length by $2^5=128$ times through pooling operations, this module effectively compresses the input information while retaining essential features. The TF module is specifically designed to extract cell-type-specific features by taking the expression of transcription factors in the given context, as well as their corresponding motif score into account. This module helps capture the unique characteristics of each cell type by considering the binding status of TFs involved in gene regulation. In the transformer module, each token corresponds to a genomic bin in the original DNA sequence and has hybrid features derived from both sequence and TFs. The module leverages self-attention mechanisms to learn the comprehensive relationships among the input bins, enabling the model to make predictions of multiple chromatin states under the given context cellular. By taking advantage of this approach, EpiGePT provides a powerful tool for predicting multiple chromatin states and enables researchers to gain insights into the underlying regulatory mechanisms of the genome.

EpiGePT enables genome-wide prediction of chromatin states

To assess the predictive performance for epigenomic signals of EpiGePT, specifically in predicting chromatin accessibility, a comprehensive evaluation was conducted. We first applied EpiGePT to predict the chromatin accessibility based on the widely available public DNase-seq[14] data across diverse cell types or tissues. In brief, DNase-seq data across 129 cell types were collected from the ENCODE[12] project. After data preprocessing and normalization (see Methods), 1,175,374 genomic regions were extracted where each pair of cell type and genomic regions constitutes a training instance. We meticulously devised comprehensive experimental settings by partitioning the training and test sets based on either genomic regions or cell types. In detail, we employed the following three data partitioning settings for a comprehensive evaluation (Fig. S1, Text S1). For "cross-cell-type" prediction, we partitioned the data into training and testing sets based on cell types. For "cross-region" prediction, we partitioned the data into training and testing sets based on the genomic regions. For "cross-both" prediction, we conducted rigorous data split to ensure that both the cell types and genomic regions in the test stage are unseen during the training process. We employed three evaluation metrics, namely Pearson correlation coefficient, Spearman correlation coefficient and prediction square error, to assess the similarity between the predicted and true values of DNase signals (See Methods). It is shown that EpiGePT consistently outperforms other competing methods, including Enformer[11], BIRD[15], and ChromDragoNN[16] by a relatively large margin under the above experimental settings (**Fig. 2A** and Fig. S2). EpiGePT achieves 5.0%, 8.9%, and 5.2 % higher performance than Enformer, the best baseline method, in terms of the mean Pearson correlation coefficient under three data-partition settings, respectively (**Fig. 2B**). Besides the chromatin

accessibility regression task, we also designed binary chromatin accessibility status prediction

task by assessing whether a peak exists within the corresponding genomic bin (>50% overlap).

We made slight adjustments to the regression model by modifying the activation function and

loss function to accommodate the binary classification task (See methods). The results show

that EpiGePT achieves an average auPRC (area under the precision-recall curve) of 0.767

compared to 0.727 of Enformer[11], 0.623 of DeepCAGE[17] and 0.476 of ChromDragoNN[16] (**Fig.**

2C).

Next, we extended the chromatin state prediction task from a single target to multiple targets

by predicting multiple chromatin states, including chromatin accessibility, CTCF[18], ChIP-seq,

and six types of different histone modifications[19] (See Methods). When considering eights

different chromatin states, only 28 cell types have the corresponding available data

simultaneously. After preprocessing, 13,300 genomic regions each with a length of 128 kbp

were extracted, which cover 56.7% of the whole genome. However, compared to the data in

the DNase-only prediction experiment, the correlation coefficient was reduced due to the

prevalence of a substantial number of zero signals in the genomic regions being predicted.

Using a similar data split strategy as the single target for cross-cell-type prediction, EpiGePT

demonstrated a mean Pearson correlation coefficient between 0.259 to 0.566 of different

chromatin states in the test cell types (**Fig. 2D**). Specifically, EpiGePT achieves remarkably

high performance in predicting chromatin state signals for certain cell types, such as the colon

tissue, with a Pearson correlation coefficient of 0.888. Furthermore, as it shown in Fig. S3C, it

significantly outperformed Enformer in terms of performance across these tested cell types and

different signals (one-side p-value < 2.79e-10 under binomial hypothesis test). To make the

chromatin states prediction task more illustrative, several tracks of predicted chromatin states

and the corresponding ground truth chromatin states were displayed. For instance, at the

position of from 61,056,000 to 61,184,000 on chromosome 20, we used the UCSC genome

browser[20] to show the predicted values and true values of CTCF (Pearson correlation coefficient

of 0.518) and DNase signals (Pearson correlation coefficient of 0.869), as well as the regulatory

relationships within this region. (Fig. S2A). In addition, we also compared EpiGePT with

ChromDragoNN[16] on binary and quaternary classification tasks based on ChromHMM[21]

annotations (Fig. S3). EpiGePT achieved an average auROC (area under the receiver operating

characteristic curve) of 0.855 in binary classification, significantly higher than that of

ChromDragoNN[16] (0.774). In quaternary classification, EpiGePT achieved a macro-auROC of

0.879, also significantly higher than ChromDragoNN[16] (0.856, one-side p-value < 0.001).

These results demonstrate the effectiveness and accuracy of EpiGePT in predicting multiple

chromatin states, leveraging the four modules. The effectiveness and prediction power achieved,

in conjunction with the self-attention mechanism, lays the foundation for deciphering

regulatory relationships.

To further verify the roles of the main modules in the model, we conducted the ablation

experiments on the model architecture. For TF module ablation, the above experimental results

compared to EpiGePT without TF module (EpiGePT-seq) and EpiGePT have demonstrated that

EpiGePT outperforms EpiGePT-seq in cross-cell-type prediction of DNase signals, with an

average Pearson correlation coefficient of 0.714 and a median of 0.74 for EpiGePT-seq, while

EpiGePT achieves 0.756 (average) and 0.787 (median). In addition, the inclusion of the TF

module enables EpiGePT to predict chromatin states at the locus level for different cell types.

However, like Enformer, EpiGePT-seq predicts the same values for different cell types at the

same locus, resulting in a zero correlation for cross-cell-type prediction. We examined the

impact of the TF module on multi-task prediction by employing three methods, namely

replacing TF scores with zero, adding random noise to TF, and removing motif binding scores.

The results indicated that when the expression of TFs was set to zero, the prediction of H3K27ac

yielded a Pearson correlation coefficient of 0.190. However, incorporating the TF module

significantly improved the coefficient to 0.543, demonstrating a beneficial impact of the TF

module.

For sequence module ablation, we randomly subsampled 10,000 genomic bins and 20 cell

types to train a TF-only model. The results indicated that removing the sequence

module resulted in an average decrease of 0.084 in the Pearson correlation coefficients

of the eight signals on a cell-type wise basis, and with a particularly significant decrease

of 0.13 in predicting H3K4me3 signals (Fig. S4A).

For multi-task module, we predicted the eight chromatin states involved in training

using eight individual models for single-task prediction. The results were evaluated on

a cross-cell type prediction manner. In the case of predicting the signal of H3K4me1,

the average Pearson correlation decreased from 0.408 to 0.329. When predicting the

H3K4me1 signal, the average Pearson correlation coefficient decreased from 0.408 to

0.329. Similarly, the overall prediction performance for the eight signals declined by

0.074 (**Fig. 4B**). This decrease may be attributed to the intricate nature of gene

regulation. The distinct chromatin states can complement and synergize with each other

through multi-task learning, allowing the model to gain deeper biological insights

compared to a single-target prediction model.

Furthermore, we performed additional experiments to investigate the effect of the

number of cell lines on the prediction performance. Specifically, we focused on DNase

predictions and randomly downsampled the training cell types from 103 to 75, 50, 25

for each of the five folds in the cross-validation experiment. The results demonstrated

a strong positive correlation between the number of cell lines and the prediction

performance. Under five-fold cross-validation, the median Pearson correlation

coefficient on the test set across 129 cell types decreased from 0.793 to 0.790, 0.761,

and 0.732, respectively. These findings suggest that our current model has potential

room for improvement and additional training with more cell lines will lead to even

better predictive performance, thereby offering more comprehensive insights into the

regulatory mechanisms for researchers (Fig. S4C). In summary, EpiGePT demonstrated

superior performance in predicting both single and multiple epigenomic signals over

existing methods, providing a robust foundation for decoding the complex landscape

of gene regulation.

EpiGePT facilitates long-range chromatin interaction identification

We examined the capacity of EpiGePT for predicting long-range chromatin interactions, which

play a critical role in preserving chromatin architecture and elucidating 3D contacts between

distal regulatory elements and target genes. Traditional methods typically take short DNA

sequence (e.g., 1kbp) as input, thus cannot take the long-range chromatin interactions into

consideration. During the training of EpiGePT model, the self-attention mechanism in the

transformer module plays an important role in capturing the potential interactions between

different DNA bins. We utilized the cell-type specific self-attention scores to predict chromatin

interactions, including enhancer-promoter and silencer-promoter interactions (see Methods).

We initially investigated whether EpiGePT can differentiate experimentally validated enhancer-

promoter interactions from other interactions. Two datasets containing 664 and 5,091 candidate

enhancer-promoter interactions or element-TSS interactions obtained by CRISPRi[22]

experiments were used and further filtered and stratified by the distance. In the Gasperini et

al[23]. dataset, EpiGePT consistently outperform EpiGePT-seq and Enformer by achieving the

highest auPRC in all groups. For instance, EpiGePT achieved auPRC of 0.949, 0.726, and 0.810

for identifying enhancer-promoter pairs in the 0-3 kbp, 3-20 kbp, and 20-64 kbp ranges,

respectively (**Fig. 3A and** Fig. S8). In the Fulco et al.[24] dataset, EpiGePT obtains better

performance than EpiGePT-seq in most groups, which illustrates the positive benefit of the cell-

type-specificity brought in the TF module. EpiGePT consistently outperforms Enformer across

different groups by a relatively large margin. As shown in Fig. 3A, EpiGePT achieves an auPRC

of 0.618, compared to 0.531 of Enformer, and 0.568 of EpiGePT-seq in 0-12kbp group.

Next, we explored whether EpiGePT is also capable of predicting the promoter-silencer

interactions. Since there is very limited experiment-validated silencer-promoter interactions,

we downloaded putative silencers from the SilencerDB[25] and used the promoter of annotated

nearest gene as the potential target. For negative silencer-promoter pairs, we selected the same

promoter and equidistant genomic regions in the opposite direction to ensure the consistency

of distance distributions between positive and negative sample pairs at different distance levels.

As a result, EpiGePT achieves a better performance in discerning positive silencer-promoter

pairs from negative pairs than Enformer by a relatively large margin. For instance, EpiGePT

displays an auROC of 0.575 in long-range interactions (32-64kbp) with positive-to-negative

ratio 1:1 setting, compared to 0.547 of EpiGePT-seq, and 0.483 of Enformer (**Fig. 3B**).

According to these results, the self-attention mechanism significantly enhances the ability to

identify potential chromatin interactions and increases the interpretability of the model.

The HiChIP[26] sequencing technology provides unprecedented opportunities to uncover 3D

genomic interactions. We aim to investigate the predictive performance of EpiGePT on 3D

genome interaction based on HiChIP data. Here, we employ the same strategy as described

above to calculate attention scores for the regulatory element-promoter pairs and collected

HiChIP loops on K562 and GM12878 cell lines from the HiChIPdb[27]. The results demonstrate

that incorporating TF expression data into EpiGePT leads to enhanced predictive performance

for HiChIP loops compared to the pure sequence models Enformer and EpiGePT-seq, across

diverse distance ranges and in two distinct cell lines. Specifically, within the 20-40 kbp distance

range on K562 cell line, EpiGePT achieves an auROC of 0.599 for the 1:1 positive-to-negative

ratio, surpassing Enformer's performance of 0.545 (**Fig. 3E**). These findings suggest that, even

without any fine-tuning, EpiGePT's attention scores encompass more accurate and

comprehensive biological information, underscoring its potential for capturing intricate

genomic interactions.

To better understand the self-attention mechanism of EpiGePT and bridge the gap between the

model and its interpretability, we visualized the attention matrices after normalization (**Fig. 3C**).

The visualization shows prominent scores between certain genomic bins, indicating the

potential presence of interactions. We centered on the transcription start site (TSS) of the CHD4

gene and calculated the self-attention scores between the genomic bins within its upstream and

downstream 128kbp. The attention scores exhibited peaks near the regulatory elements in the

vicinity of the TSS, which further validates the feasibility and accuracy of our prediction of

enhancer-promoter interactions (**Fig. 3D**).

EpiGePT improves variant effect prediction

One of the most essential tasks for EpiGePT is to dissect the effect of genetic variants that occur

in different genomic regions. As most of the variants identified by the GWAS studies lie in the

non-coding regions of the genome, which makes it difficult to interpret the variant effect, most

sequence-based computational models directly take the alleles sequence as input and compare

the difference in the predicted regulatory activity. The advantage of EpiGePT model comes

from the TF module where variant effect can be estimated under any given cellular context.

This is extremely helpful when predicting the effect of the disease- or phenotype-associated

SNPs. To test the ability of EpiGePT in variant effect prediction, we first collected an eQTLs

dataset[28] that contains 20,913 causal and non-causal variant-gene pairs in total across 49

different tissues from the supplementary data of Wang et al[28]. EpiGePT and EpiGePT-seq are

then applied to estimate the log-odds scores (LOS) given both the reference and alterative DNA

sequence and the corresponding relevant TF profile (see Methods, **Fig. 4A**). Finally, a random

forest classifier is trained based on the LOS scores across different chromatin states. The

experimental results show that in the lung tissue, EpiGePT demonstrates and auPRC of 0.922,

compared to 0.873 of Enformer in distinguishing casual SNPs. To verify the effectiveness of

TF module, we replace the TF profile of lung with stomach, which is much less relevant to the

lung tissue. The auPRC decreases from 0.922 to 0.892 (**Fig. 4B**). Similarly, EpiGePT-seq can

achieve an average auPRC of 0.910, compared to 0.898 of Enformer using 5-fold cross-validation for predicting causal variants on 48 extracted tissues (Fig. S3D). In the adrenal gland tissue, EpiGePT-seq demonstrates an average auPRC of 0.883, compared to only 0.842 of Enformer. The above experiments show the predictive power of EpiGePT in estimating the variant effect.

To further evaluate the performance of EpiGePT in predicting disease-associated variants, we extracted 52, 876 pathogenic SNPs from the ClinVar[29] database and 418, 863 benign SNPs from the ClinVar database, also with 84, 095 benign SNPs from the ExAC database[30] as positive and negative sets, respectively. We defined a 64kbp region surrounding each pathogenic SNP as the risk region. We screened all benign and likely benign SNPs that fall within the risk region from the negative sets for classification. As the relevant tissue or cell type information is not available, we concatenated the LOS of the eight epigenomic signals and self-attention scores across 28 cell types into a single 252-dimensional vector and then train a classifier for predicting whether the given SNP is pathogenic (see Methods). To assess the whether the 252-dimensional features are beneficial in predicting pathogenic SNPs, we concatenated it with 52 annotations from CADD[31], resulting in a comprehensive feature vector. Subsequently, we compared the performance of this combined feature vector with that of the individual features derived exclusively from CADD. We then utilized these two sets of features to train multi-layer perceptron (MLP) classifiers separately. The results demonstrate that incorporating EpiGePT's variant effect features from multiple cell types significantly enhances the performance of the classifier in predicting pathogenic SNPs. Specifically, when the positive-to-negative sample ratio was set to 1:1, the average auROC increased from 0.772 to 0.806, and the average accuracy

increased from 0.690 to 0.723 (**Fig. 4C**). This observation indicates that features extracted by

EpiGePT provide a valuable complement to CADD annotation, enabling a more comprehensive

depiction of variant characteristics, and thereby facilitating the discovery of disease-associated

variants.

EpiGePT prioritizes potential SNPs associated with comorbidities of COVID-19

We investigated whether the ability of EpiGePT to predict variant effects could help in the

discovery of key SNPs related to COVID-19. COVID-19 is an infectious disease caused by the

SARS-CoV-2 virus, which emerged in late 2019 and quickly spread around the world, causing

a global pandemic[32]. In order to validate the ability of EpiGePT in identifying key SNPs, we

collected GWAS data from the COVID-19 host genetics[33], including 9,484 variants. These

variants were derived from 4,933 patients with confirmed severe respiratory symptoms and

1,398,672 control individuals without COVID-19 symptoms. To validate the ability of the

model to identify COVID-19-related SNPs, we firstly defined a risk region around the selected

COVID-19-related SNPs and computed the rank of the variant score of pathogenic SNPs within

the surrounding benign SNPs from the ClinVar database. The expected rank for random

guessing (uniform distribution) is 0.5. Interestingly, we found that the average rank of COVID-

19-related SNPs was significantly lower than 0.5 across several tissues or cell types (**Fig. 4D**).

For instance, when lung expression data was employed and a 6-kbp risk region was examined,

the median rank was 0.250, and when expression data of esophagus squamous epithelium was

used the median rank was 0.333, significantly lower than 0.5 (one-side p-value of 0.013 under

one-sided Binomial Test). However, when we employed the expression data from smooth

muscle cells, which are a more widespread cell type with lower relevance to COVID-19, the

median rank exhibited a notable decrease to 0.381. Notably, when focusing on the 40-kbp risk

region, the median rank further declined to 0.850, higher than 0.5. These findings suggest that

EpiGePT model is able to prioritize the COVID-19-related SNPs thus shedding lights on

finding the potential disease-associated variants with our pretrained large language model.

Next, based on the aforementioned findings, we aimed to use EpiGePT to identify genes that

are highly related to COVID-19. Since the genetic pathology of COVID-19 is not yet clear and

the earliest lesion is in the lungs, we ranked all 9,484 possible SNPs using lung expression data.

We then identified the SNPs with the highest ranks and performed gene ontology enrichment

analysis on nearest genes of the 100 top ranked SNPs (**Fig. 4E**). The enrichment results revealed

potential biological processes that are relevant to COVID-19, such as the regulation of

glucokinase activity which is associated with the homeostasis of human blood glucose[34].

Notably, diabetes mellitus, a condition closely associated with hyperglycemia, is a typical

comorbidity of COVID-19[35]. Besides, among the top 10 potential genes that scored the highest,

we identified that the TBC1D4 gene, which regulates glucose homeostasis, is potentially

associated with COVID-19 comorbidities. Our findings are consistent with previous research

by Pellegrina et al.[36] and highlights the potential of our EpiGePT approach in discovering new

genetic markers that may be implicated in the pathogenesis of COVID-19. Overall, our

EpiGePT model provides new perspectives for understanding how the genetic variants could

contribute to the COVID-19 susceptibility and severity.

Fine-tuning on EpiGePT enables accurate prediction of regulatory interactions

Fine-tuning is an strategy that transfers the knowledge of a pretrained model to new tasks,

which is particularly prevalent in language models such as GPT[37] and BERT[38]. Here, we

finetuned a pretrained EpiGePT model on a new task for predicting the 3D genome interaction.

Given the HiChIP H3K27ac data from K562 and GM12878 cell lines, the features of two

anchors were extracted from a pretrained EpiGePT model and then fed to a finetune network

to predict whether it is a HiChIP loop. We compared EpiGePT with pretrain and finetuning

strategy to two baselines, DeepTACT[39] and a k-mer frequency[40] based method (see Methods).

The results illustrate that EpiGePT exhibits a superior classification performance across diverse

distance ranges compared to baselines. For example, in the GM12878 cell line within the 20-

40kbp distance range, EpiGePT demonstrates a significantly improved predictive performance

with an auROC of 0.949, surpassing 0.866 of DeepTACT [39] and 0.771 of Kmer (**Fig. 4F** and

Fig. S7). This significant improvement achieved through fine-tuning EpiGePT on a limited

dataset aligns well with the concept of few-shot learners[1], highlighting the power of the

pretrained EpiGePT model.

EpiGePT encompasses the regulatory relationships between TFs and target genes.

One of the key differentiating factors of EpiGePT compared to other sequence models lies in

its integration of TF binding status and TF expression. This unique feature empowers EpiGePT

to capture potential regulatory relationships embedded within the genomic sequence. In this

study, we specifically aimed to validate whether EpiGePT learns the regulatory relationships

between TFs and target genes (TGs). We defined gradient importance scores (GIS) based on

the absolute gradient values of predicted epigenomic signals w.r.t. TF profile to rank TFs give

a TG (see Methods). Particularly, we collected 15 TFs that play critical regulatory roles in

embryonic stem cells (ESC)[41, 42] and validated their interaction relationships using computed

GIS. As an example, for a target gene STAT3 that plays essential role for ESC pluripotency[43],

we computed GIS for each core TF across 1000 genomic bins and find other key TFs in ESC

ranked 1st (REST) and 2nd (POU5F1) at specific bins (**Fig. 5A-B**). Interestingly, the GO terms

enriched by the top 10% prioritized TF coding genes also included biological processes of

embryonic cell differentiation and development when we focus on the genomic bin that

POU5F1 ranked 2nd (**Fig. 5C**). We selected a TF as the target gene and calculated the integrated

GIS (IGIS) score for another key TF across eight epigenomic signals. Multiple TF-gene pairs

identified by IGIS that showed significant associations with POU5F1, such as ESRRB-

POU5F1[44] (rank 2nd), and ETV5-POU5F1[45] (rank 5th). Furthermore, we use TF-TG

relationships from either ChIP-seq data or external databases as ground truth to validate

whether IGIS is effective in prioritizing the TFs given a TG. First, we defined potential TF-

target gene pairs based on TF ChIP-seq data specific to certain cell types among all human

genes (see Methods). The results demonstrated a significant difference in rank between TF-

target gene pairs and TF-non-target gene pairs based on the IGIS score (**Fig. 5D**), with the

former exhibiting considerably higher ranks (one-side p-value < 0.001). Second, we collected

TF-target regulatory network data from two publicly available databases. We obtained a total

of 1,066 TF-gene pairs from the GRNdb[46] database based on liver-specific GTEx data, and

2,705 TF-gene pairs from the TRRUST[47] database after filtering. Then we calculated the rank

of each TF based on GIS of the TF expression and a genomic bin-level mask for each pair.

Interestingly, when using liver expression data, we found that the average rank of TFs from

TRRUST was 7.9%, significantly lower than the rank based on expression values (one-side p-

value < 1e-5). Similarly, based on the GRNdb data, the majority of TF-gene pairs obtained had

TF ranks within the range of 20%, and the mean of this distribution was significantly lower

than the rank based on expression values with one-side *p*-value < 1e-5 (**Fig. 5E**). For instance,

TMEM55B plays a significant role in regulating lysosome movement, and is regulated by sterol

response element binding factor 2 (SREBF2)[48], while GIS enable the identification of SREBF2

as the top-ranked TF associated with TMEM55B, further validating the role of GIS in

prioritizing functional TFs. The comprehensive validation from both ChIP-seq datasets and

external databases further support the effectiveness of GIS in identifying context-specific TF-

TG relationships.

Online prediction tool for EpiGePT

In order to facilitate the utilization of EpiGePT for the prediction of multiple chromatin

states of any cellular context and any genomic regions, especially for research

personnel who lack coding expertise, we have developed a user-friendly online web

server, named EpiGePT-online (https://health.tsinghua.edu.cn/epigept/) (Text S2). The

online web server was developed using PHP, JavaScript and HTML, which provides an

interactive web interface for online prediction of 8 chromatin states of specific genomic regions

(**Fig. 6**). Users can obtain the predicted signals of multiple genomic regions by submitting a

region file and a TF expression file of 711 selected TFs (Supplementary Table S3), or obtain

predicted signals of specific regions directly by selecting genomic locus. As to the two types of

input files, we provide example files to demonstrate their formats, and accept expression files

stored in either numpy or csv formats, to increase the universality of the web server (Fig. S5).

The web server outputs the results in a web summary html, which saves significant amount of

time for installation and implementation. Furthermore, we provide a detailed tutorial to enable

users to quickly learn how to use our website. We anticipate that this web server will assist

researchers in predicting chromatin states of specific cell types and further deepening

their understanding of gene regulatory mechanisms.

Discussion

In this paper, we introduced EpiGePT, a transformer-based large language model, for predicting

the chromatin states given any cellular context. Compared with the existing machine learning

based computational methods, EpiGePT takes transcription factor profile and DNA sequence

information as inputs by multitask learning and self-attention mechanism within a unified

model. With these two types of input information and four modules of network architecture,

EpiGePT overcomes the limitation of the existing models and demonstrates state-of-art

performance in prediction of multiple chromatin signals in diverse experimental settings. With

the superior predictive performance of EpiGePT, we are able to investigate one of the

fundamental questions in functional genomics: how transcription factors and cis-regulatory

elements regulate gene activity. In this work, we investigated this question from two aspects:

1) identifying the interactions of cis-regulatory elements and their target genes with the help of

self-attention mechanism in EpiGePT; 2) estimating the variant effect based on the LOS scores

computed by the outputs of EpiGePT to assist in discovering human disease-associated SNPs.

First, the self-attention scores between tokens can provides us with an intuitive and quantitative

measure of the interaction level between different genomic regions, which offers new

opportunity to discover the target gene of cis-regulatory elements and find the interpretability

of EpiGePT. Second, the LOS scores of the multiple chromatin signals from different tissue or

cell lines are complementing each other, which provides us with a more comprehensive

characterization of the variant and enables accurate prediction of the variant impact. Such

variant effect prediction by EpiGePT establishes a foundation for understanding the underlying

relationship between genetic variations and disease mechanisms.

There exist several extensions and refinements that can be applied to further improve the

EpiGePT model. Firstly, the incorporation of chromatin regulators (CRs) as trans-acting factors

into the TF module could enhance the modeling of regulated transcription processes, thereby

increasing the accuracy of the predictions. Secondly, the inclusion of high-order interactions

between TFs in the framework could provide a more comprehensive representation of the

regulatory relationships, and potentially enhance the predictive performance. Third, the

application of EpiGePT to single-cell genomics could enable the profiling of chromatin signals

at single-cell resolution, facilitating a holistic understanding of regulatory heterogeneity in

different cell subpopulations for researchers.

Based on EpiGePT, users are able to predict multiple chromatin profiles in different cell lines

or tissues, which could provide a foundation for biological discovery, decoding transcriptional

regulation mechanisms, and investigating disease mechanisms. We anticipate that EpiGePT can

provide valuable insights to researchers in understanding regulatory mechanisms.

Methods

Data processing

Chromatin accessibility data and Expression data We used three different datasets in the experiments. For chromatin accessible data, we downloaded DNase bam files and narrow peaks across 129 human biosamples from ENCODE[12] project (Supplementary table S1 and S2). We divided the human hg19 genome into 200bp non-overlapping locus (use bin instead), and we assigned the label for each locus in each cell type. For the regression design, we pooled the bam files of multiple replicates for a cell type (Supplementary table S1 and S2), and obtain the raw read count n_{lk} for locus l in cell type k. We normalized the raw read count in order to eliminate the effect of sequencing depths, in the form of $\tilde{n}_{lk} = N n_{lk}/N_k$, where N_k denotes the total number of pooled reads for cell type k and $N = \min_{k} N_k$ denotes the minimal number of pooled reads across all cell types. The normalized read counts are further log transformed with pseudo count 1, which represent the continuous level of chromatin accessibility. For binary classification design, we assigned a binary label y_{lk} to 1 if the number of raw read counts of the locus l in the cell type k greater than 30, which represent the locus is an accessible region in this cell type, resulting in the identification of regions as accessible in 13% on average and 8% at median in the screened genomic regions across 129 cell types. The proportion of open regions varies among different cell types, and the average openness level mentioned above is generally consistent with that maintained in ChromDragoNN[16].

RNA-seq data of the 711 human transcription factors were downloaded and extracted from the ENCODE project (Supplementary table S5 and S6). We perform log transformation with

pseudo count 1 and quantile normalization based on TPM values. The normalized TPM values were averaged across replicates and mean expression profile of each cell type was finally used to calculated of the transcription feature.

Multiple chromatin signals data DNase-seq, RNA-seq and ChIP-seq data were also downloaded from ENCODE project (Supplementary table S3, S4 and S6). We applied the same process to these data as above, and finally we obtained the 8 chromatin signals of 13,300,000 bins of 128bp in 28 cell types. The continuous level of chromatin signals we extracted were 'DNase', 'CTCF', 'H3K27ac', 'H3K4me3', 'H3K36me3', 'H3K27me3', 'H3K9me3' and 'H3K4me1', which includes crucial epigenetic modifications and markers for gene regulation and transcription.

Chromatin states data We downloaded the 15-state ChromHMM[21] annotations across 127 epigenomes from the ROADMAP project. The state of chromatin is annotated for each 200bp bin in a specific cell type. RNA-seq data across 56 cell types of TFs was download and extracted from the ROADMAP[49] project (Supplementary table S7 and S8). Subsequently, we mapped these 711 transcription factors to the downloaded RNA-seq data, resulting in the identification of RNA-seq data for 642 transcription factors. In the subsequent experiments, we utilized the expression data of these 642 transcription factors. We finally calculated the normalized TPM values of the 642 TFs on 56 cell types we extracted for the using in the classification model. For coarse grain chromatin state prediction, we took the state 'Quies' as low signal regions and other states as signal regions. For fine grain chromatin state prediction, we extracted the state 'TssA', 'TssAFlnk', 'TssBiv' and 'BivFlnk' as TSS regions, state 'EnhG', 'Enh' and 'EnhBiv' as enhancer regions, 'Quies' as low signal regions and other state as other regions. To balance the

number of different chromatin states, we downsampled the low signal regions and obtained

921,074 locus each cell line finally.

Model architecture

Sequence module and Transformer module

As shown in Figure 1 and Fig. S6A, the sequence module receives a one-hot matrix (A =

[0,0,0,1], C = [0,1,0,0], G = [0,0,1,0], T = [0,0,0,1]) of size (128000,4) as input, representing a

sequence of 128 kilobase pairs (kbps) and contains five 1-dimentional convolutional blocks to

extract DNA sequence features. Each block includes a convolutional layer and a maxpooling

layer (Fig. S6B). The first convolutional layer considers the input channels as 4 and performs

convolution along the sequence direction. The input sequence features are one-hot embeddings

of size $L \times 4$, where L denotes the length of the input long range DNA sequence. After 5

maxpooling layers, the output size of sequence feature is $L/N \times C$, where C denotes the hyper-

parameter for sequence embedding and N denotes the length of locus to predict. We set C to

256 in the pre-training stage of chromatin accessibility prediction experiments. Rectified linear

units (ReLU) are used after each convolution operation for keeping positive activations and

setting negative activation values to zeros. Sequence features were then concatenated with

transcriptomics features, and we finally obtained a vector of size $L/N \times (C + n_{TF})$, where n_{TF}

denotes the dimension of the transcription factors features after padding. In our model, after

adding padding to the 711 TFs, the n_{TF} is set to 712. Therefore, the input token number for the

transformer module is 1000, and each token embedding has a dimensionality of 968.

We utilize the transformer module to integrate information from both the sequence and

transcription factors (TFs), enabling the capturing of long-range interactions between genomic

bins. We applied N_t layers of Transformer encoder with n_{head} different attention heads to the

token embedding sequence. The input X of the transformer encoder is a genomic bin sequence

with dimensions $(Sequence\ length, embedding\ dim)$. Specifically, this dimension is (1000,

968) in EpiGePT, indicating that input genomic bin sequence has a length of 1000, and each

genomic bin has an embedded representation that combines the sequence information with cell-

type-specific features with dimension of 968. Each Transformer encoder includes a multi-head

self-attention mechanism and a feed-forward neural network. For self-attention in each head,

the calculation is based on the matrix operation.

$$Attention(Q, K, V) = softmax(\frac{QK^T}{d_k})V$$

531 For multi-head attention, Transformer encoder learns parameter matrices $W_i^Q \in$

532 $\mathbb{R}^{d_{model} \times d_Q}, W_i^K \in \mathbb{R}^{d_{model} \times d_K}$ and $W_i^V \in \mathbb{R}^{d_{model} \times d_V}$ for the i_{th} head and concatenate the

533 multiple heads to do the projection.

$$Q_i = X \times W_i^Q, K_i = X \times W_i^K, Q_i = X \times W_i^V$$

Where d_{model} denotes the dimension of token in the input sequence X, which is 968 in

EpiGePT and $d_Q = d_K = d_V = 512$. The matrices Q, K, and V are obtained by the application

of mapping functions represented by W_i^Q, W_i^K and W_i^V, followed by concatenating of Q_i to Q,

K_i to K, and V to V. These mapping functions serve to transform the concatenated embeddings

into the resulting matrices. We set N_t to 16 for the chromatin accessible prediction experiments,

N_t to 12 for the chromatin state classification and multiple chromatin signals prediction

experiments, and set n_{head} to 8 for all experiments.

The regression model, the output layer uses a linear transformation and use mean square error

(MSE) as the loss function. For classification model, the output layer uses a linear

transformation combined with a sigmoid function, and use the cross-entropy loss for

classification experiments.

TF module

For binding status, we scanned the input bins for potential binding sites for a set of 711 human

transcription factors from HOCOMOCO database[50] with the tool Homer[51] (Table S5). We then

selected the maximum score of reported binding status for each transcription factor to obtain a

vector of 711 dimensions as the motif feature for each DNA bin. For gene expression, we

focused on log-transformed TPM values of the 711 transcription factors and obtained a vector

of 711 dimensions after quantile normalization as the expression feature. With these data, we

combined the two vectors of motif and expression features by taking the element-wise product,

and we concatenated the result to the output of sequence module.

Model evaluation

To evaluate our model, we applied five-fold cross-validation in the different experiments on

cell-type level. For chromatin accessible experiments, the 129 cell lines are partitioned into a

training set and a testing set randomly.

Cell-type-wise metrics are defined to evaluate our method in different experiments, which were

calculate with the data within a test cell type across all genomic locus. For binary classification

design, we used cell-type-wise auPRC and auROC to evaluate our EpiGePT. Let $Y_{L \times K}$ and $\hat{Y}_{L \times K}$

be the true and predicted matrix, where L denotes the number of locus and K denotes the number

of test cell types. We calculated the auPRC and auROC for each $(y_{1i}, y_{2i}, \cdots, y_{Li})$ and

$(\hat{y}_{1i}, \hat{y}_{2i}, \cdots, \hat{y}_{Li})$. For multiple classification, we use macro average of the auPRC and auROC

to evaluate the classification performance, which compute the metric independently for each

class and then take the average hence treating all classes equally. For regression design, we

used two metrics for model evaluation, which are cell-type-wise Pearson correlation coefficient

and prediction squared error. Prediction square error (PSR) is calculated as $PSR = 1 -$

$\sum_k \sum_l (y_{lk} - \hat{y}_{lk})^2 / (y_{lk} - \bar{y}_{*k})^2$, where $\bar{y}_{*k} = \sum_l y_{lk}/L$ denotes the mean of the true level of

the response in the cell type k.

To compare the performance of our method with other baseline methods, we conducted

hypothesis testing on the metrics based on cell types. Since the metrics on a given cell type

across different methods are paired data and the statistical distribution is unknown, we

employed both Binomial and Wilcoxon tests, with the alternative hypothesis being that

EpiGePT outperforms the other methods. If we reject the null hypothesis, it provides

compelling evidence to support the claim that EpiGePT performs better than the other methods.

To evaluate the computational efficiency, we recorded the running time of a single epoch of

EpiGePT and baseline methods (Supplementary Text S3). Compared to traditional CNN models

such as DeepCAGE[17] and ChromDragoNN[16], as well as larger sequence models like Enformer,

EpiGePT demonstrates a balance between high computational efficiency and performance.

Model fine-tunning

For the fine-tuning process, we kept the parameters of the pre-trained model fixed without

making any updates. For the specific fine-tuning task of chromatin interaction prediction based

on HiChIP data, the multi-task module was replaced with a two-layer MLP network, containing

256 hidden nodes for each layer. During the training process, only the weights in the MLP

network were updated. Notably, when utilizing HiChIP data at a resolution of 5k, both the

enhancer and promoter anchors spanned 5kbp. Then we input a region extending 128kbp from

the center of the anchor of the neighboring gene into the EpiGePT. Consequently, a 968-

dimensional feature vector for each genomic bin was derived from the output of the last

transformer encoder layer. These feature vectors from all bins within the two anchors were

concatenated, resulting in a high-dimensional vector of size 76,472.

Baseline methods

Four baselines were introduced for epigenetic signals prediction. BIRD[15] is a multiple linear

regression model that only takes gene expression data as input and makes predictions on a fixed

locus. ChromDragoNN[16] is a deep neural network that takes gene expression of 1630 TFs and

DNA sequence as input. Specifically, ChromDragoNN[16] uses a ResNet[52] to extract sequence

features and use linear transformation to combine the TF gene expression feature and sequence

feature to make the final prediction. DeepCAGE[17] Integrating regulatory DNA sequence is a

deep densely connected convolutional network for predicting chromatin accessibility. The

dense-connected neural network architecture used by DeepCAGE[17] may struggle to capture the

complex interactions between genomic regions. Enformer[11] is a deep neural network that

integrates convolutional neural network and transformer, and only takes DNA sequence as input.

Enformer takes DNA sequence of length 196kbp as input to predict 5,313 genomic tracks of

human and 1,643 tracks of mouse genome simultaneously. However, one of the limitations of

Enformer is that it can only model and predict cell types in the training data and cannot be

applied to new cell types. In order to ensure the fairness of the benchmark experiment, we

retrained the Enformer model with the same input and output data as EpiGePT when reproduce

the Enformer model (Text S4).

Two baseline methods were introduced for predicting HiChIP interaction. DeepTACT[39] is a

deep learning method for predicting 3D chromatin contacts using both DNA sequence and

chromatin accessibility. We adopted the structure of DeepTACT[39] and kept the anchor length at

5k. The input to the model consists of two anchor sequences represented as one-hot matrices

and the two openness scores of the two anchors on the corresponding cell type extracted from

OpenAnnotate[53]. Regarding the Kmer features[40], K is chosen as 5 to extract sequence features.

For each anchor, a vector of dimension $4^5 = 1024$ was obtained. Further training was

performed using an MLP with a hidden layer dimension of 256.

Enhancer, Silencer and HiChIP loop prioritization

We collected cis-regulatory elements-gene pairs in K562 cells from other studies and public

database to demonstrate the interpretability of self-attention mechanisms in the EpiGePT.

Enhancers and silencers are typical *cis*-regulatory elements known play important roles in

transcriptional control during normal development and disease. For enhancers, we downloaded

enhancer-gene pairs from two studies: Gasperini et al.[23] and Fulco et al.[24], both of which were

tested using a CRISPRi[22] assay perturbation. Two datasets contain 664 and 5,091 enhancer-

promoter interactions or element-TSS interactions. For silencers, we obtained and random

sampled 831 validated silencers-gene pairs with distance within 64kbp in K562 cells curated

from high-throughput experiments from SilencerDB. As there are no experimentally validated

interaction relationships between these silencers and genes, we generated silencer-gene pairs

by associating the nearest neighbor genes for classification purposes. Similarly, negative

samples were generated by constructing DNase-seq, ATAC-seq and nearest genes using the

same approach. Ultimately, we obtained a dataset comprising 1,662 silencer-gene pairs,

encompassing both positive and negative instances.

To obtain scores for regulatory element-gene pairs, we first used the region extending 128kbp

from the center of the enhancer as input and extracted the token where the interacting genes

reside, so that we could filter out regulatory element-gene pairs that were located further than

64kbp apart. Subsequently, we stratified the remaining pairs based on their distance. Since the

positive and negative sample ratios varied across datasets, we adopted different stratification

strategies for different distance ranges (**Fig. 3**). Next, we averaged the attention matrices of the

Transformer encoder across all layers and heads. The summed attention scores from other

tokens to the key token containing the gene TSS were used as the attention score of this element-

gene pair. This score represents the attention value that the enhancer-centered region receives

for the transcription start site (TSS) of the gene. We also calculated the attention score from the

bin containing the center of the regulatory element to the bin containing the TSS, which only

slightly affects the experimental results of regulatory element prioritization.

We collected 5k resolution data from the HiChIPdb (http://health.tsinghua.edu.cn/hichipdb/)

database, specifically from K562 and GM12878 cell lines. We filtered the data to include only

loops where at least one anchor falls within a gene region. We stratified the loops based on

distance into three categories: 0-20kbp, 20-40kbp, and 40-64kbp. For each distance category,

we selected 2000 positive pairs with most significant q-value. To ensure consistency in the

distance distribution, we selected negative pairs by fixing a gene and choosing anchors at

equidistant locations in the opposite direction.

Gradient importance scores

EpiGePT possesses the capability to assign priority rankings to transcription factors by utilizing

gradient importance scores (GIS), taking into account specific cell types and chromatin regions.

The GIS were employed to identify potential functional relationships between specific

transcription factors (TFs) and target genes. Specifically, for a given TF-target gene pair, the

transcription start sites (TSS) of genes were used as central loci, and the regions spanning 128

kbp upstream and downstream of the TSS were selected as input. Next, we filtered out bins

with motif binding scores indicating potential binding for the given TF. For these selected bins,

we calculated the GIS for the predictions of eight epigenomic signals across the 711 core TFs.

$$GIS_{ijk} = \frac{1}{|\zeta|} \sum_{l \in \zeta} |\frac{\partial \hat{y}_{ljk}}{\partial tf_{ij}}|$$

Where, i denotes the ith TF in the set of core TFs, j denotes the jth cell type, k denotes the kth

predicted epigenomic signal, and ζ denotes the set of genomic bins that have binding for the

given TF. In the calculation of the gradient, $\hat{y}_{ljk}$ denotes the predicted value of the kth

epigenomic signal by the model using the expression in the jth cell type at the lth bin. On the

other hand, tf_{ij} denotes the product of the expression of ith TF in the jth cell type and the

corresponding TF binding score.

If we consider the GIS for the prediction of all 8 epigenomic signals simultaneously, we can

prioritize the TFs by calculating their ranks based on each signal separately. Then, we can

calculate an integrated gradient importance score (IGIS) for each TF by aggregating the ranks

from all 8 signals.

671
$$IGIS_{ij} = \frac{1}{8} \sum_{k} rank(GIS_{ijk})$$

Both the GIS and the IGIS are capable of capturing the significance of a transcription factor

(TF) in regulating a specific gene within the context of a specific cell type. Consequently, these

scores hold potential value in the discovery of TFs that play crucial roles in the regulation of

specific genes, thereby contributing to our understanding of essential regulatory mechanisms.

In the context of validating TF-TG pairs in the GRNdb and TRRUST databases, we opted to

utilize liver expression data as a representative example due to the unavailability of cell type

information for TRRUST. Furthermore, in this experimental setup, the tf_{ij} denotes the

expression of ith TF in the jth cell type and ζ denotes the set of genomic bins that have binding

for the TF of the given TF-target gene pair.

Potential TF-target gene pairs from ChIP-seq data

In this study, we utilized three distinct cell types to conduct a comprehensive screening of TF-

target gene pairs and non-target gene pairs across the human genome. Initially, we obtained the

narrow peak files (ENCFF388AJH, ENCFF717IXP, and ENCFF885KLR) from ChIP-seq

experiments across three cell types from the ENCODE project. Subsequently, we meticulously

examined the number of peaks within a 128kbp region both upstream and downstream of the

transcription start site (TSS) for each gene. Different thresholds were applied to the ChIP-seq

data of various TFs. Genes lacking any peaks within the defined region were classified as non-

target genes, while genes surpassing the threshold in terms of peak counts were designated as

target genes. Specifically, for the aforementioned three cell types, threshold values of 10, 15,

and 6 were respectively employed. Finally, the IGIS approach was employed to determine the

corresponding ranks of TFs in the TF-target gene pairs.

Pathogenic SNPs prioritization

We collected single nucleotide polymorphisms (SNPs) data from the ClinVar and ExAC

databases, which include both potentially pathogenic and benign SNPs. To evaluate the ability

of EpiGePT to predict variant effects, we computed the log-odds scores (LOS) for multiple

chromatin signals using EpiGePT on these SNPs. Subsequently, we utilized these scores to

distinguish between pathogenic and benign SNPs. The LOS score for each chromatin signal

was defined by computing a forward pass through the model using the reference and alternative

alleles.

$$\Delta O_{signal} = \log \left(\frac{output(I_{alt})}{output(I_{ref})} \right)$$

Each chromatin epigenomic profile in each cell line or tissue predicted by EpiGePT can be used

to compute a specific variant score. We did not take the absolute value in this calculation, so

the resulting LOS score indicates the direction of change in the model output after the

appearance of the variant. In addition to the predicted chromatin signals output by the eight

models, attention score changes based on self-attention are also noteworthy. We computed the

log-odds scores for attention by summing the attention scores of the 10 bins upstream and

downstream of the locus of the SNP, to evaluate the effect of the variant.

$$\Delta O_{attention} = \sum_{i=-5}^{5} |log\left(\frac{attn(bin_i)_{I(alt)}}{attn(bin_i)_{I(ref)}}\right)|$$

Where i represents the index of the neighboring bins relative to the locus of the SNP. To avoid

the variant effects of different bins from cancelling each other out during the summation process,

we computed the absolute value of the change in attention scores for each bin and then summed

the scores of the 10 adjacent bins centered at the SNP position. For the classification of

pathogenic SNPs, we calculated these nine LOS for attention separately for each of the 28

tissues or cell lines in training data. As a result, we obtained a feature vector of 252 dimensions

for each SNP. Then a classifier with 252 features computed by EpiGePT and 52 annotations

from CADD score as inputs are used to predict pathogenic SNPs against benign or likely benign

SNPs. Here, we employed MLP as classifier to validate the effectiveness of the features we

obtained. A five-fold cross-validation experiment is employed for validation, and we utilize two

different positive-to-negative sample ratios, namely 1:1 and 1:2. For each sample ratio, we

randomly sample 32,000 positive samples. The effectiveness of the variant score in identifying

pathogenic SNPs is evaluated using the area under the auROC and the auPRC. Additionally,

we also utilized the logistic regression (LR) as the classifier, consistent with the LR classifier

used in CADD, and found a similar improvement when predicting pathogenic SNPs.

We applied the same method to calculate the LOS scores of the 8 predicted chromatin signals

for the COVID-19 GWAS data. The absolute values of the scores were summed as the overall

score for each SNP. For each significant SNP associated with COVID-19 severity obtained

from the GWAS data, we selected normal SNPs within a 64kb region around the SNP as

background to calculate the rank of the LOS score for the COVID-19 associated SNP in this

730 region. Furthermore, we calculated the LOS scores for all 9, 484 COVID-19 associated SNPs

731 and ranked them accordingly. The top 10 SNPs with the highest LOS scores were selected,

732 which are considered to have potential genetic associations with COVID-19 severity and

733 complications.

GTEx classification

We collected eQTL data from the supplementary materials of Wang et al[28]. In their study, the

authors identified causal eQTLs through statistical fine-mapping, using a posterior inclusion

probability (PIP) threshold of >0.9 for putative causal variants based on expression modifier

score (EMS), and a PIP threshold of <0.9 for putative non-causal variants. To validate the ability

of EpiGePT to distinguish potential causal variants, we perform a classification task on these

variants. For each variation, 128kbp sequence regions near it were selected as the input of the

model, and a score of variation was given by EpiGePT model. For each variant under each

tissue, we can obtain an 8-dimensional vector of genomic features including DNase, CTCF and

other ChIP-seq signals. Based on the LOS score, separate random forest classifiers consisting

of 10 decision trees are trained for each tissue in order to distinguish between causal and non-

causal variants. The models are evaluated using 5-fold validation on each tissue, with area under

the auPRC and auROC as metrics for assessing their ability to distinguish between causal and

non-causal variants.

Code availability

749 All components of EpiGePT are freely available at https://github.com/ZjGaothu/EpiGePT.

750 Here, users can access the code for reproducing EpiGePT, as well as the data collection and

751 preprocessing pipelines used for model training in benchmark experiments.

Data availability

Information and processed data of multiple chromatin signals of whole genome, motif binding

status and expression data of TFs in the corresponding cell lines/tissues, which are used in

EpiGePT are available at Supplementary Tables. The information about the cell lines/tissues

used and the 711 filtered transcription factors is available in the supplementary table. The High

throughput validated silencers of K562 cell line are download from SilencerDB

(http://health.tsinghua.edu.cn/silencerdb) database. The HiChIP data of K562 cell line and

GM12878 cell line are downloaded from HiChIPdb (http://health.tsinghua.edu.cn/hichipdb/)

database. The DNase-seq peak and ATAC-seq peak data are obtained from the ENCODE

project. Enhancer-gene pairs of CRISPRi[24] experiments are obtained from the supplementary

information of Gasperini et al. and Fulco et al. The regulatory network data for transcription

factors and target genes were obtained from the TRRUST[47] database

(https://www.grnpedia.org/trrust/) and the GRNdb[46] database (http://www.grndb.com). The

annotated chromatin states for whole genome are downloaded from the ROADMAP

epigenomics project (https://egg2.wustl.edu/roadmap/web_portal/chr_state_learning.html).

The RNA-seq read counts matrix for protein coding genes used for the prediction of the

chromatin 15-states annotated by ChromHMM are downloaded from the ROADMAP project

(https://egg2.wustl.edu/roadmap/data/byDataType/rna/expression/57epigenomes.N.pc.gz).

The GWAS data of COVID-19 are download from the COVID-19 Host Genetics Initiative

(https://www.covid19hg.org/).

Competing interests

The authors have declared no competing interests.

Acknowledgments

Z.G and R.J. was supported by the National Key Research and Development Program of China [2021YFF1200902] and the National Natural Science Foundation of China [62203236 and 62273194]. Q.L., W.Z and W.H.W were supported by NIH grants R01 HG010359, P50 HG007735 and NSF DMS 1952386.

Figures

Figure 1

Fig. 1 Overview of the EpiGePT model for multiple epigenomic signals prediction. The model consists of four modules, namely the Sequence module, the TF module, the Transformer module, and the Multi-task module. The sequence module comprises multiple layers of convolution applied to the one-hot encoded DNA sequence input. The input sequence length consists of 1000 genomic bins of 128bp for the prediction of multiple signals and 50 bins of 200bp for the prediction of DNase signal alone. The TF module encompasses the binding status and expression of 711 transcription factors. The Transformer module consists of a series of consecutive transformer encoders, while the multi-task module is composed of a fully connected layer.

 Figure 2

 Fig. 2 Performance of EpiGePT and baseline methods on the benchmark experiment. (A)

EpiGePT and baseline methods were compared in terms of their regression performance for

DNase signal regression across cell types, genomic regions, and combined cell type and

genomic regions. (B) Comparison of EpiGePT and Enformer performance. Each point in the

scatter plot represents the performance of Enformer on the data of a specific cell type (x-axis)

compared to the performance of EpiGePT (y-axis). (C) EpiGePT and baseline methods'

performance on binary prediction of DNase-seq signals. (D) EpiGePT demonstrates excellent

performance in predicting diverse epigenetic signals across various cell types, including

DNase-seq, CTCF, and histone modifications. (E) EpiGePT predictions compared to

experimental signals visualized for a representative example. Genome-wide multi-signal

predictions (black) and DNase-specific predictions (orange).

Figure 3

Fig. 3 Application of self-attention mechanism in EpiGePT for long-range chromatin interaction identification. (A) The performance (auPRC) of attention score of EpiGePT in distinguishing enhancer-gene pairs at different distance ranges on two different datasets. (B) The performance (auROC and auPRC) of attention score of EpiGePT in distinguishing silencer-gene pairs at different distance ranges based on the data from SilencerDB[25]. (C) Heatmap of the self-attention matrix of each attention head centered at the TSS of the CHD4 gene, the (*i, j*) element in the matrix denotes the average attention score between the *i*th genomic bin and the

810   *j*th genomic bin across all layers. (D) Attention scores centered at the TSS of the CHD4 gene,

811   and putative enhancer regions in its vicinity. (E) The performance (auROC and auPRC) of

812   attention score of EpiGePT in distinguishing HiChIP loops of H3K27ac at different distance

813   ranges on K562 cell line and GM12878 cell line.

Figure 4

Fig. 4 Variant effect prediction of EpiGePT. (A) The LOS score for each epigenomic signal

is calculated by the log change fold of the predicted epigenomic signal for reference genome

and WGS genome. (B) The performance of EpiGePT and Enformer in discriminating causal

SNPs on the Lung tissue. (C) The three subplots from left to right respectively depict the

classification results for disease-related SNPs and benign SNPs down-sampled sourced from

the ClinVar database, with balanced positive and negative samples (1:1 and 1:2 ratio), as well

as normal SNPs sourced from the ExAC database with a MLP classifier. (D) The ranked

position of COVID-19 related GWAS data among surrounding benign SNPs based on their

LOS scores, as determined using different tissue or cell-type expression data. The results were

stratified based on the distance range of the risk region. The resulting mean and median ranks

were both below 0.5. (E) Enrichment result (Biological process, Cellular component and

Molecular function) of the nearest genes of the COVID-19 associated SNPs with the max LOS

scores. (F) The performance (auROC and auPRC) of the fine-tuned EpiGePT model and

baseline methods (DeepTACT and Kmer) in distinguishing enhancer-gene pairs at various

distance ranges (0-20 kbp, 20-40 kbp and 40-64 kbp) on K562 cell line.

Figure 5

 Fig. 5 Gradient importance scores (GIS) uncover regulatory transcription factors. (A)

 Genomic regions around TSS of the STAT3 gene and expression data on ESC were fed into

 EpiGePT. The scatter plot represents the GIS scores of core TFs on each genomic bin. Each dot

 represents the GIS score of a core TF on a specific genomic bin. In specific bins, key TFs in

 ESC, such as NR5A2 and POU1F5 highlighted in the figure, exhibit high ranks in the GIS

 scores. (B) Heatmap of GIS for 10% important TFs surrounding STAT3 gene, specifically

 focusing on bins with POU1F5 ranked 3rd. Each row represents a predicted epigenomic signal,

and the TFs are sorted based on their GIS on DNase signals. Bar plot of the top 15 TFs with the

highest GIS scores. (C) Based on the top 10% ranked TFs mentioned above, gene ontology

enrichment analysis revealed significant enrichment in biological processes related to

embryonic development and cellular differentiation. (D) Based on TF ChIP-seq data, all 23,635

human genes were classified into target genes and non-target genes. The results revealed that

TFs exhibited significantly higher ranks on potential target genes compared to non-target genes.

(E) The distribution of the rank of TFs in the GIS and expression value among the 2,705 TF-

gene pairs from the TRRUST database and 1,066 TF-gene pairs derived from genotype-tissue

expression (GTEx) data of the liver sourced from the GRNdb database. The analysis reveals

that the median rank of TFs from the TRRUST database is significantly lower than 0.06 (one-

side p-value < 3.12 e-18) and the median rank of TFs obtained from the GRNdb database is

significantly lower than 0.5 (one-side p-value < 2.50 e-140).

Figure 6

Fig. 6 Overview of the online prediction web server of EpiGePT. We collected eight types

of epigenetic genome modification signals and corresponding expression data of transcription

factors in different cell types or tissues from the ENCODE project. Based on these data, we

trained the EpiGePT model and deployed it as a built-in kernel on an Apache server. Users

without much coding experience can also access the web server in two ways to obtain the eight

types of epigenetic genome modification signals for specified cell types and genomic regions

without programming or installation.

Reference

1. Brown, T. et al. Language models are few-shot learners. *Advances in neural information processing systems* **33**, 1877-1901 (2020).

2. Ramesh, A. et al. in International Conference on Machine Learning 8821-8831 (PMLR, 2021).

3. Alexander, R.P., Fang, G., Rozowsky, J., Snyder, M. & Gerstein, M.B. Annotating non-coding regions of the genome. *Nature Reviews Genetics* **11**, 559-571 (2010).

4. O'Malley, R.C. et al. Cistrome and epicistrome features shape the regulatory DNA landscape. *Cell* **165**, 1280-1292 (2016).

5. Yenduri, G. et al. Generative Pre-trained Transformer: A Comprehensive Review on Enabling Technologies, Potential Applications, Emerging Challenges, and Future Directions. *arXiv preprint arXiv:2305.10435* (2023).

6. Liu, Q., Xia, F., Yin, Q. & Jiang, R. Chromatin accessibility prediction via a hybrid deep convolutional neural network. *Bioinformatics* **34**, 732-738 (2018).

7. Min, X., Zeng, W., Chen, N., Chen, T. & Jiang, R. Chromatin accessibility prediction via convolutional long short-term memory networks with k-mer embedding. *Bioinformatics* **33**, i92-i101 (2017).

8. Yin, Q., Wu, M., Liu, Q., Lv, H. & Jiang, R. DeepHistone: a deep learning approach to predicting histone modifications. *BMC genomics* **20**, 11-23 (2019).

9. Liu, Q., Lv, H. & Jiang, R. hicGAN infers super resolution Hi-C data with generative adversarial networks. *Bioinformatics* **35**, i99-i107 (2019).

10. Zeng, W., Wu, M. & Jiang, R. Prediction of enhancer-promoter interactions via natural language processing. *BMC Genomics* **19**, 84 (2018).

11. Avsec, Ž. et al. Effective gene expression prediction from sequence by integrating long-range interactions. *Nature methods* **18**, 1196-1203 (2021).

12. Consortium, E.P. An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**, 57 (2012).

13. Vaswani, A. et al. in Advances in neural information processing systems 5998-6008 (2017).

14. Song, L. & Crawford, G.E. DNase-seq: a high-resolution technique for mapping active gene regulatory elements across the genome from mammalian cells. *Cold Spring Harbor Protocols*

888   **2010**, pdb. prot5384 (2010).

889   15.   Zhou, W. et al. Genome-wide prediction of DNase I hypersensitivity using gene expression.

890   *Nature communications* **8**, 1-17 (2017).

891   16.   Nair, S., Kim, D.S., Perricone, J. & Kundaje, A. Integrating regulatory DNA sequence and gene

892   expression to predict genome-wide chromatin accessibility across cellular contexts.

893   *Bioinformatics* **35**, i108-i116 (2019).

894   17.   Liu, Q., Hua, K., Zhang, X., Wong, W.H. & Jiang, R. DeepCAGE: incorporating transcription

895   factors in genome-wide prediction of chromatin accessibility. *Genomics, Proteomics &*

896   *Bioinformatics* **20**, 496-507 (2022).

897   18.   Holwerda, S.J.B. & de Laat, W. CTCF: the protein, the binding partners, the binding sites and

898   their chromatin loops. *Philosophical Transactions of the Royal Society B: Biological Sciences*

899   **368**, 20120369 (2013).

900   19.   Bannister, A.J. & Kouzarides, T. Regulation of chromatin by histone modifications. *Cell*

901   *research* **21**, 381-395 (2011).

902   20.   Karolchik, D. et al. The UCSC genome browser database. *Nucleic acids research* **31**, 51-54

903   (2003).

904   21.   Ernst, J. & Kellis, M. Chromatin-state discovery and genome annotation with ChromHMM.

905   *Nature protocols* **12**, 2478-2492 (2017).

906   22.   Larson, M.H. et al. CRISPR interference (CRISPRi) for sequence-specific control of gene

907   expression. *Nature protocols* **8**, 2180-2196 (2013).

908   23.   Gasperini, M. et al. A genome-wide framework for mapping gene regulation via cellular genetic

909   screens. *Cell* **176**, 377-390. e319 (2019).

910   24.   Fulco, C.P. et al. Activity-by-contact model of enhancer–promoter regulation from thousands

911   of CRISPR perturbations. *Nature genetics* **51**, 1664-1669 (2019).

912   25.   Zeng, W. et al. SilencerDB: a comprehensive database of silencers. *Nucleic acids research* **49**,

913   D221-D228 (2021).

914   26.   Mumbach, M.R. et al. HiChIP: efficient and sensitive analysis of protein-directed genome

915   architecture. *Nature methods* **13**, 919-922 (2016).

916   27.   Zeng, W., Liu, Q., Yin, Q., Jiang, R. & Wong, W.H. HiChIPdb: a comprehensive database of

917   HiChIP regulatory interactions. *Nucleic Acids Research* **51**, D159-D166 (2023).

28.   Wang, Q.S. et al. Leveraging supervised learning for functionally informed fine-mapping of cis-

eQTLs identifies an additional 20,913 putative causal eQTLs. *Nature Communications* **12**, 3394

(2021).

29.   Landrum, M.J. et al. ClinVar: public archive of interpretations of clinically relevant variants.

Nucleic acids research **44**, D862-D868 (2016).

30.   Karczewski, K.J. et al. The ExAC browser: displaying reference data information from over 60

000 exomes. *Nucleic acids research* **45**, D840-D845 (2017).

31.   Rentzsch, P., Witten, D., Cooper, G.M., Shendure, J. & Kircher, M. CADD: predicting the

deleteriousness of variants throughout the human genome. *Nucleic acids research* **47**, D886-

D894 (2019).

32.   Li, J., Lai, S., Gao, G.F. & Shi, W. The emergence, genomic diversity and global spread of

SARS-CoV-2. *Nature* **600**, 408-418 (2021).

33.   org, C.-H.G.I.a.b. The COVID-19 host genetics initiative, a global initiative to elucidate the role

of host genetic factors in susceptibility and severity of the SARS-CoV-2 virus pandemic.

European Journal of Human Genetics **28**, 715-718 (2020).

34.   Agius, L. Targeting hepatic glucokinase in type 2 diabetes: weighing the benefits and risks.

Diabetes **58**, 18-20 (2009).

35.   Singh, A.K., Gupta, R., Ghosh, A. & Misra, A. Diabetes in COVID-19: Prevalence,

pathophysiology, prognosis and practical considerations. *Diabetes & Metabolic Syndrome:*

Clinical Research & Reviews **14**, 303-310 (2020).

36.   Pellegrina, D., Bahcheli, A.T., Krassowski, M. & Reimand, J. Human phospho-signaling

networks of SARS-CoV-2 infection are rewired by population genetic variants. *Molecular*

Systems Biology **18**, e10823 (2022).

37.   Radford, A., Narasimhan, K., Salimans, T. & Sutskever, I. Improving language understanding

by generative pre-training. (2018).

38.   Devlin, J., Chang, M.-W., Lee, K. & Toutanova, K. Bert: Pre-training of deep bidirectional

transformers for language understanding. *arXiv preprint arXiv:1810.04805* (2018).

39.   Li, W., Wong, W.H. & Jiang, R. DeepTACT: predicting 3D chromatin contacts via bootstrapping

deep learning. *Nucleic acids research* **47**, e60-e60 (2019).

40.   Chor, B., Horn, D., Goldman, N., Levy, Y. & Massingham, T. Genomic DNA k-mer spectra:

models and modalities. *Genome biology* **10**, 1-10 (2009).

41.   Zhou, Q., Chipperfield, H., Melton, D.A. & Wong, W.H. A gene regulatory network in mouse

embryonic stem cells. *Proceedings of the National Academy of Sciences* **104**, 16438-16443

(2007).

42.   Sharov, A.A. et al. Identification of Pou5f1, Sox2, and Nanog downstream target genes with

statistical confidence by applying a novel algorithm to time course microarray and genome-

wide chromatin immunoprecipitation data. *BMC genomics* **9**, 1-19 (2008).

43.   Raz, R., Lee, C.-K., Cannizzaro, L.A., d'Eustachio, P. & Levy, D.E. Essential role of STAT3 for

embryonic stem cell pluripotency. *Proceedings of the National Academy of Sciences* **96**, 2846-

2851 (1999).

44.   van den Berg, D.L. et al. An Oct4-centered protein interaction network in embryonic stem cells.

Cell stem cell **6**, 369-381 (2010).

45.   Zhang, J. et al. The oncogene Etv5 promotes MET in somatic reprogramming and orchestrates

epiblast/primitive endoderm specification during mESCs differentiation. *Cell death & disease*

9, 224 (2018).

46.   Fang, L. et al. GRNdb: decoding the gene regulatory networks in diverse human and mouse

conditions. *Nucleic acids research* **49**, D97-D103 (2021).

47.   Han, H. et al. TRRUST v2: an expanded reference database of human and mouse transcriptional

regulatory interactions. *Nucleic acids research* **46**, D380-D386 (2018).

48.   Willett, R. et al. TFEB regulates lysosomal positioning by modulating TMEM55B expression

and JIP4 recruitment to lysosomes. *Nature communications* **8**, 1580 (2017).

49.   Bernstein, B.E. et al. The NIH roadmap epigenomics mapping consortium. *Nature*

biotechnology **28**, 1045-1048 (2010).

50.   Kulakovskiy, I.V. et al. HOCOMOCO: towards a complete collection of transcription factor

binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic acids research*

46, D252-D259 (2018).

51.   Heinz, S. et al. Simple combinations of lineage-determining transcription factors prime cis-

regulatory elements required for macrophage and B cell identities. *Molecular cell* **38**, 576-589

(2010).

52.   He, K., Zhang, X., Ren, S. & Sun, J. in Proceedings of the IEEE conference on computer vision

and pattern recognition 770-778 (2016).

53.   Chen, S. et al. OpenAnnotate: a web server to annotate the chromatin accessibility of genomic

regions. *Nucleic Acids Research* **49**, W483-W490 (2021).

Supryware Materials

Supplementary Materials

Fig. S1. Three data partitioning strategies for model training and testing.

Fig. S2. EpiGePT's performance in predicting DNase-seq and other epigenetic signals.

Fig. S3. Performance of EpiGePT and baseline methods on chromatin states classification,

multiple epigenomic profiles prediction and causal variants classification.

Fig. S4. Ablation analysis of the EpiGePT model.

Fig. S5. Case application of the EpiGePT-online.

Fig S6. Model architecture of EpiGePT for multiple epigenomic signals prediction.

Fig. S7. The fine-tuning performance of the EpiGePT model on predicting potential enhancer-

promoter regulatory networks.

Fig. S8. The performance (auROC) of attention score of EpiGePT in distinguishing regulatory

element-gene pairs at different distance ranges.

Text S1. Data splitting strategy for model training.

Text S2. System design and implementation of the web server.

Text S3. Running time of the EpiGePT and baseline methods.

Text S4. Implementation of Enformer model and Enformer+.

Table S1. The information of DNase-seq bam file across 129 biosamples from the ENCODE

project.

Table S2. The information of RNA-seq tab-separated values (tsv) file across 129 biosamples

from the ENCODE project.

Table S3. The information of DNase-seq, CTCF and other six Histone markers bam file across

28 cell lines or tissues from the ENCODE project.

Table S4. The information of RNA-seq tab-separated values (tsv) file across 28 cell lines or

tissues from the ENCODE project.

Table S5. The preprocessed expression data of 711 human transcription factors from the

ENCODE project across 129 biosamples.

Table S6. The preprocessed expression data of 711 human transcription factors from the

ENCODE project across 28 cell lines or tissues.

Table S7. The order and names of epigenomes of the expression matrices across 56 epigenomes

from the ROADMAP project.

Table S8. The preprocessed expression data of 642 human transcription factors across 56

epigenomes from the ROADMAP project.