Title:

Nonlinear feedback modulation contributes to the optimization of flexible decision-making

Xuanyu Wu[1] & Yang Zhou[1,2,3,4]

1. Peking-Tsinghua Center for Life Sciences, Peking University, Beijing, 100871, China
2. School of Psychological and Cognitive Sciences, Peking University, Beijing, 100871, China
3. PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing, 100871, China
4. Department of Neurobiology, The University of Chicago, Chicago, Illinois 60637

* **Corresponding author:** Yang Zhou

Email: yangzhou1@pku.edu.cn

Author Contributions: Y.Z. designed the experiments and wrote the manuscript. Y.Z. trained monkeys and collected the behavioral and neuronal data, Y.Z. and X.W. analyzed the data and made the figures. Y.Z. and X.W. designed the network models, X.W. trained the RNNs, analyzed the RNNs and implemented the RNN inactivation experiment. Y.Z. supervised the project.

Competing Interest Statement: The authors declare no competing financial interests.

Keywords: Decision-making, Monkey electrophysiology, Posterior parietal cortex, Feedback modulation, Artificial neural network. attractor basin

This PDF file includes:

      Main Text
      Figures 1 to 6

Abstract

Neural activity in the primate brain correlates with both sensory evaluation and action selection aspects of decision-making. However, the intricate interaction between these distinct neural processes and their impact on decision behaviors remains unexplored. Here, we examined the interplay of these decision processes in posterior parietal cortex (PPC) when monkeys performed a flexible decision task. We found that the PPC activity related to monkeys' abstract decisions about visual stimuli was nonlinearly modulated by monkeys' following saccade choices directed outside each neuron's response field. Recurrent neural network modeling indicated that the feedback connections, matching the learned stimuli-response associations during the task, might mediate such feedback modulation. Further analysis on network dynamics revealed that selectivity-specific feedback connectivity intensified the attractor basins of population activity underlying saccade choices, thereby increasing the reliability of flexible decisions. These results highlight an iterative computation between different decision processes, mediated primarily by precise feedback connectivity, contributing to the optimization of flexible decision-making.

Main Text

Introduction

Perceptual decisions typically involve the transformation of sensory inputs to discrete motor responses. Consequently, sensory evaluation and action selection emerge as two fundamental processes essential for implementing perceptual decision behavior[1-7]. Neural activity associated with either the sensory or motor aspects of decision-making is widely distributed across different brain areas, such as PPC [8-22], frontal cortex [18,23-28], superior colliculus [29,30], and caudate [31]. A recent inactivation study has shown that the lateral intraparietal area (LIP), a subregion of PPC, plays a causal role in flexible visuomotor decisions, with preferential involvement in sensory evaluation rather than action selection [14]. Meanwhile, areas outside PPC, like frontal eye field and superior colliculus, have been shown to be causally involved in the action selection aspect of decision-making [32,33]. These findings suggest that sensory evaluation and action selection likely involve distinct neural processing in the primate brain during flexible decision-making.

However, previous reports have also shown that neural activity related to sensory evaluation and action selection overlaps temporally and spatially in the brain [34]. This leaves open the possibility of interplay between these distinct processes during decision-making. From the perspective of information processing, the abstract result of sensory evaluation should be transmitted to the motor planning circuits to guide the action selection process in a feedforward manner, although both processes could proceed in parallel in the brain. However, whether and how action selection processes might exert a feedback influence on sensory evaluation have not been explicitly studied.

In this study, we examined the activities of single LIP neurons during a flexible decision task in which monkeys needed to report their decisions about a motion stimulus with a saccade to one of two color targets. Specifically, we arranged the motion stimuli inside each neuron's response field (RF), and positioned the saccade targets in the direction perpendicular to the axis of neural RFs. We found that LIP activity responded to visual motion was nonlinearly modulated by the monkeys' following saccade choice direction relative to the recorded brain hemisphere. Notably, this modulation was aligned precisely with the functional properties of each neuron, as well as specifically impacted the decision-correlated but not the stimulus-correlated activity of LIP neurons. This suggests a precise 'feedback' modulation from action selection process to the neural processing of sensory evaluation during decision-making.

RNN models trained on complex behavioral tasks have shown promise for understanding
neural computations and circuit mechanisms underlying cognitive functions [35-37]. To further
explore how action selection can modulate sensory evaluation during decision-making, we trained
multi-module RNNs, which consist of different hemispheres and RF structures, to perform the
same decision task and analyzed the activity of network units during task performance. These
RNNs exhibited similar behavioral performance and neural activity patterns as observed in the
monkey electrophysiology experiment, including patterns of nonlinear feedback modulation from
action selection to sensory evaluation. Combining analysis on network activity and connectivity
with projection-specific inactivation experiments in the RNNs, we further showed that the precise
feedback connections between units that showed matched functional properties within different
network modules were the key circuit mechanism for mediating the modulation of sensory
evaluation by action selection. Such feedback modulation significantly enhanced the consistency
of the RNNs' decisions by strengthening the attractor basins of network dynamics underlying
saccade choices.
107
108　**Results**
109
Nonlinear modulation of saccade choice on visual motion selectivity in LIP
To test the mechanisms underlying the feedback modulation of action selection on sensory
evaluation, we trained two monkeys to perform a reaction-time version of the flexible visual-
motion discrimination (FVMD) task, in which they needed to choose one of two colored saccade
targets based on their decisions about the motion direction of a sample stimulus shown at
different coherence levels (**Figure 1A**). Monkeys learned the mappings between the two motion
directions (315° and 135°) and two target colors (red and green) at the start of training, and the
mappings were fixed across the study. Because the locations of the red and green targets were
randomly interleaved across trials, neither of the motion directions was directly linked with one
specific saccade direction. Both the performance accuracy and the reaction time (RT) changed
systematically as a function of motion coherence levels (**Figure 1B-1E**): as the motion coherence
increased, both monkeys chose the correct target more frequently and more rapidly.
122
We recorded single-neuron activity from the LIP while the monkeys performed the FVMD
task. We used the memory-guided saccade (MGS) task, which is commonly employed in LIP
studies, to map the receptive fields (RFs) of all isolated LIP neurons. Specifically, we mapped
both the visual and memory RFs of each neuron by analyzing their activity during the target
presentation and delay periods of the MGS task (see Methods). To examine the neural activity
related to the evaluation of stimulus motion, we presented the motion stimuli within the RF of
each neuron, while positioning the saccade targets at locations orthogonal to the line connecting
the center of the RF (which also marks the center of the motion stimulus) and the fixation dot. In
total, 104 of 194 visually responsive LIP neurons (monkey M: 50/78; monkey B: 54/116) showed
significant direction selectivity (DS) to the motion stimuli (one-way ANOVA, $P < 0.01$). To
examine the influence of action selection on sensory evaluation, we analyzed data from the
subset of sessions in which the saccade targets were aligned more closely with the horizontal
direction than the vertical direction (83 of 104 neurons). In these sessions, the motor planning
corresponding to a saccade to either target would be mediated primarily by one brain
hemisphere. We therefore defined the conditions under which the correct target was contralateral
or ipsilateral to the recorded hemisphere as the contralateral target (CT) condition or ipsilateral
target (IT) condition, respectively (**Figure 1F** and **1G**).
Figure 2A-2F shows three example LIP neurons that exhibited significant motion
coherence correlated DS. Surprisingly, LIP neurons showed greater DS in the CT condition than
in the IT condition, even though the same motion stimuli were used in the same spatial location
for both conditions. The averaged population activity showed this DS difference between CT and
IT conditions for all four coherence levels (**Figure 2G, 2H**). During presentation of their preferred
motion direction, LIP neurons showed significantly elevated activity in the CT relative to the IT at
all coherence levels (**Figure S1A, S1B**, nested ANOVA: $P_{(high)} = 0.0326$, $F = 4.65$; $P_{(medium)} =$

0.0088, F = 7.03; $P_{(low)}$ = 0.0076, F = 7.32; $P_{(zero)}$ = 0.0124, F = 6.4), and a trend toward lower
activity to the nonpreferred direction for CT vs. IT (**Figure S1C, S1D,** nested ANOVA: $P_{(high)}$ =
0.0994, F = 2.75; $P_{(medium)}$ = 0.0649, F = 3.12; $P_{(low)}$ = 0.0311, F = 4.73; $P_{(zero)}$ = 0.0273, F = 4.96).
Most of the LIP neurons (48 of 83) showed such opposing trends in activity modulation between
the preferred and nonpreferred directions (**Figure 2I**). These results indicated a nonlinear
modulation of saccade choice on motion DS in LIP, aligned precisely with the response property
of each neuron. This is unlikely to be driven by a linear gain modulation of saccade direction
selectivity. Receiver operating characteristic (ROC) analysis further confirmed significantly
greater motion DS in the CT condition than in the IT condition (**Figure 2J** and **2K**; nested
ANOVA: $P_{(high)}$ = 5.0e-4, F= 12.44; $P_{(medium)}$ = 9.53e-6, F = 20.91; $P_{(low)}$ = 9.33e-7, F = 26.03; $P_{(zero)}$
= 2.56e-8, F= 34.3). Such DS differences were observed even before stimulus onset. Moreover,
LIP neurons exhibited similar levels of mean activity between different saccade directions (CT vs.
IT) before monkeys' saccade choice (**Figure 2L**), further supporting that saccade direction
selectivity did not significantly contribute to the observed modulation of LIP neurons' responses to
motion stimuli.
162
To better quantify the influence of saccade direction on neuronal responses to motion
stimuli, we calculated a modulation index for each neuron (STAR Methods). We then compared
the modulation indices across neurons with different direction preferences and found no
systematic relationship between direction preference and saccade-related modulation in LIP
neurons at the population level (**Figure S2**). Furthermore, a demixed principal component
analysis on the pseudo-population activity (STAR Methods) revealed that the saccade direction
related representation was a substantial component of LIP activity, as the saccade direction and
motion–saccade interaction together explained a similar amount of variance in the population
activity as the stimulus motion direction did (**Figure 2M** and **Figure S3**). In contrast, we did not
observe significant DS differences between the CT and IT conditions in the data sessions in
which the saccade targets were aligned close to the vertical direction (21 of 104 neurons, **Figure
S4**).
175
Decision-correlated but not stimulus-correlated activity was modulated in LIP
We then examined the impact of nonlinear feedback modulation on the correlation between LIP
DS and monkeys' choice behavior. We found that LIP DS was more decision-correlated in the CT
condition than in the IT condition. Illustrated in **Figure 2K**, the LIP DS on zero-coherence trials,
which reflected monkeys' trial-by-trial categorical choice, was significantly greater in the CT
condition than in the IT condition (nested ANOVA: P = 2.56e-8, F= 34.3). We also quantified the
correlation between LIP neural activity and the trial-by-trial categorical choice or the physical
motion direction by comparing LIP neural activity on correct versus incorrect trials. We used low-
coherence trials, as monkeys made enough errors in these trials. In the CT condition but not the
IT condition, LIP DS at the population level was significantly reversed in sign on incorrect trials as
compared to correct trials (**Figure 3A** and **3B**; nested ANOVA: $P_{(CT)}$ = 0.0045, F = 8.32).
Accordingly, in the CT condition but not in the IT condition, LIP activity correlated more closely
with the monkeys' trial-by-trial abstract decisions about motion direction, as opposed to the
physical motion direction (**Figure 3C** and **3D**; nested ANOVA, $P_{(CT)}$ = 0.0016, F = 10.31; $P_{(IT)}$ =
0.443, F = 0.59). Meanwhile, LIP activity correlated more closely with the trial-by-trial abstract
decisions during the CT condition relative to the IT condition on the low coherence trials (nested
ANOVA: P = 1.95e-8, F = 34.94). Furthermore, we used partial correlation analysis to examine
decision- and stimulus-related components of DS (i.e., r-decision and r-stimulus, **Figure 3E** and
3F) using all four coherence levels. The decision-related component of LIP DS was significantly
greater in the CT condition than in the IT condition (**Figure 3E**; nested ANOVA: P = 1.07e-6, F=
25.72), and this difference emerged even before motion stimulus onset. This suggests that the
LIP DS was more closely correlated with monkeys' decisions in the CT condition than in the IT
condition. The upregulation in r-decision for contralateral choices may reflect the monkeys'
internal choice bias or expectation (choice between two motion directions) prior to stimulus
presentation, which could influence their subsequent decisions more in the CT condition.

However, the stimulus-related component of the LIP DS was similar at the population level
between the two conditions (**Figure 3F**; nested ANOVA: P = 0.7606, F = 0.09), suggesting that
the modulation of LIP DS by the later saccade choice was primarily related to a decision process
rather than basic sensory processing.
205
We further examined the impact of nonlinear feedback modulation on the correlation
between LIP DS and the monkeys' RTs. Shortly after motion stimulus onset, LIP neurons showed
greater population response to their preferred motion direction on shorter versus longer RT trials
for all motion coherence levels in the CT condition (**Figure S5**) but not in the IT condition (**Figure**
S6). On zero-coherence trials (**Figure 3G** and **3H**), DS evolved more rapidly on shorter vs. longer
RT trials in the CT condition but not in the IT condition (bootstrap: $P_{(CT)} < 0.01$, $P_{(IT)} = 0.87$).
Together, these results indicated that the feedback modulation of saccade choice on sensory
evaluation predominantly impacted the decision-correlated activity in LIP.
214
Trained multi-module RNNs replicated the nonlinear feedback modulation
Our electrophysiology data show that action selection nonlinearly modulates the neural
processing of sensory evaluation, which indicates a complex, iterative computation for flexible
decision-making. Training RNNs on behavioral tasks used in experimental neurophysiological
studies has proven to be helpful for exploring putative circuit computations underlying cognitive
tasks[28,36,37]. Therefore, we trained multi-module RNNs to perform the FVMD task to examine the
circuits and computation mechanisms underlying the interplay of different decision processes. We
adopted simple neurobiological principles to constrain the connection structure but not the
functional roles to generate recurrently connected modules (STAR Methods). Because our
neurophysiological results showed that the modulation effect of action selection on sensory
evaluation depended on the saccade direction relative to the recorded brain hemisphere, we
implemented RNNs composed of two main modules organized in parallel to simulate two brain
hemispheres. Each main module consisted of two nominal modules, with each nominal module
receiving the visual motion input (motion module) or the target color input (target module),
corresponding to the two neuron populations whose RFs covered the motion stimulus or saccade
targets, respectively (**Figure 4A**).
231
We independently trained 50 such networks with randomly initialized weights and
identical hyperparameters to perform the FVMD task. The FVMD task setup used for training
RNNs was tailored to match the monkey experiments. Specifically, the RNNs were trained using
motion stimuli with two high-coherence levels and were tested using stimuli with another four
different coherence levels (high, medium, low, and zero). After training, all 50 networks
converged to perform the FVMD task with high accuracies (>99%) on the training coherence and
exhibited coherence-dependent performances when tested with untrained stimuli (**Figure 4B** and
4C): accuracy increased and RT decreased with motion coherence.
240
We then analyzed the unit activity in different modules when the trained RNNs were
tested with untrained motion stimuli. Across all networks, the units in the motion module and
target modules exhibited activity corresponding to sensory evaluation and action selection,
respectively. Specifically, units in the motion modules showed coherence-correlated motion DS
(**Figure 4D** and **4F** and **Figure S7** and **S8**; mean r = 0.29; one-sample t-test: P = 4.09e-42, t(49)
= 46.31), and neuronal activity on zero-coherence motion trials reflected the trial-by-trial abstract
decisions of the networks (t-test: $P_{(example\ RNN)}$ = 5.97e-27, t(91) = 15.33; $P_{(population)}$ = 1.05e-31,
t(49) = 27.86). Such motion DS resembled our electrophysiology data when the motion stimuli
were presented within the RFs of the LIP neurons. Meanwhile, units in the target modules
showed coherence-correlated saccade DS (**Figure 4E** and **4G** and **Figure S7** and **S8**), and the
neuronal activity on zero-coherence motion trials also showed significant choice probability (t-test:
$P_{(example\ RNN)}$ = 7.06e-8, t(68) =15.83; $P_{(population)}$ = 2.08e-22, t(49) = 17.21). Such saccade DS was
consistent with the commonly observed decision-related activity in previous studies in which the
saccade target was presented within the RFs of LIP neurons[9-11].

255
Furthermore, we found that the motion DS of units in the motion modules was
significantly greater in the CT condition than in the IT condition for the majority of RNNs (**Figure
4H** and **4I** and **Figure S9A** and **S9B**): in the CT condition relative to the IT condition, activity was
significantly greater for the preferred motion direction but weaker for the nonpreferred direction
(**Figure S10**). ROC analysis confirmed that the motion DS in the motion module was significantly
greater for all four coherence levels in the CT vs. IT conditions in most RNNs (**Figure 4J, 4K** and
4N and **Figure S9C** and **S9D**; two-way ANOVA: $P_{(example RNN)}$ = 1.53e-11, F = 47.49; $P_{(population)}$ =
3.50e-77, F = 660.84). Similar to the monkey electrophysiology data, the decision-correlated
motion DS (r-decision) in the motion module was dramatically reduced in the IT versus CT
condition (paired t-test: P = 6.84e-17, t(49)= -12.53), and this difference was substantially greater
than the difference in the stimulus-related (r-stimulus) motion DS between the two conditions
(paired t-test: P = 2.02e-14, t(49) = -10.70; **Figure 4L, 4M** and **4O** and **Figure S9E** and **S9F**).
These results indicated that action selection also nonlinearly modulated the neural processing of
sensory evaluation during decision-making in the RNNs.
270
Selectivity-specific feedback connections mediate the modulation of sensory evaluation
by action selection
Our multi-module RNNs showed a remarkable similarity to both the behavioral performances and
the neural activity patterns in the monkey electrophysiology data, suggesting that these networks
may have adopted mechanisms similar to those in the monkey brain to mediate the decision
processes in the FVMD task. Therefore, we explored the potential circuit mechanisms underlying
the modulation of action selection on sensory evaluation in RNNs. We hypothesized that this
modulation might result from the feedback connections from the target module to the motion
module. To test this hypothesis, we examined the recurrent weights of the cross-module
connections (STAR Methods). Both the feedback and feedforward cross-module connection
weights were significantly greater between the selectivity-matched unit pairs (e.g., units in the
motion module preferred to encode 315° and units in the target module preferred to encode red)
than the selectivity-nonmatched unit pairs (e.g., units in the motion module preferred to encode
315° while units in the target module preferred to encode green) (**Figure 5A**; paired t-test:
$P_{(feedforward)}$ = 7.18e-16, t(49) = -11.76; $P_{(feedback)}$ = 5.26e-18, t(49) = -13.40). In particular, the
average feedback connectivity was primarily excitatory between the selectivity-matched unit pairs
but was inhibitory on average between the selectivity-nonmatched unit pairs. Furthermore, most
of the networks exhibited positive correlations between the connection weight and the match
extent of the neural selectivity of each cross-module unit pair (**Figure 5B**; mean r = 0.11; one-
sample t-test: P = 6.39e-16; t(49) = 11.79; STAR Methods), indicating that units that exhibited
stronger encoding preference to one motion direction in the motion module were more likely to
receive more extensive feedback projections from the units that exhibited stronger encoding on
the matched target color in the target module. These results suggested precise feedback
connections between RNN modules that were aligned with the functional properties of different
units.
296
To test the causal contribution of the selectivity-specific feedback connections to the
modulation of action selection on sensory evaluation, we conducted three projection-specific
inactivation experiments in the RNNs (STAR Methods). First, we performed nonspecific ablation
of all the feedback projections from the target module to the motion module when testing with
untrained motion stimuli. In most of the tested RNNs, this caused a dramatic reduction in their
performance in all nonzero-coherence stimulus conditions (**Figure 5C**; two-way ANOVA: P =
4.74e-80, F = 704.65) as well as significantly prolonged RTs for high- and medium-coherence
stimulus conditions (**Figure 5D**; two-way ANOVA: P = 2.45e-4, F = 13.79). The mean RTs
became more dispersed across all the trained RNNs and did not differ among different motion
coherences after ablation (one-way ANOVA: P = 0.68, F = 0.50). Ablating feedback connections
significantly affected the motion DS in the motion module. On average, although the averaged
motion DS did not decrease after ablation (**Figure S11A** and **S11C**), the difference in the motion

DS between the CT and IT conditions vanished (**Figure 5E-5G** and **Figure S12A** and **S12B**; two-
way ANOVA: $P_{(CT\ vs.\ IT)}$ = 0.14, F = 2.21). These results indicated a dramatic reduction in the
nonlinear feedback modulation on sensory evaluation by action selection in the RNNs. To further
test the importance of different patterns of feedback connectivity for the RNNs to solve the FVDM
task, we next performed pattern-specific ablation of the feedback connections in RNNs. We
separated the feedback projections in each RNN into specific (i.e., the feedback connection
weights were positive or negative between the selectivity-matched or selectivity-nonmatched unit
pairs, respectively) and nonspecific groups. Despite similar numbers and weights of the feedback
connections in these two groups (**Figure S13A** and **S13B**), ablating the feedback connection in
the specific group versus the nonspecific group resulted in much more severe impairments in the
RNNs' behavior performance (**Figure S13C** and **S13D**). These results indicated that the feedback
connections that reflected the learned stimulus–response association was crucial for the RNNs to
solve the FVDM task.
322
Second, we disrupted the selectivity-specific feedback connectivity without changing the
total strength of the feedback connections by randomly rearranging the feedback connection
weights. This also resulted in a significant reduction in the RNNs' performance accuracy for
nonzero-coherence motion stimuli (**Figure 5C**; two-way ANOVA: P = 1.38e-90, F = 883.61) and
prolonged RTs for high- and medium-coherence motion stimuli (**Figure 5D**; two-way ANOVA: P =
1.94e-12, F = 54.06). The changes of RNN units' activity patterns after scrambling feedback
connections were similar to the effects after feedback connectivity was fully ablated (**Figure
S11B**, **S11C, S14A, S14B, S14D** and **S14E**). Particularly, the levels of nonlinear modulation of
sensory evaluation by saccade selection dramatically decreased, as evident by the diminished
difference in the motion DS between the CT and IT conditions (**Figure 5H**, two-way ANOVA: P =
5.81e-12, F = 51.52).
334
Third, we trained another 50 RNNs without feedback connectivity to learn the FVMD task.
Similar to the full-model RNNs, all 50 networks converged to perform the FVDM task with high
accuracies (>99%) after training and exhibited coherence-dependent performances when tested
with untrained stimuli (**Figure S15A** and **S15B**). Meanwhile, units in the motion module and target
modules exhibited coherence-dependent motion DS and saccade DS, respectively (**Figure S15C**
and **S15D**). Consistent with the above connectivity ablation experiment, units in the motion
module of these RNNs did not exhibit different levels of motion DS between CT and IT conditions
(**Figure S15E-S15G**; two-way ANOVA: P = 0.63, F = 0.24), indicating no noticeable nonlinear
feedback modulation on sensory evaluation by action selection. Importantly, the performance
accuracies of these RNNs decreased significantly when tested with untrained nonzero-coherence
stimuli (**Figure S15H**; two-way ANOVA: P = 1.46e-14, F = 65.64), as compared to the full-model
RNNs. Furthermore, we trained an additional 50 RNNs without feedback connections to learn the
FVMD task. These RNNs were initialized with greater recurrent connection probabilities than the
full-model RNNs, such that the number of total trainable connection weights matched that in the
full-model RNNs. Interestingly, these RNNs still exhibited significantly lower performance
accuracies when tested with high- and medium-coherence stimuli relative to the full-model RNNs
(**Figure S16**; two-way ANOVA: P = 8.24e-6, F = 20.60). Together, these results suggested that
the nonlinear feedback modulation, which was mediated by the selectivity-specific feedback
connections, were important for the RNNs to efficiently solve flexible decision tasks.
354
Nonlinear feedback modulation enhanced the decision consistency by strengthening the
attractor basins of network dynamics
Our RNN modeling indicates a crucial role for nonlinear feedback modulation in optimizing flexible
decision-making. Subsequently, we delved into the computational mechanisms embodied by
nonlinear feedback modulation during the decision-making process. Specifically, we examined
the unit activity in the target module, a direct contributor to the decision outputs of RNNs.
Strikingly, the projection-specific inactivation experiments led to a significant reduction in the
magnitude of saccade DS in the nonzero-coherence trials (**Figure 6A-6C**, $P_{(ablating\ feedback)}$ = 6.90e-

5, F = 16.29; $P_{(shuffling\ feedback)}$ = 1.2e-19, F = 95.08; two-way ANOVA). Additionally, saccade DS
emerged earlier than the motion stimulus onset in the nonzero-coherence trials, and its positive
correlation with the motion coherence levels diminished after the projection-specific inactivation
(**Figure 6A-C** and **Figure S12C, S14C** and **S14F**; mean $r_{(ablating\ feedback)}$ = -0.034; mean $r_{(shuffling}$
$_{feedback)}$ = −0.030). These disrupted patterns of saccade DS observed in the target module
following projection-specific inactivation aligned with the decreased decision consistency of
RNNs, where decision consistency reflects the degree of agreement in the model's choices under
specific task conditions. This suggests a diminished reliance on sensory input and an increased
dependence on internal noise in the decision-making process.
372
Previous studies have demonstrated that attractor dynamics in networks can explain
choice consistency, with steeper landscapes around attractor basins reflecting consistent
decisions[38-42]. Consequently, we investigated attractor dynamics in the state space of network
activity that underlies decision-making variability in RNNs with different types of feedback
connectivity. Similar to prior studies[40], we examined the neural dynamics underlying decisions in
a 1-D neural subspace using unit activity in the target module responsible for saccade choices
(see methods). Beginning with the reconstruction of a numerical approximation of the energy
landscape in this 1-D subspace, we explored changes in this landscape after different types of
projection-specific inactivation. This revealed a dramatic reduction in the depth of attractor basins
in the energy landscape of population activity after ablating all feedback connectivity (**Fig. 6D-E,**
paired t-test: P = 2.28e-11, t(49) = -8.61). Moreover, ablating the feedback connections led to a
more severe reduction in the depth of attractor basins in the network dynamics within the specific
group compared to the nonspecific groups (**Fig. 6F-G**, paired t-test: P = 1.59e-6 , t(49) =-5.46).
386
We further examined how the energy landscape in the 1-D subspace changed in relation
to task difficulty (motion coherence). Consistent with prior findings[40], trials with lower decision
consistency (trials using lower motion coherence) exhibited shallower attractor basins at the time
of decision for all types of RNNs (**Fig. 6H-K**). However, both the depth and the positional
separation of attractor basins in the network dynamics significantly decreased for all non-zero
motion coherence levels after the ablation of all feedback connections (comparing **Figure 6I** with
Figure 6H; $P_{(depth)}$ = 5.20e-25, F = 122.80; $P_{(position)}$ = 1.82e-27, F = 137.75; two-way ANOVA).
Notably, this reduction in basin depth and separation was more pronounced in the specific group
compared to the nonspecific groups after ablating the feedback connections (comparing **Figure
6J** with **Figure 6K**; $P_{(depth)}$ = 2.65e-13, F =57.35; $P_{(position)}$ = 3.73e-14, F = 61.79; two-way
ANOVA). These results might underlie the computational mechanisms that explain the observed
reduction in the decision consistency of RNNs following projection-specific inactivation: the
shallower and closer attractor basins after ablating feedback connections resulted in less
consistent decisions. This happened because the variability in neural activity made it more likely
for population activity to stochastically shift out of the shallower basins and into nearby alternative
ones.
403
404
Discussion
Here we show that primate LIP activity related to evaluating the visual motion stimulus was
nonlinearly modulated by an impending saccade choice that was used to report decisions during
a FVMD task, even though the saccades were toward non-RF locations. This suggests that the
sensory evaluation and action selection may proceed iteratively during decision-making. This
view is also consistent with the common observation that action selection related activity arises at
the early stage of evidence accumulation during decision-making[3,9,34], as well as our observation
that the modulation of action selection on sensory evaluation emerged during the early task
period even before motion stimulus onset in the FVMD task. Brain areas that are related to
different decision processes, although spatially separated[14,21,33], likely form a real-time associated
network to solve current decision tasks. The instantaneous result of sensory evaluation may be
transmitted to the brain areas responsible for action selection during decision-making in a

feedforward manner, and the action selection process may also exert a real-time modulation on
the neural processing of sensory evaluation in a feedback manner.
419
The feedback modulation we observed during decision-making was distinct from the
modulation of feature-based attention in early sensory cortex. Although both modulation effects
match the stimulus tuning of recipient cells[43-47], only the feedback modulation observed in the
current study reflects the neural processing of decision-making. This is because only the
decision-related component, not the stimulus-related component, of LIP activity was modulated
by the monkeys' later saccade choice during the FVMD task (**Figure 3**). This suggests that the
feedback modulation from action selection to LIP activity was primarily related to the decision
process rather than low-level sensory processing.
428
A recent study demonstrated that neurons in the middle temporal area responded more
strongly to motion stimuli when monkeys saccaded toward their RFs in a standard decision task
with a fixed mapping between motion stimuli and saccade directions[48]. This modulation emerged
through the training process and contributed causally to the monkeys' following saccade choices.
Consistently, we found that the response of LIP neurons to motion stimuli was more strongly
correlated with the monkeys' decisions in the CT condition (saccades toward RFs) than in the IT
condition, in a more flexible decision task. Together, these results suggest that the modulation of
action selection on sensory processing may be a general process in perceptual decision-making.
However, the observed modulation of saccade direction on LIP neurons' responses to motion
stimuli cannot be simply explained by saccade direction selectivity. Several lines of evidence
argue against this possibility. First, the modulation effect was nonlinear; specifically, neuronal
firing rates increased for preferred motion directions but decreased for non-preferred directions
(**Figure 2I** and Figure S1). This pattern is unlikely to be driven by a linear gain modulation based
on saccade directions. Second, we found that LIP neurons exhibited similar levels of activity in
both the CT and IT conditions (Figure 2M), which is inconsistent with the presence of clear
saccade direction selectivity.
445
The selectivity-specific feedback modulation from action selection to sensory evaluation
emerged in our multi-module RNNs during training, although the cross-module connections were
randomly initialized before training. Importantly, our causal in-silico experiments demonstrated
that these precise feedback connections played an essential role in mediating the modulation of
action selection on sensory evaluation while solving the FVDM task. Previous studies showed
that both the corticocortical feedback from the secondary visual cortex to the primary visual cortex
and the corticogeniculate feedback in primates are organized into parallel streams resembling the
reciprocal feedforward pathways[49-51], suggesting a potential functionally specific feedback
connection in visual processing. However, to the best of our knowledge, a similar segregation of
feedback that reflects the encoding properties of the target neurons has not been evident in
decision network in vivo. Therefore, it will be important for future studies to examine the
connectivity and correlation of neural activity among neural groups related to different decision
processes through sophisticated anatomical experimental approaches as well as multi-channel
recordings in vivo.
460
Our brain includes extensive feedback connections across different brain areas, which, in
some cases, even outnumber the feedforward connections[51,52]. However, compared to the
feedforward connections, much less is known about the function of feedback connections.
Feedback modulation has been implicated in top-down modulation of neuronal responses in the
early sensory cortex, such as attention and expectation, which facilitate the processing of
important stimuli and suppress distractors[53-55]. Furthermore, both experimental and computational
studies have shown that weak decision-correlated neural activity (i.e., choice probability) as well
as the correlated firing of pairs of neurons (i.e., noise correlation) in the early sensory cortex
might partially result from feedback input in decision tasks[56-58], suggesting a potential role for
feedback connection in modulating early sensory processing during decision-making.

Consistently, the nonlinear feedback modulation, mediated primarily by selectivity-specific
feedback connections, were important for our multi-module RNNs to efficiently solve the FVMD
task, as the RNNs' decision behavior became more stochastic following ablating/disrupting the
feedback connectivity. This was evident not only in the diminished behavioral performance of the
RNNs but also in the disrupted patterns of activity related to saccade choice in the target module.
Notably, the nonlinear feedback modulation intensified the attractor basins of the population
activity associated with saccade choice, led to more reliable decisions based on sensory input.
Our results unveil a novel pattern of feedback modulation during the neural processing of
decision-making, and suggest a potentially critical role that feedback modulation plays in
increasing the consistency of flexible sensorimotor decisions.
481
Two-state attractor models have been widely employed to elucidate two-alternative
forced decision-making processes[38-42,59,60]. In these models, decisions arise when network activity
falls into one of two attractor basins. Previous studies have posited that decision consistency is
influenced by the configuration of the energy landscape surrounding these attractor basins[40,42,60].
Essentially, steep-sided and deep-basin energy landscapes contribute to consistent decision-
making, as it becomes challenging for internal noise to shift activity between basins. Conversely,
when energy landscapes are relatively flat and attractor basins are shallow, decisions are
generated more stochastically, as it is easier for internal noise to drive activity out of the attractor
basins and into alternative ones. Consistently, our RNNs modelling showed that shallower and
less separated attractor basins were associated with decreased decision consistency in more
difficult trials. Building upon previous research, our results provide in-silico evidence supporting
the notion that the energy landscape within the state space of population activity causally
influences decision consistency. Moreover, our findings present the initial circuit mechanism
through which the configuration of the energy landscape for decision-related activity is modulated
by feedback modulation, to our knowledge. Future studies are needed to explore the connections
between different types of circuit connectivity and population activity dynamics in cognitive
processing in vivo.
499
Our results are consistent with predictive coding theories of feedback function, which
propose that an internal model of the world is generated in the brain based on sensory data and
prior experience and is refined by incoming sensory data[61-64]. In terms of the architecture of
hierarchical predictive coding schemes, different neuronal ensembles encode different
attributes/choices in either the action selection or sensory evaluation process at each level of the
cortical hierarchy during flexible decision-making. Meanwhile, the conditionally independent
expectations are functionally segregated, so that descending predictions from the action selection
process are limited to the sensory evaluation process on the matched\associated stimulus
attribute but not the opposite attribute. The selectivity-specific feedback connectivity in our RNNs
fits the functional segregation of expectation very well during the interplay between distinct neural
processes of decision-making. Specifically, the neural ensembles related to either sensory
evaluation or action selection processes for the same choice might form precise recurrently
connected loops. The prediction signal carried by the precise feedback connections might
facilitate sensory evaluation of the associated stimulus but prohibit evaluation of the nonmatched
stimulus. This could, in turn, amplify the task-relevant sensory input, facilitate the sensorimotor
transformation, and ultimately result in faster and more accurate action choice in flexible
decisions. In future work, it will be important to use techniques such as projection-specific
inactivation and microstimulation in vivo to test the causal contribution of feedback connectivity in
decision-making and other cognitive functions.
519
520
521 **Materials and Methods**
522
Behavioral task, stimulus display, and animal preparation

The flexible visual motion discrimination (FVMD) task (**Figure 1A**) has been reported previously[14]
and is briefly summarized below. In this task, monkeys were required to saccade to either the
green or red targets based on the direction of the visual motion stimulus. Two motion directions
(135°, 315°) were used, each with four different coherence levels (0%, 9%, 18%, 36% for monkey
M; and 0%, 13%, 25%, 50% for monkey B) were tested. If the sample direction was 135°, the
monkeys must saccade to the green target to receive a juice reward, whereas the 315° direction
was associated with the red target. The rewarded target (red or green) was randomly chosen
(with 50% probability) on each zero-coherence trial. The positions of red and green targets were
randomly chosen between the two positions on each trial at each recording session. Therefore,
there is no fixed mapping between motion stimulus and saccade direction.
534
To initiate each trial, monkeys must hold a touch-bar and acquire gaze fixation. They then
needed to maintain fixation within a 2.0-2.5° window throughout the trial before their saccade
choice. After a 500ms fixation period, two colored saccade targets appeared simultaneously at
opposite positions relative to the fixation point with equal eccentricities (8° and 9° for Monkey M
and B, respectively). 400ms later, a sample motion stimulus was presented at a location
orthogonal to the axis of, but at the same eccentricity as, the saccade targets. We used motion
stimuli that were full contrast, 8° diameter, random-dot movies composed of 190 dots per frame,
and the dots moved at 10°/s. Monkeys needed to saccade to either red or green targets within a
60-2000ms window after sample stimulus onset. The two saccade targets were equidistant from
the stimulus, with the distance typically ranging from 12 to 15 degrees. A juice reward would be
delivered to the monkeys if they made correct saccade choice.
546
Two male monkeys (Macaca mulatta, 15~16 years old, 8–14 kg) were trained on the
FVMD task and implanted with a head post as well as a recording chamber positioned over PPC.
Our surgical, behavioral, and neurophysiological approach has been described in detail in a
previous study[14] . Stereotaxic coordinates for chamber placement were determined from
magnetic resonance imaging (MRI) scans obtained before chamber implantation. LIP chambers
were centered on the intraparietal sulcus (IPS), 4.0 mm posterior to the intra-aural line and 1.0
mm lateral from the midline for monkey M, and 0 mm anterior to the intra-aural line and 15.0 mm
lateral from the midline for monkey B. Monkeys were housed in individual cages under a 12-hour
light/dark cycle. Behavioral training and experimental recordings were conducted during the light
portion of the cycle. Monkeys sat comfortably while head-fixed in a custom-made primate chair
inside a dark experiment rig. The task stimuli were displayed on a 21-inch color CRT monitor
(1280*1024 resolution, 75 Hz refresh rate, 57 cm viewing distance). Both monkeys were tested
with identical stimuli and timing. A solenoid-operated reward system was used to deliver juice
reward to the monkeys. Monkeys' eye positions were monitored by an optical eye tracker (SR
Research) at a sampling rate of 1 kHz and stored for offline analysis. Stimulus presentation, task
events, rewards, and behavioral data acquisition were accomplished using an Intel-based PC
equipped with MonkeyLogic software running in MATLAB (http://www.monkeylogic.net). All
experimental and surgical procedures were in accordance with the University of Chicago Animal
Care and Use Committee and National Institutes of Health guidelines.
566
Electrophysiological recording
We used either 75-μm tungsten microelectrodes (FHC, ~1 MΩ) or 16-channel V-Probes (Plexon)
to record single neuron activity in LIP. Neurophysiological signals were amplified, digitized, and
stored for offline spike sorting (Plexon) to verify the quality and stability of neuronal isolations. We
recorded neuronal activity in a memory-guided saccade task to map LIP RF locations before each
FVMD recording session.
573
Our LIP recordings targeted different hemispheres in the two monkeys (monkey M: left
hemisphere; monkey Q: right hemisphere). Therefore, the contralateral target (CT) condition and
ipsilateral target (IT) conditions in **Figure 1F** and **1G** referred to opposite saccade directions
(target locations) between the two monkeys. For monkey M, the CT condition corresponded to

the trials in which the correct saccade target was on the right visual field, whereas the CT
condition for monkey Q referred to the trials in which the correct saccade was toward the left
visual field. We localized LIP in each monkey according to the patterns of neuronal activity in the
MGS task (i.e., spatial selectivity during stimulus presentation and delay). All neurons included in
the dataset were recorded from the same locations (the same grid holes and similar depths: ~5-
10 mm from the cortical surface) where we encountered spatial selectivity in the MGS task. LIP
neurons were also identified based on anatomical criteria, such as the location of each electrode
track relative to that expected from the MRI scans, the pattern of gray–white matter transitions
encountered on each electrode penetration, and the relative depths of each neuron.
587
We aimed to present the visual motion stimulus inside the RFs of the identified neurons
during each recording session. For single-channel electrode recording, only neurons exhibited
visual responses to the motion stimuli during prescreening (tested with the FVMD task) were
recorded with sufficient trials (~300-600) of the FVDM task. For neurons exhibiting clear spatial
RFs during the MGS task, we presented the motion stimulus inside LIP neurons' RFs; while for
those neurons which did not show a clear RF during the MGS task, we presented motion stimuli
in the positions (always in the visual field contralateral to the recorded hemisphere) in which
neurons exhibited the strongest response to the motion stimuli. For the multi-channel recordings,
we recorded all neurons isolated across all channels and presented the motion stimulus inside
one of the isolated neurons' RFs. Because adjacent recording sites were located 100μm apart,
nearby neurons typically showed similar RF locations.
599
600 ***Data analysis***
601
Neuronal pre-screening
While part of the neural data was presented in a previous report[14], the current analysis focuses
on a different phenomenon which was not examined previously. We included all neurons
recorded from single-channel electrodes for the analysis. For multi-channel recordings, we only
included the neurons which showed significant modulation (different from fixation period activity,
one-way ANOVA, $p < 0.01$) of their averaged activity across all motion stimuli because the
stimulus could not always be presented within the RF of all the recorded neurons. The low-firing
neurons whose maximum firing rates were less than 2.0 spikes/s (to the direction producing
greater average responses) during stimulus presentation were also excluded for further analysis.
We then applied a one-way ANOVA test to compare activity between the two different motion
directions during the period following motion stimulus onset (50-250 ms after motion stimulus
onset) to select neurons that showed significant motion direction selectivity (DS) during the
decision period. Only neurons that showed significant ($p < 0.01$) DS were used for further
analysis.
616
Receiver operating characteristic (ROC) analysis
To quantify each neuron's DS in the FVMD task, we applied an ROC analysis to the distribution
of firing rates within sliding windows (100 ms width, 5 ms steps). The ROC value, which ranges
between 0.0 and 1.0, indicates the performance of an ideal observer in assigning motion direction
based on each neuron's trial-by-trial firing rates. Values of 1.0 and 0.0 correspond to perfect
classification (i.e., strong DS), while a value of 0.5 indicates chance-level classification
performance (i.e., no DS). For trials with zero coherence motion, we assigned direction labels on
each trial based on the monkey's choice.
625
In **Figure 3**, in order to test whether DS reflected monkeys' trial-by-trial choices, we used
an ROC analysis to quantify whether LIP activity correlated with monkeys' trial-by-trial categorical
choices more than the physical direction of motion stimulus by analyzing both correct and error
trials. We only included low coherence trials in this analysis because there were sufficient
numbers of error trials (average performance: Monkey M: 74% correct, Monkey B: 67% correct).
LIP neuronal activity was analyzed by ROC according to either the monkey's trial-by-trial

categorical choices or the direction of the sample stimulus on each trial. Furthermore, for this
analysis, we only used neurons for which we recorded sufficient trials (for both correct and error
trials, N>4) for the low coherence condition of each motion direction (n = 45).
635
Partial correlation analysis
We performed a partial correlation analysis to quantify the correlation between LIP neural activity
and the monkeys' trial-by-trial categorical choices or the physical motion direction of the stimulus
using all trials. For each trial, we obtained three parameters for the calculation:　the stimulus
direction, the pre-choice neuronal activity, and the monkeys' choice. We assigned the stimulus
directions with different values for different directions and coherence levels: positive and negative
values were used for preferred and nonpreferred directions, respectively; while 4, 2, 1 and 0 are
used for code the high, medium, low, and zero coherence levels. We also assigned different
values to different choice directions (–2 for choosing the preferred direction, and + 2 for choosing
the nonpreferred direction). Two measures were then calculated: r stimulus = r (neuronal activity,
stimulus direction| choice direction), the partial correlation between neuronal activity and stimulus
direction given the monkeys' choices; and r choice = r (neuronal activity, choice direction |
stimulus direction), the partial correlation between neuronal activity and the monkeys' categorical
choice given the stimulus direction. In **Figure 3**, we used the mean activity within a sliding
window (100ms width, sliding with 5ms steps) for each neuron to perform the partial correlation
analysis.
652
653　*Modulation Index*
654　We have calculated a modulation index for each neuron to reflect the influence of saccade
655　direction on neuron's response to visual stimuli. The modulation index is calculated as:

$$\delta_{\text{contra}} = r_{pref}^{contra} - r_{non-pref}^{contra}$$

$$\delta_{ipsi} = r_{pref}^{ipsi} - r_{non-pref}^{ipsi}$$

$$\text{MI} = \frac{\delta_{\text{contra}} - \delta_{\text{ipsi}}}{\delta_{\text{contra}} + \delta_{\text{ipsi}}}$$

656　, where r_{pref}^{contra} represents the average firing rate from 50ms to 250ms after sample onset for all
657　contralateral saccade trails with a neuron's preferred moving direction of visual stimuli. The
658　naming conventions are the same for $r_{non-pref}^{contra}$, r_{pref}^{ipsi} , and $r_{non-pref}^{ipsi}$. An MI value between 0 and
659　1 indicate higher modulation in contralateral saccade trials, and an MI value between -1 and 0
660　indicates higher modulation in ipsilateral saccade trials.
661
dPCA analysis
We conducted demixed principal component analysis using the methodology and the code from a
previous study[65] (http://github.com/machenslab/dPCA) to reduce the dimensionality of the
population activity as the standard PCA and demixes all task variables. Specifically, we tested
how much each task variable (motion direction of sample stimuli, saccade direction, motion-
saccade interaction, timing) contributes to the LIP population activity during the DMC task.
668
669　　　As demonstrated in the previous study, the dPCA finds separate decoder (F) and
670　encoder (D) matrices for each task variable (∅) by minimizing the loss function:

$$L_{dPCA} = \sum_0 \left\| X_0 - F_0 D_0 X \right\|^2$$

671
672　where X is a linear decomposition of the data matrix, which contains the instantaneous firing rate
673　of the recorded neurons, into variable-specific averages:

$$X = \sum_0 X_0 + X_{noise}$$

675 Here, we decomposed the neural activities into four parts: condition-independent, stimulus-
676 dependent (two motion directions with four coherence levels), saccade-dependent (two saccade
677 directions), dependent on the stimulus-saccade interaction, and noise. The decoder and encoder
678 axes permit us to reduce the data into a few components capturing the majority of the variance of
679 the data dependent on each task variable.
680
681 ***Recurrent neural network (RNN) training***
682
Network Implementation
We trained biologically inspired networks to perform the FVMD task using methodology similar to
previous studies[28,36]. We implemented multiple modules in the networks through constraints on
the input/output structure and the initial recurrent connectivity of the hidden layer. Specifically, we
built excitatory-inhibitory networks with 200 hidden units divided into two equal-size main modules
to simulate two brain hemispheres. Each main module was further divided into two nominal
modules, with each one only receiving the visual motion input (motion module) or target color
input (target module), respectively. This design was intended to simulate the two neuron
populations whose RFs covered the motion stimulus or saccade target in the monkey
electrophysiology experiment (**Figure 4A**). Every nominal module was allocated one quarter of
the excitatory units (40) and one quarter of the inhibitory units (10) in the overall network to
ensure that the modules did not differ in their balance of excitation/inhibition prior to training.
Meanwhile, in order to simulate the local and long-range connection structures in the brain, we
set different levels of recurrent connectivity within and between different modules: the local
(connectivity within each nominal module) recurrent connection density (probability) was the
highest (50%); the across-RF (connectivity between the two nominal modules within the same
main module) connection density was set to be in the medium level (25%); the cross-hemisphere
(connectivity between either the motion modules or the target modules across different main
modules) connection density was the lowest (10%). Only excitatory neurons could have "cross-
hemisphere" projections to the corresponding nominal module (e.g., from the motion module of
the "left hemisphere" to the motion module of the "right hemisphere"). Only the connection weight
within the hidden layer was updated during training following methods described previously[28]. In
addition, the connection weights between network units are endowed with short-term synaptic
plasticity, which is aimed to provide connection weights with activity-dependent fluctuation over
short timescales within each trial.
708
709 The input to the network consisted of two parts: 9 motion input units and 8 target color
710 input units. The preferred directions of motion input units were evenly spaced across 360°, with
711 response tuning distributed according to a von Mises function. The value of the nth motion input
712 unit was set to

$$r_n = \frac{2}{3}\left(ace^{\left(\kappa cos(\theta - \theta_{pref})\right)} + b\mathcal{N}(0,2) + \sqrt{\frac{2}{\alpha}}\mathcal{N}(0, \sigma_{in}) \right) + d$$

713 , where $\alpha = \frac{\Delta t}{\tau_{mem}}$, θ is the direction of motion stimulus in radian, θ_{pref} is the preferred direction of
714 this motion input unit, $a = \frac{4}{e^{\kappa}}$ after stimulus onset but 0 elsewhere, $c \in [0, 1]$ is the coherence of
715 the dot motion stimulus, b is a binary value that equals 0 before stimulus onset and 1 after that,
716 and d is a constant visual input signal with an amplitude of 2 that decays with time. Specifically,
717 for zero-coherence input, since $ace^{\left(\kappa cos(\theta - \theta_{pref})\right)}$ would be zero, we set this term as $0.4max(c)$.
718
The target input units were initially evenly divided into two groups projected to the two
target modules of hidden units. Each group was further divided into red and green subgroups.
The color subgroup was set as active with an amplitude of $\frac{4}{3}$ to represent the color within the
projected target module, while the other color group remained silent (amplitude =0). An

exponential decay filter was also used to fit the activity of sensory neurons in the early sensory
cortex across time:

$$f(x) = \left(\frac{1}{3}\right)^t x$$

725 , where x is the constant visual input for motion input and the color signal for color input, t is time
726 in miliseconds.
727
The two output units of the network simulate two different saccade directions. In order to
simulate the oculomotor control in the brain, each brain hemisphere (main module) projects much
more densely to the contralateral response units (probability = 0.32) than the ipsilateral response
units (probability = 0.08). To generate a probability distribution over output values at each time
point, we also applied a softmax function to scale the response unit activities in the output layer.
To reduce the stochasticity of the network activity brought by input and output connections, we
randomly sampled input and output weights from a normal distribution ($\mathcal{N}(0.2, 0.05)$). In addition,
the input weights were re-initialized if there were extreme values (the minimum value was smaller
than ¾ of the maximum value) after multiplying the stimulus signal and weight of motion or target
input. Similarly, the output weights were re-initialized if there were extreme values in the output
weight values. The input units had excitatory and random projections to the recurrent units with a
probability of 0.32. However, only the excitatory units in the target modules of each "hemisphere"
could project to the output units. Both the input and output weights were fixed for each network
during task training.
742
Network training
The networks were trained using BrainPy[66] on an Intel(R) Core(TM) i9-9900K CPU (3.60GHz, 8
cores). The network was optimized using backpropagation through time (BPTT) and stochastic
gradient descent with an Adam optimizer (default setting, first moment estimates decay rate=0.9,
second moment estimates decay rate =0.999) to minimize a loss function. The network
parameters (recurrent weights/biases) were optimized to minimize a loss function with three parts
as in a previous study[28]: (a) a performance loss; (b) a metabolic cost on mean neuronal activity;
and (c) a metabolic cost on connectivity. The gradient was clipped to a maximum L2 value of 0.1
to avoid the exploding gradient problem. During training, only the hidden unit related parameters
(weights, bias, and initial activities) were updated. The training process was terminated when the
network's performance accuracy reached 99% or till the maximum number of iterations (2000).
754
During each trial of the training and testing of the FVMD task, networks were first
presented with two color targets through the target input units, and then presented with the
motion direction through the motion input units. Both the motion and target inputs persisted until
the end of each trial. A short time after the motion input (100ms, 5 time steps), the networks were
required to report the direction of the stimulus motion by choosing the saccade target with the
appropriate color. Specifically, each element of the task design in the FVMD task that the models
were trained to perform was tailored to match those used in the monkey experiments: the
directions of the motion stimuli, the target colors, as well as the task (stimulus) durations were the
same as that used in monkey electrophysiology experiments. Trials were programmatically
generated by constructing motion/target inputs to the networks at each timestep and the desired
response (left saccade, right saccade) at each timestep. In total, we trained 50 networks to
perform the FVDM task. All networks were first trained using motion directions with two
coherence levels (60% and 90%) and were then tested using motion stimuli with another four
different coherence levels (75%, 55%, 35%, and zero), which had never been presented during
training. Besides the difference in the coherence of motion input, the color inputs remained the
same during the testing period. All the networks achieved consistently high performance by the
end of training (> 99% accuracy). All the important model hyperparameters were listed in **Table
S1**.
773
Quantification of the networks' behavioral performance

To test whether our multi-module RNNs exhibited decision behavior similar to monkey subjects,
we examined their performance accuracies and RTs when tested by novel stimuli with different
coherence levels. The performance accuracy was defined as whether the output of the response
units (starting from 100ms after stimulus onset to the trial end) matched the desired output.
Furthermore, we defined the networks' RTs as the time point from which the differences between
two output units were greater than a threshold (0.8) for three consecutive time points after the
stimulus onset. During some trials, the threshold was never reached until the end of the trial
(500ms after the motion stimulus onset). In these cases, we artificially set the RT to be 600ms in
these trials for further analysis.
784
Analysis of RNN activity
We performed the same analyses on the RNN units as we did on the neurophysiology data. In
Figure 4-5, data from both the example network and averaged results across all the networks
were shown. For every network, we only included the units which exhibited at least one kind of
task-related modulation during the test period (i.e., motion direction selectivity or saccade
direction selectivity, one-way ANOVA test, $p < 0.01$). In order to quantify the activity related to
evaluating motion stimulus, we analyzed the motion DS only including the units in the motion
modules, whereas only the saccade DS of units in the target modules were calculated to quantify
the activity related to saccade selection during the decision-making.
794
Analysis of RNN connectivity
To examine the potential circuit mechanisms underlying the modulation of action selection on
sensory evaluation during decision-making in the RNNs, we examined the cross-module
connection weights in the trained networks. Specifically, we defined the feedforward connection
as the projection from units in the motion module to units in the target module within the same
main module (hemisphere). We also defined the feedback connection as the projection from units
in the target module to units in the motion module within the same main module. Furthermore, we
grouped the cross-module connections into the selectivity-matched and selectivity-mismatched
groups based on the preferences of the neural encoding of units in the motion and target
modules. The selectivity-matched group included two types of unit pairs: 1) units in the motion
module preferred to encode 315° and units in the target module preferred to encode red, 2) units
in the motion module preferred to encode 135° and units in the target module preferred to encode
green. The selectivity-mismatched group includes the other two types of unit pairs: 3) units in the
motion module preferred to encode 315° while units in the target module preferred to encode
green, and 4) units in the motion module preferred to encode 135°while units in the target module
preferred to encode red. The values of connection weights projected from excitatory and
inhibitory units were set as positive and negative, respectively.
812
We also examined the correlation between the similarity of neural selectivity and
connection strength for different unit pairs in the RNNs based on the following steps: First, both
the preference and strength of motion direction encoding or target color encoding were quantified
for units in the motion module or target module, respectively. This was done by calculating the
averaged differential activity to the 315° and 135° motion stimuli after motion stimulus onset or
calculating the averaged differential activity to the red and green targets after target onset. In this
case, the unit pairs that exhibited matched selectivity would show selectivity values with the same
sign (positive or negative), whereas the unit pairs that exhibited mismatched selectivity would
show selectivity values with the opposite signs. Second, the units in both the motion module and
target modules were ranked based on selectivity values. Third, the selectivity similarity of each
cross-module unit pair was quantified by calculating the reverse value of the absolute differences
between the target color selectivity rank and the motion direction selectivity rank. The rank of the
weight value of each pair was calculated and then reversed to give the lower weight value a
higher rank score. Finally, we calculated the Pearson correlation between weight rank and
selectivity similarity. A positive correlation indicates that units that exhibited stronger encoding

preference to one motion direction in the motion module were more likely to connect with the
units in the target module that exhibited stronger encoding on the matched target color.
830
Inactivation experiments in silico
Our examination of the network connectivity within the hidden layer of the RNNs revealed that the
selectivity-specific top-down connections originated from the target module to the motion module.
To assess the causal contribution of such precise feedback connections to the modulation of
action selection on sensory evaluation, we performed two in-silico analogues of neuronal
inactivation experiments similar to a previous study[28]. We hypothesized that ablating or disrupting
the precise feedback connection from the target module to the motion module would significantly
impact the activity patterns of the units in the motion module. Specifically, such inactivation on
connectivity was expected to reduce the difference in motion DS in the notion module between
CT and IT conditions. To test this hypothesis, we performed the projection-specific inactivation
experiments in the RNNs in three different ways. First, we ablated all the feedback projections
from the target module to the motion module while keeping all the other network parameters
unchanged after the networks were fully trained. We then tested the networks with the untrained
motion stimuli used in the normal experiment. The connectivity ablation was implemented by
directly setting the connection weights from units in the target module to units in the motion
module to zero. Second, we randomly rearranged the connection weights from the target module
to the motion module after the network was fully trained by shuffling the corresponding weights in
the weight matrix. Then, we tested the networks with the testing motion stimuli. This was aimed to
disrupt the selectivity-specific feedback connection without changing the total strength of the
feedback connection. We repeated this random feedback weight rearrangement process 100
times for each of the 50 trained networks. Subsequently, we measured the impact on the
networks' behavior performance and the activity patterns in both the motion and saccade
modules. In particular, for each network, we tested whether there were still different levels of
motion direction selectivity of units in the motion module between CT and IT conditions after
inactivating the feedback connectivity.
856
We further performed pattern-specific ablation of the feedback connections in the RNNs,
in order to examine the causal roles of different patterns of feedback connectivity in solving the
FVDM task. Specifically, we separated the feedback projections in each RNN into the specific
group and the nonspecific group. The specific group included two conditions: 1) the feedback
connection weight was positive between the selectivity-matched unit pairs, and 2) the feedback
connection weight was negative between the selectivity-nonmatched unit pairs; whereas the
nonspecific group included the rest of the feedback conditions in each RNN. We then tested the
effects of the pattern-specific connectivity inactivation on the RNNs' behavior performance when
tested with the untrained motion stimuli.
866
In the third inactivation experiment, we trained 50 additional RNNs without feedback
connections to learn the FVMD task, in order to test the importance of feedback connectivity in
solving the flexible decision task. These networks were initialized using identical hyperparameters
and trained with the same methodology as the normal RNNs, except that the feedback
connections from the target module to the motion module were ablated before training. We then
examined these RNNs' behavior performance and unit activity when testing with the untrained
motion stimuli used in the normal experiment. Furthermore, we trained another 50 RNNs without
feedback connection to learn the FVMD task. These RNNs were initialized with higher recurrent
connection probabilities than the full-model RNNs to keep the number of the total trainable
connection weight comparable to that in the full-model RNNs. We raised the recurrent connection
probability by a factor of 1.176 after removing feedback connections (average number of
connections in 50 normal RNNs = 8193, average number of connections in 50 networks without
feedback = 8184, p = 0.53, independent sample t-test). Except for the recurrent connection
probability, all the other hyperparameters and the training methodology were the same as the

normal RNNs. We then compared the behavior performance of these RNNs with that of the
normal RNNS when testing with the untrained motion stimuli.
883
Analysis of neural landscape
For each network, we adopted a methodology similar to a previous study (cite) to reconstruct the
energy landscape of population activity within the target module. Initially, units in the target
module exhibiting saccade direction selectivity were identified. Subsequently, we employed
principal component analysis (PCA) to reduce the dimensionality of the population activity.
Specifically, PCA was performed on the average activity across these units, considering each
stimulus motion direction, coherence level, and choice. We retained the first five principal
components (PCs) for further analysis, as they explained over 90% of the total variance in
population activity. Following this, the trial-by-trial population activities were transformed into 5-
dimensional data using the aforementioned five PCs.
894
To predict trial-by-trial choice, we employed a linear Support Vector Machine (SVM)
classifier with 10-fold cross-validation. Specifically, we utilized the average activity of the last 100
ms of each trial as input for the classifier. The choice axis was determined by selecting the
normal vector of the separating hyperplane from the SVM classifier that demonstrated the best
performance. Subsequently, the 5-dimensional data points were projected onto the choice axis,
resulting in 1-dimensional projections. These projections represented the positions along the
choice axis, with the intersection between the normal vector and the hyperplane serving as the
zero point.
The potential in this context is determined by integrating the spatial component of the
time derivative of the population activity, denoted as X_t , for a given condition. To be more
precise, we began by computing the expected value across trials of the time derivative of the unit
activities for each choice. Subsequently, we employed the potential function $V(x, t)$ to to evaluate
the potential at position x and time t,

$$V(x,t) = \left(\sum_{\mathrm{p}<x} \frac{dX_t}{dt} | X_t = p \right) \cdot (-\Delta x)$$

908 Crucially, we computed the potential separately for each choice, establishing the potential
909 at position 0 as the zero-potential reference. To provide a holistic evaluation of the potential
910 landscape, we combined the potentials of both choices at position 0.
911
Furthermore, we expanded the computation of potentials to encompass each stimulus
coherence level using the aforementioned approach. Similar to the prior analysis, the time
derivatives were still derived from the 1-D projections of each data point onto the choice axis.
However, in this instance, the conditions were based on the stimulus coherence level rather than
the choice. For a representative potential value for each model, we averaged the potentials at the
time point of 300ms after the stimulus onset. Additionally, recognizing that the positions visited by
each model might differ, we ensured the accuracy of standard error estimation by retaining only
the potential values at positions continuously visited by four or more models.
920
921
922 **Acknowledgments**
923

924 We thank Tianming Yang, Matthew Rosen, Siyu Wang, Cheng Xue, Gongcheng Yu, Mingze Li
925 and Ou Zhu for their comments on an earlier version of this manuscript. We also thank the
926 veterinary staff at The University of Chicago Animal Resources Center for expert assistance.
927
928 This study was supported by STI2030-Major Projects (2021ZD0203800), NSFC32171036.

929
930
References

1　O'Connell, R. G., Shadlen, M. N., Wong-Lin, K. & Kelly, S. P. Bridging Neural and
Computational Viewpoints on Perceptual Decision-Making. *Trends Neurosci* **41**, 838-852,
doi:10.1016/j.tins.2018.06.005 (2018).
2　Freedman, D. J. & Assad, J. A. Neuronal Mechanisms of Visual Categorization: An
Abstract View on Decision Making. *Annu Rev Neurosci* **39**, 129-147,
doi:10.1146/annurev-neuro-071714-033919 (2016).
3　Gold, J. I. & Shadlen, M. N. The neural basis of decision making. *Annu Rev Neurosci* **30**,
535-574, doi:10.1146/annurev.neuro.29.051605.113038 (2007).
4　Huk, A. C., Katz, L. N. & Yates, J. L. The Role of the Lateral Intraparietal Area in (the Study
of) Decision Making. *Annu Rev Neurosci* **40**, 349-372, doi:10.1146/annurev-neuro-
072116-031508 (2017).
5　Hanks, T. D. & Summerfield, C. Perceptual Decision Making in Rodents, Monkeys, and
Humans. *Neuron* **93**, 15-31, doi:10.1016/j.neuron.2016.12.003 (2017).
6　Shadlen, M. N. & Kiani, R. Decision making as a window on cognition. *Neuron* **80**, 791-
806, doi:10.1016/j.neuron.2013.10.047 (2013).
7　Freedman, D. J. & Assad, J. A. A proposed common neural mechanism for categorization
and perceptual decisions. *Nat Neurosci* **14**, 143-146, doi:10.1038/nn.2740 (2011).
8　Platt, M. L. & Glimcher, P. W. Neural correlates of decision variables in parietal cortex.
Nature **400**, 233-238, doi:10.1038/22268 (1999).
9　Roitman, J. D. & Shadlen, M. N. Response of neurons in the lateral intraparietal area
during a combined visual discrimination reaction time task. *J Neurosci* **22**, 9475-9489
(2002).
10　Shadlen, M. N. & Newsome, W. T. Motion perception: seeing and deciding. *Proceedings*
of the National Academy of Sciences of the United States of America **93**, 628-633,
doi:10.1073/pnas.93.2.628 (1996).
11　Shadlen, M. N. & Newsome, W. T. Neural basis of a perceptual decision in the parietal
cortex (area LIP) of the rhesus monkey. *J Neurophysiol* **86**, 1916-1936,
doi:10.1152/jn.2001.86.4.1916 (2001).
12　Sugrue, L. P., Corrado, G. S. & Newsome, W. T. Matching behavior and the
representation of value in the parietal cortex. *Science* **304**, 1782-1787,
doi:10.1126/science.1094765 (2004).
13　Yates, J. L., Park, I. M., Katz, L. N., Pillow, J. W. & Huk, A. C. Functional dissection of signal
and noise in MT and LIP during decision-making. *Nat Neurosci* **20**, 1285-1292,
doi:10.1038/nn.4611 (2017).
14　Zhou, Y. & Freedman, D. J. Posterior parietal cortex plays a causal role in perceptual and
categorical decisions. *Science* **365**, 180-185, doi:10.1126/science.aaw8347 (2019).
15　Freedman, D. J. & Assad, J. A. Experience-dependent representation of visual categories
in parietal cortex. *Nature* **443**, 85-88, doi:10.1038/nature05078 (2006).
16　Freedman, D. J. & Assad, J. A. Distinct encoding of spatial and nonspatial visual
information in parietal cortex. *J Neurosci* **29**, 5671-5680, doi:10.1523/JNEUROSCI.2878-
08.2009 (2009).
17　Bennur, S. & Gold, J. I. Distinct representations of a perceptual decision and the
associated oculomotor plan in the monkey lateral intraparietal area. *J Neurosci* **31**, 913-
921, doi:10.1523/JNEUROSCI.4417-10.2011 (2011).

18   Swaminathan, S. K. & Freedman, D. J. Preferential encoding of visual categories in
parietal cortex compared with prefrontal cortex. *Nat Neurosci* **15**, 315-320,
doi:10.1038/nn.3016 (2012).
19   Zhou, Y., Mohan, K. & Freedman, D. J. Abstract Encoding of Categorical Decisions in
Medial Superior Temporal and Lateral Intraparietal Cortices. *J Neurosci* **42**, 9069-9081,
doi:10.1523/JNEUROSCI.0017-22.2022 (2022).
20   Kiani, R. & Shadlen, M. N. Representation of confidence associated with a decision by
neurons in the parietal cortex. *Science* **324**, 759-764, doi:10.1126/science.1169405
(2009).
21   Katz, L. N., Yates, J. L., Pillow, J. W. & Huk, A. C. Dissociated functional significance of
decision-related activity in the primate dorsal stream. *Nature* **535**, 285-288,
doi:10.1038/nature18617 (2016).
22   Zhou, Y., Zhu, O. & Freedman, D. J. Posterior Parietal Cortex Plays a Causal Role in
Abstract Memory-Based Visual Categorical Decisions. *J Neurosci* **43**, 4315-4328,
doi:10.1523/JNEUROSCI.2241-22.2023 (2023).
23   Kim, J. N. & Shadlen, M. N. Neural correlates of a decision in the dorsolateral prefrontal
cortex of the macaque. *Nat Neurosci* **2**, 176-185, doi:10.1038/5739 (1999).
24   Rossi-Pool, R. *et al.* Decoding a Decision Process in the Neuronal Population of Dorsal
Premotor Cortex. *Neuron* **96**, 1432-1446 e1437, doi:10.1016/j.neuron.2017.11.023
(2017).
25   Ding, L. & Gold, J. I. Neural correlates of perceptual decision making before, during, and
after decision commitment in monkey frontal eye field. *Cereb Cortex* **22**, 1052-1067,
doi:10.1093/cercor/bhr178 (2012).
26   Ferrera, V. P., Yanike, M. & Cassanello, C. Frontal eye field neurons signal changes in
decision criteria. *Nat Neurosci* **12**, 1458-1462, doi:10.1038/nn.2434 (2009).
27   Heitz, R. P. & Schall, J. D. Neural mechanisms of speed-accuracy tradeoff. *Neuron* **76**,
616-628, doi:10.1016/j.neuron.2012.08.030 (2012).
28   Zhou, Y. *et al.* Distributed functions of prefrontal and parietal cortices during sequential
categorical decisions. *Elife* **10**, doi:10.7554/eLife.58782 (2021).
29   Horwitz, G. D. & Newsome, W. T. Separate signals for target selection and movement
specification in the superior colliculus. *Science* **284**, 1158-1161,
doi:10.1126/science.284.5417.1158 (1999).
30   Peysakhovich, B. *et al.* Primate superior colliculus is engaged in abstract higher-order
cognition. *bioRxiv*, doi:10.1101/2023.01.17.524416 (2023).
31   Ding, L. & Gold, J. I. Caudate encodes multiple computations for perceptual decisions. *J
Neurosci* **30**, 15747-15759, doi:10.1523/JNEUROSCI.2894-10.2010 (2010).
32   Gold, J. I. & Shadlen, M. N. Representation of a perceptual decision in developing
oculomotor commands. *Nature* **404**, 390-394, doi:10.1038/35006062 (2000).
33   Jun, E. J. *et al.* Causal role for the primate superior colliculus in the computation of
evidence for perceptual decisions. *Nat Neurosci* **24**, 1121-1131, doi:10.1038/s41593-
021-00878-6 (2021).
34   Shushruth, S., Mazurek, M. & Shadlen, M. N. Comparison of Decision-Related Signals in
Sensory and Motor Preparatory Responses of Neurons in Area LIP. *J Neurosci* **38**, 6350-
6365, doi:10.1523/JNEUROSCI.0668-18.2018 (2018).

1021	35	Engel, T. A., Chaisangmongkon, W., Freedman, D. J. & Wang, X. J. Choice-correlated
1022		activity fluctuations underlie learning of neuronal category representation. *Nat Commun*
1023		**6**, 6454, doi:10.1038/ncomms7454 (2015).
1024	36	Masse, N. Y., Yang, G. R., Song, H. F., Wang, X. J. & Freedman, D. J. Circuit mechanisms
1025		for the maintenance and manipulation of information in working memory. *Nat Neurosci*
1026		**22**, 1159-1167, doi:10.1038/s41593-019-0414-3 (2019).
1027	37	Song, H. F., Yang, G. R. & Wang, X. J. Training Excitatory-Inhibitory Recurrent Neural
1028		Networks for Cognitive Tasks: A Simple and Flexible Framework. *PLoS Comput Biol* **12**,
1029		e1004792, doi:10.1371/journal.pcbi.1004792 (2016).
1030	38	Deco, G., Rolls, E. T., Albantakis, L. & Romo, R. Brain mechanisms for perceptual and
1031		reward-related decision-making. *Prog Neurobiol* **103**, 194-213,
1032		doi:10.1016/j.pneurobio.2012.01.010 (2013).
1033	39	Finkelstein, A. *et al.* Attractor dynamics gate cortical information flow during decision-
1034		making. *Nat Neurosci* **24**, 843-850, doi:10.1038/s41593-021-00840-6 (2021).
1035	40	Wang, S., Falcone, R., Richmond, B. & Averbeck, B. B. Attractor dynamics reflect decision
1036		confidence in macaque prefrontal cortex. *Nat Neurosci* **26**, 1970-1980,
1037		doi:10.1038/s41593-023-01445-x (2023).
1038	41	Wang, X. J. Probabilistic decision making by slow reverberation in cortical circuits.
1039		*Neuron* **36**, 955-968, doi:10.1016/s0896-6273(02)01092-9 (2002).
1040	42	Wong, K. F., Huk, A. C., Shadlen, M. N. & Wang, X. J. Neural circuit dynamics underlying
1041		accumulation of time-varying evidence during perceptual decision making. *Front*
1042		*Comput Neurosci* **1**, 6, doi:10.3389/neuro.10.006.2007 (2007).
1043	43	Motter, B. C. Neural correlates of attentive selection for color or luminance in
1044		extrastriate area V4. *J Neurosci* **14**, 2178-2189, doi:10.1523/JNEUROSCI.14-04-
1045		02178.1994 (1994).
1046	44	Motter, B. C. Neural correlates of feature selective memory and pop-out in extrastriate
1047		area V4. *J Neurosci* **14**, 2190-2199, doi:10.1523/JNEUROSCI.14-04-02190.1994 (1994).
1048	45	Martinez-Trujillo, J. C. & Treue, S. Feature-based attention increases the selectivity of
1049		population responses in primate visual cortex. *Curr Biol* **14**, 744-751,
1050		doi:10.1016/j.cub.2004.04.028 (2004).
1051	46	Bichot, N. P., Rossi, A. F. & Desimone, R. Parallel and serial neural mechanisms for visual
1052		search in macaque area V4. *Science* **308**, 529-534, doi:10.1126/science.1109676 (2005).
1053	47	Maunsell, J. H. & Treue, S. Feature-based attention in visual cortex. *Trends Neurosci* **29**,
1054		317-322, doi:10.1016/j.tins.2006.04.001 (2006).
1055	48	Laamerad, P., Liu, L. D. & Pack, C. C. Decision-related activity and movement selection in
1056		primate visual cortex. *Sci Adv* **10**, eadk7214, doi:10.1126/sciadv.adk7214 (2024).
1057	49	Federer, F., Ta'afua, S., Merlin, S., Hassanpour, M. S. & Angelucci, A. Stream-specific
1058		feedback inputs to the primate primary visual cortex. *Nat Commun* **12**, 228,
1059		doi:10.1038/s41467-020-20505-5 (2021).
1060	50	Briggs, F. & Usrey, W. M. Parallel processing in the corticogeniculate pathway of the
1061		macaque monkey. *Neuron* **62**, 135-146, doi:10.1016/j.neuron.2009.02.024 (2009).
1062	51	Briggs, F. Role of Feedback Connections in Central Visual Processing. *Annu Rev Vis Sci* **6**,
1063		313-334, doi:10.1146/annurev-vision-121219-081716 (2020).
1064	52	Harris, K. D. & Mrsic-Flogel, T. D. Cortical connectivity and sensory coding. *Nature* **503**,
1065		51-58, doi:10.1038/nature12654 (2013).

53　Moore, T. & Armstrong, K. M. Selective gating of visual signals by microstimulation of
frontal cortex. *Nature* **421**, 370-373, doi:10.1038/nature01341 (2003).
54　Gilbert, C. D. & Li, W. Top-down influences on visual processing. *Nature reviews.*
Neuroscience **14**, 350-363, doi:10.1038/nrn3476 (2013).
55　McManus, J. N., Li, W. & Gilbert, C. D. Adaptive shape processing in primary visual
cortex. *Proceedings of the National Academy of Sciences of the United States of America*
108, 9739-9746, doi:10.1073/pnas.1105855108 (2011).
56　Nienborg, H. & Cumming, B. G. Decision-related activity in sensory neurons reflects
more than a neuron's causal effect. *Nature* **459**, 89-92, doi:10.1038/nature07821 (2009).
57　Bondy, A. G., Haefner, R. M. & Cumming, B. G. Feedback determines the structure of
correlated variability in primary visual cortex. *Nat Neurosci* **21**, 598-606,
doi:10.1038/s41593-018-0089-1 (2018).
58　Wimmer, K. *et al.* Sensory integration dynamics in a hierarchical network explains choice
probabilities in cortical area MT. *Nat Commun* **6**, 6177, doi:10.1038/ncomms7177
(2015).
59　Prat-Ortega, G., Wimmer, K., Roxin, A. & de la Rocha, J. Flexible categorization in
perceptual decision making. *Nat Commun* **12**, 1283, doi:10.1038/s41467-021-21501-z
(2021).
60　Wong, K. F. & Wang, X. J. A recurrent network mechanism of time integration in
perceptual decisions. *J Neurosci* **26**, 1314-1328, doi:10.1523/JNEUROSCI.3733-05.2006
(2006).
61　Rao, R. P. & Ballard, D. H. Predictive coding in the visual cortex: a functional
interpretation of some extra-classical receptive-field effects. *Nat Neurosci* **2**, 79-87,
doi:10.1038/4580 (1999).
62　Friston, K. & Kiebel, S. Predictive coding under the free-energy principle. *Philos Trans R*
Soc Lond B Biol Sci **364**, 1211-1221, doi:10.1098/rstb.2008.0300 (2009).
63　Bastos, A. M. *et al.* Canonical microcircuits for predictive coding. *Neuron* **76**, 695-711,
doi:10.1016/j.neuron.2012.10.038 (2012).
64　Shipp, S. Neural Elements for Predictive Coding. *Front Psychol* **7**, 1792,
doi:10.3389/fpsyg.2016.01792 (2016).
65　Kobak, D. *et al.* Demixed principal component analysis of neural population data. *Elife* **5**,
doi:10.7554/eLife.10989 (2016).
66　Chaoming Wang, X. C., Tianqiu Zhang, Si Wu. BrainPy: a flexible, integrative, efficient,
and extensible framework towards general-purpose brain dynamics programming.
bioRxiv, doi:https://doi.org/10.1101/2022.10.28.514024 (2022).

1101

Figures

1104
1105
Figure 1. Behavioral task.
(**A**) The flexible visual-motion discrimination (FVMD) task. Monkeys needed to report their
decision about the direction of the visual motion stimuli by choosing either the green or red
saccade target. The appearance of the two color targets preceded that of the visual-motion
stimulus, and the target positions were randomly chosen on each trial to avoid fixed mapping
between motion stimulus and saccade direction. Monkeys could initiate their saccade as soon as
they had made their decision. (**B-C**) Psychometric curves for the two monkeys. The averaged
performance accuracy from all recording sessions for each monkey is plotted as the proportion of
trials in which 315° was chosen as a function of the directions and coherence levels of the motion
stimuli. (**D-E**) Chronometric curves are shown separately for the two monkeys. (**F-G**) Two trial
conditions, contralateral (F) and ipsilateral (G), defined according to the spatial configurations of
task stimuli during neural recording.
1118
1119
1120

Figure 2. Nonlinear feedback modulation of saccade choice on sensory evaluation in LIP.
(**A**) The activity of one example neuron in the CT condition of the FVMD task is shown for each motion-coherence level. The zero-coherence trials were grouped based on the monkey's choices. The two vertical dashed lines denote the time of target and motion stimulus onset, respectively. (**B**) The same example neuron's activity in the IT condition of the FVMD task. (**C-F**) The activities of two more example neurons. (**G-H**) The averaged population activities of all direction-selective neurons (n = 83) that were collected during the recording sessions in which the saccade targets were arranged in either horizontal or oblique directions. The activity to each motion direction and coherence level is shown separately for the CT condition (G) and IT condition (H). (**I**) The activity differences between CT and IT conditions (CT minus IT) of single LIP neurons were plotted for both preferred and nonpreferred motion directions. Each dot represents the activity of a single neuron. The histograms in horizontal and vertical axis represent the distribution of activity difference between CT and IT conditions for preferred and nonpreferred motion directions, respectively. (**J**) An ROC analysis was used for quantifying the motion DS for both CT (solid) and IT (dashed) conditions. The colored dots denote the time points for which there was a significant difference between the CT and IT conditions (paired t-test: P < 0.01). (**K**) The average DS for low- and zero-coherence trials is shown as in (J). (**L**) The mean activity (averaged across all motion directions and coherence levels) was compared between the two saccade directions (CT vs. IT) at the population level. Activity was aligned to either motion stimulus onset (left panel) or saccade onset (right panel).There was no significant difference between CT and IT conditions before monkeys made saccade choices. (**M**) Variance in LIP population activity as explained by the individual demixed principal components. Each bar shows the proportion of total explained variance that was contributed by the four task variables. The pie chart shows the total variance explained by each task variable. H, high; M, medium; L, low; 0, zero; P, preferred; NP nonpreferred.

1148
1149

1150
1151
Figure 3. Feedback modulation specifically impacted the decision-correlated activity .
(**A-B**) Averaged population activities on low-coherence trials in the CT condition (A) and IT
condition (B) are shown separately for correct (corr) and error (err) trials. (**C-D**) An ROC analysis
quantified the stimulus-related and decision-related LIP DS on low-coherence trials. The
correlations between LIP neural activity and the monkeys' decisions about motion direction (red)
or the physical direction of the motion stimulus (blue) in both the CT (C) and IT (D) conditions are
shown over time. (**E-F**) Partial correlation analysis revealed the decision-related and stimulus-
related components of LIP activity. The values for r-decision (E) (the partial correlation between
neuronal activity and monkeys' choice, given the stimulus direction) and r-stimulus (F) (partial
correlation between neuronal activity and stimulus direction, given the monkeys' choices) were
compared between the IT and CT conditions. (**G-H**) Correlation between LIP DS and the
monkeys' RTs on zero-coherence trials. The choice selectivity on zero-coherence trials is shown
for the faster RT and slower RT trials. Shaded areas denote ±SEM. The black stars indicate time
periods in which there was a significant difference (paired t-test: P < 0.01).
1166
1167

1168
1169

Figure 4. Multi-module RNNs trained with the FVMD task.
(**A**) Model schematic of the RNNs. Each RNN consists of nine motion direction tuned input units,
eight color tuned target input units, 200 hidden units, and two response units. The hidden layer of
each RNN consists of two main modules. Each main module consists of two nominal modules,
each of which receives either the visual motion input (motion module) or the target color input
(target module). Only the target modules project to the two response units, and each main
module projects primarily to one response unit. All four nominal modules were assigned with
equal number of units (25%) in the network, which consisted of 80% excitatory and 20% inhibitory
units. (**B-C**) The performance accuracies (B) and RTs (C) of all 50 trained RNNs are shown
separately for each motion coherence level. (**D-E**) Two example units from an example RNN. (**D**)
The neural activity of an example unit from the motion module is shown for each motion direction
and coherence level. (**E**) The neural activity of an example unit from the target module is shown
for each saccade direction and coherence level. (**F-G**) The motion DS for the motion module (F)
and saccade DS for the target module (G) for the example RNN. (**H-I**) The averaged population
activities of all direction-selective units in the motion module of the example RNN are shown for
the CT condition (H) and IT condition (I). (**J-K**) The averaged motion DS in the motion module of
the example RNN for both the CT (solid) and IT (dashed) conditions was quantified by ROC
analysis. (**L-M**) Partial correlation analysis. The values for r-decision (L) and r-stimulus (M) are
compared between the IT and CT conditions for the example RNN. (**N**) A comparison of the
motion DS in the motion module of all trained RNNs between the CT and IT conditions is shown
for each coherence level. (**O**) A comparison of the r-decision and r-stimulus between the IT and
CT conditions is shown for all trained RNNs. (Paired t-test: ***, P < 0.001).

1192
1193

Figure 5. The circuit mechanisms underlying the nonlinear feedback modulation in RNNs.
(**A**) The averaged cross-module connection weights are shown for both feedforward and
feedback connections. Units in the motion module (M) and target module (T) were grouped based
on their preferences for motion DS (315° vs. 135°) and target color selectivity (red [r] vs. green
[g]), respectively. (**B**) The correlation between the match extent of the neural encoding between
units in the motion module and those in target modules and the connection weights between
them. Each dot denotes data from one RNN. (**C-D**) A comparison of the performance accuracy
(C) and RT (D) of the full-model RNNs, RNNs without feedback connections, and RNNs with
shuffled feedback connections. (**E-F**) The averaged activity of units in the motion module of the
example RNN after the feedback connections were ablated is shown for CT (E) and IT (F)
conditions. (**G**) The comparison of the averaged motion DS between CT and IT conditions for all
trained RNNs after feedback connections were ablated. An ROC analysis was used to quantify
the motion DS. (**H**) The differences in motion DS between CT and IT conditions were compared
between full-model RNNs and RNNs with shuffled feedback connectivity.

1210
1211

1212

Figure 6. The computational mechanisms underlying the nonlinear feedback modulation in decision-making.

(**A**) The averaged activity of units in the target module of the example RNN, after the feedback connections were ablated, is shown separately for different saccade directions and coherence levels. (**B**) The averaged saccade DS in the target module of the example RNN after full feedback ablation. (**C**) The averaged saccade DS in the target module of the full-model RNNs, RNNs without feedback connections, and RNNs with shuffled feedback connections. (Paired t-test: *, P < 0.05; **, P < 0.01; ***, P < 0.001; n.s., not significant). (**D-G**) The evolution of the averaged energy landscapes was shown over time. A numerical approximation of the energy landscape in the 1-D decision (saccade choice) subspace is constructed for both full-model RNNs and RNNs with various types of projection-specific inactivation. Each plot represents the averaged results of 50 RNNs, where Position 0 signifies the SVM decision boundary, and the vertical dashed line marks the time of motion stimulus onset. Unvisited portions of the state space are left blank as there is no gradient or potential estimate. (**H-k**) Averaged numerical estimate of energy landscapes for trials with different task difficulty levels (motion coherence). Results from two different saccade choices were average together. Only the potential values at positions continuously visited by four or more models were retained. Shaded areas denote ±SEM.

1230