Title:

Nonlinear feedback modulation contributes to the optimization of flexible decision-making

Xuanyu Wu[1] & Yang Zhou[1,2,3,4]

1. Peking-Tsinghua Center for Life Sciences, Peking University, Beijing, 100871, China
2. School of Psychological and Cognitive Sciences, Peking University, Beijing, 100871, China
3. PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing, 100871, China
4. Department of Neurobiology, The University of Chicago, Chicago, Illinois 60637

* **Corresponding author:** Yang Zhou

Email: yangzhou1@pku.edu.cn

Author Contributions: Y.Z. designed the experiments and wrote the manuscript. Y.Z. trained monkeys and collected the behavioral and neuronal data, Y.Z. and X.W. analyzed the data and made the figures. Y.Z. and X.W. designed the network models, X.W. trained the RNNs, analyzed the RNNs and implemented the RNN inactivation experiment. Y.Z. supervised the project.

Competing Interest Statement: The authors declare no competing financial interests.

Keywords: Decision-making, Monkey electrophysiology, Posterior parietal cortex, Feedback modulation, Artificial neural network. attractor basin

This PDF file includes:

      Main Text
      Figures 1 to 6

Abstract

Neural activity in the primate brain correlates with both sensory evaluation and action selection aspects of decision-making. However, the intricate interaction between these distinct neural processes and their impact on decision behaviors remains unexplored. Here, we examined the interplay of these decision processes in posterior parietal cortex (PPC) when monkeys performed a flexible decision task. We found that the PPC activity related to monkeys' abstract decisions about visual stimuli was nonlinearly modulated by monkeys' following saccade choices directing outside each neuron's response field. Recurrent neural network modeling indicated that the feedback connections, matching the learned stimuli-response associations during the task, mediated such feedback modulation. Further analysis on network dynamics revealed that selectivity-specific feedback connectivity intensified the attractor basins of population activity underlying saccade choices, thereby increasing the consistency of flexible decisions. These results highlight an iterative computation between different decision processes, mediated primarily by precise feedback connectivity, contributing to the optimization of flexible decision-making.

Main Text

Introduction

Perceptual decisions typically involve the transformation of sensory inputs to discrete motor responses. Consequently, sensory evaluation and action selection emerge as two fundamental processes essential for implementing perceptual decision behavior[1-7]. Neural activity associated with either the sensory or motor aspects of decision-making is widely distributed across different brain areas, such as PPC [8-22], frontal cortex [18,23-28], superior colliculus [29,30], and caudate [31]. A recent inactivation study has shown that the lateral intraparietal area (LIP), a subregion of PPC, plays a causal role in flexible visuomotor decisions, with preferential involvement in sensory evaluation rather than action selection [14]. Meanwhile, areas outside PPC, like frontal eye field and superior colliculus, have been shown to be causally involved in the action selection aspect of decision-making [32,33]. These findings suggest that sensory evaluation and action selection likely involve distinct neural processing in the primate brain during flexible decision-making.

However, previous reports have also shown that neural activity related to sensory evaluation and action selection overlaps temporally and spatially in the brain [34]. This leaves open the possibility of interplay between these distinct processes during decision-making. From the perspective of information processing, the abstract result of sensory evaluation should be transmitted to the motor planning circuits to guide the action selection process in a feedforward manner, although both processes could proceed in parallel in the brain. However, whether and how action selection processes might exert a feedback influence on sensory evaluation have not been explicitly studied.

In this study, we examined the activities of single LIP neurons during a flexible decision task in which monkeys needed to report their decisions about a motion stimulus with a saccade to one of two color targets. Specifically, we arranged the motion stimuli inside each neuron's response field (RF), and positioned the saccade targets in the direction perpendicular to the axis of neural RFs. We found that LIP activity responded to visual motion was nonlinearly modulated by the monkeys' following saccade choice direction relative to the recorded brain hemisphere. Notably, this modulation was aligned precisely with the functional properties of each neuron, as well as specifically impacted the decision-correlated but not the stimulus-correlated activity of LIP neurons.

90   This suggests a precise 'feedback' modulation from action selection process to the neural
91   processing of sensory evaluation during decision-making.
92
RNN models trained on complex behavioral tasks have shown promise for understanding
neural computations and circuit mechanisms underlying cognitive functions [35-37]. To further explore
how action selection can modulate sensory evaluation during decision-making, we trained multi-
module RNNs, which consist of different hemispheres and RF structures, to perform the same
decision task and analyzed the activity of network units during task performance. These RNNs
exhibited similar behavioral performance and neural activity patterns as observed in the monkey
electrophysiology experiment, including patterns of nonlinear feedback modulation from action
selection to sensory evaluation. Combining analysis on network activity and connectivity with
projection-specific inactivation experiments in the RNNs, we further showed that the precise
feedback connections between units that showed matched functional properties within different
network modules were the key circuit mechanism for mediating the modulation of sensory
evaluation by action selection. Such feedback modulation significantly enhanced the consistency
of the RNNs' decisions by strengthening the attractor basins of network dynamics underlying
saccade choices.
107
108 **Results**
109
Nonlinear feedback modulation of saccade choice on visual motion selectivity in LIP
To test the mechanisms underlying the feedback modulation of action selection on sensory
evaluation, we trained two monkeys to perform a reaction-time version of the flexible visual-motion
discrimination (FVMD) task, in which they needed to choose one of two colored saccade targets
based on their decisions about the motion direction of a sample stimulus shown at different
coherence levels (**Figure 1A**). Monkeys learned the mappings between the two motion directions
(315° and 135°) and two target colors (red and green) at the start of training, and the mappings
were fixed across the study. Because the locations of the red and green targets were randomly
interleaved across trials, neither of the motion directions was directly linked with one specific
saccade direction. Both the performance accuracy and the reaction time (RT) changed
systematically as a function of motion coherence levels (**Figure 1B-1E**): as the motion coherence
increased, both monkeys chose the correct target more frequently and more rapidly.
122
We recorded single-neuron activity from the LIP while the monkeys performed the FVMD
task. To examine the neural activity related to the evaluation of stimulus motion, we presented the
motion stimuli within the RF of each neuron but arranged the saccade targets at locations
orthogonal to the axis of the neural RF. In total, 104 of 194 visually responsive LIP neurons (monkey
M: 50/78; monkey B: 54/116) showed significant direction selectivity (DS) to the motion stimuli (one-
way ANOVA, $P < 0.01$). To examine the influence of action selection on sensory evaluation, we
analyzed data from the subset of sessions in which the saccade targets were aligned more closely
with the horizontal direction than the vertical direction (83 of 104 neurons). In these sessions, the
motor planning corresponding to a saccade to either target would be mediated primarily by one
brain hemisphere. We therefore defined the conditions under which the correct target was
contralateral or ipsilateral to the recorded hemisphere as the contralateral target (CT) condition or
ipsilateral target (IT) condition, respectively (**Figure 1F** and **1G**).
135
Figure 2A-2F shows three example LIP neurons that exhibited significant motion
coherence correlated DS. Surprisingly, LIP neurons showed greater DS in the CT condition than in
the IT condition, even though the same motion stimuli were used in the same spatial location for
both conditions. The averaged population activity showed this DS difference between CT and IT
conditions for all four coherence levels (**Figure 2G, 2H**). During presentation of their preferred
motion direction, LIP neurons showed significantly elevated activity in the CT relative to the IT at
all coherence levels (**Figure S1A, S1B**, nested ANOVA: $P_{(high)} = 0.0326$, $F = 4.65$; $P_{(medium)} = 0.0088$,
$F = 7.03$; $P_{(low)} = 0.0076$, $F = 7.32$; $P_{(zero)} = 0.0124$, $F = 6.4$), and a trend toward lower activity to the

nonpreferred direction for CT vs. IT (**Figure S1C, S1D,** nested ANOVA: $P_{(high)}$ = 0.0994, F = 2.75; $P_{(medium)}$ = 0.0649, F = 3.12; $P_{(low)}$ = 0.0311, F = 4.73; $P_{(zero)}$ = 0.0273, F = 4.96). Most of the LIP neurons (48 of 83) showed such opposing trends in activity modulation between the preferred and nonpreferred directions (**Figure 2I**). Receiver operating characteristic (ROC) analysis further confirmed significantly greater motion DS in the CT condition than in the IT condition (**Figure 2J** and **2K**; nested ANOVA: $P_{(high)}$ = 5.0e-4, F= 12.44; $P_{(medium)}$ = 9.53e-6, F = 20.91; $P_{(low)}$ = 9.33e-7, F = 26.03; $P_{(zero)}$ = 2.56e-8, F= 34.3). Such DS differences were observed even before stimulus onset. Meanwhile, LIP neurons exhibited similar levels of mean activity between different saccade directions (CT vs. IT) before monkeys' saccade choice (**Figure S1E**). These results indicated a nonlinear modulation of saccade choice on motion DS in LIP, aligned precisely with the response property of each neuron. Furthermore, a demixed principal component analysis on the pseudo-population activity (STAR Methods) revealed that the saccade direction related representation was a substantial component of LIP activity, as the saccade direction and motion–saccade interaction together explained a similar amount of variance in the population activity as the stimulus motion direction did (**Figure 2L** and **Figure S2**). In contrast, we did not observe significant DS differences between the CT and IT conditions in the data sessions in which the saccade targets were aligned close to the vertical direction (21 of 104 neurons, **Figure S3**).

Feedback modulation specifically impacted the decision-correlated activity in LIP
We then examined the impact of nonlinear feedback modulation on the correlation between LIP DS and monkeys' choice behavior. We found that LIP DS was more decision-correlated in the CT condition than in the IT condition. Illustrated in **Figure 2K**, the LIP DS on zero-coherence trials, which reflected monkeys' trial-by-trial categorical choice, was significantly greater in the CT condition than in the IT condition (nested ANOVA: P = 2.56e-8, F= 34.3). We also quantified the correlation between LIP neural activity and the trial-by-trial categorical choice or the physical motion direction by comparing LIP neural activity on correct versus incorrect trials. We used low-coherence trials, as monkeys made enough errors in these trials. In the CT condition but not the IT condition, LIP DS at the population level was significantly reversed in sign on incorrect trials as compared to correct trials (**Figure 3A** and **3B**; nested ANOVA: $P_{(CT)}$ = 0.0045, F = 8.32). Accordingly, in the CT condition but not in the IT condition, LIP activity correlated more closely with the monkeys' trial-by-trial abstract decisions about motion direction, as opposed to the physical motion direction (**Figure 3C** and **3D**; nested ANOVA, $P_{(CT)}$ = 0.0016, F = 10.31; $P_{(IT)}$ = 0.443, F = 0.59). Meanwhile, LIP activity correlated more closely with the trial-by-trial abstract decisions during the CT condition relative to the IT condition on the low coherence trials (nested ANOVA: P = 1.95e-8, F = 34.94). Furthermore, we used partial correlation analysis to examine decision- and stimulus-related components of DS (i.e., r-decision and r-stimulus, **Figure 3E** and **3F**) using all four coherence levels. The decision-related component of LIP DS was significantly greater in the CT condition than in the IT condition (**Figure 3E**; nested ANOVA: P = 1.07e-6, F= 25.72), and this difference emerged even before motion stimulus onset. This suggests that the LIP DS was more closely correlated with monkeys' decisions in the CT condition than in the IT condition. However, the stimulus-related component of the LIP DS was similar at the population level between the two conditions (**Figure 3F**; nested ANOVA: P = 0.7606, F = 0.09), suggesting that the modulation of LIP DS by the later saccade choice was primarily related to a decision process rather than basic sensory processing.

We further examined the impact of nonlinear feedback modulation on the correlation between LIP DS and the monkeys' RTs. Shortly after motion stimulus onset, LIP neurons showed greater population response to their preferred motion direction on shorter versus longer RT trials for all motion coherence levels in the CT condition (**Figure S4**) but not in the IT condition (**Figure S5**). On zero-coherence trials (**Figure 3G** and **3H**), DS evolved more rapidly on shorter vs. longer RT trials in the CT condition but not in the IT condition (bootstrap: $P_{(CT)}$ < 0.01, $P_{(IT)}$ = 0.87).

194   Together, these results indicated that the feedback modulation of saccade choice on sensory
195   evaluation predominantly impacted the decision-correlated activity in LIP.
196
Trained multi-module RNNs replicated the nonlinear feedback modulation
Our electrophysiology data show that action selection nonlinearly modulates the neural processing
of sensory evaluation, which indicates a complex, iterative computation for flexible decision-
making. Training RNNs on behavioral tasks used in experimental neurophysiological studies has
proven to be helpful for exploring putative circuit computations underlying cognitive tasks[28,36,37].
Therefore, we trained multi-module RNNs to perform the FVMD task to examine the circuits and
computation mechanisms underlying the interplay of different decision processes. We adopted
simple neurobiological principles to constrain the connection structure but not the functional roles
to generate recurrently connected modules (STAR Methods). Because our neurophysiological
results showed that the modulation effect of action selection on sensory evaluation depended on
the saccade direction relative to the recorded brain hemisphere, we implemented RNNs composed
of two main modules organized in parallel to simulate two brain hemispheres. Each main module
consisted of two nominal modules, with each nominal module receiving the visual motion input
(motion module) or the target color input (target module), corresponding to the two neuron
populations whose RFs covered the motion stimulus or saccade targets, respectively (**Figure 4A**).
212
We independently trained 50 such networks with randomly initialized weights and identical
hyperparameters to perform the FVMD task. The FVMD task setup used for training RNNs was
tailored to match the monkey experiments. Specifically, the RNNs were trained using motion stimuli
with two high-coherence levels and were tested using stimuli with another four different coherence
levels (high, medium, low, and zero). After training, all 50 networks converged to perform the FVMD
task with high accuracies (>99%) on the training coherence and exhibited coherence-dependent
performances when tested with untrained stimuli (**Figure 4B** and **4C**): accuracy increased and RT
decreased with motion coherence.
221
We then analyzed the unit activity in different modules when the trained RNNs were tested
with untrained motion stimuli. Across all networks, the units in the motion module and target
modules exhibited activity corresponding to sensory evaluation and action selection, respectively.
Specifically, units in the motion modules showed coherence-correlated motion DS (**Figure 4D** and
4F and **Figure S6** and **S7**; mean r = 0.29; one-sample t-test: P = 4.09e-42, t(49) = 46.31), and
neuronal activity on zero-coherence motion trials reflected the trial-by-trial abstract decisions of the
networks (t-test: $P_{(example RNN)}$ = 5.97e-27, t(91) = 15.33; $P_{(population)}$ = 1.05e-31, t(49) = 27.86). Such
motion DS resembled our electrophysiology data when the motion stimuli were presented within
the RFs of the LIP neurons. Meanwhile, units in the target modules showed coherence-correlated
saccade DS (**Figure 4E** and **4G** and **Figure S6** and **S7**), and the neuronal activity on zero-
coherence motion trials also showed significant choice probability (t-test: $P_{(example RNN)}$ = 7.06e-8,
t(68) =15.83; $P_{(population)}$ = 2.08e-22, t(49) = 17.21). Such saccade DS was consistent with the
commonly observed decision-related activity in previous studies in which the saccade target was
presented within the RFs of LIP neurons[9-11].
236
Furthermore, we found that the motion DS of units in the motion modules was significantly
greater in the CT condition than in the IT condition for the majority of RNNs (**Figure 4H** and **4I** and
Figure S8A and **S8B**): in the CT condition relative to the IT condition, activity was significantly
greater for the preferred motion direction but weaker for the nonpreferred direction (**Figure S9**).
ROC analysis confirmed that the motion DS in the motion module was significantly greater for all
four coherence levels in the CT vs. IT conditions in most RNNs (**Figure 4J, 4K** and **4N** and **Figure
S8C** and **S8D**; two-way ANOVA: $P_{(example RNN)}$ = 1.53e-11, F = 47.49; $P_{(population)}$ = 3.50e-77, F =
660.84). Similar to the monkey electrophysiology data, the decision-correlated motion DS (r-
decision) in the motion module was dramatically reduced in the IT versus CT condition (paired t-
test: P = 6.84e-17, t(49)= -12.53), and this difference was substantially greater than the difference
in the stimulus-related (r-stimulus) motion DS between the two conditions (paired t-test: P = 2.02e-

14, t(49) = -10.70; **Figure 4L, 4M** and **4O** and **Figure S8E** and **S8F**). These results indicated that
action selection also nonlinearly modulated the neural processing of sensory evaluation during
decision-making in the RNNs.
251
Selectivity-specific feedback connections mediate the modulation of sensory evaluation by
action selection
Our multi-module RNNs showed a remarkable similarity to both the behavioral performances and
the neural activity patterns in the monkey electrophysiology data, suggesting that these networks
may have adopted mechanisms similar to those in the monkey brain to mediate the decision
processes in the FVMD task. Therefore, we explored the potential circuit mechanisms underlying
the modulation of action selection on sensory evaluation in RNNs. We hypothesized that this
modulation might result from the feedback connections from the target module to the motion
module. To test this hypothesis, we examined the recurrent weights of the cross-module
connections (STAR Methods). Both the feedback and feedforward cross-module connection
weights were significantly greater between the selectivity-matched unit pairs (e.g., units in the
motion module preferred to encode 315° and units in the target module preferred to encode red)
than the selectivity-nonmatched unit pairs (e.g., units in the motion module preferred to encode
315° while units in the target module preferred to encode green) (**Figure 5A**; paired t-test:
$P_{(feedforward)}$ = 7.18e-16, t(49) = -11.76; $P_{(feedback)}$ = 5.26e-18, t(49) = -13.40). In particular, the average
feedback connectivity was primarily excitatory between the selectivity-matched unit pairs but was
inhibitory on average between the selectivity-nonmatched unit pairs. Furthermore, most of the
networks exhibited positive correlations between the connection weight and the match extent of
the neural selectivity of each cross-module unit pair (**Figure 5B**; mean r = 0.11; one-sample t-test:
P = 6.39e-16; t(49) = 11.79; STAR Methods), indicating that units that exhibited stronger encoding
preference to one motion direction in the motion module were more likely to receive more extensive
feedback projections from the units that exhibited stronger encoding on the matched target color in
the target module. These results suggested precise feedback connections between RNN modules
that were aligned with the functional properties of different units.
276
To test the causal contribution of the selectivity-specific feedback connections to the
modulation of action selection on sensory evaluation, we conducted three projection-specific
inactivation experiments in the RNNs (STAR Methods). First, we performed nonspecific ablation of
all the feedback projections from the target module to the motion module when testing with
untrained motion stimuli. In most of the tested RNNs, this caused a dramatic reduction in their
performance in all nonzero-coherence stimulus conditions (**Figure 5C**; two-way ANOVA: P = 4.74e-
80, F = 704.65) as well as significantly prolonged RTs for high- and medium-coherence stimulus
conditions (**Figure 5D**; two-way ANOVA: P = 2.45e-4, F = 13.79). The mean RTs became more
dispersed across all the trained RNNs and did not differ among different motion coherences after
ablation (one-way ANOVA: P = 0.68, F = 0.50). Ablating feedback connections significantly affected
the motion DS in the motion module. On average, although the averaged motion DS did not
decrease after ablation (**Figure S10A** and **S10C**), the difference in the motion DS between the CT
and IT conditions vanished (**Figure 5E-5G** and **Figure S11A** and **S11B**; two-way ANOVA: $P_{(CT\ vs.}$
$_{IT)}$ = 0.14, F = 2.21). These results indicated a dramatic reduction in the nonlinear feedback
modulation on sensory evaluation by action selection in the RNNs. To further test the importance
of different patterns of feedback connectivity for the RNNs to solve the FVDM task, we next
performed pattern-specific ablation of the feedback connections in RNNs. We separated the
feedback projections in each RNN into specific (i.e., the feedback connection weights were positive
or negative between the selectivity-matched or selectivity-nonmatched unit pairs, respectively) and
nonspecific groups. Despite similar numbers and weights of the feedback connections in these two
groups (**Figure S12A** and **S12B**), ablating the feedback connection in the specific group versus
the nonspecific group resulted in much more severe impairments in the RNNs' behavior
performance (**Figure S12C** and **S12D**). These results indicated that the feedback connections that

reflected the learned stimulus–response association was crucial for the RNNs to solve the FVDM
task.
302
Second, we disrupted the selectivity-specific feedback connectivity without changing the
total strength of the feedback connections by randomly rearranging the feedback connection
weights. This also resulted in a significant reduction in the RNNs' performance accuracy for
nonzero-coherence motion stimuli (**Figure 5C**; two-way ANOVA: P = 1.38e-90, F = 883.61) and
prolonged RTs for high- and medium-coherence motion stimuli (**Figure 5D**; two-way ANOVA: P =
1.94e-12, F = 54.06). The changes of RNN units' activity patterns after scrambling feedback
connections were similar to the effects after feedback connectivity was fully ablated (**Figure S10B**,
S10C, S13A, S13B, S13D and **S13E**). Particularly, the levels of nonlinear modulation of sensory
evaluation by saccade selection dramatically decreased, as evident by the diminished difference
in the motion DS between the CT and IT conditions (**Figure 5H**, two-way ANOVA: P = 5.81e-12, F
= 51.52).
314
Third, we trained another 50 RNNs without feedback connectivity to learn the FVMD task.
Similar to the full-model RNNs, all 50 networks converged to perform the FVMD task with high
accuracies (>99%) after training and exhibited coherence-dependent performances when tested
with untrained stimuli (**Figure S14A** and **S14B**). Meanwhile, units in the motion module and target
modules exhibited coherence-dependent motion DS and saccade DS, respectively (**Figure S14C**
and **S14D**). Consistent with the above connectivity ablation experiment, units in the motion module
of these RNNs did not exhibit different levels of motion DS between CT and IT conditions (**Figure**
S14E-S14G; two-way ANOVA: P = 0.63, F = 0.24), indicating no noticeable nonlinear feedback
modulation on sensory evaluation by action selection. Importantly, the performance accuracies of
these RNNs decreased significantly when tested with untrained nonzero-coherence stimuli (**Figure**
S14H; two-way ANOVA: P = 1.46e-14, F = 65.64), as compared to the full-model RNNs.
Furthermore, we trained an additional 50 RNNs without feedback connections to learn the FVMD
task. These RNNs were initialized with greater recurrent connection probabilities than the full-model
RNNs, such that the number of total trainable connection weights matched that in the full-model
RNNs. Interestingly, these RNNs still exhibited significantly lower performance accuracies when
tested with high- and medium-coherence stimuli relative to the full-model RNNs (**Figure S15**; two-
way ANOVA: P = 8.24e-6, F = 20.60). Together, these results suggested that the nonlinear
feedback modulation, which was mediated by the selectivity-specific feedback connections, were
important for the RNNs to efficiently solve flexible decision tasks.
334
Nonlinear feedback modulation enhanced the decision consistency by strengthening the
attractor basins of network dynamics
Our RNN modeling indicates a crucial role for nonlinear feedback modulation in optimizing flexible
decision-making. Subsequently, we delved into the computational mechanisms embodied by
nonlinear feedback modulation during the decision-making process. Specifically, we examined the
unit activity in the target module, a direct contributor to the decision outputs of RNNs. Strikingly,
the projection-specific inactivation experiments led to a significant reduction in the magnitude of
saccade DS in the nonzero-coherence trials (**Figure 6A-6C**, $P_{(ablating\ feedback)}$ = 6.90e-5, F = 16.29;
$P_{(shuffling\ feedback)}$ = 1.2e-19, F = 95.08; two-way ANOVA). Additionally, saccade DS emerged earlier
than the motion stimulus onset in the nonzero-coherence trials, and its positive correlation with the
motion coherence levels diminished after the projection-specific inactivation (**Figure 6A-C** and
Figure S11C, S13C and **S13F**; mean $r_{(ablating\ feedback)}$ = -0.034; mean $r_{(shuffling\ feedback)}$ = −0.030). These
disrupted patterns of saccade DS observed in the target module following projection-specific
inactivation aligned with the decreased decision consistency of RNNs, indicating a diminished
reliance on sensory input and an increased dependence on internal noise in the decision-making
process.
351
Previous studies have demonstrated that attractor dynamics in networks can explain
choice consistency, with steeper landscapes around attractor basins reflecting consistent

decisions[38-42]. Consequently, we investigated attractor dynamics in the state space of network
activity that underlies decision-making variability in RNNs with different types of feedback
connectivity. Similar to prior studies[40], we examined the neural dynamics underlying decisions in a
1-D neural subspace using unit activity in the target module responsible for saccade choices (see
methods). Beginning with the reconstruction of a numerical approximation of the energy landscape
in this 1-D subspace, we explored changes in this landscape after different types of projection-
specific inactivation. This revealed a dramatic reduction in the depth of attractor basins in the
energy landscape of population activity after ablating all feedback connectivity (**Fig. 6D-E,** paired
t-test: P = 2.28e-11, t(49) = -8.61). Moreover, ablating the feedback connections led to a more
severe reduction in the depth of attractor basins in the network dynamics within the specific group
compared to the nonspecific groups (**Fig. 6F-G**, paired t-test: P = 1.59e-6 , t(49) =-5.46).
365
We further examined how the energy landscape in the 1-D subspace changed in relation
to task difficulty (motion coherence). Consistent with prior findings[40], trials with lower decision
consistency (trials using lower motion coherence) exhibited shallower attractor basins at the time
of decision for all types of RNNs (**Fig. 6H-K**). However, both the depth and the positional separation
of attractor basins in the network dynamics significantly decreased for all non-zero motion
coherence levels after the ablation of all feedback connections ($P_{(depth)}$ = 5.20e-25, F = 122.80;
$P_{(position)}$ = 1.82e-27, F = 137.75; two-way ANOVA). Notably, this reduction in basin depth and
separation was more pronounced in the specific group compared to the nonspecific groups after
ablating the feedback connections (**Fig. 6J-K**, $P_{(depth)}$ = 2.65e-13, F =57.35; $P_{(position)}$ = 3.73e-14, F
= 61.79; two-way ANOVA). These results might underlie the computational mechanisms that
explain the observed reduction in the decision consistency of RNNs following projection-specific
inactivation: the shallower and closer attractor basins after ablating feedback connections resulted
in less consistent decisions. This happened because the variability in neural activity made it more
likely for population activity to stochastically shift out of the shallower basins and into nearby
alternative ones.
381
382
Discussion
Here we show that primate LIP activity related to evaluating the visual motion stimulus was
nonlinearly modulated by an impending saccade choice that was used to report decisions during a
FVMD task, even though the saccades were toward non-RF locations. This suggests that the
sensory evaluation and action selection may proceed iteratively during decision-making. This view
is also consistent with the common observation that action selection related activity arises at the
early stage of evidence accumulation during decision-making[3,9,34], as well as our observation that
the modulation of action selection on sensory evaluation emerged during the early task period even
before motion stimulus onset in the FVMD task. Brain areas that are related to different decision
processes, although spatially separated[14,21,33], likely form a real-time associated network to solve
current decision tasks. The instantaneous result of sensory evaluation may be transmitted to the
brain areas responsible for action selection during decision-making in a feedforward manner, and
the action selection process may also exert a real-time modulation on the neural processing of
sensory evaluation in a feedback manner.
397
The feedback modulation we observed during decision-making was distinct from the
modulation of feature-based attention in early sensory cortex. Although both modulation effects
match the stimulus tuning of recipient cells[43-47], only the feedback modulation observed in the
current study reflects the neural processing of decision-making. This is because only the decision-
related component, not the stimulus-related component, of LIP activity was modulated by the
monkeys' later saccade choice during the FVMD task (**Figure 3**). This suggests that the feedback
modulation from action selection to LIP activity was primarily related to the decision process rather
than low-level sensory processing.
406

The selectivity-specific feedback modulation from action selection to sensory evaluation
emerged in our multi-module RNNs during training, although the cross-module connections were
randomly initialized before training. Importantly, our causal in-silico experiments demonstrated that
these precise feedback connections played an essential role in mediating the modulation of action
selection on sensory evaluation while solving the FVDM task. Previous studies showed that both
the corticocortical feedback from the secondary visual cortex to the primary visual cortex and the
corticogeniculate feedback in primates are organized into parallel streams resembling the
reciprocal feedforward pathways[48-50], suggesting a potential functionally specific feedback
connection in visual processing. However, to the best of our knowledge, a similar segregation of
feedback that reflects the encoding properties of the target neurons has not been evident in
decision network in vivo. Therefore, it will be important for future studies to examine the connectivity
and correlation of neural activity among neural groups related to different decision processes
through sophisticated anatomical experimental approaches as well as multi-channel recordings in
vivo.
421
Our brain includes extensive feedback connections across different brain areas, which, in
some cases, even outnumber the feedforward connections[50,51]. However, compared to the
feedforward connections, much less is known about the function of feedback connections.
Feedback modulation has been implicated in top-down modulation of neuronal responses in the
early sensory cortex, such as attention and expectation, which facilitate the processing of important
stimuli and suppress distractors[52-54]. Furthermore, both experimental and computational studies
have shown that weak decision-correlated neural activity (i.e., choice probability) as well as the
correlated firing of pairs of neurons (i.e., noise correlation) in the early sensory cortex might partially
result from feedback input in decision tasks[55-57], suggesting a potential role for feedback connection
in modulating early sensory processing during decision-making. Consistently, the nonlinear
feedback modulation, mediated primarily by selectivity-specific feedback connections, were
important for our multi-module RNNs to efficiently solve the FVMD task, as the RNNs' decision
behavior became more stochastic following ablating/disrupting the feedback connectivity. This was
evident not only in the diminished behavioral performance of the RNNs but also in the disrupted
patterns of activity related to saccade choice in the target module. Notably, the nonlinear feedback
modulation intensified the attractor basins of the population activity associated with saccade choice,
led to more reliable decisions based on sensory input. Our results unveil a novel pattern of feedback
modulation during the neural processing of decision-making, and suggest a potentially critical role
that feedback modulation plays in increasing the consistency of flexible sensorimotor decisions.
441
Two-state attractor models have been widely employed to elucidate two-alternative forced
decision-making processes[38-42,58,59]. In these models, decisions arise when network activity falls
into one of two attractor basins. Previous studies have posited that decision consistency is
influenced by the configuration of the energy landscape surrounding these attractor basins[40,42,59].
Essentially, steep-sided and deep-basin energy landscapes contribute to consistent decision-
making, as it becomes challenging for internal noise to shift activity between basins. Conversely,
when energy landscapes are relatively flat and attractor basins are shallow, decisions are
generated more stochastically, as it is easier for internal noise to drive activity out of the attractor
basins and into alternative ones. Consistently, our RNNs modelling showed that shallower and less
separated attractor basins were associated with decreased decision consistency in more difficult
trials. Building upon previous research, our results provide in-silico evidence supporting the notion
that the energy landscape within the state space of population activity causally influences decision
consistency. Moreover, our findings present the initial circuit mechanism through which the
configuration of the energy landscape for decision-related activity is modulated by feedback
modulation, to our knowledge. Future studies are needed to explore the connections between
different types of circuit connectivity and population activity dynamics in cognitive processing in
vivo.
459

Our results are consistent with predictive coding theories of feedback function, which
propose that an internal model of the world is generated in the brain based on sensory data and
prior experience and is refined by incoming sensory data[60-63]. In terms of the architecture of
hierarchical predictive coding schemes, different neuronal ensembles encode different
attributes/choices in either the action selection or sensory evaluation process at each level of the
cortical hierarchy during flexible decision-making. Meanwhile, the conditionally independent
expectations are functionally segregated, so that descending predictions from the action selection
process are limited to the sensory evaluation process on the matched\associated stimulus attribute
but not the opposite attribute. The selectivity-specific feedback connectivity in our RNNs fits the
functional segregation of expectation very well during the interplay between distinct neural
processes of decision-making. Specifically, the neural ensembles related to either sensory
evaluation or action selection processes for the same choice might form precise recurrently
connected loops. The prediction signal carried by the precise feedback connections might facilitate
sensory evaluation of the associated stimulus but prohibit evaluation of the nonmatched stimulus.
This could, in turn, amplify the task-relevant sensory input, facilitate the sensorimotor
transformation, and ultimately result in faster and more accurate action choice in flexible decisions.
In future work, it will be important to use techniques such as projection-specific inactivation and
microstimulation in vivo to test the causal contribution of feedback connectivity in decision-making
and other cognitive functions.
479
480
481   **Materials and Methods**
482
Behavioral task, stimulus display, and animal preparation
The flexible visual motion discrimination (FVMD) task (**Figure 1A**) has been reported previously[14]
and is briefly summarized below. In this task, monkeys were required to saccade to either the green
or red targets based on the direction of the visual motion stimulus. Two motion directions (135°,
315°) were used, each with four different coherence levels (0%, 9%, 18%, 36% for monkey M; and
0%, 13%, 25%, 50% for monkey B) were tested. If the sample direction was 135°, the monkeys
must saccade to the green target to receive a juice reward, whereas the 315° direction was
associated with the red target. The rewarded target (red or green) was randomly chosen (with 50%
probability) on each zero-coherence trial. The positions of red and green targets were randomly
chosen between the two positions on each trial at each recording session. Therefore, there is no
fixed mapping between motion stimulus and saccade direction.
494
To initiate each trial, monkeys must hold a touch-bar and acquire gaze fixation. They then
needed to maintain fixation within a 2.0-2.5° window throughout the trial before their saccade
choice. After a 500ms fixation period, two colored saccade targets appeared simultaneously at
opposite positions relative to the fixation point with equal eccentricities (8° and 9° for Monkey M
and B, respectively). 400ms later, a sample motion stimulus was presented at a location orthogonal
to the axis of, but at the same eccentricity as, the saccade targets. We used motion stimuli that
were full contrast, 8° diameter, random-dot movies composed of 190 dots per frame, and the dots
moved at 10°/s. Monkeys needed to saccade to either red or green targets within a 60-2000ms
window after sample stimulus onset. A juice reward would be delivered to the monkeys if they made
correct saccade choice.
505
Two male monkeys (Macaca mulatta, 15~16 years old, 8–14 kg) were trained on the FVMD
task and implanted with a head post as well as a recording chamber positioned over PPC. Our
surgical, behavioral, and neurophysiological approach has been described in detail in a previous
study[14] . Stereotaxic coordinates for chamber placement were determined from magnetic
resonance imaging (MRI) scans obtained before chamber implantation. LIP chambers were
centered on the intraparietal sulcus (IPS), 4.0 mm posterior to the intra-aural line and 1.0 mm lateral
from the midline for monkey M, and 0 mm anterior to the intra-aural line and 15.0 mm lateral from
the midline for monkey B. Monkeys were housed in individual cages under a 12-hour light/dark

cycle. Behavioral training and experimental recordings were conducted during the light portion of
the cycle. Monkeys sat comfortably while head-fixed in a custom-made primate chair inside a dark
experiment rig. The task stimuli were displayed on a 21-inch color CRT monitor (1280*1024
resolution, 75 Hz refresh rate, 57 cm viewing distance). Both monkeys were tested with identical
stimuli and timing. A solenoid-operated reward system was used to deliver juice reward to the
monkeys. Monkeys' eye positions were monitored by an optical eye tracker (SR Research) at a
sampling rate of 1 kHz and stored for offline analysis. Stimulus presentation, task events, rewards,
and behavioral data acquisition were accomplished using an Intel-based PC equipped with
MonkeyLogic software running in MATLAB (http://www.monkeylogic.net). All experimental and
surgical procedures were in accordance with the University of Chicago Animal Care and Use
Committee and National Institutes of Health guidelines.
525
Electrophysiological recording
We used either 75-μm tungsten microelectrodes (FHC, ~1 MΩ) or 16-channel V-Probes (Plexon)
to record single neuron activity in LIP. Neurophysiological signals were amplified, digitized, and
stored for offline spike sorting (Plexon) to verify the quality and stability of neuronal isolations. We
recorded neuronal activity in a memory-guided saccade task to map LIP RF locations before each
FVMD recording session.
532
Our LIP recordings targeted different hemispheres in the two monkeys (monkey M: left
hemisphere; monkey Q: right hemisphere). Therefore, the contralateral target (CT) condition and
ipsilateral target (IT) conditions in **Figure 1F** and **1G** referred to opposite saccade directions (target
locations) between the two monkeys. For monkey M, the CT condition corresponded to the trials in
which the correct saccade target was on the right visual field, whereas the CT condition for monkey
Q referred to the trials in which the correct saccade was toward the left visual field. We localized
LIP in each monkey according to the patterns of neuronal activity in the MGS task (i.e., spatial
selectivity during stimulus presentation and delay). All neurons included in the dataset were
recorded from the same locations (the same grid holes and similar depths: ~5-10 mm from the
cortical surface) where we encountered spatial selectivity in the MGS task. LIP neurons were also
identified based on anatomical criteria, such as the location of each electrode track relative to that
expected from the MRI scans, the pattern of gray–white matter transitions encountered on each
electrode penetration, and the relative depths of each neuron.
546
We aimed to present the visual motion stimulus inside the RFs of the identified neurons
during each recording session. For single-channel electrode recording, only neurons exhibited
visual responses to the motion stimuli during prescreening (tested with the FVMD task) were
recorded with sufficient trials (~300-600) of the FVDM task. For neurons exhibiting clear spatial
RFs during the MGS task, we presented the motion stimulus inside LIP neurons' RFs; while for
those neurons which did not show a clear RF during the MGS task, we presented motion stimuli in
the positions (always in the visual field contralateral to the recorded hemisphere) in which neurons
exhibited the strongest response to the motion stimuli. For the multi-channel recordings, we
recorded all neurons isolated across all channels and presented the motion stimulus inside one of
the isolated neurons' RFs. Because adjacent recording sites were located 100μm apart, nearby
neurons typically showed similar RF locations.
558
559 ***Data analysis***
560
Neuronal pre-screening
While part of the neural data was presented in a previous report[14], the current analysis focuses on
a different phenomenon which was not examined previously. We included all neurons recorded
from single-channel electrodes for the analysis. For multi-channel recordings, we only included the
neurons which showed significant modulation (different from fixation period activity, one-way
ANOVA, $p < 0.01$) of their averaged activity across all motion stimuli because the stimulus could
not always be presented within the RF of all the recorded neurons. The low-firing neurons whose

maximum firing rates were less than 2.0 spikes/s (to the direction producing greater average
responses) during stimulus presentation were also excluded for further analysis. We then applied
a one-way ANOVA test to compare activity between the two different motion directions during the
period following motion stimulus onset (50-250 ms after motion stimulus onset) to select neurons
that showed significant motion direction selectivity (DS) during the decision period. Only neurons
that showed significant ($p < 0.01$) DS were used for further analysis.
574
Receiver operating characteristic (ROC) analysis
To quantify each neuron's DS in the FVMD task, we applied an ROC analysis to the distribution of
firing rates within sliding windows (100 ms width, 5 ms steps). The ROC value, which ranges
between 0.0 and 1.0, indicates the performance of an ideal observer in assigning motion direction
based on each neuron's trial-by-trial firing rates. Values of 1.0 and 0.0 correspond to perfect
classification (i.e., strong DS), while a value of 0.5 indicates chance-level classification performance
(i.e., no DS). For trials with zero coherence motion, we assigned direction labels on each trial based
on the monkey's choice.
583
In **Figure 3**, in order to test whether DS reflected monkeys' trial-by-trial choices, we used
an ROC analysis to quantify whether LIP activity correlated with monkeys' trial-by-trial categorical
choices more than the physical direction of motion stimulus by analyzing both correct and error
trials. We only included low coherence trials in this analysis because there were sufficient numbers
of error trials (average performance: Monkey M: 74% correct, Monkey B: 67% correct). LIP
neuronal activity was analyzed by ROC according to either the monkey's trial-by-trial categorical
choices or the direction of the sample stimulus on each trial. Furthermore, for this analysis, we only
used neurons for which we recorded sufficient trials (for both correct and error trials, N>4) for the
low coherence condition of each motion direction (n = 45).
593
Partial correlation analysis
We performed a partial correlation analysis to quantify the correlation between LIP neural activity
and the monkeys' trial-by-trial categorical choices or the physical motion direction of the stimulus
using all trials. For each trial, we obtained three parameters for the calculation: the stimulus
direction, the pre-choice neuronal activity, and the monkeys' choice. We assigned the stimulus
directions with different values for different directions and coherence levels: positive and negative
values were used for preferred and nonpreferred directions, respectively; while 4, 2, 1 and 0 are
used for code the high, medium, low, and zero coherence levels. We also assigned different values
to different choice directions (−2 for choosing the preferred direction, and + 2 for choosing the
nonpreferred direction). Two measures were then calculated: r stimulus = r (neuronal activity,
stimulus direction| choice direction), the partial correlation between neuronal activity and stimulus
direction given the monkeys' choices; and r choice = r (neuronal activity, choice direction | stimulus
direction), the partial correlation between neuronal activity and the monkeys' categorical choice
given the stimulus direction. In **Figure 3**, we used the mean activity within a sliding window (100ms
width, sliding with 5ms steps) for each neuron to perform the partial correlation analysis.
609
dPCA analysis
We conducted demixed principal component analysis using the methodology and the code from a
previous study[64] (http://github.com/machenslab/dPCA) to reduce the dimensionality of the
population activity as the standard PCA and demixes all task variables. Specifically, we tested how
much each task variable (motion direction of sample stimuli, saccade direction, motion-saccade
interaction, timing) contributes to the LIP population activity during the DMC task.
616
617   As demonstrated in the previous study, the dPCA finds separate decoder (F) and encoder
618 (D) matrices for each task variable (∅) by minimizing the loss function:

$$L_{dPCA} = \sum_0 \left\| X_0 - F_0 D_0 X \right\|^2$$

620 where X is a linear decomposition of the data matrix, which contains the instantaneous firing rate
621 of the recorded neurons, into variable-specific averages:

$$X = \sum_0 X_0 + X_{noise}$$

622
623 Here, we decomposed the neural activities into four parts: condition-independent, stimulus-
624 dependent (two motion directions with four coherence levels), saccade-dependent (two saccade
625 directions), dependent on the stimulus-saccade interaction, and noise. The decoder and encoder
626 axes permit us to reduce the data into a few components capturing the majority of the variance of
627 the data dependent on each task variable.

628
629 ***Recurrent neural network (RNN) training***

630
Network Implementation
We trained biologically inspired networks to perform the FVMD task using methodology similar to
previous studies[28,36]. We implemented multiple modules in the networks through constraints on the
input/output structure and the initial recurrent connectivity of the hidden layer. Specifically, we built
excitatory-inhibitory networks with 200 hidden units divided into two equal-size main modules to
simulate two brain hemispheres. Each main module was further divided into two nominal modules,
with each one only receiving the visual motion input (motion module) or target color input (target
module), respectively. This design was intended to simulate the two neuron populations whose
RFs covered the motion stimulus or saccade target in the monkey electrophysiology experiment
(**Figure 4A**). Every nominal module was allocated one quarter of the excitatory units (40) and one
quarter of the inhibitory units (10) in the overall network to ensure that the modules did not differ in
their balance of excitation/inhibition prior to training. Meanwhile, in order to simulate the local and
long-range connection structures in the brain, we set different levels of recurrent connectivity within
and between different modules: the local (connectivity within each nominal module) recurrent
connection density (probability) was the highest (50%); the across-RF (connectivity between the
two nominal modules within the same main module) connection density was set to be in the medium
level (25%); the cross-hemisphere (connectivity between either the motion modules or the target
modules across different main modules) connection density was the lowest (10%). Only excitatory
neurons could have "cross-hemisphere" projections to the corresponding nominal module (e.g.,
from the motion module of the "left hemisphere" to the motion module of the "right hemisphere").
Only the connection weight within the hidden layer was updated during training following methods
described previously[28]. In addition, the connection weights between network units are endowed
with short-term synaptic plasticity, which is aimed to provide connection weights with activity-
dependent fluctuation over short timescales within each trial.

655
656 The input to the network consisted of two parts: 9 motion input units and 8 target color input
657 units. The preferred directions of motion input units were evenly spaced across 360°, with response
658 tuning distributed according to a von Mises function. The value of the nth motion input unit was set
659 to

$$r_n = \frac{2}{3}\left(ace^{\left(\kappa\cos(\theta - \theta_{pref})\right)} + b\mathcal{N}(0,2) + \sqrt{\frac{2}{\alpha}}\mathcal{N}(0, \sigma_{in}) \right) + d$$

661 , where $\alpha = \frac{\Delta t}{\tau_{mem}}$, θ is the direction of motion stimulus in radian, θ_{pref} is the preferred direction of
662 this motion input unit, $a = \frac{4}{e^\kappa}$ after stimulus onset but 0 elsewhere, $c \in [0, 1]$ is the coherence of the
663 dot motion stimulus, b is a binary value that equals 0 before stimulus onset and 1 after that, and d
664 is a constant visual input signal with an amplitude of 2 that decays with time. Specifically, for zero-
665 coherence input, since $ace^{\left(\kappa\cos(\theta - \theta_{pref})\right)}$ would be zero, we set this term as $0.4max(c)$.
666

667   The target input units were initially evenly divided into two groups projected to the two
668 target modules of hidden units. Each group was further divided into red and green subgroups. The
669 color subgroup was set as active with an amplitude of $\frac{4}{3}$ to represent the color within the projected
670 target module, while the other color group remained silent (amplitude =0). An exponential decay
671 filter was also used to fit the activity of sensory neurons in the early sensory cortex across time:

672
$$\mathrm{f}(x) = \left(\frac{1}{3}\right)^{t} \mathrm{x}$$

673 , where x is the constant visual input for motion input and the color signal for color input, t is time
674 in miliseconds.
675
The two output units of the network simulate two different saccade directions. In order to
simulate the oculomotor control in the brain, each brain hemisphere (main module) projects much
more densely to the contralateral response units (probability = 0.32) than the ipsilateral response
units (probability = 0.08). To generate a probability distribution over output values at each time
point, we also applied a softmax function to scale the response unit activities in the output layer.
To reduce the stochasticity of the network activity brought by input and output connections, we
randomly sampled input and output weights from a normal distribution ($\mathcal{N}(0.2, 0.05)$). In addition,
the input weights were re-initialized if there were extreme values (the minimum value was smaller
than ¾ of the maximum value) after multiplying the stimulus signal and weight of motion or target
input. Similarly, the output weights were re-initialized if there were extreme values in the output
weight values. The input units had excitatory and random projections to the recurrent units with a
probability of 0.32. However, only the excitatory units in the target modules of each "hemisphere"
could project to the output units. Both the input and output weights were fixed for each network
during task training.
690
Network training
The networks were trained using BrainPy[65] on an Intel(R) Core(TM) i9-9900K CPU (3.60GHz, 8
cores). The network was optimized using backpropagation through time (BPTT) and stochastic
gradient descent with an Adam optimizer (default setting, first moment estimates decay rate=0.9,
second moment estimates decay rate =0.999) to minimize a loss function. The network parameters
(recurrent weights/biases) were optimized to minimize a loss function with three parts as in a
previous study[28]: (a) a performance loss; (b) a metabolic cost on mean neuronal activity; and (c) a
metabolic cost on connectivity. The gradient was clipped to a maximum L2 value of 0.1 to avoid
the exploding gradient problem. During training, only the hidden unit related parameters (weights,
bias, and initial activities) were updated. The training process was terminated when the network's
performance accuracy reached 99% or till the maximum number of iterations (2000).
702
During each trial of the training and testing of the FVMD task, networks were first presented
with two color targets through the target input units, and then presented with the motion direction
through the motion input units. Both the motion and target inputs persisted until the end of each
trial. A short time after the motion input (100ms, 5 time steps), the networks were required to report
the direction of the stimulus motion by choosing the saccade target with the appropriate color.
Specifically, each element of the task design in the FVMD task that the models were trained to
perform was tailored to match those used in the monkey experiments: the directions of the motion
stimuli, the target colors, as well as the task (stimulus) durations were the same as that used in
monkey electrophysiology experiments. Trials were programmatically generated by constructing
motion/target inputs to the networks at each timestep and the desired response (left saccade, right
saccade) at each timestep. In total, we trained 50 networks to perform the FVDM task. All networks
were first trained using motion directions with two coherence levels (60% and 90%) and were then
tested using motion stimuli with another four different coherence levels (75%, 55%, 35%, and zero),
which had never been presented during training. Besides the difference in the coherence of motion
input, the color inputs remained the same during the testing period. All the networks achieved

consistently high performance by the end of training (> 99% accuracy). All the important model
hyperparameters were listed in **Table S1**.
720
Quantification of the networks' behavioral performance
To test whether our multi-module RNNs exhibited decision behavior similar to monkey subjects, we
examined their performance accuracies and RTs when tested by novel stimuli with different
coherence levels. The performance accuracy was defined as whether the output of the response
units (starting from 100ms after stimulus onset to the trial end) matched the desired output.
Furthermore, we defined the networks' RTs as the time point from which the differences between
two output units were greater than a threshold (0.8) for three consecutive time points after the
stimulus onset. During some trials, the threshold was never reached until the end of the trial (500ms
after the motion stimulus onset). In these cases, we artificially set the RT to be 600ms in these trials
for further analysis.
731
Analysis of RNN activity
We performed the same analyses on the RNN units as we did on the neurophysiology data. In
Figure 4-5, data from both the example network and averaged results across all the networks were
shown. For every network, we only included the units which exhibited at least one kind of task-
related modulation during the test period (i.e., motion direction selectivity or saccade direction
selectivity, one-way ANOVA test, p < 0.01). In order to quantify the activity related to evaluating
motion stimulus, we analyzed the motion DS only including the units in the motion modules,
whereas only the saccade DS of units in the target modules were calculated to quantify the activity
related to saccade selection during the decision-making.
741
Analysis of RNN connectivity
To examine the potential circuit mechanisms underlying the modulation of action selection on
sensory evaluation during decision-making in the RNNs, we examined the cross-module
connection weights in the trained networks. Specifically, we defined the feedforward connection as
the projection from units in the motion module to units in the target module within the same main
module (hemisphere). We also defined the feedback connection as the projection from units in the
target module to units in the motion module within the same main module. Furthermore, we
grouped the cross-module connections into the selectivity-matched and selectivity-mismatched
groups based on the preferences of the neural encoding of units in the motion and target modules.
The selectivity-matched group included two types of unit pairs: 1) units in the motion module
preferred to encode 315° and units in the target module preferred to encode red, 2) units in the
motion module preferred to encode 135° and units in the target module preferred to encode green.
The selectivity-mismatched group includes the other two types of unit pairs: 3) units in the motion
module preferred to encode 315° while units in the target module preferred to encode green, and
4) units in the motion module preferred to encode 135°while units in the target module preferred to
encode red. The values of connection weights projected from excitatory and inhibitory units were
set as positive and negative, respectively.
759
We also examined the correlation between the similarity of neural selectivity and
connection strength for different unit pairs in the RNNs based on the following steps: First, both the
preference and strength of motion direction encoding or target color encoding were quantified for
units in the motion module or target module, respectively. This was done by calculating the
averaged differential activity to the 315° and 135° motion stimuli after motion stimulus onset or
calculating the averaged differential activity to the red and green targets after target onset. In this
case, the unit pairs that exhibited matched selectivity would show selectivity values with the same
sign (positive or negative), whereas the unit pairs that exhibited mismatched selectivity would show
selectivity values with the opposite signs. Second, the units in both the motion module and target
modules were ranked based on selectivity values. Third, the selectivity similarity of each cross-
module unit pair was quantified by calculating the reverse value of the absolute differences between
the target color selectivity rank and the motion direction selectivity rank. The rank of the weight

value of each pair was calculated and then reversed to give the lower weight value a higher rank
score. Finally, we calculated the Pearson correlation between weight rank and selectivity similarity.
A positive correlation indicates that units that exhibited stronger encoding preference to one motion
direction in the motion module were more likely to connect with the units in the target module that
exhibited stronger encoding on the matched target color.
777
Inactivation experiments in silico
Our examination of the network connectivity within the hidden layer of the RNNs revealed that the
selectivity-specific top-down connections originated from the target module to the motion module.
To assess the causal contribution of such precise feedback connections to the modulation of action
selection on sensory evaluation, we performed two in-silico analogues of neuronal inactivation
experiments similar to a previous study[28]. We hypothesized that ablating or disrupting the precise
feedback connection from the target module to the motion module would significantly impact the
activity patterns of the units in the motion module. Specifically, such inactivation on connectivity
was expected to reduce the difference in motion DS in the notion module between CT and IT
conditions. To test this hypothesis, we performed the projection-specific inactivation experiments
in the RNNs in three different ways. First, we ablated all the feedback projections from the target
module to the motion module while keeping all the other network parameters unchanged after the
networks were fully trained. We then tested the networks with the untrained motion stimuli used in
the normal experiment. The connectivity ablation was implemented by directly setting the
connection weights from units in the target module to units in the motion module to zero. Second,
we randomly rearranged the connection weights from the target module to the motion module after
the network was fully trained by shuffling the corresponding weights in the weight matrix. Then, we
tested the networks with the testing motion stimuli. This was aimed to disrupt the selectivity-specific
feedback connection without changing the total strength of the feedback connection. We repeated
this random feedback weight rearrangement process 100 times for each of the 50 trained networks.
Subsequently, we measured the impact on the networks' behavior performance and the activity
patterns in both the motion and saccade modules. In particular, for each network, we tested
whether there were still different levels of motion direction selectivity of units in the motion module
between CT and IT conditions after inactivating the feedback connectivity.
802
We further performed pattern-specific ablation of the feedback connections in the RNNs,
in order to examine the causal roles of different patterns of feedback connectivity in solving the
FVDM task. Specifically, we separated the feedback projections in each RNN into the specific group
and the nonspecific group. The specific group included two conditions: 1) the feedback connection
weight was positive between the selectivity-matched unit pairs, and 2) the feedback connection
weight was negative between the selectivity-nonmatched unit pairs; whereas the nonspecific group
included the rest of the feedback conditions in each RNN. We then tested the effects of the pattern-
specific connectivity inactivation on the RNNs' behavior performance when tested with the
untrained motion stimuli.
812
In the third inactivation experiment, we trained 50 additional RNNs without feedback
connections to learn the FVMD task, in order to test the importance of feedback connectivity in
solving the flexible decision task. These networks were initialized using identical hyperparameters
and trained with the same methodology as the normal RNNs, except that the feedback connections
from the target module to the motion module were ablated before training. We then examined these
RNNs' behavior performance and unit activity when testing with the untrained motion stimuli used
in the normal experiment. Furthermore, we trained another 50 RNNs without feedback connection
to learn the FVMD task. These RNNs were initialized with higher recurrent connection probabilities
than the full-model RNNs to keep the number of the total trainable connection weight comparable
to that in the full-model RNNs. We raised the recurrent connection probability by a factor of 1.176
after removing feedback connections (average number of connections in 50 normal RNNs = 8193,
average number of connections in 50 networks without feedback = 8184, p = 0.53, independent
sample t-test). Except for the recurrent connection probability, all the other hyperparameters and

the training methodology were the same as the normal RNNs. We then compared the behavior
performance of these RNNs with that of the normal RNNS when testing with the untrained motion
stimuli.
829
Analysis of neural landscape
For each network, we adopted a methodology similar to a previous study (cite) to reconstruct the
energy landscape of population activity within the target module. Initially, units in the target module
exhibiting saccade direction selectivity were identified. Subsequently, we employed principal
component analysis (PCA) to reduce the dimensionality of the population activity. Specifically, PCA
was performed on the average activity across these units, considering each stimulus motion
direction, coherence level, and choice. We retained the first five principal components (PCs) for
further analysis, as they explained over 90% of the total variance in population activity. Following
this, the trial-by-trial population activities were transformed into 5-dimensional data using the
aforementioned five PCs.
840
To predict trial-by-trial choice, we employed a linear Support Vector Machine (SVM)
classifier with 10-fold cross-validation. Specifically, we utilized the average activity of the last 100
ms of each trial as input for the classifier. The choice axis was determined by selecting the normal
vector of the separating hyperplane from the SVM classifier that demonstrated the best
performance. Subsequently, the 5-dimensional data points were projected onto the choice axis,
resulting in 1-dimensional projections. These projections represented the positions along the
choice axis, with the intersection between the normal vector and the hyperplane serving as the zero
point.
The potential in this context is determined by integrating the spatial component of the time
derivative of the population activity, denoted as X_t , for a given condition. To be more precise, we
began by computing the expected value across trials of the time derivative of the unit activities for
each choice. Subsequently, we employed the potential function $V(x, t)$ to to evaluate the potential
at position x and time t,

854
$$V(x,t) = \left(\sum_{p<x} \frac{dX_t}{dt} | X_t = p \right) \cdot (-\Delta x)$$

855   Crucially, we computed the potential separately for each choice, establishing the potential
856 at position 0 as the zero-potential reference. To provide a holistic evaluation of the potential
857 landscape, we combined the potentials of both choices at position 0.
858
Furthermore, we expanded the computation of potentials to encompass each stimulus
coherence level using the aforementioned approach. Similar to the prior analysis, the time
derivatives were still derived from the 1-D projections of each data point onto the choice axis.
However, in this instance, the conditions were based on the stimulus coherence level rather than
the choice. For a representative potential value for each model, we averaged the potentials at the
time point of 300ms after the stimulus onset. Additionally, recognizing that the positions visited by
each model might differ, we ensured the accuracy of standard error estimation by retaining only
the potential values at positions continuously visited by four or more models.
867
868
869 **Acknowledgments**
870

871 We thank Tianming Yang, Matthew Rosen, Siyu Wang, Cheng Xue, Gongcheng Yu, Mingze Li
872 and Ou Zhu for their comments on an earlier version of this manuscript. We also thank the
873 veterinary staff at The University of Chicago Animal Resources Center for expert assistance.
874
875 This study was supported by STI2030-Major Projects (2021ZD0203800), NSFC32171036.

876
877

878 **References**
879

1   O'Connell, R. G., Shadlen, M. N., Wong-Lin, K. & Kelly, S. P. Bridging Neural and
Computational Viewpoints on Perceptual Decision-Making. *Trends Neurosci* **41**, 838-852,
doi:10.1016/j.tins.2018.06.005 (2018).
2   Freedman, D. J. & Assad, J. A. Neuronal Mechanisms of Visual Categorization: An
Abstract View on Decision Making. *Annu Rev Neurosci* **39**, 129-147,
doi:10.1146/annurev-neuro-071714-033919 (2016).
3   Gold, J. I. & Shadlen, M. N. The neural basis of decision making. *Annu Rev Neurosci* **30**,
535-574, doi:10.1146/annurev.neuro.29.051605.113038 (2007).
4   Huk, A. C., Katz, L. N. & Yates, J. L. The Role of the Lateral Intraparietal Area in (the Study
of) Decision Making. *Annu Rev Neurosci* **40**, 349-372, doi:10.1146/annurev-neuro-
072116-031508 (2017).
5   Hanks, T. D. & Summerfield, C. Perceptual Decision Making in Rodents, Monkeys, and
Humans. *Neuron* **93**, 15-31, doi:10.1016/j.neuron.2016.12.003 (2017).
6   Shadlen, M. N. & Kiani, R. Decision making as a window on cognition. *Neuron* **80**, 791-
806, doi:10.1016/j.neuron.2013.10.047 (2013).
7   Freedman, D. J. & Assad, J. A. A proposed common neural mechanism for categorization
and perceptual decisions. *Nat Neurosci* **14**, 143-146, doi:10.1038/nn.2740 (2011).
8   Platt, M. L. & Glimcher, P. W. Neural correlates of decision variables in parietal cortex.
Nature **400**, 233-238, doi:10.1038/22268 (1999).
9   Roitman, J. D. & Shadlen, M. N. Response of neurons in the lateral intraparietal area
during a combined visual discrimination reaction time task. *J Neurosci* **22**, 9475-9489
(2002).
10   Shadlen, M. N. & Newsome, W. T. Motion perception: seeing and deciding. *Proceedings
of the National Academy of Sciences of the United States of America* **93**, 628-633,
doi:10.1073/pnas.93.2.628 (1996).
11   Shadlen, M. N. & Newsome, W. T. Neural basis of a perceptual decision in the parietal
cortex (area LIP) of the rhesus monkey. *J Neurophysiol* **86**, 1916-1936,
doi:10.1152/jn.2001.86.4.1916 (2001).
12   Sugrue, L. P., Corrado, G. S. & Newsome, W. T. Matching behavior and the
representation of value in the parietal cortex. *Science* **304**, 1782-1787,
doi:10.1126/science.1094765 (2004).
13   Yates, J. L., Park, I. M., Katz, L. N., Pillow, J. W. & Huk, A. C. Functional dissection of signal
and noise in MT and LIP during decision-making. *Nat Neurosci* **20**, 1285-1292,
doi:10.1038/nn.4611 (2017).
14   Zhou, Y. & Freedman, D. J. Posterior parietal cortex plays a causal role in perceptual and
categorical decisions. *Science* **365**, 180-185, doi:10.1126/science.aaw8347 (2019).
15   Freedman, D. J. & Assad, J. A. Experience-dependent representation of visual categories
in parietal cortex. *Nature* **443**, 85-88, doi:10.1038/nature05078 (2006).
16   Freedman, D. J. & Assad, J. A. Distinct encoding of spatial and nonspatial visual
information in parietal cortex. *J Neurosci* **29**, 5671-5680, doi:10.1523/JNEUROSCI.2878-
08.2009 (2009).
17   Bennur, S. & Gold, J. I. Distinct representations of a perceptual decision and the
associated oculomotor plan in the monkey lateral intraparietal area. *J Neurosci* **31**, 913-
921, doi:10.1523/JNEUROSCI.4417-10.2011 (2011).

18 Swaminathan, S. K. & Freedman, D. J. Preferential encoding of visual categories in
parietal cortex compared with prefrontal cortex. *Nat Neurosci* **15**, 315-320,
doi:10.1038/nn.3016 (2012).
19 Zhou, Y., Mohan, K. & Freedman, D. J. Abstract Encoding of Categorical Decisions in
Medial Superior Temporal and Lateral Intraparietal Cortices. *J Neurosci* **42**, 9069-9081,
doi:10.1523/JNEUROSCI.0017-22.2022 (2022).
20 Kiani, R. & Shadlen, M. N. Representation of confidence associated with a decision by
neurons in the parietal cortex. *Science* **324**, 759-764, doi:10.1126/science.1169405
(2009).
21 Katz, L. N., Yates, J. L., Pillow, J. W. & Huk, A. C. Dissociated functional significance of
decision-related activity in the primate dorsal stream. *Nature* **535**, 285-288,
doi:10.1038/nature18617 (2016).
22 Zhou, Y., Zhu, O. & Freedman, D. J. Posterior Parietal Cortex Plays a Causal Role in
Abstract Memory-Based Visual Categorical Decisions. *J Neurosci* **43**, 4315-4328,
doi:10.1523/JNEUROSCI.2241-22.2023 (2023).
23 Kim, J. N. & Shadlen, M. N. Neural correlates of a decision in the dorsolateral prefrontal
cortex of the macaque. *Nat Neurosci* **2**, 176-185, doi:10.1038/5739 (1999).
24 Rossi-Pool, R. *et al.* Decoding a Decision Process in the Neuronal Population of Dorsal
Premotor Cortex. *Neuron* **96**, 1432-1446 e1437, doi:10.1016/j.neuron.2017.11.023
(2017).
25 Ding, L. & Gold, J. I. Neural correlates of perceptual decision making before, during, and
after decision commitment in monkey frontal eye field. *Cereb Cortex* **22**, 1052-1067,
doi:10.1093/cercor/bhr178 (2012).
26 Ferrera, V. P., Yanike, M. & Cassanello, C. Frontal eye field neurons signal changes in
decision criteria. *Nat Neurosci* **12**, 1458-1462, doi:10.1038/nn.2434 (2009).
27 Heitz, R. P. & Schall, J. D. Neural mechanisms of speed-accuracy tradeoff. *Neuron* **76**,
616-628, doi:10.1016/j.neuron.2012.08.030 (2012).
28 Zhou, Y. *et al.* Distributed functions of prefrontal and parietal cortices during sequential
categorical decisions. *Elife* **10**, doi:10.7554/eLife.58782 (2021).
29 Horwitz, G. D. & Newsome, W. T. Separate signals for target selection and movement
specification in the superior colliculus. *Science* **284**, 1158-1161,
doi:10.1126/science.284.5417.1158 (1999).
30 Peysakhovich, B. *et al.* Primate superior colliculus is engaged in abstract higher-order
cognition. *bioRxiv*, doi:10.1101/2023.01.17.524416 (2023).
31 Ding, L. & Gold, J. I. Caudate encodes multiple computations for perceptual decisions. *J
Neurosci* **30**, 15747-15759, doi:10.1523/JNEUROSCI.2894-10.2010 (2010).
32 Gold, J. I. & Shadlen, M. N. Representation of a perceptual decision in developing
oculomotor commands. *Nature* **404**, 390-394, doi:10.1038/35006062 (2000).
33 Jun, E. J. *et al.* Causal role for the primate superior colliculus in the computation of
evidence for perceptual decisions. *Nat Neurosci* **24**, 1121-1131, doi:10.1038/s41593-
021-00878-6 (2021).
34 Shushruth, S., Mazurek, M. & Shadlen, M. N. Comparison of Decision-Related Signals in
Sensory and Motor Preparatory Responses of Neurons in Area LIP. *J Neurosci* **38**, 6350-
6365, doi:10.1523/JNEUROSCI.0668-18.2018 (2018).

35   Engel, T. A., Chaisangmongkon, W., Freedman, D. J. & Wang, X. J. Choice-correlated
activity fluctuations underlie learning of neuronal category representation. *Nat Commun*
6, 6454, doi:10.1038/ncomms7454 (2015).

36   Masse, N. Y., Yang, G. R., Song, H. F., Wang, X. J. & Freedman, D. J. Circuit mechanisms
for the maintenance and manipulation of information in working memory. *Nat Neurosci*
22, 1159-1167, doi:10.1038/s41593-019-0414-3 (2019).

37   Song, H. F., Yang, G. R. & Wang, X. J. Training Excitatory-Inhibitory Recurrent Neural
Networks for Cognitive Tasks: A Simple and Flexible Framework. *PLoS Comput Biol* **12**,
e1004792, doi:10.1371/journal.pcbi.1004792 (2016).

38   Deco, G., Rolls, E. T., Albantakis, L. & Romo, R. Brain mechanisms for perceptual and
reward-related decision-making. *Prog Neurobiol* **103**, 194-213,
doi:10.1016/j.pneurobio.2012.01.010 (2013).

39   Finkelstein, A. *et al.* Attractor dynamics gate cortical information flow during decision-
making. *Nat Neurosci* **24**, 843-850, doi:10.1038/s41593-021-00840-6 (2021).

40   Wang, S., Falcone, R., Richmond, B. & Averbeck, B. B. Attractor dynamics reflect decision
confidence in macaque prefrontal cortex. *Nat Neurosci* **26**, 1970-1980,
doi:10.1038/s41593-023-01445-x (2023).

41   Wang, X. J. Probabilistic decision making by slow reverberation in cortical circuits.
Neuron **36**, 955-968, doi:10.1016/s0896-6273(02)01092-9 (2002).

42   Wong, K. F., Huk, A. C., Shadlen, M. N. & Wang, X. J. Neural circuit dynamics underlying
accumulation of time-varying evidence during perceptual decision making. *Front*
Comput Neurosci **1**, 6, doi:10.3389/neuro.10.006.2007 (2007).

43   Motter, B. C. Neural correlates of attentive selection for color or luminance in
extrastriate area V4. *J Neurosci* **14**, 2178-2189, doi:10.1523/JNEUROSCI.14-04-
02178.1994 (1994).

44   Motter, B. C. Neural correlates of feature selective memory and pop-out in extrastriate
area V4. *J Neurosci* **14**, 2190-2199, doi:10.1523/JNEUROSCI.14-04-02190.1994 (1994).

45   Martinez-Trujillo, J. C. & Treue, S. Feature-based attention increases the selectivity of
population responses in primate visual cortex. *Curr Biol* **14**, 744-751,
doi:10.1016/j.cub.2004.04.028 (2004).

46   Bichot, N. P., Rossi, A. F. & Desimone, R. Parallel and serial neural mechanisms for visual
search in macaque area V4. *Science* **308**, 529-534, doi:10.1126/science.1109676 (2005).

47   Maunsell, J. H. & Treue, S. Feature-based attention in visual cortex. *Trends Neurosci* **29**,
317-322, doi:10.1016/j.tins.2006.04.001 (2006).

48   Federer, F., Ta'afua, S., Merlin, S., Hassanpour, M. S. & Angelucci, A. Stream-specific
feedback inputs to the primate primary visual cortex. *Nat Commun* **12**, 228,
doi:10.1038/s41467-020-20505-5 (2021).

49   Briggs, F. & Usrey, W. M. Parallel processing in the corticogeniculate pathway of the
macaque monkey. *Neuron* **62**, 135-146, doi:10.1016/j.neuron.2009.02.024 (2009).

50   Briggs, F. Role of Feedback Connections in Central Visual Processing. *Annu Rev Vis Sci* **6**,
313-334, doi:10.1146/annurev-vision-121219-081716 (2020).

51   Harris, K. D. & Mrsic-Flogel, T. D. Cortical connectivity and sensory coding. *Nature* **503**,
51-58, doi:10.1038/nature12654 (2013).

52   Moore, T. & Armstrong, K. M. Selective gating of visual signals by microstimulation of
frontal cortex. *Nature* **421**, 370-373, doi:10.1038/nature01341 (2003).

53   Gilbert, C. D. & Li, W. Top-down influences on visual processing. *Nature reviews.*
Neuroscience **14**, 350-363, doi:10.1038/nrn3476 (2013).
54   McManus, J. N., Li, W. & Gilbert, C. D. Adaptive shape processing in primary visual
cortex. *Proceedings of the National Academy of Sciences of the United States of America*
108, 9739-9746, doi:10.1073/pnas.1105855108 (2011).
55   Nienborg, H. & Cumming, B. G. Decision-related activity in sensory neurons reflects
more than a neuron's causal effect. *Nature* **459**, 89-92, doi:10.1038/nature07821
(2009).
56   Bondy, A. G., Haefner, R. M. & Cumming, B. G. Feedback determines the structure of
correlated variability in primary visual cortex. *Nat Neurosci* **21**, 598-606,
doi:10.1038/s41593-018-0089-1 (2018).
57   Wimmer, K. *et al.* Sensory integration dynamics in a hierarchical network explains choice
probabilities in cortical area MT. *Nat Commun* **6**, 6177, doi:10.1038/ncomms7177
(2015).
58   Prat-Ortega, G., Wimmer, K., Roxin, A. & de la Rocha, J. Flexible categorization in
perceptual decision making. *Nat Commun* **12**, 1283, doi:10.1038/s41467-021-21501-z
(2021).
59   Wong, K. F. & Wang, X. J. A recurrent network mechanism of time integration in
perceptual decisions. *J Neurosci* **26**, 1314-1328, doi:10.1523/JNEUROSCI.3733-05.2006
(2006).
60   Rao, R. P. & Ballard, D. H. Predictive coding in the visual cortex: a functional
interpretation of some extra-classical receptive-field effects. *Nat Neurosci* **2**, 79-87,
doi:10.1038/4580 (1999).
61   Friston, K. & Kiebel, S. Predictive coding under the free-energy principle. *Philos Trans R*
Soc Lond B Biol Sci **364**, 1211-1221, doi:10.1098/rstb.2008.0300 (2009).
62   Bastos, A. M. *et al.* Canonical microcircuits for predictive coding. *Neuron* **76**, 695-711,
doi:10.1016/j.neuron.2012.10.038 (2012).
63   Shipp, S. Neural Elements for Predictive Coding. *Front Psychol* **7**, 1792,
doi:10.3389/fpsyg.2016.01792 (2016).
64   Kobak, D. *et al.* Demixed principal component analysis of neural population data. *Elife* **5**,
doi:10.7554/eLife.10989 (2016).
65   Chaoming Wang, X. C., Tianqiu Zhang, Si Wu. BrainPy: a flexible, integrative, efficient,
and extensible framework towards general-purpose brain dynamics programming.
bioRxiv, doi:https://doi.org/10.1101/2022.10.28.514024 (2022).

1047

1048 **Figures**
1049

1050
1051
Figure 1. Behavioral task.
(**A**) The flexible visual-motion discrimination (FVMD) task. Monkeys needed to report their decision
about the direction of the visual motion stimuli by choosing either the green or red saccade target.
The appearance of the two color targets preceded that of the visual-motion stimulus, and the target
positions were randomly chosen on each trial to avoid fixed mapping between motion stimulus and
saccade direction. Monkeys could initiate their saccade as soon as they had made their decision.
(**B-C**) Psychometric curves for the two monkeys. The performance accuracy for each monkey is
plotted as the proportion of trials in which 315° was chosen as a function of the directions and
coherence levels of the motion stimuli. (**D-E**) Chronometric curves are shown separately for the
two monkeys. (**F-G**) Two trial conditions, contralateral (F) and ipsilateral (G), defined according to
the spatial configurations of task stimuli during neural recording.
1063
1064
1065

1066
1067

Figure 2. Nonlinear feedback modulation of saccade choice on sensory evaluation in LIP.
(**A**) The activity of one example neuron in the CT condition of the FVMD task is shown for each motion-coherence level. The zero-coherence trials were grouped based on the monkey's choices. The two vertical dashed lines denote the time of target and motion stimulus onset, respectively. (**B**) The same example neuron's activity in the IT condition of the FVMD task. (**C-F**) The activities of two more example neurons. (**G-H**) The averaged population activities of all direction-selective neurons (n = 83) that were collected during the recording sessions in which the saccade targets were arranged in either horizontal or oblique directions. The activity to each motion direction and coherence level is shown separately for the CT condition (G) and IT condition (H). (**I**) The activity differences between CT and IT conditions (CT minus IT) of single LIP neurons were plotted for both preferred and nonpreferred motion directions. Each dot represents the activity of a single neuron. The histograms in horizontal and vertical axis represent the distribution of activity difference between CT and IT conditions for preferred and nonpreferred motion directions, respectively. (**J**) An ROC analysis was used for quantifying the motion DS for both CT (solid) and IT (dashed) conditions. The colored dots denote the time points for which there was a significant difference between the CT and IT conditions (paired t-test: P < 0.01). (**K**) The average DS for low- and zero-coherence trials is shown as in (J). (**L**) Variance in LIP population activity as explained by the individual demixed principal components. Each bar shows the proportion of total explained variance that was contributed by the four task variables. The pie chart shows the total variance explained by each task variable. H, high; M, medium; L, low; 0, zero; P, preferred; NP nonpreferred.

1088
1089

1090
1091
Figure 3. Feedback modulation specifically impacted the decision-correlated activity .
(**A-B**) Averaged population activities on low-coherence trials in the CT condition (A) and IT
condition (B) are shown separately for correct (corr) and error (err) trials. (**C-D**) An ROC analysis
quantified the stimulus-related and decision-related LIP DS on low-coherence trials. The
correlations between LIP neural activity and the monkeys' decisions about motion direction (red)
or the physical direction of the motion stimulus (blue) in both the CT (C) and IT (D) conditions are
shown over time. (**E-F**) Partial correlation analysis revealed the decision-related and stimulus-
related components of LIP activity. The values for r-decision (E) (the partial correlation between
neuronal activity and monkeys' choice, given the stimulus direction) and r-stimulus (F) (partial
correlation between neuronal activity and stimulus direction, given the monkeys' choices) were
compared between the IT and CT conditions. (**G-H**) Correlation between LIP DS and the monkeys'
RTs on zero-coherence trials. The choice selectivity on zero-coherence trials is shown for the faster
RT and slower RT trials. Shaded areas denote ±SEM. The black stars indicate time periods in
which there was a significant difference (paired t-test: P < 0.01).
1106
1107

1108
1109
Figure 4. Multi-module RNNs trained with the FVMD task.
(**A**) Model schematic of the RNNs. Each RNN consists of nine motion direction tuned input units,
eight color tuned target input units, 200 hidden units, and two response units. The hidden layer of
each RNN consists of two main modules. Each main module consists of two nominal modules,
each of which receives either the visual motion input (motion module) or the target color input
(target module). Only the target modules project to the two response units, and each main module
projects primarily to one response unit. All four nominal modules were assigned with equal number
of units (25%) in the network, which consisted of 80% excitatory and 20% inhibitory units. (**B-C**)
The performance accuracies (B) and RTs (C) of all 50 trained RNNs are shown separately for each
motion coherence level. (**D-E**) Two example units from an example RNN. (**D**) The neural activity of
an example unit from the motion module is shown for each motion direction and coherence level.
(**E**) The neural activity of an example unit from the target module is shown for each saccade
direction and coherence level. (**F-G**) The motion DS for the motion module (F) and saccade DS for
the target module (G) for the example RNN. (**H-I**) The averaged population activities of all direction-
selective units in the motion module of the example RNN are shown for the CT condition (H) and
IT condition (I). (**J-K**) The averaged motion DS in the motion module of the example RNN for both
the CT (solid) and IT (dashed) conditions was quantified by ROC analysis. (**L-M**) Partial correlation
analysis. The values for r-decision (L) and r-stimulus (M) are compared between the IT and CT
conditions for the example RNN. (**N**) A comparison of the motion DS in the motion module of all
trained RNNs between the CT and IT conditions is shown for each coherence level. (**O**) A
comparison of the r-decision and r-stimulus between the IT and CT conditions is shown for all
trained RNNs. (Paired t-test: ***, P < 0.001).
1132
1133

1134
1135
Figure 5. The circuit mechanisms underlying the nonlinear feedback modulation in RNNs.
(**A**) The averaged cross-module connection weights are shown for both feedforward and feedback
connections. Units in the motion module (M) and target module (T) were grouped based on their
preferences for motion DS (315° vs. 135°) and target color selectivity (red [r] vs. green [g]),
respectively. (**B**) The correlation between the match extent of the neural encoding between units
in the motion module and those in target modules and the connection weights between them. Each
dot denotes data from one RNN. (**C-D**) A comparison of the performance accuracy (C) and RT (D)
of the full-model RNNs, RNNs without feedback connections, and RNNs with shuffled feedback
connections. (**E-F**) The averaged activity of units in the motion module of the example RNN after
the feedback connections were ablated is shown for CT (E) and IT (F) conditions. (**G**) The
comparison of the averaged motion DS between CT and IT conditions for all trained RNNs after
feedback connections were ablated. An ROC analysis was used to quantify the motion DS. (**H**) The
differences in motion DS between CT and IT conditions were compared between full-model RNNs
and RNNs with shuffled feedback connectivity.
1150
1151

1152

Figure 6. The computational mechanisms underlying the nonlinear feedback modulation in decision-making.
(**A**) The averaged activity of units in the target module of the example RNN, after the feedback connections were ablated, is shown separately for different saccade directions and coherence levels. (**B**) The averaged saccade DS in the target module of the example RNN after full feedback ablation. (**C**) The averaged saccade DS in the target module of the full-model RNNs, RNNs without feedback connections, and RNNs with shuffled feedback connections. (Paired t-test: *, P < 0.05; **, P < 0.01; ***, P < 0.001; n.s., not significant). (**D-G**) The evolution of the averaged energy landscapes was shown over time. A numerical approximation of the energy landscape in the 1-D decision (saccade choice) subspace is constructed for both full-model RNNs and RNNs with various types of projection-specific inactivation. Each plot represents the averaged results of 50 RNNs, where Position 0 signifies the SVM decision boundary, and the vertical dashed line marks the time of motion stimulus onset. Unvisited portions of the state space are left blank as there is no gradient or potential estimate. (**H-k**) Averaged numerical estimate of energy landscapes for trials with different task difficulty levels (motion coherence). Results from two different saccade choices were average together. Only the potential values at positions continuously visited by four or more models were retained. Shaded areas denote ±SEM.

1170