Selectivity-specialized feedback connections mediate the interaction of distinct neural processes during flexible decision-making

Authors and author addresses:

Xuanyu Wu[1] & Yang Zhou[1,2,3]

1. Peking-Tsinghua Center for Life Sciences, Peking University, Beijing, 100871, China
2. School of Psychological and Cognitive Sciences, Peking University, Beijing, 100871, China
3. PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing, 100871, China

* Corresponding author: Yang Zhou, yangzhou1@pku.edu.cn

Lead contact: Yang Zhou, Ph.D., School of Psychological and Cognitive Sciences, Peking University, Beijing, 100871, China.

Email: yangzhou1@pku.edu.cn

Tel: +86 18638526400

Abstract:

Neural activity in the primate brain correlates with both sensory evaluation and action selection aspects of decision-making. However, the neural mechanisms underlying the interplay between these processes remain unclear. Here, we examined the modulation of sensory evaluation by action selection in posterior parietal cortex (PPC) when monkeys used saccade choices to report their decisions about visual stimuli in a flexible decision task. We found that the PPC activity related to monkeys' abstract decisions about the visual stimuli was significantly modulated by monkeys' following saccade choices directing outside each neuron's response field. Modeling with recurrent neural networks trained to perform the same task indicated that the feedback connections that matched the learned stimuli-response associations during the task mediated the interplay of these decision processes and were necessary for efficiently generating flexible decisions. These results indicate an iterative computation between different decision processes that could be mediated primarily by precise feedback connectivity, and potentially provide a novel framework for decision-making.

Introduction:

Decision-making usually involves evaluating the behavioral relevance of incoming sensory stimuli (i.e., sensory evaluation) and selecting appropriate motor responses (i.e., action selection)[1-7]. Neural activity related to either the sensory or motor aspects of decision-making is widely distributed across different brain areas, such as PPC[8-21], frontal cortex[18,22-26], superior colliculus[27,28], and caudate[29]. A recent inactivation study has shown that the lateral intraparietal area (LIP), a subregion of PPC, plays a causal role in flexible visuomotor decisions, with preferential involvement in sensory evaluation rather than action selection[14]. Meanwhile, areas outside PPC, such as frontal eye field and superior colliculus, have been shown to be causally involved in the action selection aspect of decision-making[30,31]. These findings suggest that sensory evaluation and action selection likely involve distinct neural processing in the primate brain during flexible decision-making.

However, previous reports have also shown that neural activity related to sensory evaluation and action selection overlap temporally and spatially in the brain[32]. This leaves open the possibility of interplay between the two distinct processes during decision-making. From the perspective of information processing, the abstract result of sensory evaluation should be transmitted to the motor planning circuits to guide the action selection process in a feedforward manner, although both processes could proceed in parallel in the brain. However, whether and how action selection processes might exert a feedback influence on sensory evaluation have not been explicitly studied.

In this study, we examined the activities of single LIP neurons during a flexible decision task in which monkeys needed to report their decisions about a motion stimulus with a saccade to one of two color targets. Specifically, we arranged the motion stimuli

inside each neuron's response field (RF), and positioned the saccade targets in the direction
perpendicular to the axis of neural RFs. We found that LIP activity related to evaluate the
motion stimuli was significantly modulated by the monkeys' following saccade choice
direction relative to the recorded brain hemisphere. Such modulation was not a simple
addictive gain modulation but aligned with the functional properties of each neuron. This
suggests a precise 'feedback' modulation from action selection process to the neural
processing of sensory evaluation during decision-making.

RNN models trained on complex behavioral tasks have shown promise for
understanding neural computations and circuit mechanisms underlying cognitive
functions[33-35]. To further explore how action selection can modulate sensory evaluation
during decision-making, we trained multi-module RNNs, which consist of different
hemispheres and RF structures, to perform the same decision task and analyzed the activity
of network units during task performance. These RNNs exhibited similar behavioral
performance and neural activity patterns as observed in the monkey electrophysiology
experiment, including patterns of feedback modulation from action selection to sensory
evaluation. Combining network connectivity analysis with projection-specific inactivation
experiments in the RNNs, we further showed that the precise feedback connections
between units that showed matched functional properties within different network modules
were the key circuit mechanism for mediating the modulation of sensory evaluation by
action selection, which facilitated the RNNs to solve the flexible decision task accurately
and rapidly.

Results:

LIP neurons' selectivity for visual motion is modulated by saccade choices made away
from their response fields (RFs)

To test the mechanisms underlying the modulation of action selection on sensory
evaluation, we trained two monkeys to perform a reaction-time version of the flexible
visual-motion discrimination (FVMD) task, in which they needed to choose one of two
colored saccade targets based on their decisions about the motion direction of a sample
stimulus shown at different coherence levels (**Fig. 1a**). Monkeys learned the mappings
between the two motion directions (315° and 135°) and two target colors (red and green)
at the start of training, and the mappings were fixed across the study. Because the locations
of the red and green targets were randomly interleaved across trials, neither of the motion
directions was directly linked with one specific saccade direction. Both the performance
accuracy and the reaction time (RT) changed systematically as a function of motion
coherence levels (**Fig. 1b-e**): as the motion coherence increased, both monkeys chose the
correct target more frequently and more rapidly.

We recorded single-neuron activity from the LIP while the monkeys performed the
FVMD task. To examine the neural activity related to the evaluation of stimulus motion,
we presented the motion stimuli within the RF of each neuron but arranged the saccade

targets at locations orthogonal to the axis of the neural RF. In total, 104 of 194 visually responsive LIP neurons (monkey M: 50/78; monkey B: 54/116) showed significant direction selectivity (DS) to the motion stimuli (one-way ANOVA, $P < 0.01$). To examine the influence of action selection on sensory evaluation, we analyzed data from the subset of sessions in which the saccade targets were aligned more closely with the horizontal direction than the vertical direction (83 of 104 neurons). In these sessions, the motor planning corresponding to a saccade to either target would be mediated primarily by one brain hemisphere. We therefore defined the conditions under which the correct target was contralateral or ipsilateral to the recorded hemisphere as the contralateral target (CT) condition or ipsilateral target (IT) condition, respectively (**Fig. 1f,g**).

Fig. 2a-f shows three example LIP neurons that exhibited significant motion coherence correlated DS. Surprisingly, LIP neurons showed greater DS in the CT condition than in the IT condition, even though the same motion stimuli were used in the same spatial location for both conditions. The averaged population activity showed this DS difference between CT and IT conditions for all four coherence levels (**Fig. 2g,h** and **Extended data fig. 1a-d**). During presentation of their preferred motion direction, LIP neurons showed significantly elevated activity in the CT relative to the IT at all coherence levels (nested ANOVA: $P_{(high)} = 0.0326$, $F = 4.65$; $P_{(medium)} = 0.0088$, $F = 7.03$; $P_{(low)} = 0.0076$, $F = 7.32$; $P_{(zero)} = 0.0124$, $F = 6.4$), and a trend toward lower activity to the nonpreferred direction for CT vs. IT (nested ANOVA: $P_{(high)} = 0.0994$, $F = 2.75$; $P_{(medium)} = 0.0649$, $F = 3.12$; $P_{(low)} = 0.0311$, $F = 4.73$; $P_{(zero)} = 0.0273$, $F = 4.96$). Most of the LIP neurons (48 of 83) showed such opposing trends in activity modulation between the preferred and nonpreferred directions (**Extended data fig. 1e**). Receiver operating characteristic (ROC) analysis further confirmed significantly greater motion DS in the CT condition than in the IT condition (**Fig. 2j,k**; nested ANOVA: $P_{(high)} = 5.0e-4$, $F = 12.44$; $P_{(medium)} = 9.53e-6$, $F = 20.91$; $P_{(low)} = 9.33e-7$, $F = 26.03$; $P_{(zero)} = 2.56e-8$, $F = 34.3$). Such DS differences were observed even before stimulus onset. Meanwhile, LIP neurons exhibited similar levels of mean activity between different saccade directions (CT vs. IT) before monkeys' saccade choice (**Fig. 2i**). These results indicated that the influence of saccade direction on LIP neural activity was not a simple additive gain process but was instead a precise modulation aligned with the response property of each neuron. Furthermore, a demixed principal component analysis on the pseudo-population activity (see Methods) revealed that the saccade direction related representation was a substantial component of LIP activity, as the saccade direction and motion–saccade interaction together explained a similar amount of variance in the population activity as the stimulus motion direction did (**Fig. 2l** and **Extended data fig. 2**). In contrast, we did not observe significant DS differences between the CT and IT conditions in the data sessions in which the saccade targets were aligned close to the vertical direction (21 of 104 neurons, **Extended data fig. 3**).

LIP motion DS correlates more closely with monkeys' decisions in the CT condition than in the IT condition

We then examined whether the LIP DS was more decision-correlated in the CT condition
than in the IT condition. As shown in **Fig. 2j**, the LIP DS on zero-coherence trials, which
reflected monkeys' trial-by-trial categorical choice, was significantly greater in the CT
condition than in the IT condition (nested ANOVA: P = 2.56e-8, F= 34.3). We then
quantified the correlation between LIP neural activity and the trial-by-trial categorical
choice or the physical motion direction by comparing LIP neural activity on correct versus
incorrect trials. We used low-coherence trials, as monkeys made enough errors in these
trials. In the CT condition but not the IT condition, LIP DS at the population level was
significantly reversed in sign on incorrect trials as compared to correct trials (**Fig. 3a,b**;
nested ANOVA: $P_{(CT)} = 0.0045$, F = 8.32). Accordingly, in the CT condition but not in the
IT condition, LIP activity correlated more closely with the monkeys' trial-by-trial abstract
decisions about motion direction, as opposed to the physical motion direction (**Fig. 3c,d**;
nested ANOVA, $P_{(CT)} = 0.0016$, F = 10.31; $P_{(IT)} = 0.443$, F = 0.59). Meanwhile, LIP activity
correlated more closely with the trial-by-trial abstract decisions during the CT condition
relative to the IT condition on the low coherence trials (nested ANOVA: P = 1.95e-8, F =
34.94). Furthermore, we used partial correlation analysis to examine decision- and
stimulus-related components of DS (i.e., r-decision and r-stimulus, **Fig. 3e,f**) using all four
coherence levels. The decision-related component of LIP DS was significantly greater in
the CT condition than in the IT condition (**Fig. 3e**; nested ANOVA: P = 1.07e-6, F= 25.72),
and this difference emerged even before motion stimulus onset. This suggests that the LIP
DS was more closely correlated with monkeys' decisions in the CT condition than in the
IT condition. However, the stimulus-related component of the LIP DS was similar at the
population level between the two conditions (**Fig. 3f**; nested ANOVA: P = 0.7606, F =
0.09), suggesting that the modulation of LIP DS by the later saccade choice was primarily
related to a decision process rather than basic sensory processing.

We next examined the correlation between LIP DS and the monkeys' RTs. Shortly
after motion stimulus onset, LIP neurons showed greater population response to their
preferred motion direction on shorter versus longer RT trials for all motion coherence
levels in the CT condition (**Extended data fig. 4**) but not in the IT condition (**Extended
data fig. 5**). On zero-coherence trials (**Fig. 3g,h**), DS evolved more rapidly on shorter vs.
longer RT trials in the CT condition but not in the IT condition (bootstrap: $P_{(CT)} < 0.01$,
$P_{(IT)} = 0.87$). Thus, LIP DS was more closely correlated with decision behavior in the CT
condition than the IT condition, including the trial-by-trial categorical choice and RTs of
the monkeys.

**Trained multi-module RNNs exhibited similar behavioral performance and neural
activity patterns as observed in the monkey electrophysiology experiment**

Our electrophysiology data show that action selection significantly modulates the neural
processing of sensory evaluation, which indicates a complex, iterative computation for
flexible decision-making. Training RNNs on behavioral tasks used in experimental
neurophysiological studies has proven to be helpful for exploring putative circuit
computations underlying cognitive tasks[34-36]. Therefore, we trained multi-module RNNs to

perform the FVMD task to examine the circuit mechanisms underlying the interplay of
different decision processes. We adopted simple neurobiological principles to constrain the
connection structure but not the functional roles to generate recurrently connected modules
(see Methods). Because our neurophysiological results showed that the modulation effect
of action selection on sensory evaluation depended on the saccade direction relative to the
recorded brain hemisphere, we implemented RNNs composed of two main modules
organized in parallel to simulate two brain hemispheres. Each main module consisted of
two nominal modules, with each nominal module receiving the visual motion input (motion
module) or the target color input (target module), corresponding to the two neuron
populations whose RFs covered the motion stimulus or saccade targets, respectively (**Fig.
4a**).

We independently trained 50 such networks with randomly initialized weights and
identical hyperparameters to perform the FVMD task. The FVMD task setup used for
training RNNs was tailored to match the monkey experiments. Specifically, the RNNs were
trained using motion stimuli with two high-coherence levels and were tested using stimuli
with another four different coherence levels (high, medium, low, and zero). After training,
all 50 networks converged to perform the FVMD task with high accuracies (>99%) on the
training coherence and exhibited coherence-dependent performances when tested with
untrained stimuli (**Fig. 4b,c**): accuracy increased and RT decreased with motion coherence.

We then analyzed the unit activity in different modules when the trained RNNs
were tested with untrained motion stimuli. Across all networks, the units in the motion
module and target modules exhibited activity corresponding to sensory evaluation and
action selection, respectively. Specifically, units in the motion modules showed coherence-
correlated motion DS (**Fig. 4d,f** and **Extended data fig. 6,7**; mean r = 0.29; one-sample t-
test: P = 4.09e-42, t(49) = 46.31), and neuronal activity on zero-coherence motion trials
reflected the trial-by-trial abstract decisions of the networks. Such motion DS resembled
our electrophysiology data when the motion stimuli were presented within the RFs of the
LIP neurons. Meanwhile, units in the target modules showed coherence-correlated saccade
DS (**Fig. 4e,g** and **Extended data fig. 6,7**), and the neuronal activity on zero-coherence
motion trials also showed significant choice probability (t-test: $P_{(example\ RNN)}$ = 7.06e-8, t(68)
=15.83; $P_{(population)}$ = 2.08e-22, t(49) = 17.21). Such saccade DS was consistent with the
commonly observed decision-related activity in previous studies in which the saccade
target was presented within the RFs of LIP neurons[9-11].

Furthermore, we found that the motion DS of units in the motion modules was
significantly greater in the CT condition than in the IT condition for the majority of RNNs
(**Fig. 4h,i** and **Extended data fig. 8a,b**): in the CT condition relative to the IT condition,
activity was significantly greater for the preferred motion direction but weaker for the
nonpreferred direction (**Extended data fig. 9**). ROC analysis confirmed that the motion
DS in the motion module was significantly greater for all four coherence levels in the CT
vs. IT conditions in most RNNs (**Fig. 4j,k,n** and **Extended data fig. 8c,d**; two-way
ANOVA: $P_{(example\ RNN)}$ = 1.53e-11, F = 47.49; $P_{(population)}$ = 3.50e-77, F = 660.84). Similar

to the monkey electrophysiology data, the decision-correlated motion DS (r-decision) in
the motion module was dramatically reduced in the IT versus CT condition (paired t-test:
P = 6.84e-17, t(49)= -12.53), and this difference was substantially greater than the
difference in the stimulus-related (r-stimulus) motion DS between the two conditions
(paired t-test: P = 2.02e-14, t(49) = -10.70; **Fig. 4l,m,o** and **Extended data fig. 8e,f**). These
results indicated that action selection also markedly modulated the neural processing of
sensory evaluation during decision-making in the RNNs.

Selectivity-specialized feedback connections mediate the modulation of sensory
evaluation by action selection

Our multi-module RNNs showed a remarkable similarity to both the behavioral
performances and the neural activity patterns in the monkey electrophysiology data,
suggesting that these networks may have adopted mechanisms similar to those in the
monkey brain to mediate the decision processes in the FVMD task. Therefore, we explored
the potential circuit mechanisms underlying the modulation of action selection on sensory
evaluation in RNNs. We hypothesized that this modulation might result from the feedback
connections from the target module to the motion module. To test this hypothesis, we
examined the recurrent weights of the cross-module connections (see Methods). Both the
feedback and feedforward cross-module connection weights were significantly greater
between the selectivity-matched unit pairs (e.g., units in the motion module preferred to
encode 315° and units in the target module preferred to encode red) than the selectivity-
nonmatched unit pairs (e.g., units in the motion module preferred to encode 315° while
units in the target module preferred to encode green) (**Fig. 5a**; paired t-test: $P_{(feedforward)}$ =
7.18e-16, t(49) = -11.76; $P_{(feedback)}$ = 5.26e-18, t(49) = -13.40). In particular, the average
feedback connectivity was primarily excitatory between the selectivity-matched unit pairs
but was inhibitory between the selectivity-nonmatched unit pairs. Furthermore, most of the
networks exhibited positive correlations between the connection weight and the match
extent of the neural selectivity of each cross-module unit pair (**Fig. 5b**; mean r = 0.11; one-
sample t-test: P = 6.39e-16; t(49) = 11.79; see Methods), indicating that units that exhibited
stronger encoding preference to one motion direction in the motion module were more
likely to receive more extensive feedback projections from the units that exhibited stronger
encoding on the matched target color in the target module. These results suggested precise
feedback connections between RNN modules that were aligned with the functional
properties of different units.

To test the causal contribution of the selectivity-specialized feedback connections to
the modulation of action selection on sensory evaluation, we conducted three projection-
specific inactivation experiments in the RNNs (see Methods). First, we performed
nonspecific ablation of all the feedback projections from the target module to the motion
module when testing with untrained motion stimuli. In most of the tested RNNs, this caused
a dramatic reduction in their performance in all nonzero-coherence stimulus conditions
(**Fig. 5c**; two-way ANOVA: P = 4.74e-80, F = 704.65) as well as significantly prolonged
RTs for high- and medium-coherence stimulus conditions (**Fig. 5d**; two-way ANOVA: P

= 2.45e-4, F = 13.79). The mean RTs became more dispersed across all the trained RNNs
and did not differ among different motion coherences after ablation (one-way ANOVA: P
= 0.68, F = 0.50). Ablating feedback connections also significantly affected the motion DS
in the motion module. On average, although the averaged motion DS did not significantly
decrease after ablation (**Extended data fig. 10a,c**), the difference in the motion DS
between the CT and IT conditions vanished (**Fig. 5e-g** and **Extended data fig. 11a,b**; two-
way ANOVA: $P_{(CT\ vs.\ IT)}$ = 0.14, F = 2.21). Strikingly, ablating feedback connections also
resulted in significant changes in the patterns of saccade DS in the target module. Without
feedback connections, the magnitude of the saccade DS in the target module significantly
decreased in nonzero-coherence trials (**Fig. 5i-k**; two-way ANOVA: P = 6.90e-5, F = 16.29)
and did not change with motion coherence (one-way ANOVA: P = 0.12, F = 1.98).

To further test the importance of different patterns of feedback connectivity for the
RNNs to solve the FVDM task, we next performed pattern-specific ablation of the feedback
connections in RNNs. We separated the feedback projections in each RNN into specialized
(i.e., the feedback connection weights were positive or negative between the selectivity-
matched or selectivity-nonmatched unit pairs, respectively) and nonspecialized groups.
Despite similar numbers and weights of the feedback connections in these two groups
(**Extended data fig. 12a,b**), ablating the feedback connection in the specialized group
versus the nonspecialized group resulted in much more severe impairments in the RNNs'
behavior performance (**Extended data fig. 12c,d**). This indicated that the feedback
connections that reflected the learned stimulus–response association might be more
important for the RNNs to solve the FVDM task than the other feedback connections.

Second, we disrupted the selectivity-specialized feedback connectivity without
changing the total strength of the feedback connections by randomly rearranging the
feedback connection weights. This also resulted in a significant reduction in the RNNs'
performance accuracy for nonzero-coherence motion stimuli (**Fig. 5c**; two-way ANOVA:
P = 1.38e-90, F = 883.61) and prolonged RTs for high- and medium-coherence motion
stimuli (**Fig. 5d**; two-way ANOVA: P = 1.94e-12, F = 54.06). The changes of RNN units'
activity patterns after scrambling feedback connections were similar to the effects after
feedback connectivity was fully ablated: the difference in the motion DS between the CT
and IT conditions dramatically decreased (**Fig. 5h** and **Extended data fig. 10b,c and
13a,b,d,e**; two-way ANOVA: P = 5.81e-12, F = 51.52), and the weakened saccade DS in
the target module was no longer positively correlated with motion coherence (**Fig. 5k** and
Extended data fig. 13c,f; mean r = –0.03; one-sample t-test: P = 3.56e-5, t(49) = -4.55).

Third, we trained another 50 RNNs without feedback connectivity to learn the FVMD
task. Similar to the full-model RNNs, all 50 networks converged to perform the FVMD
task with high accuracies (>99%) after training and exhibited coherence-dependent
performances when tested with untrained stimuli (**Extended data fig. 14a,b**). Meanwhile,
units in the motion module and target modules exhibited coherence-dependent motion DS
and saccade DS, respectively (**Extended data fig. 14c,d**). Consistent with the above
connectivity ablation experiment, units in the motion module of these RNNs did not exhibit

different levels of motion DS between CT and IT conditions (**Extended data fig. 14e-g**;
two-way ANOVA: P = 0.63, F = 0.24). Importantly, the performance accuracies of these
RNNs decreased significantly when tested with untrained nonzero-coherence stimuli
(**Extended data fig. 14h**; two-way ANOVA: P = 1.46e-14, F = 65.64), as compared to the
full-model RNNs. Furthermore, we trained an additional 50 RNNs without feedback
connections to learn the FVMD task. These RNNs were initialized with greater recurrent
connection probabilities than the full-model RNNs, such that the number of total trainable
connection weights matched that in the full-model RNNs. Interestingly, these RNNs still
exhibited significantly lower performance accuracies when tested with high- and medium-
coherence stimuli relative to the full-model RNNs (**Extended data fig. 15**; two-way
ANOVA: P = 8.24e-6, F = 20.60). Together, these results suggested that the selectivity-
specialized feedback connections were important for the RNNs to efficiently solve flexible
decision tasks.

Discussion:

Here we show that primate LIP activity related to evaluating the visual motion stimulus
was precisely modulated by an impending saccade choice that was used to report decisions
during a FVMD task, even though the saccades were toward non-RF locations. This
suggests that the sensory evaluation and action selection may interact, and may even
proceed iteratively, during decision-making. This view is also consistent with the common
observation that action selection related activity arises at the early stage of evidence
accumulation during decision-making[3,9,32], as well as our observation that the modulation
of action selection on sensory evaluation emerged during the early task period even before
motion stimulus onset in the FVMD task. Brain areas that are related to different decision
processes, although spatially separated[14,21,31], likely form a real-time associated network to
solve current decision tasks. The instantaneous result of sensory evaluation may be
transmitted to the brain areas responsible for action selection during decision-making in a
feedforward manner, and the action selection process may also exert a real-time modulation
on the neural processing of sensory evaluation in a feedback manner.

The feedback modulation we observed during decision-making was distinct from
the modulation of feature-based attention in early sensory cortex. Although both
modulation effects match the stimulus tuning of recipient cells[37-41], only the feedback
modulation observed in the current study reflects the neural processing of decision-making.
This is because only the decision-related component, not the stimulus-related component,
of LIP activity was modulated by the monkeys' later saccade choice during the FVMD task
(**Fig. 3**). This suggests that the feedback modulation from action selection to LIP activity
was primarily related to the decision process rather than low-level sensory processing.

The neural selectivity-specialized feedback modulation from action selection to
sensory evaluation emerged in our multi-module RNNs during training, although the cross-
module connections were randomly initialized before training. Importantly, our causal in-

silico experiments demonstrated that these precise feedback connections played an essential role in mediating the modulation of action selection on sensory evaluation while solving the FVDM task. Previous studies showed that both the corticocortical feedback from the secondary visual cortex to the primary visual cortex and the corticogeniculate feedback in primates are organized into parallel streams resembling the reciprocal feedforward pathways[42-44], suggesting a potential functionally specialized feedback connection in visual processing. However, to the best of our knowledge, a similar segregation of feedback that reflects the encoding properties of the target neurons has not been evident in decision network. Therefore, it will be important for future studies to examine the connectivity and correlation of neural activity among neural groups related to different decision processes through sophisticated anatomical experimental approaches as well as multi-channel recordings in vivo.

Our brain includes extensive feedback connections across different brain areas, which, in some cases, even outnumber the feedforward connections[44,45]. However, compared to the feedforward connections, much less is known about the function of feedback connections. Feedback modulation has been implicated in top-down modulation of neuronal responses in the early sensory cortex, such as attention and expectation, which facilitate the processing of important stimuli and suppress distractors[46-48]. Furthermore, both experimental and computational studies have shown that weak decision-correlated neural activity (i.e., choice probability) as well as the correlated firing of pairs of neurons (i.e., noise correlation) in the early sensory cortex might partially result from feedback input in decision tasks[49-51], suggesting a potential role for feedback connection in modulating early sensory processing during decision-making. Consistently, selectivity-specialized feedback connections were important for our multi-module RNNs to efficiently solve the FVMD task, as the RNNs' decision behavior became more stochastic after ablating/disrupting the feedback connectivity. This was evident in the diminished behavioral performance of the RNNs and disrupted patterns of activity related to saccade choice in the target module. Our results suggest that selectivity-specialized feedback modulation might facilitate brain networks to efficiently generate flexible sensorimotor decisions based on sensory input.

Our results are consistent with predictive coding theories of feedback function, which propose that an internal model of the world is generated in the brain based on sensory data and prior experience and is refined by incoming sensory data[52-55]. In terms of the architecture of hierarchical predictive coding schemes, different neuronal ensembles encode different attributes/choices in either the action selection or sensory evaluation process at each level of the cortical hierarchy during flexible decision-making. Meanwhile, the conditionally independent expectations are functionally segregated, so that descending predictions from the action selection process are limited to the sensory evaluation process on the matched\associated stimulus attribute but not the opposite attribute. The selectivity specialized feedback connectivity in our RNNs fits the functional segregation of expectation very well during the interplay between distinct neural processes of decision-making. Specifically, the neural ensembles related to either sensory evaluation or action

selection processes for the same choice might form precise recurrently connected loops.
The prediction signal carried by the precise feedback connections might facilitate sensory
evaluation of the associated stimulus but prohibit evaluation of the nonmatched stimulus.
This could, in turn, amplify the task-relevant sensory input, speed up the sensorimotor
transformation, and ultimately result in faster and more accurate action selection. In future
work, it will be important to use techniques such as projection-specific inactivation and
microstimulation in vivo to test the causal contribution of feedback connectivity in
decision-making and other cognitive functions.

409
Online Methods

Behavioral task, stimulus display, and animal preparation

The flexible visual motion discrimination (FVMD) task (**Fig 1a**) has been reported
previously[14] and is briefly summarized below. In this task, monkeys were required to
saccade to either the green or red targets based on the direction of the visual motion
stimulus. Two motion directions (135°, 315°) were used, each with four different
coherence levels (0%, 9%, 18%, 36% for monkey M; and 0%, 13%, 25%, 50% for monkey
B) were tested. If the sample direction was 135°, the monkeys must saccade to the green
target to receive a juice reward, whereas the 315° direction was associated with the red
target. The rewarded target (red or green) was randomly chosen (with 50% probability) on
each zero-coherence trial. The positions of red and green targets were randomly chosen
between the two positions on each trial at each recording session. Therefore, there is no
fixed mapping between motion stimulus and saccade direction.

To initiate each trial, monkeys must hold a touch-bar and acquire gaze fixation.
They then needed to maintain fixation within a 2.0-2.5° window throughout the trial before
their saccade choice. After a 500ms fixation period, two colored saccade targets appeared
simultaneously at opposite positions relative to the fixation point with equal eccentricities
(8° and 9° for Monkey M and B, respectively). 400ms later, a sample motion stimulus was
presented at a location orthogonal to the axis of, but at the same eccentricity as, the saccade
targets. We used motion stimuli that were full contrast, 8° diameter, random-dot movies
composed of 190 dots per frame, and the dots moved at 10°/s. Monkeys needed to saccade
to either red or green targets within a 60-2000ms window after sample stimulus onset. A
juice reward would be delivered to the monkeys if they made correct saccade choice.

Two male monkeys (Macaca mulatta, 15~16 years old, 8–14 kg) were trained on
the FVMD task and implanted with a head post as well as a recording chamber positioned
over PPC. Our surgical, behavioral, and neurophysiological approach has been described
in detail in a previous study[14] . Stereotaxic coordinates for chamber placement were
determined from magnetic resonance imaging (MRI) scans obtained before chamber
implantation. LIP chambers were centered on the intraparietal sulcus (IPS), 4.0 mm
posterior to the intra-aural line and 1.0 mm lateral from the midline for monkey M, and 0

mm anterior to the intra-aural line and 15.0 mm lateral from the midline for monkey B.
Monkeys were housed in individual cages under a 12-hour light/dark cycle. Behavioral
training and experimental recordings were conducted during the light portion of the cycle.
Monkeys sat comfortably while head-fixed in a custom-made primate chair inside a dark
experiment rig. The task stimuli were displayed on a 21-inch color CRT monitor
(1280*1024 resolution, 75 Hz refresh rate, 57 cm viewing distance). Both monkeys were
tested with identical stimuli and timing. A solenoid-operated reward system was used to
deliver juice reward to the monkeys. Monkeys' eye positions were monitored by an optical
eye tracker (SR Research) at a sampling rate of 1 kHz and stored for offline analysis.
Stimulus presentation, task events, rewards, and behavioral data acquisition were
accomplished using an Intel-based PC equipped with MonkeyLogic software running in
MATLAB (http://www.monkeylogic.net). All experimental and surgical procedures were
in accordance with the University of Chicago Animal Care and Use Committee and
National Institutes of Health guidelines.

Electrophysiological recording

We used either 75-μm tungsten microelectrodes (FHC, ~1 MΩ) or 16-channel V-Probes
(Plexon) to record single neuron activity in LIP. Neurophysiological signals were amplified,
digitized, and stored for offline spike sorting (Plexon) to verify the quality and stability of
neuronal isolations. We recorded neuronal activity in a memory-guided saccade task to
map LIP RF locations before each FVMD recording session.

Our LIP recordings targeted different hemispheres in the two monkeys (monkey M:
left hemisphere; monkey Q: right hemisphere). Therefore, the contralateral target (CT)
condition and ipsilateral target (IT) conditions in **Fig 1f-g** referred to opposite saccade
directions (target locations) between the two monkeys. For monkey M, the CT condition
corresponded to the trials in which the correct saccade target was on the right visual field,
whereas the CT condition for monkey Q referred to the trials in which the correct saccade
was toward the left visual field. We localized LIP in each monkey according to the patterns
of neuronal activity in the MGS task (i.e., spatial selectivity during stimulus presentation
and delay). All neurons included in the dataset were recorded from the same locations (the
same grid holes and similar depths: ~5-10 mm from the cortical surface) where we
encountered spatial selectivity in the MGS task. LIP neurons were also identified based on
anatomical criteria, such as the location of each electrode track relative to that expected
from the MRI scans, the pattern of gray–white matter transitions encountered on each
electrode penetration, and the relative depths of each neuron.

We aimed to present the visual motion stimulus inside the RFs of the identified
neurons during each recording session. For single-channel electrode recording, only
neurons exhibited visual responses to the motion stimuli during prescreening (tested with
the FVMD task) were recorded with sufficient trials (~300-600) of the FVDM task. For
neurons exhibiting clear spatial RFs during the MGS task, we presented the motion
stimulus inside LIP neurons' RFs; while for those neurons which did not show a clear RF
during the MGS task, we presented motion stimuli in the positions (always in the visual

field contralateral to the recorded hemisphere) in which neurons exhibited the strongest
response to the motion stimuli. For the multi-channel recordings, we recorded all neurons
isolated across all channels and presented the motion stimulus inside one of the isolated
neurons' RFs. Because adjacent recording sites were located 100μm apart, nearby neurons
typically showed similar RF locations.

Data analysis

Neuronal pre-screening

While part of the neural data was presented in a previous report[14], the current analysis
focuses on a different phenomenon which was not examined previously. We included all
neurons recorded from single-channel electrodes for the analysis. For multi-channel
recordings, we only included the neurons which showed significant modulation (different
from fixation period activity, one-way ANOVA, $p < 0.01$) of their averaged activity across
all motion stimuli because the stimulus could not always be presented within the RF of all
the recorded neurons. The low-firing neurons whose maximum firing rates were less than
2.0 spikes/s (to the direction producing greater average responses) during stimulus
presentation were also excluded for further analysis. We then applied a one-way ANOVA
test to compare activity between the two different motion directions during the period
following motion stimulus onset (50-250 ms after motion stimulus onset) to select neurons
that showed significant motion direction selectivity (DS) during the decision period. Only
neurons that showed significant ($p < 0.01$) DS were used for further analysis.

Receiver operating characteristic (ROC) analysis

To quantify each neuron's DS in the FVMD task, we applied an ROC analysis to the
distribution of firing rates within sliding windows (100 ms width, 5 ms steps). The ROC
value, which ranges between 0.0 and 1.0, indicates the performance of an ideal observer in
assigning motion direction based on each neuron's trial-by-trial firing rates. Values of 1.0
and 0.0 correspond to perfect classification (i.e., strong DS), while a value of 0.5 indicates
chance-level classification performance (i.e., no DS). For trials with zero coherence motion,
we assigned direction labels on each trial based on the monkey's choice.

In **Fig 3**, in order to test whether DS reflected monkeys' trial-by-trial choices, we
used an ROC analysis to quantify whether LIP activity correlated with monkeys' trial-by-
trial categorical choices more than the physical direction of motion stimulus by analyzing
both correct and error trials. We only included low coherence trials in this analysis because
there were sufficient numbers of error trials (average performance: Monkey M: 74%
correct, Monkey B: 67% correct). LIP neuronal activity was analyzed by ROC according
to either the monkey's trial-by-trial categorical choices or the direction of the sample
stimulus on each trial. Furthermore, for this analysis, we only used neurons for which we
recorded sufficient trials (for both correct and error trials, N>4) for the low coherence
condition of each motion direction (n = 45).

Partial correlation analysis

We performed a partial correlation analysis to quantify the correlation between LIP neural
activity and the monkeys' trial-by-trial categorical choices or the physical motion direction
of the stimulus using all trials. For each trial, we obtained three parameters for the
calculation: the stimulus direction, the pre-choice neuronal activity, and the monkeys'
choice. We assigned the stimulus directions with different values for different directions
and coherence levels: positive and negative values were used for preferred and
nonpreferred directions, respectively; while 4, 2, 1 and 0 are used for code the high,
medium, low, and zero coherence levels. We also assigned different values to different
choice directions (-2 for choosing the preferred direction, and $+2$ for choosing the
nonpreferred direction). Two measures were then calculated: r stimulus = r (neuronal
activity, stimulus direction| choice direction), the partial correlation between neuronal
activity and stimulus direction given the monkeys' choices; and r choice = r (neuronal
activity, choice direction | stimulus direction), the partial correlation between neuronal
activity and the monkeys' categorical choice given the stimulus direction. In **Fig 3**, we used
the mean activity within a sliding window (100ms width, sliding with 5ms steps) for each
neuron to perform the partial correlation analysis.

dPCA analysis

We conducted demixed principal component analysis using the methodology and the code
from a previous study[56] (http://github.com/machenslab/dPCA) to reduce the
dimensionality of the population activity as the standard PCA and demixes all task
variables. Specifically, we tested how much each task variable (motion direction of sample
stimuli, saccade direction, motion-saccade interaction, timing) contributes to the LIP
population activity during the DMC task.

As demonstrated in the previous study, the dPCA finds separate decoder (F) and encoder
(D) matrices for each task variable ($\emptyset$) by minimizing the loss function:

$$L_{dPCA} = \sum_0 \left\| X_0 - F_0 D_0 X \right\|^2$$

546 where X is a linear decomposition of the data matrix, which contains the instantaneous firing rate
547 of the recorded neurons, into variable-specific averages:

$$X = \sum_0 X_0 + X_{noise}$$

Here, we decomposed the neural activities into four parts: condition-independent, stimulus-
dependent (two motion directions with four coherence levels), saccade-dependent (two
saccade directions), dependent on the stimulus-saccade interaction, and noise. The decoder
and encoder axes permit us to reduce the data into a few components capturing the majority
of the variance of the data dependent on each task variable.

Recurrent neural network (RNN) training

Network Implementation

We trained biologically inspired networks to perform the FVMD task using methodology similar to previous studies[34,36]. We implemented multiple modules in the networks through constraints on the input/output structure and the initial recurrent connectivity of the hidden layer. Specifically, we built excitatory-inhibitory networks with 200 hidden units divided into two equal-size main modules to simulate two brain hemispheres. Each main module was further divided into two nominal modules, with each one only receiving the visual motion input (motion module) or target color input (target module), respectively. This design was intended to simulate the two neuron populations whose RFs covered the motion stimulus or saccade target in the monkey electrophysiology experiment (**Fig 4a**). Every nominal module was allocated one quarter of the excitatory units (40) and one quarter of the inhibitory units (10) in the overall network to ensure that the modules did not differ in their balance of excitation/inhibition prior to training. Meanwhile, in order to simulate the local and long-range connection structures in the brain, we set different levels of recurrent connectivity within and between different modules: the local (connectivity within each nominal module) recurrent connection density (probability) was the highest (50%); the across-RF (connectivity between the two nominal modules within the same main module) connection density was set to be in the medium level (25%); the cross-hemisphere (connectivity between either the motion modules or the target modules across different main modules) connection density was the lowest (10%). Only excitatory neurons could have "cross-hemisphere" projections to the corresponding nominal module (e.g., from the motion module of the "left hemisphere" to the motion module of the "right hemisphere"). Only the connection weight within the hidden layer was updated during training following methods described previously[36]. In addition, the connection weights between network units are endowed with short-term synaptic plasticity, which is aimed to provide connection weights with activity-dependent fluctuation over short timescales within each trial.

The input to the network consisted of two parts: 9 motion input units and 8 target color input units. The preferred directions of motion input units were evenly spaced across 360°, with response tuning distributed according to a von Mises function. The value of the nth motion input unit was set to

$$r_n = \frac{2}{3}\left(ace^{\left(\kappa\cos(\theta-\theta_{pref})\right)} + b\mathcal{N}(0,2) + \sqrt{\frac{2}{\alpha}}\mathcal{N}(0,\sigma_{in}) \right) + d$$

, where $\alpha = \frac{\Delta t}{\tau_{mem}}$, θ is the direction of motion stimulus in radian, θ_{pref} is the preferred direction of this motion input unit, $a = \frac{4}{e^\kappa}$ after stimulus onset but 0 elsewhere, $c \in [0, 1]$ is the coherence of the dot motion stimulus, b is a binary value that equals 0 before stimulus onset and 1 after that, and d is a constant visual input signal with an amplitude of 2 that decays with time. Specifically, for zero-coherence input, since $ace^{\left(\kappa\cos(\theta-\theta_{pref})\right)}$ would be zero, we set this term as $0.4max(c)$.

The target input units were initially evenly divided into two groups projected to the
two target modules of hidden units. Each group was further divided into red and green
subgroups. The color subgroup was set as active with an amplitude of $\frac{4}{3}$ to represent the
color within the projected target module, while the other color group remained silent
(amplitude =0). An exponential decay filter was also used to fit the activity of sensory
neurons in the early sensory cortex across time:

$$f(x) = \left(\frac{1}{3}\right)^t x$$

, where x is the constant visual input for motion input and the color signal for color input,
t is time in seconds.

The two output units of the network simulate two different saccade directions. In
order to simulate the oculomotor control in the brain, each brain hemisphere (main module)
projects much more densely to the contralateral response units (probability = 0.32) than the
ipsilateral response units (probability = 0.08). To generate a probability distribution over
output values at each time point, we also applied a softmax function to scale the response
unit activities in the output layer.

To reduce the stochasticity of the network activity brought by input and output
connections, we randomly sampled input and output weights from a normal distribution
($\mathcal{N}(0.2, 0.05)$). In addition, the input weights were re-initialized if there were extreme
values (the minimum value was smaller than ¾ of the maximum value) after multiplying
the stimulus signal and weight of motion or target input. Similarly, the output weights were
re-initialized if there were extreme values in the output weight values. The input units had
excitatory and random projections to the recurrent units with a probability of 0.32.
However, only the excitatory units in the target modules of each "hemisphere" could
project to the output units. Both the input and output weights were fixed for each network
during task training.

Network training

The networks were trained using BrainPy[57] on an Intel(R) Core(TM) i9-9900K CPU
(3.60GHz, 8 cores). The network was optimized using backpropagation through time
(BPTT) and stochastic gradient descent with an Adam optimizer (default setting, first
moment estimates decay rate=0.9, second moment estimates decay rate =0.999) to
minimize a loss function. The network parameters (recurrent weights/biases) were
optimized to minimize a loss function with three parts as in a previous study[36]: (a) a
performance loss; (b) a metabolic cost on mean neuronal activity; and (c) a metabolic cost
on connectivity. The gradient was clipped to a maximum L2 value of 0.1 to avoid the
exploding gradient problem. During training, only the hidden unit related parameters
(weights, bias, and initial activities) were updated. The training process was terminated
when the network's performance accuracy reached 99% or till the maximum number of
iterations (2000).

During each trial of the training and testing of the FVMD task, networks were first
presented with two color targets through the target input units, and then presented with the
motion direction through the motion input units. Both the motion and target inputs persisted
until the end of each trial. A short time after the motion input (100ms, 5 time steps), the
networks were required to report the direction of the stimulus motion by choosing the
saccade target with the appropriate color. Specifically, each element of the task design in
the FVMD task that the models were trained to perform was tailored to match those used
in the monkey experiments: the directions of the motion stimuli, the target colors, as well
as the task (stimulus) durations were the same as that used in monkey electrophysiology
experiments. Trials were programmatically generated by constructing motion/target inputs
to the networks at each timestep and the desired response (left saccade, right saccade) at
each timestep. In total, we trained 50 networks to perform the FVDM task. All networks
were first trained using motion directions with two high coherence levels and were then
tested using motion stimuli with another four different coherence levels (high, medium,
low, and zero), which had never been presented during training. Besides the difference in
the coherence of motion input, the color inputs remained the same during the testing period.
All the networks achieved consistently high performance by the end of training (> 99%
accuracy).

Hyperparameter	Value	Hyperparameter	Value
Training batch size	1024	Decision threshold	0.8
Testing batch size	2048	Training stimuli coherence levels	[0.6, 0.9]
Learning rate	0.02	Testing stimuli coherence levels	[0, 0.35, 0.55, 0.75]
Number of input/hidden/output units	17(N_{in}), 200(N_{rec}), 2(N_{out})	Fixation/Target/Stimulus/Decision Time	500ms(T_{fix}), 400ms(T_{targ}), 500ms(T_{stim}), 100ms(T_{dec})
Time step	20	Spike cost	0.004
Membrane time constant	100	Weight cost	1
STSP neurotransmitter time constant	200ms / 1000ms (facilitating / depressing)	Input noise distribution std	0.07 (σ_{in})
STSP neurotransmitter utilization	2000ms / 200ms (facilitating / depressing)	Noise distribution std of hidden unit	0.08 (σ_{rec})

STSP neurotransmitter increment	0.1 / 0.3 (facilitating / depressing)

Table1. Important model hyperparameters

Quantification of the networks' behavioral performance

To test whether our multi-module RNNs exhibited decision behavior similar to monkey
subjects, we examined their performance accuracies and RTs when tested by novel stimuli
with different coherence levels. The performance accuracy was defined as whether the
output of the response units (starting from 100ms after stimulus onset to the trial end)
matched the desired output. Furthermore, we defined the networks' RTs as the time point
from which the differences between two output units were greater than a threshold (0.8)
for three consecutive time points after the stimulus onset. During some trials, the threshold
was never reached until the end of the trial (500ms after the motion stimulus onset). In
these cases, we artificially set the RT to be 600ms in these trials for further analysis.

Analysis of RNN activity

We performed the same analyses on the RNN units as we did on the
neurophysiology data. In **Fig 4-5**, data from both the example network and averaged results
across all the networks were shown. For every network, we only included the units which
exhibited at least one kind of task-related modulation during the test period (i.e., motion
direction selectivity or saccade direction selectivity, one-way ANOVA test, $p < 0.01$). In
order to quantify the activity related to evaluating motion stimulus, we analyzed the motion
DS only including the units in the motion modules, whereas only the saccade DS of units
in the target modules were calculated to quantify the activity related to saccade selection
during the decision-making.

Analysis of RNN connectivity

To examine the potential circuit mechanisms underlying the modulation of action
selection on sensory evaluation during decision-making in the RNNs, we examined the
cross-module connection weights in the trained networks. Specifically, we defined the
feedforward connection as the projection from units in the motion module to units in the
target module within the same main module (hemisphere). We also defined the feedback
connection as the projection from units in the target module to units in the motion module
within the same main module. Furthermore, we grouped the cross-module connections into
the selectivity-matched and selectivity-mismatched groups based on the preferences of the
neural encoding of units in the motion and target modules. The selectivity-matched group
included two types of unit pairs: 1) units in the motion module preferred to encode 315°
and units in the target module preferred to encode red, 2) units in the motion module
preferred to encode 135° and units in the target module preferred to encode green. The

selectivity-mismatched group includes the other two types of unit pairs: 3) units in the
motion module preferred to encode 315° while units in the target module preferred to
encode green, and 4) units in the motion module preferred to encode 135°while units in the
target module preferred to encode red. The values of connection weights projected from
excitatory and inhibitory units were set as positive and negative, respectively.

We also examined the correlation between the similarity of neural selectivity and
connection strength for different unit pairs in the RNNs based on the following steps: First,
both the preference and strength of motion direction encoding or target color encoding
were quantified for units in the motion module or target module, respectively. This was
done by calculating the averaged differential activity to the 315° and 135° motion stimuli
after motion stimulus onset or calculating the averaged differential activity to the red and
green targets after target onset. In this case, the unit pairs that exhibited matched selectivity
would show selectivity values with the same sign (positive or negative), whereas the unit
pairs that exhibited mismatched selectivity would show selectivity values with the opposite
signs. Second, the units in both the motion module and target modules were ranked based
on selectivity values. Third, the selectivity similarity of each cross-module unit pair was
quantified by calculating the reverse value of the absolute differences between the target
color selectivity rank and the motion direction selectivity rank. The rank of the weight
value of each pair was calculated and then reversed to give the lower weight value a higher
rank score. Finally, we calculated the Pearson correlation between weight rank and
selectivity similarity. A positive correlation indicates that units that exhibited stronger
encoding preference to one motion direction in the motion module were more likely to
connect with the units in the target module that exhibited stronger encoding on the matched
target color.

Inactivation experiments in silico

Our examination of the network connectivity within the hidden layer of the RNNs revealed
that the selectivity-specialized top-down connections originated from the target module to
the motion module. To assess the causal contribution of such precise feedback connections
to the modulation of action selection on sensory evaluation, we performed two in-silico
analogues of neuronal inactivation experiments similar to a previous study[36]. We
hypothesized that ablating or disrupting the precise feedback connection from the target
module to the motion module would significantly impact the activity patterns of the units
in the motion module. Specifically, such inactivation on connectivity was expected to
reduce the difference in motion DS in the notion module between CT and IT conditions.
To test this hypothesis, we performed the projection-specific inactivation experiments in
the RNNs in three different ways. First, we ablated all the feedback projections from the
target module to the motion module while keeping all the other network parameters
unchanged after the networks were fully trained. We then tested the networks with the
untrained motion stimuli used in the normal experiment. The connectivity ablation was
implemented by directly setting the connection weights from units in the target module to
units in the motion module to zero. Second, we randomly rearranged the connection

weights from the target module to the motion module after the network was fully trained by shuffling the corresponding weights in the weight matrix. Then, we tested the networks with the testing motion stimuli. This was aimed to disrupt the selectivity-specialized feedback connection without changing the total strength of the feedback connection. We repeated this random feedback weight rearrangement process 100 times for each of the 50 trained network. Subsequently, we measured the impact on the networks' behavior performance and the activity patterns in both the motion and saccade modules. In particular, for each network, we tested whether there were still different levels of motion direction selectivity of units in the motion module between CT and IT conditions after inactivating the feedback connectivity.

We further performed pattern-specific ablation of the feedback connections in the RNNs, in order to examine the causal roles of different patterns of feedback connectivity in solving the FVDM task. Specifically, we separated the feedback projections in each RNN into the specialized group and the nonspecialized group. The specialized group included two conditions: 1) the feedback connection weight was positive between the selectivity-matched unit pairs, and 2) the feedback connection weight was negative between the selectivity-nonmatched unit pairs; whereas the nonspecialized group included the rest of the feedback conditions in each RNN. We then tested the effects of the pattern-specific connectivity inactivation on the RNNs' behavior performance when tested with the untrained motion stimuli.

In the third inactivation experiment, we trained 50 additional RNNs without feedback connections to learn the FVMD task, in order to test the importance of feedback connectivity in solving the flexible decision task. These networks were initialized using identical hyperparameters and trained with the same methodology as the normal RNNs, except that the feedback connections from the target module to the motion module were ablated before training. We then examined these RNNs' behavior performance and unit activity when testing with the untrained motion stimuli used in the normal experiment. Furthermore, we trained another 50 RNNs without feedback connection to learn the FVMD task. These RNNs were initialized with higher recurrent connection probabilities than the full-model RNNs to keep the number of the total trainable connection weight comparable to that in the full-model RNNs. We raised the recurrent connection probability by a factor of 1.176 after removing feedback connections (average number of connections in 50 normal RNNs = 8193, average number of connections in 50 networks without feedback = 8184, $p = 0.53$, independent sample t-test). Except for the recurrent connection probability, all the other hyperparameters and the training methodology were the same as the normal RNNs. We then compared the behavior performance of these RNNs with that of the normal RNNS when testing with the untrained motion stimuli.

Acknowledgements:

We thank Tianming Yang, Matthew Rosen, Cheng Xu, Gongcheng Yu, Mingze Li and Ou Zhu for their comments on an earlier version of this manuscript. We also thank the veterinary staff at The University of Chicago Animal Resources Center for expert assistance.

Funding:

This study was supported by STI2030-Major Projects (2021ZD0203800, NSF32171036.

Author contributions:

Y.Z. designed the experiments and wrote the manuscript. Y.Z. trained monkeys and collected the behavioral and neuronal data, Y.Z. and X.W. analyzed the data and made the figures. Y.Z. and X.W. designed the network models, X.W. trained the RNNs, analyzed the RNNs and implemented the RNN inactivation experiment. Y.Z. supervised the project.

Competing interests:

The authors declare no competing financial interests.

Reference:

O'Connell, R. G., Shadlen, M. N., Wong-Lin, K. & Kelly, S. P. Bridging Neural and Computational Viewpoints on Perceptual Decision-Making. *Trends Neurosci* **41**, 838-852, doi:10.1016/j.tins.2018.06.005 (2018).

Freedman, D. J. & Assad, J. A. Neuronal Mechanisms of Visual Categorization: An Abstract View on Decision Making. *Annu Rev Neurosci* **39**, 129-147, doi:10.1146/annurev-neuro-071714-033919 (2016).

Gold, J. I. & Shadlen, M. N. The neural basis of decision making. *Annu Rev Neurosci* **30**, 535-574, doi:10.1146/annurev.neuro.29.051605.113038 (2007).

Huk, A. C., Katz, L. N. & Yates, J. L. The Role of the Lateral Intraparietal Area in (the Study of) Decision Making. *Annu Rev Neurosci* **40**, 349-372, doi:10.1146/annurev-neuro-072116-031508 (2017).

Hanks, T. D. & Summerfield, C. Perceptual Decision Making in Rodents, Monkeys, and Humans. *Neuron* **93**, 15-31, doi:10.1016/j.neuron.2016.12.003 (2017).

Shadlen, M. N. & Kiani, R. Decision making as a window on cognition. *Neuron* **80**, 791-806, doi:10.1016/j.neuron.2013.10.047 (2013).

Freedman, D. J. & Assad, J. A. A proposed common neural mechanism for categorization and perceptual decisions. *Nat Neurosci* **14**, 143-146, doi:10.1038/nn.2740 (2011).

Platt, M. L. & Glimcher, P. W. Neural correlates of decision variables in parietal cortex. *Nature* **400**, 233-238, doi:10.1038/22268 (1999).

9   Roitman, J. D. & Shadlen, M. N. Response of neurons in the lateral intraparietal area
during a combined visual discrimination reaction time task. *J Neurosci* **22**, 9475-9489
(2002).
10   Shadlen, M. N. & Newsome, W. T. Motion perception: seeing and deciding. *Proceedings*
of the National Academy of Sciences of the United States of America **93**, 628-633,
doi:10.1073/pnas.93.2.628 (1996).
11   Shadlen, M. N. & Newsome, W. T. Neural basis of a perceptual decision in the parietal
cortex (area LIP) of the rhesus monkey. *J Neurophysiol* **86**, 1916-1936,
doi:10.1152/jn.2001.86.4.1916 (2001).
12   Sugrue, L. P., Corrado, G. S. & Newsome, W. T. Matching behavior and the
representation of value in the parietal cortex. *Science* **304**, 1782-1787,
doi:10.1126/science.1094765 (2004).
13   Yates, J. L., Park, I. M., Katz, L. N., Pillow, J. W. & Huk, A. C. Functional dissection of signal
and noise in MT and LIP during decision-making. *Nat Neurosci* **20**, 1285-1292,
doi:10.1038/nn.4611 (2017).
14   Zhou, Y. & Freedman, D. J. Posterior parietal cortex plays a causal role in perceptual and
categorical decisions. *Science* **365**, 180-185, doi:10.1126/science.aaw8347 (2019).
15   Freedman, D. J. & Assad, J. A. Experience-dependent representation of visual categories
in parietal cortex. *Nature* **443**, 85-88, doi:10.1038/nature05078 (2006).
16   Freedman, D. J. & Assad, J. A. Distinct encoding of spatial and nonspatial visual
information in parietal cortex. *J Neurosci* **29**, 5671-5680, doi:10.1523/JNEUROSCI.2878-
08.2009 (2009).
17   Bennur, S. & Gold, J. I. Distinct representations of a perceptual decision and the
associated oculomotor plan in the monkey lateral intraparietal area. *J Neurosci* **31**, 913-
921, doi:10.1523/JNEUROSCI.4417-10.2011 (2011).
18   Swaminathan, S. K. & Freedman, D. J. Preferential encoding of visual categories in
parietal cortex compared with prefrontal cortex. *Nat Neurosci* **15**, 315-320,
doi:10.1038/nn.3016 (2012).
19   Zhou, Y., Mohan, K. & Freedman, D. J. Abstract Encoding of Categorical Decisions in
Medial Superior Temporal and Lateral Intraparietal Cortices. *J Neurosci* **42**, 9069-9081,
doi:10.1523/JNEUROSCI.0017-22.2022 (2022).
20   Kiani, R. & Shadlen, M. N. Representation of confidence associated with a decision by
neurons in the parietal cortex. *Science* **324**, 759-764, doi:10.1126/science.1169405
(2009).
21   Katz, L. N., Yates, J. L., Pillow, J. W. & Huk, A. C. Dissociated functional significance of
decision-related activity in the primate dorsal stream. *Nature* **535**, 285-288,
doi:10.1038/nature18617 (2016).
22   Kim, J. N. & Shadlen, M. N. Neural correlates of a decision in the dorsolateral prefrontal
cortex of the macaque. *Nat Neurosci* **2**, 176-185, doi:10.1038/5739 (1999).
23   Rossi-Pool, R. *et al.* Decoding a Decision Process in the Neuronal Population of Dorsal
Premotor Cortex. *Neuron* **96**, 1432-1446 e1437, doi:10.1016/j.neuron.2017.11.023
(2017).
24   Ding, L. & Gold, J. I. Neural correlates of perceptual decision making before, during, and
after decision commitment in monkey frontal eye field. *Cereb Cortex* **22**, 1052-1067,
doi:10.1093/cercor/bhr178 (2012).
25   Ferrera, V. P., Yanike, M. & Cassanello, C. Frontal eye field neurons signal changes in
decision criteria. *Nat Neurosci* **12**, 1458-1462, doi:10.1038/nn.2434 (2009).

26   Heitz, R. P. & Schall, J. D. Neural mechanisms of speed-accuracy tradeoff. *Neuron* **76**,
616-628, doi:10.1016/j.neuron.2012.08.030 (2012).
27   Horwitz, G. D. & Newsome, W. T. Separate signals for target selection and movement
specification in the superior colliculus. *Science* **284**, 1158-1161,
doi:10.1126/science.284.5417.1158 (1999).
28   Peysakhovich, B. *et al.* Primate superior colliculus is engaged in abstract higher-order
cognition. *bioRxiv*, doi:10.1101/2023.01.17.524416 (2023).
29   Ding, L. & Gold, J. I. Caudate encodes multiple computations for perceptual decisions. *J*
Neurosci **30**, 15747-15759, doi:10.1523/JNEUROSCI.2894-10.2010 (2010).
30   Gold, J. I. & Shadlen, M. N. Representation of a perceptual decision in developing
oculomotor commands. *Nature* **404**, 390-394, doi:10.1038/35006062 (2000).
31   Jun, E. J. *et al.* Causal role for the primate superior colliculus in the computation of
evidence for perceptual decisions. *Nat Neurosci* **24**, 1121-1131, doi:10.1038/s41593-
021-00878-6 (2021).
32   Shushruth, S., Mazurek, M. & Shadlen, M. N. Comparison of Decision-Related Signals in
Sensory and Motor Preparatory Responses of Neurons in Area LIP. *J Neurosci* **38**, 6350-
6365, doi:10.1523/JNEUROSCI.0668-18.2018 (2018).
33   Engel, T. A., Chaisangmongkon, W., Freedman, D. J. & Wang, X. J. Choice-correlated
activity fluctuations underlie learning of neuronal category representation. *Nat Commun*
6, 6454, doi:10.1038/ncomms7454 (2015).
34   Masse, N. Y., Yang, G. R., Song, H. F., Wang, X. J. & Freedman, D. J. Circuit mechanisms
for the maintenance and manipulation of information in working memory. *Nat Neurosci*
22, 1159-1167, doi:10.1038/s41593-019-0414-3 (2019).
35   Song, H. F., Yang, G. R. & Wang, X. J. Training Excitatory-Inhibitory Recurrent Neural
Networks for Cognitive Tasks: A Simple and Flexible Framework. *PLoS Comput Biol* **12**,
e1004792, doi:10.1371/journal.pcbi.1004792 (2016).
36   Zhou, Y. *et al.* Distributed functions of prefrontal and parietal cortices during sequential
categorical decisions. *Elife* **10**, doi:10.7554/eLife.58782 (2021).
37   Motter, B. C. Neural correlates of attentive selection for color or luminance in
extrastriate area V4. *J Neurosci* **14**, 2178-2189, doi:10.1523/JNEUROSCI.14-04-
02178.1994 (1994).
38   Motter, B. C. Neural correlates of feature selective memory and pop-out in extrastriate
area V4. *J Neurosci* **14**, 2190-2199, doi:10.1523/JNEUROSCI.14-04-02190.1994 (1994).
39   Martinez-Trujillo, J. C. & Treue, S. Feature-based attention increases the selectivity of
population responses in primate visual cortex. *Curr Biol* **14**, 744-751,
doi:10.1016/j.cub.2004.04.028 (2004).
40   Bichot, N. P., Rossi, A. F. & Desimone, R. Parallel and serial neural mechanisms for visual
search in macaque area V4. *Science* **308**, 529-534, doi:10.1126/science.1109676 (2005).
41   Maunsell, J. H. & Treue, S. Feature-based attention in visual cortex. *Trends Neurosci* **29**,
317-322, doi:10.1016/j.tins.2006.04.001 (2006).
42   Federer, F., Ta'afua, S., Merlin, S., Hassanpour, M. S. & Angelucci, A. Stream-specific
feedback inputs to the primate primary visual cortex. *Nat Commun* **12**, 228,
doi:10.1038/s41467-020-20505-5 (2021).
43   Briggs, F. & Usrey, W. M. Parallel processing in the corticogeniculate pathway of the
macaque monkey. *Neuron* **62**, 135-146, doi:10.1016/j.neuron.2009.02.024 (2009).
44   Briggs, F. Role of Feedback Connections in Central Visual Processing. *Annu Rev Vis Sci* **6**,
313-334, doi:10.1146/annurev-vision-121219-081716 (2020).

45   Harris, K. D. & Mrsic-Flogel, T. D. Cortical connectivity and sensory coding. *Nature* **503**,
51-58, doi:10.1038/nature12654 (2013).
46   Moore, T. & Armstrong, K. M. Selective gating of visual signals by microstimulation of
frontal cortex. *Nature* **421**, 370-373, doi:10.1038/nature01341 (2003).
47   Gilbert, C. D. & Li, W. Top-down influences on visual processing. *Nature reviews.*
Neuroscience **14**, 350-363, doi:10.1038/nrn3476 (2013).
48   McManus, J. N., Li, W. & Gilbert, C. D. Adaptive shape processing in primary visual
cortex. *Proceedings of the National Academy of Sciences of the United States of America*
108, 9739-9746, doi:10.1073/pnas.1105855108 (2011).
49   Nienborg, H. & Cumming, B. G. Decision-related activity in sensory neurons reflects
more than a neuron's causal effect. *Nature* **459**, 89-92, doi:10.1038/nature07821
(2009).
50   Bondy, A. G., Haefner, R. M. & Cumming, B. G. Feedback determines the structure of
correlated variability in primary visual cortex. *Nat Neurosci* **21**, 598-606,
doi:10.1038/s41593-018-0089-1 (2018).
51   Wimmer, K. *et al.* Sensory integration dynamics in a hierarchical network explains choice
probabilities in cortical area MT. *Nat Commun* **6**, 6177, doi:10.1038/ncomms7177
(2015).
52   Rao, R. P. & Ballard, D. H. Predictive coding in the visual cortex: a functional
interpretation of some extra-classical receptive-field effects. *Nat Neurosci* **2**, 79-87,
doi:10.1038/4580 (1999).
53   Friston, K. & Kiebel, S. Predictive coding under the free-energy principle. *Philos Trans R*
Soc Lond B Biol Sci **364**, 1211-1221, doi:10.1098/rstb.2008.0300 (2009).
54   Bastos, A. M. *et al.* Canonical microcircuits for predictive coding. *Neuron* **76**, 695-711,
doi:10.1016/j.neuron.2012.10.038 (2012).
55   Shipp, S. Neural Elements for Predictive Coding. *Front Psychol* **7**, 1792,
doi:10.3389/fpsyg.2016.01792 (2016).
56   Kobak, D. *et al.* Demixed principal component analysis of neural population data. *Elife* **5**,
doi:10.7554/eLife.10989 (2016).
57   Chaoming Wang, X. C., Tianqiu Zhang, Si Wu. BrainPy: a flexible, integrative, efficient,
and extensible framework towards general-purpose brain dynamics programming.
bioRxiv, doi:https://doi.org/10.1101/2022.10.28.514024 (2022).

Fig. 1. Behavioral task. **a**, The flexible visual-motion discrimination (FVMD) task. Monkeys needed to report their decision about the direction of the visual motion stimuli by choosing either the green or red saccade target. The appearance of the two color targets preceded that of the visual-motion stimulus, and the target positions were randomly chosen on each trial to avoid fixed mapping between motion stimulus and saccade direction. Monkeys could initiate their saccade as soon as they had made their decision. **b-c**, Psychometric curves for the two monkeys. The performance accuracy for each monkey is plotted as the proportion of trials in which 315° was chosen as a function of the directions and coherence levels of the motion stimuli. **d-e**, Chronometric curves are shown separately for the two monkeys. **f-g**, Two trial conditions, contralateral (**f**) and ipsilateral (**g**), defined according to the spatial configurations of task stimuli during neural recording.

Fig. 2. LIP activity related to evaluating motion stimuli was modulated by monkeys' saccade choice towards non-RF locations. **a**, The activity of one example neuron in the CT condition of the FVMD task is shown for each motion-coherence level. The zero-coherence trials were grouped based on the monkey's choices. The two vertical dashed lines denote the time of target and motion stimulus onset, respectively. **b**, The same example neuron's activity in the IT condition of the FVMD task. **c-f**, The activities of two more example neurons. **g-i**, The averaged population activities of all direction-selective neurons (n = 83) that were collected during the recording sessions in which the saccade targets were arranged in either horizontal or oblique directions. The activity to each motion direction and coherence level is shown separately for the CT condition (**g**) and IT condition (**h**). In (**i**), the mean activity (averaged across all motion directions and coherence levels) was compared between the two saccade directions (CT vs. IT) at the population level. **j**, An ROC analysis was used for quantifying the motion DS for both CT (solid) and IT (dashed) conditions. The colored dots denote the time points for which there was a significant difference between the CT and IT conditions (paired t-test: P < 0.01). **k**, The average DS for low- and zero-coherence trials is shown as in j. **l**, Variance in LIP population activity as explained by the individual demixed principal components. Each bar shows the proportion of total explained variance that was contributed by the four task variables. The pie chart shows the total variance explained by each task variable. H, high; M, medium; L, low; 0, zero; P, preferred; NP nonpreferred.

Fig. 3. LIP DS was more decision-correlated in the CT condition than in the IT condition. **a-b**, Averaged population activities on low-coherence trials in the CT condition (**a**) and IT condition (**b**) are shown separately for correct (corr) and error (err) trials. **c-d**, An ROC analysis quantified the stimulus-related and decision-related LIP DS on low-coherence trials. The correlations between LIP neural activity and the monkeys' decisions about motion direction (red) or the physical direction of the motion stimulus (blue) in both the CT (**c**) and IT (**d**) conditions are shown over time. **e-f**, Partial correlation analysis revealed the decision-related and stimulus-related components of LIP activity. The values for r-decision (**e**) (the partial correlation between neuronal activity and monkeys' choice, given the stimulus direction) and r-stimulus (**f**) (partial correlation between neuronal activity and stimulus direction, given the monkeys' choices) were compared between the IT and CT conditions. **g-h**, Correlation between LIP DS and the monkeys' RTs on zero-coherence trials. The choice selectivity on zero-coherence trials is shown for the faster RT and slower RT trials. Shaded areas denote ±SEM. The black stars indicate time periods in which there was a significant difference (paired t-test: P < 0.01).

Fig. 4. Multi-module RNNs trained with the FVMD task. **a**, Model schematic of the RNNs. Each RNN consists of nine motion direction tuned input units, eight color tuned target input units, 200 hidden units, and two response units. The hidden layer of each RNN consists of two main modules. Each main module consists of two nominal modules, each of which receives either the visual motion input (motion module) or the target color input (target module). Only the target modules project to the two response units, and each main module projects primarily to one response unit. All four nominal modules were assigned with equal number of units (25%) in the network, which consisted of 80% excitatory and 20% inhibitory units. **b-c**, The performance accuracies (**b**) and RTs (**c**) of all 50 trained RNNs are shown separately for each motion coherence level. **d-e**, Two example units from an example RNN. **d**, The neural activity of an example unit from the motion module is shown for each motion direction and coherence level. **e**, The neural activity of an example unit from the target module is shown for each saccade direction and coherence level. **f-g**, The motion DS for the motion module (**f**) and saccade DS for the target module (**g**) for the example RNN. **h-i**, The averaged population activities of all direction-selective units in the motion module of the example RNN are shown for the CT condition (**h**) and IT condition (**i**). **j-k**, The averaged motion DS in the motion module of the example RNN for both the CT (solid) and IT (dashed) conditions was quantified by ROC analysis. **l-m**, Partial correlation analysis. The values for r-decision (**l**) and r-stimulus (**m**) are compared between the IT and CT conditions for the example RNN. **n**, A comparison of the motion DS in the motion module of all trained RNNs between the CT and IT conditions is shown for each coherence level. **o**, A comparison of the r-decision and r-stimulus between the IT and CT conditions is shown for all trained RNNs. (Paired t-test: ***, P < 0.001).

Fig. 5. The circuit mechanisms underlying the interaction between different decision processes in RNNs. **a**, The averaged cross-module connection weights are shown for both feedforward and feedback connections. Units in the motion module (M) and target module (T) were grouped based on their preferences for motion DS (315° vs. 135°) and target color selectivity (red [r] vs. green [g]), respectively. **b**, The correlation between the match extent of the neural encoding between units in the motion module and those in target modules and the connection weights between them. Each dot denotes data from one RNN. **c-d**, A comparison of the performance accuracy (**c**) and RT (**d**) of the full-model RNNs, RNNs without feedback connections, and RNNs with shuffled feedback connections. **e-f**, The averaged activity of units in the motion module of the example RNN after the feedback connections were ablated is shown for CT (**e**) and IT (**f**) conditions. **g**, The comparison of the averaged motion DS between CT and IT conditions for all trained RNNs after feedback connections were ablated. An ROC analysis was used to quantify the motion DS. **h**, The differences in motion DS between CT and IT conditions were compared between full-model RNNs and RNNs with shuffled feedback connectivity. **i**, The averaged activity of units in the target module of the example RNN after the feedback connections were ablated is shown separately for different saccade directions and coherence levels. **j**, The averaged saccade DS in the target module of the example RNN after the feedback connections were ablated. **k**, The averaged saccade DS in the target module of the full-model RNNs, RNNs without feedback connections, and RNNs with shuffled feedback connections. (Paired t-test: *, P < 0.05; **, P < 0.01; ***, P < 0.001; n.s., not significant).